Computational Model to Quantify the Growth of Antibiotic Resistant Bacteria in Wastewater

Indorica Sutradhar[1] Carly Ching[1], Darash Desai[1], Mark Suprenant[1], Emma Briars[2,3], Zachary Heins[1,3], Ahmad S. Khalil[1,3,4] & Muhammad H. Zaman[1, 5*]

[1]Department of Biomedical Engineering, Boston University, Boston, MA 02215, US;

[2]Department of Bioinformatics, Boston University, Boston, MA 02215, US;

[3]Biological Design Center, Boston University, Boston, MA 02215, US;

[4]Wyss Institute for Biologically Inspired Engineering, Harvard University, Boston, MA 02115, US;

[5]Howard Hughes Medical Institute, Boston University, Boston, MA 02215, USA

*Corresponding author: zaman@bu.edu

Abstract

Although wastewater and sewage systems are known to be significant reservoirs of
antibiotic resistant bacterial populations and periodic outbreaks of drug resistant infection, there
is little quantitative understanding of the drivers behind resistant population growth in these
settings. In order to fill this gap in quantitative understanding of the development of antibiotic
resistant infections in wastewater, we have developed a mathematical model synthesizing many
known drivers of antibiotic resistance in these settings to help predict the growth of resistant
populations in different environmental scenarios. A number of these drivers of drug resistant
infection outbreak including antibiotic residue concentration, antibiotic interaction, chromosomal
mutation and horizontal gene transfer, have not previously been integrated into a single
computational model. We validated the outputs of the model with quantitative studies conducted
on the eVOLVER continuous culture platform. Our integrated model shows that low levels of
antibiotic residues present in wastewater can lead to increased development of resistant
populations, and the dominant mechanism of resistance acquisition in these populations is
horizontal gene transfer rather than acquisition of chromosomal mutations. Additionally, we
found that synergistic antibiotic interactions lead to increased resistant population growth. These
findings, consistent with recent experimental and field studies, provide new quantitative
knowledge on the evolution of antibiotic resistant bacterial reservoirs, and the model developed
herein can be adapted for use as a prediction tool in public health policy making, particularly in
low income settings where water sanitation issues remain widespread and disease outbreaks
continue to undermine public health efforts.

Importance

The rate at which antimicrobial resistance (AMR) has developed and spread throughout the world has increased in recent years, and according to the Review on Antimicrobial Resistance in 2014 it is suggested that the current rate will lead several million people AMR-related deaths by 2050[25]. One major reservoir of resistant bacterial populations that has been linked to outbreaks of drug resistant bacterial infections, but is not well understood, is in wastewater settings, where antibiotic pollution is often present. Using ordinary differential equations incorporating several known drivers of resistance in wastewater, we find that interactions between antibiotic residues and horizontal gene transfer significantly affect the growth of resistant bacterial reservoirs.

Introduction

Wastewater and sewage systems are a major reservoir of resistant bacterial populations due to the collection of antibiotic waste from humans and animals, inappropriate drug disposal, and effluent from drug manufacturers, hospitals and agricultural/veterinary settings (1). In some cases, environmental concentrations can reach, or even exceed, minimal inhibitory concentrations of certain antibiotics (2). This problem is particularly pertinent in low- and middle-income countries (LMICs) where cases of antibiotic resistant infections have been rising and 70% of sewage produced is estimated to enter the environment untreated (1). For example, in 2016, an outbreak of extensively drug resistant (XDR) typhoid cases emerged in in the southern Sindh province of Pakistan, and geospatial mapping revealed that the XDR *Salmonella* Typhi infections spread around sewage lines in the city of Hyderabad (3). Such outbreaks put the health and safety of surrounding populations at risk and, due to the communicable nature of these infections, increase risk for populations beyond the immediate vicinity of wastewater and sewage lines. Antibiotic pollution in wastewater and sewage systems results in a complex environment with many interacting antibiotic residues and this pollution is a major driver of the antibiotic resistance that is leading to outbreaks in surrounding communities (4, 5). One reason for this is that selective pressure from antibiotics in the environment is known to promote chromosomal resistance mutations (2, 6, 7). Furthermore, the interactions between different antibiotic residues may also affect resistance due to the synergistic and antagonistic effects changing the selective pressure on the bacteria (8). Additionally, the concentrations of antibiotic residues and disinfectants in sewage may be able to support and promote horizontal transfer of resistance genes among bacteria (9-12). These mobile resistance genes have previously been measured at high levels in wastewater and sewage (2). Though AMR and its drivers have been studied in sewage and wastewater settings to some extent (2, 12, 13), one of the largest gaps in understanding the emergence of AMR within a sewage environment is the limited quantitative understanding of the effects and interactions of the many biological and environmental mechanisms at work (14). Quantitative understanding of resistant population growth in wastewater settings is critical in order to predict the development of large resistant populations in wastewater that may pose a risk to local populations of resistant infection outbreak. Furthermore, quantitative tools for understanding of resistance can be used to develop and model strategies for preventing the emergence of large resistant populations and therefore influence policy decisions. Therefore, there is a critical need for developing quantitative methods of probing AMR in wastewater environments. Our study is a step in filling this gap in knowledge.

Mathematical modelling has been an important tool in quantitatively studying AMR development at both an epidemiologic and mechanistic level (6, 8, 12, 15-17). Epidemiological studies have included approaches that investigate bacterial population dynamics in a number of biological contexts including biofilms (15). At the population level, much of AMR modeling deals with disease states in which resistance patterns are long established, with significant gaps in the study of resistance emergence in new pathogens and the role of environmental factors in the development of AMR (16). At the mechanistic level, modelling has been used as a tool to

understand the individual roles of both mutational and plasmid-mediated resistances (8, 12).
Models based on resistance developed from chromosomal mutations have been used to study the
role of synergistic and antagonistic antibiotic interactions on the emergence of resistant
populations, finding that increased synergy increases the likelihood of resistance acquisition (8).
Additionally, studies using modelling as a tool to understand horizontal gene transfer (HGT) as a
driver of AMR have found HGT, specifically through the mechanism of conjugation, to be a
significant mode of resistance acquisition for *Escherichia coli* in settings including agricultural
slurry (12). However, much of the work done recently has been focused on fitting data in the
absence of antibiotic exposure (17). Understanding how bacteria evolve upon antibiotic exposure
is important to develop strategies that prevent the emergence of resistance (6). Thus, there
remains a need for quantitative modeling of both chromosomal mutation acquisition and HGT
under selective pressure from antibiotics to fully understand AMR development in settings such
as sewage and wastewater, where antibiotic residues are often present. Here we present a model
of emergence and growth of drug resistant bacteria in wastewater, integrating acquisition of
resistance through both chromosomal mutations and HGT while also incorporating the effects of
antibiotic interactions.
99
Methods
Our mathematical model of the growth of antibiotic resistant bacterial populations in wastewater
builds on prior studies (8, 12, 18) and extends it to incorporate a variety of critical, but overlooked,
input factors specific to the bacterial species, antibiotics and environments of interest. The inputs
can be broadly classified into bacterial parameters, environmental parameters and antibiotic
parameters. Bacteria specific input factors include the growth rates of antibiotic susceptible and
resistant strains, mutation rates, and rates of HGT. The antibiotic specific inputs, such as
bactericidal activity and degree of synergy, allow for the study of the effects of drug quality and
antibiotic pollution on the development of resistance. Additionally, environmental inputs,
including physical fluid inflow and outflow rates and antibiotic residue concentration, allow for
the modelling of resistance development in a variety of settings of interest. These input parameters
can be used to model an output of resistant bacterial populations over time, thus allowing for the
prediction of resistant population development (Figure 1).

Figure 1. Inputs and outputs for preliminary model of antibiotic resistance development in a single bacterial species in a wastewater setting

The model consists of ordinary differential equations governing the concentration of two antibiotics (C_1 and C_2) over time as well as equations modelling the susceptible population (*S*) and populations resistant to both antibiotics 1 and 2 from chromosomal mutation (R_m) or from HGT, specifically a resistance-conferring plasmid (R_p) over time (Eq set 1) (Table 1). Additionally, two populations of bacteria that are resistant through chromosomal mutation to each antibiotic individually are modelled (R_1 and R_2). Each antibiotic is modelled with terms for the antibiotic residue concentration in the environment (*E*) and the antibiotic clearance rate (k_e). Each bacterial population is modelled with terms for growth rate (α), which is limited by the carrying capacity (N_{max}). The bactericidal activities of the antibiotics are modelled by the term for killing rate in response to each antibiotic ($\delta_{max,1}$ and $\delta_{max,2}$). These killing terms are modified by the antibiotic concentration where the killing action is half its maximum value for either susceptibility (C_S^{50}) or resistance to each antibiotic ($C_{R,1}^{50}$, $C_{R,2}^{50}$). These killing rates and half max concentration terms are assumed to be equal for each antibiotic. Additionally, the killing rates are modified by a synergy term (*syn*) with *syn*<1 indicating antagonistic interaction and *syn*>1 indicating synergistic interaction. Also included in the model are terms for bacterial inflow (g_s, g_{Rm}, g_{R1}, g_{R2}, g_{Rp}) and outflow (k_T) rates (Table 2). These inflow and outflow rates are determined by the physical flow rates of the system of interest, and the outflow terms were validated experimentally as described below. As susceptible and resistant bacterial inflow and outflow rates for wastewater settings are not known, the model simulations assume bacterial inflow and outflow rates of zero. However, these parameters are still included due to their relevance for the modeling of systems such as

wastewater where bacterial inflow and outflow will affect resistant population development and
they may be quantified in future field studies. Chromosomal mutation is modelled through terms
for mutation rates under antibiotic pressure (m) which are concentration dependent. Parameters
governing mutation rate and antibiotic interaction are based on a previously developed model from
Michel et al (8). HGT is modelled assuming the mechanism of plasmid conjugation and a HGT
rate (β). HGT model terms and parameter values are adapted from a model of HGT in agricultural
waste from Baker et al (12). Other model parameters are based on experimentally derived
parameters of *E.* coli in piglet studies (18, 19). We note that the incorporation of all of these
parameters, which have previously not been studied at once, into one single model can provide
insights into their roles in the emergence and development of resistant populations. These
equations were coded and solved in MATLAB (R2018a, Mathworks Inc).

$$1)\quad \frac{dC_1}{dt} = E_1 - k_e C_1$$

$$2)\quad \frac{dC_2}{dt} = E_2 - k_e C_2$$

$$3)\quad \frac{dS}{dt} = \alpha_S \left(1 - \frac{R_m + R_1 + R_2 + R_p + S}{N_{max}}\right) S + g_s - k_T S - syn * \delta_{max,1} \left(\frac{C_1}{C_1 + C_S^{50}}\right) S -$$
$$syn * \delta_{max,2} \left(\frac{C_2}{C_2 + C_S^{50}}\right) S - \frac{\beta S R_p}{R_m + R_p + S}$$

$$4)\quad \frac{dR_m}{dt} = \alpha_R \left(1 - \frac{R_m + R_1 + R_2 + R_p + S}{N_{max}}\right) R_m + g_{Rm} - k_T R_m - syn * \delta_{max,1} \left(\frac{C_1}{C_1 + C_{R,1}^{50}}\right) R_m -$$
$$syn * \delta_{max,2} \left(\frac{C_2}{C_2 + C_{R,2}^{50}}\right) R_m + syn * m_T(C_1, C_2) S + R_1 m_2(C_2) + R_2 m_1(C_1)$$

$$5)\quad \frac{dR_1}{dt} = \alpha_{R,1} \left(1 - \frac{R_m + R_1 + R_2 + R_p + S}{N_{max}}\right) R_1 + g_{R1} - k_T R_1 - syn * \delta_{max,1} \left(\frac{C_1}{C_1 + C_{R,1}^{50}}\right) R_1 -$$
$$syn * \delta_{max,2} \left(\frac{C_2}{C_2 + C_S^{50}}\right) R_1 + m_1(C_1) S$$

$$6)\quad \frac{dR_2}{dt} = \alpha_{R,2} \left(1 - \frac{R_m + R_1 + R_2 + R_p + S}{N_{max}}\right) R_2 + g_{R2} - k_T R_2 - syn * \delta_{max,1} \left(\frac{C_1}{C_1 + C_S^{50}}\right) R_2 -$$
$$syn * \delta_{max,2} \left(\frac{C_2}{C_2 + C_{R,2}^{50}}\right) R_2 + m_2(C_2) S$$

$$7)\quad \frac{dR_p}{dt} = \alpha_{Rp} \left(1 - \frac{R_m + R_1 + R_2 + R_p + S}{N_{max}}\right) R_p + g_{Rp} - k_T R_p - syn * \delta_{max,1} \left(\frac{C_1}{C_1 + C_{R,1}^{50}}\right) R_p -$$
$$syn * \delta_{max,2} \left(\frac{C_2}{C_2 + C_{R,2}^{50}}\right) R_p + \frac{\beta S R_p}{R_m + R_p + S}$$

147   **Eq Set 1.** Sensitive and resistant populations under selective pressure from antimicrobial
148   combination therapy including chromosomal mutation and HGT resistance mechanisms
149

Variable	Definitions
C_1	Antibiotic 1 Concentration (ug/mL)
C_2	Antibiotic 2 Concentration (ug/mL)
S	Susceptible (cells)
R_m	Resistant to both Antibiotic 1 and Antibiotic 2 from Chromosomal Mutation (cells)
R_1	Resistant to only Antibiotic 1 from Chromosomal Mutation (cells)
R_2	Resistant to only Antibiotic 2 from Chromosomal Mutation (cells)
R_p	Resistant to Antibiotic 1 and Antibiotic 2 from MDR Plasmid (cells)
E	Environmental Concentration of Antibiotic((ug/mL)/hr)
syn	Synergy Parameter (non-dimensional)

152 **Table 1.** Model variables and definitions

Parameter	Value	Description	Source
k_e	1.97	Antibiotic Clearance (1/hr)	18
α_S	13.66	Growth Rate of Susceptible Bacteria (1/hr)	18
α_{Rm}	1.9	Growth Rate of Bacteria Resistant from Mutation (1/hr)	18
α_{Rp}	2.1	Growth Rate of Bacteria Resistant from Plasmid (1/hr)	18, 12, 26
N_{max}	10^9	Carrying Capacity (cells/mL)	23
$g_s, g_{Rm},$ g_{R1}, g_{R2}, g_{Rp}	0	Bacterial Influx Rates (cells/hr)	N/A
k_T	0	Bacterial Efflux Rate (1/hr)	N/A
$\delta_{max,1}$, $\delta_{max,2}$	27.14	Bacterial Killing Rate in Response to Antibiotic 1 and Antibiotic 2 (1/hr)	18
$C_S^{50}, C_{R,1}^{50},$ $C_{R,2}^{50}$	12.5, 100, 100	Antibiotic Concentration where the Killing Action is Half its Maximum Value (ug/mL)	Experimentally Determined
$m_1(C_1)$	$2.35*10^{-6}$ (C_1)	Mutation Frequency under Antibiotic 1 (1/hr)	8
$m_2(C_2)$	$2.35*10^{-8}$ (C_2)	Mutation Frequency under Antibiotic 2 (1/hr)	8
β	.001	Gene Transfer Rate (1/hr)	12

Table 2. Model parameter values and descriptions

Experimental validation of the model was done using the eVOLVER system (24). The

experiment was initialized with inoculation of LB media with *E. coli* MG1655 in static

conditions at 37°C. Then, inflow and outflow of Rifampicin-containing LB media at three

concentrations (6, 8 and 12 mg/L) were started at a flow rate of 16 mL/hr. Following the

beginning of inflow/outflow, each culture condition was sampled daily and the concentration of

total bacteria was monitored through OD measurements taken through the eVOLVER system

and resistance bacteria were calculated through plating on selective LB agar containing 200

mg/L Rifampicin. This Rifampicin concentration is eight-fold greater than the experimentally

determined MIC for Rifampicin (25 mg/L).

Results and Discussion

Model Validation

In order to validate the developed model, experiments were conducted with a simplified

experimental system including only one antibiotic (Rifampicin) and no plasmid-containing

bacteria such that there was only chromosomal mutation as a mechanism for resistance. This

experimental validation was done on the eVOLVER, a continuous culture platform which allows

for automated, highly flexible and scalable microbial growth and lab evolution. eVOLVER

allows for continuous flow conditions and independent, precise and multiparameter control of

growth conditions such as temperature and flow rate (24). This allows us to precisely recreate the

model parameters in vitro in a high throughput and repeatable manner. For these eVOLVER

studies, we set certain inflow and outflow rates, initial bacterial populations, and antibiotic

concentrations, as required as inputs for the computational model and then measure susceptible

and resistant bacterial populations over the course of the continuous evolutions experiment. The

outputs of the study can then be directly compared to the outputs of the model. The results of

these studies are shown in Figure 2a-c. The experiments showed that at 6 mg/L and 8 mg/L

Rifampicin (where the MIC of Rifampicin is 25 mg/L), the resistant population developed to a

steady state around the same order of magnitude of the steady state population of susceptible

bacterial. This steady state resistant population was larger for the 8 mg/L Rifampicin condition.

At 12 mg/L, the experiments showed that the resistant bacteria dominated the population and the

susceptible population dropped to zero. After the conclusion of the experiment, model equations

and parameter values were adjusted to match the experimental behavior (Eq set 2) (Table 3).

Notably, the mutation rates were increased by several orders of magnitude to match the rapid *in*

vitro development of resistance. Additionally, the experimental results led to the addition of an

equation for a theorized semi-resistant population with a lower level of resistance (MIC=100

mg/L) (S_1). This was added to match the experimental finding that resistance can coexist at a

steady state with the susceptible population. We theorized that the observed susceptible

population also contained a subpopulation of semi-resistant cells which were not detected by the

8X MIC resistance cutoff rate of the experiment. These semi-resistant cells would allow for a

sustained susceptible population in conditions with constant antibiotic pressure.

$$1) \quad \frac{dC_1}{dt} = E_1 - k_e C_1$$

$$2) \quad \frac{dS}{dt} = \alpha_S \left(1 - \frac{R_1 + S + S_1}{N_{max}}\right) S + g_S - k_T S - \delta_{max,1} \left(\frac{C_1}{C_1 + C_{S,1}^{50}}\right) S - m_1(C_1)S - ms_1(C_1)S$$

$$3) \quad \frac{dR_1}{dt} = \alpha_{R,1} \left(1 - \frac{R_1 + S + S_1}{N_{max}}\right) R_1 + g_R - k_T R_1 - \delta_{max,1} \left(\frac{C_1}{C_1 + C_{R,1,1}^{50}}\right) R_1 + m_1(C_1)S + m_1(C_1)S_1$$

$$4) \quad \frac{dS_1}{dt} = \alpha_{S,1} \left(1 - \frac{R_1 + S + S_1}{N_{max}}\right) R_1 + g_{S_1} - k_T S_1 - \delta_{max,1} \left(\frac{C_1}{C_1 + C_{S_1}^{50}}\right) S_1 + ms_1(C_1)S - m_1(C_1)S_1$$

200 **Eq Set 2.** Augmented model of sensitive, semi-resistant, and resistant populations under
201 selective pressure from Rifampicin therapy including only chromosomal mutation mechanisms
202
203

Parameter	Value	Description
k_e	1.97	Antibiotic Clearance (1/hr)
α_S	13.66	Growth Rate of Susceptible Bacteria (1/hr)
$\alpha_{R,1}$	4.99	Growth Rate of Bacteria Resistant from Mutation (1/hr)
$\alpha_{S,1}$	8.3	Growth Rate of Bacteria Semi-Resistant from Mutation (1/hr)
N_{max}	$3*10^8$	Carrying Capacity (cells/mL)
$g_S, g_{R,1}, g_{S,1}$	0	Bacterial Influx Rates (cells/hr)
k_T	0.8	Bacterial Efflux Rate (1/hr)
$\delta_{max,1}, \delta_{max,2}$	27.14	Bacterial Killing Rate in Response to Antibiotic 1 and Antibiotic 2 (1/hr)
$C_S^{50}, C_{S1}^{50}, C_{R,1}^{50}$	12.5, 100, 200	Antibiotic Concentration where the Killing Action is Half its Maximum Value (ug/mL)
$m_1(C_1)$	$8.5*10^{-3}(C_1)$	Mutation Frequency of Full Resistance Mutation (1/hr)
$ms_1(C_2)$	$8.5*10^{-2}(C_1)$	Mutation Frequency of Low Resistance Mutation (1/hr)

205 **Table 3.** Adjusted model parameter values and descriptions based on eVOLVER experimental
206 conditions and results
207

208 After adjustments, the resulting model outputs matched the general behavior of the resistant and
209 susceptible bacterial populations in response to the selected concentrations of Rifampicin (Figure
210 2d-f). This established the ability of the model to match in vitro behavior allowing for the
211 development of a model with predictive capabilities.
212

Figure 2. Measured susceptible and resistant populations from eVOLVER experiments of E. coli grown in continuous culture in media containing a.) 6 mg/L Rifampicin, b.) 8 mg/L Rifampicin, c.) 12 mg/L Rifampicin. Post-adjustment model outputs for susceptible and resistant populations of E. coli grown in media containing d.) 6 mg/L Rifampicin, e.) 8 mg/L Rifampicin, f.) 12 mg/L Rifampicin.

Following this initial study, the adjusted model was used to make predictions for a second experiment with different Rifampicin concentrations (Figure 3a-b). This second experiment was again conducted using the eVOLVER with the same growth conditions and the results of the experiment qualitatively matched the behavior predicted by the model (Figure 3c-d). This predictive capability demonstrated the validity of the modelling assumptions and prompted further simulation studies using the model.

Figure 3. Post-adjustment model predictions for susceptible and resistant populations of *E. coli* grown in media containing a.) 3 mg/L Rifampicin, b.) 10 mg/L Rifampicin. Measured susceptible and resistant populations from eVOLVER experiments of E. coli grown in continuous culture in media containing c.) 3 mg/L Rifampicin, d.) 10 mg/L Rifampicin.

Model Simulations

   Initial model simulations were done to compare the outputs of our model with known scenarios under low antibiotic pressure. In the scenario of no HGT, as expected, only mutational resistance was observed (Figure 4a). Likewise, for a scenario in which there is no chromosomal mutation (Chromosomal Mutation Rate = Initial Mutant Population = 0), only resistance from HGT was observed and due to an assumed higher growth rate due to a lower fitness cost for bacteria with resistance-conferring plasmids than those with chromosomal resistance mutations (26), this resistance overtook the susceptible population at a higher rate (Figure 4b).

Figure 4. Sensitive and resistant populations under **a.)** resistance only from chromosomal mutations; **b.)** resistance only from the HGT of resistance conferring plasmids (MDR plasmid)

Very low antibiotic residue concentration can promote resistant population growth

We then turned our attention to simulating realistic scenarios in wastewater and sewage. We studied the effects of changing antibiotic residue concentration on the development of antibiotic resistance (Figure 5). Low subinhibitory concentrations (0.5 - 2 ug/ml), as would be present in wastewater settings (4, 5), were studied. These concentrations are an order of magnitude lower than the minimum inhibitory concentrations of each of the two simulated antibiotics, which have been assumed as 25 ug/mL based on our experimental results with rifampicin. At these low concentrations, increased antibiotic residue concentration increased the rate of resistance development and decreased the time to resistant population dominance (defined as time when $R_m+R_p> S$). This is in agreement with several prior studies linking subtherapeutic antibiotic levels with both chromosomal resistance mutation development and HGT (6, 7, 9-12). Future studies will look at higher antibiotic concentrations and the point of inflection where this increase in antibiotic concentration begins to prevent resistance through bactericidal activity. Furthermore, HGT was observed to be the dominant mode of resistance, in agreement with patterns reported in studies of *E. coli* in other settings (20).

Figure 5. Sensitive and resistant populations under antibiotic residue concentrations of a.) 0.5 ug/ml antibiotic 1 + 0.5 ug/ml antibiotic 2; b.) 1 ug/ml antibiotic 1 + 1 ug/ml antibiotic 2; c.) 2 ug/ml antibiotic 1 + 2 ug/ml antibiotic 2

Increase in HGT Rate increases resistance at low concentrations

Additionally, the effect of HGT was modelled by increasing the HGT rate, β (Figure 6). Prior studies have indicated that HGT rate is a less significant driver of resistance frequencies (21). However, our model shows that at very low subinhibitory concentrations of antibiotic, increased HGT rate significantly decreases the time to resistant population domination. This result indicates that increasing HGT rate allows resistance to be acquired at very low selective pressures where there are infrequent chromosomal mutations. This result is significant as it shows that decreasing antibiotic levels in wastewater to low levels is not sufficient in preventing the growth of resistant populations. At very low concentrations of antibiotics, the HGT rate of antibiotic resistance genes in wastewater can significantly affect the proliferation of resistant populations in bacteria with high gene transfer rates. As the HGT rates of the multitude of bacterial species in wastewater is not a commonly monitored parameter, this may be a factor to be mindful of in wastewater surveillance, as these rates can have a significant effect on the evolution of resistant population reservoirs even at low antibiotic concentrations.

281 **Figure 6** Effect of horizontal gene transfer rate on time to resistant population dominance for
282 subinhibitory antibiotic residue concentrations (defined as time when $R_m+R_p> S$)
283
Effect of Bacterial Killing Rate
Because wastewater settings can often have low concentrations of antibiotics from
various polluting factors (4, 5), we probed the specific effects of antibiotic residues on the
development of resistant populations. Individual antibiotics have different killing rates based on
factors such as their modes of action. Thus, we investigated the effect of bacterial killing rate on
resistant population growth (Figure 7). Increasing the killing rate terms ($\delta_{max,1}$ *and* $\delta_{max,2}$) was
observed to decrease the time it takes for the resistant population to dominate under
subinhibitory concentrations. This result is in agreement with previous studies showing increased
selective pressure from antibiotics at subinhibitory concentrations can increase resistance
development (6, 7).
294
295

Figure 7 Effect of bacterial killing rate on time to resistant population dominance for subinhibitory antibiotic residue concentrations (defined as time when $R_m+R_p> S$)

Synergistic antibiotic interactions increase resistant population growth

      The interaction between two antibiotics has previously been shown to affect resistance acquisition (22). Synergy is the interaction of multiple drugs to have a greater killing action than the sum of their parts. Antibiotic synergy has also been shown to increase the likelihood of resistance acquisition at subtherapeutic doses (8). However, the effects of antibiotic interaction on the growth of resistant populations in wastewater settings, where many antibiotic residues can be present, has not previously been observed or modeled. In order to fill this gap, we have probed the effects of synergistic effects between antibiotic residues on the system by varying the synergy parameter, *syn*. Results from our model show that increased synergy between different antibiotics resulted in a decrease in the time to resistant population elimination at suprainhibitory concentrations (Figure 8a). This result may initially seem counterintuitive, but is in fact consistent with previous studies. This is because the synergistic action between the antibiotics increases the killing action to the point where it is effective on the resistant populations at lower concentrations than for antagonistic pairs. Hence, the synergy between the two antibiotics allows for greater bactericidal activity at lower concentrations. However, at subinhibitory concentrations, increasing synergy between antibiotics decreased the time to resistant population dominance (Figure 8b). This is in agreement with other models in which synergy between antibiotics was observed to increase resistance acquisition (8). Our model further shows that these effects of synergy are observable even with low levels of antibiotic, such as the antibiotic residue concentrations present in wastewater. While the effects of antibiotic interaction on resistance acquisition have been previously modeled, low subinhibitory antibiotic concentrations

320 such as those found in wastewater have not been well studied and are critical for understanding
321 resistance development in wastewater settings (17).

322 **Figure 8** Effect of synergy of antibiotic combinations on **a.)** time to resistant population
323 elimination for suprainhibitory antibiotic residue concentrations **b.)** time to resistant population
324 dominance for subinhibitory antibiotic residue concentrations (defined as time when $R_m+R_p > S$)

Conclusion

Though we have been able to draw a number of conclusions on the relative effect of a
variety of factors affecting resistance development in wastewater, we note that our model does
have limitations. First, our model is based on parameter values from literature, not all of which
have been experimentally validated against wastewater conditions. Additionally, other
phenomena related to microbial growth, such as dormancy, biofilm formation and the Long-
Term Survival phase, have not been included in this initial model and the addition of these terms
in future modelling studies may allow for better quantification of resistance development.
However, despite these limitations, experimental validation demonstrated the ability of our
model to qualitatively predict *in vitro* behavior of bacterial resistance development in response to
multiple subinhibitory concentrations of Rifampicin (Figure 3). We have also assumed a
simplified case where the two mechanisms of resistance acquisition that are modelled are
independent, that as interaction between HGT and chromosomal mutation and the behavior of
cells having both are unknown. This assumption can be tested through future experimental work.
Our model also does not account for multiple bacterial species or a multitude of antibiotic
residues, as would be present in a complex environment like wastewater. Interaction between the
multiple bacterial species as well as quorum sensing within bacteria of the same species could
have an effect on resistant population size and is a future area of expansion for this model.
Additional improvements to the accuracy and robustness of the model could be made through
parameter initialization from field data, and experimental validation of model inputs and outputs
under specific wastewater conditions.

Despite these limitations, our model provides new, and important quantitative insight on
the evolution of resistant bacterial reservoirs. It also provides an integrated framework to
incorporate several aspects of resistance acquisition and growth previously lacking from models
focused on AMR in wastewater settings. In terms of results, we have shown that HGT rate can
be a significant driver of resistant population growth at very low antibiotic concentrations
(Figures 5 and 6). This indicates that HGT rates of bacteria in wastewater may be important to
monitor in addition to antibiotic residue concentration. We have also shown that synergy
between the antibiotic residues present in wastewater can increase the rate of resistant population
growth, even at the low concentrations present in wastewater (Figure 8). Thus, antibiotic residues
in wastewater may pose a greater risk than might be expected without taking these antibiotic
interactions into consideration. This has important implications for determining acceptable
antibiotic levels in wastewater post treatment, as determining levels without considering
antibiotic interactions may lead to overestimating the permissible level of antibiotic and allow
for the proliferation of antibiotic resistant bacterial populations. This model can be adapted for
use as a prediction tool for public health policy makers and be used to predict resistant
population emergence in different sewage and wastewater conditions. Additionally, it can be
expanded to be used to model different resistant outbreak prevention strategies in sewage and
wastewater treatment.
364

Author Contributions

366 I. Sutradhar designed the model, conducted experiments, and analyzed the data. C. Ching andD.
367 Desai provided guidance on model design and verification. M. Suprenant provided guidance on
368 data analysis. A. Khalil provided facilities for experimental work with the eVOLVER and E.
369 Briars and Z. Heins provided assistance on experimental design. I. Sutradhar and M. H. Zaman
370 wrote the article.
371

Acknowledgements

373 This research was supported in part by the National Institutes of Health training grant at Boston
374 University, T32 EB006359. A.S.K. also acknowledges funding from the NIH Director's New
375 Innovator Award (1DP2AI131083). The content is solely the responsibility of the authors and
376 does not necessarily represent the official views of the National Institutes of Health.

References

1. Nadimpalli, M. L., S. J. Marks, M. C. Montealegre, R. H. Gilman, M. J. Pajuelo, M. Saito, P. Tsukayama, S. M. Njenga, J. Kiiru, J. Swarthout, M. A. Islam, T. R. Julian, and A. J. Pickering. 2020. Urban informal settlements as hotspots of antimicrobial resistance and the need to curb environmental transmission. Nat. Microbiol. 5:787–795.

2. Bengtsson-Palme, J., and D. Larsson. 2016. Concentrations of antibiotics predicted to select for resistant bacteria: Proposed limits for environmental regulation. Environ Int. 86:140-149.

3. Klemm, E. J., S. Shakoor, A. J. Page, F. N. Qamar, K. Judge, D. K. Saeed, V. K. Wong, T. J. Dallman, S. Nair, S. Baker, G. Shaheen, S. Qureshi, M. T. Yousafzai, M. K. Saleem, Z. Hasan, G. Dougan, and R. Hasan. 2018. Emergence of an extensively drug-resistant Salmonella enterica serovar typhi clone harboring a promiscuous plasmid encoding resistance to fluoroquinolones and third-generation cephalosporins. MBio 9:1–10.

4. Gothwal, R. and T. Shashidhar. 2015. Antibiotic Pollution in the Environment: A Review. CLEAN-Soil, Air, Water. 43(4):479-489.

5. Tran, N. H., M. Reinhard, and K. Y. Gin. 2017. Occurrence and fate of emerging contaminants in municipal wastewater treatment plants from different geographical regions-a review. Water Res. 133:182-207.

6. Andersson, D. I. and D. Hughes. 2014. Microbiological effects of sublethal levels of antibiotics. Nat. Rev. Microbiol. 12: 465–478.

7. Ching, C. and M. H. Zaman. 2020. Development and selection of low- level multi-drug resistance over an extended range of sub-inhibitory ciprofloxacin concentrations in Escherichia coli. Sci. Rep. 1–9

8. Michel, J. B., P. J. Yeh, R. Chait, R. C. Moellering, and R. Kishony. 2008. Drug interactions modulate the potential for evolution of resistance. Proc. Natl. Acad. Sci. U. S. A. 105:14918–14923.

9. Hastings, P. J., S. M. Rosenberg, and A. Slack. 2004. Antibiotic-induced lateral transfer of antibiotic resistance. Trends Microbiol. 12:399–401.

10. Zhang, Y., A. Z. Gu, M. He, D. Li, and J. Chen. 2017. Subinhibitory Concentrations of Disinfectants Promote the Horizontal Transfer of Multidrug Resistance Genes within and across Genera. Environ. Sci. Technol. 51:570–580.

11. Lopatkin, A. J., S. Huang, R. P. Smith, J. K. Srimani, T. A. Sysoeva, S. Bewick, D. K Karig, and L. You. 2016. Antibiotics as a selective driver for conjugation dynamics. Nat. Microbiol. 1:1–8.

12. Baker, M., J. L. Hobman, C. E. R. Dodd, S. J. Ramsden, and D. J Stekel. 2016. Mathematical modelling of antimicrobial resistance in agricultural waste highlights importance of gene transfer rate. FEMS Microbiol. Ecol. 92:1–10.

13. Brinch, C., P. Leekitcharoenphon, A. S. R. Duarte, C. A. Sevendsen, J. D. Jensen, and F. M Aarestrup. 2020. Long-Term Temporal Stability of the Resistome in Sewage from Copenhagen. mSystems. 5:1-20

14. Birkegård, A. C., T. Halasa, N. Toft, A. Folkesson, and K. Græsbøll, K. 2018. Send more data: a systematic review of mathematical models of antimicrobial resistance. Antimicrob. Resist. Infect. Control 7:1–12.

15. Acemel, R. D., F. Govantes, and A. Cuetos. 2018. Computer simulation study of early bacterial biofilm development. Sci. Rep. 8:1–9.

16. Niewiadomska, A. M., B. Jayabalasingham, J. C. Seidman, L. Willem, B. Grenfell, D. Spiro, and C. Viboud. 2019. Population-level mathematical modeling of antimicrobial resistance: A systematic review. BMC Med. 17:1–20.

17. Leclerc, Q. J., J. A. Lindsay, and G. M. Knight. 2019. Mathematical modelling to study the horizontal transfer of antimicrobial resistance genes in bacteria: Current state of the field and recommendations. J. R. Soc. Interface 16.

18. Blanquart, F. 2019. Evolutionary epidemiology models to predict the dynamics of antibiotic resistance. Evol Appl. 12:365-383.

19. Nguyen, T. T., J. Guedj, E. Chachaty, J. de Gunzburg, A. Andremont, and F Mentré. 2014. Mathematical Modeling of Bacterial Kinetics to Predict the Impact of Antibiotic Colonic Exposure and Treatment Duration on the Amount of Resistant Enterobacteria Excreted. PloS Comput Biol. 10.

20. Frazão, N., A. Sousa, M. Lässig, and I. Gordo. 2019. Horizontal gene transfer overrides mutation in Escherichia coli colonizing the mammalian gut. Proc. Natl. Acad. Sci. U. S. A. 116:17906–17915.

21. Lehtinen, S., C. Chewapreecha, J. Lees, W. Hanage, M. Lipsitch, N. Croucher, S. Bentley, P. Turner, C. Fraser, and R. Mostowy. 2020. Horizontal gene transfer rate is not the primary determinant of observed antibiotic resistance frequencies in Streptococcus pneumoniae. Sci Adv. 6.

22. Weinstein, Z. B., N. Kuru, S. Kiriakov, A. C. Palmer, A. S. Khalil, P. A. Clemons, M. H. Zaman, F. P. Roth, and M. Cokol. 2018. Modeling the impact of drug interactions on therapeutic selectivity. Nat Commun. 9:1-9.

23. Kram, K. and S. Finkle. 2014. Culture volume and vessel affect long-term survival, mutation frequency, and oxidative stress of Escherichia coli. Applied and Environmental Microbiology. 80:1732-1738.

24. Wong, B. G., Mancuso, C. P., Kiriakov, S., Bashor, C. J. and A. S. Khalil. 2018. Precise, automated control of conditions for high-throughput growth of yeast and bacteria with eVOLVER. Nat. Biotechnol. 36, 614–623.

25. O'Neill J. 2014. Review on Antimicrobial Resistance Antimicrobial Resistance: Tackling a crisis for the health and wealth of nations. London: Review on Antimicrobial Resistance;

26. Vogwill, T., and R. C. MacLean. (2015). The genetic basis of the fitness costs of antimicrobial resistance: a meta-analysis approach. Evolutionary applications, 8(3), 284–295. https://doi.org/10.1111/eva.12202