The transcription factor CLAMP is required for neurogenesis in *Drosophila melanogaster*.

Maria A. Tsiarli [1, 2, 6 *,] James A. Kentro[1,5, 6], Ashley M. Conard [1, 3, 4], Lucy Xu [1], Erica Nguyen[1], Kate

O'Connor-Giles[5,6], Erica N. Larschan [1,6 *]

[1] Department of Molecular Biology, Celluiar Biology and Biochemistry, Brown University, Providence,

RI 02912, USA

[2] Department of Neurobiology, Harvard Medical School, Boston, MA Department 02115, USA

[3] Center for Computational and Molecular Biology, Brown University, Providence, RI, 02912;

[4] Computer Science Department, Brown University, Providence, RI, 02912

[5] Department of Neuroscience, Brown University, Providence, RI, USA

[6] Carney Institute for Brain Science, Brown University, Providence, RI, USA

* corresponding author

Contact information

Maria A. Tsiarli: maria_tsiarli@brown.edu; maria_tsiarli@hms.harvard.edu

Ashley M. Conard: ashley_conard@brown.edu

Lucy Xu: lucy_xu@alumni.brown.edu

Erica Nguyen: erica_nguyen@alumni.brown.edu

Erica N. Larschan: erica_larschan@brown.edu

23
24
25

Abstract

Neural stem cell (NSC) differentiation is controlled by cell-intrinsic and external signals from the stem cell

niche including niche surface glia (SG). However, the mechanisms by which transcription factors drive

NSC differentiation within the niche remain largely unknown. Here, we show that the *Drosophila*

melanogaster transcription factor, Chromatin-linked adaptor for MSL proteins (CLAMP) is required for

regulation of stemness and proliferation of NSCs, especially of the optic lobe (OL). CLAMP promotes

transcription of genes involved in stemness, proliferation, and glial development and represses transcription

of genes involved in neurogenesis and niche survival. Consistent with transcriptional changes, CLAMP

promotes NSC proliferation and niche SG production, while lack of CLAMP severely and specifically

impacts OL development. To identify potential mechanisms by which CLAMP may regulate brain

development, we examined CLAMP motifs and available CLAMP ChIP-seq data to determine which genes

may be direct versus indirect targets. CLAMP motifs are present at many target genes including the glial-

determining gene, *glial cells missing,* while *Tailless,* the master regulator of OL- development is directly

bound by CLAMP. In accordance to these results, in larval OL NSCs lacking CLAMP, Tailless levels are

decreased dramatically, suggesting that CLAMP controls OL neurogenesis *via* Tailless. Overall, our results

suggest that CLAMP regulates a transcriptional program which drives NSC proliferation and differentiation

via cell-intrinsic and niche-dependent mechanisms that involve transcriptional regulation of *Tailless* and

niche glia.

Introduction

Brain development is an extremely complex yet carefully orchestrated process, regulated along spatiotemporal axes. For the formation of the functional brain circuitry to take place, neural stem cells (NSCs) must undergo multiple rounds of divisions and lineage decisions at each step of the process are governed by specific transcription factors (TFs) (KONSTANTINIDES *et al.* 2018). In addition to the intrinsic developmental program, external metabolic and paracrine signals constitute another layer of developmental control over NSCs. Specifically, NSCs are part of a specialized microenvironment, the stem cell niche (SCADDEN 2006), which plays an architectural and nurturing role for the NSCs and their progeny. Moreover, the niche serves as a hub for the relay of metabolic and developmental signals that regulate differentiation of NSCs in order for the proper sequence of developmental events to be maintained (SPEDER AND BRAND 2018). In mammals, the stem cell niche is composed of various cell populations including neurons, glial cells, resident immune cells, the microglia, blood vessels which along with endothelial cells and glial cells form the blood-brain-barrier (DANI *et al.* 2019). In *Drosophila*, the stem cell niche consists of NSCs, NSC progeny and glial cells of various types that are vital to the maintenance of the homeostatic balance required for proper neurogenesis to take place (HINDLE AND BAINTON 2014; SPÉDER AND BRAND 2014; BAILEY *et al.* 2015; KANAI *et al.* 2018; SPEDER AND BRAND 2018; RAMON-CAÑELLAS *et al.* 2019; YUAN *et al.* 2020). However, little is understood about the gene regulatory mechanisms that generate the stem cell niche.

In *Drosophila melanogaster* neurogenesis occurs at two distinct stages: the embryonic and post-embryonic stages (HARTENSTEIN AND CAMPOS-ORTEGA 1984; PROKOP AND TECHNAU 1991). The *Drosophila* central nervous system (CNS) is composed of the central brain (CB), the optic lobe (OL) and the ventral nerve cord (VNC). During the embryonic stage, CB neural stem cells (neuroblasts in *Drosophila*) proliferate to generate primary neurons until they become quiescent at the end of the embryonic stage. The optic lobe

primordium which would form the OL, is generated by epithelial cells of the posterior procephalic region

between embryonic stages 11-13 (RAMON-CAÑELLAS *et al.* 2018). Between stages 12 and 13, the OL

primordium invaginates and attaches at the ventrolateral surface of the brain and embryonic OL neuroblasts

(EONs) are formed (HAKES *et al.* 2018). EONs proliferate briefly to generate glial and neuronal progeny

and also enter quiescence (HAKES *et al.* 2018; CONTRERAS *et al.* 2019b). During the early larval stage,

concurrent with the initiation of larval feeding, metabolic signals impinge on the NSCs and cause exit from

quiescence and initiation of neurogenesis.

Neural stem cells divide asymmetrically, such that in each division they generate a new NSC and a daughter

cell which will generate glial and/or neuronal progeny (EGGER *et al.* 2008). For the appropriate asymmetric

division to take place, polarity determinants have to be sequestered on the apical and basal sides of the

mother NSC: the apical complexes are inherited by the new NSC and maintain the proliferating state,

whereas basal proteins are inherited by the daughter cell and will direct it to generate differentiated lineages.

The Par protein complex which is apically localized also directs the localization of the basal determinants.

Moreover, Par complex proteins such as the *Inscuteable* (Insc) and *Partner of Inscuteable (Pins)* regulate

the orientation of the mitotic spindle which determines how the cleavage furrow will form during the

division of the NSC (GALLAUD *et al.* 2017; CREWS 2019). Additionally, in the optic lobe, sequential

expression of transcription factors in proliferating NSCs, including of *Tailless* and *Notch*, regulates the

temporal generation of diverse progeny types (LI *et al.* 2013; BERTET *et al.* 2014).

In the *Drosophila,* five distinct CNS glial types are identified based on morphological, positioning and

molecular profile criteria: perineurial (PG), subperineurial glia (SPG), cortex glia (CG), astrocyte-like and

ensheathing glia (YILDIRIM *et al.* 2019). Perineural glia ~~are found on the outside layer of the CNS and~~ serve

a nurturing role for the development and function of the NSCs and also regulate NSC proliferation (KANAI

et al. 2018). Reciprocally, the proliferating NSCs regulate their proliferation and the survival of PG *via* a

mechanism that implicates secretion the Glass bottom boat protein. With the initiation of larval feeding,

SPG and PG contribute to the reactivation of quiescent NSCs (SOUSA-NUNES *et al.* 2011; SPÉDER AND

BRAND 2014; KANAI *et al.* 2018) *via* fat body signaling and induction of insulin secretion from SPG. In

conditions of oxidative stress and hypoxia, SPG and CG form lipid droplets (LDs) which protect NSCs

from oxidative stress (BAILEY *et al.* 2015; KIS *et al.* 2015). CG forms a bespoke membrane chamber around

dividing NSCs and their progeny, thus providing protection to the proliferating NSCs and their niche, and

serving as individual spatial units for each expanding NSC and its progeny (SPEDER AND BRAND 2018). A

special subtype of CG, *mir8*$^+$ surface-associated CG, controls neurogenesis in the OL (MORANTE *et al.*

2013; RAMON-CAÑELLAS *et al.* 2018; YILDIRIM *et al.* 2019). Thus, niche glia are important for the function

of NSCs. However, the molecular mechanisms that direct the functional and nurturing relationship between

niche glia and NSCs remain poorly understood.

Here we define a role for the essential CLAMP zinc finger TF (SORUCO *et al.* 2013b) in neurogenesis which

is partly mediated by transcriptional regulation of key TFs controlling stemness and proliferation, and partly

via niche glia. CLAMP was discovered as part of a screen for factors that are necessary for the function of

the dosage compensation machinery in *Drosophila melanogaster* males (LARSCHAN *et al.* 2012). CLAMP

is essential for viability of both males and females and has transcriptional targets on all chromosomes, with

a stronger binding affinity on the X chromosome (SORUCO *et al.* 2013b; URBAN 2017). Since its first

characterization as a required link for the MSL complex to bind to the X-chromosome (SORUCO *et al.*

2013a; TIKHONOVA *et al.* 2019), CLAMP is emerging as a multifaceted transcription factor with multiple

roles including chromatin organization (KAYE *et al.* 2017), histone locus function (RIEDER *et al.* 2017) and

regulation of Pol II (URBAN *et al.* 2017). The identification of *Clamp* (CG1832), as a functional interactor

of *embryonic lethal abnormal visual system* (*elav*) and *erect wings* (*ewg),* critical regulators of embryonic

and post-embryonic neural development, suggested a potential neuronal role of CLAMP (HAUSSMANN *et*

al. 2008; HAUSSMANN AND SOLLER 2010b). Therefore, we hypothesized that CLAMP is a TF which

regulates neuronal development and NSC biology.

To test our hypothesis, we investigated the role of CLAMP in nervous system development. We found that

CLAMP is important both for neuronal and glial cell generation and regulates NSC proliferation *via* cell-

intrinsic and niche-dependent mechanisms. We used a CRISPR/Cas9-generated frameshift mutant in the

clamp gene which eliminates the expression of the CLAMP protein (URBAN 2017). Using this approach,

we observed that lack of CLAMP results in brain development defects. Specifically, *clamp* null mutants

have a smaller brain than controls, with the most prominent defect observed in the optic lobe which is

almost entirely lost. To determine the transcriptional networks affected by the absence of CLAMP, we

performed mRNA-sequencing data of *clamp* mutant embryos and larval brains, followed by gene ontology

(GO) analysis, which revealled that CLAMP positively regulates stemness/NSC proliferation genes and

glial-development genes.

To specifically determine which of the differentially expressed genes (DEGs) are under direct CLAMP

regulation, we performed binding motif analysis which revealed that CLAMP motifs are present at key

neurogenesis genes such as *bazooka, gbb* and the glial-development regulator, *glial cells missing.* To further

validate direct regulation of the identified DEGs by CLAMP, we proceeded with mining available CLAMP

embryo ChIP-seq data. Importantly, this analysis revealed that *Tailless,* the master regulator of OL

development, is both regulated and bound by CLAMP, suggesting that it is a direct target and can underlie

the profound defects in OL development seen in *clamp* null mutants.

To further explore the relation between CLAMP and Tailless, we proceeded with immunostaining in *clamp*

null mutant brains using NSC and glia markers and Tailless. This revealed that *clamp* null mutants exhibit

a brain-wide reduction in proliferating NSCs and surface glia and a profound defect in the development of

the OL, similar to the phenotype seen in *Tailless* knock-down brains (GUILLERMIN *et al.* 2015).

Furthermore, knock-down of *clamp* in glia but not in neurons, revealed that NSC proliferation and glia

generation are partly regulated *via* glial CLAMP.

Our results support a model in which CLAMP regulates a transcriptional program which drives NSC

proliferation and differentiation *via* cell-intrinsic and niche-dependent mechanisms that involve direct

transcriptional regulation of *Tailless* and niche glia.

Materials and Methods

Fly Strains

Fly stocks used in this study were as follows: y^1w^1 wild type controls, $clamp^2$ null, and *CLAMP* rescue lines, which were generated by Urban et al.(2017) and described therein *(URBAN 2017)*. Briefly, the *clamp* null, is a CRISPR-Cas9 generated frameshift mutation which abolishes CLAMP protein expression, whereas the *clamp* rescue was generated by introduction of a full CLAMP rescue construct on chromosome 3L. For the RNAi experiments the following lines were used: *repo-GAL4*, *elav^c155* (*elav-GAL4*) and UAS-*clamp-RNAi,* all obtained from the Bloomington Stock Center. All crosses were done at 25ºC and a 12-hour light/dark cycle.

Immunostaining

The following antibodies were used: Rabbit anti-CLAMP (1:200, Abcam custom made), mouse anti-Repo (1:50, Developmental Studies Hybridoma Bank (DSHB) 8D12), rabbit anti-DCP1 (1:100, Cell Signaling), rat anti-Deadpan (Dpn, 1:100, Abcam), rat anti-Elav (1:50, DSHB), mouse anti-Prospero (Pros, 1:30, DSHB), PCNA (PC10, Abcam), Nc82 (1:10, DSHB), rabbit anti-Tailless (Kindly provided by Dr James Sharpe, EMBL-Barcelona). Secondary antibodies used were Alexa Fluor Conjugated A405, A488, A568, A647 (1:1000, Invitrogen). A standard staining protocol was used.

Quantification of marker-positive and superficial glia cells

To quantify superficial glia, 3-μm projections (z-projections of three 1-μm optical sections) of the dorsal surface of third instar larval brain hemispheres stained for Repo were used and quantified manually using Adobe Photoshop, as exemplified by Kanai et al. (2018). Six hemispheres were quantified for each

genotype. The same approach was used to quantify marker positive cells i.e., Dpn, Pros and EdU. To ensure

consistency between samples of control and the *clamp* mutant brains basic morphological features based

on DAPI staining in combination with stack depth were monitored.

Image acquisition and processing

Fluorescent images were acquired using an inverted Olympus FLUOVIEW FV1000 confocal laser

scanning microscope or a Leica TCS SP2 AOBS DMIRE2 inverted scanning confocal microscope. Adobe

Photoshop was used to adjust brightness and contrast in images and ImageJ (or Fiji (Schindelin et al., 2012))

was used to analyzed images, while Adobe Illustrator was used to compile figures.

EdU Incorporation

EdU incorporation in proliferative brain cells, was performed using Click-It Edu Alexa Fluor A488

(Thermo Fisher Scientific, MA, USA) in an adaptation of Tastan and Liu (2015) (TASTAN AND LIU 2015)

method. Briefly, samples were dissected in ice-cold Graces media and then transferred in Graces media

containing 5μg/ml of EdU, for 2 hours at room temperature. Then, samples were washed in 0.1% PBST

and fixed in 4% PFA in 0.1% PBST, for 20 min at room temperature. Next, tissues were washed in 5%

HINGS in 0.1 % PBST and then permealized in 0.5% PBST for 20 min (RT). After this step, the EdU Click

It steps were followed according to the manufacturer's instructions and then additional antibody staining

was followed as described above.

Larval Staging

Third Instar larvae crawling on the vial walls were selected to be mid/late stage using the bromophenol blue

method adjusted from Maroni and Stamey (1983) (MARONI AND STAMEY 1983), using 0.05% w/v

bromophenol blue-containing fly food. To control for variations in brain size due to crowding, experiments

were done under controlled density conditions. Specifically, a set number of mated males and females were

allowed to mate for 5 hours and then removed from the vial. Additionally, the progeny number was

evaluated to validate that we did not have large deviations between control and *clamp* mutant larvae

numbers.

RNA extraction and RNA sequencing

Mid/late third instar larvae were staged using the bromophenol blue method and larval brains were dissected

in ice-cold PBS, collected in TRIzol until RNA extraction was performed using the phenol-chloroform

method. Each biological replicate was generated by pooling at least 12 larval brains. mRNA-focused

sequencing libraries were constructed from total RNA using the Illumina Truseq RNA Library kit v2

according to the manufacturer's instructions (no strand-information). Paired-end 100 bp or 50 bp was

performed on an Illumina HiSeq2500 sequencer in High-Output mode at the MGH NextGen Sequencing

Core. Four biological samples were used per experimental condition. As the control group, we used the

CLAMP rescue line in which the wild type CLAMP gene was inserted into an attP site under the control of

its endogenous promoter (URBAN 2017).

RNA sequencing data analysis and Differential Gene Expression Analysis

RNA-seq reads were processed using FastQC (ANDREWS 2010) with default parameters to check the

quality of raw sequence data and filter out sequences flagged for poor quality, and aligned using HISAT2

(KIM *et al.* 2015). Then read counts per gene were generated via HTSeq (ANDERS *et al.* 2015) using the

Drosophila melanogaster genome dm6 (BDGP6) genome as reference. Differential gene expression

analysis was performed using DESeq2 (LOVE *et al.* 2014). *gProfiler* was used for GO analysis (RAUDVERE

et al. 2019). Using TIMEOR (CONARD *et al.* 2020) the temporally differentially expressed genes where

displayed in   interactive expression cluster-maps along the developmental time scale (see Supplementary

files: *Embryo_heatmap* and *L3_brain_heatmap*), from which additional, GO analysis was performed by

TIMEOR *via* clusterProfiler (YU *et al.* 2012) to evaluate for possible gene networks.

Clamp Binding motifs

We used FIMO (GRANT *et al.* 2011) to identify individual CLAMP motif occurrences within ± 1KB from

the transcription start and end sites for all genes within each gene trajectory cluster. The position weight

matrix used (KAYE *et al.* 2018) highlights the CLAMP motif independently of GAF binding potential. The

specific command used is: *(fimo --oc $OUTPUT_DIR/ CLAMP_PWM.txt $geneList)*

Average profiles

The average control-normalized read depth across a set of regions was plotted using deepTools

computeMatrix and plotProfile. The read depth source are ChIP-seq data in the ENCODE database for

CG1832   (Clamp)   from   mixed   sex   embryos   (0-24hrs),   part   of   the   modERN   project

(https://www.encodeproject.org/experiments/ENCSR858HYS/). The regions were chosen as each set of

differentially expressed genes (DEGs) from the *clamp* null versus *clamp* rescue RNA-seq experiments in

embryo and third instar larvae. These were kept as all DEGs in an experiment and as separate images for

up or down regulated genes in the absence of *clamp*. The gene regions were scaled to match positions of

the transcription start sites (TSS) and transcription end sites (TES) with 3K base pairs shown on either side

of the gene.

Western Blotting

Larval brains were dissected from wandering third instar larvae in ice-cold 0.7% saline and transferred in

PBS + 0.3% Triton X-100 on ice. Liquid was carefully removed, then samples were frozen in liquid

nitrogen and stored at -80°C until use. Approximately ten brains were collected per biological replicate.

For protein extraction, 50 µL of ice-cold RIPA buffer (150 µL1M Tris-HCl pH 8, 12 µL 0.5M EDTA, 0.003

g sodium deoxycholate, 15 µL 20% SDS, 90 µL 5M NaCl, 30 µL 50x protease inhibitor, and 2700 µL

water) was added to each sample. Samples were homogenized using an electric handheld homogenizer and

disposable plastic pestle. After homogenization, the pestle was rinsed with an additional 50 µL of RIPA for

a total volume of 100 µL per tube. Samples were rotated at 4°C for 1-2 hours. Protein concentration was

quantified using the Qubit Protein Assay kit (Thermo Fisher Scientific).

Statistical Analysis

Excel and GraphPad Prism were used for statistical analysis. Bar graphs were generated using the mean

and standard error of the mean (SEM) for each sample. Statistical significance was determined using an

unpaired Student's t-test.

Data Availability Statement & Correspondence

Data and materials generated in this study are available upon request from the corresponding authors.

Transcriptomic data have been deposited in Gene Expression Omnibus (GEO) Database, under the

accession number GSE148704.

Results

CLAMP is expressed throughout the larval nervous system and in proliferating neuroblasts.

CLAMP is maternally deposited at both the protein and RNA levels and therefore *clamp* transcript and

CLAMP protein levels are present at high levels during embryogenesis (RIEDER *et al.* 2017). Near the end

of embryogenesis (16-18 hours), during tissue differentiation and the cessation of the first wave of

neurogenesis, *clamp* transcript levels show a reduction followed by a sharp increase between 20-24 hours

and then a decrease throughout the post-embryonic neurogenesis period (Fig. 1a). Tissue-specific

expression data from the modENCODE Development RNA-Seq Collection (GRAVELEY *et al.* 2011;

BROWN *et al.* 2014)/Flybase (THURMOND *et al.* 2019) show that *clamp* is highly expressed in the third (L3)

central nervous system (Fig.1b).

To define the role of CLAMP in neural development, we mapped its expression in the developing L3 central

nervous system using immunofluorescence. We found that CLAMP is ubiquitously expressed in the nucleus

throughout the larval nervous system, both in cells of the brain and the VNC (Figs.1c and d). Interestingly,

a subset of neuronal processes in the VNC also show protein localization of CLAMP, suggesting that it

may be transported to neuronal arborizations (Fig. 1c, axonal processes indicated by arrows). Co-staining

of proliferating cell nuclear antigen (PCNA) which labels proliferating cells, revealed that CLAMP is

expressed in proliferating NSCs (Figs.1e-h), both in the central brain and the OL (Figs.1f-h, cyan and yellow

arrows, respectively). CLAMP colocalizes with PCNA both in CB neuroblasts and ganglion mother cells,

as identified based on their size difference (Figs.1f-h, cyan and magenta arrows, respectively).

Loss of CLAMP disrupts larval development

To determine the role of CLAMP in nervous system development, we characterized brain development in

clamp² null mutants that we previously generated in our lab using CRISPR (URBAN 2017). In these mutants,

a frameshift mutation in the *clamp* gene abolishes expression of CLAMP protein (URBAN 2017) (from here

on referred to as *clamp* null). Due to CLAMP's early role in the recruitment of the dosage compensation

complex in males (SORUCO *et al.* 2013b), the majority of male larvae die at the L2 stage. In contrast, female

larvae die at the late L3/prepupa stage (URBAN 2017). Because of our interest in the essential role of

CLAMP independent of dosage compensation and due to the small number of male *clamp* null larvae, we

focused on female L3 larvae. Comparison of *clamp* null L3 larvae to stage-matched controls (see Materials

and Methods) revealed that *clamp* larvae are significantly smaller than controls (Figs.2a and b).

Furthermore, macroscopic examination of the central nervous system (CNS) of *clamp* null larvae revealed

that their brains are also significantly smaller than controls (Figs.2c and d). Western blotting for CLAMP

in *clamp* null larvae brains verified the absence of CLAMP protein in L3 brains (Fig.2e).

Loss of CLAMP disrupts brain development

To evaluate CNS development in *clamp null* mutants, we focused on the mid/late L3 stage (~120 hours

after larval hatching (ALH)), because the *clamp* mutants die during late L3/early pupa stages. To gain

temporal insight, we also examined early L3 larvae brains. The overall morphology of the CNS, as observed

with DAPI staining, is different between nulls and controls (Figs.3a and e). As neural development

progresses through the larval stages, the neuroepithelial (NE) cells of the optic lobe (OL) start to

differentiate into neuroblasts (NBs), which will subsequently divide to give rise to neurons and glia

(CONTRERAS *et al.* 2019a). We observed that the *clamp* null OL is profoundly reduced in size. Specifically,

the clamp *null* OL neuroepithelia at 120 hours ALH resemble the control neuroepithelia at 72 hours ALH

(i.e., early L3) (Figs. 3a, b, e and f). Similarly, the *clamp* mutant central brain (CB) of the mid/late L3, has

an underdeveloped morphology and resembles the early control CB (Figs.3f and a).

Next, we examined NSCs in *clamp* null and control brains using an antibody against Deadpan (Dpn), which labels all NSCs. Full z-stacks of the Dpn-positive NSCs in early and mid/late L3 brains (Figs. 3c, d, g and h), revealed that in the *clamp* null, the development of NBs, both in the CB and OL, is considerably delayed from the early L3 stage and remains so until the mid/late L3 stage when the animals die. The OL NB population is greatly affected as the area containing Dpn-positive NSCs is much smaller in *clamp* nulls than controls in mid/late L3 brains (Figs 3d and h, dashed outlines). These observations suggest that the deficits observed in L3 *clamp* null brains may be due to an earlier arrest of NEs, resulting in the generation of fewer NSCs. Comparison of early L3 *clamp* heterozygotes (*clamp*$^{+/-}$) and *clamp* null brains indicates that a single copy of the *clamp* gene is sufficient to maintain the overall morphology of the developing OL (Supplementary Fig. 1).

Developmental transcriptomic analysis of clamp null mutants reveals a role for CLAMP in nervous system development

The neurodevelopmental deficits observed in the larval *clamp* null mutant brain suggest that the function of CLAMP initiates earlier than the L3 stage. CLAMP is a ubiquitously-expressed TF, and is present throughout the nervous system from early embryonic stages due to the deposition of maternal transcripts into the egg (RIEDER *et al.* 2019) (Fig.1a). Each neurodevelopmental stage is characterized by unique transcriptomic events controlled by specific TF factor networks (KONSTANTINIDES *et al.* 2018). Thus, to understand how CLAMP regulates nervous system development on the transcriptomic level, we conducted RNA-sequencing at the following key developmental stages in *clamp* null mutants: A) late stage embryos (16-18 hours), a time point where neurogenesis pauses (HAKES *et al.* 2018) and most NBs enter quiescence (Fig.4a); and B) mid/late L3 stage brains (Fig.4b), when neuronal and glial cell production are at their maximum (AVET-ROCHEX *et al.* 2012). The selection of the specific embryonic time point was based on studies by Haussmann et al. (2008, 2010), where *clamp* was identified as a functional interactor of *embryonic lethal abnormal vision* (*elav*)and *erect wing* (*ewg*), in 16-18 hour embryos (HAUSSMANN *et al.*

2008; HAUSSMANN AND SOLLER 2010a). Both *elav* and *ewg* are critical to nervous system development

(ROBINOW AND WHITE 1988; DESIMONE AND WHITE 1993), further underscoring a potential role of

CLAMP in the process. To control for non-specific transcriptional changes due to other mutations that

could be in the genetic background, we used the *CLAMP* rescue line as a control (URBAN 2017). In the

CLAMP rescue line, *clamp* is inserted into an attP site under the control of the endogenous promoter.

Analysis of our transcriptomic data revealed that many genes were significantly differentially expressed

(DEGs) both in the embryo and the L3 brain (padj ≤ 0.05, Figs 4a-c). Interestingly, a large proportion of

DEGs were shared between embryo and L3 brain data sets (Fig.4c). The embryo dataset contains 5917

DEGs (padj ≤ 0.05 and Supplementary Table 1a), while the L3 contains 4465 DEGs (padj ≤ 0.05 and

Supplementary Table 2a). Of these DEGs, 2363 are shared between the two stages representing 40% of the

embryonic DEGs and 53% of the L3 DEGs. The size of this shared DEGs dataset supports the hypothesis

that *clamp* regulates gene networks controlling neurodevelopment form embryonic to L3 stages.

Analysis of the transcriptomic data revealed that a great number of genes was significantly and differentially

expressed (DEGs) both in the embryo and the L3 brain (padj ≤ 0.05, Figs 4a-c). Interestingly, a large

proportion of the DEGs was shared between the embryo and the *clamp* null L3 brain (Fig.4c). Accordingly,

the embryo dataset contains 5917 DEGs (padj ≤ 0.05 and Supplementary Table 1a), whereas the L3, 4465

DEGs (padj ≤ 0.05 and Supplementary Table 2a). Of these DEGs, 2363 are shared between the two stages

representing 40% of the embryonic DEGs and 53% of the L3 genes. The size of this shared DEGs dataset,

support the hypothesis that *clamp* deletion results in major changes in gene networks controlling

neurodevelopment, beginning form the embryonic stage and propagated through the L3 stage.

GO analysis revealed statistically significant enrichment for developmental/stem cell-related GO

categories, including neural development, at both the embryonic and larval developmental stages (Figs.4d,

e). Due to the neurodevelopmental defects in the *clamp* null mutant, we then specifically mined our

expression data for differential expression of genes that are implicated in cell cycle attributes, nervous system development and neural stem cell biology. Enrichment was also observed for GO categories involving metabolic pathways. This is interesting as post-embryonic neurogenesis is largely orchestrated by signals originating from nutrient availability and insulin signaling (BRITTON AND EDGAR 1998; BAILEY *et al.* 2015; GALAGOVSKY *et al.* 2018).

Functional analysis of the embryonic DEG dataset revealed highly statistically significant enrichment in categories such as cell cycle regulation, cell cycle checkpoints, mitotic cell cycle phase transition, neurogenesis and nervous system development, metabolic processes, synaptic development and signaling (Supplementary Table 1b). We focused our analysis on two major highly enriched GO categories: i). *Cell Cycle Regulation* [GO:0051726, padj: 6.22E-11] and ii). *Nervous System Development* [GO:0007399, padj: 4.32261E-05].

For the <u>*GO:0051726 -Cell Cycle Regulation*</u> category (Supplementary Table 1c), functional analysis of identified DEGs suggests specific roles, including in G2/M transition of the mitotic cell cycle, regulation of chromosome segregation, cell cycle G1/S phase transition, negative regulation of nervous system development and of insulin signaling (Fig.4d).

Expression of *elav,* an RNA-binding protein and master regulator of neural development, was increased in the absence of CLAMP [L2FC: 0.44; padj: 0.045]. ELAV regulates the expression of *ewg* and both are critical for nervous system development, and neuronal maturation (ZAHARIEVA *et al.* 2015). Haussmann et al. (2008, 2010) (HAUSSMANN *et al.* 2008; HAUSSMANN AND SOLLER 2010b) reported that *clamp* along with a group of nine functionally-related genes which includes *groucho,* another critical developmental regulator (TURKI-JUDEH AND COUREY 2012), are positively regulated by EWG and restrict synaptic growth (Table 1). Notably, 9 out of 10 DEGs regulated by EWG are also DEGs in the *clamp* null embryos.

Therefore, it is possible that a synergistic relationship exists between the CLAMP and ELAV/EWG

pathways, in which they mutually regulate each other's expression to ensure proper brain development.

Meta-analysis of the _GO:0007399: Nervous System development_ category (Supplementary Table 1d),

revealed strong enrichment in the sub-categories of neurogenesis and central nervous system, visual system

development, and glial cell development and differentiation (Fig.4e). Several DEGs critical for neural

development and NSC biology were identified: _Notch_ [L2FC: -0.5, padj: 0.01], _lola_ [L2FC: -0.6, padj:

0.004], and components of the Brahma chromatin remodeling complex such as _moira_ [L2FC: -1.019, padj:

6.21E-05] (NEUMÜLLER _et al._ 2011; JANSSENS _et al._ 2014; SOUTHALL _et al._ 2014). Importantly, many

DEGs are part of specific TF networks that regulate different aspects of neural stem cell renewal identified

by Neumuller et al. (2011): _Su(z)12, Su(var)2-10_, the Hdac1 _Rpd3_ [L2FC: -0.93, padj: 0.0004], and _lola_.

Notch signaling is critical for the balance between proliferation and differentiation in NSCs and maintains

the balance between the symmetrically and asymmetrically dividing stem cell pools in the optic lobe

(EGGER _et al._ 2010), neuroblast selection and delamination (AREFIN _et al._ 2019) and is a key regulator of

asymmetric divisions throughout the nervous system (CONTRERAS _et al._ 2018; HAKES AND BRAND 2019).

Furthermore, additional DEGs in the _Nervous System development_ GO category are: 1) _polo, aPKC, par6,_

pins, aurA and _baz,_ all components of the Par-complex which regulates asymmetric neuroblast divisions;

2) _PCNA, RFC38, dacapo,_ and _Reptin_ (part of the Tip60 complex), which participate in the molecular

network controlling cell polarity, asymmetric NSC division and cell fate (NEUMÜLLER _et al._ 2011); 3)

CSN1b, CSN4 and _non-stop_ which are implicated in lamina glia cell migration and optic lobe development

(WEAKE _et al._ 2008).

Categorization of the embryonic DEGs allowed us to identify potential pathways in which CLAMP is likely

to regulate target genes. Overall, lack of CLAMP results in dysregulation of three inter-dependent gene/TF

networks: 1) neurogenesis/gliogenesis; 2) cell cycle/asymmetric division; and 3) neuronal

arborization/synaptogenesis.

Transcriptomic analysis of overlapping embryo and L3 *clamp* null DEGs implicates CLAMP in

regulating embryonic and post-embryonic neurogenesis.

The transition from the embryonic to the post-embryonic stage is accompanied by many physiological

changes with the most notable being the introduction of feeding which brings about a cascade of metabolic

signal that induces activation of quiescent NSCs and neurogenesis. Comparison of the *clamp* null embryo

and L3 mRNA-seq results revealed that many DEGs were shared amongst the two stages (2363 DEGs)

(Fig.4b). Meta-analysis of the top two GO categories arising from the DEGs shared between embryos and

L3 DEGs revealed that identified genes have key roles in nervous system development, neurogenesis, and

mitotic cell cycle, suggesting continuing roles in neural development pre- and post-embryonically.

The fact that cell cycle-related categories were enriched in our transcriptomic analysis is interesting because

CLAMP maintains its expression in L3 proliferating NSCs and co-localizes with PCNA (Figs.1i and j),

which has critical roles in the cell cycle (KURKI *et al.* 1986). The expression of *pcna* is downregulated in

both developmental stages (*clamp* null embryo L2FC: -1.692 and L3 larva L2FC: -0.824, Supplementary

Tables 1a and 2a) along with other cell cycle regulators/chromosome dynamics factors such as *AurB*, *atms*

and *baz*, suggesting that CLAMP may regulate cell cycle progression and dynamics directly at a subset of

its targets.

CLAMP regulates genes controlling NSC attributes, glia development and function

The L3 stage encompasses a dramatic shift in the establishment of the nervous system. A plethora of

signaling pathways, including metabolic signals stimulated by larval feeding (CHELL AND BRAND

2010)'(BRITTON AND EDGAR 1998; LANET *et al.* 2013), are integrated at the L3 stage to allow NSCs to

promote maturation of the nervous system. Consistent with this shift in NSC function, GO analysis of the

L3 DEGs (Supplementary Table 2b) revealed an enrichment in several broad categories, including

neurogenesis, mitotic DNA replication, and neuroblast differentiation. We find expression changes in many

genes critical for nervous system development, including those in the following categories: 1) neuroblast

attributes; 2) cell cycle dynamics; 3) glial cell development; 4) optic lobe development; 5) ecdysone &

insulin signaling dynamics; 6) lipid metabolism; 7) CLAMP-related interactors; 8) Synapse function &

structure.

Next, we classified DEGs as downregulated or upregulated in the *clamp* null mutant (downregulated vs.

upregulated, Tables 2 and 3, respectively). Downregulated genes are putatively targets activated by

CLAMP in wild-type animals. These genes could be classified into the following categories: 1) *NSC*

attributes (cell-cycle progression, cell polarity, and survival); 2) *control* of *NSC evolution/differentiation*;

3) *regulation of gliogenesis*; 4) *optic lobe development-specific*. In contrast, upregulated genes that are

putatively repressed by CLAMP are part of the following categories: 1) *lipid metabolism/promoting*

mobilization of lipids; 2) *ecdysone and insulin signaling in glia cells*; 3) *pro-arborization/synapse*

formation and function; 4) *pro-neural/suppression of proliferation*. Based on these data, we hypothesized

that loss of *clamp* leads to deficits in neural stem cell attributes and promotes changes that lead to the

precocious acquisition of a neuronal fate at the expense of gliogenesis. Our expression data also suggest

that loss of *clamp* may cause concomitant activation of lipid metabolism/mobilization as a compensatory

response from the stem cell-niche. Of great interest, we observed a prominent and strong downregulation

in genes orchestrating OL development such as *eya, tll, dac* and *optix* (See Table 2 and Supplementary

Tables 2d and Figs.4b). The downregulation of OL development master regulatory genes in the absence of

clamp aligns with the greatly reduced size of the OL (Fig.3) in the L3 *clamp* null mutants. Collectively,

these data are consistent with a critical role for CLAMP in the regulation of OL NSCs compared with the

rest of the brain.

CLAMP binds major neurodevelopment genes

To evaluate the possibility of direct regulation of DEGs by CLAMP, we determined where CLAMP binding

sites are present upstream of target genes using FIMO analysis (GRANT *et al.* 2011). Previously

characterized CLAMP binding motifs were used for this analysis (KAYE *et al.* 2018). Interestingly, genes

that contain putative CLAMP binding sites and are dysregulated in *clamp* nulls in both the embryonic and

L3 stages include : 1) the important proliferation/mitosis regulators *bazooka (baz), separation anxiety* (*san*),

a N-terminal acetyltransferase required for mitotic chromosome segregation (RIBEIRO *et al.* 2016), *cyclin*

dependent kinase 7 (*cdk7*) vital for mitosis (LAROCHELLE *et al.* 1998), *PCNA* and *Notch*; 2) *sobremesa*

(*sbm*) an SLC7 amino acid transporter expressed in glia and regulating brain development *via* ecdysone

signaling(GALAGOVSKY *et al.* 2018); and *Negative elongation factor E* (*nelf-E*) which is a known interactor

of CLAMP (URBAN 2017) (Fig.4f, Embryo and L3 panel).

Developmental stage-specific genes that contain putative CLAMP binding sites include: A) <u>*Embryo stage*</u>:

anaphase promoting complex subunit 7 (*APC7*)*; pontin (pont)* which is part of the TIP60 complex which

regulates NSC attributes*; aurora A (aurA)* a kinase regulating asymmetric protein localization and thus

required for NSC proliferation (LEE *et al.* 2006)*; extra sex combs (esc)*, a component of the PRC2 complex

which regulates NSC proliferation and neural development (CURT *et al.* 2019) and *fruitless (fru)*, implicated

in the formation of axonal projections in the embryonic CNS (SONG *et al.* 2002) (Fig.4e, Embryo panel).

B) <u>*L3 stage*</u>: Even more interesting during the L3 stage (Fig.4f, L3 panel), CLAMP binding motifs are

found in *gcm,* an essential regulator of glial development*; glass bottom boat* (*gbb*) which regulates both

NB proliferation and surface glia survival; tailless (*tll),* a key regulator of optic lobe development and

anaplastic lymphoma kinase (Alk) which is implicated in cell cycle regulation and NSC proliferation; and

Dichaete (*D*), a TF regulating neuronal fate in the OL.

CLAMP directly binds and regulates Tailless and Notch

To further examine and validate which of the FIMO-identified CLAMP binding targets have been experimentally-confirmed, thus provide the basis for their direct binding and regulation by CLAMP, we mined available embryonic 0-24 hour CLAMP ChIP-seq data generated by modENCODE (CONSORTIUM *et al.* 2010) and L3 whole larval data (KUZU *et al.* 2016) using the Integrative Genomics Viewer (IGV) (ROBINSON *et al.* 2011). This analysis revealed that CLAMP has been experimentally been validated to directly bind to *baz*, *Notch and, Nelf-E,* the latter of which serves as a positive control to our results as it is a validated target of CLAMP (URBAN 2017) (Fig.4g). Importantly, CLAMP directly binds and regulates *Tll* expression (Fig.4g), which indicates that the defects in OL development seen in the *clamp* null mutants is due to downregulation of *tll* expression in the absence of CLAMP.

To further assess direct regulation by CLAMP, we quantified CLAMP binding to DEGs in both our data sets compared to a control gene set of neuronally expressed genes. In support of a direct regulatory role for CLAMP at a subset of its target genes, we found that genes that are downregulated in *clamp* nulls show higher average read depth around the TSS (Fig. 4h) and have a greater binding frequency, particularly in larvae.

CLAMP promotes NSC proliferation and prevents precocious neurogenesis

Our transcriptomics analysis results demonstrate that CLAMP is exerting direct control on neural development by directly regulating the transcription of key effectors that function in multiple processes including asymmetric cell division, cell cycle and optic lobe development (DANIEL *et al.* 1999; DAS AND BHADRA 2020), both in embryos and larvae (Figs.4f, g). Importantly, we observed a marked decrease in glial-development and cell cycle dynamics genes in the absence of CLAMP (Table 2).Therefore, we

hypothesized that lack of CLAMP disturbs the balance between neuronal and glia formation through the

L3 stage, by impinging neural stem cell proliferation, especially in the OL.

To test our hypothesis, we evaluated mid-late L3 brains for proliferating NSCs using the S-phase marker

EdU, the NSC marker Dpn and a quantification approach introduced by Kanai et al.(KANAI *et al.* 2018) in

which the superficial layers of the brain are evaluated. We chose this approach due to the importance of

superficial glia in the overall biology of neuroblasts and used it throughout our study. Consistent with our

initial observations (Fig.3), the OL in *clamp* null larvae brains was dramatically reduced in size compared

to controls, and both the CB and OL contained fewer proliferating NSCs (EdU$^+$Dpn$^+$, Figs.5a and c and

Supplementary Figs.2 a, b). Quantification of total proliferating NSCs (EdU+) from the superficial layers

of the brain (KANAI *et al.* 2018), showed a dramatic ~90% reduction compared to controls (Fig. 5e).

Moreover, we observe an overall reduced number of proliferating cells compared to controls, suggesting a

reduced proliferation of stem cells and/or change in cell cycle dynamics in the absence of CLAMP (EdU+

Fig.5e).

Genes encoded by many pro-neural TFs are upregulated in the absence of CLAMP (Table 3), in addition

to TFs that promote cell cycle regulation in proliferating stem cells. Specifically, in the L3 *clamp* null brain

dpn, which regulates NSC proliferation, was significantly downregulated (Table 2; L2FC: -0.665, padj:

0.00245). The combination of reduced *dpn* transcript levels and the reduced numbers of proliferating NSCs

in the *clamp* null brains suggest that CLAMP may regulate NSC differentiation.

The TF Prospero (Pros), acts as a master regulator driving the decision of stem cells to either self-renew or

differentiate (CHOKSI *et al.* 2006), with nuclear Pros signaling the initiation of neuronal differentiation

(KNOBLICH AND JAN 1995; CHOKSI *et al.* 2006). Thus, to evaluate changes in neuronal production in the

clamp null brain we performed Pros immunostaining. We found that the *clamp* null brain contained more

clusters of nuclear Pros immunostaining than matched controls, thus indicating a rapid change in NSC cell

fate attributes (COLONQUES *et al.* 2011) (Figs.5b and d, arrows indicate Pros+ clusters). Interestingly, while

the total number of NSCs in the *clamp* null brain was reduced by approximately 70% (Dpn$^+$, Fig.5f), Pros+

cell numbers were increased by approximately 70% (Pros$^+$, Fig.5f), indicating that stem cells are acquiring

a precocious neuronal fate. The reduction in NSCs is particularly evident in the OL, where very few Dpn+

cells were observed, contrary to a widespread presence of Pros+ cells ~~were present~~ (Figs.5a-d, outlines in

Figs.5b and d indicate the OL). Pros, acts a transcriptional repressor of self-renewal genes and induces

differentiation (CHOKSI *et al.* 2006; LAI AND DOE 2014). While a significantly higher number of cells have

nuclear Pros (Pros+, Sup.Fig.2c) in the *clamp* null brains, most of these cells have not incorporated EdU

(Pros+EdU-, Sup.Fig.2c). In combination with the finding that in the *clamp* null brain, a lower number of

cells incorporate EdU within the pulse window, and thus a smaller number of NSCs are actively

proliferating (Fig.5e), the Pros immunostaining suggests that in the absence of CLAMP, NSCs undergo

fewer cell division cycles and/or precociously exit the cell cycle to acquire a differentiated fate.

To further evaluate changes in the neuronal pool of cells, we performed Elav staining to label neurons

(ROBINOW AND WHITE 1988), which revealed that *clamp* null brains contain a higher density of Elav$^+$

clusters compared to controls (Figs.5g and h). However, even though the overall Elav+ cluster density

increased the Elav$^+$ zone in the OL was reduced compared to controls (Figs. 5g and h, yellow outlines).

Immunostaining with the apoptotic marker Dcp1 revealed increased and specific presence of Dcp1$^+$ nuclei

in the OL in *clamp* nulls, compared to the controls which instead, had more widespread apoptosis (Figs. 5i

and j). Moreover, in more dorsal optical sections large Dcp1+ clusters were visible in the medulla and

lamina area (Supplementary Figs.3a and b). These results suggest that in the absence of CLAMP a pro-

differentiation pathway is activated in stem cells at the expense of their self-renewal which especially

impacts the OL.

Expression of the optic lobe development master regulator Tailless, is severely reduced in the clamp null

Our results thus far revealed a direct control of the key OL development gene, *tailless* by CLAMP,

culminating in a severely underdeveloped OL in the *clamp* null and downregulation of *tll* in the *clamp* null

(Fig.4 and Fig.6). Interestingly, in *tll* knockdown animals, the OL is severely underdeveloped or absent,

OL neurogenesis is affected and OL neuroblasts undergo increased apoptosis, as seen in the *clamp* null OLs

(GUILLERMIN *et al.* 2015). To examine whether the *clamp* null OL phenotype can be attributed to a loss of

the Tll protein, we performed immunofluorescence. This approach revealed a consistency between the

dramatic reduction in the *tll* transcript identified by mRNA-seq and the protein, with the latter being almost

absent in the OL neuroepithelium (Fig.6d). *tll* expression is required for the survival of the OL NSCs since

its absence leads to apoptosis (GUILLERMIN *et al.* 2015), which is in agreement with the increased and

specific cell-death we observed in the OL region (Figs. 5i and j and Supplementary Figs.3a and b).

Furthermore, it highlights that CLAMP has a direct control in OL development through binding to the *tll*

promoter to regulate *tll* expression.

Lack of CLAMP results in reduction in number of surface glia

632
OL development is highly dependent on glia-neuron interactions because glia form a supportive niche for

neural stem cells to function (PEREANU *et al.* 2005; RAMON-CAÑELLAS *et al.* 2018). Furthermore,

neuroepithelial cells of the OL, give rise to hundreds of precursor/progenitor cells of different lineages,

including neuroglioblasts and glial progenitor cells. For the lineage diversification to take place an intricate

temporal expression of fate-determining TFs occurs in the precursor cells, including *Notch* and *Tll,* both of

which are under direct CLAMP control (Fig.4g). In OL medulla NBs, Tll is expressed in the oldest NBs

which go on to generate neuronal progeny or express *glia cells missing (gcm)* and gradually turn off <u>*Tll*</u> to

express *Repo* as glial cells (LI *et al.* 2013).

Our transcriptomic analysis supports the hypothesis that lack of CLAMP promotes neuronal differentiation

at the expense of gliogenesis from the late embryo stage. Therefore, the defects observed in OL

development of the L3 *clamp* null could be attributed to a combination of deficits in gliogenesis, precocious

neurogenesis as a result of altered *Tll and/or Notch* expression. It was recently shown that surface glia (SG)

are crucial for NB proliferation (KANAI *et al.* 2018), an interplay which involves niche and autocrine

signaling. Specifically, SG regulate NSC proliferation *via* the secretion of the proteoglycan Dally-like,

(Dlp); 2) Stem cells secrete the BMP-ligand Glass bottom boat (Gbb) which regulates their proliferation

and both the stem cell and SG survival. Interestingly, the *gbb* gene contains CLAMP-binding motifs

(Fig.4f) and is downregulated in the *clamp* null L3 brains (Supplementary Table 2d). Furthermore, we

observed enrichment of the BMP-signaling GO category in the *clamp* null (Fig.4d). Therefore, a loss of the

dynamic interplay between SG and NSCs resulting from the altered expression of key fate-determining

TFs, *Tll* and *Notch* in the absence of CLAMP could also contribute the deficits in the *clamp* null brain.

To test this hypothesis, we first examined whether a change in SG numbers is seen in the *clamp* null. We

used the glia-specific marker Repo (XIONG *et al.* 1994b; YUASA *et al.* 2003) to mark glial cells, and

quantified SG in L3 brains using the approach by Avet-Rochex et al. (2012) (AVET-ROCHEX *et al.* 2012).

This revealed a ~25 % reduction in SG number in the *clamp* null, compared to the control (Figs.7 a-d).

Additionally, the distribution and numbers of glial cells throughout the brain (Supplementary Figs. 3c and

d) and especially of the OL and the lobula plug was altered in terms of topological organization

(Supplementary Figs. 3e and f). Overall, these data suggest that CLAMP regulates gliogenesis and glia

distribution especially in the OL.

To determine if the effect on SG number was due to the lack of CLAMP in glia specifically, we used the

GAL4-UAS system to induce *clamp*-RNAi knock down (KD) using the glia-specific driver, *repo-GAL4*

(XIONG et al. 1994a). Interestingly, we find that in the *repo>clamp RNAi* brains, the number of SG was

reduced to a similar extent as in the *clamp* nulls (Repo+ cells, ~30% reduction, Figs. 7a-d). To determine

whether glia-specific or neuronal-specific *clamp* depletion has differential effects on the proliferation

(EdU+) and the number of NSCs (Dpn+ cells), we also performed neuronal *clamp* KD by crossing to an

elav-GAL4 fly line (LIN AND GOODMAN 1994). Comparison with the data we already obtained from the

clamp null and control brains (see Figs.5e and f) revealed that *clamp* KD in neuronal lineages did not result

in significant changes either in EdU+ or Dpn+ cells, while glia-specific *clamp* KD caused a reduction in

the number of total proliferating cells (EdU+ cells, ~15% reduction compared to control). Thus, while glial

clamp KD results in the same amount of reduction in Repo+ SG number as in the *clamp* null, the effect on

proliferating NSCs is smaller compared to the null. This suggests that downregulation of glial CLAMP

mainly affects the proliferation of glia-producing NSCs likely *via* both cell autonomous mechanisms (e.g.,

reduction in Dpn+ gliogenic NSCs) and *via* non-cell autonomous mechanisms (regulation of proliferation

by niche glia), although it does not recapitulate the full phenotype seen in the *clamp* null leaving room for

additional mechanisms.

In summary, our results suggest that CLAMP controls NSC stemness, proliferation and lineage output,

especially in the OL, by directly regulating the expression of key TFs such as *Notch* and *Tailless,* and by

affecting neural stem cell niche homeostasis. Ultimately, lack of CLAMP leads to imbalance in neuronal

over glial generation and leads to precocious differentiation of neural stem cells.

Discussion

The understanding of the gene regulatory mechanisms by which neural stem cells function to ultimately

build the brain is critical to provide insight into normal. Signaling relays originating from the stem niche

play an instructive role in the neurogenetic process, and highlight the fact that the environmental context

acts synergistically with intrinsic developmental signals (SCADDEN 2006; SPEDER AND BRAND 2018). In

this study, we have defined a new role for the essential transcription factor, CLAMP, in nervous system

development and NSC-niche interactions, especially in optic lobe development. We show that CLAMP

positively directly regulates *Notch* and *Tailless*, which regulate NSC stemness, proliferation and progeny

output, while *Tailless* is a master regulator of OL development. The multilayered control of CLAMP over

NSC function is evidenced in *clamp* null brains, where the OL is severely underdeveloped, niche glial cell

generation is reduced in favor of precocious neurogenesis, while concurrently the niche protective

mechanisms are overactivated. Furthermore, lack of *clamp* affects the expression of key NSC

proliferation/division genes such as *baz*. In summary, our study reveals that CLAMP regulates brain

development *via* cell-intrinsic and NSC niche-dependent mechanisms that involve niche glia.

The formation of the OL is profoundly impaired in *clamp* null mutants, a phenotype accompanied by

reduction of EdU$^+$ NSCs (Figs.5a-d). We also observed increased numbers of Pros$^+$ cells in the null OL

(Figs. 5b and d, outlined area) and throughout the brain, consistent with lack of CLAMP promoting a

neurogenic over a gliogenic fate. While glial *clamp* KD results in a partial reduction in the number of

proliferating NSCs (EdU$^+$, Fig.6e), it does not affect the number of Dpn$^+$ NSCs/neuroblasts. These results

suggest an effect on the Dpn$^-$ progeny of the NSCs, such as the ganglion mother cells (GMCs) and common

progenitor cells (CPCs) of the OL. This is an intriguing scenario as CPCs divide asymmetrically to generate

Lawf precursor cells and glial precursor cells (GPCs); glial *clamp* KD resulted in the approximately the

same reduction in Repo+ glial cells, as seen in the *clamp* null. Although, Pros+ cells were not counted in

the glial *clamp* KD brains, evaluation of the mean signal intensity of the Pros channel suggested an

increased signal compared to the control (not shown) and suggesting the lack of CLAMP alters the balance

between neuronal and glial-progenies at the precursor level. Also, it is possible that there are changes in the

cell cycle kinetics that cause fewer NSCs to be actively proliferating. This possibility is consistent with the

GO analysis which identified genes involved in cell cycle kinetics/transitions (Fig4.c).

Moreover, as CLAMPs' specific role in OL development is evidenced by the fact that CLAMP directly

binds and positively regulates *tll* expression, which is also downregulated both at the transcript and protein

levels in the OL, in the absence of CLAMP (Fig.4). Furthermore, other transcription factors critical for OL

neurogenesis such as *eya, dac* and *optix* are downregulated in the *clamp* null mutant brains (see Table 2).

In the post-embryonic nervous system, *gcm* and *gcm2* are expressed in OL NSCs destined both for glia and

neuron generation which they regulate *via* downstream mechanisms involving Dachshund (Dac)(CHOTARD

et al. 2005). The expression of g*cm, gcm2* and *dac* is reduced in the *clamp* null (Table 2). Moreover, both

gcm and *gcm2* transcript levels were greatly reduced in the *clamp* null L3 brain, while *gcm* contains CLAMP

binding sites (Fig.4e). The OL contains a special population of NSCs, the common progenitor cells (CPCs),

which generate both glia precursor cells (GPCs) and neurons (Lawfs), under the control of Notch signaling

(CHEN *et al.* 2016). Moreover, Notch is a key regulator which maintains the balance between

symmetric/self-renewing and asymmetric NSCs divisions; loss of Notch function promotes precocious

differentiation and premature neurogenesis (EGGER *et al.* 2010). In the absence of CLAMP, *Notch* is at

least two-fold downregulated both in the embryonic and L3 stages (Supplementary Table 1a and Table 2),

and we determined that Notch has a CLAMP binding site and is a direct target for CLAMP (Figs.4f and g).

Overall, our data support a model in which CLAMP functions upstream of *Notch* and *Tll* signaling, which

it directly binds, and regulates NSC proliferation and differentiation in the OL.

Neural development and NSC function are intricately linked to niche glia and depend on support and

nutrition from surface glia *via* Gbb-Dlp signaling, which sustains NSC proliferation and survival but also

ensures the survival of surface glia (KANAI *et al.* 2018). Thus, there is an interdependency between NSCs

and niche glia, ensuring overall homeostasis in the stem cell niche. In *clamp* null and *repo>clamp* RNAi

larvae brains, surface glia numbers are reduced to the same extent indicating that lack of glial CLAMP

affects gliogenesis (Fig.7d). Furthermore, the glial master gene, *gcm,* contains CLAMP binding motifs and

is significantly downregulated in the *clamp* null brain, indicating that CLAMP may directly regulate

gliogenesis through *gcm*-related mechanisms (Fig.4f). Supportive of a glial role for CLAMP is the fact that

glial-specific *clamp* KD by *repo*-GAL4, resulted in specific reduction of SG cells, to the same extend seen

in the *clamp* null. In contrast, a neuronal specific *clamp* KD (*elav> clamp* RNAi) did not affect NSC

proliferation, number or, glial cell number (Fig.6). A small reduction in cycling cells (~15%, EdU+) was

observed in *repo>clamp* RNAi conditions compared to a massive 85% reduction in the *clamp* null brain.

This difference can be attributed to the fact that in the *repo>clamp* RNAi conditions, the reduction in *clamp*

levels will only take place after *repo* expression is established during terminal differentiation of glial cells,

and through direct induction by Gcm (YUASA *et al.* 2003).

In the future, it will be interesting to determine whether this reduction in glial cell number can be attributed

specifically to glia-producing cells in which *gcm* activity is still high and indicative of their proliferation

status. In the *clamp* null mutants, *clamp* is depleted before *repo* starts to be expressed and thus may result

in a greater impact on NSC proliferation and neuronal production in the OL because both *gcm* and *gcm2*

regulate glial and neuronal development in the OL via *Dac* (CHOTARD *et al.* 2005) which is also

downregulated in the *clamp* null brain (Table 2). Thus, it is possible if we used *gcm*-GAL4 to drive *clamp*

RNAi at earlier stages, we would unmask a greater effect both on NSC proliferation and NSC number.

We also observed that the *foxO* gene (Forkhead box, class O) is part of many of the cell cycle sub-categories

that we identified from GO analysis, including the *insulin signaling category*, which is a known target of

FOXO (KRAMER *et al.* 2003). FOXO belongs to a family of highly dynamic and conserved transcription

factors and regulates gene expression during processes such as apoptosis, cell-cycle transitions including

negative regulation of G1/S and G2/M (HUANG AND TINDALL 2007), both of which are highly enriched in

our GO analysis. Overexpression of FOXO during larval development results in reduction both in cell size

and number in the developing eye (KRAMER *et al.* 2003). In addition, it was shown that FOXO affects the

temporal sequence and cell fate specification in a subset of olfactory neurons in *Drosophila* (WEI *et al.*

2019). Interestingly, in the *clamp* null embryo the *foxO* transcript is upregulated (L2FC: 0.46846,

padj:0.000283, Supplementary Table 1a). Therefore, it is possible that some of the phenotypic effects that

we observed in the *clamp* null mutant are also related to dysregulation of *foxO* gene expression (KRAMER

et al. 2003; WEI *et al.* 2019) .

Clamp is maternally deposited and its expression is high during the majority of embryogenesis (RIEDER *et*

al. 2017) (Fig.1a). While the persistence of maternal *clamp* was not specifically evaluated in late embryos,

CLAMP protein deposited by *clamp* heterozygous mothers persists beyond the embryonic stage and is

present in the histone locus body throughout the development of the larval salivary gland (RIEDER *et al.*

2017). Thus, it is possible that a trace amount of maternally deposited *clamp* could still be present at 16-18

hour embryos and reduces the strength of the zygotic *clamp* nulls phenotype compared with embryos that

are depleted of maternal CLAMP which die early in embryogenesis soon after zygotic genome activation.

Glial cells can arise from different mechanisms, either from dedicated glioblasts or from NBs that have

both glial and neuronal potential (BOSSING *et al.* 1996; BERNARDONI *et al.* 1999; SCHMID *et al.* 1999;

UDOLPH *et al.* 2001). Our results show that CLAMP regulates a transcriptional program which drives NSC

proliferation and differentiation *via* cell-intrinsic and niche-dependent mechanisms that involve

transcriptional regulation of *Tailless, Notch* and niche glia.

799   REFERENCES
800

Anders, S., P. T. Pyl and W. Huber, 2015 HTSeq—a Python framework to work with high-throughput

sequencing data. Bioinformatics 31**:** 166-169.

Andrews, S., 2010 FastQC: a quality control tool for high throughput sequence data, pp. Babraham

Bioinformatics, Babraham Institute, Cambridge, United Kingdom.

Arefin, B., F. Parvin, S. Bahrampour, C. B. Stadler and S. Thor, 2019 Drosophila Neuroblast Selection Is

Gated by Notch, Snail, SoxB, and EMT Gene Interplay. Cell reports 29**:** 3636-3651. e3633.

Avet-Rochex, A., A. K. Kaul, A. P. Gatt, H. McNeill and J. M. Bateman, 2012 Concerted control of

gliogenesis by InR/TOR and FGF signalling in the Drosophila post-embryonic brain. Development

139**:** 2763-2772.

Bailey, Andrew P., G. Koster, C. Guillermier, Elizabeth M. A. Hirst, James I. MacRae *et al.*, 2015

Antioxidant Role for Lipid Droplets in a Stem Cell Niche of Drosophila. Cell 163**:** 340-353.

Bernardoni, R., M. Kammerer, J.-L. Vonesch and A. Giangrande, 1999 Gliogenesis depends on glide/gcm

through asymmetric division of neuroglioblasts. Developmental biology 216**:** 265-275.

Bertet, C., X. Li, T. Erclik, M. Cavey, B. Wells *et al.*, 2014 Temporal Patterning of Neuroblasts Controls

Notch-Mediated Cell Survival through Regulation of Hid or Reaper. Cell 158**:** 1173-1186.

Bossing, T., G. Udolph, C. Q. Doe and G. M. Technau, 1996 The Embryonic Central Nervous System

Lineages ofDrosophila melanogaster: I. Neuroblast Lineages Derived from the Ventral Half of the

Neuroectoderm. Developmental biology 179**:** 41-64.

Britton, J. S., and B. A. Edgar, 1998 Environmental control of the cell cycle in Drosophila: nutrition

activates mitotic and endoreplicative cells by distinct mechanisms. Development 125**:** 2149-2158.

Brown, J. B., N. Boley, R. Eisman, G. E. May, M. H. Stoiber *et al.*, 2014 Diversity and dynamics of the

Drosophila transcriptome. Nature 512**:** 393-399.

Chell, J. M., and A. H. Brand, 2010 Nutrition-responsive glia control exit of neural stem cells from

quiescence. Cell 143**:** 1161-1173.

Chen, Z., A. D. V. Rodriguez, X. Li, T. Erclik, V. M. Fernandes *et al.*, 2016 A unique class of neural

progenitors in the Drosophila optic lobe generates both migrating neurons and glia. Cell reports 15**:**

774-786.

Choksi, S. P., T. D. Southall, T. Bossing, K. Edoff, E. de Wit *et al.*, 2006 Prospero acts as a binary switch

between self-renewal and differentiation in Drosophila neural stem cells. Developmental cell 11**:**

775-789.

Chotard, C., W. Leung and I. Salecker, 2005 glial cells missing and gcm2 cell autonomously regulate both

glial and neuronal development in the visual system of Drosophila. Neuron 48**:** 237-251.

Colonques, J., J. Ceron, H. Reichert and F. J. Tejedor, 2011 A transient expression of Prospero promotes

cell cycle exit of Drosophila postembryonic neurons through the regulation of Dacapo. PLoS One

6**:** e19342.

Conard, A. M., N. Goodman, Y. Hu, N. Perrimon, R. Singh *et al.*, 2020 TIMEOR: a web-based tool to

uncover temporal regulatory mechanisms from multi-omics data. bioRxiv.

Consortium, m., S. Roy, J. Ernst, P. V. Kharchenko, P. Kheradpour *et al.*, 2010 Identification of functional

elements and regulatory circuits by Drosophila modENCODE. Science 330**:** 1787-1797.

Contreras, E. G., B. Egger, K. S. Gold and A. H. Brand, 2018 Dynamic Notch signalling regulates neural

stem cell state progression in the Drosophila optic lobe. Neural development 13**:** 1-11.

Contreras, E. G., J. Sierralta and C. Oliva, 2019a Novel strategies for the generation of neuronal diversity:

lessons from the fly visual system. Frontiers in Molecular Neuroscience 12**:** 452461.

Contreras, E. G., J. A. Sierralta and C. Oliva, 2019b Novel strategies for the generation of neuronal

diversity: lessons from the fly visual system. Frontiers in Molecular Neuroscience 12**:** 140.

Crews, S. T., 2019 Drosophila Embryonic CNS Development: Neurogenesis, Gliogenesis, Cell Fate, and

Differentiation. Genetics 213**:** 1111-1144.

Curt, J. R., B. Y. Salmani and S. Thor, 2019 Anterior CNS expansion driven by brain transcription factors.

Elife 8**:** e45274.

Dani, N., R. H. Herbst, N. Habib, J. Head, D. Dionne *et al.*, 2019 A cellular and spatial map of the choroid

plexus across brain ventricles and ages. BioRxiv**:** 627539.

Daniel, A., K. Dumstrei, J. A. Lengyel and V. Hartenstein, 1999 The control of cell fate in the embryonic

visual system by atonal, tailless and EGFR signaling. Development 126**:** 2945-2954.

Das, P., and M. P. Bhadra, 2020 Histone deacetylase (Rpd3) regulates Drosophila early brain development

via regulation of Tailless. Open Biol 10**:** 200029.

DeSimone, S. M., and K. White, 1993 The Drosophila erect wing gene, which is important for both neuronal

and muscle development, encodes a protein which is similar to the sea urchin P3A2 DNA binding

protein. Molecular and cellular biology 13**:** 3641-3649.

Egger, B., J. M. Chell and A. H. Brand, 2008 Insights into neural stem cell biology from flies. Philosophical

Transactions of the Royal Society B: Biological Sciences 363**:** 39-56.

Egger, B., K. S. Gold and A. H. Brand, 2010 Notch regulates the switch from symmetric to asymmetric

neural stem cell division in the Drosophila optic lobe. Development 137**:** 2981-2987.

Galagovsky, D., A. Depetris-Chauvin, G. Manière, F. Geillon, M. Berthelot-Grosjean *et al.*, 2018

Sobremesa L-type Amino Acid Transporter Expressed in Glia Is Essential for Proper Timing of

Development and Brain Growth. Cell reports 24**:** 3156-3166. e3154.

Gallaud, E., T. Pham and C. Cabernard, 2017 Drosophila melanogaster neuroblasts: a model for asymmetric

stem cell divisions, pp. 183-210 in *Asymmetric Cell Division in Development, Differentiation and*

Cancer. Springer.

Grant, C. E., T. L. Bailey and W. S. Noble, 2011 FIMO: scanning for occurrences of a given motif.

Bioinformatics 27**:** 1017-1018.

Graveley, B., G. May, A. Brooks, J. Carlson, L. Cherbas *et al.*, 2011 The D. melanogaster transcriptome:

modENCODE RNA-Seq data for dissected tissues. Retrieved from website: http://www.

modencode. org/Celniker. shtml.

Guillermin, O., B. Perruchoud, S. G. Sprecher and B. Egger, 2015 Characterization of tailless functions

during Drosophila optic lobe formation. Developmental Biology 405**:** 202-213.

Hakes, A. E., and A. H. Brand, 2019 Neural stem cell dynamics: the development of brain tumours. Current

Opinion in Cell Biology 60**:** 131-138.

Hakes, A. E., L. Otsuki and A. H. Brand, 2018 A newly discovered neural stem cell population is generated

by the optic lobe neuroepithelium during embryogenesis in *Drosophila melanogaster*.

Development 145**:** dev166207.

Hartenstein, V., and J. A. Campos-Ortega, 1984 Early neurogenesis in wild-typeDrosophila melanogaster.

Wilhelm Roux's archives of developmental biology 193**:** 308-325.

Haussmann, I. U., and M. Soller, 2010a Differential activity of EWG transcription factor isoforms identifies

a subset of differentially regulated genes important for synaptic growth regulation. Developmental

biology 348**:** 224-230.

Haussmann, I. U., and M. Soller, 2010b Differential activity of EWG transcription factor isoforms identifies

a subset of differentially regulated genes important for synaptic growth regulation. Dev Biol 348**:**

224-230.

Haussmann, I. U., K. White and M. Soller, 2008 Erect wing regulates synaptic growth in Drosophila by

integration of multiple signaling pathways. Genome biology 9**:** R73.

Hindle, S. J., and R. J. Bainton, 2014 Barrier mechanisms in the Drosophila blood-brain barrier. Frontiers

in neuroscience 8**:** 414.

Huang, H., and D. J. Tindall, 2007 Dynamic FoxO transcription factors. J Cell Sci 120**:** 2479-2487.

Janssens, D. H., H. Komori, D. Grbac, K. Chen, C. T. Koe *et al.*, 2014 Earmuff restricts progenitor cell

potential by attenuating the competence to respond to self-renewal factors. Development 141**:**

1036-1046.

Kanai, M. I., M.-J. Kim, T. Akiyama, M. Takemura, K. Wharton *et al.*, 2018 Regulation of neuroblast

proliferation by surface glia in the Drosophila larval brain. Scientific reports 8**:** 1-15.

Kaye, E. G., M. Booker, J. V. Kurland, A. E. Conicella, N. L. Fawzi *et al.*, 2018 Differential occupancy of

two GA-binding proteins promotes targeting of the drosophila dosage compensation complex to

the male X chromosome. Cell reports 22**:** 3227-3239.

Kaye, E. G., A. Kurbidaeva, D. Wolle, T. Aoki, P. Schedl *et al.*, 2017 Drosophila dosage compensation
loci associate with a boundary-forming insulator complex. Molecular and cellular biology 37**:**
e00253-00217.

Kim, D., B. Langmead and S. Salzberg, 2015 hisat2. Nature methods.

Kis, V., B. Barti, M. Lippai and M. Sass, 2015 Specialized cortex glial cells accumulate lipid droplets in
Drosophila melanogaster. PLoS One 10**:** e0131250.

Knoblich, J. A., and Y. N. Jan, 1995 Asymmetric segregation of Numb and Prospero during cell division.
Nature 377**:** 624-627.

Konstantinides, N., K. Kapuralin, C. Fadil, L. Barboza, R. Satija *et al.*, 2018 Phenotypic convergence:
distinct transcription factors regulate common terminal features. Cell 174**:** 622-635. e613.

Kramer, J. M., J. T. Davidge, J. M. Lockyer and B. E. Staveley, 2003 Expression of Drosophila FOXO
regulates growth and can phenocopy starvation. BMC developmental biology 3**:** 5.

Kurki, P., M. Vanderlaan, F. Dolbeare, J. Gray and E. Tan, 1986 Expression of proliferating cell nuclear
antigen (PCNA)/cyclin during the cell cycle. Experimental cell research 166**:** 209-219.

Kuzu, G., E. G. Kaye, J. Chery, T. Siggers, L. Yang *et al.*, 2016 Expansion of GA dinucleotide repeats
increases the density of CLAMP binding sites on the X-chromosome to promote Drosophila dosage
compensation. PLoS genetics 12**:** e1006120.

Lai, S.-L., and C. Q. Doe, 2014 Transient nuclear Prospero induces neural progenitor quiescence. eLife 3**:**
e03363.

Lanet, E., A. P. Gould and C. Maurange, 2013 Protection of neuronal diversity at the expense of neuronal
numbers during nutrient restriction in the Drosophila visual system. Cell reports 3**:** 587-594.

Larochelle, S., J. Pandur, R. P. Fisher, H. K. Salz and B. Suter, 1998 Cdk7 is essential for mitosis and for
in vivo Cdk-activating kinase activity. Genes & development 12**:** 370-381.

Larschan, E., M. M. Soruco, O. K. Lee, S. Peng, E. Bishop *et al.*, 2012 Identification of chromatin-
associated regulators of MSL complex targeting in Drosophila dosage compensation. PLoS Genet
8**:** e1002830.

Lee, C.-Y., R. O. Andersen, C. Cabernard, L. Manning, K. D. Tran *et al.*, 2006 Drosophila Aurora-A kinase
inhibits neuroblast self-renewal by regulating aPKC/Numb cortical polarity and spindle orientation.
Genes & development 20**:** 3464-3474.

Li, X., T. Erclik, C. Bertet, Z. Chen, R. Voutev *et al.*, 2013 Temporal patterning of Drosophila medulla
neuroblasts controls neural fates. Nature 498**:** 456-462.

Lin, D. M., and C. S. Goodman, 1994 Ectopic and increased expression of Fasciclin II alters motoneuron
growth cone guidance. Neuron 13**:** 507-523.

Love, M., S. Anders and W. Huber, 2014 Differential analysis of count data–the DESeq2 package. Genome
Biol 15**:** 10.1186.

Maroni, G., and S. Stamey, 1983 Use of blue food to select synchronous, late third instar larvae. Dros. Inf.
Serv 59**:** 142-143.

Morante, J., D. M. Vallejo, C. Desplan and M. Dominguez, 2013 Conserved miR-8/miR-200 defines a glial
niche that controls neuroepithelial expansion and neuroblast transition. Developmental cell 27**:**
174-187.

Neumüller, Ralph A., C. Richter, A. Fischer, M. Novatchkova, Klaus G. Neumüller *et al.*, 2011 Genome-
Wide Analysis of Self-Renewal in Drosophila Neural Stem Cells by Transgenic RNAi. Cell Stem
Cell 8**:** 580-593.

Pereanu, W., D. Shy and V. Hartenstein, 2005 Morphogenesis and proliferation of the larval brain glia in
Drosophila. Developmental biology 283**:** 191-203.

Prokop, A., and G. M. Technau, 1991 The origin of postembryonic neuroblasts in the ventral nerve cord of
Drosophila melanogaster. Development 111**:** 79-88.

Ramon-Cañellas, P., H. P. Peterson and J. Morante, 2018 From early to late neurogenesis: Neural
progenitors and the glial niche from a fly's point of view. Neuroscience.

Ramon-Cañellas, P., H. P. Peterson and J. Morante, 2019 From early to late neurogenesis: Neural
progenitors and the glial niche from a fly's point of view. Neuroscience 399**:** 39-52.

Raudvere, U., L. Kolberg, I. Kuzmin, T. Arak, P. Adler *et al.*, 2019 g:Profiler: a web server for functional
enrichment analysis and conversions of gene lists (2019 update). Nucleic Acids Research 47**:**
W191-W198.

Ribeiro, A. L., R. D. Silva, H. Foyn, M. N. Tiago, O. S. Rathore *et al.*, 2016 Naa50/San-dependent N-
terminal acetylation of Scc1 is potentially important for sister chromatid cohesion. Scientific
reports 6**:** 1-16.

Rieder, L. E., W. T. Jordan and E. N. Larschan, 2019 Targeting of the Dosage-Compensated Male X-
Chromosome during Early Drosophila Development. Cell Reports 29**:** 4268-4275.e4262.

Rieder, L. E., K. P. Koreski, K. A. Boltz, G. Kuzu, J. A. Urban *et al.*, 2017 Histone locus regulation by the
Drosophila dosage compensation adaptor protein CLAMP. Genes & development 31**:** 1494-1508.

Robinow, S., and K. White, 1988 The locus elav of Drosophila melanogaster is expressed in neurons at all
developmental stages. Developmental biology 126**:** 294-303.

Robinson, J. T., H. Thorvaldsdóttir, W. Winckler, M. Guttman, E. S. Lander *et al.*, 2011 Integrative
genomics viewer. Nature Biotechnology 29**:** 24-26.

Scadden, D. T., 2006 The stem-cell niche as an entity of action. Nature 441**:** 1075-1079.

Schmid, A., A. Chiba and C. Q. Doe, 1999 Clonal analysis of Drosophila embryonic neuroblasts: neural
cell types, axon projections and muscle targets. Development 126**:** 4653-4689.

Song, H.-J., J.-C. Billeter, E. Reynaud, T. Carlo, E. P. Spana *et al.*, 2002 The fruitless gene is required for
the proper formation of axonal tracts in the embryonic central nervous system of Drosophila.
Genetics 162**:** 1703-1724.

Soruco, M. M., J. Chery, E. P. Bishop, T. Siggers, M. Y. Tolstorukov *et al.*, 2013a The CLAMP protein
links the MSL complex to the X chromosome during Drosophila dosage compensation. Genes &
development 27**:** 1551-1556.

Soruco, M. M., J. Chery, E. P. Bishop, T. Siggers, M. Y. Tolstorukov *et al.*, 2013b The CLAMP protein
links the MSL complex to the X chromosome during Drosophila dosage compensation. Genes Dev
27**:** 1551-1556.

Sousa-Nunes, R., L. L. Yee and A. P. Gould, 2011 Fat cells reactivate quiescent neuroblasts via TOR and

glial insulin relays in Drosophila. Nature 471**:** 508-512.

Southall, T. D., C. M. Davidson, C. Miller, A. Carr and A. H. Brand, 2014 Dedifferentiation of neurons

precedes tumor formation in Lola mutants. Developmental cell 28**:** 685-696.

Speder, P., and A. H. Brand, 2018 Systemic and local cues drive neural stem cell niche remodelling during

neurogenesis in Drosophila. Elife 7**:** e30413.

Spéder, P., and A. H. Brand, 2014 Gap junction proteins in the blood-brain barrier control nutrient-

dependent reactivation of Drosophila neural stem cells. Developmental Cell 30**:** 309-321.

Tastan, Ö. Y., and J.-L. Liu, 2015 CTP synthase is required for optic lobe homeostasis in Drosophila.

Journal of genetics and genomics 42**:** 261-274.

Thurmond, J., J. L. Goodman, V. B. Strelets, H. Attrill, L. S. Gramates *et al.*, 2019 FlyBase 2.0: the next

generation. Nucleic acids research 47**:** D759-D765.

Tikhonova, E., A. Fedotova, A. Bonchuk, V. Mogila, E. N. Larschan *et al.*, 2019 The simultaneous

interaction of MSL2 with CLAMP and DNA provides redundancy in the initiation of dosage

compensation in Drosophila males. Development 146**:** dev179663.

Turki-Judeh, W., and A. J. Courey, 2012 Groucho: a corepressor with instructive roles in development, pp.

65-96 in *Current topics in developmental biology*. Elsevier.

Turner, F. R., and A. P. Mahowald, 1977 Scanning electron microscopy of Drosophila melanogaster

embryogenesis: II. Gastrulation and segmentation. Developmental biology 57**:** 403-416.

Udolph, G., P. Rath and W. Chia, 2001 A requirement for Notch in the genesis of a subset of glial cells in

the Drosophila embryonic central nervous system which arise through asymmetric divisions.

Development 128**:** 1457-1466.

Urban, J., G. Kuzu, S. Bowman, B. Scruggs, T. Henriques *et al.*, 2017 Enhanced chromatin accessibility of

the dosage compensated Drosophila male X-chromosome requires the CLAMP zinc finger protein.

PloS one 12**:** e0186855.

Urban, J. A., Doherty, C. A, Jordan, W. T, Bliss, J. E, Feng, J., Soruco, M. M, Rieder, L. E, Tsiarli, M. A,

1005   Larschan, E. N, 2017 The essential Drosophila CLAMP protein differentially regulates non-coding

1006   roX RNAs in male and females. Chromosome Research 25**:** 101-113.

1007   Weake, V. M., K. K. Lee, S. Guelman, C. H. Lin, C. Seidel *et al.*, 2008 SAGA-mediated H2B

1008   deubiquitination controls the development of neuronal connectivity in the Drosophila visual

1009   system. The EMBO journal 27**:** 394-405.

1010   Wei, J.-Y., P.-C. Chung, S.-Y. Chu and H. H. Yu, 2019 FOXO regulates cell fate specification of

1011   Drosophila ventral olfactory projection neurons. Journal of neurogenetics 33**:** 33-40.

1012   Xiong, W.-C., H. Okano, N. H. Patel, J. A. Blendy and C. Montell, 1994a repo encodes a glial-specific

1013   homeo domain protein required in the Drosophila nervous system. Genes & development 8**:** 981-

1014   994.

1015   Xiong, W. C., H. Okano, N. H. Patel, J. A. Blendy and C. Montell, 1994b repo encodes a glial-specific

1016   homeo domain protein required in the Drosophila nervous system. Genes & Development 8**:** 981-

1017   994.

1018   Yildirim, K., J. Petri, R. Kottmeier and C. Klämbt, 2019 Drosophila glia: Few cell types and many

1019   conserved functions. Glia 67**:** 5-26.

1020   Yu, G., L.-G. Wang, Y. Han and Q.-Y. He, 2012 clusterProfiler: an R package for comparing biological

1021   themes among gene clusters. Omics: a journal of integrative biology 16**:** 284-287.

1022   Yuan, X., C. W. Sipe, M. Suzawa, M. L. Bland and S. E. Siegrist, 2020 Dilp-2–mediated PI3-kinase

1023   activation coordinates reactivation of quiescent neuroblasts with growth of their glial stem cell

1024   niche. Plos Biology 18**:** e3000721.

1025   Yuasa, Y., M. Okabe, S. Yoshikawa, K. Tabuchi, W. C. Xiong *et al.*, 2003 Drosophila homeodomain

1026   protein REPO controls glial differentiation by cooperating with ETS and BTB transcription factors.

1027   Development 130**:** 2419-2428.

1028   Zaharieva, E., I. U. Haussmann, U. Brauer and M. Soller, 2015 Concentration and Localization of

1029   Coexpressed ELAV/Hu Proteins Control Specificity of mRNA Processing. Mol Cell Biol 35**:** 3104-

1030   3115.

1031

Acknowledgements

We thank Dr. Edward A. Kravitz for providing MAT laboratory space and contributing to supply costs for

completion of this study, Dr. Yick-Bun Chan for experimental support and assistance with DEG analysis

and Dr. Yiannis Savva for experimental support. We would also like to thank Dr. Séverine Trannoy for

critical reading of this manuscript. We thank the Harvard Medical School Neurobiology Department and

the Neurobiology Imaging Facility for consultation and instrument availability that supported this work.

This facility is supported in part by the Neural Imaging Center as part of an NINDS P30 Core Center grant

#NS072030. This work was supported in part by a Harvard Medical School Fix Fund Postdoctoral

Fellowship to M.A.T and by US National Institute of Health grants to E.N.L and E.A.K (R35 GM126994-

01 and R35 GM118137).

1042

Contributions

MAT and E.N.L conceived the study and M.A.T designed the study. M.A.T performed

immunohistochemistry and imaging, collection of RNA-seq samples and analysis of RNA-seq DEG results.

L.X contributed to sample collection for RNA-seq experiments and immunohistochemistry, E.N

contributed in size experiments, A.M.C contributed in RNA-seq data analysis and JK in ChIP-seq data

analysis. M.A.T prepared figures and wrote the original draft while M.A.T and E.N.L curated the

manuscript with input from all authors.

1050

1051   **Declaration of Interests**

1052   The authors declare no competing interests.

1053

1054

1055

Figure 1 CLAMP is ubiquitously expressed throughout the L3 nervous system and is present in

proliferating neuroblasts. (a) Temporal expression data from modEncode (Flybase) reveal that *clamp*

expression levels decrease as embryonic development proceeds and as the maternal transcripts are used-up

and reach a nadir at 18-20 hours after egg lay, which is then followed by a sharp increase just before

embryonic development is completed. During the larval stages *clamp* expression decreases until L3, where

it slowly starts to increase again before the pupa stage. (b) Tissue specific L3 expression data, reveal that

clamp is most highly expressed in the developing imaginal disks, the CNS and the fat body.

(c-d) CLAMP is expressed both in the L3 central nervous system, specifically in the optic lobe (OL, yellow

dashed outline), central brain, (CB) and in the ventral nerve cord (VNC). CLAMP is present in the nucleus

throughout the nervous system as seen by co-staining with DAPI and Nc82, which marks the synaptic

processes. In the VNC it is also observed in the neuronal processes (c, arrows). Grayscale panels show

individual staining channels. (e-h) Higher resolution images reveal high expression of CLAMP in the

developing CB and OL and in proliferating NSCs marked by Proliferating cell nuclear antigen (PCNA). (f-

h) Magnification of the boxed area in panel e, reveals that CLAMP is present in CB cycling neuroblasts

and mother ganglion cells (magenta and cyan arrows, respectively). Scale bars: 50 μm and 25 μm in j.

1071

1072

1073

1074

1075

1076

1077

1078

1079

Figure 2 Lack of CLAMP results in growth deficits both in the body and the nervous system (a, b)

Mutant *clamp* null L3 larvae are considerably smaller in size, compared to control L3s (Mean length ±

SEM; Control: 3.369 ± 0.06031, n=14; *clamp* null: 3.041 ± 0.04371, n=20; ttest p< 0.0001). (c, d) This

reduction is maintained at the CNS level of *clamp* null L3 larvae (Mean normalized brain area ± SEM;

Control: 1.001 ± 0.0485, n=20; *clamp* null: 0.601 ± 0.04052, n=20; ttest p< 0.0001). (e) Western blot for

CLAMP in samples from L3 brain lysates from control and *clamp* null L3 showing the absence of CLAMP

protein (n=2 per condition). Scale bars: 1 mm in a; 50 μm in c.

1087

Figure 3 Lack of CLAMP greatly affects brain development and causes defects in optic lobe

formation. (a, b and e, f) Brain development in the *clamp* null L3 is severely impaired from the early L3

stage, while even at the mid/late L3 stage, *clamp* null brains resemble early L3 control brains. (c, d and g,h)

Full z-stacks of L3 brains showing the distribution of Deadpan (Dpn) -positive NSCs areas, reveals that

NSCs development both in the central brain (CB) and the optic lobe (OL) of the null are impaired, starting

from the early L3 stage and continuing to the mid/late L3 stage. Especially impaired is the OL development

as seen by the reduction in the Dpn+ area indicated with the white dashed outlined areas. EAD: eye antenna

disk. Panels a, b, e and f are 3 optical sections whereas c, d, g and h are full z-stacks throughout the brain.

Scale bars: 50 μm.

Developmental RNA-sequencing

a 16-18 hours embryo

b mid/late L3 Brain

GO Functional Analysis

c DEGs

d **e**

f CLAMP Binding Motif Analysis (FIMO)

Gene	Function
pont	NB attributes
aurA	NB self-renewal; spindle orientation
apc7	Chromosome segregation
esc	NB proliferation
fru	Axonal projections
san	Sister chromatin cohesion
baz	Asymmetric cell division
pcna	Cell cycle dynamics
notch	NB attributes; stem cell division
cdk7	Mitosis regulator
Nelf-E	Transcriptional pausing
sbm	Neural development regulation *via* ecdysone signaling
gcm	Master regulator of gliogenesis
tll	Optic lobe development
gbb	NB proliferation and surface glia survival
D	Regulation of neuronal fate in the OL
Alk	Cell cycle regulation and NB proliferation
wun	Regulates bioactive lipid levels

g IGV Analysis for Clamp Binding

baz

notch

Nelf-E

tll

h Average Profile Analysis

Embryo

L3

1097

Figure 4 CLAMP is implicated in transcriptomic networks that regulate the proliferation and
differentiation of NSCs. (a, b) Experimental workflow for RNA sequencing data generation and DEG
meta-analysis. Embryo panel is scanning electron image of a late stage embryo from(TURNER AND
MAHOWALD 1977). (c) Venn diagram of DEGs in the *clamp* null embryo and L3. (d,e) GO analysis of the
clamp null DEGs revealed high enrichment in categories related to cell cycle dynamics and neural
development. (f) Find Individual Motif Occurrences (FIMO) analysis of the DEGs for the CLAMP binding
motif, revealed that critical regulators of neural development and mitosis are putative targets for CLAMP
binding both at the embryonic and L3 stage, (g) while analysis of embryo ChIP-seq data revealed a direct
regulation of *Tll, Notch, baz* and *Nelf-E* by CLAMP. (h) Average binding profiles for the upregulated
and downregulated DEGs in embryo and L3 *clamp* null. Gene regions were scaled to match
positions of the transcription start sites (TSS) and transcription end sites (TES) with 3k base pairs
shown on either side of the gene. For comparison in each image, the average read depth is also
plotted for 2571 gene regions that show expression in the nervous system but are not differentially
expressed in any of these RNA-seq experiments (NONs). The shaded region around each line is
the standard error of the mean. The genes downregulated in the absence of clamp show higher
average read depth around the TSS, particularly in larvae.

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

Figure 5 Lack of CLAMP impairs neural stem cell proliferation and promotes precocious

differentiation. (a-d) Proliferation of NSCs as seen by EdU labeling, is highly reduced in the *clamp* null

brain, with the most evident impact being on OL NSCs (panels a and c, dashed lines indicate the OL). In

parallel, nuclear Pros+ clusters indicate acquisition of neuronal fate, are more evident in the *clamp* null

brains (panels b and d, arrows). (e) Both the total number of proliferating stem cells (EdU$^+$ Mean ± SEM,

Control: 323.6 ± 4.679; *clamp* null: 50.40 ± 3.174; p<0.0001; n=10) and of proliferating Dpn$^+$ NSCs is

reduced (Dpn$^+$EdU$^+$ Mean ± SEM; Control 102.6 ± 2.141; *clamp* null 11.50 ± 0.3727; n=10). (f) The

reduction in total Dpn+ cells (Dpn$^+$ Mean ± SEM, Control: 191.9 ± 5.313; *clamp* null: 57.90 ± 1.402; p

<0.0001; n = 10), is accompanied by a respective increase in Pros+ cells in the *clamp* null brains (Pros$^+$

Mean ± SEM, Control: 47.1 ± 1.716; *clamp* null: 148.3 ± 2.629; p <0.0001; n = 10). (g, h) Additionally,

despite having an increase in the neuronal marker Elav in the *clamp* null brains, the OL development is

impaired compared to controls (outlines). (i, j) Apoptotic cells marked with DCP1 are more in the OL of

the null compared to a more ubiquitous distribution in the control. CB, central brain; OL, optic lobe; VNC:

ventral nerve cord. Scale bars: a-d, 100 μm; g and h, 50 μm.

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157 **Figure 6 Direct regulation of *Tailless* by CLAMP causes an almost complete absence of its expression**

1158 **in the OL.** (a-c) Analysis of embryo ChIP-seq data reveal that CLAMP directly binds to the *Tailless* gene

1159 in putative promoter areas. (d,e). Expression of the Tailless protein is almost absent in the clamp null optic

1160 (OL) lobe which is severely underdeveloped. Central brain (CB). Scale bars, 50 μm.

1161

Figure 7 Glial-CLAMP surface glia generation and NSC proliferation (a-e) Surface glia (SG) is reduced by ~25% in the *clamp* null brain compared to controls (% SG Repo$^+$ ± SEM, *Control*: 99.98 ± 5.328; *clamp* null: 75.37 ± 1.843; n=6) whereas, (e-i) proliferating NSCs (EdU$^+$ ± SEM, *Control:* 323.6 ± *4.679; clamp* null: 50.40 ± *3.174*) and NSCs (Dpn$^+$ ± SEM, *Control:* 191.9 ± *5.313; clamp* null: 57.90 ± *1.402*) are also reduced in numbers. Glial-specific *clamp* knock-down (*repo>clamp RNAi*) promotes a similar reduction in SG numbers ((% SG Repo$^+$ ± SEM, *repo>clamp RNAi* 131.0 ± 3.162) and promotes a reduction in proliferating NSCs (*repo>clamp RNAi* EdU$^+$ ± SEM*:* 277 ± 6.572; Dpn$^+$ ± SEM: 181 ± 3.964), but to a lesser extend as the one seen in the *clamp* null. Neuronal-specific clamp KD (*elav>clamp RNAi*)

1170 does not affect NSC proliferation and numbers NSCs (*elav>clamp RNAi* EdU$^+$ ± SEM*:* 316. ± 9.519; Dpn$^+$

1171 ± SEM: 217 ± 14.45). Scale bars 100 µm.

1172

1173

1174

Gene Name	log2FC	padj
elav	0.44	0.043
ewg	NA	NA
clamp	-1	6.39E-14
CG12299	-0.70	0.0001
gro	-0.88	2.45E-06
CG8924	-1.12	1.15E-10
JIL-1	NA	NA
osk	-4.76	1.67E-11
ac3	0.54	4.87E-05
cdc16	-0.83	0.005
Ibf1	-0.78	0.0002
Bcl7-like	-0.93	2.92E-11

1189 *Table 1: Functional interactors of clamp as identified by Haussman et al., 2008*

Gene ID	Gene Name	Log2FC	padj	Function
FBgn0044047	Dilp6	-3.028	4.38E-42	OL development/Glia Neuroblast interactions/Insulin Signaling
FBgn0003720	tll	-1.71	1.89E-31	OL development
FBgn0031745	rau	-1.373	3.42E-18	Glial cell development/differentiation of neurons and glia in the developing eye.
FBgn0014179	gcm	-1.281	5.90E-12	Glial cell development-promotes gliogenesis over neurogenesis
FBgn0010433	ato	-1.246	3.88E-15	OL development
FBgn0000529	bsh	-1.155	1.85E-24	OL development
FBgn0019809	gcm2	-1.154	3.97E-20	Has a redundant role with the product of gcm in the differentiation of glial cells and subset of neurons(CHOTARD et al. 2005).
FBgn0031375	erm	-1.116	2.76E-19	Cell fate regulator during neurogenesis.
FBgn0000163	baz	-1.105	6.44E-10	Bazooka forms a complex with par-6 and aPKC. Functions in cell polarization and ACD.
FBgn0021776	mira	-0.99	3.16E-07	Neuroblast proliferation.
FBgn0000137	ase	-0.972	1.54E-05	Type I NB attributes.
FBgn0263512	Vsx2	-0.956	9.10E-23	OL development.
FBgn0010382	CycE	-0.951	0.00131	Cell cycle dynamics.
FBgn0001983	wor	-0.94	3.01E-05	Worniu, neuroblast attributes.
FBgn0000320	eya	-0.909	2.05E-08	OL development /insulin producing cells differentiation.
FBgn0000404	CycA	-0.855	0.010084	Cell cycle progression including G2.
FBgn0025739	pon	-0.847	3.66E-07	Contributes to asymmetric localization of numb and the subsequent suppression of Notch signaling.
FBgn0002973	numb	-0.845	1.67E-15	Acts via Notch signaling, asymmetric division control.
FBgn0005655	PCNA	-0.824	0.00019	Cell cycle dynamics.
FBgn0011674	insc	-0.788	1.44E-06	Spindle orientation; asymmetric divisions.
FBgn0037913	fabp	-0.72	1.85E-06	Expressed in lipid droplet-forming glial cells.
FBgn0013469	klu	-0.713	0.00013	Klumpfuss (klu) is a temporal fate determination.
FBgn0005677	dac	-0.707	4.97E-11	OL development.
FBgn0010109	dpn	-0.665	0.00245	Neuroblast self-renewal -interacts with klu and erm.
FBgn0004647	Notch	-0.642	0.0018	OL development/Stem cell proliferation.
FBgn0000411	Dichaete (D)	-0.634	2.56E-08	Nervous system development and OL development.
FBgn0044324	Chro	-0.61	0.00021	Neuroblast proliferation. Functions downstream of InR/PI3K/TOR signaling and regulates expression of quiescence-inducing Pros to permit NSC reactivation.
FBgn0005427	ewg	-0.533	1.06E-06	Neurogenesis/Synaptogenesis.
FBgn0040080	pins	-0.52	0.008985	Partner of Inscutable; involved in ACD and establishment of cell polarity.

1196　　　　　　　　　　*Table 2: Down-regulated genes in the clamp null L3 brain*

Gene ID	Gene Name	Log2Fc	padj	Function
FBgn0000014	abd-A	2.983	0.00019	Controls the end of neuroblast proliferation.
FBgn0025879	Timp	2.299	9.57E-23	Coordinates synaptogenesis
FBgn0001258	Ldh	2.353	2.43E-22	Lipid metabolism.
FBgn0003117	pnr	2.106	0.008812	Activator of *proneural* achaete-scute complex genes.
FBgn0011746	ana	1.831	3.97E-20	Encodes a secreted glycoprotein that is expressed in glia and at lower levels in neuroblasts, <u>suppresses neuroblast proliferation</u>.
FBgn0032006	Pvr	1.742	8.56E-22	Glia development /neuroblast proliferation.
FBgn0016032	lbm	1.576	3.84E-21	facilitates synapse assembly
FBgn0037989	ATP8B	1.664	2.18E-20	Flippase, translocates phospholipids from one side of a membrane to the other forming new membrane structures, especially vesicles.
FBgn0031461	daw	1.502	2.82E-05	Regulates insulin release from IPC neurons.
FBgn0041087	wun2	1.249	3.17E-20	Encodes a lipid phosphate phosphatase; expressed in astrocytes.
FBgn0086677	jeb	1.224	1.37E-06	Functions as signaling ligand for the product of <u>Alk</u>.
FBgn0066101	LpR1	0.899	3.91E-21	Cellular uptake of neutral lipids from the circulating protein encoded by apolpp.
FBgn0000416	Sap-r	0.889	2.23E-29	Lysosomal Lipid homeostasis.
FBgn0000546	EcR	0.847	6.72E-06	Ecdysone Receptor.
FBgn0040505	Alk	0.692	0.00014	Employs Ras/ERK and PI3K signalling pathways, functions in multiple contexts including synaptic signaling.
FBgn0001297	kay	0.699	7.62E-12	Specifically expressed in glia.
FBgn0039332	alrm	0.635	0.00038	Glial related.
FBgn0000370	crc	0.61	3.83E-17	Coactivator of the product EcR, promotes expression the molting peptide hormone encoded by ETH.
FBgn0016078	wun	0.558	2.85E-07	Lipid phosphate phosphatase.
FBgn0013342	nSyb	0.506	3.24E-05	Synaptic signaling.
FBgn0030574	sbm	0.504	0.002373	L-type amino acid transporter expressed in glia; essential for proper timing of development and brain growth (GALAGOVSKY *et al.* 2018).

1197   *Table 3: UP-regulated genes in the clamp null L3 brain*

1198

1199

1200

1201

1202

1203

1204

1205

 Supplementary figures

Supplementary Figure 1: A single dose of the *Clamp* gene is sufficient to allow for normal brain development. NE, neuroepithelium; OL, optic lobe; CB, central brain. Scale bars 100 μm

1220

1221

1222

1223

1224

1225 **Supplementary Figure 2: Proliferation on neural stem cells is impaired in the absence of CLAMP.**

1226 OL, optic lobe; CB, central brain. Scale bars 100 μm

1227

1228

1229

1230

1231

1232 **Supplementary figure 3: Absence of CLAMP results in increased cell death, especially in the optic**

1233 **lobe's Lobula plug.** Lp, Lobula plug; OL, optic lobe; CB, central brain. Scale bars 100 µm

1234