The landscape of antibody binding to SARS-CoV-2

Anna S. Heffron[1], Sean J. McIlwain[2,3], David A. Baker[1], Maya F. Amjadi[4], Saniya Khullar[2],
Ajay K. Sethi[5], Miriam A. Shelef[4,6], David H. O'Connor[1,7], Irene M. Ong[2,3,8]*

[1]Department of Pathology and Laboratory Medicine, University of Wisconsin-Madison,
Madison, WI, United States of America
[2]Department of Biostatistics and Medical Informatics, University of Wisconsin-Madison,
Madison, WI, United States of America
[3]University of Wisconsin Carbone Comprehensive Cancer Center, University of Wisconsin-
Madison, Madison, WI, United States of America
[4]Department of Medicine, University of Wisconsin-Madison, Madison, WI, United States of
America
[5]Department of Population Health Sciences, University of Wisconsin-Madison, Madison, WI,
United States of America
[6]William S. Middleton Memorial Veterans Hospital, Madison, WI, United States of America
[7]Wisconsin National Primate Research Center, University of Wisconsin-Madison, Madison,
Wisconsin, United States of America
[8]Department of Obstetrics and Gynecology, University of Wisconsin-Madison, Madison, WI,
United States of America

*Corresponding author

Abstract

The search for potential antibody-based diagnostics, vaccines, and therapeutics for pandemic
severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has focused almost exclusively
on the spike (S) and nucleocapsid (N) proteins[1–8]. Coronavirus membrane (M), orf3a, and orf8
proteins are also humoral immunogens in other coronaviruses (CoVs)[8–11] but remain largely
uninvestigated for SARS-CoV-2. Here we show that SARS-CoV-2 infection induces robust
antibody responses to epitopes throughout the SARS-CoV-2 proteome, particularly in M, in
which one epitope achieved near-perfect diagnostic accuracy. We map 79 B cell epitopes
throughout the SARS-CoV-2 proteome and demonstrate that anti-SARS-CoV-2 antibodies
appear to bind homologous peptide sequences in the 6 known human CoVs. Our results
demonstrate previously unknown, highly reactive B cell epitopes throughout the full proteome of
SARS-CoV-2 and other CoV proteins, especially M, which should be considered in diagnostic,
vaccine, and therapeutic development.

Introduction

Antibodies mediate protection from coronaviruses (CoVs) including SARS-CoV-2[1–8], severe
acute respiratory syndrome coronavirus (SARS-CoV)[8,12–15] and Middle Eastern respiratory
syndrome coronavirus (MERS-CoV)[8,16–19]. All CoVs encode 4 main structural proteins, spike
(S), envelope (E), membrane (M), and nucleocapsid (N), as well as multiple non-structural
proteins and accessory proteins[20]. In SARS-CoV-2, anti-S and anti-N antibodies have received
the most attention to date[1–8], including in serology-based diagnostic tests[1–5] and vaccine
candidates[6–8]. However, anti-S antibodies have been linked to antibody dependent enhancement
for SARS-CoV-2 and other CoVs[7,21–26], and prior reports observed that not all individuals
infected with SARS-CoV-2 produce detectable antibodies against S or N[1–5], indicating a need for

expanded antibody-based options. Much less is known about antibody responses to other SARS-
CoV-2 proteins, though data from other CoVs suggest they may be important. Antibodies against
SARS-CoV M can be more potent than antibodies against SARS-CoV S[9–11], and some
experimental SARS-CoV and MERS-CoV vaccines elicit responses to M, E, and orf8[8].
Additionally, previous work has demonstrated humoral cross-reactivity between CoVs[7,14,27–30]
and suggested it could be protective[24,30], although full-proteome cross-reactivity has not been
investigated. We designed a peptide microarray tiling the proteome of SARS-CoV-2 and 8 other
human and animal CoVs in order to assess antibody epitope specificity and potential cross-
reactivity with other CoVs, and we used this microarray to profile IgG antibody responses in 40
COVID-19 convalescent patients and 20 SARS-CoV-2-naive controls.
57
CoV reactivity in uninfected controls
Greater than 90% of adult humans are seropositive for the "common cold" CoVs (CCCoVs:
HCoV-HKU1, HCoV-OC43, HCoV-NL63, and HCoV-229E)[31,32], but it is unknown how these
pre-existing antibodies might affect reactivity to SARS-CoV-2 or other CoVs. We measured IgG
reactivity in sera from 20 SARS-CoV-2-naïve control subjects to CoV linear peptides,
considering reactivity that was >3 standard deviations above the mean for the $\log_2$-quantile
normalized array data to be indicative of antibody binding. All sera exhibited binding in known
epitopes of at least 1 of the control non-CoV strains (poliovirus vaccine and rhinovirus; Fig. 1,
Extended data 1, Extended data 2) and were collected in Wisconsin, USA, where exposure to
SARS-CoV or MERS-CoV was extremely unlikely. We found that at least one epitope in
structural or accessory proteins had binding in 100% of controls for HCoV-HKU1, 85% of
controls for HCoV-OC43, 65% for HCoV-NL63, and 55% for HCoV-229E (Fig. 2, Extended
data 2). Apparent cross-reactive binding was observed in 45% of controls for MERS-CoV, 50%
for SARS-CoV, and 50% for SARS-CoV-2.
72
SARS-CoV-2 proteome humoral profiling
We aimed to map the full extent of binding of antibodies induced by SARS-CoV-2 infection and
to rank the identified epitopes in terms of likelihood of importance and immunodominance. We
defined epitope recognition as antibody binding to contiguous peptides in which the average
$\log_2$-normalized intensity for patients was at least 2-fold greater than for controls with *t*-test
statistics yielding adjusted *p*-values <0.1. We chose these criteria, rather than the 3 standard
deviation cut-off, in order to ensure that binding detected would be greater than background
binding seen in controls (2-fold greater) and to remove regions of binding that were not at least
weakly significantly different from controls (adjusted *p*<0.1). These criteria identified 79 B cell
epitopes (Fig. 3, Extended data 3) in S, M, N, orf1ab, orf3a, orf6, and orf8. We ranked these
epitopes by minimum adjusted *p*-value for any 16-mer in the epitope in order to determine the
greatest likelihood of difference from controls as a proxy for immunodominance. The highest-
ranking epitope occurred in the N-terminus of M (1-M-24). Patient sera showed high-magnitude
reactivity (up to 6.7 fluorescence intensity units) in other epitopes in S, M, N, and orf3a, with
lower-magnitude reactivity (<3.3 fluorescence intensity units) epitopes in other proteins. The
epitopes with the greatest reactivity in S occurred in the fusion peptide (residues 788-806), with
less reactivity in the receptor binding domain (residues 319-541)[6] (Fig. 3). Four detected
epitopes (553-S-26, 624-S-23, 807-S-26, and 1140-S-25) have previously been shown to be
potently neutralizing[33–35], and all 4 of these ranked within the top 10 epitopes. Forty-two of our
detected epitopes (including 1-M-24, 553-S-26, 624-S-23, 807-S-26, and 1140-S-25; Extended

data 3) confirm bioinformatic predictions of antigenicity based on SARS-CoV and MERS-
CoV[7,8,36–38], with all top-ranking epitopes confirming bioinformatic predictions.
95
The highest specificity (100%) and sensitivity (98%), determined by linear discriminant analysis
leave-one-out cross-validation, for any individual peptide was observed for a 16-mer within the
1-M-24 epitope: ITVEELKKLLEQWNLV (Extended data 4). Fifteen additional individual
peptides in M, S, and N had 100% measured specificity and at least 80% sensitivity.
Combinations of 1-M-24 with 1 of 5 other epitopes (384-N-33, 807-S-26, 6057-orf1ab-17, 227-
N-17, 4451-orf1b-16) yielded an area under the curve receiver operating characteristic of 1
(Extended data 5) based on linear discriminant analysis leave-one-out-cross-validation.
103
Human, animal CoV cross-reactivity
We defined cross-reactivity as binding by antibodies in COVID-19 convalescent sera to non-
SARS-CoV-2 peptides at an average $\log_2$-normalized intensity at least 2-fold greater than in
controls with *t*-test statistics yielding adjusted *p*-values <0.1. Antibodies in COVID-19-
convalsecent sera appeared to be cross-reactive with homologous epitopes in S, M, N, orf1ab,
orf3, orf6, and orf8 in other CoVs (Fig. 4, Extended data 6, Extended data 7, Extended data 8).
The greatest number of cross-reactive epitopes (70) were in the RaTG13 bat betacoronavirus (*β*-
CoV), the closest known relative of SARS-CoV-2 (96% nucleotide identity)[39,40], then the
pangolin CoV (51 epitopes, 85% nucleotide identity with SARS-CoV-2)[41], then SARS-CoV (40
epitopes, 78% identity[39]). One region, corresponding to SARS-CoV-2 epitope 807-S-26, was
cross-reactive across all CoVs, and one, corresponding to SARS-CoV-2 epitope 1140-S-25, was
cross-reactive across all *β*-CoVs (Fig. 4). Epitope 807-S-26 includes the CoV S fusion peptide,
and 1140-S-25 is immediately adjacent to the heptad repeat region 2, both of which are involved
in membrane fusion[42].
118
Discussion
M proteins are the most abundant proteins in CoV virions[20]. The N-terminus of M is known in
other CoVs to be a small, glycosylated ectodomain that protrudes outside the virion and interacts
with S, N, and E[20], while the rest of M resides within the viral particle. Full-length SARS-CoV
M has been shown to induce protective antibodies[11,43], and patterns of antibodies binding to
SARS-CoV M are similar to those we found in SARS-CoV-2[36]. SARS-CoV anti-M antibodies
can synergize with anti-S and anti-N antibodies[11,43], and M has been used in protective SARS-
CoV and MERS-CoV vaccines[8]. However, the mechanism of protection of anti-M antibodies
remains unknown, and this protein remains largely understudied and underutilized as an antigen.
Other groups have not previously identified the high magnitude binding we observed in M,
though that may be due to using earlier sample timepoints or different techniques or
algorithms[44,45]. Our results, in concert with prior knowledge of anti-SARS-CoV antibodies,
strongly suggest that M, particularly the 1-M-24 epitope, as well as other novel epitopes that we
identified should be investigated further as potential targets in SARS-CoV-2 diagnostics,
vaccines, and therapeutics.
134
We also found that antibodies produced in response to SARS-CoV-2 infection appeared to cross-
react with homologous epitopes throughout the proteomes of other human and non-human CoVs.
Hundreds of CoVs have been discovered in bats and other species[24,39–41,46,47], making future
spillovers inevitable. The broad cross-reactivity we observed in some homologous peptide

sequences may help guide the development of pan-CoV vaccines[18], especially given that
antibodies binding to 807-S-26 and 1140-S-25, epitope motifs cross-reactive across all CoVs and
all β-CoVs, respectively, are known to be potently neutralizing[33,34]. We cannot determine
whether the increased IgG binding to CCCoVs in COVID-19 convalescent sera is due to newly
developed cross-reactive antibodies or the stimulation of a memory response against the original
CCCoV antigens. However, cross-reactivity of anti-SARS-CoV-2 antibodies with SARS-CoV or
MERS-CoV is likely real, since our population was very unlikely to have been exposed to those
viruses. A more stringent assessment of cross-reactivity as well as functional investigations into
these cross-reactive antibodies will be vital in determining their capacity for cross-protection.
Further, our methods efficiently detect antibody binding to linear epitopes[48], but their sensitivity
for detecting parts of conformational epitopes is unknown, and additional analyses will be
required to determine whether epitopes identified induce neutralizing or otherwise protective
antibodies.
152
153 Many questions remain regarding the biology and immunology of SARS-CoV-2. Our extensive
154 profiling of epitope-level resolution antibody reactivity in COVID-19 convalescent subjects
155 provides new epitopes that could serve as important targets in the development of improved
156 diagnostics, vaccines, and therapeutics against SARS-CoV-2 and dangerous human CoVs that
157 may emerge in the future.
158

 Figures

Figure 1. Patients and control subjects show reactivity to a poliovirus control. Sera from 20
control subjects collected before 2019 were assayed for IgG binding to the full proteome of
human poliovirus 1 on a peptide microarray. Binding was measured as reactivity that was >3
standard deviations above the mean for the $\log_2$-quantile normalized array data. Patients and
controls alike showed reactivity to a well-documented linear poliovirus epitope (start position
613 [IEDB.org]; orange shading in line plot).

Figure 2. Control sera show reactivity frequently to CCCoVs and rarely to SARS-CoV,
MERS-CoV, and SARS-CoV-2. Sera from 20 control subjects collected before 2019 were
assayed for IgG binding to the full proteomes of 9 CoVs on a peptide microarray. Viral proteins
are shown aligned to the SARS-CoV-2 proteome with each virus having an individual panel;
SARS-CoV-2 amino acid (aa) position is represented on the x-axis. Binding was measured as

 reactivity that was >3 standard deviations above the mean for the $\log_2$-quantile normalized array
176 data.

178
Figure 3. Anti-SARS-CoV-2 antibodies bind throughout the viral proteome. Sera from 40
COVID-19 convalescent subjects were assayed for IgG binding to the full SARS-CoV-2
proteome on a peptide microarray. B cell epitopes were defined as peptides in which patients'
average $\log_2$-normalized intensity (black lines in line plots) is 2-fold greater than controls' (gray
lines in line plots) and *t*-test statistics yield adjusted *p*-values < 0.1; epitopes are identified by
orange shading in the line plots.

185
186

Figure 4. Anti-SARS-CoV-2 antibodies cross-react with other CoVs. Sera from 40 COVID-19 convalescent patients were assayed for IgG binding to 9 CoVs on a peptide microarray; averages for all 40 are shown. Viral proteins are aligned to the SARS-CoV-2 proteome; SARS-CoV-2 amino acid (aa) position is represented on the x-axis. Regions cross-reactive across all β-CoVs (*) or cross-reactive for SARS-CoV or MERS-CoV (#) are indicated. Gray shading indicates gaps due to alignment or lacking homologous proteins. Cross-reactive binding is defined as peptides in which patients' average $\log_2$-normalized intensity is 2-fold greater than controls' and t-test statistics yield adjusted p-values < 0.1.

Extended data

Extended data 1. All 40 COVID-19 convalescent patients and all 20 naïve controls reacted to known epitopes in at least one control virus (rhinovirus and poliovirus strains).

Extended data 2. Percentages of the 40 COVID-19 convalescent patients and 20 naïve controls reacted to known epitopes in at least one control virus (rhinovirus and poliovirus strains).

Extended data 3. B cell epitopes in the SARS-CoV-2 proteome identified by antibody binding in 40 COVID-19 convalescent patients compared to 20 naïve controls.

209 **Extended data 4.** Specificity and sensitivity for past SARS-CoV-2 infection in 40 COVID-19
210 convalescent patients compared to 20 naïve controls of individual 16-mer peptides comprising
211 epitopes throughout the full SARS-CoV-2 proteome.

212
213 **Extended data 5.** Epitopes paired with the 1-M-24 epitope obtained an area under the receiver
214 operating characteristic curve (AUC-ROC) of 1.0 for SARS-CoV-2 infection in 40 COVID-19
215 convalescent patients and 20 naïve controls using leave-one-out cross validation with linear
216 discriminant analysis.

217
218 **Extended data 6.** Alignment of epitopes in human and animal CoVs for which antibodies in sera
219 from 40 COVID-19 convalescent patients showed apparent cross-reactive binding. Alignments
220 were performed in Geneious Prime 2020.1.2 (Auckland, New Zealand).

221
222 **Extended data 7.** Cross-reactive binding of antibodies against other CoVs in 40 COVID-19
223 convalescent patients compared to 20 naïve controls.

224
225 **Extended data 8.** Cross-reactive binding of antibodies in 40 COVID-19 convalescent patients
226 compared to 20 naïve controls in protein motifs in other CoVs aligned to SARS-CoV-2.

227
228 **Extended data 9.** B cell epitopes in the SARS-CoV-2 proteome identified by antibody binding
229 in 40 COVID-19 convalescent patients compared to 20 naïve controls were differentiated using a
230 cut-off of at least a 2-fold greater magnitude reactivity in patients vs controls and *t*-test statistics
231 yielding adjusted *p*-values <0.1. Degrees of freedom, fold change, and standard deviation for
232 each peptide are given.

233
234
235 **Methods**
236
Peptide microarray design and synthesis
Viral protein sequences were selected and submitted to Nimble Therapeutics (Madison, WI,
USA) for development into a peptide microarray[48]. Sequences represented include proteomes of
all seven coronaviruses known to infect humans, proteomes of closely related coronaviruses
found in bats and pangolins, and spike proteins from other coronaviruses (accession numbers and
replicates per peptide shown in **Supplementary Table 1**). A number of proteins were included
as controls, including poliovirus, seven strains of human rhinovirus, and human cytomegalovirus
65kDa phosphoprotein. We chose these controls given that we expect most human adults will
have antibody reactivity to at least one of these proteins and proteomes. Accession numbers used
to represent each viral protein are listed in the supplemental material (accession numbers and
replicates per peptide shown in **Supplementary Table 1**). All proteins were tiled as 16 amino
acid peptides overlapping by 15 amino acids. All unique peptides were tiled in a lawn of
thousands of copies, with each unique peptide represented in at least 3 and up to 5 replicates
(**Supplementary Table 1**). The peptide sequences were synthesized in situ with a Nimble
Therapeutics Maskless Array Synthesizer (MAS) by light-directed solid-phase peptide synthesis
using an amino-functionalized support (Geiner Bio-One) coupled with a 6-aminohexanoic acid
linker and amino acid derivatives carrying a photosensitive 2-(2-nitrophenyl) propyloxycarbonyl
(NPPOC) protection group (Orgentis Chemicals). Unique peptides were synthesized in random

positions on the array to minimize impact of positional bias. Each array consists of twelve
subarrays, where each subarray can process one sample and each subarray contains up to
389,000 unique peptide sequences.
258
259 **Supplementary Table 1. Proteins represented on the peptide microarray**

	Protein(s)	GenBank accession number(s)	Number of replicates of each unique peptide
Coronavirus proteins	Severe acute respiratory syndrome coronavirus 2 proteome	NC_045512.2	4-5
	Severe acute respiratory syndrome coronavirus proteome	NC_004718.3	3
	Middle Eastern respiratory syndrome coronavirus proteome	NC_019843.3	3
	Human coronavirus HKU1 proteome	NC_006577.2	3
	Human coronavirus OC43 proteome	NC_006213.1	3
	Human coronavirus 229E proteome	NC_002645.1	3
	Human coronavirus NL63 proteome	NC_005831.2	3
	Bat coronavirus (RaTG13 isolate) proteome	MN996532.1	3
	Pangolin coronavirus proteome	MT072864.1	3
Control proteins	Human rhinovirus A1 polyprotein	NC_038311.1	3
	Human rhinovirus A7 polyprotein	DQ473503.1	3
	Human rhinovirus A16 polyprotein	L24917.1	3
	Human rhinovirus A36 polyprotein	JX074050.1	3

Human rhinovirus C2 polyprotein	EF077280.1	3
Human rhinovirus C15 polyprotein	GU219984.1	3
Human rhinovirus C41 polyprotein	KY189321.1	3
Human poliovirus 1 polyprotein	ANA67904.1	3
Human cytomegalovirus 65 kDa phosphoprotein	P06725.2	3

Human subjects

The study was conducted in accordance with the Declaration of Helsinki and approved by the Institutional Review Board of the University of Wisconsin-Madison. Clinical data and sera from subjects infected with SARS-CoV-2 were obtained from the University of Wisconsin (UW) COVID-19 Convalescent Biobank and from control subjects (sera collected prior to 2019) from the UW Rheumatology Biobank[49]. All subjects were 18 years of age or older at the time of recruitment and provided informed consent. COVID-19 convalescent subjects had a positive SARS-COV-2 PCR test at UW Health with sera collected 5-6 weeks after self-reported COVID-19 symptom resolution except blood was collected for one subject after 9 weeks. Age, sex, medications, and medical problems were abstracted from UW Health's electronic medical record (EMR). Race and ethnicity were self-reported. Hospitalization and intubation for COVID-19 and smoking status at the time of blood collection (controls) or COVID-19 were obtained by EMR abstraction and self-report and were in complete agreement. Two thirds of COVID-19 convalescent subjects and all controls had a primary care appointment at UW Health within 2 years of the blood draw as an indicator of the completeness of the medical information. Subjects were considered to have an immunocompromising condition if they met any of the following criteria: immunosuppressing medications, systemic inflammatory or autoimmune disease, cancer not in remission, uncontrolled diabetes (secondary manifestations or hemoglobin A1c $\geq 7.0\%$), or congenital or acquired immunodeficiency. Control and COVID-19 subjects were similar in regard to demographics and health (**Supplementary Table 2**). No subjects were current smokers.

Supplementary Table 2. Characteristics of COVID-19 Convalescent and Control Subjects

	COVID-19 (n=40)	Control (n=20)	*p*
Age, median (IQR) years	51 (19-83)	55 (22-83)	0.378
Sex, number female (%)	17 (42.5)	11 (55.0)	0.360

Race, number (%)			0.866
White	34 (85.0)	18 (90.0)	
Black	3 (7.5)	1 (5.0)	
Asian	3 (7.5)	1 (5.0)	
Native American	0 (0.0)	0 (0.0)	
Pacific Islander	0 (0.0)	0 (0.0)	
Ethnicity, number Hispanic (%)	5 (12.5)	1 (5.0)	0.361
Charlson comorbidity score, median (IQR)	2 (0, 3)	2 (0.5, 4)	0.551
Immunocompromised, number (%)	9 (22.5)	7 (35.0)	0.302
COVID-19 disease severity, number (%)			
Hospitalized and intubated	8 (20.0)	-	-
Hospitalized without intubation	7 (17.5)	-	-
Not hospitalized	25 (62.5)	-	-

284
285
286
287

Peptide array sample binding
Samples were diluted 1:100 in binding buffer (0.01M Tris-Cl, pH 7.4, 1% alkali-soluble casein,
0.05% Tween-20) and bound to arrays overnight at 4°C. After sample binding, the arrays were
washed 3× in wash buffer (1× TBS, 0.05% Tween-20), 10 min per wash. Primary sample
binding was detected via Alexa Fluor® 647-conjugated goat anti-human IgG secondary antibody
(Jackson ImmunoResearch). The secondary antibody was diluted 1:10,000 (final concentration
0.1 ng/µl) in secondary binding buffer (1x TBS, 1% alkali-soluble casein, 0.05% Tween-20).
Arrays were incubated with secondary antibody for 3 h at room temperature, then washed 3× in
wash buffer (10 min per wash), washed for 30 sec in reagent-grade water, and then dried by
spinning in a microcentrifuge equipped with an array holder. Fluorescent signal of the secondary
antibody was detected by scanning at 635 nm at 2 µm resolution using an Innopsys 910AL

microarray scanner. Scanned array images were analyzed with proprietary Nimble Therapeutics
software to extract fluorescence intensity values for each peptide.

Peptide microarray findings validation
We included sequences on the array of viruses which we expected all adult humans to be likely
to have been exposed to as positive controls: one poliovirus strain (measuring vaccine exposure),
and seven rhinovirus strains. Any subject whose sera did not react to at least one positive control
would be considered a failed run and removed from the analysis. All subjects in this analysis
reacted to epitopes in at least one control strain (Fig. 1, Extended data 1, Extended data 2).

Peptide microarray data analysis and data availability
The raw fluorescence signal intensity values were $\log_2$ transformed. Clusters of fluorescence
intensity of statistically unlikely magnitude, indicating array defects, were identified and
removed. Local and large area spatial corrections were applied, and the median transformed
intensity of the peptide replicates was determined. The resulting median data was cross-
normalized using quantile normalization. All peptide microarray datasets and code used in these
analyses can be downloaded from https://github.com/Ong-Research/Ong_UW_Adult_Covid-
19.git.

Statistical analysis
Statistical analyses were performed in R (v 4.0.2) using in-house scripts. For each peptide, a p-
value from a two-sided t-test with unequal variance between sets of patient and control
responses were calculated and adjusted using the Benjamini-Hochberg (BH) algorithm. To
determine whether the peptide was in an epitope (in SARS-CoV-2 proteins) or cross-reactive for
anti-SARS-CoV-2 antibodies (in non-SARS-CoV-2 proteins), we used an adjusted p-value cutoff
of <0.1 (based on multiple hypothesis testing correction for all 119,487 unique sequences on the
array) and a fold-change of greater than or equal to 2 and grouped consecutive peptides as a
represented epitope. Linear discriminant analysis leave-one-out cross validation was used to
determine specificity and sensitivity on each peptide and from each epitope using the average
signal of the component peptides.

To identify cross reactive epitopes, we used each SARS-CoV-2 epitope sequence as a query,
searched the database of proteins from the sequences in the peptide array using blastp (-word-
size 2, num-targets 4000) to find homologous sequences in the bat, pangolin, and other human
CoV strains, then determined whether the average $\log_2$-normalized intensity for these sequences
in patients was at least 2-fold greater than in controls with t-test statistics yielding adjusted p-
values <0.1. Each blast hit was then mapped back to the corresponding probe ranges.

The clinical and demographic characteristics of convalescent subjects were compared to those of
the controls using χ^2 tests for categorical variables and Wilcoxon rank-sum tests for non-
normally distributed continuous measures.

Heatmaps were created using the gridtext[50] and complexheatmap[51] packages in R. Alignments
for heatmaps were created using MUSCLE[52].

345
Funding
I.M.O. acknowledges support by the Clinical and Translational Science Award (CTSA) program, through the NIH National Center for Advancing Translational Sciences (NCATS), grants UL1TR002373 and KL2TR002374. This research was also supported by 2U19AI104317-06 (to I.M.O) and R24OD017850 (to D.H.O.) from the National Institute of Allergy and Infectious Diseases of the National Institutes of Health (www.niaid.nih.gov). A.S.H. has been supported by NRSA award T32 AI007414 and M.F.A. by T32 AG000213. S.J.M. acknowledges support by the National Cancer Institute, NIH and UW Carbone Comprehensive Cancer Center's Cancer Informatics Shared Resource (grant P30-CA-14520). This project was also funded through a COVID-19 Response Grant from the Wisconsin Partnership Program and the University of Wisconsin School of Medicine and Public Health (to M.A.S.), startup funds through the University of Wisconsin Department of Obstetrics and Gynecology (I.M.O.), and the Data Science Initiative grant from the University of Wisconsin-Madison Office of the Chancellor and the Vice Chancellor for Research and Graduate Education (with funding from the Wisconsin Alumni Research Foundation) (I.M.O.).

Acknowledgments
The authors are grateful to Dr. Christina Newman, Dr. Nathan Sherer, Dr. Thomas Friedrich, Dr. Amelia Haj, Dr. James Gern, Dr. Christine Seroogy, and Gage Moreno for their thoughtful comments and helpful discussions in preparing this manuscript.

Author contributions
A.S.H., S.J.M., D.A.B., M.F.A., M.A.S., D.H.O. and I.M.O. conceptualized this study. A.S.H., D.A.B., and I.M.O. created the array design. M.F.A. and M.A.S. selected patient and control samples. A.S.H., M.F.A., and M.A.S. collected patient demographics and characteristics by medical record chart review. A.S.H., S.J.M., D.A.B., S.K., and I.M.O. performed data normalizations, analyses, and validations, and created graphical data visualizations. A.S.H., S.J.M., D.A.B., A.K.S., and I.M.O. performed formal statistical analyses. S.J.M., D.A.B., and I.M.O. wrote the custom software scripts used. A.S.H. wrote the original draft of the manuscript with input from D.H.O. and I.M.O.. A.S.H., S.J.M., D.A.B., M.F.A., and I.M.O. wrote sections of the methods. All authors contributed to review and editing.

The authors declare the following competing interests: A.S.H., S.J.M., D.A.B., M.F.A., S.K., M.A.S., D.H.O., and I.M.O are listed as the inventors on a patent filed that is related to findings in this study. Application: 63/080568, 63/083671. Title: IDENTIFICATION OF SARS-COV-2 EPITOPES DISCRIMINATING COVID-19 INFECTION FROM CONTROL AND METHODS OF USE. Application type: Provisional. Status: Filed. Country: United States. Filing date: September 18, 2020, September 25, 2020.

References:
1. Deeks, J. J. et al. Antibody tests for identification of current and past infection with SARS-CoV-2. *Cochrane Database Syst Rev* **6**, CD013652 (2020).

2.   Liu, W. et al. Evaluation of Nucleocapsid and Spike Protein-Based Enzyme-Linked
Immunosorbent Assays for Detecting Antibodies against SARS-CoV-2. *J Clin Microbiol*
58, (2020).
3.   Tré-Hardy, M. et al. Analytical and clinical validation of an ELISA for specific SARS-
CoV-2 IgG, IgA, and IgM antibodies. *J Med Virol* (2020).
4.   Lisboa Bastos, M. et al. Diagnostic accuracy of serological tests for covid-19: systematic
review and meta-analysis. *BMJ* **370**, m2516 (2020).
5.   Ayouba, A. et al. Multiplex detection and dynamics of IgG antibodies to SARS-CoV2 and
the highly pathogenic human coronaviruses SARS-CoV and MERS-CoV. *J Clin Virol* **129**,
104521 (2020).
6.   Huang, Y., Yang, C., Xu, X. F., Xu, W. & Liu, S. W. Structural and functional properties
of SARS-CoV-2 spike protein: potential antivirus drug development for COVID-19. *Acta*
Pharmacol Sin **41**, 1141-1149 (2020).
7.   Chen, W. Promise and challenges in the development of COVID-19 vaccines. *Hum Vaccin*
Immunother 1-5 (2020).
8.   Ong, E., Wong, M. U., Huffman, A. & He, Y. COVID-19 Coronavirus Vaccine Design
Using Reverse Vaccinology and Machine Learning. *Front Immunol* **11**, 1581 (2020).
9.   Chow, S. C. et al. Specific epitopes of the structural and hypothetical proteins elicit
variable humoral responses in SARS patients. *J Clin Pathol* **59**, 468-476 (2006).
10. He, Y., Zhou, Y., Siddiqui, P., Niu, J. & Jiang, S. Identification of immunodominant
epitopes on the membrane protein of the severe acute respiratory syndrome-associated
coronavirus. *J Clin Microbiol* **43**, 3718-3726 (2005).
11. Pang, H. et al. Protective humoral responses to severe acute respiratory syndrome-
associated coronavirus: implications for the design of an effective protein-based vaccine. *J*
Gen Virol **85**, 3109-3113 (2004).
12. Sui, J. et al. Potent neutralization of severe acute respiratory syndrome (SARS) coronavirus
by a human mAb to S1 protein that blocks receptor association. *Proc Natl Acad Sci U S A*
101, 2536-2541 (2004).
13. ter Meulen, J. et al. Human monoclonal antibody as prophylaxis for SARS coronavirus
infection in ferrets. *Lancet* **363**, 2139-2141 (2004).
14. Rockx, B. et al. Structural basis for potent cross-neutralizing human monoclonal antibody
protection against lethal human and zoonotic severe acute respiratory syndrome
coronavirus challenge. *J Virol* **82**, 3220-3235 (2008).
15. Martin, J. E. et al. A SARS DNA vaccine induces neutralizing antibody and cellular
immune responses in healthy adults in a Phase I clinical trial. *Vaccine* **26**, 6338-6343
(2008).
16. Jiang, L. et al. Potent neutralization of MERS-CoV by human neutralizing monoclonal
antibodies to the viral spike glycoprotein. *Sci Transl Med* **6**, 234ra59 (2014).
17. Chen, Z. et al. Human Neutralizing Monoclonal Antibody Inhibition of Middle East
Respiratory Syndrome Coronavirus Replication in the Common Marmoset. *J Infect Dis*
215, 1807-1815 (2017).
18. Burton, D. R. & Walker, L. M. Rational Vaccine Design in the Time of COVID-19. *Cell*
Host Microbe **27**, 695-698 (2020).
19. Zhou, Y., Jiang, S. & Du, L. Prospects for a MERS-CoV spike vaccine. *Expert Rev*
Vaccines **17**, 677-686 (2018).

20. Maier, H. J., Bickerton, E. & Britton, P. *Coronaviruses : methods and protocols* (Humana
Press; Springer, New York, 2015).
21. Liu, L. et al. Anti-spike IgG causes severe acute lung injury by skewing macrophage
responses during acute SARS-CoV infection. *JCI Insight* **4**, (2019).
22. Lee, W. S., Wheatley, A. K., Kent, S. J. & DeKosky, B. J. Antibody-dependent
enhancement and SARS-CoV-2 vaccines and therapies. *Nat Microbiol* (2020).
23. Takano, T., Kawakami, C., Yamada, S., Satoh, R. & Hohdatsu, T. Antibody-dependent
enhancement occurs upon re-infection with the identical serotype virus in feline infectious
peritonitis virus infection. *J Vet Med Sci* **70**, 1315-1321 (2008).
24. Morens, D. M. & Fauci, A. S. Emerging Pandemic Diseases: How We Got to COVID-19.
Cell **182**, 1077-1092 (2020).
25. Tseng, C. T. et al. Immunization with SARS coronavirus vaccines leads to pulmonary
immunopathology on challenge with the SARS virus. *PLoS One* **7**, e35421 (2012).
26. Jaume, M. et al. SARS CoV subunit vaccine: antibody-mediated neutralisation and
enhancement. *Hong Kong Med J* **18 Suppl 2**, 31-36 (2012).
27. Tian, X. et al. Potent binding of 2019 novel coronavirus spike protein by a SARS
coronavirus-specific human monoclonal antibody. *Emerg Microbes Infect* **9**, 382-385
(2020).
28. Lv, H. et al. Cross-reactive Antibody Response between SARS-CoV-2 and SARS-CoV
Infections. *Cell Rep* **31**, 107725 (2020).
29. Pinto, D. et al. Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
antibody. *Nature* **583**, 290-295 (2020).
30. Nickbakhsh, S. et al. Epidemiology of Seasonal Coronaviruses: Establishing the Context
for the Emergence of Coronavirus Disease 2019. *J Infect Dis* **222**, 17-25 (2020).
31. Gorse, G. J., Patel, G. B., Vitale, J. N. & O'Connor, T. Z. Prevalence of antibodies to four
human coronaviruses is lower in nasal secretions than in serum. *Clin Vaccine Immunol* **17**,
1875-1880 (2010).
32. Premkumar, L. et al. The receptor binding domain of the viral spike protein is an
immunodominant and highly specific target of antibodies in SARS-CoV-2 patients. *Sci
Immunol* **5**, (2020).
33. Poh, C. M. et al. Two linear epitopes on the SARS-CoV-2 spike protein that elicit
neutralising antibodies in COVID-19 patients. *Nat Commun* **11**, 2806 (2020).
34. Zhang, B. Z. et al. Mining of epitopes on spike protein of SARS-CoV-2 from COVID-19
patients. *Cell Res* **30**, 702-704 (2020).
35. Li, Y. et al. Linear epitopes of SARS-CoV-2 spike protein elicit neutralizing antibodies in
COVID-19 patients. *Cell Mol Immunol* **17**, 1095-1097 (2020).
36. Grifoni, A. et al. A Sequence Homology and Bioinformatic Approach Can Predict
Candidate Targets for Immune Responses to SARS-CoV-2. *Cell Host Microbe* **27**, 671-
680.e2 (2020).
37. Ahmed, S. F., Quadeer, A. A. & McKay, M. R. Preliminary Identification of Potential
Vaccine Targets for the COVID-19 Coronavirus (SARS-CoV-2) Based on SARS-CoV
Immunological Studies. *Viruses* **12**, (2020).
38. Crooke, S. N., Ovsyannikova, I. G., Kennedy, R. B. & Poland, G. A. Immunoinformatic
identification of B cell and T cell epitopes in the SARS-CoV-2 proteome. *Sci Rep* **10**,
14179 (2020).

39.   Zhou, P. et al. A pneumonia outbreak associated with a new coronavirus of probable bat
origin. *Nature* **579**, 270-273 (2020).

40.   Ge, X. Y. et al. Coexistence of multiple coronaviruses in several bat colonies in an
abandoned mineshaft. *Virol Sin* **31**, 31-40 (2016).

41.   Xiao, K. et al. Isolation of SARS-CoV-2-related coronavirus from Malayan pangolins.
Nature **583**, 286-289 (2020).

42.   Tang, T., Bidon, M., Jaimes, J. A., Whittaker, G. R. & Daniel, S. Coronavirus membrane
fusion mechanism offers a potential target for antiviral development. *Antiviral Res* **178**,
104792 (2020).

43.   Shi, S. Q. et al. The expression of membrane protein augments the specific responses
induced by SARS-CoV nucleocapsid DNA immunization. *Mol Immunol* **43**, 1791-1798
(2006).

44.   Shrock, E. et al. Viral epitope profiling of COVID-19 patients reveals cross-reactivity and
correlates of severity. *Science* (2020).

45.   Mishra, N. et al. Immunoreactive peptide maps of SARS-CoV-2 and other human
coronaviruses. *bioRxiv*
https://www.biorxiv.org/content/10.1101/2020.08.13.249953v1.full.pdf (2020).

46.   Woo, P. C., Lau, S. K., Huang, Y. & Yuen, K. Y. Coronavirus diversity, phylogeny and
interspecies jumping. *Exp Biol Med (Maywood)* **234**, 1117-1127 (2009).

47.   Anthony, S. J. et al. Global patterns in coronavirus diversity. *Virus Evol* **3**, vex012 (2017).

48.   Heffron, A. S. et al. Antibody responses to Zika virus proteins in pregnant and non-
pregnant macaques. *PLoS Negl Trop Dis* **12**, e0006903 (2018).

49.   Holmes, C. L. et al. Reduced IgG titers against pertussis in rheumatoid arthritis: Evidence
for a citrulline-biased immune response and medication effects. *PLoS One* **14**, e0217221
(2019).

50.   Wilke, C. O. gridtext: Improved Text Rendering Support for 'Grid' Graphics. R package
version 0.1.1. (2020).

51.   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847-2849 (2016).

52.   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res* **32**, 1792-1797 (2004).