Title page

Characterizing the spatial signal of environmental DNA in river systems using a community

ecology approach

Isabel Cantera[1]*, Jean-Baptiste Decotte[3], Tony Dejean[3,4], Jérôme Murienne[1,] Régis Vigouroux[5],

Alice Valentini[4] and Sébastien Brosse[1]

[1]Laboratoire Evolution et Diversité Biologique, UMR5174, CNRS, IRD, Université Toulouse III

Paul Sabatier, 118 route de Narbonne, F-31062 Toulouse, France.

isa_cantera@hotmail.com, jerome.murienne@univ-tlse3.fr, sebastien.brosse@univ-tlse3.fr

[3]VIGILIFE, 17 rue du Lac Saint-André Savoie Technolac - BP 274, Le Bourget-du-Lac 73375,

France. jb.decotte@vigilife.org

[4]SPYGEN, 17 rue du Lac Saint-André Savoie Technolac - BP 274, Le Bourget-du-Lac 73375,

France. tony.dejean@spygen.com, alice.valentini@spygen.com

[5]HYDRECO, Laboratoire Environnement de Petit Saut, B.P 823, F-97388, Kourou Cedex,

French Guiana. regis.vigouroux@hydrecolab.com

Abstract

Environmental DNA (eDNA) is gaining a growing popularity among scientists but its

applicability to biodiversity research and management remains limited in river systems by the

lack of knowledge about the spatial extent of the downstream transport of eDNA.

Here, we assessed the ability of eDNA inventories to retrieve spatial patterns of fish assemblages

along two large and species rich Neotropical rivers. We first examined overall community

variation with distance through the distance decay of similarity and compared this pattern to

capture-based samples. We then considered previous knowledge on individual species

distributions, and compared it to the eDNA inventories for a set of 53 species.

eDNA collected from 28 sites in the Maroni and 25 sites in the Oyapock rivers permitted to

retrieve a decline of species similarity with distance between sites. The distance decay of

similarity derived from eDNA was similar, and even more pronounced, than that obtained with

capture-based methods (gil-nets). In addition, the species upstream-downstream distribution

range derived from eDNA matched to the known distribution of most species.

Our results demonstrate that environmental DNA does not represent an integrative measure of

biodiversity across the whole upstream river basin but provide a relevant picture of local fish

assemblages. Importantly, the spatial signal gathered from eDNA was therefore comparable to

that gathered with local capture based methods, which describes fish fauna over a few hundred

metres.

Keywords: Neotropical fishes, eDNA, spatial signal, distance decay, distribution range,

freshwater.

Introduction

In aquatic ecosystems, DNA molecules flowing in the water after being separated from their source organisms can be assigned to taxa using the environmental DNA (eDNA) metabarcoding approach. The method consists on the sampling, extraction, amplification and sequencing of free DNA suspended in the water or linked to suspended particles. Subsequently, the obtained sequences are taxonomically assigned to species based on comparisons to a reference molecular database (Taberlet, Bonin, Zinger, & Coissac, 2018). This approach has proven to be particularly efficient at measuring the diversity of fishes in aquatic ecosystems (Civade et al., 2016; Hänfling et al., 2016; Olds et al., 2016; Valentini et al., 2016; Evans et al., 2017; Pont et al., 2018; Cantera et al., 2019). Indeed, in those studies, the eDNA samples provided similar or more complete inventories than traditional survey methods, such as netting, electrofishing or visual records, at the same sites. The great detection performance of the method may be due to its high sensitivity to detect species that are rare and difficult to sample with traditional methods (Ficetola, Miaud, Pompanon, & Taberlet, 2008; Goldberg, Pilliod, Arkle, & Waits, 2011; Jerde, Mahon, Chadderton, & Lodge, 2011). In fact, the species list obtained with one eDNA sampling session was often comparable to cumulative traditional sampling sessions and historical records (Civade *et al.* 2016; Hänfling *et al.* 2016; Nakagawa *et al.* 2018; Pont *et al.* 2018; Cantera et al. 2019). Due to its sampling efficiency to reconstruct whole aquatic communities, eDNA methods are expected to revolutionize survey methods for ecological research and biodiversity monitoring on aquatic ecosystems (Keck, Vasselon, Tapolczai, Rimet, & Bouchez, 2017; Rees, Gough, Middleditch, Patmore, & Maddison, 2015; Rees, Maddison, Middleditch, Patmore, & Gough, 2014; Zinger et al., 2020). This is particularly true for large rivers where sampling the fauna can

be very difficult, as it is time consuming, costly and limited to specific habitats (Pont et al.,

2018).

The application of eDNA has been however limited in rivers because the species

detection recovered by eDNA sampling might be decoupled from the species' physical habitat

due to downstream eDNA transport. The network structure of rivers is governed by a continuous

and directional flow (McCluney et al., 2014) that can transport eDNA downstream from the DNA

source while it degrades (Barnes & Turner, 2016). The spatial extent of downstream transport

depends on the degradation rate of eDNA (Barnes & Turner, 2016) combined with the discharge

rates of the water system (Jane et al., 2015). For instance, fast-flow systems may transport eDNA

downstream over large distances before its degradation and dilution. Moreover, the identity and

density of target species could also influence eDNA detection distance (Deiner & Altermatt,

2014; Pilliod, Goldberg, Arkle, & Waits, 2014; Pont et al., 2018).

In aquaria and mesocosms, eDNA remained detectable from a few days to a few weeks

after its release in the water (see Table 1, Barnes and Turner 2016). Besides being heterogeneous,

these experimental results are difficult to relate with water velocity in the field to estimate eDNA

detection distance, given that natural ecosystems are more complex and local environmental

conditions, which influence eDNA degradation rate (Barnes & Turner, 2016), are covarying. For

instance, Takahara *et al.* (2012) found a significant relationship between eDNA production from

common carp and water temperature in a natural lagoon but not in aquaria. Moreover, in

experimental designs the DNA of target species are usually in higher concentrations compared to

natural ecosystems while DNA concentration was shown to influence degradation rate (Barnes &

Turner, 2016).

Deiner *et al.* (2016) advocated that eDNA-based inventories represent a spatially

integrated measure of biodiversity describing the diversity at the catchment scale and considered

that rivers act as conveyor belts for eDNA. Therefore, eDNA collected at one sampling point may

represent the combination of local and upstream diversity from this point. Indeed, the assumption

that eDNA of entire communities will accumulate downstream omits the phenomenon of DNA

degradation and sedimentation once it has been shed from an organism (Barnes and Turner 2016;

Pont et al. 2018). The detection distance of eDNA was investigated more explicitly in natural

rivers and streams in temperate ecosystems. Although the studies agreed that eDNA from a

species can be detected downstream from where the species occurs and that eDNA signal decays

over distance, the estimation of detection distance displays a great variability among studies. The

introduction of caged animals in streams revealed that eDNA was detected 5 m but not 50 m

downstream from caged Idaho giant salamander (Pilliod et al., 2014) and up to 1 km downstream

from caged brook trout (Wilcox et al., 2016). Contrastingly, eDNA from lake species has been

found in downstream streams over greater distances. DNA from a lacustrine fish species was

detected up to 3 km downstream from the outlet of a lake by Civade *et al.* (2016), and Deiner and

Altermatt (2014) detected two lake invertebrate species (a planktonic crustacean and a mollusc)

up to 12.3 km downstream from a lake. Nakagawa *et al.* (2018) assessed detection distance by

comparing eDNA fish inventories with observational data within intervals from 1 to 10 km

upstream from the eDNA sampling site, and showed eDNA inventories were more similar with

observational data when comparing with traditional inventories within 6 km upstream from the

eDNA sampling site. Pont *et al.* (2018) detected eDNA of Whitefish, an abundant species in Lake

Geneva, up to 130 km downstream from the lake, whereas the detection of less abundant species

was restricted to a few kilometres, suggesting that abundant species are detected farther

downstream than rare ones.

The variability among detection distances may reflect the specific local environmental

conditions of the studied water bodies (Barnes & Turner, 2016; Jane et al., 2015). Local

conditions such as acidity, oxygen demand, primary production, water temperature, solar

radiation, organic materials and the activity of microorganisms, have been shown to influence

eDNA degradation rate (Barnes & Turner, 2016; Eichmiller, Best, & Sorensen, 2016; Seymour et

al., 2018). As all the above-mentioned characteristics as well as discharge rate vary strongly

between rivers, the spatial extent of downstream transport of eDNA may be highly dependent of

the studied system. For example, by treating eDNA downstream transport as the vertical transfer

of fine particulate organic matter from the water column to the riverbed, Pont *et al.* (2018)

suggested that detection distance varies with river size, ranging from a few km in small streams

to more than 100 km in large rivers.

Overall, the discrepancy of detection distances among species and rivers reported in the

literature makes difficult to know whether the species recorded in an eDNA sample reflect a

local fish assemblage or a more spatially extended synthesis of the species present in the whole

river basin located upstream from eDNA sampling sites. Such uncertainty is reinforced in tropical

environments, which host rich, but understudied, fish communities. Tropical rivers are also

characterised by distinct environmental conditions than temperate and cold river ecosystems,

where most developments of eDNA method were achieved (Zinger *et al.* 2020). Moreover, given

the deep human-induced changes of freshwater fish biodiversity throughout the world (Su et al.,

2021), it is now urgent to determine if the fast and non-destructive eDNA samples provide

similar biodiversity measures to the traditional time consuming and often destructive capture-

based methods.

Here, we aimed at characterizing the spatial signal of eDNA in two large Neotropical

rivers through two complementary approaches considering: i) spatial patterns of the whole fish

community and ii) the individual spatial range of species along the upstream-downstream river

continuum. Considering the whole fish community, we sought to describe the decay of similarity

between communities with increasing distance between sampling sites. This widely recognised

pattern in ecology postulates that the similarity between local measures of biodiversity should

decline with the distance between observations (Soininen *et al.* 2007; Morlon *et al.* 2008, Nekola

and White 1999). This pattern should be retrieved if eDNA detection distance is short, whereas it

should be blurred under the hypothesis of a long-distance transport of eDNA. We test this

hypothesis in the Maroni and Oyapock rivers, two species rich Neotropical rivers differing in

their anthropization levels. We predict a steeper distance decay in the Maroni river due to an

upstream-downstream increase of anthropic disturbances (Gallay *et al.* 2018), which co-varies

with environmental changes along the course of the river. We also compared the eDNA distance

decay pattern to capture-based data available from the Maroni river. We here hypothesise that

under a short eDNA detection distance, distance decay of similarity between eDNA and capture

based methods should be the same. Regarding each fish species, we compared the distribution

range of detected species along the upstream-downstream gradient of the Maroni river to the

distribution range recorded by capture-based data in the same river. Again, under the hypothesis

of a local eDNA detection of the species, the distribution range of species retrieved using eDNA

should fall within the known species range, while under the hypothesis of a long-distance

transport, species should be detected downstream from their known distribution range.

Materials and methods

Study sites

This study was conducted in two tropical rivers located in the Northern East of the Amazonian

region (*sensu lato,* including Guiana shield and Amazon river drainage) (Figure 1). The Maroni

river measures 612 km from the source to the estuary and its drainage basin covers a surface of

more than 68 000 km² in Suriname and French Guiana. The Oyapock river (404 km long and

26 800 km²) flows over Amapa (Brazil) and French Guiana. Both rivers are characterized by a

low topographic gradient (mean slope =0.4% and 0.5% for Maroni and Oyapock respectively),

warm water (30.5 ± 0.23 °C and 28.4 ± 0.35 °C) and neutral pH (7 ± 0.05 and 7.2 ± 0.03). Both

rivers host typical freshwater Amazonian fauna with more than 270 and 190 described fish

species in the Maroni and Oyapock, respectively (Le Bail et al., 2012). Among these species, 141

and 119 strictly freshwater species (estuarine species were excluded) have been recorded in the

main channel of the Maroni and Oyapock rivers, respectively (Planquette *et al.* 1996; Keith *et al.*

2000). Despite being located in one of the largest unfragmented rainforest areas in the world, the

Maroni river is facing an unprecedented rise of land use changes due mainly to gold-mining

activities but also to agriculture and urbanization (Gallay et al., 2018; Rham et al., 2017). Such

anthropogenic disturbances show an increasing gradient along the course of the Maroni.

Upstream areas are free of human settlements and part of a National Park, whereas human

densities as well as deforestation and mining intensities are increasing toward the downstream

part of the river (Gallay et al., 2018). Thus, environmental and anthropogenic gradients are co-

varying along the Maroni river. Contrastingly, the Oyapock river remains less impacted, with low

human population density and gold mining activities limited to a few tributaries (Gallay et al.,

2018).

eDNA sampling

We collected environmental DNA samples in 53 sites along the main channel of two rivers

(Figure 1), with 28 sites sampled on the Maroni main channel and 25 sites sampled on the

Oyapock main channel. The estuary sections, from the sea to the upstream area influenced by the

tide, were not considered in this study, because tidal water movements could bias our results

through downstream-upstream movement of eDNA during the rising tide. The sampling was

achieved during a 2 weeks field session during the dry season (October- November) in 2017 and

2018 for the Maroni and Oyapock, respectively. In all sites, the river was wider than 20 meters

and deeper than one meter (Strahler orders 4 to 8). The distance between adjacent sites ranged

from 1.6 to 31.8 km (mean: 13.8 and SE: 1.4) in the Maroni, and from 1.2 to 23.8 km (mean: 10.8

and SE: 1.5) in the Oyapock. In both rivers, sites were sequentially sampled from downstream to

upstream, with 1 to 4 sites sampled per day, according to the travel time between sites. All the

metadata associated with the samples are available through the CEBA geoportal (vmcebagn-

dev.ird.fr) or the French Guiana geoportal (geoguyane.fr) under reference 5617a9ff-d0aa-48a9-

b2c2-cb7fd5b92692 (Murienne et al., 2019).

190   Following the protocol implemented by Cantera et al. (2019), we filtered two samples of

34 liters of water at each site to collect eDNA. A peristaltic pump (Vampire sampler, Burlke,

Germany) and a single-use tubing were used to pump the water into a single-use filtration capsule

(VigiDNA 0.45 µm; SPYGEN, le Bourget-du-Lac, France). All the material used to collect water

was single use, and therefore replaced between each sample. Single use gloves were also used to

avoid contamination. To collect eDNA, the input part of the tubing was placed 10 to 20

centimetres below the surface in areas with high water flow as recommended by Cilleros *et al.*

(2019). Sampling was achieved in turbulent area (rapid hydromorphologic unit) to ensure an

optimal homogenization of the eDNA throughout the water column. To avoid DNA

contamination among sites, the operator always remained downstream from the filtration area and

stayed on emerging rocks. At the end of the filtration, the filtration capsule was emptied of water,

filled with 80 mL of CL1 conservation buffer (SPYGEN) and stored in individual sterile plastic

bags kept in the dark.

Capture-based sampling

The fish captures were performed using a standardized gill-net protocol designed by Tejerina-

Garro and De Mérona (2001) and used as a routine biodiversity monitoring protocol by the

French Ministry of Environment (DGTM), as part of the European Water Framework directive.

Five sites located on the Maroni main channel were sampled yearly, during the dry season, from

2007 to 2016. Briefly, the protocol consist in sampling fish overnight using ten gill-nets

measuring 25 metre long and 1.5 metre height and with mesh sizes ranging from 10 to 70

millimetres. The 10 nets are set before down, over a river reach measuring 300 to 500 metres.

Nets were removed 12 hours later, after down, and all captured fish were identified to the species

level. This protocol allows capturing both diurnal and nocturnal species, and most of the fish size

ranges except the smallest species. Yearly data were pooled to provide a broad characterisation of

the fish fauna in each site. This permitted to deal with the limited efficiency of gill-nets to collect

the entire fauna. Indeed, gill-nets are strongly dependent of fish activity and need repeated

sampling effort to provide realistic species community inventories (Murphy and Willis 1996;

Pont *et al.* 2018).

eDNA laboratory and bioinformatic analyses

For DNA extraction, each filtration capsule was agitated for 15 min on an S50 shaker (cat

Ingenieurbüro™) at 800 rpm and then emptied into a 50 mL tube before being centrifuged for 15

min at 15,000×g. The supernatant was removed with a sterile pipette, leaving 15 mL of liquid at

the bottom of the tube. Subsequently, 33 mL of ethanol and 1.5 mL of 3M sodium acetate were

added to each 50 mL tube and stored for at least one night at -20°C. Tubes were centrifuged at 15

000 ×g for 15 min at 6°C, and the supernatants were discarded. After this step, 720 µL of ATL

buffer from the DNeasy Blood & Tissue Extraction Kit (Qiagen) was added. The tubes were then

vortexed, and the supernatants were transferred to 2 mL tubes containing 20 µL of proteinase K.

The tubes were finally incubated at 56°C for two hours. Afterwards, DNA extraction was

performed using NucleoSpin Soil (Macherey-Nagel GmbH & Co., Düren Germany) starting from

step six and following the manufacturer's instructions. The elution was performed by adding 100

µL of SE buffer twice. Four negative extraction controls were also performed. They were

amplified and sequenced in parallel to the field samples to monitor possible laboratory

contaminants. After the DNA extraction, the samples were tested for inhibition by qPCR (Biggs

et al. (2015). If the sample was considered inhibited, it was diluted 5-fold before the

amplification.

We performed DNA amplifications in a final volume of 25 μL including 1 U of AmpliTaq

Gold DNA Polymerase (Applied Biosystems, Foster City, CA), 10 mM of Tris-HCl, 50 mM of

KCl, 2.5 mM of $MgCl_2$, 0.2 mM of each dNTP, 0.2 μM of "teleo" primers (Valentini et al., 2016)

and 3 μL of DNA template. We also added human blocking primer for the "teleo" primers with a

final concentration of 4 μM and 0.2 μg/μL of bovine serum albumin (BSA, Roche Diagnostic,

Basel, Switzerland) to the mixture. We performed 12 PCR replicates per field sample. The

forward and reverse primer tags were identical within each PCR replicate. The PCR mixture was

denatured at 95°C for 10 min, followed by 50 cycles of 30 s at 95°C, 30 s at 55°C and 1 min at 72

°C and a final elongation step at 72°C for 7 min. This step was done in a room dedicated to

amplified DNA with negative air pressure and physical separation from the DNA extraction

rooms (with positive air pressure). We also amplified 14 negative extraction controls and five

PCR negatives controls and sequenced them in parallel with the PCR replicates. We pooled the

purified PCR products in equal volumes to achieve an expected sequencing depth of 500,000

reads per sample before the libraries preparation. Ten libraries were prepared using a PCR-free

library protocol (https://www.fasteris.com/metafast), at Fasteris facilities (Geneva, Switzerland).

Four libraries were sequenced using an Illumina HiSeq 2500 (2x125 bp) (Illumina, San Diego,

CA, USA) and the HiSeq SBS Kit v4, three using a MiSeq (2x125 bp) (Illumina, San Diego, CA,

USA) and the MiSeq Flow Cell Kit Version3 (Illumina, San Diego, CA, USA) and three using a

NextSeq (2x150 bp+8bp) (Illumina, San Diego, CA, USA) and the NextSeq Mid kit (Illumina,

San Diego, CA, USA). The libraries ran on the NextSeq were equally distributed in four lanes.

Sequencing were performed following the manufacturer's instructions at Fasteris facilities

(Geneva, Switzerland).

The sequence reads were analyzed using the OBITools package (Boyer et al., 2016)

following the protocol described in Valentini et al. (2016). The `ecotag` program was used for

the taxonomic assignment of molecular operational taxonomic units (MOTUs). We kept only the

MOTUS with a similarity to our reference database above 98%. Our local reference database

represent an updated version of the reference database available in Cilleros *et al.* (2019), resulting

in 251 Guianese fish species. The GenBank nucleotide database was checked but Guianese fishes

being poorly referenced (most of the sequences are from Cilleros *et al.* (2019)), it did not

provided additional information in our case. We discarded all MOTUs with a frequency of

occurrence below 0.001 per library in each sample, considered as tag-jumps (Schnell, Bohmann,

& Gilbert, 2015). These thresholds were empirically determined to clear all reads from the

extraction and PCR negative controls included in our global data production procedure as

suggested by De Barba et al. (2014) and Taberlet et al. (2018). For the samples sequenced on a

NextSeq platform, only species present in at least two lanes were kept. The entire procedure was

repeated for each sample, giving rise to two species lists per field site. The two obtained species

lists per site were finally pooled to obtain a single species inventory per site, as Cantera *et al.*

(2019) showed that pooling two replicate field samples was sufficient to inventory most of the

fauna in highly diverse tropical aquatic ecosystems from the same region.

eDNA data ability to reveal spatial patterns in ecology

We first tested the ability of eDNA data to reveal patterns of decay of similarity in species

composition between pairs of sites with increasing geographic distance. Presence/absence

matrices of the species detections in each site were used to build a species compositional distance

matrix between pairs of sites, for each river separately with the *vegdist* function. Pairwise

compositional distances between sites were calculated using the Jaccard index. We used linear

regressions to model the relationship between species similarity distances and watercourse

distances between sites (in km) for each river. Considering that the environment have been shown

to influence the rate of similarity decay (Nekola & White, 1999), differences in the rates of decay

in similarity with distance between rivers were tested. To do this, the slopes and intercepts of the

obtained decay relationships were compared by using a permutation procedure with 999

iterations based on linear regressions. The permutation procedure was done with the *diffslope* and

diffic functions of the 'simba' R package for the slope and intercept comparisons, respectively.

This function compares the observed difference between the slopes or intercepts of each linear

regression from each river with the permuted slope differences by randomly interchanging the

values between the two river datasets. The p-value is computed as the ratio between the number

of cases where the differences in slopes exceed the difference in slope of the actual configuration.

The distance decay pattern obtained with eDNA samples between the Oyapock and

Maroni rivers was compared. Then, we did the same between eDNA and capture-based data from

the Maroni river (Figure 1). The comparison between the eDNA and capture-based distance

decay profiles was first made with all the species detected (or captured) by the two methods. To

deal with differences in species capturability between methods, and particularly with the low

capturability of small (less than 10 centimetres long) or non-mobile species using gill-nets.

Finally, we restricted the comparison to the species detected (or captured) with both methods.

For the species level assessment, the spatial upstream-downstream range of each detected

species was compared to the known distribution range of the species derived from the capture-

based method. We used the watercourse distance from the river mouth (in km) of the sampled

sites to represent the spatial distribution range along the upstream-downstream gradient. To keep

the spatial range comparable between the two sampling methods, we discarded the most upstream

sites sampled by eDNA (S1 to S6, see Figure 1), as this part of the Maroni river has not been

investigated using gill nets. Moreover, gill-net being selective for species (*e.g.* small and/or

elongated species are not captured) we restricted this analysis to the species detected by both

eDNA and by gill-nets in at least two sites or sampling events. Indeed, the 11 species detected

only once over the 9 years' gill-net surveys represent "bycatches" of small species (<10 cm) such

as *Characidium zebra* or *Hemigrammus surinamensis,* which are hardly captured using gill-nets.

We here predict that under a short eDNA detection distance, species detections using eDNA

should not systematically occur downstream from the known distribution range of the species as

established by gill-net data.

Results

Fish species were detected in all the eDNA replicates. The number of species per site ranged

from 50 to 71 (mean = 62) in the Oyapock sites and 30 to 94 (mean = 57) in the Maroni sites. We

detected a total of 160 species, with 116 and 96 species in the Oyapock and Maroni rivers,

respectively. They account for more than 80% (80.7% and 82.3% for the Oyapock and Maroni,

respectively) of the fish species known from the main channel of both river basins, excluding

estuarine species (Le Bail, Keith, & Planquette, 2000; Planquette, Keith, P, & Le Bail, 1996).

Among the five Maroni sites sampled with capture-based methods (gill-nets), a total of 94 fish

species were captured, with species richness ranging from 50 to 66 per site. The comparison of

eDNA (22 sites, sites S7 to S28, Figure 1) and gill-nets (5 sites) inventories revealed that 53

species were detected by both methods in the Maroni river.

Results from the linear models showed that similarity decreased significantly with

watercourse distance between sites pairs sampled with eDNA data in both Oyapock ($p<0.001$,

slope= -0.0011, $R^2=0.5$) and Maroni ($p<0.001$, slope= -0.00125, $R^2=0.6$) rivers. As sites became

more distant, their corresponding fish communities became more dissimilar in species (Figure

2**A**). For both rivers, the similarity decay with distance relationship fitted well without obvious

change of the slope of the regression in any distance value. The permutation procedure showed

that the negative effect of spatial distance on species similarity was more marked in the Maroni

than in the Oyapock river, with a significantly more negative slope (permutation test, $p = 0.02$)

on the Maroni river.

The distance decay of similarity for the sites sampled with gill-nets in the Maroni river,

was not significant ($p = 0.08$, slope= -0.0005, $R^2=0.3$), despite a trend of similarity decrease with

distance between site pairs (Figure 2**B**). The slope of this trend was significantly less marked than

the slope of the similarity decay reported with eDNA data ($p<0.001$). Restricting the species data

to the species detected (or captured) by both sampling methods provided a significant decay of

similarity with distance in both eDNA ($p<0.001$, slope= -0.0009, $R^2=0.6$) and capture-based

inventories ($p=0.022$, slope= -0.0006, $R^2=0.47$). The decreasing slope of the relationship obtained

with the eDNA data was significantly more marked ($p<0.001$) than the slope obtained with

captured based data (Figure 2C).

The comparison of the species spatial ranges along the main channel of the Maroni river

provided by the eDNA approach and by gill-net captures revealed a global match of the species

distribution between methods (Figure 3). Among the 53 species detected by both methods, 74.5%

of the fauna (38 species) was not detected by eDNA downstream from capture-based records.

Indeed, distribution ranges were similar between methods for 20 species (39% of the fauna, e.g.

Hoplias aimara, HAIM; *Eigenmannia virescens,* EVIR; *Brycon falcatus,* BFAL; Figure 3) and

for 18 species (35.3% of the fauna) the capture-based method detected the species downstream

from the distribution range found using eDNA (*e.g. Moenkhausia georgiae,* MGEO;

Pristobrycon eigenmanni, SHUM; Figure 3). Only 25.5% of the considered fauna (13 species)

were detected using eDNA downstream from the distribution provided by capture-based records

(*e.g. Electrophorus electricus,* EELE; *Metaloricaria paucidens,* MPAU; *Geophagus harreri,*

GHAR*, Guianacara owroewefi,* GOWR; Figure 3). Furthermore, eDNA and capture-based

records retrieved consistent wide distribution ranges along the main channel of the river (*i.e.*

species detected in almost all the eDNA and gill-net sites) for 21 species (*e.g. H. aimara,* HAIM;

Figure 3). Nevertheless, six species (*G. harreri,* GHAR; *P. barbatus,* PBAR; *Acnodon*

oligacanthus, AOLI; *Pseudoplatystoma fasciatum,* PFAS; *E. electricus,* EELE) had a wide

distribution range in eDNA records, but not in capture-based records.

Discussion

As opposed to the 'conveyor belt' hypothesis (Deiner et al. 2016) which implies that

communities are homogenized across an entire basin because of eDNA transport, our results

revealed that eDNA data allowed to retrieve expected spatial patterns of fishes in large tropical

rivers at both community and species levels. The decay of similarity between local communities

with increasing distance as well as most species distribution ranges were similar to those reported

from capture-based samples.

Species similarity in both Maroni and Oyapock rivers was maximal between nearby fish

communities, and declined when increasing distances between sampling sites, thus fitting the

distance decay theory, a general rule in ecology (Nekola and White 1999). This rule has already

been verified in temperate rivers where it reflects the dynamic and heterogeneous architecture of

river systems that shape community composition at different spatial scales along the river

continuum (Muneepeerakul et al., 2008). However, the distance decay of similarity has rarely

been examined in Neotropical aquatic diversity (Araújo et al., 2013). Importantly, the fact that

eDNA data was able to reveal distance decay of faunistic similarity patterns necessarily implies a

short downstream transport of eDNA. Indeed, according to Nekola and White (1999), the grain

size, corresponding to the contiguous area over which a single observation is made

(corresponding to the eDNA integration distance in the present study), should be smaller than the

whole spatial extent of the study. This allows capturing variation in similarities among

comparisons and thus observing distance decay patterns. In our case, this means that the spatial

signal of eDNA to detect one fish community need to be smaller than the sampled portion of the

river, which is verified for the two rivers, and remains true even when considering only a

restricted section of each river. Therefore, the hypothesis of eDNA being transported over long-

distances and cumulating downstream was here rejected. Nevertheless, we cannot exclude that

the eDNA of some species, probably the most abundant ones, can be transported over long

distances as shown by Pont *et al.* (2018) in the Rhone river (France). In the case that this occurs,

it would certainly be limited to a few species in our samples as this potential bias did not blurred

the significant patterns of distance decay of similarity with geographic distance.

The slopes of similarity decay differed between the two rivers, with a more marked

decline in the Maroni river, suggesting a marked decrease of environmental similarities between

sites and/or more dispersal constraints with distance in this river compared to the Oyapock

(Nekola & White, 1999; Soininen et al., 2007). It can be expected that the decrease of

environmental similarities and the increase of dispersal constraints are reinforced by an upstream-

downstream gradient of human impacts in the Maroni river, resulting in a higher rate of similarity

decay with distance. Indeed, the Maroni river faces six-fold higher deforestation levels than the

Oyapock river, (0.37% *versus* 0.06% of catchment area, Gallay *et al.* 2018). Deforestation was

mainly caused by gold-mining activities that deeply impact sediment fluxes in the Maroni River

(Gallay et al., 2018) and thus strengthen environmental constrains for species (Brosse,

Grenouillet, Gevrey, Khazraie, & Tudesque, 2011; Hammond, Gond, Thoisy, Forget, & DeDijn,

2007). This difference between river basins was significantly observed in our eDNA samples,

testifying for the relevance of ecological patterns retrieved using eDNA.

Similar patterns of distance decay of similarity in the Maroni river were obtained with

eDNA and capture-based data, a pattern which was verified and even reinforced when only the

fish species captured by both methods were considered. These results validate, with fish captures,

the relevance of the distance decay pattern revealed by eDNA data. Moreover, obtaining similar

(and even more marked) distance decay between methods strongly suggests a similar spatial

sampling range between methods, and thus reveal that eDNA detection distance is equivalent or

even smaller to the size of the reach sampled with gill-nets (*i.e.* from 200 to 500 metres).

Interestingly, the negative slope of the eDNA distance decay pattern was significantly more

marked than the one retrieved for capture based data. This might either be due to a detection

distance for eDNA shorter than the spatial extent of a gill-net sample, or more likely to a higher

completeness of eDNA inventories compared to gill-net samples. Indeed, gill-nets are

particularly selective in species (Murphy and Willis 1996) and therefore only provide a partial

image of the fauna occurring in each site (Cilleros et al., 2019).

Comparing species distribution ranges retrieved from eDNA and capture samples using

gill-nets revealed a global match in more than 70% of the species, which corroborates a limited

spatial signal of the eDNA detection of the species. More importantly, only 25% of the species

were detected using eDNA downstream from their distribution range recorded using capture-

based methods. Although this could be considered as species detections due to a downstream

drift of eDNA, 10 out of these 13 species have an extended spatial distribution in the Maroni

drainage basin and are known from the downstream reaches of the Maroni (Planquette et al.

1996; Keith et al. 2000). The absence of those species in the recent gill-net captures is probably

explained by the low capturability of those species due to their morphological and ecological

characteristics (Murphy and Willis 1996). For instance, the anguilliform body shape of *E.*

electricus (EELE) makes the species difficult to capture with gill-nets, and the sedentary or

territorial behaviour of *M. paucidens* (MPAU), *G. harreri* (GHAR), *G. owroewefi* (GOWR) and

K. itany (KITA) reduces the probability to capture them with passive sampling gears such as gill-

nets. Conversely, the 20 species having consistent distributions between methods are species

having high capture probability with gill-nets, such as large bodied (*e.g. H. aimara*, HAIM),

mobile and gregarious species (*e.g. Brycon falcatus*, BFAL; *Hemiodus unimaculatus,* HEUN), or

species with spines and ossified fins that easily tangle in the nets (*e.g. Acuchenipterus nuchalis*,

ANUC; *Doras micropoeus* DMIC, *Pimelodus ornatus*, PORN).

Conclusions

We here demonstrated that fish inventories achieved using eDNA provide an assessment of the

fauna at a limited spatial grain, making this biodiversity data appropriate to describe spatial

patterns of fish communities and test ecological theories. The scale of eDNA detection certainly

does not account for the local habitat of the species that has to be measured over a few square

meters. Nevertheless, a detection distance encompassing several hundred of meters appears as a

reasonable sample grain size for studying communities or to consider anthropic impacts on fish

communities (Jackson, Peres-Neto, & Olden, 2001; Tonn, 1990). Moreover, we show that eDNA

can inventory fish fauna at a similar spatial grain than gill-net capture sampling protocols. It

might therefore be possible to compare historical and current data on fish faunas, and thus to

follow biodiversity alterations experienced by freshwater faunas though current global changes as

reported by (Su et al., 2021, p. 202). Moreover, freshwater fish are recognized as one of the most

threatened taxa (Hughes, 2021) and the expected species extinction by the end of the century

might deeply impact the functional properties of fish assemblages, as well as the services they

provide to human societies (Carmona et al., 2021). Under this environmental urgency, eDNA

inventories represent a fast and easy to handle fish inventory method providing equivalent and

even more precise data than capture based methods.

Acknowledgements

This work was funded by DGTM Guyane (project DISTDET), Investissement d'Avenir grant

DRIIHM (ANR-11-LABX-0010), through the OHM Oyapock, and the VigiLIFE project. The

References

459
Araújo, E. S., Marques, E. E., Freitas, I. S., Neuberger, A. L., Fernandes, R., & Pelicice, F. M.

(2013). Changes in distance decay relationships after river regulation: Similarity among

fish assemblages in a large Amazonian river. *Ecology of Freshwater Fish*, *22*(4), 543–

552. doi: 10.1111/eff.12054

Barnes, M. A., & Turner, C. R. (2016). The ecology of environmental DNA and implications for

conservation genetics. *Conservation Genetics*, *17*(1), 1–17. doi: 10.1007/s10592-015-

0775-4

Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C., Dejean, T., Griffiths, R. A., … Dunn, F.

(2015). Using eDNA to develop a national citizen science-based monitoring programme

for the great crested newt (Triturus cristatus). *Biological Conservation*, *183*, 19–28. doi:

10.1016/j.biocon.2014.11.029

Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P., & Coissac, E. (2016). obitools: A

unix-inspired software package for DNA metabarcoding. *Molecular Ecology Resources*,

16(1), 176–182.

Brosse, S., Grenouillet, G., Gevrey, M., Khazraie, K., & Tudesque, L. (2011). Small-scale gold

mining erodes fish assemblage structure in small Neotropical streams. *Biodiversity and*

Conservation, *20*(5), 1013–1026. doi: 10.1007/s10531-011-0011-6

Cantera, I., Cilleros, K., Valentini, A., Cerdan, A., Dejean, T., Iribar, A., … Brosse, S. (2019).

Optimizing environmental DNA sampling effort for fish inventories in tropical streams

and rivers. *Scientific Reports*, *9*(1). doi: 10.1038/s41598-019-39399-5

Carmona, C. P., Tamme, R., Pärtel, M., de Bello, F., Brosse, S., Capdevila, P., … Toussaint, A.

(2021). Erosion of global functional diversity across the tree of life. *Science Advances*,

7(13), eabf2675. doi: 10.1126/sciadv.abf2675

Cilleros, K., Valentini, A., Allard, L., Dejean, T., Etienne, R., Grenouillet, G., … Brosse, S.

(2019). Unlocking biodiversity and conservation studies in high-diversity environments

using environmental DNA (eDNA): A test with Guianese freshwater fishes. *Molecular*

Ecology Resources, *19*(1), 27–46. doi: https://doi.org/10.1111/1755-0998.12900

Civade, R., Dejean, T., Valentini, A., Roset, N., Raymond, J.-C., Bonin, A., … Pont, D. (2016).

Spatial representativeness of environmental DNA metabarcoding signal for fish

biodiversity assessment in a natural freshwater system. *PloS One*, *11*(6), e0157366.

De Barba, M., Miquel, C., Boyer, F., Mercier, C., Rioux, D., Coissac, E., & Taberlet, P. (2014).

DNA metabarcoding multiplexing and validation of data accuracy for diet assessment:

Application to omnivorous diet. *Molecular Ecology Resources*, *14*(2), 306–323. doi:

10.1111/1755-0998.12188

Deiner, K., & Altermatt, F. (2014). Transport distance of invertebrate environmental DNA in a

natural river. *PLoS One*, *9*(2), e88786.

Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J.-C., & Altermatt, F. (2016). Environmental

DNA reveals that rivers are conveyer belts of biodiversity information. *Nature*

Communications, *7*, 12544. doi: 10.1038/ncomms12544

Eichmiller, J. J., Best, S. E., & Sorensen, P. W. (2016). Effects of Temperature and Trophic State

on Degradation of Environmental DNA in Lake Water. *Environmental Science &*

Technology, *50*(4), 1859–1867. doi: 10.1021/acs.est.5b05672

Evans, N. T., Li, Y., Renshaw, M. A., Olds, B. P., Deiner, K., Turner, C. R., … Pfrender, M. E.

(2017). Fish community assessment with eDNA metabarcoding: Effects of sampling

design and bioinformatic filtering. *Canadian Journal of Fisheries and Aquatic Sciences*,

74(9), 1362–1374. doi: 10.1139/cjfas-2016-0306

Ficetola, G. F., Miaud, C., Pompanon, F., & Taberlet, P. (2008). Species detection using

environmental DNA from water samples. *Biology Letters*, *4*(4), 423–425. doi:

10.1098/rsbl.2008.0118

Gallay, M., Martinez, J., Allo, S., Mora, A., Cochonneau, G., Gardel, A., … Laraque, A. (2018).

Impact of land degradation from mining activities on the sediment fluxes in two large

rivers of FRENCH GUIANA. *Land Degradation & Development*, *29*(12), 4323–4336. doi:

10.1002/ldr.3150

Goldberg, C. S., Pilliod, D. S., Arkle, R. S., & Waits, L. P. (2011). Molecular Detection of

Vertebrates in Stream Water: A Demonstration Using Rocky Mountain Tailed Frogs and

Idaho Giant Salamanders. *PLoS ONE*, *6*(7), e22746. doi: 10.1371/journal.pone.0022746

Hammond, D. S., Gond, V., Thoisy, B. de, Forget, P.-M., & DeDijn, B. P. E. (2007). Causes and

consequences of a tropical forest gold rush in the Guiana Shield, South America. *AMBIO:*

A Journal of the Human Environment, *36*(8), 661–670. doi: 10.1579/0044-

7447(2007)36[661:CACOAT]2.0.CO;2

Hänfling, B., Lawson Handley, L., Read, D. S., Hahn, C., Li, J., Nichols, P., … Winfield, I. J.

(2016). Environmental DNA metabarcoding of lake fish communities reflects long-term

data from established survey methods. *Molecular Ecology*, *25*(13), 3101–3119. doi:

10.1111/mec.13660

Hughes, K. (2021). *The world's forgotten fishes*. UICN report.

Jackson, D. A., Peres-Neto, P. R., & Olden, J. D. (2001). *What controls who is where in*

freshwater fish communities—The roles of biotic, abiotic, and spatial factors. *58*, 14.

Jane, S. F., Wilcox, T. M., McKelvey, K. S., Young, M. K., Schwartz, M. K., Lowe, W. H., …

Whiteley, A. R. (2015). Distance, flow and PCR inhibition: EDNA dynamics in two

headwater streams. *Molecular Ecology Resources*, *15*(1), 216–227. doi: 10.1111/1755-

0998.12285

Jerde, C. L., Mahon, A. R., Chadderton, W. L., & Lodge, D. M. (2011). "Sight-unseen" detection

of rare aquatic species using environmental DNA: EDNA surveillance of rare aquatic

species. *Conservation Letters*, *4*(2), 150–157. doi: 10.1111/j.1755-263X.2010.00158.x

Keck, F., Vasselon, V., Tapolczai, K., Rimet, F., & Bouchez, A. (2017). Freshwater

biomonitoring in the Information Age. *Frontiers in Ecology and the Environment*, *15*(5),

266–274. doi: 10.1002/fee.1490

Le Bail, P. Y., Covain, R, Jegu, M, Fisch-Muller, S, Vigouroux, R, & Keith, P. (2012). *Updated*

checklist of the freshwater and estuarine fishes of French Guiana (Cybium).

Le Bail, P. Y., Keith, P., & Planquette, P. (2000). *Atlas des poissons d'eau douce de Guyane.*

Tome 2, fascicule II☐: Siluriformes. Paris, France: M.N.H.N./S.P.N.

McCluney, K. E., Poff, N. L., Palmer, M. A., Thorp, J. H., Poole, G. C., Williams, B. S., …

Baron, J. S. (2014). Riverine macrosystems ecology: Sensitivity, resistance, and resilience

of whole river basins with human alterations. *Frontiers in Ecology and the Environment*,

12(1), 48–58. doi: 10.1890/120367

Morlon, H., Chuyong, G., Condit, R., Hubbell, S., Kenfack, D., Thomas, D., … Green, J. L.

(2008). A general framework for the distance-decay of similarity in ecological

communities. *Ecology Letters*, *11*(9), 904–917. doi: 10.1111/j.1461-0248.2008.01202.x

Muneepeerakul, R., Bertuzzo, E., Lynch, H. J., Fagan, W. F., Rinaldo, A., & Rodriguez-Iturbe, I.

(2008). Neutral metacommunity models predict fish diversity patterns in Mississippi–

Missouri basin. *Nature*, *453*(7192), 220–222. doi: 10.1038/nature06813

Murienne, J., Cantera, I., Cerdan, A., Cilleros, K., Decotte, J.-B., Dejean, T., … Brosse, S.

(2019). Aquatic eDNA for monitoring French Guiana biodiversity. *Biodiversity Data*

Journal, *7*, e37518. doi: 10.3897/BDJ.7.e37518

Murphy, B. R., & Willis, D. W. (1996). *Fisheries techniques*. American Fisheries Society

Bethesda, Maryland.

Nakagawa, H., Yamamoto, S., Sato, Y., Sado, T., Minamoto, T., & Miya, M. (2018). Comparing

local- and regional-scale estimations of the diversity of stream fish using eDNA

metabarcoding and conventional observation methods. *Freshwater Biology*, *63*(6), 569–

580. doi: 10.1111/fwb.13094

Nekola, J. C., & White, P. S. (1999). The distance decay of similarity in biogeography and

ecology.   *Journal   of   Biogeography*,   *26*(4),   867–878.   doi: 10.1046/j.1365-

2699.1999.00305.x

Olds, B. P., Jerde, C. L., Renshaw, M. A., Li, Y., Evans, N. T., Turner, C. R., … Lamberti, G. A.

(2016). Estimating species richness using environmental DNA. *Ecology and Evolution*,

6(12), 4214–4226. doi: 10.1002/ece3.2186

Pilliod, D. S., Goldberg, C. S., Arkle, R. S., & Waits, L. P. (2014). Factors influencing detection

of eDNA from a stream-dwelling amphibian. *Molecular Ecology Resources*, *14*(1), 109–

116. doi: 10.1111/1755-0998.12159

Planquette, P., Keith, P, & Le Bail, P. Y. (1996). *Atlas des poissons d'eau douce de Guyane*

(tome 1) (Vol. 22). Paris, France: IEBG - M.N.H.N., INRA, CSP, Min. Env.

Pont, D., Rocle, M., Valentini, A., Civade, R., Jean, P., Maire, A., … Dejean, T. (2018).

Environmental DNA reveals quantitative patterns of fish biodiversity in large rivers

despite its downstream transportation. *Scientific Reports*, *8*(1). doi: 10.1038/s41598-018-

28424-8

Rees, H. C., Gough, K. C., Middleditch, D. J., Patmore, J. R. M., & Maddison, B. C. (2015).

Applications and limitations of measuring environmental DNA as indicators of the

presence of aquatic animals. *Journal of Applied Ecology*, *52*(4), 827–831. doi:

10.1111/1365-2664.12467

Rees, H. C., Maddison, B. C., Middleditch, D. J., Patmore, J. R. M., & Gough, K. C. (2014).

REVIEW: The detection of aquatic animal species using environmental DNA - a review

of eDNA as a survey tool in ecology. *Journal of Applied Ecology*, *51*(5), 1450–1459. doi:

10.1111/1365-2664.12306

Rham, M., Thibault, P., Shapiro, A., Smartt, T., Paloeng, C., Crabbe, S., … Joubert, P. (2017).

Monitoring the impact of gold mining on the forest cover and freshwater in the Guiana

Shield. (p. 20).

Schnell, I. B., Bohmann, K., & Gilbert, M. T. P. (2015). Tag jumps illuminated—Reducing

sequence-to-sample misidentifications in metabarcoding studies. *Molecular Ecology*

Resources, *15*(6), 1289–1303. doi: 10.1111/1755-0998.12402

Seymour, M., Durance, I., Cosby, B. J., Ransom-Jones, E., Deiner, K., Ormerod, S. J., … Creer,

S. (2018). Acidity promotes degradation of multi-species environmental DNA in lotic

mesocosms. *Communications Biology*, *1*(1), 4. doi: 10.1038/s42003-017-0005-3

Soininen, J., McDonald, R., & Hillebrand, H. (2007). The distance decay of similarity in

ecological communities. *Ecography*, *30*(1), 3–12. doi: 10.1111/j.0906-7590.2007.04817.x

Su, G., Logez, M., Xu, J., Tao, S., Villéger, S., & Brosse, S. (2021). Human impacts on global

freshwater fish biodiversity. *Science*, *371*(6531), 835. doi: 10.1126/science.abd3369

Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). *Environmental DNA for biodiversity*

research and monitoring. Oxford, U.K.: Oxford University Press.

Takahara, T., Minamoto, T., Yamanaka, H., Doi, H., & Kawabata, Z. (2012). Estimation of Fish

Biomass Using Environmental DNA. *PLoS ONE*, *7*(4), e35868. doi:

10.1371/journal.pone.0035868

Tejerina-Garro, F. L., & De MéRona, B. (2001). Gill net sampling standardisation in large rivers

of French Guiana (South America). *Bulletin Français de La Pêche et de La Pisciculture*,

(357–360), 227–240. doi: 10.1051/kmae/2001046

Tonn, W. M. (1990). Climate Change and Fish Communities: A Conceptual Framework.

Transactions of the American Fisheries Society, *119*(2), 337–352. doi: 10.1577/1548-

8659(1990)119<0337:CCAFCA>2.3.CO;2

Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., … Dejean, T.

(2016). Next-generation monitoring of aquatic biodiversity using environmental DNA

metabarcoding. *Molecular Ecology*, *25*(4), 929–942. doi: 10.1111/mec.13428

Wilcox, T. M., McKelvey, K. S., Young, M. K., Sepulveda, A. J., Shepard, B. B., Jane, S. F., …

Schwartz, M. K. (2016). Understanding environmental DNA detection probabilities: A

case study using a stream-dwelling char Salvelinus fontinalis. *Biological Conservation*,

194, 209–216. doi: 10.1016/j.biocon.2015.12.023

Zinger, L., Donald, J., Brosse, S., Gonzalez, M. A., Iribar, A., Leroy, C., … Lopes, C. M. (2020).

Advances and prospects of environmental DNA in neotropical rainforests. In *Advances in*

Ecological Research (Vol. 62, pp. 331–373). Elsevier. doi: 10.1016/bs.aecr.2020.01.001

617
618

Authors' contributions

IC and SB conceived the ideas, designed methodology and led the writing of the manuscript; SB, JBD, JM and RV collected the data; IC analyzed the data; AV and TD conducted the laboratory work and bio-informatic analyses.

Conflict of interest

Teleo primers and the use of the amplified fragment for identifying fish species from environmental samples are patented by the CNRS and the Université Grenoble Alpes. This patent only restricts commercial applications and has no implications for the use of this method by academic researchers. SPYGEN owns a licence for this patent. A.V. and T.D. are research scientists at a private company specialising in the use of eDNA for species detection.

Figures

Figure 1: Map of the study area indicating the eDNA sampling sites in the
Maroni (red dots) and Oyapock (blue dots) rivers. Grey triangles
correspond to sites sampled with a capture-based method (gill-nets). Inset
map on the top indicate the location of the study area in South America.

Figure 2: Decay of fish species similarity (Jaccard index) with watercourse
distance between pairs of sites. The solid lines and 95% confidence intervals
(shaded areas) were fitted through linear regression. (**A**) Distance decay for
the Oyapock (blue) and Maroni (red) sites sampled with eDNA. (**B**) Distance
decay in the Maroni for eDNA (red dots) and capture-based (black triangles)
samples considering all captured (or detected) fauna. (**C**) Distance decay in
the Maroni for eDNA (red dots) and capture-based (black triangles) samples
considering the species captured (or detected) with both capture-based and
eDNA methods.

Figure 3: Species distribution ranges along the upstream-downstream gradient of the Maroni river derived from capture-based samples (grey
boxplots) and eDNA (red boxplots). Species are ordered according to the mean distance from the river mouth of the sites where they were detected
using eDNA (increasing mean from left to right). The species names corresponding to the codes are in Table S1.