A parental transcriptional response to microsporidia infection induces inherited immunity in offspring

Alexandra R. Willis[1*], Winnie Zhao[1*], Ronesh Sukhdeo[1], Lina Wadi[1], Hala Tamim El Jarkass[1], Julie M. Claycomb[1] and Aaron W. Reinke[1#]

[1] Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada

*These authors contributed equally

#Corresponding author

aaron.reinke@utoronto.ca

Abstract

Inherited immunity is an emerging field and describes how the transfer of immunity from parents to offspring can promote progeny survival in the face of infection. The mechanisms of how inherited immunity is induced are mostly unknown. The intracellular parasite *Nematocida parisii* is a natural microsporidian pathogen of *Caenorhabditis elegan*s. Here, we show that *N. parisii*-infected worms produce primed offspring that are resistant to microsporidia infection. We find that immunity is induced in a dose dependent manner and lasts for a single generation. Intergenerational immunity prevents host cell invasion by *N. parisii* and also enhances survival to the bacterial pathogen *Pseudomonas aeruginosa*. Further, we show that inherited immunity is triggered by the host transcriptional response to infection, which can also be induced through maternal somatic depletion of negative regulators PALS-22 and the retinoblastoma protein ortholog LIN-35. We show that other biotic and abiotic stresses, such as viral infection and cadmium exposure, that induce a similar transcriptional response to microsporidia can also induce immunity in progeny. Our results demonstrate that distinct stimuli can induce inherited immunity to provide resistance against multiple classes of pathogens. These results show that activation of an innate immune response can provide protection against pathogens not only within a generation, but also in the next generation.

Introduction

Animals have evolved diverse immune mechanisms to limit the negative impact of pathogens and parasites on host fitness. While immunological memory is typically considered a hallmark of the antibody-mediated adaptive immune system, memory of pathogen exposure has now been documented in animals lacking adaptive immunity. Although, invertebrates only posses innate immunity, at least 20 species, including insects, crustaceans, and molluscs, have now been shown to transfer protective immunity to their progeny following infection[1]. Although this epigenetically inherited immunity can protect offspring against a variety of bacterial, fungal and viral pathogens, it is largely unclear how immunity is induced. Several reports have described the deposition of bacterial cell-wall fragments in offspring following parental infection, as well as immune genes being upregulated in both parents and their progeny[2,3]. Immune priming can be specific, whereby immunity is only active against the same strain of bacteria that the parents were infected with[4,5]. Conversely, it may be broad; for example, mealworm beetles primed with either fungi or a Gram-positive or Gram-negative bacteria induce immunity against Gram-positive pathogens[6]. Although the effectors that provide immunity in the progeny are mostly unknown, antimicrobial peptides are often upregulated in offspring[1,2,6].

Studies in the genetically tractable nematode *C. elegans* have enabled fundamental immune advances and shed light on many epigenetically inherited and stress-induced phenotypes[7–9]. As such, *C. elegans* has recently become a powerful model for the study of inherited innate immunity[10]. In this host, antiviral immunity, mediated by the small RNA mediated silencing of viral transcripts, was shown to last for at least three generations and be dependent on RNA interference (RNAi) pathways[11]. Although there are conflicting reports about whether the natural Orsay virus can induce heritable immunity in *C. elegans*, injection with vesicular stomatitis virus was able to protect progeny against reinfection[12–14]. Several studies have shown that learned bacterial avoidance can be transferred to progeny. In one case, heritable avoidance to *Pseudomonas aeruginosa* bacteria was dependent on RNAi pathways and a bacterial RNA used by *C. elegans* to inhibit host gene expression[15–17]. Parental exposure to pathogenic bacteria can

also protect offspring by increasing the likelihood of progeny adopting a stress-resistant dauer

phenotype[18]. Finally, resistance can be induced by upregulating immune genes in offspring. For example,

intergenerational immunity to pathogenic *Pseudomonas vranovensis* is dependent on the induced

expression of cysteine synthases[19,20].

Microsporidia are a large phylum of fungal-related parasites that infect most animal species[21]. These

pathogens can be lethal to immunocompromised humans, and can cause huge economic losses by

infecting agriculturally important animals such as honey bees, silk worms and shrimp[22]. *Nematocida*

parisii is a natural microsporidia parasite that commonly infects *C. elegans* in the wild. Infection of *C.*

elegans by *N. parisii* begins when microsporidia spores are ingested[23]. Spores inside the intestinal lumen

then fire a unique polar-tube structure which is used to transfer the cellular contents of the spore

(sporoplasm), including the parasite's genetic material into the host intestinal cell[24]. The pathogen then

replicates intracellularly to form meronts, and spreads from cell to cell by fusing the cells and forming

syncytia[25]. These meronts ultimately differentiate into mature spores which exit infected cells non-lytically,

and can result in the shedding of up to 200,000 spores during infection[23,26]. Microsporidia are pathogenic

to *C. elegans*, resulting in reduced fecundity and ultimately death[25,27].

Several mechanisms of innate immunity against microsporidia have been described in invertebrates.

Microsporidia infection typically induces a strong host transcriptional response, which often includes

upregulation of a suite of different antimicrobial peptides[28]. Though antimicrobial peptides are commonly

upregulated in other invertebrates, and *C. elegans* possess many different families of these proteins, so

far only C-type lectins have been shown to be upregulated upon *N. parisii* infection[29,30]. Instead, *C.*

elegans possesses a novel stress/immune pathway called the 'intracellular pathogen response' (IPR)

that is induced upon infection by both microsporidia and Orsay virus[29,31]. The IPR includes upregulation

of ubiquitin adapter proteins and is thought is to be involved in clearing intracellular parasites[29,32,33].

In this study, we reveal robust resistance to microsporidia in the offspring of *N. parisii*-infected *C. elegans*. We show that inherited immunity dramatically reduces host-cell invasion by *N. parisii* and also confers resistance to the bacterial pathogen *P. aeruginosa*. We find that immunity is conferred in a dose-dependent manner, lasts throughout development, and is maintained for a single generation. Using tissue-specific depletion or expression of negative regulators of the IPR (LIN-35 and PALS-22), we demonstrate that immunity in progeny is dependent on a parental transcriptional response, and that this immunity is transferred from maternal somatic tissues to the progeny. Remarkably, the host transcriptional response and thus inherited immunity, can be induced by both biotic and abiotic stresses which mimic the microsporidial response. Together, these results provide insight into how inherited immune responses can be induced to provide immunity against pathogenic infection.

2. Results

2.1. Parental infection by *N. parisii* confers immunity to *C. elegans* progeny in a dose-dependent manner

Infection of *C. elegans* with the natural microsporidian pathogen *N. parisii* delays development and impacts worm fertility[24,34]. To quantify the effects of infection, we exposed early larval stage (L1) animals to varying doses of *N. parisii* spores. At 72 hours post infection (hpi), animals were fixed and stained with the chitin-binding dye DY96 (Figures S1A and S1B). DY96 stains both microsporidia spores and worm embryos, allowing us to determine both parasite burden and host reproductive development. Worms infected with higher doses of spores exhibited greater parasite burden and a smaller body size at 72 hpi (Figures S1C and S1D). Higher infection doses also correlated with a reduction in the percentage of gravid adults (animals carrying embryos), as well as a reduction in the number of embryos per animal (Figures S1E and S1F). In agreement with these findings, infection of transgenic animals expressing a fluorescent protein in the germline revealed germline deformities in worms exposed to high doses of spores (Figure S1G).

To determine the effects of parental infection on offspring, we infected parental (P0) generations at the

L1 stage with a 'very low', 'low' or 'moderate' dose of *N. parisii* that resulted in a ~10-50% reduction in

the number of gravid adults (Figure S1E). At 72 hpi, infected adults were treated with a sodium

hypochlorite solution to release the F1 embryos from adults and destroy microsporidia spores. As *N.*

parisii infection is not vertically transmitted, resultant larvae are not infected with microsporidia (Figure

S2)[27]. F1 generations of synchronized L1s were then exposed to a high dose of spores. Remarkably,

imaging revealed a quantifiable reduction in the parasite burden of primed worms from infected parents,

as compared to naïve worms from uninfected parents (Figures 1A and 1B). Immunity in F1 progeny was

dose-dependent; parents experiencing a higher infection burden gave rise to offspring that were more

resistant to *N. parisii*. In agreement with our data for parasite burden, we saw that primed worms under

infection conditions were larger and produced significantly more embryos than their naïve counterparts

(Figures 1C-1E). Primed worms showed no fitness advantage under non-infection conditions and were

indeed smaller and contained fewer embryos than worms from uninfected parents (Figures S3A and

S3B). Together, these data reveal a robust and dose-dependent immunity phenotype in the offspring of

N. parisii-infected worms.

2.2 Inherited immunity prevents microsporidia invasion events by restricting spores in the

intestine

To resist microsporidia infection, animals may (i) prevent invasion of intestinal cells or (ii) destroy the

invaded pathogen[24,33]. To determine by which mechanism inherited immunity confers resistance to *N.*

parisii, we first performed invasion assays. Naïve or primed L1 worms were exposed to a high dose of

spores and fixed 30 minutes post infection (mpi) or 3 hpi. To visualize invasion events in these animals,

we performed fluorescence *in situ* hybridization (FISH) to detect *N. parisii* 18S ribosomal RNA in host

intestinal cells, thus indicating the presence of intracellular sporoplasms. Our analysis revealed

significantly fewer invasion events in primed animals at both time points, including an over 98% reduction

in infectious events at 30 mpi (Figures 2A, 2B and S4A). To determine if spores were present in the

intestinal lumen, animals were also stained with DY96. Strikingly, we also observed far fewer spores in

the guts of primed animals (Figures 2C, 2D and S4B).

To test whether a reduction of spores in our primed worms was a result of reduced feeding, we allowed

naïve and primed L1 stage worms to feed on fluorescent beads. After 30 min or 3 h, we fixed the

populations and quantified fluorescence in individual animals. We failed to detect any reduction in feeding

in primed worms at either time point (Figures 2E and S4C). At the 3 h time point, primed animals actually

fed significantly more than naïve worms, indicating that microsporidia invasion results in reduced feeding

in *C. elegans* (Figure S4C).

To test whether enhanced clearance of intracellular infection might also contribute to microsporidia

resistance in primed animals, we performed pulse-chase experiments[24]. Here, naïve and primed animals

were maintained in the presence of high concentrations of *N. parisii* spores for 3 h, before washing

thoroughly to remove any microsporidia spores not inside the animals. The population was then split; half

was immediately fixed to represent the initial infection, and the other half maintained in the absence of

spores until a 24 h end point when they were fixed to assess clearance. Detection of sporoplasms using

18S RNA FISH and subsequent quantifications showed no evidence of intracellular pathogen clearance

in the primed animals (Figure S4D). These data suggest that limiting invasion is the principal way by

which inherited immunity provides protection against microsporidia.

To provide insight into the mechanisms underlying protection in primed animals, we next tested the

specificity of the immune response. For this, we assayed resistance to the extracellular Gram-negative

bacterial pathogen *Pseudomonas aeruginosa* (Strain PA14) using a well-established 'slow killing'

protocol[7]. Here, naïve or *N. parisii*-primed animals were plated on wild-type PA14 at the L1 stage and

survival assayed at 84 hpi. Remarkably, primed worms were significantly less susceptible to *P.*

aeruginosa infection than naïve animals (Figure 2F). Slow killing by *P. aeruginosa* occurs as a result of

bacterial accumulation in the gut[35]. To visualize bacterial burden in our naïve and primed worms, we

performed infection assays using dsRed-expressing PA14. At 48 hpi, we fixed and quantified fluorescence in our naïve and primed populations. In agreement with the data from survival assays, the bacterial burden in microsporidia-primed animals was significantly reduced (Figures 2G, 2H and S4E). Together, these results suggest that a host intestinal factor may be protecting primed worms from infection by destroying microsporidia spores and other pathogenic microbes within the intestinal lumen.

To test whether inherited immunity could protect against all classes of pathogen, we tested the response of *N. parisii*-primed animals to the intracellular intestinal pathogen Orsay virus. Surprisingly, FISH staining revealed that primed worms were similarly susceptible to Orsay virus as their naïve counterparts (Figure 2I). This result is particularly notable given the significant overlap between the transcriptional responses induced by Orsay virus and *N. parisii*, with both activating IPR genes[36]. PALS-22 is a negative regulator of the IPR and a loss-of-function mutation in *pals-22* results in animals with a constitutively activated IPR response that are resistant to viral and microsporidia infection, but not to *P. aeruginosa* (Figures 2I and S4F)[32]. Mutants deficient for *pals-22* limit microsporidia infection by preventing invasion as well as clearing invaded parasites (Figure S4F). Thus, IPR-mediated immunity is distinct in several ways from the immunity found in primed animals.

2.3 Inherited immunity to *N. parisii* lasts a single generation and persists throughout development

We next sought to understand the kinetics of the inherited immune response to microsporidia infection. To determine whether immunity could be transmitted over multiple generations, we tested resistance to microsporidia in both the F1 and F2 progeny of *N. parisii*-infected worms. We observed resistance to microsporidia only in the F1 population, indicating that this response lasts for a single generation (Figures 3A and 3B, S5A and S5B). To test the longevity of the inherited immune response within the F1 generation, naïve and primed worms were infected with *N. parisii* either at the L1 stage, 24 h later at the L2/L3 stage, or 48 h later at the L4 stage. After 30 mpi, worms were fixed, stained with a FISH probe to detect *N. parisii*, and the number of sporoplasms per individual quantified. While immune-primed worms

continued to show some resistance to infection at the L4 stage, resistance was strongest at the L1 stage

of infection (Figure 3C).

We next tested whether a greater ancestral history of infection might enhance immunity phenotypes and

potentially enable the immune response to transmit over multiple generations. Animals were infected for

one, two, or three sequential generations with *N. parisii* (P0s, F1s and F2s). Susceptibility to microsporidia

was assessed in F3 animals, as well as in F4 animals following a single generation of rest without infection

(Figures S5C-E). We found that an increased history of ancestral infections did not enhance immunity

phenotypes among the F3 populations (Figures 3D and 3E). Furthermore, resistance was seen

exclusively in the F3 animals and no resistance was observed in the F4 generations (Figures S5F and

S5G).

We next wanted to determine for how long a parent must be infected with *N. parisii* in order to transmit

immunity to progeny. For this, parental generations were infected as previously at the L1 stage for 72 h

of total infection, or rested on their typical *E. coli* food source and subsequently infected as L2/L3s for 48

h total infection, or as L4s for 24 h total infection. Infection assays on resultant F1 progenies showed that

parental worms infected only briefly as L4s were still able to confer immunity phenotypes to offspring,

though to a lesser extent than animals from parents infected for longer periods (Figures 3F and 3G). As

N. parisii takes over 48 hours to sporulate[25], these results indicate that the early stages of microsporidian

infection alone (i.e. invasion and replication) are sufficient to induce immunity in progeny.

Together, these data show that inherited immunity protects the F1 progeny from infected parents

throughout development and supports a model in which the immediate parental environment is the most

important factor in determining the immune competency of offspring.

2.4 The parental transcriptional response to infection triggers inherited immunity in offspring

N. parisii is an intestinal parasite that does not come into direct contact with the *C. elegans* germline, and

how information is transferred from the soma to the germline is not known (Figure S6A). To explore this,

we tested whether mutants defective in small-RNA inheritance and histone modification were able to

transmit immunity phenotypes to offspring. We found that the offspring of *N. parisii* infected mutants still

became protected against infection (Figure S7A-D)[19,37]. In addition, we found that the master immune

regulator PMK-1 was not required for the induction of immunity, consistent with this pathway not being

involved in immunity to *N. parisii* (Figure S7C-D)[27]. These data indicate that the small RNA, histone

modification, and PMK-1 pathways are not required for transmission of immunity.

To determine whether infection itself, or merely exposure to spores, is required for the transmission of

inherited immunity from parent to progeny, we used heat-killed spores. Parent populations were either

uninfected, exposed to live *N. parisii*, or exposed to heat-killed spores. Unlike live spores, heat-killed

spores fail to induce the IPR response, as demonstrated using transcriptional reporters for key IPR genes

(Figures S6B and S6C) [36]. Consistent with a requirement for infection and induction of the IPR to initiate

inherited immunity, the offspring of parents exposed to heat-killed spores showed no enhanced protection

against *N. parisii* infection (Figures 4A and 4B).

We next tested whether other environmental conditions that induce a similar transcriptional response as

microsporidia infection could also induce immunity in progeny. When analysing mRNA sequencing data,

we confirmed a previously noted similarity to the Orsay virus response, and also noticed that the *C.*

elegans response to *N. parisii* infection overlaps significantly with that of worms exposed to the heavy

metal cadmium (Figure S6D).

To explore a role for the transcriptional response in inducing inherited immunity, we first performed *N.*

parisii infection assays on the offspring of untreated parents, or parents exposed to either *N. parisii* or

Orsay virus. We saw that the Orsay-primed F1 progeny of loss-of-function *rde-1* mutants, but not N2

worms, showed significantly reduced infection and improved fitness under microsporidia infection

conditions, as compared to naïve controls (Figures 4C and D). This difference between the *rde-1*

genotype and N2 can be attributed to the increased susceptibility of *rde-1* mutants to Orsay virus, allowing

the P0s to have an enhanced viral response (Figure S6E)[38].

We next assayed the susceptibility of cadmium-primed animals to *N. parisii* infection. Strikingly,

resistance to microsporidia was observed in the offspring of parents exposed to cadmium (Figures 4E

and 4F). Additionally, cadmium-primed animals exposed to *P. aeruginosa* had significantly lower bacterial

burden than naïve animals, supporting of a role for the transcriptional response in transmitting immunity

against both these classes of pathogens (Figure 4G). Conversely, both infection-primed and cadmium-

primed animals were less fit and displayed reduced gravidity when faced with a high concentration of

cadmium (Figure S6F). Similarly, we found that while the progeny of osmotically stressed parents were

better able to cope with a high salt stress, the offspring of *N. parisii*-infected animals were less viable

under this stress condition (Figure 4H). These data indicate that primed animals are not simply more

resistant to any given stress, but specifically to pathogenic *N. parisii* and *P. aeruginosa*. Furthermore,

these data support a role for the transcriptional response in transmitting inherited immunity to offspring,

and highlight the complex relationships that underlie responses to abiotic and biotic stresses across

generations.

2.5. Progeny of animals with an artificially-activated transcriptional response are resistant to

infection

In addition to the previously characterized *pals-22* mutant, we also observed that mutants of *lin-35*, the

C. elegans ortholog of retinoblastoma protein (RB), induce a similar transcriptional response to

microsporidia infection (Figure S6D)[39,40]. These mutants share extensive similarity with the transcriptional

response to *N. parisii*, with 242 shared genes upregulated at least two-fold in both *lin-35* and *pals-22* as

well as *N. parisii* infected worms (Figure S8). To determine whether activating this host response in the

absence of infection or environmental stimuli would induce immunity, we performed infection assays in

pals-22 and *lin-35* mutants. Both *pals-22* and *lin-35* mutants have defects resulting in fewer embryos,

even when grown under normal conditions (Figure 5A). Thus, the number of embryos per worm in these

mutants during infection conditions is similar to wild type. However, in agreement with a role for the IPR

response in restricting infection, both *pals-22* and *lin-35* mutants had a significantly lower parasite burden

than wild-type animals (Figure 5B).

Next, to determine if transcriptional response activation in these mutants could induce immunity in

progeny, we performed mating assays. We first examined the cross-progeny of *pals-22* and *lin-35* mutant

hermaphrodites mated with wild-type males to determine if parents with an upregulated transcriptional

response generated resistant F1s (Figure 5C). The heterozygous cross-progeny of both mutants

produced a similar number of embryos in uninfected conditions as wild-type, as the developmental timing

and brood size are recessive traits of *pals-22* and *lin-35* mutants. When infected with *N. parisii*, cross-

progeny produced more embryos and have a lower pathogen load than wild-type animals (Figures 5D

and 5E). On average, 57% of *pals-22* and 77% of *lin-35* maternal cross-progeny became gravid under

infection conditions, compared to less than 10% of wild-type animals (Figures 5F and 5G). In contrast,

paternal cross-progeny i.e. the offspring of wild-type hermaphrodites mated with males carrying either

pals-22 or *lin-35* mutations, do not exhibit improved reproductive fitness under infection conditions

(Figures 5F and 5G). These results reveal that inherited immunity to microsporidia is maternally

transferred and can be induced by transcriptional activation alone.

2.6. The signal for offspring resistance can originate in multiple somatic tissues

To identify the tissues required to induce the transcriptional response and thereby transmit inherited

immunity, we used two methods. First, we infected the maternal cross-progeny of *pals-22* mutants

carrying a transgene for wild-type *pals-22* under an endogenous promoter, or under a promoter specific

for a single tissue type where *pals-22* is typically expressed[41]. While offspring of the *pals-22* endogenous

rescue strain were no longer resistant to infection, neuronal and hypodermal-specific rescue strains still

produced resistant progeny, but to a lesser extent than *pals-22* mutants (Figure 6A). This indicates that

a signal in the neuronal and hypodermal tissues may contribute to the transmission of inherited immunity,

but may not be crucial. The intestinal-specific rescue produced progeny that were less resistant than

pals-22 mutants, though this was not statistically significant, indicating that an intestinal signal could also contribute to transmission of immunity. The maternal cross-progeny of *lin-35* mutants with wild-type *lin-35* expressed under either an endogenous or intestine-specific promoter were no longer resistant to *N. parisii*, indicating that an intestinal signal was important for transmission of immunity in this case. Furthermore, addition of *lin-35* under the *pie-1* germline promoter was seen to partially reduce resistance (Figure 6B).

In a second approach, we used the auxin inducible degradation (AID) system[42] to degrade either PALS-22 or LIN-35 in a tissue-specific manner in the parental generation only, and assessed F1 progenies for resistance to microsporidia (Figure S9). Targeted degradation of PALS-22, and thus signal induction, in the adult intestine alone was sufficient to enhance immunity in progeny, while degradation in any other single tissue was unable to provide protection (Figure 6C). Additionally, degradation of LIN-35 in somatic tissues only, was sufficient to confer resistance to offspring (Figure 6D). Taken together, these results suggest that induced expression of the IPR either in the adult intestine only, or in multiple somatic tissues is sufficient to transmit inherited immunity to offspring.

3. Discussion

Inherited immunity is a nascent and rapidly growing field of research, with important consequences for our understanding of health and evolution. Multiple studies have now demonstrated that parental exposure to one pathogen can protect offspring against subsequent exposure to the same pathogen[1,13,19,20]. Several studies have also shown that parents exposed to a particular stress can produce progeny that are protected against the same stress, and that abiotic stress can provide transgenerational resistance to pathogenic infection[5,6,43,44]. Critically, it is not yet clear how the environment of the parent determines which stresses the offspring are protected against. Here, we show that inherited immunity to microsporidia can be activated by microsporidia and viral infection, as well as heavy metal stress, which all elicit a similar transcriptional response in the parents. Inherited immunity can also be activated through maternal somatic depletion of negative regulators of the transcriptional

response. Indeed, activation of the transcriptional response within just a single generation in the intestine

is sufficient to induce inherited immunity in offspring. Taken together, we have demonstrated that the

signal for initiating inherited immunity against microsporidia is somatic induction of a shared maternal

transcriptional stress response.

Inherited immunity phenotypes are often costly and can result in primed animals being more sensitive to

other stresses[1,45]. Populations of *C. elegans* often develop within the same immediate environment, so

parental infection is a good indicator of progeny exposure to microsporidia[46]. Here we find that progeny

from microsporidia infected parents are smaller, carry fewer embryos, and are more sensitive to osmotic

and heavy metal stress. While inherited immunity in response to microsporidia infection lasts only a single

generation, it can be induced in every generation, and this may be a strategy to limit fitness trade-offs.

Immunity to microsporidia in *C. elegans* is thought to function through activation of the IPR and associated

pathogen clearance[33,36]. Upregulation of the IPR results in immunity to both microsporidia and virus, but

not bacteria[32]. Here, inherited immunity protects against microsporidia and bacteria, but not virus.

Additionally, constant induction of the IPR genes in *pals-22* and *lin-35* mutants also greatly impacts

animal development, whereas offspring with activated inherited immunity have less severe growth

defects[32,47]. Our results suggest that although induction of the IPR is sufficient to activate inherited

immunity, inherited immunity and IPR-mediated immunity are separate immune responses. We also

show that inhibition of the RB ortholog *lin-35* provides immunity against microsporidia, both for the parents

and their offspring. Although *lin-35* has been implicated in stress responses and negatively regulation of

immune gene expression, this is the first example of *lin-35* mutants providing pathogen resistance[48,49].

RB is evolutionarily conserved, but in mammals acts as a positive regulator of antiviral immunity and

immune cell development[50,51].

Once the signal for inherited immunity is induced in the soma, the response must be transferred to

developing progeny. Several inherited multigenerational responses that last for more than two

generations are dependent on RNAi pathways[16,52]. Consistent with recent reports of responses that last

only one or two generations being transmitted independent of these pathways, heritable immunity to

microsporidia (which lasts a single generation) is not reliant on RNAi machinery[19,37].

Being able to harness inherited immunity would provide a way to prevent infection of invertebrates[53].

Inherited immunity can be induced without the pathogen itself, by molecules that activate an immune

response, and thus a similar approach could be used to combat microsporidia infections[54–56]. Inherited

immunity could be employed to block infection of beneficial insects such as honey bees, and inhibition of

this immunity could be used to improve the efficacy of microsporidia as biocontrol agents for locusts and

other pests[57]. Although this is the first report of inherited immunity to microsporidia, mosquitoes whose

parents were infected with microsporidia contained significantly fewer malaria parasites. This suggests

that manipulation of these pathways could also be used to prevent invertebrate vectors of human disease

from transmitting infection[58,59].

Materials and Methods

4.1. Worm maintenance

C. elegans strains were maintained at 21°C on nematode growth media (NGM) plates seeded with 10X

OP50-1 *Escherichia coli*, as previously described (Brenner, 1974). Strains used in this study are listed in

Supplemental Table 1. For maintenance and infection assays, 10X concentrates of OP50-1 were

prepared by growing cultures to saturation in lysogeny broth (LB) at 37°C for 16-18 h. Populations were

synchronized by washing worms off plates with M9 solution and bleaching with sodium hypochlorite / 1M

NaOH until the embryos of gravid adults were released into solution. Eggs were washed three times with

M9, resuspended in 5 ml M9, and rotated at 21°C for 18-24 h to allow embryos to hatch into L1s. For

pelleting of live worms, animals were centrifuged in microcentrifuge tubes for 30 s at 1400xg.

Supplemental Table 1. *C. elegans* strains used in this study

Strain name	Genotype	Source
N2	Wild-type, Bristol strain	Caenorhabditis Genetics Center (CGC)
JMC101	*csr-1(tor67[csr-1 exon2::GFP::Flag IV:7958598])IV* *csr-1(tor67[gfp::3xflag::csr-1]) IV*	(Ouyang et al., 2019)
DA465	*eat-2(ad465)* II	(Raizen et al., 1995)
ERT054	*jyIs8[C17H1.6p::GFP, myo-2::mCherry]* X	(Bakowski et al., 2014)
ERT071	*jyIs14[F26F2.1p::GFP, myo-2::mCherry]*	(Bakowski et al., 2014)
ERT356	*pals-22(jy1)* III	(Reddy et al., 2017)
WM27	*rde-1(ne219)* V	CGC
MT10430	*lin-35(n745)* I	CGC
ERT304	*pals-22(jy1) III; jyIs8[C17H1.6p::GFP, myo-2p::mCherry]* X	(Reddy et al., 2017)
AWR33	*lin-35(n745)* I; *jyIs8[C17H1.6p::GFP, myo-2::mCherry]* X	This study
ERT526	*jySi37[pals-22p::pals-22::GFP, unc-119 (+)]* II; *pals-22(jy1) unc-119(ed3)* III	(Reddy et al., 2017)
ERT527	*jySi38[vha-6p::pals-22::GFP, unc-119 (+)]* II; *pals-22(jy1) unc-119(ed3)* III	(Reddy et al., 2017)
ERT534	*jySi39[unc-119p::pals-22::GFP, unc-119 (+)]* II; *pals-22(jy1) unc-119(ed3)* III	(Reddy et al., 2017)
ERT539	*jySi41[dpy-7p::pals-22::GFP, unc-119(+)]* II; *pals-22(jy1) unc-119(ed3)* III	(Reddy et al., 2017)
AWR35	*lin-35(n745)* I; *unc-119(ed3)* III; *vrIs55[ges-1p::lin-35::GFP::FLAG::lin-35 3'UTR]*	This study
AWR36	*lin-35(n745)* I; *unc-119(ed3)* III; *vrIs60[lin-35p::lin-35::GFP::FLAG::lin-35 3'UTR]*	This study
AWR37	*lin-35(n745)* I; *unc-119(ed3)* III; *vrIs93[mex-5p::lin-35::GFP::FLAG::lin-35 3'UTR]*	This study
AWR38	*lin-35(n745)* I; *unc-119(ed3)* III; *vrIs56[pie-1p::lin-35::GFP::FLAG::lin-35 3'UTR]*	This study
AWR45	*pals-22(kea8[pals-22::GFP::degron])* III	This study
AWR59	*keaSi10[rpl-28p::TIR1::mRuby::unc-54 3'UTR +Cbr-unc-119(+)]* II; *pals-22(kea8[pals-22::GFP::degron])* III	This study
AWR61	*keaSi11[vit-2p::TIR1::mRuby::unc-54 3'UTR +Cbr-unc-119(+)]* II; *pals-22(kea8[pals-22::GFP::degron])* III	This study
AWR64	*kea15[rgef-1p::TIR1::mRuby::unc-54 3'UTR + Cbr-unc-119(+)]* II; *pals-22(kea8[pals-22::GFP::degron])* III	This study

AWR62	*keaSi9[myo-3p::TIR1::mRuby::unc-54 3'UTR +Cbr-unc-119(+)]* II; *pals-22(kea8[pals-22::GFP::degron])* III	This study
AWR63	*keaSi12[dpy-7p::TIR1::mRuby::unc-54 3'UTR + Cbr-unc-119(+)]* II; *pals-22(kea8[pals-22::GFP::degron])* III	This study
AWR41	*lin-35(kea7[lin-35p::degron::GFP::lin-35])* I	This study
AWR54	*lin-35(kea7[lin-35p::degron::GFP::lin-35])* I; *ieSi57[eft-3p::TIR1::mRuby::unc-54 3'UTR + Cbr-unc-119(+)]* II	This study
AWR56	*lin-35(kea7[lin-35p::degron::GFP::lin-35])* I; *ieSi64[gld-1p::TIR1::mRuby::gld-1 3'UTR + Cbr-unc-119(+)]* II	This study
YY538	*hrde-1(tm1200)* III	CGC
WM156	*R04A9.2(tm1116)* X	CGC
YY1083	*wago-4(tm1019)* II	(Xu et al. 2018)
WM160	*sago-1(tm1195)* V	CGC
WM154	*sago-2(tm894)* I	CGC
Unnamed	*ppw-1 (tm914)* I	(The C. elegans Deletion Mutant Consortium et al., 2012)
NL2098	*rrf-1(pk1417)* I	CGC
MT17463	*set-25(n5021)* III	CGC
AU0066	*pmk-1(km25)* IV	CGC

4.2. Construction of transgenic strains

For strains with tissue-specific expression of *lin-35* in a *lin-35* mutant background, MT10430 was crossed

to DP38. The resulting *lin-35(n745)* I; *unc-119(ed3)* III double mutant was then crossed to YL398, YL402,

YL409, and YL468[47]

SapTrap was used to construct a repair template for introducing the auxin inducible degron tag via

CRISPR[60]. Briefly, the ~500-600 bp regions immediately upstream of the *pals-22* start codon or

downstream of the *lin-35* stop codon were PCR amplified from genomic N2 DNA to act as homology

arms. These were cloned into the pDD379 backbone, along with sgRNA and other SapTrap-SEC kit

plasmids (Addgene)[61]. The repair construct and co-injection markers were microinjected into N2 worms

and hygromycin resistant rollers were selected. The SEC was then excised via heat shock[62].

Strains carrying *vit-2p*::TIR1, *rgef-1p*::TIR1, *myo-3p*::TIR1, *dpy-7p*::TIR1, and *rpl-28p*::TIR1 were

generated by MosSCI[63]. Repair constructs were generated using Multisite LR Gateway cloning using

pCFJ150 as LR entry vector and microinjected into strain EG6699.

All strains generated for this study were outcrossed to N2 at least three times before use. Plasmids and

primers used in this study are listed in Supplemental Table 2.

4.3. Preparation of microsporidia spores

N. parisii spores were prepared as described previously[26]. In brief, microsporidia spores were used to

infect large populations of *C. elegans* N2 worms. Infected worms were then harvested, mechanically

disrupted using 1 mm diameter Zirconia beads (BioSpec), and the lysate filtered through 5 μm filters

(Millipore Sigma™) to remove nematode debris. Spores preparations were assayed for bacterial growth

and those that were free of contaminating bacteria were stored at -80°C. Each assay was performed

using spores of the same batch. In total, five batches were employed in this study.

4.4. Microsporidia infection assays

4.4.1. Basic infection and priming assays

Synchronized populations of L1 worms were mixed with 1 ml of 10X OP50-1 alone as uninfected controls,

or 1 ml of 10X OP50-1 supplemented with *N. parisii* spores, and plated on 10 cm NGM plates (doses

defined in Supplemental Table 3 below). For priming assays, 2,500 P0 animals were infected with a 'low

dose' of *N. parisii* such that >90% of animals were infected at 72 hpi and >80% of animals were fertile in

order to generate a sufficient yield of primed F1 embryos for subsequent experiments (Figure S1). At 72

hpi, worms were collected and washed three times in 1 ml M9. To measure infection in the parental

generation, 10% of P0 worms were fixed and parasite burden and gravidity assessed. The remaining

90% of worms were bleached to obtain naïve or primed embryos from uninfected or infected adults,

respectively.

Following hatching in M9, 1,000 naïve or primed F1 animals were challenged at the L1 stage on a 6 cm

plate with a high dose of *N. parisii*. This resulted in ~10-20% of naïve animals being gravid, so that fitness

increases or decreases would be easily detectable. At 72 hpi, worms were fixed, stained with DY96, and

gravidity and infection assessed by microscopy.

Supplemental Table 3. Doses of *N. parisii* used in this study

N. parisii dose	Plate concentration (spores/cm^2)	Millions of spores per 6 cm plate
Very low	17,700	0.5
Low	35,400	1.0
Moderate	70,700	2.0
High	88,400	2.5
Very high	106,000	3.0
Maximal	212,000	6.0

4.4.2. Assessment of parental infection duration on inherited immunity

To test how long animals must be infected to transmit immunity to offspring, 2,500 P0 animals were either

not infected or infected with a low dose of *N. parisii* at (i) the L1 stage, (ii) the L2/L3 stage after 24 h rest

on 10X OP50-1 or (iii) the L4 stage after 48 h rest on 10X OP50-1. At 72 h post L1, P0 populations were

bleached to release F1 embryos. Naïve and primed F1 animals were tested for inherited immunity as

described in **Section 4.4.1**.

4.4.3. Assessment of inherited immunity through development

To test inherited immunity throughout development, naïve or primed worms were obtained as described

in **Section 4.4.1**. Next, 1,000 naïve or primed animals were plated on 10X OP50-1 supplemented with a

maximal dose of *N. parisii* spores at (i) the L1 stage, (ii) the L2/L3 stage after 24 h rest on 10X OP50-1

or (iii) the L4 stage after 48 h rest on 10X OP50-1. At 30 mpi, animals were fixed, stained with MicroB

FISH probe to detect *N. parisii* 18S RNA, and the number of sporoplasms quantified by microscopy.

4.4.4. Assessment of spores in the gut and host cell invasion by *N. parisii*

To assay host cell invasion by microsporidia and the presence of spores in the gut, we conducted short

infection assays [24]. Naïve or primed F1 animals were obtained as described in **Section 4.4.1**. Next, 1,000

naïve or primed animals were mixed with 10X OP50-1 and a maximal dose of *N. parisii* spores at the L1

stage and plated on NGM. At 30 mpi, animals were fixed and stained with DY96 and MicroB FISH probe,

and the number of spores or sporoplasms quantified by microscopy.

4.4.5. Assessment of host clearance of *N. parisii*

To assay host clearance of microsporidia, we conducted pulse infection assays. For this, naïve or primed

F1 animals were obtained as described in **Section 4.4.1**. Next, 2,000 naïve or primed animals were mixed

with 10X OP50-1 and a very high dose of *N. parisii* spores at the L1 stage and plated on NGM. At 3 hpi,

populations were split and half the animals fixed. The remaining animals were maintained in the absence

of spores until a 24 h end point before fixing. Animals were stained with MicroB FISH probe, and the

number of sporoplasms at each time point quantified by microscopy.

4.4.6. Assessment of *N. parisii*-induced immunity over multiple generations

To assess immunity to microsporidia over multiple generations, we performed an ancestral infection

assay (Figure S5C). Here, 2,500 animals were infected at the L1 stage with *N. parisii* for one, two or three

successive generations (P0, F1, F2). Each infection period lasted 72 h before treating with sodium

hypochlorite solution to obtain the next generation of embryos. F3 L1 populations were split and either

subject to infection testing or maintained under non-infection conditions for the collection and subsequent

testing of F4 offspring. To test immunity, F3 and F4 larvae were exposed to a high dose of *N. parisii* at

the L1 stage. At 72 hpi, F3 and F4 animals were fixed and stained with DY96 to visualize *N. parisii* spores.

4.5. Heat-killed spore infection assay

Synchronized populations of 2,500 L1 worms were mixed with (i) 10X OP50-1 alone, or 10X OP50-1

supplemented with a low dose of either (ii) live or (iii) heat-killed *N. parisii* spores, and plated on NGM.

For heat killing, live spores were treated at 65°C for 10 min. At 72 hpi, parental generations were bleached

to obtain naïve embryos or embryos primed with heat-killed or live spores. Inherited immunity to *N. parisii*

was tested as described in **Section 4.4.1**.

4.6. *P. aeruginosa* infection assays

Strains of wild-type or dsRed-expressing *P. aeruginosa* PA14 were used to assay survival and bacterial

burden, respectively. Strains were grown in 3 ml LB overnight from a single bacterial colony. A volume

of 20 µl of bacterial culture was used to seed 3.5 cm slow-killing plates for survival assays[35]. A volume of

50 µl of bacterial culture was used to seed 6 cm slow-killing plates for bacterial burden assays. To prevent

C. elegans pathogen avoidance behaviour [64], bacterial culture was spread to ensure plates were fully

covered with *P. aeruginosa*. Seeded plates were incubated for 24 h at 37°C for growth of bacterial lawns

and maintained for a further 24 h at room temperature (RT) prior to infection assays. Naïve and primed

worms were obtained as in **Section 4.4.1**. To assay survival, 20-30 naïve or primed L1 worms were

transferred to 3.5 cm wild-type PA14-seeded plates and incubated at 25°C for 84 h. Worms that failed to

respond to pressure from a metal pick were considered non-viable. To assay bacterial burden, 1,000

naïve or primed L1 worms were plated on 6 cm dsRed PA14-seeded plates, and incubated at 25°C for

48 h. At 48 hpi, worms were collected from plates, washed three times in M9, and fixed and bacterial

burden assessed by microscopy.

4.7. Orsay virus infection assays

Orsay virus filtrate was prepared as described previously[29]. In brief, plates of Orsay virus infected animals

were maintained until starvation. Virus shed by infected worms was collected by washing plates with M9,

passing through 0.22 μm filters (Millipore Sigma™), and stored at -80°C. For infections to test resistance

to Orsay virus, naïve and primed worms were obtained as in **Section 4.4.1.** Next, 1,000 naïve or primed

L1s were mixed with 100 μl of 10X OP50-1 and 500 μl of the viral filtrate, then plated on 6 cm NGM. At

16 hpi, animals were fixed and FISH stained to assess infection status.

To generate Orsay-primed animals, 2,500 L1s were mixed with 1 ml of 10x OP50-1 and 500 μl of viral

filtrate. At 72 hpi, animals were collected and bleached to obtain primed embryos. Inherited immunity to

N. parisii was tested as described in **Section 4.4.1**.

4.8. Bead-feeding assay

Naïve or primed L1 worms were obtained as described in **Section 4.4.1**. Next,1,000 animals were placed

on 6 cm plates in 400 μl total volume of M9 containing 10% (v/v) 10X OP50-1 and 4% (v/v) 0.2 μm green

fluorescent polystyrene beads (Degradex Phosphorex). Where spores were included for the 3 h time

point, a maximal dose of *N. parisii* spores was used. After 30 min or 3 h, animals were fixed, and bead

ingestion assessed by microscopy.

4.9. Osmotic stress assays

Osmotic stress adaptation assays were performed as previously described[65]. Synchronized populations

of 2,500 L1 worms were plated on either standard NGM (50 mM salt) plates together with (i) 10X OP50-

1 alone or (ii) 10X OP50-1 supplemented with a low dose of *N. parisii*, or (iii) on NGM containing an

elevated concentration of salt (250 mM) together with 10X OP50-1. At 80 hpi, parent populations were

bleached to obtain naïve, infection-primed or salt-primed embryos. To test resistance to osmotic stress,

1,000 embryos were plated on NGM containing a high concentration of salt (420 mM). The percentage

of embryos that had hatched by 48 h was quantified using light microscopy.

4.10. Cadmium assays

We first confirmed that cadmium exposure induced IPR gene expression in *C. elegans*. This was determined by plating ERT054 and ERT071 fluorescent reporter worm strains on 50 mM calcium chloride and observing increased GFP expression in exposed worms, relative to unexposed controls. Synchronized populations of 2,500 L4 worms were plated on either (i) standard NGM together with 10X OP50-1 alone or (ii) 10X OP50-1 supplemented with a low dose of *N. parisii*, or (iii) NGM containing 50 mM cadmium chloride together with 10X OP50-1. After 24 h, parent populations were bleached to obtain naïve, infection-primed or cadmium-primed embryos. Immunity to *N. parisii* was tested as described in **Section 4.4.1.** Immunity to *P. aeruginosa* was tested as described in **Section 4.6**. To test for protection against cadmium stress, 1,000 F1 larvae were maintained on standard NGM with 10X OP50-1 until L4 stage, then plated on NGM containing 50 mM cadmium chloride. After 24 h, animals were fixed and stained with DY96 for embryo counting.

4.11. Auxin-inducible depletion experiments

Auxin plates were prepared by adding auxin stock solution (400 mM auxin [Alfa Aesar] in ethanol) to NGM, for a final concentration of 200 uM auxin, immediately before pouring plates. Control plates were prepared by adding ethanol to NGM, for a final concentration of 0.15%. Auxin plates were stored in the dark at 4°C and used within one month.

Embryos obtained from bleaching gravid adults were plated on auxin or ethanol control plates following M9 washes. 10X OP50-1 was added to plates 18-24 h after plating to allow embryos to hatch and synchronize. Worms were bleached 72 h post L1 arrest and F1 immunity to *N. parisii* tested as described in **Section 4.4.1**.

4.12. Cross-progeny generation

For testing maternal effects, 50 L4 ERT054 males and 30 L4 hermaphrodites were plated on a 3.5 cm NGM plate for mating. To test paternal effects, L4 mutant males carrying the *jyIs8* transgene were

crossed with L4 N2 hermaphrodites (Figure 5C). After 22-24 h, animals were bleached to obtain embryos

and F1 immunity to *N. parisii* tested as described in **Section 4.4.1**. During quantification, the presence of

myo-2p::mCherry was used to distinguish cross-progeny from self-progeny.

4.13. Fixation

For visualization of germlines of JMC101 (GFP::3xFLAG::CSR-1) animals, bead-feeding assays, and

staining of *N. parisii* or Orsay virus with FISH probes, worms were fixed in 1 ml 4% paraformaldehyde

(PFA) in phosphate buffered saline (PBS) containing 0.1% Tween-20 (PBST), for 20 min at RT or

overnight at -20°C. For *P. aeruginosa* burden assays and DY96 staining, worms were fixed in 1 ml

acetone for 10 min at RT, or overnight at 4°C. For pelleting of worms during fixing protocols, animals

were centrifuged in microcentrifuge tubes for 30 s at 10,000xg

4.14. Staining with Direct Yellow 96 (DY96)

To assay parasite burden and worm gravidity, microsporidia spores and embryos were visualized with

the chitin-binding dye DY96. Acetone-fixed animals were washed twice in 1 ml PBST, resuspended in

500 µl staining solution (PBST, 0.1% sodium dodecyl sulfate [SDS], 20 ug/ml DY96), and rotated at 21°C

for 30 min in the dark. Stained worms were resuspended in 20 µl EverBrite™ Mounting Medium (Biotium)

and mounted on slides for imaging. For pelleting of worms during staining protocols, animals were

centrifuged in microcentrifuge tubes for 30 s at 10,000xg.

4.15. Fluorescence in Situ Hybridization (FISH) assays

For FISH staining of *N. parisii* 18S rRNA or Orsay virus RNA, worms were fixed in PFA as above and

washed twice in 1 ml PBST. Worms were then washed once in 1 ml hybridization buffer (900 mM NaCl,

20 mM Tris pH 8.0, 0.01% SDS), and incubated overnight at 46°C in 100 µl hybridization buffer containing

5-10 ng/µl of FISH probe conjugated to a Cal Fluor 610 dye (LGC Biosearch Technologies). 5 ng/µl

MicroB (ctctcggcactccttcctg)[27] was used to detect *N. parisii* 18S RNA. 10 ng/µl of Orsay 1

(gacatatgtgatgccgagac) and Orsay 2 (gtagtgtcattgtaggcagc) mixed 50:50 was used to detect Orsay virus.

Stained animals were washed once in 1 ml wash buffer (hybridization buffer containing 5 mM EDTA) and

incubated in 500 µl fresh wash buffer for a further 30 min at 46°C. Worms were resuspended in 20 µl

EverBrite™ Mounting Medium (Biotium) and mounted on slides for imaging. To pellet worms during

staining protocols, animals were centrifuged in microcentrifuge tubes for 30 s at 10,000xg. For co-staining

of *N. parisii* RNA with DY96, wash buffer was supplemented with 20 ug/ml DY96 for the final 30 min

incubation.

4.16. Microscopy & image analysis

For measurement of worm body size, quantification of embryos and gravid or infected animals, as well

as *N. parisii* or *P. aeruginosa* burden, worms were imaged using an Axioimager 2 (Zeiss). Worms carrying

1 or more embryos were considered gravid, worms possessing any quantity of DY96 stained

microsporidia spores were considered infected. Bead ingestion and precise pathogen burdens were

determined using ImageJ/FIJI[66]; here, each worm was defined as an individual 'region of interest' and

fluorescence from GFP (DY96-stained microsporidia, or GFP beads) or dsRed (dsRed-expressing PA14)

subject to 'threshold' and 'measure area percentage' functions on ImageJ. For DY96-stained samples.

images were thresholded to capture the brighter signal from microsporidia spores, while eliminating the

dimmer GFP signal from worm embryos. Final values are given as % fluorescence for single animals.

4.17. Transcriptional analyses

WormExp[67] was used to search for published expression data sets that have a significant overlap with

the set of genes enriched in *N. parisii* infected animals.

FPKM values from RNA-Seq data were obtained for *pals-22*[32] mutants and five *N. parisii* infection time

points[36] and fold changes were calculated. Fold-change values were obtained for *lin-35* mutant

microarray data[68] and replicates averaged. Log$_2$ fold-changes greater than 2 or less than -2 were used

to determine differentially expressed genes. Genes that were either up or down regulated in *lin-35, pals-*

22 and at least one infection time point were plotted as a heatmap with dendrograms using heatmap.2

function from gplots package in R with all arguments[69] set to default except for trace which was set to

none.

4.18. Statistical analyses

594   Unless otherwise stated, p-values were determined by 2-tailed unpaired Student's t-test. All p-values not

595   meeting significance requirements are displayed on Figures for clarity. These p-values were calculated

596   using Prism software (GraphPad Software Inc.). Statistical significance is defined as p-value < 0.05,

597   unless otherwise stated (i.e. when using Bonferroni correction for multiple testing).

Acknowledgements

We thank Emily R. Troemel for her mentorship and support in allowing A.W.R to start this project as a

postdoctoral fellow in her lab. We thank Malina A. Bakowski and Lianne B. Cohen for sharing initial results

on the shared transcriptional similarity between *lin-35* mutants and microsporidia infection. We thank

Andy P. Ryan and Kirthi C. Reddy for sharing initial findings on the shared transcriptional similarity

between cadmium exposure and microsporidia infection. We thank Nicholas O. Burton for advice on the

osmotic stress assay. We are grateful to Calvin Mok and Nicholas O. Burton for providing helpful

comments on the manuscript. Additional *C. elegans* strains were provided by the *Caenorhabditis*

Genetics Center, which is funded by the National Institutes of Health (NIH) Office of Research

Infrastructure Programs Grant P40 OD010440. Schematics were created using BioRender.com. This

work was supported by the Natural Sciences and Engineering Research Council of Canada (Grant

#522691522691).

References

1.   Tetreau, G., Dhinaut, J., Gourbal, B. & Moret, Y. Trans-generational Immune Priming in

Invertebrates: Current Knowledge and Future Prospects. *Front Immunol* **10**, 1938 (2019).

2. Barribeau, S. M., Schmid-Hempel, P. & Sadd, B. M. Royal Decree: Gene Expression in Trans-

Generationally Immune Primed Bumblebee Workers Mimics a Primary Immune Response. *PLOS*

ONE **11**, e0159635 (2016).

3. Salmela, H., Amdam, G. V. & Freitak, D. Transfer of Immunity from Mother to Offspring Is

Mediated via Egg-Yolk Protein Vitellogenin. *PLOS Pathogens* **11**, e1005015 (2015).

4. Little, T. J., O'Connor, B., Colegrave, N., Watt, K. & Read, A. F. Maternal Transfer of Strain-

Specific Immunity in an Invertebrate. *Current Biology* **13**, 489–492 (2003).

5. Norouzitallab, P. *et al.* Environmental heat stress induces epigenetic transgenerational inheritance of

robustness in parthenogenetic Artemia model. *The FASEB Journal* **28**, 3552–3563 (2014).

6. Dubuffet, A. *et al.* Trans-generational Immune Priming Protects the Eggs Only against Gram-

Positive Bacteria in the Mealworm Beetle. *PLOS Pathogens* **11**, e1005178 (2015).

7. Kim, D. H. *et al.* A Conserved p38 MAP Kinase Pathway in Caenorhabditis elegans Innate

Immunity. *Science* **297**, 623–626 (2002).

8. Weinhouse, C., Truong, L., Meyer, J. N. & Allard, P. Caenorhabditis elegans as an emerging model

system in environmental epigenetics: C. elegans as an Environmental Epigenetics Model. *Environ.*

Mol. Mutagen. **59**, 560–575 (2018).

9. Fire, A. *et al.* Potent and specific genetic interference by double-stranded RNA in Caenorhabditis

elegans. *Nature* **391**, 806–811 (1998).

10. Willis, A. R., Sukhdeo, R. & Reinke, A. W. Remembering your enemies: mechanisms of within-

generation and multigenerational immune priming in Caenorhabditis elegans. *The FEBS Journal*

n/a,.

11. Rechavi, O., Minevich, G. & Hobert, O. Transgenerational inheritance of an acquired small RNA-

based antiviral response in C. elegans. *Cell* **147**, 1248–1256 (2011).

12. Ashe, A., Sarkies, P., Le Pen, J., Tanguy, M. & Miska, E. A. Antiviral RNA Interference against

Orsay Virus Is neither Systemic nor Transgenerational in Caenorhabditis elegans. *J Virol* **89**, 12035–

12046 (2015).

13. Gammon, D. B. *et al.* The Antiviral RNA Interference Response Provides Resistance to Lethal

Arbovirus Infection and Vertical Transmission in Caenorhabditis elegans. *Curr Biol* **27**, 795–806

(2017).

14. Sterken, M. G. *et al.* A heritable antiviral RNAi response limits Orsay virus infection in

Caenorhabditis elegans N2. *PLoS One* **9**, e89760 (2014).

15. Kaletsky, R. *et al.* C. elegans interprets bacterial non-coding RNAs to learn pathogenic avoidance.

Nature 1–7 (2020) doi:10.1038/s41586-020-2699-5.

16. Moore, R. S., Kaletsky, R. & Murphy, C. T. Piwi/PRG-1 Argonaute and TGF-beta Mediate

Transgenerational Learned Pathogenic Avoidance. *Cell* **177**, 1827-1841.e12 (2019).

17. Pereira, A. G., Gracida, X., Kagias, K. & Zhang, Y. C. elegans aversive olfactory learning generates

diverse intergenerational effects. *Journal of Neurogenetics* **0**, 1–11 (2020).

18. Palominos, M. F. *et al.* Transgenerational Diapause as an Avoidance Strategy against Bacterial

Pathogens in Caenorhabditis elegans. *mBio* **8**, mBio.01234-17, e01234-17 (2017).

19. Burton, N. O. *et al.* Cysteine synthases CYSL-1 and CYSL-2 mediate C. elegans heritable

adaptation to P. vranovensis infection. *Nat Commun* **11**, 1741 (2020).

20. Ma, C. *et al.* N6-methyldeoxyadenine is a transgenerational epigenetic signal for mitochondrial

stress adaptation. *Nat Cell Biol* **21**, 319–327 (2019).

21. Wadi, L. & Reinke, A. W. Evolution of microsporidia: An extremely successful group of eukaryotic

intracellular parasites. *PLoS Pathog* **16**, e1008276 (2020).

22. Stentiford, G. D. *et al.* Microsporidia – Emergent Pathogens in the Global Food Chain. *Trends*

Parasitol **32**, 336–348 (2016).

23. Luallen, R. J. *et al.* Discovery of a Natural Microsporidian Pathogen with a Broad Tissue Tropism in

Caenorhabditis elegans. *PLoS Pathog.* **12**, e1005724 (2016).

24. Balla, K. M., Andersen, E. C., Kruglyak, L. & Troemel, E. R. A wild C. elegans strain has enhanced

epithelial immunity to a natural microsporidian parasite. *PLoS Pathog* **11**, e1004583 (2015).

25. Balla, K. M., Luallen, R. J., Bakowski, M. A. & Troemel, E. R. Cell-to-cell spread of microsporidia

causes Caenorhabditis elegans organs to form syncytia. *Nat Microbiol* **1**, 16144 (2016).

26. Estes, K. A., Szumowski, S. C. & Troemel, E. R. Non-Lytic, Actin-Based Exit of Intracellular

Parasites from C. elegans Intestinal Cells. *PLoS Pathog* **7**, e1002227 (2011).

27. Troemel, E. R., Félix, M.-A., Whiteman, N. K., Barrière, A. & Ausubel, F. M. Microsporidia Are

Natural Intracellular Parasites of the Nematode Caenorhabditis elegans. *PLoS Biol* **6**, e309 (2008).

28. Jarkass, H. T. E. & Reinke, A. W. The ins and outs of host-microsporidia interactions during

invasion, proliferation and exit. *Cellular Microbiology* **22**, e13247 (2020).

29. Bakowski, M. A. *et al.* Ubiquitin-Mediated Response to Microsporidia and Virus Infection in C.

elegans. *PLOS Pathogens* **10**, e1004200 (2014).

30. Dierking, K., Yang, W. & Schulenburg, H. Antimicrobial effectors in the nematode Caenorhabditis

elegans: an outgroup to the Arthropoda. *Philosophical Transactions of the Royal Society B:*

Biological Sciences **371**, 20150299 (2016).

31. Reddy, K. C. *et al.* An Intracellular Pathogen Response Pathway Promotes Proteostasis in

C. elegans. *Curr. Biol.* **27**, 3544-3553.e5 (2017).

32. Reddy, K. C. *et al.* Antagonistic paralogs control a switch between growth and pathogen resistance

in C. elegans. *PLoS Pathog.* **15**, e1007528 (2019).

33. Balla, K. M., Lažetić, V. & Troemel, E. R. Natural variation in the roles of C. elegans autophagy

components during microsporidia infection. *PLOS ONE* **14**, e0216011 (2019).

34. Luallen, R. J., Bakowski, M. A. & Troemel, E. R. Characterization of Microsporidia-Induced

Developmental Arrest and a Transmembrane Leucine-Rich Repeat Protein in Caenorhabditis

elegans. *PLOS ONE* **10**, e0124065 (2015).

35. Tan, M.-W., Rahme, L. G., Sternberg, J. A., Tompkins, R. G. & Ausubel, F. M. Pseudomonas

aeruginosa killing of Caenorhabditis elegans used to identify P. aeruginosa virulence factors. *PNAS*

96, 2408–2413 (1999).

36. Bakowski, M. A. *et al.* Ubiquitin-mediated response to microsporidia and virus infection in C.

elegans. *PLoS Pathog* **10**, e1004200 (2014).

37. Lev, I., Bril, R., Liu, Y., Ceré, L. I. & Rechavi, O. Inter-generational consequences for growing

Caenorhabditis elegans in liquid. *Philosophical Transactions of the Royal Society B: Biological*

Sciences **374**, 20180125 (2019).

38. Félix, M.-A. *et al.* Natural and Experimental Infection of Caenorhabditis Nematodes by Novel

Viruses Related to Nodaviruses. *PLOS Biology* **9**, e1000586 (2011).

39. Lu, X. & Horvitz, H. R. lin-35 and lin-53, Two Genes that Antagonize a C. elegans Ras Pathway,

Encode Proteins Similar to Rb and Its Binding Protein RbAp48. *Cell* **95**, 981–991 (1998).

40. Yang, W., Dierking, K. & Schulenburg, H. WormExp: a web-based application for a Caenorhabditis

elegans-specific gene expression enrichment analysis. *Bioinformatics* **32**, 943–945 (2016).

41. Reddy, K. C. *et al.* An Intracellular Pathogen Response Pathway Promotes Proteostasis in

C. elegans. *Curr. Biol.* **27**, 3544-3553.e5 (2017).

42. Zhang, L., Ward, J. D., Cheng, Z. & Dernburg, A. F. The auxin-inducible degradation (AID) system

enables versatile conditional protein depletion in C. elegans. *Development* **142**, 4374–4384 (2015).

43. Kishimoto, S., Uno, M., Okabe, E., Nono, M. & Nishida, E. Environmental stresses induce

transgenerationally inheritable survival advantages via germline-to-soma communication in

Caenorhabditis elegans. *Nature Communications* **8**, 14031 (2017).

44. Eggert, H., Buhr, M. F. D. & Kurtz, J. A temperature shock can lead to trans-generational immune

priming in the Red Flour Beetle, Tribolium castaneum. *Ecology and Evolution* **5**, 1318–1326 (2015).

45. Sadd, B. M. & Schmid-Hempel, P. A distinct infection cost associated with trans-generational

priming of antibacterial immunity in bumble-bees. *Biology Letters* **5**, 798–801 (2009).

46. Félix, M.-A. & Duveau, F. Population dynamics and habitat sharing of natural populations of

Caenorhabditis elegans and C. briggsae. *BMC Biology* **10**, 59 (2012).

47. Kudron, M. *et al.* Tissue-specific direct targets of Caenorhabditis elegans Rb/E2F dictate distinct

somatic and germline programs. *Genome Biology* **14**, R5 (2013).

48. Cui, M., Cohen, M. L., Teng, C. & Han, M. The Tumor Suppressor Rb Critically Regulates

Starvation-Induced Stress Response in C. elegans. *Current Biology* **23**, 975–980 (2013).

49. González-Rangel, A. A. & Navarro, R. E. LIN-35 beyond its classical roles: its function in the stress

response. *Int. J. Dev. Biol.* (2020) doi:10.1387/ijdb.200194rn.

50. Hutcheson, J., Witkiewicz, A. K. & Knudsen, E. S. The RB tumor suppressor at the intersection of

proliferation and immunity: relevance to disease immune evasion and immunotherapy. *Cell Cycle*

14, 3812–3819 (2015).

51. Kirienko, N. V. & Fay, D. S. Transcriptome profiling of the C. elegans Rb ortholog reveals diverse

developmental roles. *Developmental Biology* **305**, 674–684 (2007).

52. Rechavi, O. *et al.* Starvation-induced transgenerational inheritance of small RNAs in C. elegans.

Cell **158**, 277–287 (2014).

53. Norouzitallab, P., Baruah, K., Vanrompay, D. & Bossier, P. Teaching Shrimps Self-Defense to Fight

Infections. *Trends in Biotechnology* **37**, 16–19 (2019).

54. Huang, C.-C. & Song, Y.-L. Maternal transmission of immunity to white spot syndrome associated

virus (WSSV) in shrimp (Penaeus monodon). *Developmental & Comparative Immunology* **23**, 545–

552 (1999).

55. Moret, Y. & Schmid-Hempel, P. Immune defence in bumble-bee offspring. *Nature* **414**, 506–506

(2001).

56. Roy, S., Kumar, V., Bossier, P., Norouzitallab, P. & Vanrompay, D. Phloroglucinol Treatment

Induces Transgenerational Epigenetic Inherited Resistance Against Vibrio Infections and Thermal

Stress in a Brine Shrimp (Artemia franciscana) Model. *Front. Immunol.* **10**, (2019).

57. Bjørnson, S. & Oi, D. Microsporidia Biological Control Agents and Pathogens of Beneficial Insects.

in *Microsporidia* 635–670 (John Wiley & Sons, Ltd, 2014). doi:10.1002/9781118395264.ch25.

58. Lorenz, L. M. & Koella, J. C. Maternal environment shapes the life history and susceptibility to

malaria of Anopheles gambiae mosquitoes. *Malaria Journal* **10**, 382 (2011).

59. Herren, J. K. *et al.* A microsporidian impairs Plasmodium falciparum transmission in Anopheles

arabiensis mosquitoes. *Nature Communications* **11**, 1–10 (2020).

60. Schwartz, M. L. & Jorgensen, E. M. SapTrap, a Toolkit for High-Throughput CRISPR/Cas9 Gene

Modification in *Caenorhabditis elegans*. *Genetics* **202**, 1277–1288 (2016).

61. Dickinson, D. J., Slabodnick, M. M., Chen, A. H. & Goldstein, B. SapTrap assembly of repair

templates for Cas9-triggered homologous recombination with a self-excising cassette.

microPublication Biology **2018**, (2018).

62. Dickinson, D. J., Pani, A. M., Heppert, J. K., Higgins, C. D. & Goldstein, B. Streamlined Genome

Engineering with a Self-Excising Drug Selection Cassette. *Genetics* **200**, 1035–1049 (2015).

63. Frokjaer-Jensen, C. *et al.* Single-copy insertion of transgenes in Caenorhabditis elegans. *Nat Genet*

40, 1375–1383 (2008).

64. Zhang, Y., Lu, H. & Bargmann, C. I. Pathogenic bacteria induce aversive olfactory learning in

Caenorhabditis elegans. *Nature* **438**, 179–184 (2005).

65. Burton, N. O. *et al.* Insulin-like signalling to the maternal germline controls progeny response to

osmotic stress. *Nat Cell Biol* **19**, 252–257 (2017).

66. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**,

676–682 (2012).

67. Yang, W., Dierking, K. & Schulenburg, H. WormExp: a web-based application for a Caenorhabditis

elegans-specific gene expression enrichment analysis. *Bioinformatics* **32**, 943–945 (2016).

68. Petrella, L. N. *et al.* synMuv B proteins antagonize germline fate in the intestine and ensure C.

elegans survival. *Development* **138**, 1069–1079 (2011).

69. R Development Core Team. *R: A Language and Environment for Statistical Computing*. (R

Foundation for Statistical Computing, 2019).

Figures

Figure 1. Parental infection by *N. parisii* confers immunity to the progeny of *C. elegans*.

P0 populations of N2 *C. elegans* were uninfected or exposed to varying concentrations of *N. parisii* spores

at the L1 stage (doses defined in methods). At 72 hpi, animals were treated with sodium hypochlorite

solution to release F1 embryos. F1 L1 larvae were then exposed to a high dose of *N. parisii*. At 72 hpi F1

animals were fixed and stained with DY96 to visualize both *N. parisii* spores and worm embryos. (A)

Representative images of F1 populations stained with DY96. Scale bars, 200 μm. (B) Images of DY96

stained F1 worms were analysed and fluorescence from *N. parisii* spores thresholded to determine

parasite burdens of individual worms (% of body filled with spores). Each circle represents a

measurement from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 3

independent experiments using n = 20 worms per condition per experiment. (C) Images of F1 worms

were analysed, and the area of individual worms calculated. Each circle represents a measurement from

a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments

using n = 20 worms per condition per experiment. (D) Images of DY96 stained F1 worms were analysed

and worms possessing 1 or more embryos were considered gravid. Mean ± SEM (horizontal bars) is

shown. Data pooled from 3 independent experiments using n = 98-351 worms per condition per

experiment. (E) Images of DY96 stained F1 worms were analysed and embryos per worm quantified.

Each circle represents a count from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled

from 3 independent experiments using n = 20 worms per condition per experiment. The p-values were

determined by unpaired two-tailed Student's t-test. (B-E) Significance with Bonferroni correction was

defined as $p < 0.0166$. **, $p < 0.0033$; ***, $p < 0.00033$.

Figure 2. Inherited immunity prevents microsporidia invasion and *P. aeruginosa* colonization, but not viral infection.

(A-D) P0 populations of N2 *C. elegans* were either not infected or infected with a low dose of *N. parisii* spores at the L1 stage (doses defined in methods). At 72 hpi, animals were treated with sodium hypochlorite solution to release F1 embryos. Naïve and primed F1 larvae were exposed to a maximal dose of *N. parisii* at the L1 stage. At 30 mpi, animals were fixed and stained with DY96 to detect *N. parisii* spores (green) as well as a FISH probe to detect *N. parisii* 18S RNA (red). (A) Representative images of worms stained with FISH probe to detect invaded sporoplasms, marked by asterisks. Scale bars, 25 μm. (B) The number of sporoplasms per animal was quantified by microscopy. Each circle represents a count from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n = 16-20 worms per condition per experiment. (C) Representative images of worms stained with DY96 to detect spores, marked by asterisks, in the intestinal lumen. Scale bars, 25 μm. (D) The number of spores per animal was quantified by microscopy. Each circle represents a count from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n = 20-21 worms per condition per experiment. (E) Naïve and primed F1 larvae were fed fluorescent beads at the L1 stage. After 30 min, animals were fixed and imaged. Fluorescence from beads was thresholded to determine the amount of beads eaten by each worm (% of body filled with beads). Each circle represents a measurement from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n = 24-30 worms per condition per experiment. (F) Naïve and *N. parisii*-primed F1 L1 larvae were maintained on slow-killing plates with wild-type *P. aeruginosa* and survival monitored. Animal survival at an 84 hpi end point is shown. Mean ± SEM (horizontal bars) is shown. Data pooled from 4 independent experiments each comprising 2-4 technical replicates, using n = 13-37 worms per condition per experiment. (G-H) Naïve and *N. parisii*-primed F1 L1 larvae were maintained on slow-killing plates with dsRed-*P. aeruginosa* and fixed at 48 hpi. (G) Representative images of worm populations grown on dsRed-*P. aeruginosa*. Scale bars, 200 μm. (H) Images of worms were analysed and fluorescence thresholded to determine bacterial burdens of individual worms (% of

body filled with *P. aeruginosa*). Each circle represents a measurement from a single worm. Mean ± SEM

(horizontal bars) is shown. Data pooled from 3 independent experiments using n = 21-47 worms per

condition per experiment. (I) Naïve and primed F1, and *pals-22* mutant L1 larvae were infected with Orsay

virus. At 16 hpi, worms were fixed and stained with FISH probe to detect Orsay virus RNA. Worms with

one or more cells stained by FISH were counted as infected. Mean ± SEM (horizontal bars) is shown.

Data pooled from 4 independent experiments with n > 52 worms per condition per experiment. The p-

values were determined by unpaired two-tailed Student's t-test. (B-F, H) Significance was defined as $p <$

0.05; **, $p < 0.01$; ***, $p < 0.001$. (I) Significance with Bonferroni correction was defined as *, $p < 0.025$;

****, $p < 0.0001$.

Figure 3. Inherited immunity in *N. parisii* primed *C. elegans* lasts a single generation and is strongest

in early larval stages.

(A-B) P0 populations of N2 *C. elegans* were either not infected or infected with a low dose of *N. parisii*

spores at the L1 stage (doses defined in methods). At 72 hpi, animals were treated with sodium

hypochlorite solution to release F1 embryos. F1 embryo populations were then split and either tested for

immunity, or maintained under non-infection conditions for the collection and subsequent testing of F2

embryos. Both naïve and primed F1 and F2 larvae were exposed to a high dose of *N. parisii* at the L1

stage. At 72 hpi, F1 and F2 animals were fixed and stained with DY96 to visualize *N. parisii* spores and

embryos. (A) Individual DY96 stained worms were imaged to determine infection status. Mean ± SEM

(horizontal bars) is shown. Data pooled from 3 independent experiments using n = 16-20 worms per

condition per experiment. (B) Images of DY96 stained worms were analysed and worms possessing 1 or

more embryos were considered gravid. Mean ± SEM (horizontal bars) is shown. Data pooled from 3

independent experiments using n = 16-20 worms per condition per experiment. (C) Naïve and primed F1

larvae were obtained as above and challenged with a maximal dose of *N. parisii* spores at either the L1,

L2/L3 or L4 stage. Animals were fixed at 30 mpi and stained with a FISH probe to detect *N. parisii* 18S

RNA. The number of sporoplasms per animal was quantified by microscopy. Shown is the average

number of sporoplasms in primed worms relative to the naïve control. Mean ± SEM (horizontal bars) is

shown. Data pooled from 3 independent experiments using n = 12-20 worms per condition per

experiment. (D-E) Briefly, N2 *C. elegans* (P0, F1, F2) were infected at the L1 stage with *N. parisii* for one,

two or three successive generations. Each infection period lasted 72 h before treating with sodium

hypochlorite solution to obtain the next generation of embryos (see schematic Figure S5C). F3 larvae

were exposed to a high dose of *N. parisii* at the L1 stage. At 72 hpi, F3 animals were fixed and stained

with DY96 to visualize *N. parisii* spores and embryos. (D) Images of DY96 stained F3 worms were

analysed and fluorescence from *N. parisii* spores thresholded to determine parasite burdens of individual

worms (% of body filled with spores). Each circle represents a measurement from a single worm. Mean

± SEM (horizontal bars) is shown. Data pooled from 2 independent experiments using n = 25 worms per

condition per experiment. (E) Images of DY96 stained F3 worms were analysed and embryos per worm

quantified. Each circle represents a count from a single worm. Mean ± SEM (horizontal bars) is shown.

Data pooled from 2 independent experiments using n = 30 worms per condition per experiment. (F-G)

P0 populations of N2 *C. elegans* were either not infected or infected with a low dose of *N. parisii* spores

at the L1 stage (for 72 h), L2/L3 stage (for 48 h) or L4 stage (for 24 h). At 72 h post L1, animals were

treated with sodium hypochlorite solution to release F1 embryos. Both naïve and primed F1 larvae were

exposed to a high dose of *N. parisii* at the L1 stage. At 72 hpi, animals were fixed and stained with DY96

to visualize *N. parisii* spores and embryos. (F) Individual DY96 stained worms were imaged to determine

infection status. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments

using n = 100-197 worms per condition per experiment. (G) Images of DY96 stained worms were

analysed and worms possessing 1 or more embryos were considered gravid. Mean ± SEM (horizontal

bars) is shown. Data pooled from 4 independent experiments using n = 100-197 worms per condition per

experiment. The p-values were determined by unpaired two-tailed Student's t-test. (A-B, G) Significance

was defined as: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. (C, F) Significance with Bonferroni correction was

defined as: *, $p < 0.025$; **, $p < 0.005$; ***, $p < 0.0005$. (D-E). Significance with Bonferroni correction was

defined as: *, $p < 0.016$.

Figure 4. The parental transcriptional response to *N. parisii* triggers inherited immunity.

(A-B) P0 populations of N2 *C. elegans* were either not infected or exposed to a low dose of heat-killed or live *N. parisii* spores at the L1 stage (doses defined in methods). At 72 hpi, animals were treated with sodium hypochlorite solution to release F1 embryos. F1 larvae were exposed to a high dose of *N. parisii* at the L1 stage. At 72 hpi, animals were fixed and stained with DY96 to visualize both *N. parisii* spores and worm embryos. (A) Individual DY96 stained worms were imaged to determine infection status. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n =100 worms per condition per experiment. (B) Images of DY96 stained F1 worms were analysed and worms possessing 1 or more embryos were considered gravid. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n = 100 worms per condition per experiment. (C-D) P0 populations of N2 and *rde-1* mutants were not infected, infected with *N. parisii* (low dose), or infected with Orsay virus at the L1 stage. At 72 hpi, animals were treated with sodium hypochlorite solution to release F1 embryos. F1 larvae were exposed to a high dose of *N. parisii* at the L1 stage, then fixed and stained at 72 hpi as above. (C) DY96 stained worms were analyzed to determine infection status. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n = 100 worms per condition per experiment. (D) Images of DY96 stained F1 worms were analysed and worms possessing 1 or more embryos were considered gravid. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n = 100 worms per condition per experiment. (E-G) P0 populations of N2 *C. elegans* were either untreated, exposed to 50 mM cadmium from the L4 stage, or infected with a low dose of *N. parisii* spores at the L4 stage. After 24 h, animals were treated with sodium hypochlorite solution to release F1 embryos. (E-F) F1 larvae were exposed to a high dose of *N. parisii* at the L1 stage. At 72 hpi, animals were fixed and stained with DY96 to visualize both *N. parisii* spores and worm embryos. (E) Individual DY96 stained worms were imaged to determine infection status. Mean ± SEM (horizontal bars) is shown. Data pooled from 4 independent experiments using n = 100 worms per condition per experiment. (F) Images of DY96 stained worms were analysed and embryos per worm quantified. Each circle represents a count from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from

4 independent experiments using n = 25-30 worms per condition per experiment. (G) Naïve and

cadmium-primed F1 L1 larvae were maintained on slow-killing plates with dsRed-*P. aeruginosa* and fixed

at 48 hpi. Images of worms were analysed and fluorescence thresholded to determine bacterial burdens

of individual worms (% of body filled with bacteria). Each circle represents a measurement from a single

worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n =

25 worms per condition per experiment. (H) P0 populations of N2 *C. elegans* were either untreated or

infected with a low dose of *N. parisii* spores at the L1 stage on regular NGM (containing 50 mM salt), or

maintained on 250 mM salt. After 72 h, animals were treated with sodium hypochlorite solution to release

F1 embryos. F1 embryos were maintained on 420 mM salt. At 48 hpi, embryo hatching was assessed by

light microscopy. Mean ± SEM (horizontal bars) is shown. Data pooled from 4 independent experiments

using n = 100 worms per condition per experiment. The p-values were determined by unpaired two-tailed

Student's t-test. (A-F, H) Significance with Bonferroni correction was defined as: *, $p < 0.025$; **, $p <$

0.005; ***, $p < 0.0005$. (G) Significance was defined as $p < 0.05$. ***, $p < 0.001$.

Figure 5. Mutants that phenocopy the transcriptional response to infection transfer immunity to

offspring through the maternal germline.

(A-B) Wild-type, *pals-22(jy1)*, and *lin-35(n745)* animals were not infected or infected with a high dose of

N. parisii (infected) from the L1 stage. After 72 h, animals were fixed and stained with DY96 to visualize

N. parisii spores and embryos. (A) Images of worms were analysed and embryos per worm quantified.

Each circle represents a count from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled

from 3 independent experiments using n = 20 worms per condition per experiment. (B) Images of worms

were analysed and fluorescence from *N. parisii* spores thresholded to determine parasite burdens of

individual worms (% of body filled with spores). Each circle represents a measurement from a single

worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n =

20 worms per condition per experiment. (C) Schematic of mating to obtain maternal and paternal cross-

progeny. *myo*-2p::mCherry used as a marker to distinguish cross-progeny from self-progeny. (D-E) P0

animals were allowed to mate for 24 h and then treated with sodium hypochlorite solution to release F1

embryos. F1 larvae were not infected or infected with a high dose of *N. parisii* at the L1 stage. At 72 hpi,

animals were fixed and stained with DY96 to visualize *N. parisii* spores and embryos. (D) Images of

worms were analysed and embryos per worm quantified. Each circle represents a count from a single

worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n =

20 worms per condition per experiment. (E) Images of worms were analysed and fluorescence from *N.*

parisii spores thresholded to determine parasite burdens of individual worms (% of body filled with

spores). Each circle represents a measurement from a single worm. Mean ± SEM (horizontal bars) is

shown. Data pooled from 3 independent experiments using n = 20 worms per condition per experiment.

Only hermaphrodite maternal cross progeny are included in the quantifications. (F-G) P0 animals were

allowed to mate for 24 h and then treated with sodium hypochlorite solution to release F1 embryos. F1

larvae were exposed to a high dose of *N. parisii* at the L1 stage. At 72 hpi, animals were fixed and stained

with DY96 to visualize *N. parisii* embryos. Images of worms were analysed and worms possessing 1 or

more embryos were considered gravid. Mean ± SEM (horizontal bars) is shown. Data pooled from 3

independent experiments using n = 13-67 worms per condition per experiment. The p-values were

determined by Kruskal-Wallis test (non-parametric one-way ANOVA) (A-B, D-E) and by ordinary one-

way ANOVA with post hoc (F-G). Significance was defined as: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$, ****,

$p < 0.0001$.

944
945

946
947
948

Figure 6. Induction of the IPR in somatic tissues induces inherited immunity.

(A-B) P0 animals were allowed to mate for 24 h and then treated with sodium hypochlorite solution to release F1 embryos. F1 larvae were exposed to a high dose of *N. parisii* at the L1 stage. At 72 hpi, animals were fixed and stained with DY96 to visualize embryos. The percentage of gravid F1 was quantified for the maternal cross progeny of (A) pals-22 mutants carrying a wild-type pals-22 transgene and (B) lin-35 mutants carrying a wild-type lin-35 transgene. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n = 17-78 worms per condition per experiment. Tissues expressing the rescue transgenes are indicated on the graph. (C-D) P0 animals were grown on either control plates or plates containing 200uM auxin from embryos to mediate degradation of PALS-22 or LIN-35. At 72 h animals were treated with sodium hypochlorite solution to release F1 embryos. F1 larvae were exposed to a high dose of *N. parisii* at the L1 stage. At 72 hpi, animals were fixed and stained with DY96 to visualize embryos. The percentage of gravid F1 was quantified for (C) degron-tagged pals-

22 strains and (D) degron-tagged lin-35 strains. Mean ± SEM (horizontal bars) is shown. Data pooled

from 3 independent experiments using n > 100 worms per condition per experiment. Tissues expressing

the rescue transgenes are indicated on the graph. The p-values were determined by ordinary one-way

ANOVA with post hoc (A-D). Significance was defined as: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****,

$p < 0.0001$.

Supplementary figures

Figure S1. Infection of *C. elegans* by *N. parisii* severely impacts worm fitness.

N2 *C. elegans* were uninfected or exposed to varying concentrations of *N. parisii* spores at the L1 stage

(doses defined in methods). (A-F) At 72 hpi, animals were fixed and stained with DY96 to visualize both

N. parisii spores and worm embryos. (A) Representative images of worm populations stained with DY96.

Scale bars, 200 μm. (B) Representative images of individual worms stained with DY96 (green) and DAPI

(blue). Left inset image shows DY96-stained embryos. Right inset image shows DY96 stained spores.

Scale bars, 100 μm. (C) Images of DY96 stained worms were analysed and fluorescence from *N. parisii*

spores thresholded to determine parasite burdens of individual worms (% of body filled with spores).

Mean ± SEM (horizontal bars) is shown. Curve generated using the hyperbola nonlinear fit line function

on GraphPad Prism software. Data pooled from 3 independent experiments using n = 20 worms per

condition per experiment. (D) Images of worms were analysed, and the area of individual worms

calculated. Each circle represents a measurement from a single worm. Mean ± SEM (horizontal bars) is

shown. Data pooled from 3 independent experiments using n = 20 worms per condition per experiment.

(E) Images of DY96 stained worms were analysed and worms possessing 1 or more embryos were

considered gravid. Mean ± SEM (horizontal bars) is shown. Data pooled from 4 independent experiments

using n = 35-195 worms per condition per experiment. (F) Images of DY96 stained worms were analysed

and embryos per worm quantified. Each circle represents a count from a single worm. Mean ± SEM

(horizontal bars) is shown. Data pooled from 3 independent experiments using n = 20 worms per condition

per experiment. The p-values were determined by unpaired two-tailed Student's t-test. (D-F) Significance

with Bonferroni correction was defined as $p < 0.01$. **, $p < 0.002$; ***, $p < 0.0002$. (G) Transgenic *C.*

elegans expressing a fluorescent germline protein (GFP::3xFLAG::CSR-1) were infected with *N. parisii*

as above. Representative images of the germlines of individual worms. Scale bars, 50 μm.

Figure S2. *N. parisii* infection of *C. elegans* is not vertically transmitted.

P0 populations of N2 *C. elegans* were either not infected or infected with a moderate dose of *N. parisii* spores at the L1 stage (doses defined in methods). At 72 hpi, animals were treated with sodium hypochlorite solution to release F1 embryos. Naïve uninfected larvae served as a negative control for *N. parisii* detection. Naïve infected larvae exposed to a maximal dose of *N. parisii* served as a positive control for *N. parisii* detection. At 2 hpi F1 animals were fixed and stained with DY96 to detect *N. parisii* spores (green) as well as a FISH probe to detect *N. parisii* 18S RNA (red). *N. parisii* was never observed in primed, uninfected animals. Representative images of F1 populations are shown. Scale bars, 50 μm.

1005

Figure S3. Progeny of *N. parisii* infected *C. elegans* are less fit when uninfected.

1006

P0 populations of N2 *C. elegans* were uninfected or exposed to varying concentrations of *N. parisii* spores

at the L1 stage (doses defined in methods). At 72 hpi, animals were treated with sodium hypochlorite

solution to release F1 embryos. Naïve or primed F1 L1 larvae were grown under non-infection conditions

for 72 h then fixed and stained with DY96 to visualize worm embryos. (A) Images of F1 worms were

analysed, and the area of individual worms calculated. Each circle represents a measurement from a

single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 2 independent experiments using

n = 17-20 worms per condition per experiment. (B) Images of DY96 stained F1 worms were analysed

and embryos per worm quantified. Each circle represents a count from a single worm. Mean ± SEM

(horizontal bars) is shown. Data pooled from 2 independent experiments using n = 20 worms per condition

per experiment. The p-values were determined by unpaired two-tailed Student's t-test. Significance with

Bonferroni correction was defined as: *, p < 0.0166; **, p < 0.0033; ***, p < 0.00033.

Figure S4. Inherited immunity protects primed larvae from infection

Figure S4. Inherited immunity reduces microsporidia and *Pseudomonas* in the intestine, whereas

***pals-22 mutants* both reduce microsporidia invasion and cause microsporidia clearance.**

P0 populations of N2 *C. elegans* were either not infected or infected with a low dose of *N. parisii* spores

at the L1 stage (doses defined in methods). At 72 hpi, animals were treated with sodium hypochlorite

solution to release F1 embryos. (A-B) Naïve and primed F1 larvae were exposed to a very high dose of

N. parisii at the L1 stage. At 3 hpi, animals were fixed and stained with DY96 to detect *N. parisii* spores

(green) as well as a FISH probe to detect *N. parisii* 18S RNA and reveal intracellular sporoplasms (red).

(A) The number of sporoplasms per animal was quantified by microscopy. Each circle represents a count

from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 2 independent

experiments using n = 20-26 worms per condition per experiment. (B) The number of spores per animal

was quantified by microscopy. Each circle represents a count from a single worm. Mean ± SEM

(horizontal bars) is shown. Data pooled from 2 independent experiments using n = 20 worms per condition

per experiment. (C) Naïve and primed F1 larvae were fed on fluorescent beads at the L1 stage. After 3

hours, animals were fixed and imaged. Fluorescence from beads was thresholded to determine the

amount of beads eaten by each worm (% of body filled with beads). Each circle represents a

measurement from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 2

independent experiments using n = 20-30 worms per condition per experiment. (D) Naïve and primed F1

larvae were exposed to a very high dose of *N. parisii* at the L1 stage. At 3 hpi, populations were split and

half of the animals fixed. The remaining animals were maintained in the absence of spores until a 24 h

end point before fixing. Animals were stained with a FISH probe to detect *N. parisii* 18S RNA (red) and

the number of sporoplasms per animal was quantified by microscopy. Each circle represents a count

from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 2 independent

experiments using n = 20 worms per condition per experiment. (E) Naïve and *N. parisii*-primed F1 L1

larvae were maintained on slow-killing plates with dsRed-*P. aeruginosa* and fixed at 48 hpi.

Representative images of worms grown on dsRed-*P. aeruginosa* are shown. Scale bars, 100 μm. (F) N2

and *pals-22* F1 larvae were exposed to a very high dose of *N. parisii* at the L1 stage. At 3 hpi, populations

were split and half of the animals fixed. The remaining animals were maintained in the absence of spores

until a 24 h end point before fixing. Animals were stained with a FISH probe to detect *N. parisii* 18S RNA

(red) and the number of sporoplasms per animal was quantified by microscopy. Three independent

experiments with n = 100 worms per experiment. The p-values were determined by unpaired two-tailed

Student's t-test. Significance with Bonferroni correction was defined as $p < 0.05$. ***, $p < 0.001$.

1050

1051

1052

Figure S5. Inherited immunity in *N. parisii* primed *C. elegans* lasts a single generation

(A-B) P0 populations of N2 *C. elegans* were either not infected or infected with a low dose of *N. parisii* spores at the L1 stage (doses defined in methods). At 72 hpi, animals were treated with sodium hypochlorite solution to release F1 embryos. F1 L1 populations were then split and either subject to infection testing, or maintained under non-infection conditions for the collection and subsequent testing of F2 embryos. Both naïve and primed F1 and F2 larvae were exposed to a high dose of *N. parisii* at the L1 stage. At 72 hpi, F1 and F2 animals were fixed and stained with DY96 to visualize *N. parisii* spores. (A) Images of DY96 stained worms were analysed and fluorescence from *N. parisii* spores thresholded to determine parasite burdens of individual worms (% of body filled with spores). Each circle represents a measurement from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n = 15-20 worms per condition per experiment. (B) Images of worms were analysed, and the area of individual worms calculated. Each circle represents a measurement from a single worm. Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n = 15-20 worms per condition per experiment. (C) Schematic of ancestral infection history assay. N2 *C. elegans* (P0, F1, F2) were infected at the L1 stage with *N. parisii* for one, two or three successive generations. Each infection period lasted 72 h before treating with sodium hypochlorite solution to obtain the next generation of embryos. F3 L1 populations were split and either tested for immunity or maintained under non-infection conditions for the collection and subsequent testing of F4 offspring. For infection testing, F3 and F4 larvae were exposed to a high dose of *N. parisii* at the L1 stage. At 72 hpi, F3 and F4 animals were fixed and stained with DY96 to visualize *N. parisii* spores. (D) As the offspring of infected parents are resistant to infection, second (F1) or third (F2) generation doses were increased to ensure these primed animals still became infected. To determine infection status of F2 populations prior to testing of next generations, individual DY96 stained worms were imaged to determine infection status. Mean ± SEM (horizontal bars) is shown. Data pooled from 2 independent experiments using n = 35-133 worms per condition per experiment. (E) Images of DY96 stained F2 worms were analysed and embryos per worm quantified. Each circle represents a count from a single worm. Mean ± SEM (horizontal bars) is

shown. Data pooled from 2 independent experiments using n = 25-30 worms per condition per

experiment. (F) Individual DY96 stained F4 worms were imaged to determine infection status. Mean ±

SEM (horizontal bars) is shown. Data pooled from 2 independent experiments using n = 37-200 worms

per condition per experiment. (G) Images of DY96 stained F4 worms were analysed and embryos per

worm quantified. Each circle represents a count from a single worm. Mean ± SEM (horizontal bars) is

shown. Data pooled from 2 independent experiments using n = 30 worms per condition per experiment.

The p-values were determined by unpaired two-tailed Student's t-test. (A-B) Significance was defined as:

*, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. (F-G) Significance with Bonferroni correction was defined as $p <$

0.016.

1089

D

Category	Dataset	Source	# of genes overlapping with UP by *N. parisii* datasets (Bakowski et al. 2014)				
			8 hpi, 123 genes	16 hpi, 68 genes	30 hpi, 103 genes	40 hpi, 64 genes	64 hpi, 79 genes
Microbes	UP by virus Orsay, 53 genes	(Sarkies et al. 2013)	35 (p = 9.3e-70)	31 (p = 2.4e-68)	34 (p = 5.3e-70)	18 (p = 1.6e-33)	8 (p = 1.1e-10)
Chemicals/ Stress	UP by cadmium, 232 genes	(Cui et al. 2007)	36 (p = 5.6e-45)	25 (p = 1.1e-33)	35 (p = 2.9e-46)	24 (p = 1.5e-32)	14 (p = 5.3e-14)
Mutants	Up by pals-22, 2749 genes	(Reddy et al. 2019)	105 (p = 2.0e-87)	62 (p = 3.9e-55)	78 (p = 2.6e-57)	42 (p = 7.4e-27)	38 (p = 5.5e-18)
	UP by lin-35 mutant (Petrella), 657 genes	(Petrella et al. 2011)	73 (p = 1.0e-86)	47 (p = 5.5e-60)	55 (p = 1.8e-61)	28 (p = 5.8e-28)	14 (p = 2.0e-08)

1090

1091

Figure S6. Transmission of inherited immunity requires the transcriptional response to

microsporidia, which is mimicked by other environmental conditions and mutants.

(A) Transgenic *C. elegans* expressing a fluorescent germline protein (GFP::3xFLAG::CSR-1) were

infected with a low dose of *N. parisii* spores at the L1 stage (doses defined in methods). At 72 hpi, animals

were fixed and stained with a FISH probe to detect *N. parisii* 18S RNA (red). Representative images of

the germline of an infected worm are shown. Inset images: *Np*, *N. parisii*; (1) DG, distal gonad; (2) PG,

posterior gonad; (3) E, embryo. Scale bars, 20 μm. (B-C) Transgenic *C. elegans* expressing GFP under

IPR gene promoters were either uninfected or exposed to a low dose of heat-killed or live *N. parisii* spores

at the L1 stage. At 5 hpi, animals were imaged. Scale bars, 100 μm. (B) Representative images of

transgenic *pals-5p::gfp* worms. (C) Representative images of transgenic *f26f2.1p::gfp* worms. (D) Table

comparing genes previously reported to be upregulated in *N. parisii* infected animals with gene

expression changes in other published data sets. p-values calculated using Fisher's Exact test. (E) N2

and *rde-1* mutant L1 larvae were infected with Orsay virus and fixed at 72 hpi. Representative images of

worms stained with FISH probe to detect Orsay virus RNA. Scale bars, 400 μm. (F) P0 populations of N2

C. elegans were either untreated, exposed to 50 mM cadmium from the L4 stage, or infected with a low

dose of *N. parisii* spores at the L4 stage. After 24 h, animals were treated with sodium hypochlorite

solution to release F1 embryos. F1 larvae were exposed to 50 mM cadmium at the L4 stage. After 24 h,

animals were fixed and stained with DY96 to visualize worm embryos. Images of DY96 stained worms

were analysed and embryos per worm quantified. Each circle represents a count from a single worm.

Mean ± SEM (horizontal bars) is shown. Data pooled from 3 independent experiments using n = 25-26

worms per condition per experiment. The p-values were determined by unpaired two-tailed Student's t-

test. Significance with Bonferroni correction was defined as $p < 0.025$. ***, $p < 0.0005$.

1114

1115

Figure S7. Intergenerational transmission of immunity does not depend on small-RNA inheritance

factors, a histone methyltransferase, or the P38 MAP kinase pathway.

(A-D) N2 or mutant P0 animals were either not infected or infected with a moderate dose of *N. parisii*

spores at the L1 stage (doses defined in methods). At 72 hpi, animals were treated with sodium

hypochlorite solution to release F1 embryos. Naïve or primed F1 L1 larvae were then infected with a high

dose of *N. parisii*. Percentage of infected animals (A, C) and gravid animals (B, D) were quantified in the

infected F1s at 72hpi. Mean ± SEM (horizontal bars) is shown. 2-5 independent experiments with n > 100

worms each. The p-values were determined by unpaired two-tailed Student's t-test. Significance was

defined as: *, p < 0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001

1125

1126

Figure S8. *N. parisii* infection induces many genes that are also upregulated in both *lin-35* and *pals-22* mutants. A shared transcriptional response was identified by determining genes that were either up or down regulated in both *lin-35* and *pals-22* mutants and at least one *N. parisii* infection time point. (A) Heat map showing cluster analysis of the shared transcriptional response with the fold change of each cell corresponding to scale at the top. White cells in heatmap represent gene not determined to be differentially expressed. (B) Fraction of shared genes that are either up or down regulated.

1133

1134

1135

1136

1137 **Figure S9. Degradation of *lin-35* in somatic tissues.**

1138 Degron::GFP::*lin-35* worms with somatic TIR1 expression were grown on control plates, containing no

1139 auxin, or 200 uM auxin plates for degradation of LIN-35. Images of representative animals show GFP

1140 expression 48 h post-hatch on their respective plates. Asterisks mark intestinal nuclei GFP. Scale bars,

1141 100 μm.

1142

1143