Running Head: MEDIAL PFC AND EMOTIONAL MEMORY CONSOLIDATION

Medial prefrontal cortex has a causal role in selectively enhanced consolidation of emotional memories after a 24-hour delay: An iTBS study

Nicholas Yeh[1]*, Jessica D. Payne[1], Sara Y. Kim[1], Elizabeth A. Kensinger[2], Joshua D. Koen[1], Nathan S. Rose[1]

[1] Department of Psychology, University of Notre Dame, Notre Dame, IN, USA
[2] Department of Psychology, Boston College, Chestnut Hill, MA, USA

*Corresponding Author:
Nicholas Yeh
Department of Psychology
University of Notre Dame
Notre Dame, IN 46556
Email: nyeh@nd.edu

Acknowledgments:
Preparation of this manuscript was supported by the National Science Foundation (Grant BCS-1539361 awarded to J.D.P and E.A.K., and CAREER Grant 1848440 awarded to N.S.R) and from the University of Notre Dame William P. and Hazel B. Collegiate Chair funds (awarded to N.S.R). The authors would like to thank Katie McGuckin, Willian De Faria, Clare Hannon, Shannon O'Hara, and Frank Calabrese for contributing to data collection and analysis.

Conflict of interest:
The authors declare no conflicts of interest.

34   **Abstract**

35   Previous research points to an association between retrieval-related activity in the medial

prefrontal cortex (mPFC) and preservation of emotional information compared to co-occurring

neutral information following sleep. Although the role of the mPFC in emotional memory likely

begins at encoding, little research has examined how mPFC activity during encoding interacts

with consolidation processes to enhance emotional memory. This issue was addressed in the

present study using transcranial magnetic stimulation in conjunction with an emotional memory

paradigm. Healthy young adults encoded negative and neutral scenes while undergoing

concurrent TMS with a short modified intermittent theta burst stimulation (iTBS) protocol.

Participants received stimulation to either the mPFC or an active control site (motor cortex)

during the encoding phase. Recognition memory for scene components (objects and

backgrounds) was assessed after a short (30-minute) and a long delay (24-hour, including a night

of sleep) to obtain measures of specific and gist-based memory processes. The results

demonstrated that, relative to control stimulation, iTBS to the mPFC enhanced memory for

negative objects on the long delay test (collapsed across specific and gist-based memory

measures). mPFC stimulation had no discernable effect on memory for objects on the short delay

test nor on the background images at either test. These results suggest that mPFC activity

occurring during encoding interacts with consolidation processes to preferentially preserve

negatively salient information.

Significance Statement

Understanding how emotional information is remembered over time is critical to understanding memory in the real world. The present study uses transcranial magnetic stimulation (TMS) to investigate the interplay between mPFC activity that occurs during memory encoding and its subsequent interactions with post-encoding consolidation processes. Excitatory TMS delivered to the mPFC during encoding enhanced memory for negatively valenced pictures on a test following a 24-hr delay, with no such effect on a test occurring shortly after the encoding phase. These results are consistent with the hypothesis that emotional aspects of memories are differentially subjected to consolidation processes, and that the mPFC might contribute to this "tag-and-capture" mechanism during the initial formation of such memories.

Introduction

Emotional memory is often characterized by "trade-off effects", where superior

recollection of emotional experiences, or emotional parts of an experience, occur at the expense

of memory for neutral aspects (Kensinger et al., 2007a; Payne et al., 2008). One account of

emotional memory trade-offs proposes that synaptic processes operating near the initial learning

event can "tag" emotionally salient aspects of an experience and set the stage for downstream

preferential consolidation of these emotional aspects (Kim & Payne, 2020; Payne & Kensinger,

2018; also see, Richter-Levin & Akirav, 2003). This "tag and capture" mechanism has been

proposed to explain how encoding and subsequent consolidation processes interact to promote

later stabilization of memories. While the synaptic tag has yet to be directly examined in

humans, recent studies have observed network-level neural processes (via functional magnetic

resonance imaging; Tambini, Rimmele, Phelps, & Davachi, 2017) and behavioral outcomes

(Ballarini, Martinez, Diaz Perez, Moncada, & Viola, 2013; Dunsmoor, Murty, Davachi, &

Phelps, 2015; Patil, Murty, Dunsmoor, Phelps, & Davach, 2016) that are consistent with the

theorized tagging mechanism.

An open question in memory research is whether these network-level neural processes

can be utilized to causally manipulate the preferential encoding and consolidation of emotional

information. Neuroimaging studies have established that increased levels of activation of the

medial temporal lobe (MTL; including the amygdala and hippocampus) and the medial

prefrontal cortex (mPFC) are associated with successful emotional encoding of negative

information (cf. Murty et al., 2010; Payne & Kensinger, 2010, 2011, 2018; see also Goto &

Grace, 2008). Communication between the MTL and mPFC (i.e., functional connectivity)

appears to be critical for successful emotional memory encoding as well (Berkers et al., 2016;

Kensinger & Corkin, 2004).

There is also evidence that longer consolidation delays (e.g., 24-hour delays) are crucial

for emotion-related memory enhancements (Dunsmoor et al., 2015; see also Patil et al., 2016 for

reward-related memory). One probable reason for this delay-dependency is the presence of sleep

during the consolidation interval. Sleep-based consolidation processes play a pivotal role in

emotional memory trade-off effects by selectively preserving negative information (Payne et al.,

2008, 2012; Payne & Kensinger, 2011), especially gist-based aspects of negative information

(Payne et al., 2008; Hu et al., 2006). Evidence from fMRI studies suggests that sleep leads to a

refinement in the neural networks engaged in emotional memory retrieval. For example, studies

have revealed that initial broad network activation observed during encoding is refined into a

smaller network centered on the amygdala, hippocampus, and ventral mPFC following a sleep-

filled consolidation delay, with greater connectivity among these regions correlating with

enhanced memory for negative information compared to wakefulness (Bennion et al., 2015;

Payne & Kensinger, 2011; Sterpenich et al., 2009). This refinement in emotional memory

retrieval networks presumably occurs as memories are consolidated.

Importantly, the processes (e.g., tag and capture mechanisms) that lead to refinements in

emotional memory retrieval networks are established near the time of encoding. Payne and

Kensinger (2018) propose that successful emotional memory will be optimal with increased

MTL and PFC activity near encoding, with sleep occurring shortly after. Critically, emotional

tags can be set via stress- and arousal-related neuromodulators and reflect initial encoding

activity between the MTL and PFC but require a consolidation delay containing sleep to ensure

that these tags persist (Payne & Kensinger, 2018), suggesting that post-learning sleep is essential

for transforming temporary synaptic changes into long-lasting systems-level ones, and for

linking these distributed tags into an integrated memory trace.

Thus, by modulating neural activity at encoding, we may be able to affect how emotional

information is subsequently consolidated. The present work therefore sought to understand how

the mPFC might influence interactions between encoding and consolidation processes that are

thought to lead to selective preservation of negative memories (Payne & Kensinger, 2018).

During the encoding phase of an emotional memory trade-off task (Payne et al., 2009), we

applied intermittent theta burst stimulation (iTBS), which induces excitatory neural effects in the

targeted and functionally connected areas (Hermiller, Chen, Parrish, & Voss, 2020; Huang et al.,

2005; Tang et al., 2019), to the mPFC or an active control site (motor cortex; MC). A recognition

test assessed memory following a short (30-minute) delay and a long (24-hour) delay that

included sleep.

Under the assumption that mPFC activity at encoding interacts with subsequent

consolidation processes to refine the MTL-mPFC network associated with memory for negative

information, we expected that iTBS-induced excitation of the mPFC during an emotional

memory encoding task would, in turn, modulate activity within an emotion–cognition network,

including changes in MTL-mPFC connectivity. Altered behavioral outcomes (i.e., emotional

memory performance) at the short delay would be suggestive of differences in encoding and

early consolidation processes due to mPFC stimulation. On the other hand, altered behavioral

outcomes at the long delay, especially in the absence of differences at the short delay, would

indicate that TMS-related modulation of this network prioritized certain items during encoding

for *later* consolidation (e.g., during sleep). Thus, specifically, our *a priori* hypothesis was that,

relative to control stimulation, mPFC stimulation would preferentially enhance memory for

negative, but not neutral, information following a long delay. Furthermore, in light of research

showing dissociable relationships between sleep-based consolidation and specific vs. gist-like

emotional memories (Bovy et al., 2020; Hu et al., 2006; Payne et al., 2008), this study also

examined the effects of mPFC stimulation during encoding on specificity for these emotional

memory trade-offs.

139   **Materials and Methods**

140   *Participants*

141   Forty-five participants (aged 18-24; $M = 19.71$) were recruited from the University of

Notre Dame. Participants identified as 71% Caucasian, 11% Asian, 9% African American, 7%

Hispanic, and 2% who declined to state. All participants were fluent English speakers, had no

history of sleep disturbances, and did not take medication known to impact sleep or present

contraindications for TMS. The current sample size was selected to be in line with prior TMS

work (Sack et al., 2009) that has detected significant behavioral effects when targeting cortical

regions based on the 10-20 system (see TMS section below). Participants were randomly

assigned to receive iTBS either to the mPFC ($N = 23$) or to motor cortex (MC; $N = 22$), which

served as an active control site.

150   Participants were screened for contraindications for TMS following published criteria

(Rossi et al., 2009), and were excluded if they self-reported any history of seizures, brain injuries

(e.g., stroke, aneurysms, concussions, traumatic brain injury), severe headaches/migraines,

fainting, metal implants, neurological/psychiatric disorders, potential pregnancy, or psychoactive

medication use. Additionally, participants were instructed to get no less than six hours of sleep

the night before and refrain from caffeine, alcohol, and tobacco three hours prior to the study.

Upon arrival, participants provided informed consent. Following completion of the study, they

received either course credit or monetary compensation ($15/hour).

Materials

The stimuli have been used in previous work (Payne et al., 2008) with 448 total critical

scene components (objects and backgrounds) selected in this study. The critical scene

components were then partitioned into old (256) and new items (192). The encoding task

consisted of 128 complex valenced scenes (64 negative, 64 neutral) that were created by placing

a negative (e.g., snake) or neutral object (e.g., chipmunk) on a plausible neutral background (e.g.,

forest). The short and long delayed recognition tests each contained a unique set of old and new

scene components (i.e., objects and background in isolation). Studied scenes consisted of the

objects and the background from "whole" scenes, i.e., objects on backgrounds that were

presented together during the study phase. New scene components were drawn from scenes that

were not presented during the study phase. Each recognition test consisted of 224 total scene

components: 32 previously viewed ("same") objects (16 negative, 16 neutral), 32 similar objects

(16 negative, 16 neutral), 64 foil objects (32 negative, 32 neutral), 32 same backgrounds (16

shown with negative object, 16 shown with a neutral object), 32 similar backgrounds (16 similar

objects were shown with a negative object, and 16 with a neutral object), and 32 foil

backgrounds (all neutral). Same components were defined as the exact same background or

object studied during encoding (e.g., identical snake). Similar components were defined as an

alternative version of a background or object that differed in any number of visual features from

those studied during encoding (e.g., a snake that differed in color, positioning, shape, species,

etc.). Note, since backgrounds were always non-emotional, there are no negative backgrounds.

The scene components were never presented twice (i.e., in both tests) and were counterbalanced.

Additionally, participants never saw both a same (e.g., same snake) and similar (e.g., similar

snake) scene during the one test.

Overview of Experimental Procedure

Participants completed the experiment over the course of two sessions separated by

approximately 24-hours. The first session included determination of active motor threshold (see

section Transcranial Magnetic Stimulation below), the encoding task, and a recognition test after

a 30-minute delay ("short delay" test). iTBS was administered during the encoding phase with a

30-minute delay occurring from the completion of the encoding phase to the onset of the

recognition test. Note that simultaneous electroencephalography recordings were obtained during

the encoding and short-delay recognition tasks, but not in the final (24-hour) delayed recognition

test. These data are not the focus of the present manuscript and therefore will not be discussed

further. After the first session, participants returned to the lab approximately 24-hours later to

complete a recognition test on the remaining untested items ("long delay" test).

Emotional Memory Trade-off Task

Encoding Phase (Figure 1a). Participants were instructed that they would be presented

with a series of scenes and to imagine that they were coming across the scenes in real life.

Participants viewed 128 negative and neutral complex scenes. For each scene, they made an

approach/avoid judgment using a 7-point scale (1 = move closer, 7 = move away) to facilitate

incidental encoding. The scenes were presented pseudorandomized in 16 blocks with 8 images

per block (4 negative, 4 neutral). Each trial began with a 100-200ms jitter prior to a 1000ms

fixation cross. Following the fixation cross, a valenced scene was presented for 3000ms and was

replaced with a self-paced avoid/approach judgment scale. Following the 8[th] trial of each block,

participants underwent 2s of iTBS (30 pulses), for a total of 480 pulses over the entire encoding

phase.

Recognition Memory Test. Participants were instructed to make same, similar, or new

judgments to old and new scene components on both the short and long delayed recognition

tests. Participants completed a short practice phase to familiarize themselves with the recognition

test and the same, similar, and new judgments before the recognition test (see Figure 1b). Same

judgments were defined as the exact same background or object studied during encoding (e.g.,

identical snake). Similar judgments were defined as an alternative version of a background or

object that differed in a specific visual detail from those studied during encoding (e.g., a snake

that differed in color, positioning, species, etc.).

Figure 1. (a) Encoding phase. (i) Encoding trial structure and timing. Participants viewed a
fixation cross (1s), followed by a valenced image (3s) and an approach/avoid scale (self-paced).
Stimulation occurred after every 8[th] image for a total of 128 images and 16 blocks. Participants
underwent stimulation to only one site, which was randomly assigned. (ii) The Fz and C3
electrodes were targeted for mPFC (N = 23) and motor control stimulation (N = 22),
respectively. The Fz electrode was selected based on its proximity to the dorsal mPFC region
(MNI: 4, 52, 46) that is functionally connected with the ventral mPFC region (MNI; 4, 56, -8;

indicated by the green crosshair) that was linked to the selective preservation of negative objects
in memory from Payne and Kensinger (2011). **(b)** Recognition task. Same, similar, and new
objects (negative and neutral) and backgrounds (neutral) were presented individually.
Participants responded "same", "similar", or "new". Scene components were presented in
randomized order.
226

Each trial began with a 100-200ms (10ms steps) jittered interval prior to a 1000ms

fixation cross. Following the fixation cross, participants made self-paced same, similar, or new

judgments of individually presented negative objects, neutral objects, or neutral backgrounds.

Objects and backgrounds were presented randomly, but only once, in either the short or long

delay recognition test.

Transcranial magnetic stimulation

A PowerMag EEG 100 TMS stimulator (Mag & More GmbH., Munich, Germany) and a

70 mm figure-eight coil (PMD70-pCool) were used for administration of TMS. Active motor

threshold was obtained by placing the coil over the left motor cortex (near the C3 electrode) and

locating the site that produced visible movement in the right thumb from single pulses of TMS.

Active motor threshold was defined as the percentage of stimulator output that elicited visible

movement in the right thumb on 5 out of 10 trials while participants maintained contraction of

the right thumb and index finger. We determined active motor threshold and iTBS intensity

while participants were wearing the EEG cap, so the coil distance from the stimulation site was

the same during active motor thresholding and the encoding task iTBS. The mean motor

threshold was matched between participants receiving mPFC and MC stimulation (mPFC: $M =$

59%, $SD = 10\%$; MC: $M = 59\%$, $SD = 11\%$; $t(42) = -.25$, $p = .805$). As is standard, the iTBS

protocol was administered at 80% of active motor threshold.

The stimulation site for the dorsal mPFC was determined using a large-scale meta-

analysis of 340 fMRI datasets using Neurosynth.org (Yarkoni, Poldrack, Nichols, Van Essen, &

Wager, 2011). The Fz electrode was selected as the dorsal mPFC stimulation site based on the

Neurosynth meta-analysis that showed the MNI coordinate under the Fz electrode (MNI: 4, 52,

46; Okamoto et al., 2004; Okamoto & Dan, 2005) had strong functional connectivity and

coactivation with ventral mPFC coordinates identified by Payne and Kensinger (2011; MNI = 4,

56, -8; Figure 1b), which was associated with enhanced negative object recognition following

sleep compared to wakefulness. The rationale for targeting the dorsal mPFC is that TMS

modulates neural activity in both the target site and its associated networks (for an example with

TMS stimulation to angular gyrus modulating the hippocampus, which are part of the same

network, see Nilakantan, Bridge, Gagnon, VanHaurents, & Voss, 2017). Thus, we reasoned that

stimulating the dorsal mPFC would also affect the ventral mPFC activity, which is not readily

accessible to TMS. For the control stimulation site, we targeted the MC under the C3 electrode

because it is a common control site (e.g., Daskalakis et al., 2008; Fecchio et al., 2017) and,

importantly, it does not have strong connectivity with the ventral mPFC region that has been

implicated in emotional memory.

The typical offline protocol for iTBS follows 2s on/8s off cycles that deliver 600 pulses

with each 2s of stimulation corresponding to 3 pulses at 50Hz with an inter stimulus interval of

200ms. We adopted a slightly modified iTBS approach with identical 2s periods of stimulation

that was applied in a similar 2s on/8s off manner that was interleaved (between each encoding

block) throughout the encoding phase. Specifically, following each encoding block (8 images),

participants were instructed to close their eyes and relax while the TMS coil was positioned to

the mPFC or MC stimulation site location. For the mPFC condition, the coil was positioned at 0°

from the midline pointing posteriorly to target the Fz electrode. For the MC condition, the coil

was positioned at 45° from the sagittal plane to target the C3 electrode (Figure 1b). Following 2s

of iTBS, participants were instructed to continue with the encoding task with a self-paced button

press. The average time from the last TMS pulse to the onset of the first and last image in the

following encoding block was 8.28 seconds (range: 6.14 - 19.66) and 48.71 seconds (range:

45.62 - 60.01), respectively.

A similar approach has previously been shown to facilitate memory and, importantly, to

not have cumulative effects on performance (Demeter et al., 2016). Prior to the start of the

experimental task, participants received one 2s train of iTBS to become familiarized with the

procedure. No participants reported any adverse response to iTBS.

Data Analysis

For the recognition data, participants' memory scores were calculated for general

recognition (gist-familiarity) and specific recognition (recollection). The contrast between

general and specific recognition scores was our focus because prior research suggests that

overnight sleep exerts its strongest effects on gist-based familiarity, rather than specific

recollection of memory details (Payne et al., 2008). In line with prior studies using a same-

similar-new judgment at retrieval (Garoff et al., 2005; Kensinger et al., 2007b), general

recognition parallels the independence-formulas score commonly used in Remember/Know

paradigms (Yonelinas & Jacoby, 1995) with the following formula:

$$General\ recognition\ hits = \frac{P(\text{similiar}|\text{"same"})}{(1 - P(\text{same}|\text{"same"}))}$$

$$General\ false\ alarms = \frac{P(\text{similiar}|\text{"new"})}{(1 - P(\text{same}|\text{"new"}))}$$

Thus, general hits are calculated based on when participants respond "similar" to

previously viewed items, taking into account when participants respond "same" to previously

viewed items. General false alarms are measured when participants respond "similar" to new

items, taking into account when they respond "same" to new items. Specific recognition was

scored with the following formula:

$$Specific\ recognition\ hits\ =\ P(same|"same")$$

$$Specific\ false\ alarms\ =\ P(same|"new")$$

Thus, specific hits and false alarms are computed when participants respond "same" to

previously viewed items and "same" to new items, respectively.

Since our primary aim was to investigate the effects of mPFC iTBS on emotional

memory we computed recognitions scores for negative objects, neutral objects, the neutral

backgrounds that were originally paired with a negative object, and the neutral backgrounds that

were originally paired with a neutral object. Importantly, these calculations were repeated for the

short and long delay tests. This enabled us to systematically compare how mPFC versus MC

iTBS affected memory for negative and neutral objects, backgrounds in negative and neutral

scenes, across the short and long delay tests. Analyses focused on memory performance for

"same", "similar", and "new" responses to the same and new scene components (i.e., responses

to similar objects and background scenes were not analyzed), to be consistent with previous

work using this paradigm (Kensinger et al., 2007a, 2007b; Waring et al., 2010).

There were statistically comparable 'same' (range = .038-.096) and 'similar' (range =

.153-.258) false alarm rates between the cells in the experiment (all p's > .05), which is

consistent with previous work (Payne et al., 2008). All memory performance analyses were

conducted on corrected recognition scores (hits – false alarms) to obviate issues with response

bias effects. We will refer to these measures as recognition scores from this point forward.

318   Results

319   We tested our *a priori* hypothesis that, relative to MC stimulation, mPFC stimulation

would facilitate recognition for negative objects at a long delay using a 2 (stimulation site:

mPFC, MC) x 2 (valence: negative, neutral) x 2 (scene component: object, background) x 2

(delay: short, long) x 2 (Memory score: general, specific) ANOVA. Analyses were conducted in

R (version 3.6.3.) with the afex (Singmann, Bolker, Westfall, Aust, & Ben-Shachar, 2020) and

emmeans packages (Lenth, 2020). Three participants (mPFC: 2, MC: 1) were excluded from the

analysis due to no observations in one or more response categories for the general recognition

test. Results are considered significant at an alpha level of .05 unless otherwise noted and post

hoc degrees of freedom were estimated with the Kenward-Roger method. The full results are

reported in Table 1, and the primary results of interest are reported below.

The emotional memory trade-off effect

330   In line with prior work (e.g., Payne et al., 2008), we first examined the existence of the

emotional trade-off effect and if this effect remained consistent over time. As expected, we

replicated the emotional memory trade-off effect as our results revealed a significant valence x

scene component interaction, $F(1, 40) = 43.81$ $MSE = .02$, $p < .011$, $\eta_p^2 = .523$ (*Figure 2*).

Follow up *t*-tests revealed that for negative scenes, objects were better remembered than their

accompanying backgrounds, $t(73.2) = 7.67$, $p < .001$, $d = .705$. Critically, this pattern was not

observed for neutral scenes, as memory performance did not differ between objects and

backgrounds, $t(73.2) = .14$, $p = .890$, $d = .013$. Somewhat surprisingly, the scene component x

valence x delay interaction revealed no evidence that the emotional memory trade-off differed

between a short and long delay, $F(1, 40) = 2.19$, $MSE = .02$, $p = .147$, $\eta_p^2 = .052$.

Figure 2. Corrected recognition scores (hits – false alarms) revealed an emotional memory trade-
off effect with greater memory for objects compared to backgrounds in negative scenes.
Additionally, no memory differences emerged for neutral scenes. Error bars represent 95%
confidence intervals. ***$p < .001$.
345

mPFC activity modulates the emotional memory trade-off effect at a long delay

We then examined the causal role of mPFC activity near the time of encoding interacting

with consolidation processes (e.g., modulating the emotional memory trade-off effect). Because

prior work has linked the mPFC in selectively preserving negative information following sleep,

we anticipated that negative objects would be preferentially remembered following a long delay

that included sleep in the mPFC condition relative to the MC condition. In line with our

prediction, our analysis revealed a significant stimulation site x scene component x valence x

delay interaction, which suggests that mPFC compared to MC iTBS modulated the emotional

memory trade-off effect, $F(1, 40) = 6.27$, $MSE = .02$, $p = .016$, $\eta_p^2 = .136$ (*Figure 3*).

To parse out the 4 way-interaction, follow up *t*-tests compared memory differences

between the mPFC vs. MC stimulation conditions at each of the other factors (scene component,

valence, delay). At the short delay, we observed no memory differences between the mPFC and

MC stimulation site conditions for negative objects, $t(110) = 1.30$, $p = .196$, $d = .313$, or their

backgrounds, $t(110) = -.02$, $p = .988$, $d = -.004$. Similarly, no memory differences emerged

between the mPFC and MC stimulation site condition for neutral objects, $t(110) = 1.49$, $p = .139$,

$d = .358$, or backgrounds, $t(110) = -1.46$, $p = .147$, $d = -.351$. Thus, we found no evidence that

mPFC iTBS modulated the emotional memory trade-off effect at a short delay that did not

include sleep (collapsed across memory score).

In line with our predictions, at a long delay, mPFC compared to MC stimulation

facilitated memory for negative objects, $t(110) = 2.78$, $p = .006$, $d = .669$, with no differences

emerging for their backgrounds, $t(110) = .33$, $p = .739$, $d = .081$. Importantly, we found no

evidence for memory differences for neutral objects, $t(110) = 1.47$, $p = .145$, $d = .353$, or

backgrounds, $t(110) = 1.36$, $p = .175$, $d = .328$. This pattern of results suggests that iTBS to the

mPFC (compared to the MC) modulated the emotional memory trade-off effect by selectively

preserving memory for negative objects, but not backgrounds, following a long delay that

included a night of sleep (collapsed across specific and general memory scores). This effect was

only found at the long delay, as no differences emerged for objects or backgrounds when

memory was assessed at the short delay. Importantly, for neutral scenes, no differences between

stimulation conditions emerged for objects or backgrounds when memory was assessed at either

a short or long delay.

Figure 3. Corrected recognition scores (collapsed across memory score) revealed that mPFC
compared to MC iTBS facilitated memory for negative objects at a long delay. No memory
differences were found between negative objects at a short delay for mPFC vs. MC control
stimulation. Error bars represent 95% confidence intervals. *p < .05.
381
The role of mPFC activity in memory specificity

Next, we sought to address if mPFC stimulation differentially modulated the emotional

memory trade-off effect at a short or long delay when assessed on specific vs. general memory.

The 5 way interaction found no evidence that mPFC vs. MC stimulation at a short or long delay

differentially modulated the emotional memory trade-off effect when memory was assessed on

specific and general recognition scores, $F(1, 40) = .12$, $MSE = .02$, $p = .727$, $\eta_p^2 = .003$. Lastly,

we examined if mPFC iTBS differentially modulated the emotional memory trade-off effect

collapsed across delay when corrected recognition scores were assessed on specific vs. general

information (gist). Our results revealed a marginally significant the 4-way interaction among

stimulation site, valence, scene component, and memory type, $F(1, 40) = 4.04$, $MSE = .02$, $p =$

.051, $\eta_p^2 = .092$. Although this did not reach significance (and requires further investigation),

there appeared to be a numeric trend such that mPFC stimulation modulated the emotional

memory trade-off effect for specific vs. general recognition (collapsed across short and long

delays).

Summary of Results

Collectively, the findings revealed that iTBS to the mPFC selectively facilitated

recognition of negative information after a long delay while no differences emerged for their

accompanying backgrounds. Critically, no memory differences emerged when memory was

assessed following a short delay in negative scenes or at either a short or long delay in neutral

scenes. Together, these findings suggest that modulating mPFC activity during encoding

selectively facilitated emotional memories.

404   **General Discussion**

This study investigated the causal role of the mPFC in the preferential encoding and

consolidation of emotionally salient information. Our results provide preliminary evidence that

the mPFC is causally involved in the encoding of negatively valenced scenes. Specifically,

stimulating the mPFC, but not the MC, during encoding with iTBS enhanced memory for

negative objects only after a 24-hour delay (filled with sleep). Moreover, iTBS to the mPFC had

no detectable effect on memory for negative stimuli on the short delay (30 minute) test, on

neutral information on either test, or on the background scene component in any condition.

Notably, false alarm rates did not differ between any of the conditions, which provides evidence

that our results are due to effects on recognition, rather than differences in response bias between

the mPFC and MC groups. We interpret these findings as being consistent with the idea that

mPFC activity near the time of encoding potentiates subsequent consolidation processes, which

together facilitate memory for negative information (Payne & Kensinger, 2010, 2011, 2018).

The results reported here also partially coverage with recent work suggesting a causal

role for the mPFC in gist-based false memories (Bovy et al., 2020; for related findings, see

Berkers et al., 2017). Bovy and colleagues (2020) examined the relationship between gist-based

emotional false memories and mPFC by stimulating the mPFC using an inhibitory TMS

protocol, specifically continuous theta-burst stimulation (cTBS). Prior to encoding in an

emotionally-valenced Deese-Roediger-McDermott (DRM) paradigm, participants underwent a

negative mood induction and received TMS to the mPFC using the above mentioned cTBS or a

control (i.e., 5Hz rTMS protocol). They found that cTBS to mPFC during encoding reduced false

recognition of negative critical lures on a delayed test (~24-hour) that included a sleep interval.

The authors took these findings as evidence that the mPFC plays an important role in extracting

schematic (or gist) information during encoding. Our findings extend the results described above.

Here, we observed iTBS – a TMS protocol thought to be associated with increased excitation

("long-term potentiation", Wischnewski & Schutter, 2015) of the stimulated region – delivered

to the mPFC during encoding facilitated memory for negative objects on a 24-hour delayed test.

Moreover, we demonstrate that mPFC iTBS benefits emotional episodic memory by using a

memory task consisting of complex emotional scenes as opposed to semantically related word

lists. Most importantly, in addition to the 24-hour delayed test, we also included a short (30

minute) delayed memory test to better establish that mPFC activity during encoding interacted

with subsequent consolidation processes. We observed that mPFC iTBS at encoding selectively

enhanced negative memory only on the 24-hour delayed memory test. Our findings suggest that

mPFC activity near the time of encoding interacts with downstream consolidation processes to

support retention of emotional memories.

The present results also provide support for theories of the selective consolidation of

emotional information (Kim & Payne, 2020; Payne & Kensinger, 2018; Richter-Levin & Akirav,

2003). Specifically, excitatory stimulation (i.e., neuromodulation) of the dorsal mPFC during

encoding may have affected subsequent memory performance by upregulating regions important

for emotional memory encoding and retrieval, including the ventral mPFC, amygdala and

hippocampus (Bennion et al., 2015; Kensinger & Corkin, 2004; Murty et al., 2010; Payne &

Kensinger, 2010). Future work could directly test this possibility by examining the effects of

TMS in conjunction with fMRI measures of activity and connectivity.

There are three limitations of the present study that warrant mention. First, iTBS was

interspersed throughout the encoding task to attempt to modulate brain regions, specifically the

MTL, that are functionally connected with the mPFC and critically involved in the encoding of

emotional information. Although we aimed to modulate mPFC-MTL activity during the

encoding phase it is possible that our stimulation procedure may have also altered neural activity

during early consolidation and/or retrieval processes (i.e., during the short delayed recognition

test) due to iTBS protocols having long lasting effects that extend 20-60 minutes following

stimulation. Even though we are unaware of any evidence to suggest that the short modified

iTBS protocol implemented in the present study modulated neural activity beyond the encoding

phase (e.g., modulating retrieval related processes) different stimulation protocols may provide

further insights into the role of the mPFC activity near the time of encoding in selectively

preserving negative information following a sleep filled delay. Furthermore, it is unclear if our

results would differ had we used a different TMS protocol. For example, as described above,

Bovy and colleagues found a causal role of the mPFC in gist-based false memories implementing

cTBS while our iTBS protocol revealed a causal role of the mPFC in emotional memory

(collapsed across specific and gist-based memories). Of relevance to this point, a recent meta-

analysis of TMS effects on episodic memory found that offline 1 Hz rTMS led to larger

enhancing effects on episodic memory compared to other stimulation protocols, including iTBS

(Yeh & Rose, 2019). Second, we only examined the effects of mPFC stimulation during memory

encoding. It is possible that mPFC stimulation during a different stage of memory, such as

during post-encoding periods, would lead to a different pattern of results. To draw stronger

causal claims about the role of mPFC in emotional memory, we used an active control

stimulation site (left MC) instead of another common control site (i.e., vertex) or sham

stimulation. Prior findings have suggested vertex stimulation may modulate activity in the

default mode network (Jung et al., 2016), which is involved in emotion (Sheline et al., 2009) and

memory processes (Rugg & Vilberg, 2013). Although we cannot rule out the possibility that MC

stimulation contributed to the effects on memory by impairing recognition of negative objects,

we find this possibility unlikely for two reasons. One, memory performance in the MC

stimulation group reported here was comparable to memory performance in a no-stimulation

control group in a pilot study (not reported). Two, and most importantly, MC is not associated

with emotional memory. Lastly, future work is needed to determine if different stimulation

protocols (e.g., frequency, intensity, control stimulation site, stage of memory) will contribute to

further insights about the role of encoding-consolidation interactions involving the mPFC and its

associated network.

Future work should conduct a more direct investigation of the relationship between

mPFC activity during encoding and sleep-based consolidation in emotional memory. Sleep

affords an ideal environment for offline consolidation to transform, integrate, and preserve

salient and future relevant information (Payne, 2011). Synchronous neural oscillations during

sleep (e.g., theta oscillations, slow oscillations, spindles) have been observed in the amygdala,

hippocampus, and prefrontal cortex, and may facilitate synaptic plasticity and selective memory

consolidation during sleep (Kim & Payne, 2020; Rasch & Born, 2013). Examining the macro

and microarchitecture of sleep in conjunction with causal methods like TMS will provide critical

insight into the neural mechanisms of emotional memory encoding and consolidation.

490   In conclusion, the results of this study provide preliminary evidence for a causal role of

491   the mPFC in selectively preserving negative memories. The current work moves beyond

492   correlational findings and provides initial causal support for theories suggesting that activity and

493   connectivity in an emotional memory network (e.g., mPFC-MTL) near the time of encoding

494   interact with subsequent consolidation processes (perhaps during sleep) to form long-lasting

495   emotional memories (Kim & Payne, 2020; Payne & Kensinger, 2018).

References

Ballarini, F., Martínez, M. C., Díaz Perez, M., Moncada, D., & Viola, H. (2013). Memory in elementary school children is improved by an unrelated novel experience. *PloS one*, *8*(6), e66875. https://doi.org/10.1371/journal.pone.0066875

Bennion, K. A., Mickley Steinmetz, K. R., Kensinger, E. A., & Payne, J. D. (2015). Sleep and cortisol interact to support memory consolidation. *Cerebral Cortex*, *25*(3), 646–657. https://doi.org/10.1093/cercor/bht255

Berkers, R. M. W. J., Klumpers, F., & Fernández, G. (2016). Medial prefrontal–hippocampal connectivity during emotional memory encoding predicts individual differences in the loss of associative memory specificity. *Neurobiology of Learning and Memory*, *134*, 44–54. https://doi.org/10/ggk792

Berkers, R. M. W. J., van der Linden, M., de Almeida, R. F., Müller, N. C. J., Bovy, L., Dresler, M., Morris, R. G. M., & Fernández, G. (2017). Transient medial prefrontal perturbation reduces false memory formation. *Cortex*, *88*, 42–52. https://doi.org/10.1016/j.cortex.2016.12.015

Bovy, L., Berkers, R. M. W. J., Pottkämper, J. C. M., Varatheeswaran, R., Fernández, G., Tendolkar, I., & Dresler, M. (2020). Transcranial magnetic stimulation of the medial prefrontal cortex decreases emotional memory schemas. *Cerebral Cortex*, bhz329. https://doi.org/10.1093/cercor/bhz329

Daskalakis, Z. J., Farzan, F., Barr, M. S., Maller, J. J., Chen, R., & Fitzgerald, P. B. (2008). Long-interval cortical inhibition from the dorsolateral prefrontal cortex: A TMS-EEG study. *Neuropsychopharmacology*, *33*(12), 2860–2869. https://doi.org/10.1038/npp.2008.22

Demeter, E., Mirdamadi, J. L., Meehan, S. K., & Taylor, S. F. (2016). Short theta burst

stimulation to left frontal cortex prior to encoding enhances subsequent recognition

memory. *Cognitive, Affective, & Behavioral Neuroscience*, *16*(4), 724–735.

https://doi.org/10.3758/s13415-016-0426-3

Dolcos, F., LaBar, K. S., & Cabeza, R. (2004). Dissociable effects of arousal and valence on

prefrontal activity indexing emotional evaluation and subsequent memory: An event-

related fMRI study. *NeuroImage*, *23*(1), 64–74. https://doi.org/10/ckkrtb

Dunsmoor, J. E., Murty, V. P., Davachi, L., & Phelps, E. A. (2015). Emotional learning

selectively and retroactively strengthens memories for related events. *Nature*, *520*(7547),

345–348. https://doi.org/10.1038/nature14106

Fecchio, M., Pigorini, A., Comanducci, A., Sarasso, S., Casarotto, S., Premoli, I., Derchi, C.-C.,

Mazza, A., Russo, S., Resta, F., Ferrarelli, F., Mariotti, M., Ziemann, U., Massimini, M.,

& Rosanova, M. (2017). The spectral features of EEG responses to transcranial magnetic

stimulation of the primary motor cortex depend on the amplitude of the motor evoked

potentials. *PloS One*, *12*(9), e0184910. https://doi.org/10.1371/journal.pone.0184910

Garoff, R. J., Slotnick, S. D., & Schacter, D. L. (2005). The neural origins of specific and general

memory: The role of the fusiform cortex. *Neuropsychologia*, *43*(6), 847–859.

https://doi.org/10.1016/j.neuropsychologia.2004.09.014

Goto, Y., & Grace, A. A. (2008). Limbic and cortical information processing in the nucleus

accumbens. *Trends in Neurosciences*, *31*(11), 552–558. https://doi.org/10/b4tx84

Hermiller, M. S., Chen, Y. F., Parrish, T. B., & Voss, J. L. (2020). Evidence for immediate

enhancement of hippocampal memory encoding by network-targeted theta-burst

stimulation during concurrent fMRI. *Journal of Neuroscience*, *40*(37), 7155-7168.

Hu, P., Stylos-Allan, M., & Walker, M. P. (2006). Sleep facilitates consolidation of emotional

declarative memory. *Psychological Science*, *17*(10), 891–898. https://doi.org/10/dkqxs9

Huang, Y.-Z., Edwards, M. J., Rounis, E., Bhatia, K. P., & Rothwell, J. C. (2005). Theta burst

stimulation of the human motor cortex. *Neuron*, *45*(2), 201–206.

https://doi.org/10/d2gpww

Jung, J., Bungert, A., Bowtell, R., & Jackson, S. R. (2016). Vertex stimulation as a control site

for transcranial magnetic stimulation: A Concurrent TMS/fMRI Study. *Brain*

Stimulation, *9*(1), 58–64. https://doi.org/10.1016/j.brs.2015.09.008

Kensinger, E. A., & Corkin, S. (2004). Two routes to emotional memory: Distinct neural

processes for valence and arousal. *Proceedings of the National Academy of Sciences of*

the United States of America, *101*(9), 3310–3315. https://doi.org/10/ftzx6z

Kensinger, E. A., Garoff-Eaton, R. J., & Schacter, D. L. (2007a). Effects of emotion on memory

specificity: Memory trade-offs elicited by negative visually arousing stimuli. *Journal of*

Memory and Language, *56*(4), 575–591. https://doi.org/10.1016/j.jml.2006.05.004

Kensinger, E. A., Garoff-Eaton, R. J., & Schacter, D. L. (2007b). How negative emotion

enhances the visual specificity of a memory. *Journal of Cognitive Neuroscience*, *19*(11),

1872–1887. https://doi.org/10.1162/jocn.2007.19.11.1872

Kim, S. Y., & Payne, J. D. (2020). Neural correlates of sleep, stress, and selective memory

consolidation. *Current Opinion in Behavioral Sciences*, *33*, 57–64.

https://doi.org/10/ggmtgv

Lenth, R. (2020). Emmeans: Estimated marginal means, aka least-squares means. Retrieved from

https://CRAN.R-project.org/package=emmeans

Murty, V. P., Ritchey, M., Adcock, R. A., & LaBar, K. S. (2010). fMRI studies of successful

emotional memory encoding: A quantitative meta-analysis. *Neuropsychologia*, *48*(12),

3459–3469. https://doi.org/10/dwkvv8

Nilakantan, A. S., Bridge, D. J., Gagnon, E. P., VanHaerents, S. A., & Voss, J. L. (2017).

Stimulation of the posterior cortical-hippocampal network enhances precision of memory

recollection. *Current Biology*, *27*(3), 465–470. https://doi.org/10.1016/j.cub.2016.12.042

Okamoto, M., Dan, H., Sakamoto, K., Takeo, K., Shimizu, K., Kohno, S., Oda, I., Isobe, S.,

Suzuki, T., Kohyama, K., & Dan, I. (2004). Three-dimensional probabilistic anatomical

cranio-cerebral correlation via the international 10–20 system oriented for transcranial

functional brain mapping. *NeuroImage*, *21*(1), 99–111.

https://doi.org/10.1016/j.neuroimage.2003.08.026

Okamoto, M., & Dan, I. (2005). Automated cortical projection of head-surface locations for

transcranial functional brain mapping. *NeuroImage*, *26*(1), 18–28.

https://doi.org/10.1016/j.neuroimage.2005.01.018

Patil, A., Murty, V. P., Dunsmoor, J. E., Phelps, E. A., & Davachi, L. (2016). Reward

retroactively enhances memory consolidation for related items. *Learning &*

memory, *24*(1), 65–69. https://doi.org/10.1101/lm.042978.116

Payne, J. D., Chambers, A. M., & Kensinger, E. A. (2012). Sleep promotes lasting changes in

selective memory for emotional scenes. *Frontiers in Integrative Neuroscience*, *6*.

https://doi.org/10/ggk9f7

Payne, J. D., & Kensinger, E. A. (2010). Sleep's role in the consolidation of emotional episodic

memories. *Current Directions in Psychological Science*, *19*(5), 290–295.

https://doi.org/10/b6xkvt

Payne J. D. (2011). Sleep on it!: Stabilizing and transforming memories during sleep. *Nature*

Neuroscience, 14(3), 272–274. https://doi.org/10.1038/nn0311-272

Payne, J. D., & Kensinger, E. A. (2011). Sleep leads to changes in the emotional memory trace:

Evidence from fMRI. *Journal of Cognitive Neuroscience*, *23*(6), 1285–1297.

https://doi.org/10/btvfds

Payne, J. D., & Kensinger, E. A. (2018). Stress, sleep, and the selective consolidation of

emotional memories. *Current Opinion in Behavioral Sciences*, *19*, 36–43.

https://doi.org/10/ggk8cj

Payne, J. D., Stickgold, R., Swanberg, K., & Kensinger, E. A. (2008). Sleep preferentially

enhances memory for emotional components of scenes. *Psychological Science*, *19*(8),

781–788. https://doi.org/10.1111/j.1467-9280.2008.02157.x

Rasch, B., & Born, J. (2013). About sleep's role in memory. *Physiological Reviews*, *93*(2), 681–

766. https://doi.org/10.1152/physrev.00032.2012

Richter-Levin, G., & Akirav, I. (2003). Emotional tagging of memory formation—In the search

for neural mechanisms. *Brain Research Reviews*, *43*(3), 247–256.

https://doi.org/10/c7v5xn

Rossi, S., Hallett, M., Rossini, P. M., & Pascual-Leone, A. (2009). Safety, ethical considerations,

and application guidelines for the use of transcranial magnetic stimulation in clinical

practice and research. *Clinical Neurophysiology*, *120*(12), 2008–2039.

https://doi.org/10/fwp29f

Rugg, M. D., & Vilberg, K. L. (2013). Brain networks underlying episodic memory retrieval.

Current Opinion in Neurobiology, *23*(2), 255–260.

https://doi.org/10.1016/j.conb.2012.11.005

Sack, A. T., Cohen Kadosh, R., Schuhmann, T., Moerel, M., Walsh, V., & Goebel, R. (2009).

Optimizing functional accuracy of TMS in cognitive studies: A comparison of methods.

Journal of Cognitive Neuroscience, *21*(2), 207–221.

https://doi.org/10.1162/jocn.2009.21126

Sheline, Y. I., Barch, D. M., Price, J. L., Rundle, M. M., Vaishnavi, S. N., Snyder, A. Z., Mintun,

M. A., Wang, S., Coalson, R. S., & Raichle, M. E. (2009). The default mode network and

self-referential processes in depression. *Proceedings of the National Academy of*

Sciences, *106*(6), 1942–1947. https://doi.org/10.1073/pnas.0812686106

Singmann, H., Bolker, B., Westfall, J., Aust, F., & Ben-Shachar, M. S. (2020). Afex: Analysis of

factorial experiments. Retrieved from https://CRAN.R-project.org/package=afex

Stanny, C. J., & Johnson, T. C. (2000). Effects of stress induced by a simulated shooting on

recall by police and citizen witnesses. *The American Journal of Psychology*, *113*(3), 359–

386. https://doi.org/10/cmbhhf

Sterpenich, V., Albouy, G., Darsaud, A., Schmidt, C., Vandewalle, G., Dang Vu, T. T.,

Desseilles, M., Phillips, C., Degueldre, C., Balteau, E., Collette, F., Luxen, A., & Maquet,

P. (2009). Sleep promotes the neural reorganization of remote emotional memory. *The*

Journal of Neuroscience, *29*(16), 5143–5152. https://doi.org/10/c3s6w3

Tang, Y., Jiao, X., Wang, J., Zhu, T., Zhou, J., Qian, Z., Zhang, T., Cui, H., Li, H., Tang, X., Xu,

L., Zhang, L., Wei, Y., Sheng, J., Liu, L., & Wang, J. (2019). Dynamic functional

connectivity within the fronto-limbic network induced by intermittent theta-burst

stimulation: A pilot study. *Frontiers in neuroscience*, *13*, 944.

https://doi.org/10.3389/fnins.2019.00944

Tambini, A., Rimmele, U., Phelps, E. A., & Davachi, L. (2017). Emotional brain states carry

over and enhance future memory formation. *Nature neuroscience*, *20*(2), 271–278.

https://doi.org/10.1038/nn.4468

Waring, J. D., Payne, J. D., Schacter, D. L., & Kensinger, E. A. (2010). Impact of individual

differences upon emotion-induced memory trade-offs. *Cognition & Emotion*, *24*(1), 150–

167. https://doi.org/10.1080/02699930802618918

Wischnewski, M., & Schutter, D. J. (2015). Efficacy and time course of theta burst stimulation in

healthy humans. Brain stimulation, 8(4), 685-692.

Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-

scale automated synthesis of human functional neuroimaging data. *Nature methods*, 8(8),

665-670.

Yeh, N., & Rose, N. S. (2019). How can transcranial magnetic stimulation be used to modulate

episodic memory?: A systematic review and meta-analysis. *Frontiers in Psychology*, *10*.

https://doi.org/10.3389/fpsyg.2019.00993

Yonelinas, A. P., & Jacoby, L. L. (1995). The relation between remembering and knowing as

bases for recognition: Effects of size congruency. *Journal of Memory and Language*,

34(5), 622–643. https://doi.org/10/b3gq9t

Table 1. *ANOVA results*

Effects	MSE	F(1, 40)	η_p^2	p
Main effects				
Stimulation site	.27	1.42	.034	.241
Valence	.03	6.17	.134	.017*
Scene component	.04	21.77	.352	< .001***
Delay	.03	109.87	.733	< .001***
Memory score	.09	86.98	.685	< .001
Two-way interactions				

Stimulation site x Valence	.03	.62	.015	.434
Stimulation site x Scene component	.04	7.64	.160	.009**
Stimulation site x Delay	.03	5.01	.111	.031
Stimulation site x Memory score	.09	.95	.023	.336
Valence x Scene component	.02	43.81	.523	< .001***
Valence x Delay	.02	2.79	.065	.103
Valence x Memory score	.03	5.97	.130	.019*
Scene component x Delay	.03	4.51	.101	.040*
Scene component x Memory score	.03	.35	.009	.555
Delay x Memory score	.04	34.02	.460	< .001***
Three-way interactions				
Stimulation site x Valence x Scene component	.02	.15	.004	.697
Stimulation site x Valence x Delay	.02	.30	.007	.589
Stimulation site x Valence x Memory score	.03	5.13	.114	.029*
Stimulation site x Scene component x Delay	.03	.79	.019	.381
Stimulation site x Scene component x Memory score	.03	3.43	.079	.071
Stimulation site x Delay x Memory score	.04	.20	.005	.655
Valence x Scene component x Delay	.02	2.19	.052	.147
Valence x Scene component x Memory score	.02	37.40	.483	< .001***
Valence x Delay x Memory score	.03	1.76	.042	.192
Scene component x Delay x Memory score	.03	< .01	< .001	.959
4-way interactions				
Stimulation site x Valence x Scene component x Delay	.02	6.27	.136	.016*
Stimulation site x Valence x Scene component x Memory score	.02	4.04	.092	.051
Stimulation site x Valence x Delay x Memory score	.03	.09	.002	.760
Stimulation site x Scene component x Delay x Memory score	.03	.01	< .001	.917
Valence x Scene component x Delay x Memory score	.02	1.10	.027	.301
5-way interaction				
Stimulation site x Valence x Scene component x Delay x Memory score	.02	.12	.003	.727

652 Note. Degrees of freedom were the same for all effects. * *p* < .05, ** *p* < .01,*** *p* < .001.

653