Alternate primers for whole-genome SARS-CoV-2 sequencing

Matthew Cotten*, Dan Lule Bugembe, Pontiano Kaleebu, My V.T. Phan

Author affiliations: UK Medical Research Council–Uganda Virus Research Institute
and London School of Hygiene and Tropical Medicine Uganda Research Unit,
Entebbe, Uganda (D. Bugembe, P. Kaleebu, M. Cotten); Uganda Virus Research
Institute, Entebbe (P. Kaleebu); Erasmus Medical Center, Rotterdam, the
Netherlands (M.V.T. Phan); UK Medical Research Council–University of Glasgow
Centre for Virus Research, Glasgow, Scotland, UK (M. Cotten)

*Correspondence to Matthew Cotten, MRC/UVRI & LSHTM Uganda Research
Unit, Entebbe, Uganda email: matthew.cotten@lshtm.ac.uk

Abstract

As the world is struggling to control the novel Severe Acute Respiratory Syndrome
Coronavirus 2 (SARS-CoV-2), there is an urgency to develop effective control measures.
Essential information is encoded in the virus genome sequence with accurate and complete
SARS-CoV-2 sequences essential for tracking the movement and evolution of the virus and
for guiding efforts to develop vaccines and antiviral drugs. While there is unprecedented
SARS-CoV-2 sequencing efforts globally, approximately 19 to 43% of the genomes
generated monthly are gapped, reducing their information content. The current study
documents the genome gap frequencies and their positions in the currently available data
and provides an alternative primer set and a sequencing scheme to helps improve the quality
and coverage of the genomes.

Keywords: SARS-CoV-2; COVID-19, primers; next generation sequencing;

Introduction

Since the first report on 30th December 2019 in Wuhan China and the WHO declaration of the pandemic on 12th March 2020, the novel Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) (1) and the associated disease Coronavirus Disease 2019 (COVID-19) (2)(3) have continued to spread throughout the world, causing >35 million infections and >1,000,000 death globally (4). The virus genome sequences carry important information, which can be used to interpret the virus transmission, evolution patterns and origin tracing. Furthermore, accurate and complete genomic sequences are essential for monitoring diagnostics and developing novel therapeutics and vaccines. We have seen an unprecedented amount of virus sequencing with over 130,000 complete or nearly complete genome sequences of SARS-CoV-2 now available in the GISAID database by the end of September 2020 (5). Most of the sequences have been generated by next generation sequencing using targeted amplicon methods. A scan through SARS-CoV-2 genomes from GISAID with the filter "complete genome" revealed a high frequency of gaps occurring across the genome, influencing the overall genomics quality and interpretation. Here we describe an alternate primer scheme for whole-genome sequencing to improve the genome sequence quality and coverage.

Documenting the problem

We retrieved genomes deposited to GISAID in September 2020 (9 months into the pandemic), using the "complete genome" filter and sorting the genomes by sequencing platforms information included in the metadata. Figure 1 illustrated the positions across the 30kb genome of every stretch of 200 Ns (N200; nearly the size of an amplicon) in the first genomes deposited in September 2020 using the Illumina platform (Panel A; N = 3000) versus MinION platform (Panel B; N = 3000). Histograms of gap frequencies across the genomes are shown for each platform. The gaps are not randomly distributed but occur with higher frequency in a subset of positions across the genome. Although genomes generated by the

two platforms (Illumina and MinION) show similar problem regions (nt 8000-11000, and nt

19,000-24,000 relative to the reference genome NC_045512), the patterns are not completely

identical. Given the use of several primer amplification schemes, we suspect the gaps in

coverage may be due to unexpected primer interactions, complicated sequence regions (odd

composition or secondary structure), issues with primer trimming during quality control of read

data, or some combinations of these factors.

The phenomenon is unlikely to be due to an isolated set of genomes as we observed similar N200 frequencies in genomes submitted from each month of the pandemic (Table 1), suggesting that gaps in coverage is a more general phenomenon. Of note, genomes generated using Ion Torrent show much lower levels of N200 (Table 1). The very low frequency of large gaps in the Ion Torrent data may be due to the use of a dedicated alternative primer set (6). There have been discussions and reports on the SARS-CoV-2 genome changes due to sequencing errors as well as long gaps in the genomes due to missing amplicons from the amplicon-approach sequencing (7) (8) (9). Updates of the ARTIC primers have been presented in late March 2020 to address these issues (10). However, the percentage of reported complete genomes in GISAID with 1 or more N200s continues, with 10,611 (28%) of the 38,228 genomes deposited in September 2020 having 1 or more N200 gaps (Table 1), indicating the challenges remain largely unsolved.

Table 1. Frequency of SARS-CoV-2 genomes with 1 or more 200 nt gaps (N200) by month and by sequencing platform.

Deposition period	Complete genomes	Genomes with 1 or more N200[2]	%. with 1 or more N200[3]	Illumina total[4]	Illumina % with N200[5]	MinION total[6]	MinION % with N200[7]	Ion Torrent total[8]	Ion Torrent % with N200[9]	Method unclear[10]	Method unclear % with N200[11]
1Jan20-31Jan20	54	1	1.85	9	0	2	0	0	0	43	2
1Feb20-29Feb20	126	2	1.59	44	0	9	0	5	0	65	4
1Mar20-31Mar20	2872	559	19.46	1518	16	548	32	35	6	771	18
1Apr20-30Apr20	12411	3745	30.17	4970	38	1286	27	264	0	5424	26
1May20-31May20	19787	8606	43.49	8634	52	2634	30	529	0	7990	42
1Jun20-30Jun20	21665	8723	40.26	7043	36	3844	35	629	2	10149	47
1Jul20-31Jul20	17986	4834	26.88	4965	23	1585	33	471	2	10965	29
1Aug20-31Aug20	17276	4005	23.18	11074	22	2270	26	486	0	3446	28
1Sep20-30Sep20	38227	10611	27.76	22740	23	7973	44	580	1	6934	28

1. Number of genomes with the annotation "complete" retrieved from GISAID (https://www.gisaid.org/).

2. Genomes were sorted by the presence or absence of the sequence N200.

3. ((The number of genomes with at least 1 N200)/total number of genomes)*100.

4. Number of genomes in GISAID for this period generated using any of the Illumina methods as noted in the GISAID "Sequencing technology metadata" .

5. ((The number of Illumina genomes with at least 1 N200)/total number of genomes)*100.

6.,7. Number of genomes in GISAID for this period generated using any of the MinION methods as noted in the GISAID "Sequencing technology metadata" and their percentage.

8.,9. Number of genomes in GISAID for this period generated using any of the Ion Torrent methods as noted in the GISAID "Sequencing technology metadata" and their percentage.

10.,11. Number of genomes in GISAID for this period generated using unclear methods as noted in the GISAID "Sequencing technology metadata" and their percentage.

Detailed analysis of gaps.

A more focused analysis of the frequent gaps is provided in Figure 2. The gap pattern between nt 19,000 and 24,000 (relative to the reference genome NC_045512) is shown for both MinION and Illumina sequences (first 3000 genomes of each deposited in September 2020). For reference the positions of the ARTIC primers (v.1) in the region are indicated (middle panel). A histogram of gap start positions (top panel) and the individual genome gaps (bottom panel) are also shown. The peaks are all between Forward from Amplicon X and Reverse from Amplicon X-1, for both MinION and Illumina data. Some of the gaps span the entire amplicons (the region between primerL and primerR, but the peaks of gap regions frequently span the region between Forward primer from amplicon X and Reverse primer from Amplicon X-1. Possible explanation for these gaps could be: (1) The primer pair fails at the amplification and no amplicon is available to sequence; or (2) The amplicons sequence is in

109   the raw data but a step in the data analysis, perhaps the primer trimming process, results in a

110   gap in the data.

A. MinION data

B. Illumina data

Figure 2. **Positions of 200nt gaps across SARS-CoV-2 genomes stratified by MinION or**

Illumina, in region 19000 to 24000. Genomes deposited in September 2020 as "complete

were retrieved from GISAID, sorted by sequencing platform and the presence of stretches of

N200 were monitored. For clarity only the first 3000 genomes in each set were plotted. The

upper histogram shows the frequency (in 30 nt bins) of N200 motifs across the genome region.

The middle panel shows the positions and names of ARTIC primers colored by forward or

reverse, the bottom panel shows start positions of 200 nt gap by genome, in the first 3000

genomes of each set. Panel A: MinION derived genome sequences, Panel B: Illumina-derived

genome sequences.

121
Alternate primers as a potential solution to avoid gapped genomes.

We explored an alternate set of amplification primers (termed the **Entebbe primers**)

designed using methods we had previously used for MERS-CoV (11), Norovirus (12), RSV

(13) and Yellow Fever virus (14). Important for the design were the amplicon size and the

primer placement. For their implementation, the use of primers for the reverse transcription

step, and the multiplexing of the amplicons in two staggered sets was important for the PCR.

Our experience had suggested an optimum amplicon size of around 1500 bp. The larger

amplicons reduced the total primers content of the reactions but still allowed high reverse

transcription efficiency (which, in our hands, declined beyond 1500 nt). Here we describe

primers designed for whole-genome sequencing of SARS-CoV-2, as well as sharing the

detailed laboratory methods that we used for reverse transcription, PCR amplification and

MinION library preparation to successfully sequence the SARS-CoV-2 genome.

A.

Step 1 Prepare potential primers

Input: all complete genomes (no Ns)
(n= 17220 genomes)

Slices into 33 nt fragments, 1 nt step
(n = 606389 33mers)

Counts frequency, retains value

Trims to Tm 58 °C

Removes inappropriate sequences
(bad GC, homopolymers)

Prepares unique set

Output: Potential primers
(n = 29860 primers)

Step 2 Select optimum primers

Input: Potential primers *(n = 29860)*
Reference genome,
    Choose number of amplicons,
    Choose primers per bin

Selects amplicons, primer bin positions

Selects X highest conserved primers per bin
(based on frequency)

Prepares rev/comp of reverse primers

Output: Fasta file of primers
(for 20 amplicons, 2 primers/bin = 80 primers)

B.

Figure 3. Primer design and amplicon layout. Panel A. The two main steps involving in
primers generation and selection are shown. Panel B. The layout of the 20 amplicons across
the SARS-CoV-2 genome is shown in lower panel. The blue markers indicate target positions
in the SARS-CoV-2 genome (NC_045512 used here), the grey bars indicate the resulting
amplicon.

Briefly, the primer design (Figure 3A) started with the set of complete SARS-CoV-2
genome sequences available in the GISAID database on 22 June 2020 (N = 21,687). Spaces
and disruptive characters were removed from the sequence IDs and the sequences were
further screened to remove genomes containing gaps of 6Ns or more, resulting in 17,220

clean genome sequences. Next, all sequences were sliced into 33nt strings (33mers), with a

1nt step and 606,389 unique 33mers were generated. The frequency of each 33mer was

counted to identify highly conserved 33mers. This counting method avoids the multiple

sequence alignment step commonly used in primer design and becomes prohibitive with large

and or diverse genome sets. This alignment-free approach allowed us to use all suitable

genome sequences of interest rather than a set that could be conveniently aligned. Finally,

primer-like 33mers sequences were generated by trimming the sequences to a calculated

desired melting temperature and removing any primers greater than 26nt.

In the second step we defined forward and reverse primer target regions (bins)

for the amplicons. For SARS-CoV-2, we selected 20 amplicons with an overlap of

300nt, regularly spaced across the SARS-CoV-2 genome sequence (Figure 3B). We

then selected the top conserved primer sequences (the highest frequency primers)

mapping in the 5' or 3' 185nt of each amplicon. For security, the two highest frequency

primers per bin were selected for the SARS-CoV-2 sequence, this provided some

insurance against primer failure either due to target evolution or unexpected

secondary structure. The binning and primer target locations for the final set of primers

are shown in Figure 2 and the final calculated amplicon lengths were 1495-2093nt. In

the second step we defined forward and reverse primer target regions (bins) for the

amplicons. For SARS-CoV-2, we selected 20 amplicons with an overlap of 300nt,

regularly spaced across the SARS-CoV-2 genome sequence (Figure 3B). We then

selected the top conserved primer sequences (the highest frequency primers)

mapping in the 5' or 3' 185nt of each amplicon. For security, the two highest frequency

primers per bin were selected for the SARS-CoV-2 sequence, this provided some

insurance against primer failure either due to target evolution or unexpected

secondary structure. The binning and primer target locations for the final set of primers

are shown in Figure 2 and the final calculated amplicon lengths were 1495-2093nt.

The reverse transcription, PCR amplification and library protocols were modified to

accommodate the new primers. Important changes to note are the following. Reverse

transcription was performed using the reverse primers and reverse transcription at 42°C. The

PCR cycling conditions (using Phusion enzyme) were adjusted for the new T_ms and an

increased elongation time required for the longer PCR products. Finally, the library purification

steps were adjusted to recover longer PCR and library products. A detailed step-by-step

protocol is provided in the supplementary material.

Testing the performance of primers to sequence SARS-CoV-2 using MinION

We tested the Entebbe primers performance for sequencing SARS-CoV-2 from

nucleic acid extracted from positive samples. The amplicon sizes and genome coverage are

summarised in Figure 4. In particular, panel A and B (Figure 4) illustrate the amplicon

products after the reverse transcription and PCR amplification with the expected sizes of

1400-2000 bp. These amplicons were then pooled and used for library preparation using the

MinION sequencing kits SQK-LSK109. Final libraries were quantified and sequenced using

a MinION Flow Cell (R9.4.1). The resulting read data, after quality and primer and adapter

trimming, were then mapped to the SARS-CoV-2 Wuhan1 reference genome NC_045512

(Figure 4, panels C and D) to document sequence coverage across the genome. The twenty

individual amplicons are detected in the coverage pattern with small peaks appearing where

amplicons overlap. The coverage is consistent across the genome with no missing amplicons

and the data were readily assembled into good coverage full genomes. Initial experiments

showed that amplicons 2 (spanning nt 2,400) and 16 (spanning nt 23,500) had reduced

yields (Figure 4, panel C). The primer mixes were subsequently adjusted to increase

concentrations of amplicon 1 and 16 primers for reverse transcription and PCR (see detailed

protocol in Supplementary materials), this improved yields relative to the other amplicons

(Figure 4 panel D).

Figure 4. Testing the primer performance. Panel A, PCR product size after pooling of

reaction A and B. Expected sizes of amplicons are from 1500 bp to 2093 bp before primer

trimming. Panel B, MinION reads after quality control, primer, adapter trimming. Panel C,

Reads mapped to SARS-CoV-2 reference genome, before amplicon 2 and 16 primer

boosting. Panel D, Reads mapped to SARS-CoV-2 reference genome, after amplicon 2 and

16 primer boosting.

Conclusions

Given the urgency of controlling the SARS-CoV-2 pandemic and the importance of

having good quality SARS-CoV-2 genomes, we are providing these alternative primers (the

Entebbe primers) with detailed step-by-step laboratory protocols to the community with the

hope that they benefit from the new design. The costs and efforts of sequencing SARS-CoV-

2 in the large numbers that are currently being seen is substantial and if these new primers

result in a higher proportion of gap-less genomes, this will provide added value and will

increase the utility of the resulting data.

Acknowledgements.

We thank all global SARS-CoV-2 sequencing groups for their open and rapid sharing

of sequence data and GISAID for providing an effective platform for making these data

available. We are grateful to the Oxford Nanopore Technologies and the ARTIC Network for

their support with protocols and analysis software.

We acknowledge the support of the Uganda Ministry of Health and its COVID-19

Scientific Advisory Committee, the National COVID-19 Task Force, and the staff of the

Emerging and Remerging Infections Department of the Uganda Virus Research Institute, and

the US Centers for Disease Control and Prevention. The SARS-CoV-2 diagnostic and

sequencing award is jointly funded by the UK MRC and the UK DFID under the MRC–DFID

Concordat agreement (grant agreement no. NC_PC_19060) and is also part of the European

and Developing Countries Clinical Trials Partnership 2 program supported by the European

Union. The diagnostics were also supported by the UK Research and Innovation/MRC, the

Global Fund, the Government of Uganda, the Islamic Development Bank, the World Health

Organization, GAVI, the US Centers for Disease Control and Prevention, and the Jack Ma

Foundation, among others. M.V.T.P. was supported by a Marie Sklodowska-Curie Individual

Fellowship, funded by European Union's Horizon 2020 research and innovation programme

(grant agreement no. 799417). The Uganda Medical Informatics Centre high performance

computer was supported by the UK MRC (grant no. MC_EX_MR/L016273/1) to P.K. The study

is supported by a Wellcome Epidemic Preparedness–Coronavirus grant, jointly funded by the

Wellcome Trust and UK DFID (grant agreement no. 220977/Z/20/Z) awarded to M.C.

References

1.   Edward C. Holmes, Yong-Zhen Zhang EC. Initial genome release of novel coronavirus
http://virological.org/t/initial-genome-release-of-novel-coronavirus/319. Virological.org
[Internet]. 2020; Available from: http://virological.org/t/initial-genome-release-of-novel-
coronavirus/319

2.   Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in
Wuhan, China, of Novel Coronavirus–Infected Pneumonia. N Engl J Med. 2020 Jan
29;NEJMoa2001316.

3.   Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically
ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered,
retrospective, observational study. Lancet Respir Med. 2020
Feb;S2213260020300795.

4.   Lauren Gardner et al. Coronavirus COVID-19 Global Cases by Johns Hopkins CSSE.
2020; Available from:
https://www.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b4
8e9ecf6

5.   Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data – from vision
to reality. Eurosurveillance. 2017 Mar 30;22(13):30494.

6.   Alessandrini F, Caucci S, Onofri V, Melchionda F, Tagliabracci A, Bagnarelli P, et al.
Evaluation of the Ion AmpliSeq SARS-CoV-2 Research Panel by Massive Parallel
Sequencing. Genes. 2020 Aug 12;11(8):929.

7.   Nicola De Maio, Conor Walker, Rui Borges, Lukas Weilguny, Greg Slodkowicz, Nick
Goldman. Issues with SARS-CoV-2 sequencing data. Virological.org [Internet]. 2020;
Available from: https://virological.org/t/issues-with-sars-cov-2-sequencing-data/473

8.   Page AJ, Mather AE, Le Viet T, Meader EJ, Alikhan N-FJ, Kay GL, et al. Large scale
sequencing of SARS-CoV-2 genomes from one region allows detailed epidemiology
and enables local outbreak management [Internet]. Epidemiology; 2020 Sep [cited
2020 Oct 9]. Available from:
http://medrxiv.org/lookup/doi/10.1101/2020.09.28.20201475

9.   Itokawa K, Sekizuka T, Hashino M, Tanaka R, Kuroda M. Disentangling primer
interactions improves SARS-CoV-2 genome sequencing by the ARTIC Network's
multiplex PCR [Internet]. Microbiology; 2020 Mar [cited 2020 Oct 9]. Available from:
http://biorxiv.org/lookup/doi/10.1101/2020.03.10.985150

10.   Josh Quick and Nick Loman. nCoV-2019 Version 3 Amplicon Releas. 2020; Available
from: https://community.artic.network/t/ncov-2019-version-3-amplicon-release/19

11.   Cotten M, Lam TT, Watson SJ, Palser AL, Petrova V, Grant P, et al. Full-Genome
Deep Sequencing and Phylogenetic Analysis of Novel Human Betacoronavirus. Emerg
Infect Dis [Internet]. 2013 May [cited 2020 Aug 29];19(5). Available from:
http://wwwnc.cdc.gov/eid/article/19/5/13-0057_article.htm

12.   Cotten M, Petrova V, Phan MVT, Rabaa MA, Watson SJ, Ong SH, et al. Deep
Sequencing of Norovirus Genomes Defines Evolutionary Patterns in an Urban Tropical
Setting. J Virol. 2014 Oct 1;88(19):11056–69.

13.   Agoti CN, Otieno JR, Munywoki PK, Mwihuri AG, Cane PA, Nokes DJ, et al. Local
Evolutionary Patterns of Human Respiratory Syncytial Virus Derived from Whole-
Genome Sequencing. Perlman S, editor. J Virol. 2015 Apr 1;89(7):3444–54.

14.   Phan MV, Murad SD, van der Eijk AA, Metselaar HJ, Hartog H, Harinck F, et al.
Genomic sequence of yellow fever virus from a Dutch traveller returning from the
Gambia-Senegal region, the Netherlands, November 2018. Eurosurveillance [Internet].
2019 Jan 24 [cited 2020 Aug 29];24(4). Available from:
https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2019.24.4.1800684

15.   Oxford Nanopore Technologies. Nanopore Protocol PCR tiling of COVID-19 virus.
2020; Available from: https://community.nanoporetech.com/protocols/pcr-tiling-
ncov/v/PTC_9096_v109_revH_06Feb2020

16.   Josh Quick. nCoV-2019 sequencing protocol. 2020; Available from:
https://www.protocols.io/view/ncov-2019-sequencing-protocol-bbmuik6w

17.   Hsin-Ru Chang, Jay Crespo, Kathy Lee, Kathleen McGall, Susan MacWhorter, Paul
Russell, Jeff Schatz, Donna Seid, Criss Walworth, Joe Yu. TaqMan® Assays Shipped
at Ambient Temperature Reduce Environmental Impact and Retain Their Quality and
Stability. Available from: https://assets.thermofisher.com/TFS-
Assets/LSG/brochures/cms_081489.pdfhttps://assets.thermofisher.com/TFS-
Assets/LSG/brochures/cms_081489.pdf

Supplementary Materials: 1. Detailed protocol. 2. Primer table.

We acknowledge the Oxford Nanopore Technologies and the ARTIC Network for their

extensive3 protocols which provided important background details for the method presented

here (15)(16). Potential users are strongly advised to study the original protocos carefully.

The protocol presented here assumes the users has all the ARTIC and ONT background

experience.

Detailed protocol for reverse transcription, PCR and sequencing library preparation.

Practical comments. Primers were ordered and delivered dry and were dissolved at

100 μM (= 100 pmol/μl) in PCR quality water. Dry shipped primers are reported to have better

stability at room temperature (17). Ideal the primers were allowed to dissolve in PCR grade

water for at least 3 hours at room temperature or overnight at 4 °C before diluting for use (see

below).

Prepare the following primer mixes:

RPM_A: 21 primers. 3 μl of 100 μM stock of the following Reverse primers plus PCR

grade water to 200 μl (= 300 pmol x 21 = 6300 pmol/200 μl = 32 pmol/μl. 2 μl per 20 μl RT

reaction = 3.2 pmol/μl = 3.2 μM concentration in reaction) .

Primers in RPM_A: nCoV-2019_1_LEFT, AR1_3, AR1_4, AR3_11, AR3_12,

AR5_19, AR5_20, AR7_27, AR7_28, AR9_35, AR9_36, AR11_43, AR11_44, AR13_51,

AR13_52, AR15_59, AR15_60, AR17_67, AR17_68, AR19_75, AR19_76.

RPM_B: 21 primers. 3 μl of 100 μM stock of the following Reverse primers plus PCR

grade water to 200 μl. (= 300 pmol x 21 = 6300 pmol/200 μl = 32 pmol/μl. 2 μl per 20 μl RT

reaction = 3.2 pmol/μl = 3.2 μM concentration in reaction). **Primer Boost:** For Primers BR2_7,

BR2_8, BR16_63, BR16_64 use 9 μl of 100 μM stock.

Primers in RPM_B: BR2_7, BR2_8, BR4_15, BR4_16, BR6_23, BR6_24, BR8_31,

BR8_32, BR10_39, BR10_40, BR12_47, BR12_48, BR14_55, BR14_56, BR16_63,

BR16_64, BR18_71, BR18_72, BR20_79, BR20_80, nCoV-2019_98_RIGHT.

PCR primer mix A (PPM_A): 1.5 μl of each primer (forward and reverse primer) plus

439 μl PCR grade water. = 41 primers (100 pmol/μl) = 4100 pmol in 500 μl water = 8.2 pmol/μl

in the PPM. When 2 μl PPM used per 25 μl PCR reaction = 0.66 pmol/μl in reaction (= 660

nM).

Primers in PPM_A: nCoV-2019_1_LEFT, AF1_1, AF1_2, AR1_3, AR1_4, AF3_9,

AF3_10, AR3_11, AR3_12, AF5_17, AF5_18, AR5_19, AR5_20, AF7_25, AF7_26, AR7_27,

AR7_28, AF9_33, AF9_34, AR9_35, AR9_36, AF11_41, AF11_42, AR11_43, AR11_44,

AF13_49, AF13_50, AR13_51, AR13_52, AF15_57, AF15_58, AR15_59, AR15_60,

AF17_65, AF17_66, AR17_67, AR17_68, AF19_73, AF19_74, AR19_75, AR19_76 .

PCR primer mix B (PPM_B): 1.5 μl of each primer (forward and reverse primer) plus

439 μl PCR grade water = 41 primers (100 pmol/μl) = 4100 pmol in 500 μl water = 8.2 pmol/μl

in the PPM. When 2 μl PPM used per 25 μl PCR reaction = 0.66 pmol/μl in reaction (= 660

nM). **Primer Boost:** for primers BF2_5, BF2_6, BR2_7, BR2_8, BF16_62, BF16_61,

BR16_63, BR16_64, use 4.5 μl of 100 μM stock

Primers in PPM_B: BF2_5, BF2_6, BR2_7, BR2_8, BF4_13, BF4_14, BR4_15,

BR4_16, BF6_21, BF6_22, BR6_23, BR6_24, BF8_29, BF8_30, BR8_31, BR8_32, BF10_37,

BF10_38, BR10_39, BR10_40, BF12_45, BF12_46, BR12_47, BR12_48, BF14_53,

BF14_54, BR14_55, BR14_56, BF16_61, BF16_62, BR16_63, BR16_64, BF18_69,

BF18_70, BR18_71, BR18_72, BF20_77, BF20_78, BR20_79, BR20_80, nCoV-

2019_98_RIGHT.

1. Set up Reverse Transcription reactions (two reactions per sample).

Mix the following components in a 0.2mL 8-strip tube, 1 RT_A and 1 RT-B reaction per

sample.

Component   Volume (per sample)

RT_A_Reaction

Component   **Volume**

RPM_A   1 μl

Template RNA   11μl (if using less than 11 μl viral RNA adjust to 11μl

358   with water)

359 Total Volume   12μl

361 **RT_B_Reaction**

362 **Component**   **Volume**

363 RPM_B   1 μl

364	Template RNA	11μl (if using less than 11 μl viral RNA adjust to 11μl
365		with water)
366	Total Volume	12μl

Heat at 65 °C for 5:00 mins, and quickly chill in ice/water bath for at least 1:00 min.

Viral RNA input from a clinical sample should be between Ct 18-35. If Ct is between 12-15,

then dilute the sample 100- fold in water, if between 15-18 then dilute 10-fold in water. This

will reduce the likelihood of PCR-inhibition.

Reverse Transcription

Add the following to the 12 μl annealed template RNA sample:

376	**Component**	**Volume (1 reaction)**
377	SuperScript IV 5X Buffer	4 μl
378	100mM DTT	1 μl
379	RNaseOUT RNase Inhibitor	1 μl
380	10mM dNTPs mix (10mM each)	1μl
381	SuperScript IV	1 μl
382	(add 8 μl mix per reaction)	
383	Total Volume	20 μl

2. Incubate the RT reaction as follows:

42 °C   50 minutes

70 °C   10 minutes

Hold at 4 °C

3. Set up Amplicon PCR

In the Mastermix hood set up the multiplex PCR reactions as follows in 0.2mL 8-strip PCR

tubes. Prepare one PPM_A and one PPM_B mix per sample.

393	**Component**	**1 reaction**	**8rxns_ PPM_A_Mix**	**8rxns_ PPM_B_Mix**
394	5 x Phusion HF buffer	5μl	40 μl	40 μl
395	10 mM dNTPs	0.5μl	4 μl	4 μl
396	Phusion DNA Pol	0.5μl	4 μl	4 μl
397	Primer PPM_A or PPM_B			
398	(10μM)	2 μl	16 μl	16 μl

399	Nuclease-free water	12 μl	96 μl	96 μl
400	Total	20 μl per rxn		

402 Aliquot 20 μl per tube PPM_A_Mix or PPM_B_Mix

404 In the extraction and sample addition cabinet add **5µl cDNA** to each tube and mix well by

405 pipetting.

407 **4. Run PCR**

	Step	Cycles	Temp	Time
408	**Step**	**Cycles**	**Temp**	**Time**
409	Heat Activation	1	98 °C	30sec
410	Denaturation	35	98 °C	15sec
411	Annealing	35	63 °C	30sec
412	Extension	35	72 °C	3 min
413	Polish	1	72 °C	10 min
414	Hold		4 °C	infinite

In post-PCR cabinet: After PCR, for each sample, pool the two reactions in a 1.5 ml

microcentrifuge tube.

5. Clean up using AMPure XP beads, Agencourt.

Add an equal volume (1:1) of AMPure XP beads to the sample tube and mix gently by either

flicking or pipetting. E.g. add 50 µl AMPure XP bead suspension to a 50 µl reaction.

Pulse centrifuge to collect all liquid at the bottom of the tube.

Incubate for 5 min at room temperature.

Place on magnetic rack and for 2 min

Carefully remove and discard the supernatant, being careful not to touch the bead pellet.

Ethanol wash 1: Add 200µl of room-temperature 70%volume ethanol to the pellet.

Carefully remove and discard ethanol, being careful not to touch the bead pellet.

Repeat Ethanol wash

Pulse centrifuge to collect all liquid at the bottom of the tube and carefully remove as much

residual ethanol as possible using a P10 pipette.

Open lid, allow to dry briefly (2 min)

Resuspend pellet in 30µl Elution Buffer (EB, QIAGEN)

Mix gently by flicking,

Incubate 5 min

Place on magnet and transfer sample (supernatant) to a clean 1.5mL Eppendorf tube

ensuring no beads are transferred into this tube.

6. Quantitate (A Tapestation 4200 analyzer with the D5000 assay kit as well as the Qubit

fluorometer were used.)

7. Add Barcodes (Steps 7 onward derived from the Oxford Nanopore Technologies

protocol (15)).

Set up the following reaction for each sample:

Component	Volume
DNA amplicons	5 μl*
Nuclease-free water	7.5 μl
Ultra II End Prep Reaction Buffer	1.75 μl
Ultra II End Prep Enzyme Mix	0.75 μl
Total Volume	15 μl

*Volumes usually varies based on the DNA concentration. For the 1500bp amplicons and
pool size of appx 8 we have input 50ng and for a pool size of 12 we have input 20ng
460

Incubate at room temp (20°C) for 20 min

Incubate at 65 °C for 5:00 min

Incubate on ice for 1 min

E7645S/L NEBNext Quick Ligation Module

Component	Volume
End repaired sample	15μl
Barcode vol	2.5 μl
Ultra II Ligation Master Mix	17.5 μl
Ligation Enhancer	0.5 μl
Water	1.5 μl
final volume	35.5 μl

Incubate in a thermocycler for 15 minutes at 20°C with heated lid set to 30°C (or off).

Incubate at 70 °C for 10 min

Incubate on ice for 1 min

The 70°C incubation is to inactivate the DNA ligase to prevent barcode cross-ligation when

reactions are pooled in the next step.

Pool all barcoded fragments together into a new 1.5 ml Eppendorf tube.

8. Clean up on AMPure XP beads, Agencourt.

Add an equal volume (1:1) of SPRI beads to the sample tube and mix gently by either flicking

or pipetting. E.g. add 50 µl SPRI beads to a 50 µl reaction.

Pulse centrifuge to collect all liquid at the bottom of the tube.

Incubate for 5 min at room temperature.

Place on magnetic rack and for 5 min

Carefully remove and discard the supernatant, being careful not to touch the bead pellet.

Ethanol wash 1: Add 200µl of room-temperature 70%volume ethanol to the pellet.

Carefully remove and discard ethanol, being careful not to touch the bead pellet.

Repeat Ethanol wash

Pulse centrifuge to collect all liquid at the bottom of the tube and carefully remove as much

residual ethanol as possible using a P10 pipette.

Open lid, allow to dry briefly (2 min)

Resuspend pellet in 30µl Elution Buffer (EB, QIAGEN)

Mix gently by flicking

Incubate 5 min

Place on magnet and transfer sample (supernatant) to a clean 1.5mL Eppendorf tube

ensuring no beads are transferred into this tube.

9. Quantitate as above.

10. Add AMII adapters.

AMII adapters ligation reaction:

Component	**Volume**
Barcoded amplicon pools	30 μl
NEBNext (E60562)	
Quick Ligation Reaction Buffer (5X)	10 μl
AMII adapter mix	5 μl
Quick T4 DNA Ligase	5 μl
Total	50 μl

The input of barcoded amplicon pools will depend on the number of barcoded pools and

should be between 40 ng (8 barcodes) and 160 ng (24 barcodes).

Incubate at room temperature for 15:00 min

11. Clean up on AMPure XP beads, Agencourt.

USE SFB instead of Ethanol for two washes ***CRUCIAL, Ethanol strips off motor protein

from adapter. You don't want to do this!

Clean up on AMPure XP beads, Agencourt

Add an equal volume (1:1) of SPRI beads to the sample tube and mix gently by either flicking

or pipetting. E.g. add 50 µl SPRI beads to a 50 µl reaction.

Pulse centrifuge to collect all liquid at the bottom of the tube.

Incubate for 5 min at room temperature.

Place on magnetic rack and for 2 min

Carefully remove and discard the supernatant, being careful not to touch the bead pellet.

SFB washes: Add 200µl of room-temperature **SFB** to the pellet.

resuspend

collect on magnet

Carefully remove and discard **SFB** being careful not to touch the bead pellet.

Repeat SFB wash

Pulse centrifuge to collect all liquid at the bottom of the tube and carefully remove as much

residual **SFB** as possible using a P10 pipette.

Resuspend pellet in 15 μl Elution Buffer (EB, QIAGEN)

Mix gently by flicking,

Incubate 5 min

Place on magnet and transfer sample (supernatant) to a clean 1.5mL Eppendorf tube

ensuring no beads are transferred into this tube.

12. Quantitate.

13. Proceed to MinION sequencing.

 Supplementary material Table 1. Entebbe Primers

Amplicon_no	Primer_id	Sequence[1]	Position[2]
1	AF1_1	GTCCGGGTGTGACCGAAAG	242
1	AF1_2	CCTTGTCCCTGGTTTCAACGA	274
1	AR1_3	TGCGGGAGAAAATTGATCGTACA	1894
1	AR1_4	GCACAGAATTTTGAGCAGTTTCA	1921
2	BF2_5	ACGTGCTAGCGCTAACATAGG	1540
2	BF2_6	GTTGGAGAAGGTTCCGAAGGT	1580
2	BR2_7	TAGCCTTATTTAAGGCTCCTGCAA	3477
2	BR2_8	TGCAACACCTCCTCCATGTTTAA	3459
3	AF3_9	ACAAACTGTTGGTCAACAAGACG	3220
3	AF3_10	CAACAAGACGGCAGTGAGGA	3233
3	AR3_11	TTGTGTAGATTGTCCAGAATAGGACC	4806
3	AR3_12	TCCATATGTCATTGACATGTCCACA	5014
4	BF4_13	TTTGGAAGAAGCTGCTCGGTATAT	4639
4	BF4_14	GAAACCATCTCACTTGCTGGTTC	4772
4	BR4_15	GGTTTTAGATCTTCGCAGGCAAG	6380
4	BR4_16	CCATTAGATCTGTGTGGCCAA	6534
5	AF5_17	CACACCCTCTTTTAAGAAAGGAGCTA	6190
5	AF5_18	TTGTTTGGCATGTTAACAATGCAAC	6234
5	AR5_19	CAGACGCTGATTTTGCAGATGA	7919
5	AR5_20	GCACTATCACCAACATCAGACAC	7994
6	BF6_21	AGGCTTTTGCAAACTACACAATTG	7591
6	BF6_22	TATGCTAATGGAGGTAAAGGCTTTTG	7574
6	BR6_23	CGATAGCTACAATACCACCAGCT	9409
6	BR6_24	CAATACCACCAGCTACTATAGATGCT	9397
7	AF7_25	AAGCTGGTGTTTGTGTATCTACTAGT	9243
7	AF7_26	ACCTACCTTGAAGGTTCTGTTAGAG	9164
7	AR7_27	CACTACCCAATATGGTACGTCCA	10885
7	AR7_28	TGAGTAACAACCAGTGGTGTGT	11000
8	BF8_29	CTGGAGTTCATGCTGGCACA	10560
8	BF8_30	GTTTTAGCTTGGTTGTACGCTG	10664
8	BR8_31	TTTGCCCTCTTGTCCTCAGATCTA	12313
8	BR8_32	TGGTATGACAACCATTAGTTTGGCT	12466
9	AF9_33	CTCAAGAAGCTTATGAGCAGGCT	12144
9	AF9_34	CAAGCTATAGCCTCAGAGTTTAGTTC	12089
9	AR9_35	ACGTTGACGTGATATATGTGGTACC	13770
9	AR9_36	ACGAGGTCTGCCATTGTGTATT	13802
10	BF10_37	GCGGTGTAAGTGCAGCC	13475
10	BF10_38	TCGCTTCCAAGAAAAGGACGAA	13602
10	BR10_39	CTCAATACTTGAGCACACTCATTAGC	15406
10	BR10_40	GTGACAAGCTACAACACGTTGT	15367
11	AF11_41	ATTCTATGGTGGTTGGCACAACA	15219
11	AF11_42	CAATAGCCGCCACTAGAGGAG	15173
11	AR11_43	TAGTGTAGGTGCACTTAATGGCATT	16932
11	AR11_44	GGTAAACAACAGCATCACCATAGTC	16846
12	BF12_45	AACATGTGACTGGACAAATGCTG	16566
12	BF12_46	ACTGACTTTAATGCAATTGCAACATG	16546
12	BR12_47	TACAGCAACTAGGTTAACACCTGTAG	18374
12	BR12_48	CACCCCTCGACATCGAAGC	18302

13	AF13_49	GTGGCAACTTTACAAGCTGAAAATG	18025
13	AF13_50	GACATACCTGGCATACCTAAGGAC	18160
13	AR13_51	GTTTAATGTTGCGCTTAGCCCAA	19791
13	AR13_52	GTAGTCCCAGATCACAGTATTAGCAG	19859
14	BF14_53	CGTGTATAACACGTTGCAATTTAGGT	19454
14	BF14_54	ACTTTGATGGACAACAGGGTGAA	19661
14	BR14_55	CGCGTGGTTTGCCAAGATAATTA	21285
14	BR14_56	CATAACCATCTATTTGTTCGCGTGG	21301
15	AF15_57	AGCTCATGGGACACTTCGC	21203
15	AF15_58	TTGGAGGTTCCGTGGCTATAAAG	21146
15	AR15_59	CGATTTGTCTGACTTCATCACCTC	22770
15	AR15_60	CTACCGGCCTGATAGATTTCAG	22971
16	BF16_61	TGCATCTGTTTATGCTTGGAACAG	22603
16	BF16_62	CTCTCTCAGAAACAAAGTGTACGTTG	22446
16	BR16_63	CACTTGCTGTGGAAGAAAGTGA	24371
16	BR16_64	GTTGACCACATCTTGAAGTTTTCCAA	24396
17	AF17_65	GGTGATTGCCTTGGTGATATTGC	24074
17	AF17_66	CTGTTTTGCCACCTTTGCTCA	24138
17	AR17_67	CCAGCAAAGAAAATAGTTGGCATC	25816
17	AR17_68	CGTAACAATTAGTATGCCAGCAAAGA	25830
18	BF18_69	TCCCTTTCGGATGGCTTATTGTT	25514
18	BF18_70	TGAAATCAAGGATGCTACTCCTTCAG	25446
18	BR18_71	GTTGTACCTCTAACACACTCTTGGTA	27451
18	BR18_72	CTCACAAGTAGCGAGTGTTATCAG	27418
19	AF19_73	ACGCTTTCTTATTACAAATTGGGAGC	27045
19	AF19_74	CGCTGTGACATCAAGGACCT	26994
19	AR19_75	TGGCAATGTTGTTCCTTGAGGAA	28755
19	AR19_76	CAGCCATTCTAGCAGGAGAAGTT	28885
20	BF20_77	TAGAGTATCATGACGTTCGTGTTGTT	28219
20	BF20_78	ACCCCGCATTACGTTTGGT	28309
20	BR20_79	TGGCTCTTTCAAGTCCTCCCT	29700
20	BR20_80	GCTCTTCCATATAGGCAGCTCTC	29779
	nCoV-2019_1_LEFT[3]	ACCAACCAACTTTCGATCTCTTGT	31
	nCoV-2019_98_RIGHT[3]	TTCTCCTAAGAAGCTATTAAAATCACATGG	29866

569 **Footnotes** 1. Sequence listed 5' to 3'.

570 2. Position in SARS-CoV-2 reference genome GenBank NC_045512

571 3. From original ARTIC primer set V.1 (16).