Did the Himalayan Uplift Drive Evolution of Aquatic Biodiversity Across Asia? Snowtrout (Cyprinidae: *Schizothorax*) as a Test Case

Binod Regmi[1,4], Marlis R. Douglas[1], David R. Edds[2], Karma Wangchuk[1,3], Chang Lu[3], Gopal Prasad Khanal[3], Pema Norbu[3], Sangay Norbu[3], Sonam Dorji[3], Singye Tshering[3], Zachary Angel[1], Tyler K. Chafin[1], Zachery D. Zbinden[1], Michael E. Douglas[1]

1. Department of Biological Sciences, University of Arkansas, Fayetteville Arkansas USA 72701

2. Department of Biological Sciences, Emporia State University, Emporia, Kansas USA 66801

3. National Research & Development Centre For Riverine and Lake Fisheries, Ministry of Agriculture & Forests, Royal Government of Bhutan

4. Present Address: National Institute for Arthritis, Musculoskeletal & Skin Diseases (NIAMS), National Institute of Health, Bethesda MD USA 20892

Corresponding author:

Michael Douglas (email: med1@uark.edu)

The Himalayan uplift, a tectonic event of global importance, seemingly disseminated aquatic

biodiversity broadly across Asia. But surprisingly, this hypothesis has yet to be tested. We do so

herein by sequencing 1,140 base-pair of mtDNA *cytochrome-b* for 72 tetraploid

Nepalese/Bhutanese Snowtrout (*Schizothorax spp.*), combining those data with 67 GenBank®

sequences (59 ingroup/8 outgroup), then reconstructing phylogenetic relationships using

maximum likelihood/ Bayesian analyses. Results indicate Snowtrout originated in Central Asia,

dispersed across the Qinghai-Tibetan Plateau (QTP), then into Bhutan via south-flowing

tributaries of the east-flowing Yarlung-Tsangpo River (YLTR). The headwaters of five large Asian

rivers provided dispersal corridors into southeast Asia. South of the Himalaya, the YLTR

transitions into a westward-flowing Brahmaputra River that facilitated successive colonization

of Himalayan drainages: First Bhutan, then Nepal, followed by far-western drainages

subsequently captured by the Indus River. We found greater species-divergences across rather

than within-basins, implicating vicariant evolution as a driver. The Himalaya is a component of

the "third-pole" [Earth's largest (but rapidly shrinking) glacial reservoir outside the

Arctic/Antarctic]. Its unique aquatic biodiversity must not only be recognized (as herein) but

also conserved through broad, trans-national collaborations. Our results effectively contrast

phylogeography with taxonomy as a necessary first step in this process.

18 The Himalaya is the most extensive and recently evolved mountain system on Earth

(length=2400km; width=240km; elevation=75-8800m), with a global significance underscored

20 by its large-scale lithospheric, cryospheric, and atmospheric interactions [1]. These have not

only driven global climate, but also defined the cultural and biological endemism of the region

22 [2]. Massive, tectonically derived mountain chains such as the Alps and the Himalaya are

hypothesized as being fundamental to the formation of global biodiversity gradients via

24 vicariance and local adaptation, with a significantly stronger signal in terrestrial rather than

aquatic systems [3]. Here we test how orogeny (the deformation and folding of Earth's crust by

26 lateral compression) has contributed to the diversification of freshwater fishes broadly across

Asia. We do so by evaluating the phylogeography of an endemic high-elevation fish, the

28 Snowtrout (*Schizothorax*: Cyprinidae).

30 **Himalayan orogeny**

 The Himalaya formed as a series of parallel ranges bordered by rivers on the west (Indus) and

32 east (Brahmaputra), and encompassing most of northern Pakistan, northern India (Kashmir to

the west, Assam to the east), as well as Nepal, Bhutan, and parts of China, with the extensive

34 Qinghai-Tibetan Plateau (QTP) to the north and the alluvial plains of India and Bangladesh to

the south (Fig. 1). It evolved over a 30my span, the result of a two-stage impact of the Indian

36 craton with the Eurasian landmass. The leading edge of the Indian craton was subducted, and

its upper surface essentially 'skimmed' so as to form three sequential north-to-south

38 stratigraphic zones (i.e., Tibetan, Greater Himalaya, and Lesser Himalaya [4]).

The uplift not only established the Himalaya as a topographic entity, but also profoundly

40   influenced the connectivity of its regional rivers [5,6,7]. For example, in late Oligocene–early

Miocene (26-19my), the Yarlung-Tsangpo River (YLTR; average elevation 4000m) formed at the

42   southern border of the QTP as a series of lakes with south-flowing tributaries (Fig. 11 in [8]).

The lakes eventually coalesced into a westward flowing river (early–mid Miocene; 19–15ma)

44   that eventually reversed flow in response to a continuing but disproportionate uplift. The YLTR

was captured on the eastern QTP by the Brahmaputra River in early-mid Miocene (i.e., ~18–

46   15ma) [7,9], a signature event that seemingly allowed aquatic biodiversity to disperse broadly

from east to west in the Himalaya, and to access multiple south-flowing Himalayan drainages.

Orogeny and biodiversity

50   The contemporary species richness of terrestrial biodiversity across the QTP clearly reflects the

three stratigraphic zones resulting from the uplift. Although this endemism is hypothesized as

52   being driven by tectonic activity, requisite supporting data are infrequent or lacking [10]. The

divergence of regional freshwater fishes seemingly occurred when the QTP elevated more than

54   3,000m in late-Miocene [11], with a potential result being the increased isolation of tributaries.

The climatic regime of the QTP was also greatly altered, resulting in strong selection for high-

56   altitude specialists that could seemingly subsist under harsher conditions [12].

Yet, impacts on biodiversity were surprisingly variable. For example, aquatic

58   invertebrates seemingly lose species diversity as elevation increases [13,14], whereas cold-

adapted terrestrial species are seemingly buffered by microclimates at higher altitude [15].

60   However, data for high elevation fishes remain elusive, particularly so in developing countries,

an aspect that looms large in the context of monitoring programs that are incomplete and/or

62   non-systematic [16]. This promotes an opportunity for the recognition of cryptic species not as

yet identified.

64   Here, we examined a broadly distributed Asiatic fish (Snowtrout, genus *Schizothorax*), a

component of the largest and most diverse of freshwater fish families (Cyprinidae: 1700+ valid

66   species, with 10 recognized subfamilies [17]). Previous research categorized six major

biogeographic entities within *Schizothorax* [6]: A Central Asiatic clade, followed in order by five

68   drainage-specific sister-clades – YLTR/ Irrawaddy; Indus; Irrawaddy; Mekong/Salween; Yangtze.

However, their phylogenetic interpretation provided an incomplete template for biogeography

70   and evolution of Snowtrout across the entire Himalaya and surrounding regions.

We broadened and extended these results by focusing in particular on two key

72   Himalayan regions previously opaque with regard to freshwater fish diversity: Bhutan (to the

Himalayan east) and Nepal (central). Both, in tandem, effectively link the far eastern

74   component of the Himalaya with its western terminus, allowing a more appropriate test of

tectonism, orogeny and the evolution of aquatic biodiversity. More specifically, it provides a

76   precise estimate for diversification into eastern, southern, and western Asia, and sets the stage

for subsequent climate oscillations in the Quaternary that served to isolate and modify regional

78   biodiversity [18].

80   **Contemporary significance**

The QTP (and surrounding mountains) are termed "the third pole" in that they contain the

82   greatest amount of global snow and ice beyond the Arctic/Antarctic. It is also the headwater

source for nine large Asian rivers that collectively provide ecosystem services for >1.5 billion

84   people [19]. Glaciers in the eastern Himalaya, for example, account for an estimated 14.5% of

the worldwide total, of which a quarter has been lost since 1970 [20], due largely to a global

86   warming 2x greater than average [2]. Cumulative impacts are recorded as increasing

precipitation, decreasing glacial surface area, diminishing snow cover, expanding glacial lakes,

88   and loss of permafrost [21].

The onset of these conditions has (and will continue to) significantly impact the

90   distribution of *Schizothorax*, forcing a migration into higher-elevation streams, as well as

concomitant contraction of trailing-edge habitats [22]. These are important considerations

92   given the limitations inherent with *Schizothorax* taxonomy, especially given the strong potential

for as-of-yet undiscovered biotic components (e.g. cryptic intraspecific divisions and

94   undescribed species). Regional collaborations that extend across national borders are needed

to mitigate climate-change impacts, and to foster cooperative cross-border conservation

96   strategies [21]. Such a regional mandate requires a comprehensive understanding of extant

biodiversity, to include inter-specific delineations and the nature by which these juxtapose with

98   both geographical and political boundaries. To do so requires: A) the solidification (and

unification) of taxonomies (i.e. to catalogue units for conservation); and B) an established

100 theory for the latter with which to define biogeographic/ phylogeographic expectations. A

framework that contrasts these components within a taxonomically and biologically complex

102 genus can be found herein.

Results

104 **Sequence data**

Our phylogenetic analyses incorporated 139 cytochrome-b (*cyt-b*) haplotypes spanning 1140

106   bp, with 643 of those monomorphic (=56%), 432 parsimony-informative (38%), and 65 (6%) as

singleton polymorphisms. Based on the Bayesian Information Criterion (BIC), the best fitting

108   model of sequence evolution was TN+F+I+G4 (i.e., unequal evolutionary rates among:

Purines/pyrimidines, empirical base frequencies, proportion of invariant sites, and a 4-rate

110   category discrete Gamma model). Nepalese and Bhutanese haplotypes (N=53 and N=19,

respectively) were compiled across three large, separate drainages within each country.

Phylogeography of *Schizothorax* across Asia and the Himalaya

114   To estimate independent groups in our data, we applied a Bayesian approach without *a priori*

assumptions (GENELAND V.4.0.3: [23]) and detected evidence for 13 distinct groups. An additional

116   group emerged as a 'ghost' lineage but was subsequently disregarded as an artifact of isolation-

by-distance [24]. We subsequently explored the manner by which these groups (i.e., N=13

118   geographic regions; Fig. 2) encapsulated sequence variation by employing an analysis of

molecular variance framework (AMOVA; [25]). All were found to be significantly different one

120   from another, with partitions among groups accounting for 78% of the total observed genetic

variance (Table 1). Aggregates (= clades) within groups also differed significantly, as did

122   haplotypes within aggregates (Fig. 3). Probabilities for all comparisons were based on 1,000

permutations, with significance gauged by Bonferroni-correction (P<0.005 for 13 groups,

124   p<0.0017 for 29 "aggregates").

We then visualized our groups by deriving maximum likelihood (ML) and Bayesian (BA)

126   trees, each of which resulted in largely consistent topographies (ML tree Figs. 2, 3; Bayesian

tree Fig. S1). However, all of our statistical topology tests significantly excluded the BA topology

128   ($p<0.05$), and given this, we thus focus primarily on the ML topology, other than highlighting in

a comparative sense the notable differences between the two (Table 2).

130   The majority of nodes in the ML trees [e.g., composite (Fig. 2) and extended (Fig. 3)]

were strongly supported on the basis of SH-ALRT (Shinodara-Hasegawa approximate likelihood

132   ratio test) values and ultrafast bootstrap approximations (UFBOOT2). Nodes were likewise

supported in the BA tree, with the majority displaying a posterior probability ranging from 0.9-

134   1.0 (Fig. S1).

Two surprising results emerged in both analyses: Bhutanese Snowtrout were

136   consistently non-monophyletic, nesting instead within two temporally and spatially disparate

parental clades (i.e., within the Central QTP and the Koshi drainage of Nepal). This indicates

138   multiple independent colonizations of high-altitude habitat within Bhutanese drainages. Nepali

Snowtrout, on the other hand, showed strong phylogenetic structuring east-to-west,

140   sequentially diverging in a manner consistent with the broad-scale topographies of riverine

networks, and with sub-clades being taxonomically congruent by-and-large. Below, we evaluate

142   results from the ML analysis, but underscore in the discussion those minor discrepancies

between ML and BA trees. The specifics of group membership are provided in Table S1.

144   Groups recovered using genetic clustering approaches (N=13) corresponded well with

geographic regions, concordant with groupings apparent in the ML analysis (Fig. 2). These are:

146   (1) Central Asia; (2) Central QTP; (3) YLTR West; (4) Bhutan-1A; (5) YLTR East; (6) Bhutan-1B; (7)

Eastern QTP/ Southeast Asia-Clade A (Salween-Irrawaddy drainages); (8) Eastern QTP/

148   Southeast Asia-Clade B (upper Yangtze drainage); (9) Eastern QTP/ Southeast Asia-Clade C

(Mekong-Salween drainages); (10) Bhutan-2 (via Brahmaputra River); (11) Koshi drainage

150   (eastern Nepal); (12) Gandaki drainage (central Nepal); and (13) a composite western Himalaya

that included Laka Rara and the Karnali drainage (western Nepal), the upper Ganges (northwest

152   India), and the Indus River Basin (Pakistan). The latter group (i.e., western Himalaya) was

additionally partitioned in the BA tree (Fig. S1). For discussion purposes, we subsumed a

154   significantly different single-species clade (*S. argentatus*) within the Central Asia clade (Table

S1). A more detailed phylogeny is presented in Fig. 3, with regional composites in Table S1.

156   Differences between ML and BA-trees are found in the first large geographic cluster that

is sister to Central Asia, athough the BA-tree was ubiquitously excluded across all topology tests

158   considered (Table 2). The clade containing both YLTR regions and Bhutan-1 in the ML-tree was

additionally subdivided in the BA-tree: YLTR-West was partitioned by a smaller Bhutan-1A

160   clade, whereas YLTR-East and a larger Bhutan-1B clade were collapsed.

The second large geographic cluster sister to Central Asia in the ML-tree (Fig. 2) has the

162   Eastern QTP/ Central Asia region as a basal node. It has six Himalayan regions as sister groups.

The first of these represents eight undescribed Bhutanese samples (=Bhutan-2). The second

164   depicts the Koshi River (eastern Nepal), and both are sister to the Gandaki River group (central

Nepal). Of interest is the fact that *S. richardsonii* and *S. progastus* fall together as components

166   of distinct clades, rather than clustering separately by taxonomic designation. A similar result

occurs within the Karnali, Ganges, and Indus river drainages. In the BA-tree, the Indus and

168   Ganges drainages appear as distinct clades, whereas they are less so within the ML-tree.

170   **Temporal divergences**

We employed RELTIME to derive major time-calibrated events, 13 of which are depicted in our

ML-based tree (Fig. 2; other dates in discussion). The average time for the most basally

divergent node (i.e., Central Asia) was 11.6 Ma (CI=15.4-8.7). The two large subclades sister to

Central Asia then separated at approximately 7.3 Ma (CI=11.3-4.7). In the first of these, the

Central QTP diverged at 6.2 Ma (10.2-3.7), followed by the YLTR West (4.2 Ma; CI=7.4-2.4), then

Bhutan-1A (3.3 Ma;CI=6.7-1.5), with YLTR East and Bhutan-1B at 2.7 MA (CI=5.9-1.2).

In the second large subclade sister to Central Asia, the Eastern QTP/ Southeast Asia

region diverged at 6.6 Ma (CI=10.7-4.1), followed by Bhutan-2/ Koshi River and Gandaki River/

western Himalayan clades at 4.8 Ma (CI=8.8-2.6). Our Bhutan-2 clade then separated from the

Koshi River at 0.4 Ma (CI=0.8-0.2) whereas the Gandaki separated from a composite western

Himalayan clade at 1.9 Ma (CI=3.9-0.9) (parsed more thoroughly in Discussion).

Discussion

The synergy between tectonism and climate

Orogeny and tectonism drive the evolution of aquatic ecosystems and act synergistically with

climate to impact both hydrology [26] and resident freshwater fishes [27]. For example,

tectonically derived drainage systems in western North America were gradually shaped over

evolutionary time by historic, long-term drought. The result was an erosion of genetic diversity,

with endemic fish populations repeatedly collapsing into refugia then expanding outward

during onset of more pluvial periods [28]. Episodic flooding, on the other hand, allowed

antecedent streams to down-cut, with headwaters subsequently being eroded and basins

concomitantly expanding, with large-scale dispersals being promoted [29]. These alterations

(i.e., captures, diversions, beheadings; [30]) not only extended the distribution of fishes into

194 adjacent basins [31], but also facilitated hybridization [32,33,34], a process that has confound

both taxonomists and managers (and continues to do so).

196   Flows within basins were often abridged by longitudinal barriers that are vicariant for

larger-bodied species yet serve as passive filters for those smaller and more limited in their

198 dispersal [35]. As a result, fish distributions are frequently characterized as a series of

biogeographic "islands" within and among basins, each reflecting local expansions,

200 contractions, and extinctions. As such, they provide necessary templates with which to gauge

responses by fish to similar tectonic, orogenic, and climatic impositions that have occurred

202 within analogous but less well monitored dendritic ecosystems (as herein).

The Himalaya displays sharp elevational gradients within which there are ample

204 opportunities for diversification. These, in turn, have promoted diversification, that in turn

established the region as a global hotspot of biodiversity and endemism [16,36]. Yet, broad

206 phylogeographic studies pertaining to freshwater biodiversity in the Himalaya have been

limited, with foci instead on more restricted regions, rivers, and/or mountains [37,38,39,40].

208 Our perspective was somewhat broader (per [6]), in that we sought molecular evidence for

origin and subsequent dispersal of Snowtrout across Asia, as initiated by the continued uplift of

210 the QTP. We were surprised by the robust and coherent biogeographic signal inherent in our

data, a consideration that, upon further examination, substantially extends previous results [6].

Central Asian phylogeography

214   Our phylogenetic results found five Central Asian species as sister to a significantly different *S.*

argentatus (Figs. 2, 3). Of these, S. *pseudoaksaiensis* is distributed in Lake Balkhash and its

216   catchment basin (Kazakhstan, central Asia), as well as the Ili River (northwest China) [41]), as is

S. argentatus. Similarly, *S. biddulphi* [40] and *S. eurystomus* [42] occur in the Tarim River and

218   tributaries (an endorheic basin in the Central Asiatic Desert of northwest China), whereas the

range of *S. intermediu*s is more extensive (i.e., Iran, Uzbekistan, Kazakhstan, Kyrgyzstan). It also

220   occurs in Fig. 3 as two distinct haplotypes.

In the ML tree, the Central Asia clade diverged at 11.6 Ma (CI=14.3-6.2), an estimate

222   comparable with previous results (11.4-10.5 Ma; [6]). Our timetree estimation for S. *argentatus*

(9.4 Ma; CI=14.3-6.2) also corroborates earlier estimates (9.6-8.9 Ma; [6]). It thus seems

224   reasonable to recognize Central Asia as the center of origin for schizothoracins, although we

note of this node was not placed as such in the statistically less supported BA analysis (Table 2;

226   Figure S1).

The next major split in our ML phylogeny defines two large nodes. The first

228   encompasses components of the QTP (save those in the far east) plus Bhutan (immediately

south), whereas the second incorporates the eastern QTP/ Southeast Asia plus those Himalayan

230   components that dispersed south via the Brahmaputra River then west. Our timeline is 7.3 Ma

(CI=11.3-4.7), again analogous with previously estimated values (8.3-7.6 Ma; [6]). We discuss

232   both large clades and their components below.

234   **Phylogeography of the Central QTP, YLTR, and "old Bhutan"**

The two species in the Central QTP (*S. molesworthi and S. malacanthus*) separated in early

236   Miocene (6.2 Ma; CI=10.2-3.7), whereas the YLTR West diverged late Pliocene at 4.2 Ma

(CI=7.3-2.4). Here, *S. macropogon* (3.2 Ma; CI=6.6-1.5, llate Pliocene) is sister to *S. waltoni*

238   which subsequently differentiated into western and eastern groups that are detectable via

msat DNA but not *cyt-b* [43]. It is, in turn, sister to a second YLTR West clade (*S. oconnori + S.*

240   *wangchiachii*).

The evolutionary history of *S. oconnori*, a species recognized as endemic to the YLTR

242   [41], underscores the role played by the QTP uplift in initiating Late Miocene divergence. During

this process, *S. oconnori* was subsequently separated into a western component above a

244   precipitous canyon in the YLTR, and an eastern component below [44]. Indeed, this hypothesis

is clearly sustained in our composite analyses (SH-ALRT, UFBOOT, AMOVA), with the western

246   clade (to include *S. wangchiatii*) not only significantly different from eastern *S. oconnori* (YLTR-

East), but also separated from it by an undescribed clade from Bhutan (i.e., Bhutan-1A; Fig. 3). A

248   second Bhutanese clade (Bhutan-1B) is sister to YLTR-East.

This seemingly indicates two distinct dispersals by *S. oconnori* from the ancestral YLTR.

250   One involved Bhutan-1A (the western clade), whereas a second represents the eastern clade

(Bhutan-1B). Both Bhutanese clades are currently unverified as to species but can seemingly be

252   allocated to *S. oconnori*. Quaternary climatic oscillations on the southern YTP not only yielded

evolutionarily significant units (ESUs) in both *S. waltoni* and *S. oconnori*, but also in Bhutanese

254   *Schizothorax* as well. Again, the BA-tree differs somewhat in that the YLTR lineages are further

subdivided. The YLTR-West region is broken by the Bhutan-1A clade, whereas Bhutan-1B (the

256   larger clade) becomes collapsed within the YLTR-East region.

258   **Phylogeography of the Eastern QTP and Southeast Asia**

The Eastern QTP/ Southeast Asia clade is represented by three significantly different clades (Fig.

260   3). The deepest divergence (i.e. Clade A from the remainder; Table S1) encompasses six species

from the Irrawaddy/ Salween rivers, with a timetree estimate at 5.6 Ma (CI=9.7-3.0). The

262   second (Clade B; Table S1) consists of seven species from the Yangtze River (at 1.7 Ma; CI=3.5-

0.8), whereas the third (Clade C; Table S1) involves eight species from the Mekong/ Salween

264   drainages. Divergence in the Mekong was seemingly driven by Late Pliocene orogenesis [37], an

estimate that is congruent with our derived timeline (at 2.4 Ma; CI=5.0-1.2). Three species

266   within the upper Salween River (i.e., *S. gongshanensis, S. lissolabiatus, S. nukiangensis*) display

low genetic differentiation and frequent haplotype sharing as a by-product of Quaternary

268   glaciations [39]. However, several species in clade B (such as S*. davidi, S. koslovi, S. chongli, S.

gongshanensis*) have yet to be fully described or documented, other than in general terms [40].

Dispersal of *Schizothorax* into Bhutan, Nepal, and western Himalaya

272   The YLTR originates on the QTP, flows eastward then abruptly south as it passes through the

Namche Barwa Syntaxis [5] where it was subsequently captured by the Brahmaputra River [7].

274   As such, it represents an important dispersal route from the QTP for freshwater species, and

also an important phylogenetic bookmark in that it provides an evolutionary transition between

276   species on the QTP versus those that diverged more recently, subsequent to westward

dispersal via the Lower Brahmaputra followed by northward movements into Himalayan

278   drainages that have previously down-cut to the south.

Our samples from Nepal form sister clades, each representing a distinct drainage, all of

280   which differ significantly amongst themselves, and from Southeast Asian clades (Fig. 2). Here,

additional samples from Bhutan (Bhutan-2; N=8) form a distinct clade sister to the eastern-most

282   drainage of Nepal (Koshi River Basin clade; N=11, Fig. 4, Table S1), thus effectively underscoring

the east-to-west transition mediated by dispersal from the lower Brahmaputra. Samples from

284   Bhutan-2 are currently not identified as to species, whereas those in the Koshi clade are

referred to *S. richardsonii, S. esocinus*, and *S. progastus.* The morphological similarities between

286   *S. richardsonii* and *S. progastus* are such that their discrimination in the field is problematic

[45], particularly at early life history stages. One hypothesis is that these inconsistencies may

288   contribute to the apparent phylogenetic assimilation of both species within several Koshi,

Karnali, and Gandaki river clades (Fig. 3).

290   Samples from the Gandaki and the Karnali rivers each form monophyletic clades as well.

The Gandaki (N=20) is partitioned among *S. richardsonii* (three clades), S. *progastus* (two

292   clades), and one representing a mixture of the two (Fig. 3). Samples from the Karnali River Basin

(far western Nepal, N=24; Fig. 4) fall within a composite that encompasses the Upper Ganges

294   (N=7) and Indus rivers (N=5; western Himalaya). Species from the Gandaki and Indus river

basins are *S. plagiostomus* and *S. progastus*, with *S. esocinus* also from the Indus. Once the

296   Koshi, Gandaki, and Karnali rivers reach the lowlands of India, they drain into the Ganges which

continues eastward to the Brahmaputra River then to the Bay of Bengal. (Fig. 1). The Indus,

298   however, now drains westward to the Arabian Sea.

Additional discrepancies appear between ML and BA-trees with regard to discrimination

300   among western Himalaya regions. Specifically, the BA-tree separates the Indus and Ganges into

separate clades, as sister to (and differentiated) from the Karnali River/ Rara Lake clade.

Phylogeography of Rara Lake, Nepal

304   Three morphologically differentiated and reproductively isolated *Schizothorax* (i.e., *S.*

macrophthalmus, S. nepalensis, S. raraensis) occupy separate trophic niches in Lake Rara

306   (Karnali River Basin, northwestern Nepal: [46]), with *S. raraensis* and *S. nepalensis* listed as

critically endangered (IUCN, 2017; http://www.iucnredlist.org/details/168564/0/). Of the three,

308   only *S. nepalensis* could be differentiated using mtDNA and, given this, it was hypothesized [47]

that *S. macrophthalmus* and *S. raraensis* shared mtDNA haplotypes via hybridization. Similarly,

310   two morphologically distinct Snowtrout in the upper Karnali River (i.e., *S. richardsonii* and *S.*

progastus) could not be discriminated as well, with each similarly hypothesized as exemplifying

312   interspecific mitochondrial exchange [47]. Our mitochondrial marker could not resolve this

hypothesis, thus necessitating further nuclear analysis so as to unequivocally test the role of

314   hybridization as a source of phylogenetic discordance. Such an examination, however, is further

complicated by polyploidy and ploidy variance within *Schizothorax.*

316   The geologic age of Lake Rara is estimated as ~60kya [48]. Its endemic species do not

form a monophyletic clade, but cluster instead with tributaries of the Karnali River that drain

318   the lake (Fig. 3). Haplotypes of *S. richardsonii* and *S. progastus* in the Karnali River Basin did not

differ significantly from one another yet were significantly different when compared with

320   conspecifics in the Koshi and Gandaki rivers [47]. Our analyses confirmed these results (Fig. 3).

Speciation in the Karnali River Basin could thus be poorly resolved with respect to

322 mitochondrial divergence, with a genetic signal likely obscured by the limited framework within

which ancestral alleles could sort, allowing novel haplotypes to arise. More refined molecular

324 approaches, such as reduced-representation genomics [32,33,34,49,50], could potentially

provide the necessary resolution, although it will again be compounded by the rampant

326 polyploidy found in Cyprininae.

328 **Dispersal of Snowtrout into western Himalayan tributaries**

Recent studies [6,37,51,52] suggest the current distribution of Snowtrout has been shaped by

330 Late Miocene uplift of the QTP, with vicariant events occurring within headwaters of several

rivers in the Eastern QTP/ Southeast Asia clade. However, these are localized perspectives

332 rather than broadly regional, due largely to conflicting interpretations regarding the evolution

of Himalayan paleodrainages, particularly those in the western Himalaya. A second confounding

334 factor is a deficit of molecular samples from the Central Himalaya (a crucial link between

eastern and western components of the Himalaya).

336   The southern QTP and western Himalaya are now drained by the contemporary Indus

River which extends south and west from the Karakorum Range, rather than from the High

338 Himalaya. Given its western placement, the geomorphic history of the Indus is a key component

with regards to the biogeography of *Schizothorax* in the Himalaya. The relationship of the Indus

340 with the Ganges immediately to the east is unsettled, with several proposed hypotheses [52].

Relative stasis between the two is the most conservative, whereas others involve various

342 stream-capture scenarios, such as: The Ganges initially flowing westward to the Indus then to

the Arabian Sea; The Indus flowing eastward into the Ganges then to the Bay of Bengal; and

344   lastly, an earlier capture by the Ganges of four western Himalaya rivers (i.e., Jhelum, Chenab,

Ravi, Sutlej) that are now incorporated within the contemporary Indus Basin.

346   The latter hypothesis is supported by neodymium isotope and seismic reflection data in

drill cores extracted from the detrital fan of the Indus River, as deposited for >30 Ma in the

348   Arabian Sea. These data document a major depositional shift occurring at ~5 Ma, with

Himalayan erosion superseding that deposited earlier from the Indus suture zone to the north

350   [53]. The shift is hypothesized as documenting the captured by the Indus of several large

Punjabi rivers (i.e., Jhelum, Chenab, Ravi, Sutlej), as reflected by the current basin topography).

352   However, this hypothesis is not without controversy. A subsequent comparison of sediments

from northern Pakistan with those of the Indus Fan suggests that the documented transitions in

354   the latter were a response to differential erosion rates in the source region (the Indus), rather

than a result of drainage rearrangements (as above) [52].

356   The strong east-west divergence identified in our data among Himalayan clades is the

biological data from which to interpret the geomorphic evolution of Himalayan drainage

358   systems. In this sense, our phylogeographic results are a biological corroboration of the capture

by the Indus of those drainages previously allocated to the Ganges. *Schizothorax* from the

360   Lower Ganges River diverged from a clade representing the Upper Ganges Basin and the Indus

Basin. Similar patterns have been noted for Indus and Gangetic dolphin [54]. These data also

362   suggest *Schizothorax* in the western Himalaya (i.e., Koshi, Gandaki, and Karnali) was isolated

prior to the capture of the Jhelum by the Indus.

A confused taxonomy for *Schizothorax*

366   Several species were found to be polyphyletic in our analyses, with individuals forming

biogeographically defined clusters that differed significantly from one another. In addition,

368   multiple cases were found where samples within the same region also failed to cluster with

conspecifics (e.g., Central Asia; Eastern QTP/Southeast Asia; Koshi; Gandaki; Karnali; Ganges).

370   This phenomenon highlights several trenchant issues manifested within the

schizothoracines: Incongruence between molecular and morphological phylogenies [6,37,51],

372   and convergent evolution (e.g., juxtaposition of trophic and spatial niches across species [55]).

An observed lack of association between morphological and molecular characters may be due

374   either to incomplete lineage sorting, or rapid diversification [18]. If the latter, then mtDNA as a

molecular marker will be incapable of resolving these issue [6]. These issues are amplified and

376   biological interpretations consequently hindered by the potential for DNA sequences within

online databases to be incorrectly annotated with regards to taxonomy [56].

Conclusions

380   *Schizothorax* in the Himalaya is a monophyletic clade that originated in Central Asia,

subsequently dispersed southward into Bhutan via down-cutting rivers, then westward across

382   the Eastern QTP and into Southeast Asia. Subsequent capture of the YLTR by the Brahmaputra

allowed for an east-to-west dispersal below (i.e., south of) the Himalaya, with subsequent

384   movements northward into sequential Himalaya drainages, first Bhutan, then Nepal, followed

by the western Himalaya. Our molecular data also serve to corroborate western drainages

386   previously affiliated with the Ganges as being subsequently captured by the Indus River.

Many morphologically recognized species have seemingly diverged substantially across

388   basins, suggesting the potential for cryptic speciation and/or undescribed relictual biodiversity.

Additionally, more detailed molecular studies would promote conservation planning at the

390   regional scale but will also require the analysis of populations now geographically isolated

across the Himalaya. This, in turn, will necessitate cooperation/ collaboration among

392   neighboring countries, particularly given the recognized (as well as predicted) impacts of

climate change on the Third Pole in general, and *Schizothhorax* in particular [22].

Methods

Ethics Statement

398   Bhutanese collecting permits were provided by the Royal Government's Ministry of Agriculture

and Forestry (MoAF). Sampling protocols were validated by the University of Arkansas

400   Institutional Animal Care and Use Committee (UA-IACUC).

402   **Drainages of Nepal and Bhutan**

Three major drainage basins in Nepal (east-to-west: Koshi, Gandaki, and Karnali rivers; Fig. 4),

404   support eight tributaries, several of which originate on the QTP, although we found no evidence

for historic dispersal of *Schizothorax* directly into Nepal. Mountainous slopes in the Koshi River

406   are more eroded and abruptly elevated, given that the monsoon has a stronger effect on

eastern Nepal than further west. Central Nepal is drained by the Gandaki Basin, and its western

408   tributary (Kali Gandaki) is recognized as a vicariant east-to-west break. Western Nepal, on the

other hand, is drained by the Karnali Basin, and includes one of our study sites (i.e., Lake Rara, a

410   RAMSAR site with a surface area of 10.6 km^2 at 2,990m elevation). All three basins drain south

into the east-flowing Ganges River that subsequently joins with the Brahmaputra River and

412   terminates in the Bay of Bengal.

Bhutan, on the other hand, displays five major north-south river systems (Fig. 5). The

414   Amo Chhu, Wang Chhu, and Punatsang Chhu (=Sankosh) drain western Bhutan, with the Amo

Chhu and Wang Chhu descending from the QTP and flowing southeasterly through west-central

416   Bhutan. The Punatsang Chhu drains the Great Himalayan Range. All three subsequently pass

into West Bengal (India).

418   The Manas and its several major tributaries drain central-eastern Bhutan. There, the

Dangme Chhu flows southwesterly from Arunachal Pradesh (India), whereas the Kuri Chhu

420   (from the QTP), the Chamkhar Chhu, and the Mangde Chhu drain central Bhutan. All Bhutanese

basins drain eventually into the Brahmaputra River (India), which curves south and west

422   through Bangladesh to merge with the Ganges River, then to the Bay of Bengal (Fig. 1).

424   **The origin of *Schizothorax* and its outgroups**

Snowtrout (Fig. 6) is an ideal organism to decipher the geologic processes that drove historic

426   and contemporary fish distributions in the Himalaya. It contains a diverse array of species

distributed across deserts, lakes, and rivers of the QTP, as well as the snow-fed tributaries of

428   the Himalaya [57,58]. Some 28 *Schizothorax* species are recorded in the Himalayan, sub-

Himalayan, and Tibetan regions [59].

430   Several hypotheses have been offered to explain the origin and dispersal of *Schizothorax*

across the Himalaya and Asia. Previous researchers [60] suggested the subfamily

432  Schizothoracinae initially dispersed into the QTP from southeastern Asia, and subsequently

became isolated by orogeny and climate change. Others [61] proposed that tectonism isolated

434  a primitive barbin clade on the Tibetan Plateau which then served as progenitor for a

contemporary *Schizothorax* radiation.

436   A recent revision of the subfamily Cyprininae [62] was based on a combined series of

molecular markers: *cyt-b* (N=791), mitogenomes (N=85), and nuclear RAG-1 sequences (N=97)].

438  Results retained *Schizothorax* within the monophyletic Tribe Schizothoracini that (depending

upon the marker) evolved prior to or after the Barbini. Thus, in selecting outgroups, we avoided

440  Barbini and instead fixed on tribes identified as evolving prior to the Schizothoracini. These

included (in sequence): Torini (i.e., *Tor* and *Neolissochilus*), Smilogastrini (i.e., *Puntius, Pethia*),

442  and Acrossocheilini (i.e., *Acrossocheilus*, the most immediate outgroup) (Table S1).

444  **Sampling and the amplification/ sequencing of mtDNA**

Nepali samples (N=53) represented five *Schizothorax* species. Three [i.e., *S. macrophthalmus*

446  (N=3), *S. nepalensis* (N=4), and *S. raraensis* (N=3); total N=10] represented a landlocked species-

flock within Rara Lake (Karnali River drainage; Fig. 4). Two other species [*S. richardsonii* (N=28),

448  and *S. progastus* (N=15)] are riverine and were sampled from (west-to-east): Karnali River

(N=12/N=2, respectively), Gandaki River (N=11/9), and Koshi River (N=5/4).

450   Nepali samples were supplemented with those from Bhutan (N=19) collected from four

drainage basins (west-to-east): Wang Chhu (N=1), Punatsang Chhu (N=13), Mangde Chhu (N=3),

452   Dangme Chhu (N=1), Nyera Ama Chhu (N=1); Fig. 5). Genomic DNA was extracted using the

Qiagen DNeasy kit, per manufacturer's instructions. In addition, we accessed 67 mtDNA *cyt-b*

454   sequences representing 36 species (28 *Schizothorax* plus eight outgroups [*Tor* (N=2),

*Neolissochilus (*N=1), *Puntius* (N=1), *Pethia* (N=2), *Acrossocheilus* (N=2); Table S1] (National

456   Institute of Health GENBANK®, https://www.ncbi.nlm.nih.gov/genbank/). Total number of

individuals = 139.

Polyploidy

460   The subfamily Cyprininae displays rampant polyploidy (>400 species; [62,63], an important

consideration in molecular studies, in that methodological accommodations across domains

462   (e.g., population genetics) are lacking. This is in particular problematic within a phylogenetic

context, given the greater redundancy of nuclear genes in polyploid genomes. All may

464   potentially diverge one from another over evolutionary time, such that results could differ

markedly depending upon which 'sub-genome' is accessed when polyploid molecular data are

466   phylogenetically evaluated [64].

Given this, phylogenetic analyses of polyploid species are often limited in practice either

468   to mtDNA data (due to its haploid nature), or nuclear genes cloned into bacteria and

subsequently isolated as single-copy [such as Recombination Activating Gene 1 (RAG-1); [62],

470   though the latter is recognized as more conservative evolutionarily, and without a timetree as

an option [65]. With regards to *Schizothorax,* polyploidization may represent an evolutionary

472   mechanism that facilitated an invasion of high-elevation habitats [63].

474 **Phylogenetic analyses**

We amplified mtDNA *cyt-b* using published primers [66], with products being enzymatically

476 purified, sequenced using BɪɢDʏᴇ (ver.3.1) chemistry (Applied Biosystems Inc. [ABI], Forest City,

CA), and analyzed on an ABI Prism 3700 Genetic Analyzer (W.M. Keck Center for Comparative

478 and Functional Genomics, University of Illinois, Urbana/Champaign). Sequences were

edited/aligned using Sᴇǫᴜᴇɴᴄʜᴇʀ v.5.3 (Gene Codes Inc., Ann Arbor, MI), with nucleotide

480 composition and uncorrected p-distances derived among groups (Mᴇɢᴀ X; [67].

We then developed a dataset that included only variable sites (N=365bp: DNAsp v.6;

482 [68]) which was subsequently employed in a Bayesian approach (Gᴇɴᴇʟᴀɴᴅ v.4.0.3: [23,69]) to

infer genetic clustering of individuals to infer the number of distinct groups (*K*) and designate

484 group membership. Ten runs of 100,000 MCMC iterations and a thinning interval of 100 were

used, with *K* varying between 1 and 15, but without use of optional geo-referencing, due to the

486 absence of coordinates for most sequences in the GᴇɴBᴀɴᴋ® metadata. The estimated number

of populations (K) was defined using the maximum posterior estimate. We also conducted an

488 analysis of molecular variation (AMOVA: Aʀʟᴇǫᴜɪɴ v.3.5.1.2; [25]) so as to evaluate geographic

"groups" (as informed by Gᴇɴᴇʟᴀɴᴅ results and represented in Fig. 2). Here, we defined

490 "populations" (= distinct clades) as being haplotypes within groups, per [70]. Probabilities were

ascertained using 1,000 permutations, and statistical significance was assigned using a

492 Bonferroni-correction.

We derived phylogenetic clades using a maximum likelihood (ML) analysis (IQ-Tʀᴇᴇ v2;

494 [71]), with the best-fitting substitution model generated via MᴏᴅᴇʟFɪɴᴅᴇʀ, with flexible rate

heterogeneity (e.g. gamma-distributed) across sites [72]. This was accomplished by exploring

496   model space hierarchically, adding parameters (e.g. rate categories) sequentially until the

model failed to improve. Support for the resulting tree, inferred using the selected model, was

498   explored using 1,000,000 ultrafast bootstrap approximations (UFBOOT2; [73]). We also assessed

branch support using the SH-aLRT test (Shinodara-Hasegawa approximate Likelihood Ratio

500   Test), with values ≥ 80% indicating strongly supported clades, whereas those for UFBOOT are ≥

95% (IQ-TREE v.2 Manual: http://www.iqtree.org).

502   To employ a Bayesian approach, we first derived an XML input file using the graphical

user interface (i.e., BEAUTY) for the phylogenetic software suite BEAST (v2.61; [74]), using a

504   relaxed molecular clock with an uncorrelated lognormal rate distribution and a Yule process

prior. Each analysis was run for 100 million generations, with samples taken every 1,000th

506   iteration, following a burn-in of 20%. Effective sample size (ESS) was verified as >200 across all

parameters with MCMC chain convergence visually inspected using TRACER [75]. Trees from

508   three independent runs were combined (via LOGCOMBINER) and represented as a maximum

clade credibility (MCC in TREEANNOTATOR: [74]), with nodal support representing posterior

510   probabilities. The resulting tree was visualized using FIGTREE (v.1.4.3; [76]).

Topologies from ML and BA analyses were statistically compared by assessing each with

512   respect to the original sequence alignment using IQ-TREE2 v2 [70]. To do so, we examined

support for each using RELL approximation with 10,000 replicates using: (a) Raw change in log-

514   likelihoods; (b) Bootstrap proportions [77]; (c) Kishino-Hasegawa test [78]; (d) Shimodaira-

Hasegawa test [79]; (e) Approximately Unbiased test [80]; and (f) Expected Likelihood weights

516   [81].

518 **Developing a timetree**

For molecular dating, we employed a rapid, high-performance ML-based model (RELTIME, MEGA

520 X; [67]) that incorporates a relative rate framework (RRF) within which the evolutionary rates of

sister lineages are contrasted against a minimum rate change, as diagnosed between

522 evolutionary lineages and their respective descendants. The computational speed and accuracy

of the model is appropriate for the analysis of large data sets (>100 ingroup sequences; [82]).

524 The procedural steps for RELTIME are described in [83]. The evolutionary model for our ML-

based phylogenetic tree (i.e., TN+G4+I) was also employed in RELTIME.

526 We estimated a divergence time based upon the earliest fossil recognition for the

Schizothoracini (e.g., Oligocene: *Paleoschizothorax qaidamensis*; [84]). This Qaidam Basin fossil

528 was estimated from its stratigraphic formation as being Eocene-to-Oligocene [85], and given

this, we chose an Oligocene initiation (33Ma) as our divergence time.

Acknowledgments

532 Nepali samples were obtained from the University of Kansas Natural History Museum

(http://www.kansastravel.org/lawrence/kansasnaturalhistory.htm) and represent a subset of

534 those collected by DRE during completion of a 1996 Fulbright Award. We thank Curator Leo

Smith and Collection Manager Andrew Bentley for their prompt response and support. We also

536 acknowledge the National Research Centre for Riverine and Lake Fisheries (NRCRLF, Bhutan) for

providing logistic support and personnel with which to sample fishes. Analytical resources were

538 provided by the Arkansas Economic Development Commission (Arkansas Settlement Proceeds

Act of 2000) and the Arkansas High Performance Computing Center (AHPCC). This research was

Author Contributions

548 BR, MRD, MED, DRE, KW, and ST conceived and designed the study; DRE, KW, CL, GPK, PN, SN,

SD, TKC, and ZDZ collected samples in the field; BL, ZA and MED analyzed the data; BL, MRD,

550 MED, and ZA wrote the article; All authors discussed the results and commented on the

manuscript.

Additional Information

554 **Competing Interests:** The authors declare no competing interests.

Data Availability Statement

mtDNA data: Dryad Digital Repository (http://dx.doi.org/upon-acceptance).

References

[1] Yin, A. & Harrison, T. M. Geologic evolution of the Himalayan Tibetan orogen. *Annu. Rev. Earth Planet. Sci.*, **28**, 211–280. https://doi.org/10.1146/annurev.earth.28.1.211 (2000).

[2] Xu, J. & Grumbine, R. E. Building ecosystem resilience for climate change adaptation in the Asian highlands. *WIREs Clim. Change*, **5**, 709–718. https://doi.org/10.1002/wcc.302 (2014).

[3] Pellissier, L., Heine, C., Rosauer, D. F. & Albouy, C. Are global hotspots of endemic richness shaped by plate tectonics? *Biol. J. Linn. Soc.*, **123**, 247–261. https://doi.org/10.1093/biolinnean/blx125 (2017).

[4] van Hinsbergen, D. J. *et al.* Greater India Basin hypothesis and a two-stage Cenozoic collision between India and Asia. *Proc. Nat. Acad. Sci. U.S.A.*, **109**, 7659–7664. https://doi.org/10.1073/pnas.1117262109 (2012).

[5] Brookfield, M. E. The evolution of the great river systems of southern Asia during the Cenozoic India-Asia collision: Rivers draining southwards. *Geomorphology,* **22**, 285–312. https://doi.org/10.1016/S0169-555X(97)00082-2 (1998).

[6] He, D. & Chen, Y. Biogeography and molecular phylogeny of the genus *Schizothorax* (Teleostei: Cyprinidae) in China inferred from cytochrome b sequences. *J. Biogeogr.*, **33**, 1448–1460. https://doi.org/10.1111/j.1365-2699.2006.01510.x (2006).

[7] Bracciali, L., Najman, Y., Parrish, R. R., Akhter, S. H. & Millar, I. The Brahmaputra tale of tectonics and erosion: Early Miocene river capture in the Eastern Himalaya. *Earth Planet. Sci. Lett.*, **415**, 25–37. https://doi.org/10.1016/j.epsl.2015.01.022 (2015).

[8] Li, S. *et al.* The evolution of Yarlung Tsangpo River: Constraints from the age and provenance of the Gangdese conglomerates, southern Tibet. *Gondwana Res.*, **41**, 249–266. https://doi.org/10.1016/j.gr.2015.05.010 (2017).

[9] Govin, G. *et al.* Timing and mechanism of the rise of the Shillong Plateau in the Himalayan foreland. *Geology*, **46**, 279–282. https://doi.org/10.1130/G39864.1 (2018).

[10] Favre, A. *et al.* The role of the uplift of the Qinghai-Tibetan Plateau for the evolution of Tibetan biotas. *Biol. Rev.*, **90**, 236–253. https://doi.org/10.1111/brv.12107 (2015).

[11] Deng, T. *et al.* Review: Implications of vertebrate fossils for paleo-elevations of the Tibetan Plateau. *Glob. Planet. Change,* **174**, 58–69. https://doi.org/10.1016/j.gloplacha.2019.01.005 (2019).

[12] Schluter, D. Speciation, ecological opportunity, and latitude. *Am. Nat.*, **187**, 1–18. https://doi.org/10.1086/684193 (2016).

[13] Jacobsen, D., Milner, A. M., Brown, L. E. & Dangles, O. Biodiversity under threat in glacier-fed river systems. *Nat. Clim. Change*, **2**, 361–364. https://doi.org/10.1038/NCLIMATE1435 (2012).

[14] Li, J. *et al.* Climate refugia of snow leopards in High Asia. *Biol. Conserv.* **203**, 188–196. https://doi.org/10.1016/j.biocon.2016.09.026 (2016).

[15] Scherrer, D. & Körner, C. Topographically controlled thermal-habitat differentiation buffers alpine plant diversity against climate warming. *J. Biogeogr.*, **38**, 406–416. https://doi.org/10.1111/j.1365-2699.2010.02407.x (2011).

[16] Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B. & Kent, J. Biodiversity hotspots for conservation priorities. *Nature*, **403**, 853–858. https://doi.org/10.1038/35002501 (2000).

[17] Fricke, R., Eschmeyer, W. N. & van der Laan R. (eds). *Eschmeyer's catalog of fishes: Genera, species, references.* http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp (2020).

[18] Zhao, K. I. *et al.* The youngest split in sympatric Schizothoracine fish (Cyprinidae) is shaped by ecological adaptations in a Tibetan Plateau glacier lake. *Mol. Ecol.*, **18**, 3616–3628. https://doi.org/10.1111/j.1365-294X.2009.04274.x (2009).

[19] Yao, T. *et al.* Third pole environment (TPE). *Environ. Dev.*, **3**, 52–64. https://doi.org/10.1016/j.envdev.2012.04.002 (2012).

[20] IPCC (Intergovernmental Panel on Climate Change). IPCC Special Report on the Ocean and Cryosphere in a Changing Climate (eds. Pörtner, H.-O., Roberts, D.C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A., Nicolai, M., Okem, A., Petzold, J., Rama B. & Weyer, N.M.), Glan, Switzerland (2019).

[21] Zhu, G-F., Gao, L-M. & Li, D-Z. Protect third-pole's fragile ecosystem. *Science*, **362**, 1368. https://doi.org/10.1126/science.aaw0443 (2018).

[22] Sharma, A. Dubey, V.K., Johnson, J.A., Rawal, Y.K. & Sivakumara, K. Is there always space at the top? Ensemble modeling reveals climate-driven high-altitude squeeze for the vulnerable snow trout *Schizothorax richardsonii* in Himalaya. *Ecol. Indic.,* **120**, 106900. https://doi.org/10.1016/j.ecolind.2020.106900 (2020).

[23] Guillot, G., Mortier F. & Estoup, A. GENELAND: A computer package for landscape genetics. *Mol. Ecol. Notes*, **5**, 712–715. https://doi.org/10.1111/j.1471-8286.2005.01031.x (2005).

[24] Guillot G. Inference of structure in subdivided populations at low levels of genetic differentiation—the correlated allele frequencies model revisited. *Bioinformatics*, **24**, 2222–2228. https://doi.org/10.1093/bioinformatics/btn419 (2008).

[25] Excoffier, L. & Lischer H. E. L. ARLEQUIN suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.*, **10**, 564–567. https://doi.org/10.1111/j.1755-0998.2010.02847.x (2010).

[26] Whipple, K. X. The influence of climate on the tectonic evolution of mountain belts. *Nature Geosci.*, **2**, 97–99. https://doi.org/10.1038/ngeo413 (2009).

634 [27] Hopken, M. W., Douglas, M. R. & Douglas, M. E. Stream hierarchy defines riverscape genetics of a North American desert fish. *Mol. Ecol.*, **22**, 956–971. https://doi.org/10.1111/mec.12156 (2013).

636 [28] Douglas, M. R., Brunner, P. C. & Douglas, M. E. Drought in an evolutionary context: Molecular variability in Flannelmouth Sucker (*Catostomus latipinnis*) from the Colorado

638 River Basin of western North America. *Freshw. Biol.*, **48**, 1254–1273. https://doi.org/10.1046/j.1365-2427.2003.01088.x (2003).

640 [29] Douglas, M. R. *et al.* Anthropogenic impacts drive niche and conservation metrics of a cryptic rattlesnake on the Colorado Plateau of western North America. *Royal Soc. Open*

642 *Sci.*, **3**, 160047. https://doi.org/10.1098/rsos.160047 (2016).

[30] Bishop, P. Drainage rearrangement by river capture, beheading and diversion. *Prog. Phys.*

644 *Geogr.*, **19**, 449–473. https://doi.org/10.1177/030913339501900402 (1995).

646 [31] Strange, R. M. & Burr, B. M. Intraspecific phylogeography of North American Highland fishes: A test of the Pleistocene vicariance hypothesis. *Evolution*, **51**, 885–897.

648 https://doi.org/10.1111/j.1558-5646.1997.tb03670.x (1997).

[32] Bangs, M. R., Douglas, M. R., Mussmann, S. M. & Douglas, M. E. Unraveling historical

650 introgression and resolving phylogenetic discord within *Catostomus* (Osteichthys: Catostomidae). *BMC Evol. Biol.,* **18**, 86. https://doi.org/10.1186/s12862-018-1197-y

652 (2018).

[33] Bangs, M. R., Douglas, M. R., Mussmann, S. M. & Douglas, M. E. Reticulate evolution as a

654 management challenge: Patterns of admixture with phylogenetic distance in endemic fishes of Western North America. *Evol. Applic.,* **13**, 1400–1419. https://doi.org/

656 10.1111/eva.13042 (2020).

[34] Chafin, T.K., Douglas, M.R. & Douglas M.E. Hybridization drives genetic erosion in sympatric

658 desert fishes of western North America. *Heredity,* **123**, 759–773. https://doi.org/10.1038/s41437-019-0259-2 (2019).

660 [35] Radinger, J. *et al.* The future distribution of river fish: The complex interplay of climate and land use changes, species dispersal and movement barriers. *Glob. Change Biol.*, **23**,

662 4970–4986. https://doi.org/10.1111/gcb.13760 (2017).

[36] CREP-Critical Ecosystem Partnership Fund. *The biodiversity hotspots.*

664 http://www.cepf.net/resources/hotspots/Pages/default.aspx (2017).

[37] Qi, D. *et al.* The biogeography and phylogeny of schizothoracine fishes (*Schizopygopsis*) in

the Qinghai-Tibetan Plateau. *Zool. Scr.*, **44**, 523–533. https://doi.org/10.1111/zsc.12116 (2015).

[38] Chen, W., Shen, Y., Gan, X., Wang, X. & He, S. Genetic diversity and evolutionary history of the *Schizothorax* species complex in the Lancang River (upper Mekong). *Ecology and*

Evolution, **6**, 6023–6036. https://doi.org/10.1002/ece3.2319 (2016).

[39] Chen, W., Yue, X. & He, S. Genetic differentiation of the *Schizothorax* species complex (Cyprinidae) in the Nujiang River (upper Salween). *Sci. Rep.*, **7**, 5944. https://doi.org/10.1038/s41598-017-06172-5 (2017).

[40] Wanghe, K. *et al.* Phylogeography of *Schizopygopsis stoliczkai* (Cyprinidae) in Northwest Tibetan Plateau area. *Ecol. Evol.*, **7**, 9602–9612. https://doi.org/10.1002/ece3.3452
(2017).

[41] Froese, R. & Pauly, D. (eds). FishBase 2000. World Wide Web electronic publication.
678 www.fishbase.org (2019).

[42] Huo, T. B. *et al.* Length–weight relationships of 16 fish species from the Tarim River, China.
680 *J. Appl. Ichthyol.*, **28**, 152–153. https://doi.org/10.1111/j.1439-0426.2011.01899.x (2012).

682 [43] Guo, X-Z. *et al.* Phylogeography of the threatened tetraploid fish, *Schizothorax waltoni*, in the Yarlung Tsangpo River on the southern Qinghai- Tibet Plateau: Implications for
684 conservation. *Sci. Rep.*, **9**, 2704. https://doi.org/10.1038/s41598-019-39128-y (2019).

686 [44] Guo, X-Z. *et al.* Phylogeography and population genetics of *Schizothorax o'connori*: Strong subdivision in the Yarlung Tsangpo River inferred from mtDNA and microsatellite
688 markers. *Sci. Rep.*, **6**, 29821. https://doi.org/10.1038/srep29821 (2016).

[45] Barat, A., Ali, S., Sati, J. & Sivaraman, G.K. Phylogenetic analysis of fishes of the subfamily
690 Schizothoracinae (Teleostei: Cyprinidae) from Indian Himalayas using cytochrome b gene. *Indian J. Fish.*, **59**, 43–47. https://www.researchgate.net/publication/259772138 (2012)
692

[46] Terashima, A. Three new species of the cyprinid genus *Schizothorax* from Lake Rara,
694 northwestern Nepal. *Jap. J. Ichthyol.*, **31**, 122–135. https://doi.org/10.11369/jji1950.31.122 (1984).

696 [47] Dimmick, W. W. & Edds, D. R. Evolutionary genetics of the endemic schizorathicine (Cypriniformes: Cyprinidae) fishes of Lake Rara, Nepal. *Biochem. Syst. Ecol.*, **30**, 919–929.
698 https://doi.org/10.1016/S0305-1978(02)00030-3 (2002).

[48] Zakaria, G. Late Quaternary Seismicity and Climate in Western Nepal: Himalaya. Applied
700 Geology. Unpubl. Ph.D. dissertation. Université Ghent, Department of Geology, Grenoble, France, Ghent, Belgium. https://tel.archives-ouvertes.fr/tel-
702 01935412/document (2018).

[49] Martin, B.T. *et al.* Contrasting signatures of introgression in the contact zones of North
704 American Box Turtle (*Terrapene* spp.). *Mol. Ecol.*, **2020**, 00:01–17. https://doi.org/10.1111/mec.15622 (2020).

706 [50] Mussmann, S.M., Douglas, M.R., Oakey, D.D. & Douglas M.E. Defining relictual biodiversity: Conservation units in Speckled Dace (Cyprinidae: *Rhinichthys osculus*) of the Greater
708 Death Valley Ecosystem. *Ecol. Evol.,* **2020**, 00:1–20. https://doi.org/10.1002/ece3.6736) (2020).

710   [51] Qi, D. *et al*. Convergent, parallel and correlated evolution of trophic morphologies in the subfamily Schizothoracinae from the Qinghai-Tibetan Plateau. *PLoS ONE,* **7**(3), e34070.
712   https://doi.org/10.1371/journal.pone.0034070 (2012).

    [52] Chirouze, F. *et al*. Detrital thermochronology and sediment petrology of the middle
714   Siwaliks along the Muksar Khola section in eastern Nepal. *J. Asian Earth Sci.,* **44**, 94–106. https://doi.org/10.1016/j.jseaes.2011.01.009 (2013).

716   [53] Clift, P. D. & Blusztajn, J. Reorganization of the western Himalayan river system after five million years. *Nature*, **438**, 1001–1003. https://doi.org/10.1038/nature04379 (2013).

718   [54] Braulik, G. T. *et al.* One species or two? Vicariance, lineage divergence and low mtDNA diversity in geographically isolated populations of South Asian River dolphin. *J. Mammal.*
720   *Evol.,* **22**, 111–120. https://doi.org/10.1007/s10914-014-9265-6 (2015).

    [55] Tang, Y. *et al*. Convergent evolution misled taxonomy in Schizothoracine fishes
(Cypriniformes: Cyprinidae). *Mol. Phylogenet. Evol.*, **134**, 323–337. https://doi.org/10.1016/j.ympev.2019.01.008 (2019).

[56] Vilgalys, R. Taxonomic misidentification in public DNA databases. *New Phytol.*, **160**, 4–5. https://doi.org/10.1046/j.1469-8137.2003.00894.x (2003).

[57] Edds, D. R. Fishes in Nepal: Ichthyofaunal surveys in seven nature reserves. *Ichthyol. Explor. Freshw.*, 1**8**, 277–287.
https://www.researchgate.net/publication/228787571_Fishes_in_Nepal_Ichthyofaunal _surveys_in_seven_nature_reserves (2007).

[58] Yang, J., Yang, J. X. & Chen, X. Y. A re-examination of the molecular phylogeny and biogeography of the genus *Schizothorax* (Teleostei: Cyprinidae) through enhanced
sampling with an emphasis on the species in the Yunnan-Guizhou plateau, China. *J. Zool. Syst. Evol. Res.*, **50**, 184–191. https://doi.org/10.1111/j.1439-0469.2012.00661.x (2012).

[59] Sharma, C. M. Freshwater fishes, fisheries, and habitat prospects of Nepal. *Aquat. Ecosys. Health Manag.*, **11**, 289–297. https://doi.org/10.1080/14634980802317329 (2008).

[60] Das, S. M. & Subla, B. A. The ichthyofauna of Kasmir: Part II The speciation of Kasmir fishes with two new records of species. *Ichthyologica*, **3**, 57–62 (1964).

[61] Cao, W. X., Chen, Y. Y., Wu, Y. F. & Zhu, S. Q. Origin and evolution of Schizothoracine fishes in relation to the upheaval of the Qinghai-Xizang Plateau in *Studies on the Period,*
Amplitude and Type of the Uplift of the Qinghai-Xizang Plateau (ed. Li, Q. F.) 118–129 (Beijing, China: Science Press, 1981).

[62] Yang, L. *et al.* Phylogeny and polyploidy: Resolving the classification of cyprinine fishes (Teleostei: Cypriniformes). *Mol. Phylogenet. Evol.*, **85**, 97–116.
https://doi.org/10.1016/j.ympev.2015.01.014 (2015).

    [63] Wang, X., Gan, X., Li, Y. & He, S. Cyprininae phylogeny revealed independent origins of the
746   Tibetan Plateau endemic polyploid cyprinids and their diversifications related to the Neogene uplift of the plateau. *Sci. China Life Sci.,* **59**, 1149–1165.
748   https://doi.org/10.1007/s11427-016-0007-7 (2016).

750 [64] Taylor, J.S., Braasch, I., Frickey, T., Meyer, A. & Van de Peer, Y. Genome duplication, a trait shared by 22,000 species of ray-finned fish. *Genome Res.*, **13**, 382–390. https://doi.org/10.1101/gr.640303 (2003).

752 [65] Groth, J. G. & Barrowclough, G. F. Basal divergences in birds and the phylogenetic utility of the nuclear RAG-1 gene. *Mol. Phylogenet. Evol.*, **12**, 115–123.
754 https://doi.org/10.1006/mpev.1998.0603 (1999).

[66] Houston, D. D., Evans, R. P. & Shiozawa, D. K. Evaluating the genetic status of a Great Basin
756 endemic minnow: The relict dace (*Relictus solitarius*). *Cons. Gen.*, **13**, 727–742. https://doi.org/10.1007/s10592-012-0321-6 (2012).

758 [67] Kumar, S. Stecher, G., Li, M., Knyaz, C. & Tamura, K., MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Mol. Biol. Evol.*, **35**, 1547–1549.
760 https://doi.org/10.1093/molbev/msy096 (2018).

[68] Rozas, J. *et al*. DNASP 6: DNA sequence polymorphism analysis of large datasets. *Mol. Biol.*
762 *Evol.* **34**, 3299–3302. https://doi.org/10.1093/molbev/msx248 (2017).

[69] R-Studio Team. RStudio: Integrated Development for R. RStudio, PBC, Boston, MA.
764 http://www.rstudio.com/ (2020).

[70] Solecki, A. M., Skevington, J. H., Buddle, C. M. & Wheeler, T. A. Phylogeography of higher
766 Diptera in glacial and postglacial grasslands in western North America. *BMC Ecol.*, **19**, 53. https://doi.org/10.1186/s12898-019-0266-4 (2019).

768 [71] Minh, B. Q. *et al*. IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.,* **37**, 1530–1534.
770 https://doi.org/10.1093/molbev/msaa015 (2020).

[72] Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
772 MODELFINDER: Fast model selection for accurate phylogenetic estimates *Nat. Methods*, **14**, 587–589. https://doi.org/10.1038/nmeth.4285 (2017).

774 [73] Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBOOT2: Improving the ultrafast bootstrap approximation. *Mol. Biol. Evol.*, **35**, 518–522.
776 https://doi.org/10.1093/molbev/msx281 (2018).

[74] Drummond, A. J., Suchard, M. A., Xie, D. & Rambaut, A. Bayesian phylogenetics with BEAUTI
778 and the BEAST 1.7. *Mol. Biol. Evol.*, **29**, 1969–1973. https://doi.org/10.1093/molbev/mss075 (2012).

780 [75] Rambaut, A. & Drummond, A. J. TRACER V1.4. Molecular evolution, phylogenetics and epidemiology. *Edinburgh: The University of Edinburgh, Institute of Evolutionary Biology*
782 *software.* http://tree.bio.ed.ac.uk/software/tracer/ (2007).

[76] Rambaut, A. FIGTREE v1.4. Molecular evolution, phylogenetics and epidemiology. *Edinburgh:*
784 *The University of Edinburgh, Institute of Evolutionary Biology software.* http://tree.bio.ed.ac.uk/software/figtree/ (2012).

786   [77] Kishino, H., Miyata, T. & Hasegawa, M. Maximum likelihood inference of protein phylogeny and the origin of chloroplasts. *J. Mol. Evol.*, **31**, 151–160.
788   https://doi.org/10.1007/BF02109483 (1990).

    [78] Kishino, H. & Hasegawa, M. Evaluation of the maximum likelihood estimate of the
790   evolutionary tree topologies from DNA sequence data, and the branching order in hominoidea. *J. Mol. Evol.*, **29**, 170–179. https://doi.org/10.1007/BF02100115 (1989).

792   [79] Shimodaira, H. & Hasegawa, M. Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Mol. Biol. Evol.*, **16**, 1114–1116.
794   https://doi.org/10.1093/oxfordjournals.molbev.a026201 (1999).

    [80] Shimodaira, H. An approximately unbiased test of phylogenetic tree selection. *Syst. Biol.*,
796   **51**, 492–508. https://doi.org/10.1080/10635150290069913 (2002).

    [81] Strimmer, K. & Rambaut A. Inferring confidence sets of possibly misspecified gene trees.
798   *Proc. R. Soc. London. Ser. B Biol. Sci.*, **269**, 137–142. https://doi.org/10.1098/rspb.2001.1862 (2002).

800   [82] Tamura, K., Tao, Q. & Kumar, S. Theoretical foundation of the RELTIME method for estimating divergence times from variable evolutionary rates. *Mol. Biol. Evol.*, **35**, 1770–
802   1782. https://doi.org/10.1093/molbev/msy044 (2018).

    [83] Mello, B. Estimating timetrees with MEGA and the TimeTree Resource. *Mol. Biol. Evol.*, **35**,
804   2334–2342. https://doi.org/10.1093/molbev/msy133 (2018).

    [84] Yang, T. *et al*. New schizothoracine from Oligocene of Qaidam Basin, northern Tibetan
806   Plateau, China, and its significance. *J. Vertebr. Paleontol.*, **2017**, e1442840. https://doi.org/10.1080/02724634.2018.1442840 (2018).

808   [85] Guo, J. *et al.* Three-dimensional structural model of the Qaidam Basin: Implications for crustal shortening and growth of the northeast Tibet. *Open Geosci.*, **9**,174–185.
810   https://doi.org/10.1515/geo-2017-0015 (2017).

    [86] Loucks, D.P. & van Beek, E. Water Resource Systems Planning and Management, An
812   Introduction to Methods, Models, and Applications. Deltares, UNESCO-IHE, and Springer-Verlag Open Access. https://doi.org/10.1007/978-3-319-44234-1_1 (Fig. 1.17
814   China, India, and Southeast Asia, highlighting the Tibetan Plateau: Available via license: CC BY-NC 4.0) (2017).

Table 1. Results of an analysis of molecular variance (AMOVA) based on 130 haplotypes of
820   Snowtrout (Cyprinidae: *Schizothorax*) partitioned into: (1) Geographic regions (="Among");
(2) Aggregates within clades (="A/W"); and (3) Haplotypes within aggregates (="Within").
822   Source=Grouping; D.F.=Degrees of freedom; S.S.=sums of squares; Var.=variance; %=Percent
variance. Probability values for each F_{ST}-value (derived via 1000 permutations) are significant
824   at a Bonferroni-adjusted value of $p<0.005$.

Source	D.F.	S.S.	Var.	%
Among	10	3958.7	31.4	77.9
A/W	18	466.1	5.6	13.5
Within	102	350.6	3.4	8.5
Total	130	4775.3	40.3	99.9

830 **Table 2**: Statistical topology tests comparing maximum likelihood (ML) and Bayesian (BA) phylogenetic results (=Tree). logL=Log Likelihood value; deltaL=logL difference from maximal
832 logL in the set; Bp-RELL=Bootstrap proportions using RELL method (weights sum to 1 across trees); p-KH=*p*-value of one-sided Kishino-Hasegawa test; p-SH=*p*-value of Shimodaira-
834 Hasegawa test; c-ELW=Expected Likelihood Weight (weights sum to 1 across trees); p-AU=*p*-value of Approximately Unbiased test. A plus sign (+) next to a *p*-value denote 95% confidence
836 sets, whereas a minus sign (-) denote a significant exclusion (i.e., the tree is rejected). All tests based on 1000 resamplings using the RELL method.

Tree	logL	deltaL	bp-RELL	p-KH	p-SH	c-ELW	p-AU
ML	-9526.83	0.0	0.986 +	0.986 +	1.00 +	0.978 +	0.985 +
BA	-9582.76	55.93	0.014 -	0.013 -	0.013 -	0.014 -	0.015 -

Figure captions:

Figure 1: Simplified drainage map of Asia, showing the major basins from which *Schizothorax* were obtained as GENBANK® accessions. Map available via license: CC BY-NC 4.0, and was modified from Fig. 1.17 [86].

Figure 2: A collapsed maximum likelihood tree reflecting phylogenetic relationships among geographic clades. Data were derived from sequence analysis of the cytochrome-b mitochondrial gene (1140 bp, 139 sequences, redundant haplotypes removed, clades collapsed). Red asterisk = SH-ALRT (Shinodara-Hasegawa approximate likelihood ratio test) ≥ 80% and UFBOOT (ultrafast bootstrap approximations) ≥ 95% (IQ-TREE v.2 Manual: http://www.iqtree.org). Numbers in colored triangles represent species (generalized clades) or individuals (specific clades). Numbers at nodes (N=13) represent RELTIME TIMETREE mean divergence estimates (in Ma; confidence intervals provided in text).

Figure 3: Maximum likelihood tree reflecting phylogenetic relationships among geographic clades. Data were derived from sequence analysis of the cytochrome-b mitochondrial gene (1140 bp, 140 sequences, redundant haplotypes removed). Red asterisk = SH-ALRT (Shinodara-Hasegawa approximate likelihood ratio test) ≥ 80% and UFBOOT (ultrafast bootstrap approximations) ≥ 95% (IQ-TREE v.2 Manual: http://www.iqtree.org). Numbers only at specific nodes represent UFBOOT values. Letters within composite species-triangles at tips of the tree represent species, with R=*S. richardsonii*, P=*S. progastus*, with trailing numbers representing individuals.

Figure 4: Drainage patterns of Nepal (Central Himalaya). Three major river basins merge with the Ganges River in the lowlands of India. Lake Rara (northeast Nepal) drains into the Karnali River. Details on *Schizothorax* localities are provided in Appendix 1.

Figure 5: Drainage patterns of Bhutan (Central Himalaya). Six major rivers drain southward into the Brahmaputra River (northwest India). Samples were obtained from: Amo Chhu, Wang Chhu, Punatsang Chhu/Sunkosh, and Mangde Chhu/Manas. Details on *Schizothorax* localities are provided in Appendix 1.

Figure 6: *Shizothorax sp.* (potentially *S. richardsonii*) captured at the confluence of the Berti Chhu and Mangde Chuu, Bhutan.

Humia Karnali

Rara Lake

Mugu Karnali

Drainage Map
Nepal

Seti

Karnali

Bheri

Kali
Gandaki

Trisuli

Arun

Sun Kosi

Tomur

882 **Supplementary Table S1**: Fish biodiversity evaluated in this study. Listed are: Species Name = Scientific name (A=*Acrossocheilus*; Pe=*Pethia*; Pu=*Puntius*; T=*Tor*; N=*Neolissochilus*;

884 S=*Schizothorax*) (* = Redundant haplotype); Accession# = Tissue or sequence identifier [i.e., GenBank accession number (begins with two letters); University of Kansas tissue voucher (four

886 digits); Douglas Lab DNA code (begins with 'S' or '58')]; Location = Country, geographic region or river; Group = Regions as presented in Figure 2; Group abbreviations are as follows: E-QTP/ SE

888 Asia-A=Eastern Qinghai-Tibetan Plateau/ S.E. Asia, clade A; E-QTP/ SEA-B=Eastern QTP/S.E. Asia, clade B; E-QTP/ SEA-C=Eastern QTP/S.E. Asia, clade C; YLTR=Yarlung-Tsangpo River.

Species Name	Accession #	Location	Group
A. yunnanensis	KC696535	n/a	Outgroup
A. iridescens	KC696536	n/a	Outgroup
Pu. sophore	KF574541	n/a	Outgroup
Pe. ticto	AB238969	n/a	Outgroup
Pe. conchonius	AY004751	n/a	Outgroup
T. putitora	JX204722	India: Beas River (Himachal Pradesh)	Outgroup
T. tor	KF574635	n/a	Outgroup
N. hexagonolepsis	NC026106	India: Jia Bhoreli River at Bhalukpong	Outgroup
S. argentatus	AY954269	China: Ili River (Tarim drainage)	Central Asia
S. biddulphi	FJ931466	China: Kezi River	Central Asia
S. eurystomus	AY954275	China: Tashkurgan River (Tarim)	Central Asia
S. intermedius	AY954273	China: Tashkurgan River (Tarim)	Central Asia
S. intermedius	AY954272	China: Uqturpan	Central Asia
S. pseudoaksaiensis	AY954270	China: Ili River (Tarim drainage)	Central Asia
S. malacanthus	S mal77/AY954277	China: Diantan (Irrawaddy drainage)	Central QTP
S. molesworthi	S mol29/DQ126129	China: Zayu (Tibet)	Central QTP
S. macropogon	S mac17/AY463517	China: Yarlung-Tsangpo River	YLTR_West
S. waltoni_08	MK243412.1	China: Yarlung-Tsangpo River	YLTR_West
S. waltoni_11	MK243415	China: Yarlung-Tsangpo River	YLTR_West
S. waltoni_12	MK243416	China: Yarlung-Tsangpo River	YLTR_West
S. waltoni_13	MK243417	China: Yarlung-Tsangpo River	YLTR_West
S. waltoni_14	MK243418	China: Yarlung-Tsangpo River	YLTR_West
S. waltoni_15	MK243419	China: Yarlung-Tsangpo River	YLTR_West
S. waltoni_16	MK243420	China: Yarlung-Tsangpo River	YLTR_West
S. waltoni_17	MK243421	China: Yarlung-Tsangpo River	YLTR_West
S. wangchiachii	S wan81/HQ198881	China: Yarlung-Tsangpo River	YLTR_West
S. oconnori	S oco10/KT188623	China: Yarlung-Tsangpo River	YLTR_West
S. oconnori	S oco24/KT188637	China: Yarlung-Tsangpo River	YLTR_West
S. oconnori	S oco28/KT188641	China: Yarlung-Tsangpo River	YLTR_West
S. oconnori	S oco50KT188663	China: Yarlung-Tsangpo River	YLTR_West
S. species	58Nyac01	Bhutan: Haa Chhu (Wang Chhu)	Bhutan-1A
S. species	58Haac001	Bhutan: Wang Chhu (Nya Chhu)	Bhutan-1A
S. species	58Dakp06	Bhutan: Dakpai Chhu (Mangde Chhu)	Bhutan-1A

S. species	58Dakp11	Bhutan: Dakpai Chhu (Mangde Chhu)	Bhutan-1A
S. species	58Thun03	Bhutan: Shengarong Chhu (Punatsang Chhu)	Bhutan-1A
S. oconneri	S ocon58/KT188671	China: Yarlung-Tsangpo River	YLTR_East
S. oconneri	S ocon59/KT188672	China: Yarlung-Tsangpo River	YLTR_East
S. oconneri	S ocon60/NC020781	China: Yarlung-Tsangpo River	YLTR_East
S. species	58Bert01	Bhutan: Berti Chhu (Mangde Chhu)	Bhutan-1B
S. species	58Thun008	Bhutan: Shengarong Chhu (Punatsang Chhu)	Bhutan-1B
S. species	58Danr004	Bhutan: Dang Chhu (Punatsang Chhu)	Bhutan-1B
S. species	58Khar001	Bhutan: Khardii Chhu (Dangme Chhu)	Bhutan-1B
S. species	58Pots002	Bhutan: Po Chhu (Punatsang Chhu)	Bhutan-1B
S. species	58Puza001	Bhutan: Zhawaka Chhu (Punatsang Chhu)	Bhutan-1B
S. chongi	S cho18/DQ126118	China: Jiulong River (Yangtze drainage)	E-QTP/ SE Asia-B
S. davidi	S dav13/DQ126113	China: Ya'an (Yangtze drainage)	E-QTP/ SE Asia-B
S. dolichonema	S dol17/DQ126117	China: Jiulong River (Yangtze drainage)	E-QTP/ SE Asia-B
S. dulongensis	S dul84/AY954284	China: Tengchong (Irrawaddy drainage)	E-QTP/ SE Asia-A
S. gongshanensis	S gon79/AY954279	China: Fugong (Salween drainage)	E-QTP/ SE Asia-C
S. gongshanensis	S gon80/AY954280	China: Fugong (Salween drainage)	E-QTP/ SE Asia-C
S. griseus	S gri53/AY954253	China: Yongping (Mekong draiange)	E-QTP/ SE Asia-C
S. kozlovi	S koz12/DQ126112	China: Jiulong River (Yangtze drainage)	E-QTP/ SE Asia-B
S. lantsangensis	S lan82/DQ646882	China: Lancang River (Mekong drainage)	E-QTP/ SE Asia-C
S. lissolabiatus	S lis59/KP796159	China: Nujiang River (Salween drainage)	E-QTP/ SE Asia-C
S. lissolabiatus	S lis67/KP796167	China: Nujiang River (Salween drainage)	E-QTP/ SE Asia-C
S. meridionalis	S mer82/AY954287	China: Yingjiang (Irrawaddy drainage)	E-QTP/ SE Asia-A
S. nukiangensis	Snuk25/DQ126125	China: Chalong (Tibet) (Salween drainage)	E-QTP/ SE Asia-C
S. prenanti	S pre62/AY954262	China: Fenjie (Yangtze drainage)	E-QTP/ SE Asia-B
S. prenanti	S pre61/AY954261	China: Judian (Yangtze drainage)	E-QTP/ SE Asia-B
S. rotundimaxillaris	s rot83/AY954283	China: Tengchong (Irrawaddy drainage)	E-QTP/ SE Asia-A
S. wangchiachii	S wan21/DQ126121	China: Jiulong (Yangtze drainage)	E-QTP/ SE Asia-B
S. yunnanensis	s yun71/KP796171	China: Nujiang River (Salween drainage)	E-QTP/ SE Asia-A
S. yunnanensis	S yun69/KP796169	China: Nujiang River (Salween drainage)	E-QTP/ SE Asia-A
S. yunnanensis	S yun86/ AY954286	China: Yingjiang (Irrawaddy drainage)	E-QTP/ SE Asia-A
S. yunnanensis	S yun52/AY954252	China: Yongping (Mekong drainage)	E-QTP/ SE Asia-C
S. species	58Shen006	Bhutan: Shengarong Chhu (Punatsang Chhu)	Bhutan-2
S. species	58Kame01	Bhutan: Kame Chhu (Punatsang Chhu)	Bhutan-2
S. species	58Dikc03	Bhutan: Dik Chhu (Punatsang Chhu)	Bhutan-2
S. species	58Tink001	Bhutan: Tinku Chhu (Punatsang Chhu)	Bhutan-2
S. species	58Karo003	Bhutan: Kami-Rong Chhu (Punatsang Chhu)	Bhutan-2
S. species	58Puza006	Bhutan: Zhawaka Chhu (Punatsang Chhu)	Bhutan-2
S. species	58Danr001	Bhutan: Dang Chhu (Punatsang Chhu)	Bhutan-2
S. species	58toeb12	Bhutan: Toebrong Chhu (Punatsang Chhu)	Bhutan-2
S. esocinus	S eso50/ KP712250	Nepal: Melamchi River	Koshi River
S. progastus	58_047PKO/1975	Nepal: Tumlingtar Region	Koshi River
S. progastus	58_048PKO/1976	Nepal: Tumlingtar Region	Koshi River

S. progastus	58_049PKO/1977	Nepal: Tumlingtar Region	Koshi River
S. progastus	58_050PKO/1978	Nepal: Tumlingtar Region	Koshi River
S. progastus	58_095PKO/1974	Nepal: Tumlingtar Region	Koshi River
S. richardsonii	S ric49/ KP712249	Nepal: Indrawati River	Koshi River
S. richardsonii	58_042RKO/1964	Nepal: Arun River	Koshi River
S. richardsonii	58_046RKO/1968	Nepal: Arun River	Koshi River
S. richardsonii	58_090RKO/1969	Nepal: Arun River	Koshi River
S. progastus	58_038/1955	Nepal: Kali Gandaki River	Gandaki River
S. progastus	58_039/1956	Nepal: Kali Gandaki River	Gandaki River
S. progastus	58_040/1957	Nepal: Kali Gandaki River	Gandaki River
S. progastus	58_075PGA/8822	Nepal: Rahughat River	Gandaki River
S. progastus	58_076PGA/8824	Nepal: Rahughat River	Gandaki River
S. progastus	58_077PGA/8829	Nepal: Rahughat River	Gandaki River
S. progastus	58_052PGA/1995	Nepal: Rahughat River	Gandaki River
S. progastus	58_072PGA/8814	Nepal: Rahughat River	Gandaki River
S. progastus	58_073PGA/8815	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	58_085RGA/1959	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	58_088RGA/1962	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	58_096RGA/1979	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	58_102RGA/1987	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	58_106RGA1991	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	58_113RGA/8817	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	58_114RGA/8825	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	58_115RGA/8826	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	58_117RGA/8828	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	58_120RGA/8832	Nepal: Kali Gandaki River	Gandaki River
S. richardsonii	ric20 (58_074/8820)	Nepal: Kali Gandaki River	Gandaki River
S. progastus	58_122PKA/1940	Nepal: Jhugala	Karnali River
S. progastus	58_123PKA/1941	Nepal: Jhugala	Karnali River
S. richardsonii	58_020RKA/1935	Nepal: Srikot	Karnali River
S. richardsonii	58_021RKA/1937	Nepal: Srikot	Karnali River
S. richardsonii	58_036RKA/1936	Nepal: Srikot	Karnali River
S. richardsonii	58_053RKA/2003	Nepal: Gumgarh River (Bhotechaur)	Karnali River
S. richardsonii	58_078RKA/1938	Nepal: Srikot	Karnali River
S. richardsonii	58_079RKA/1939	Nepal: Srikot	Karnali River
S. richardsonii	58_082RKA/1947	Nepal: Jhugala	Karnali River
S. richardsonii	58_083RKA/1948	Nepal: Jhugala	Karnali River
S. richardsonii	58_084RKA/1949	Nepal: Jhugala	Karnali River
S. richardsonii	58_109RKA/1999	Nepal: Gumgarh River (Bhotechaur)	Karnali River
S. richardsonii	58_110RKA/2000	Nepal: Gumgarh River (Bhotechaur)	Karnali River
S. richardsonii	58_112RKA/2002	Nepal: Gumgarh River (Bhotechaur)	Karnali River
S. macrophthalmus	>58_019MR/1933	Nepal: Rara Lake	Karnali River
S. macrophthalmus	>58_031MR/1903	Nepal: Rara Lake	Karnali River

S. macrophthalmus	>58_061MR/2011	Nepal: Rara Lake	Karnali River
S. nepalensis	>58_002NR/1793	Nepal: Rara Lake	Karnali River
S. nepalensis	>58_003NR/1794	Nepal: Rara Lake	Karnali River
S. nepalensis	>58_011NR/1906	Nepal: Rara Lake	Karnali River
S. nepalensis	>58_033NR/1917	Nepal: Rara Lake	Karnali River
S. raraensis	>58_054RR/2004	Nepal: Rara Lake	Karnali River
S. raraensis	>58_056RR/2006	Nepal: Rara Lake	Karnali River
S. raraensis	>58_057RR/2007	Nepal: Rara Lake	Karnali River
S. plagiostomus	S plaP1/KR232369	Pakistan: Neelum/Jhelum R.	Ganges River
S. progastus	58_126PKA/1944	Nepal: Jhugala (Karnali R.)	Ganges River
S. richardsonii	S ric01/JX485901	India: Upper Ganges	Ganges River
S. richardsonii	S ric02/JX485902	India: Upper Ganges	Ganges River
S. richardsonii	S ric82/JX485882	India: Upper Ganges	Ganges River
S. richardsonii	S ric83/JX485883	India: Upper Ganges	Ganges River
S. richardsonii	S ric84/JX485884	India: Upper Ganges	Ganges River
S. progastus	S proP1/KX254914	Pakistan: Neelum/ Jhelum R.	Indus River
S. progastus	S proP2/KX254915	Pakistan: Neelum/ Jhelum R.	Indus River
S. richardsonii	S ricKO/KC790369	India: Ratighat (Koshi Tributary)	Indus River
S. esocinus	S eso82/KT210882	Northern Pakistan	Indus River
S. plagiostomus	S plaP2/KT184924	Northern Pakistan	Indus River
S. plagiostomus*	S pla69/KR232369	Pakistan: Jhelum R.	Indus River
S. plagiostomus*	S pla72/KR232372	Pakistan: Jhelum R.	Indus River

Supplementary Figure S1: Phylogenetic tree derived using Bayesian methodology (Bᴇᴀsᴛ v2.61) depicting phylogenetic relationships among Snowtrout (*Schizothora;* Cyprinidae) and outgroup taxa derived from sequence analysis of the cytochrome-b mitochondrial gene (1140 bp, 139 sequences, redundant haplotypes removed). Individuals are coalesced into geographic regions, with abbreviations defined as: E-QTP/ SEA-A (Eastern Qinghai-Tibetan Plateau/ S.E. Asia, clade A); E-QTP/ SEA-B (Eastern QTP/S.E. Asia, clade B); E-QTP/ SEA-C (Eastern QTP/S.E. Asia, clade C); YLTR (Yarlung-Tsangpo River). Numbers at nodes represent Bayesian posterior probabilities. Sample metadata in Table S1.