A high-throughput amplicon sequencing approach for population-wide species diversity

and composition survey

WT Tay[1,†], LN Court[1], S Macfadyen[1], F Jacomb[1], S Vyskočilová[1,2], J Colvin[2], PJ De Barro[3]

1.   CSIRO Black Mountain Laboratories, Clunies Ross Street, ACT 2601, Australia

2.   Natural Resources Institute, University of Greenwich, Central Avenue, Chatham, Maritime Kent ME4

4TB, United Kingdom

3.   CSIRO Brisbane, Queensland 4001, Australia

†   Author for correspondence: **E-mail:** weetek.tay@csiro.au   **Tel:** +61-2-6246 4286

Abstract

Management of agricultural pests requires an understanding of pest species diversity, their interactions

with beneficial insects and spatial-temporal patterns of pest abundance. Invasive and agriculturally important

insect pests can build up very high populations, especially in cropping landscapes. Traditionally, sampling effort

for species identification involves small sample sizes and is labour intensive. Here, we describe a high throughput

sequencing (HTS) PCR amplicon method and associated analytical workflow for a rapid, intensive, high-volume

survey of pest species compositions. We demonstrate our method using the *Bemisia* pest cryptic species

complex as examples. The whiteflies *Bemisia* including the '*tabaci*' species are agriculturally important capable

of vectoring diverse plant viruses that cause diseases and crop losses. We use our HTS amplicon platform to

simultaneously process high volumes of whitefly individuals, with efficiency to detect rare (i.e., 1%) test-species

and beneficial hymenopteran parasitoid species. Field-testing our HTS amplicon method across the Tanzania,

Uganda and Malawi cassava cultivation landscapes, we identified the sub-Saharan Africa 1 *Bemisia* putative

species as the dominant pest species, with other cryptic *Bemisia* species being detected at various abundances.

We also provide evidence that *Bemisia* species compositions can be affected by sampling techniques that target

either nymphs or adults. Our method is particularly suitable to molecular diagnostic surveys of cryptic insect

species with high population densities. Our approach can be adopted to understand species biodiversity across

landscapes, with broad implications for improving trans-boundary biosecurity preparedness, thus contributing

to molecular ecological knowledge and the development of control strategies for high-density, cryptic, pest-

species complexes.

Key Words

Bemisia; African cassava whitefly; molecular diagnostics; population dynamics; Aphelinidae, parasitoids

Introduction

The ecology of a species is linked by definition to its species status. It underpins our understanding of species diversity in conservation genetics, biosecurity preparedness and developing effective pest management strategies. The correct identification of cryptic species can be challenging and often involves the use of molecular approaches. Timely identification and characterisation of pest species will greatly assist the development of effective management strategies to minimise impacts of crop losses due to pests on food security. Furthermore, techniques that enable the rapid screening and accurate identification of large numbers of individuals can accelerate research on a target pest species and lead to important insights into species ecology and biology.

The use of the maternally-inherited mitochondrial cytochrome oxidase subunit I (mtCOI) partial gene in 'DNA barcoding' via the Sanger sequencing method has contributed to molecular diagnostics of species (e.g., Berry et al., 2004; Boykin et al., 2007; Dinsdale, Cook, Riginos, Buckley, & De Barro, 2010; Elfekih, Tay, Gordon, Court, & De Barro, 2018; Tay, Evans, Boykin, & De Barro, 2012), and provided insights into the potential species diversity where cryptic species may co-exist (e.g., Hebert, Cywinska, Ball, & DeWaard, 2003; Rao, Liew, Yow, & Ratnayeke, 2018; Tay, Beckett, & De Barro, 2016). Cryptic species such as the hemipteran whitefly *Bemisia tabaci* and related 'non-*tabaci*' species are excellent examples where putative species identification can be achieved via molecular characterisation of the partial mtCOI gene (e.g., Lee, Park, Lee, Lee, & Akimoto, 2013; Mugerwa et al., 2018), provided that PCR artefacts, such as NUMTs and/or poor quality sequence trace files and/or gene regions, were first removed (e.g., Elfekih et al., 2019; Kunz, Tay, Elfekih, Gordon, & De Barro, 2019; Tay, Elfekih, Court, et al., 2017; Vyskočilová, Tay, van Brunschot, Seal, & Colvin, 2018). Within the *B. tabaci* cryptic species complex, over 40 putative *B. tabaci* cryptic species have been reported (Kunz, Tay, Elfekih, et al., 2019; Lee et al., 2013; Mugerwa et al., 2018; Vyskočilová et al., 2018). The cryptic species status of the *B. tabaci* complex reflected the significant challenges associated with the correct identification of species status (reviewed by Boykin et al., 2018; De Barro, Liu, Boykin, & Dinsdale, 2011; Tay & Gordon, 2019). Species of the *B. tabaci* complex are notorious around the world as effective vectors of viruses that cause significant diseases of crop plants (Polston, De Barro, & Boykin, 2014). In east Africa, *B. tabaci* species transmit diverse viruses that cause diseases in cassava. Cassava mosaic disease (CMD) and cassava brown streak disease (CBSD) have led to significant crop losses and loss of production by smallholder farmers (Legg et al., 2011; Legg et al., 2014).

Understanding the *Bemisia* species present in east Africa, and their patterns of abundance and distribution in

different farming landscapes, is a necessary first step to managing this pest problem (Macfadyen et al., 2018).

In African nations such as Tanzania, Malawi and Uganda, cassava is typically grown by smallholder

farming families, surrounded by non-cassava crops such as tomatoes, sweet potatoes, groundnuts, soybean,

and maize. Native ecosystems that include unmanaged grazing lands, weeds, and shrubs, are also present and

contain plants which are capable of supporting a diversity of whitefly species. Some may be novel species, which

are yet to be characterised by morphological, behavioural and molecular approaches (e.g., Mugerwa et al., 2018;

Vyskocilova, Seal, & Colvin, 2019; Vyskočilová et al., 2018). Globally, cassava is an important carbohydrate source

for over 500 million people (ARC 2014, accessed 21-April-2019), and is planted in Africa, South America, and

Asia/South East Asia. Thirteen of the top 20 global cassava producers are African countries (i.e., Nigeria, Angola,

Ghana, Democratic Republic of Congo, Malawi, United Republic of Tanzania, Cameroon, Mozambique, Benin,

Sierra Leone, Madagascar, Uganda and Rwanda (<www.worldatlas.com> up-dated 25-April-2017; accessed 21-

April-2019)). In other countries such as Thailand, Vietnam and Cambodia, it is grown as a source of high-quality

starch and is used in diverse food products (Delaquis et al., 2018; Gotz & Winter, 2016; Graziosi et al., 2016;

Kumar, Chang, Narayanan, & Ramasamy, 2017). Therefore, having techniques that enable rapid and cost-

effective surveillance to investigate pest species compositions especially in cassava (and other food crops, i.e.,

tomatoes; other cash crops, i.e., cotton, ornamental plants) production landscapes are incredibly valuable.

While identifying cryptic species diversity can best be achieved through molecular characterisation of

appropriate genomic regions, adapting the concept of high-throughput sequencing (HTS) platforms to answering

questions around species diversity and evolutionary interactions can be challenging.

The HTS platform offers a possible solution for surveillance of insect pest species, including cryptic

species that reach high population densities, provided that methods are sufficiently flexible to accommodate

study designs tailored to address targeted research questions. For example, understanding host-parasitoid

relationships, cryptic species diversity, host-plant relationships, and identifying keystone species are

fundamental ecological questions common to diverse ecosystems. Addressing these ecological and evolutionary

questions can be challenging and requires scaling up of the data collection processes. HTS platforms represent

a solution when combined with appropriate species diversity databases, flexible bioinformatics and associated

analytical pipelines, and optimised molecular methods.

The HTS concept aims to combine individuals within each population for species identification, and is

particularly susceptible to sub-optimal primer efficacies (Elfekih, Tay, et al., 2018; Mugerwa et al., 2018;

Shatters, Powell, Boykin, Liansheng, & McKenzie, 2009; Tay, Elfekih, Court, et al., 2017) that are often

encountered especially in the *B. tabaci* cryptic species complex. Partial or inconsistent Polymerase Chain

Reaction (PCR) amplification, especially when targeting co-extracted DNA of individual cryptic species that may

be found together, can lead to inaccurate conclusion of species composition within the surveyed

agroecosystems. For example, a minor species may become over-represented within a population if co-amplified

within a pool of major species but enjoys superior primer efficacies compared with the major species. Also, given

the time, effort and cost necessarily associated with the preparation of HTS DNA libraries, especially where

accurate input genomic DNA concentration is needed, the high-throughput amplicon sequencing method should

incorporate a 'back-up' step. Such step may involve a second set of primers, ensuring the amplification of the

same population, should the first set of primers fail and thereby avoiding a total PCR failure. Such a back-up

design also enables capturing of species that are missed by the first primer pairs, and vice versa. Provided that

the primer pairs developed were sufficiently robust, within a pest insect species complex such as in the whitefly

cryptic species, beneficial insects such as parasitoid species can also be investigated with appropriate sampling

of pest-insect life stages (e.g., whitefly nymphs).

Here, we describe the HTS method and primers developed for surveying the *Bemisia* cassava whitefly,

related *B. tabaci* cryptic species and associated parasitoid species complex, and the workflow for processing the

large volume of HTS amplicon sequence reads generated. Our goal was to develop a technique that enabled us

to determine the dominant *Bemisia* species in any cassava field quantitatively, using a robust sample size in

terms of numbers of individuals. We provide a verification of primer efficacies and both successful and failed

detection rates of the method, and confirmed the method and primer efficacies via Sanger sequencing of

individual field-sampled whiteflies. Our samples were collected from geographically diverse cassava cropping

landscapes in Uganda, Malawi and Tanzania, so test the limitations and usefulness of this approach especially

for countries with significant infrastructure and economic challenges for researching agricultural pests impacting

food security.

Material & Methods

Samples used in PCR primer efficacy tests

The DNA of whitefly cryptic species from Africa, Asia, South America, and Australia were individually extracted using the Qiagen Blood and Tissue DNA extraction kit (Cat #. 69506). Extracted DNA of all samples to be used for testing of PCR primer efficacies were individually eluted in 27.5 µL of EB. Whitefly species status was confirmed by PCR and Sanger sequencing using the primer set wfly-COI-F1/R1 and/or wfly-COI-F2/R2 (Suppl. Material I). Primer development and Sanger sequencing followed the methods of Elfekih, Tay, et al. (2018). Sanger sequencing was carried out at the BRF at JCSMR ANU, Canberra, ACT. Whitefly species used to verify primer efficacies and their respective GenBank accession numbers are listed in Table 1.

Table 1: *Bemisia* species used to confirm PCR primer efficacies. With the exception of MN646951, all GenBank accession numbers provided represent partial mtCOI sequences of *Bemisia* species tested where 100% sequence identity against publicly available sequences have been detected.

Species	GenBank	Field sample identifiers	Country/source
MED-ASL	MH357342[a]	60132	Malawi (Field ID M2-O3-2), crop = sweet potato Lat -12.90243413, long 34.28411598
MEAM1	KY951449[b]		Peru
B. afer	AF418673[c]	60266	Malawi, on ornamental tree cassava in township.
Bemisia sp01	MN646951[d]	30350	Field ID T1-NC1-1, Lantana weed, Tanzania, Lat -2.74227622, Long 33.02190157
SSA1	JQ286457[e]	40011	Tanzania (Field ID T2-F2), crop = cassava Lat -6.522181425, Long 38.92432862
SSA2	AY057141[f]		NRI laboratory culture
SSA3	KM377923[g]		NRI laboratory culture
NW1	MK386668[h]		Barbados
Asia II_1	MF497064[i]		Bangladesh
Asia1	MF497040[j]		Bangladesh
Lipaleyrodes (*Bemisia*)	MN056066[k]		Australia

Note: a: Vyskocilova et al. (2019); **b:** Tay, Elfekih, Court, et al. (2017), **c:** Maruthi et al. (2004); **d:** New *Bemisia* sp. (this study); **e:** Mugerwa et al. (2012); **f:** Legg, French, Rogan, Okao-Okuja, & Brown (2002); **g:** Ghosh S, Gowda M, Bouvaine S (GenBank unpublished); **h:** Kunz, Tay, Court, et al. (2019); **I, j:** Acharya, R and Kee, KY (GenBank unpublished); **k:** Kunz, Tay, Elfekih, et al. (2019).

To ascertain PCR efficacies of the high-throughput amplicon sequencing primers, various hypothetical field scenarios were simulated. These scenarios involved different genomic DNA (gDNA) that belonged to 11 *Bemisia* species (Table 2) with predetermined species status based on the standard partial mtCOI dataset (Kunz, Tay, Elfekih, et al., 2019). These hypothetical scenarios included: (I) a two-species co-occurrence scenario, in which there is a dominant *Bemisia* whitefly species (e.g., SSA1) and a minor species present in different

predetermined proportions (i.e., Table 2: Mixed-Ia (5%), Mixed-Ib (2.5%), and Mixed-Ic (1%)), and (II-IV)

scenarios assuming co-occurrence of multiple mixed *Bemisia* cryptic pest species at different proportions based

on estimates of gDNA concentration. With the exception of the test scenario 'Mixed-III' and 'Mix-IV' which were

each replicated twice, all remaining test scenarios were replicated three times to estimate error rates and

protocol robustness.

Field samples survey methods

Cassava fields in Uganda, Tanzania, and Malawi were surveyed for *Bemisia* cryptic species between

2015-2016. In total, we conducted three data collection trips across a two-year period (trip 1, 1/8/2015 -

26/08/2015, trip 2, 5/4/2016 - 25/4/2016, trip 3, 29/10/2016 - 9/11/2016). In each region, we selected up to 10

cassava fields to survey and sample for nymph and adult *Bemisia* whiteflies (for sampling details see Macfayden

et al. submitted). Briefly, up to *ca*. 50 adults were collected using an aspirator from the top leaves of cassava

plants of known variety. Nymphs were collected by selecting leaves that were lower down on the plant (and so

had visible nymphs in the second to forth instar age range on the underside of the leaf). The top lip of two

centimetre diameter circular plastic vial (5ml volume) was used to cut small leaf discs that contained nymphs.

These discs were placed in the vial filled with 99% ethanol. As a comparison, nymphs from other crops

surrounding the focal cassava field were also collected. All ethanol-preserved samples were stored in -20°C prior

to being processed in the laboratory, where nymphs were dislodged from leaves and placed in fresh Eppendorf

(1.5mL) test tubes in batches of between 20 and 40 individuals per field (note that in some cases, fields sampled

for *Bemisia* contained <20 individuals). For our research question we selected nymphs that had no obvious signs

of parasitism by hymenoptera parasitoids in order to optimize the *B. tabaci* DNA in the sample. We randomly

selected 26 field-collected *Bemisia* populations from Malawi, Uganda and Tanzania for analysis to demonstrate

the efficacies of the HTS mtCOI amplicon approach under field conditions (Table 2).

Table 2: Test scenarios to ascertain primer efficacies and species detection sensitivity by high-throughput sequencing platform. Average ± standard deviation of detection sensitivity for each category is provided based on amplicon reads as generated. 'Others' include amplicons associated with NUMTs, unknown *Bemisia* genomic regions, parasitoids (*Encarsia*, *Eretmocerus*) partial mtCOI gene region, bacteria including secondary symbionts, fungal, and of unknown origins. '0': Species not considered in test scenarios.

Detection sensitivity	Selected known *Bemisia* species											Others†
	SSA1	SSA2	SSA3	*Bemisia*. Sp01 (MN646951)	MEAM1	MED	NW1	Asia1	AsiaII-1	*Bemisia* sp. (MN056066)	*B. afer*	
Mixed-Ia (1:20)	95 (88.9±0.3)	0	0	0	0	5 (6.1±0.7)	0	0	0	0	0	0 (5±0.01)
Mixed-Ib (1:40)	97.5 (94.8±0.3)	0	0	0	0	2.5 (1.6±0.1)	0	0	0	0	0	0 (3.6±0.2)
Mixed-Ic (1:100)	99 (92±0.7)	0	0	0	1 (1.2±0.4)	0	0	0	0	0	0	0 (7.5±1.3)
Mixed-II	0	0	0	0	0	0	25 (11.4±0.7)	25 (12.7±0.9)	25 (28±1)	25 (20.2±1.2)	0	0 (27.9±2.1)
Mixed-III	0	0	0	0	0	0	30 (24.1±0.6)	50 (30.3±0.4)	10 (8.8±0.5)	10 (14±0.4)	0	0 (24.1±1.4)
Mixed-IV	58 (42.5±4.9)	8.8 (12.5±1.8)	8.8 (10.5±0.7)	6 (3±1.8)	6 (7±0.6)	6 (3.1±0.7)	0	0	0	0	6 (2.9±0.6)	0 (18.2±3.1)

Note: *Bemisia* sp01 (MN646951; From Tanzania field code: 30350). Lipaleyrodes (*Bemisia*) from Queensland, Australia (MN056066). † Only the first 10-15 amplicon contigs were searched for sequence identity by blastn (i.e., optimised for somewhat similar sequences) search against the non-redundant (nr) NCBI DNA database.

Table 3: Twenty-six field-collected whitefly samples from Malawi, Uganda and Tanzania analysed by high-throughput amplicon sequencing. Life stages were nymphs (N); and adults (A). Sample size (n). Detected individuals for each species were estimated from proportions of amplicons. *Bemisia* cryptic species detected were: *B. tabaci* Indian Ocean (IO), Mediterranean (MED), *B.* sp. unknown (U), *B. afer* (Baf), *B.* sp. 'Uganda' (Ug1), African cassava whitefly sub-Saharan Africa 1 (SSA1), sub-Saharan Africa 2 (SSA2), *Trialeurodes* sp. (T), *Encarsia* spp. (Enc), *Eretmocerus* spp. (Eret), Bacterial (Bac). Nuclear mitochondrial sequences/PCR artefacts (NUMTs) were detected in all cases. GenBank accession numbers of all novel *Erectmocerus* and *Encarsia* species were MN646915-MN646927 and MN646928-MN646950 respectively, GenBank accession numbers of all novel *Bemisia* and *Trialeurodes* whitefly species detected were MN660053-MN660056 and MN660057-MN660058, respectively. Raw HTS amplicon data can be obtained by request from the corresponding author.

Country	Site ID	Life stage	Host Plants (variety)	lat	long	Species composition	Total reads	Reads Identified (%)
Malawi	60109/60121	N, n= 26	Cassava (Gomani)	-12.639611255	34.17544665	**SSA1:** 12.7, **Baf:** 12.8, **Enc:** Y, **NUMTs:**	43,652	43,383 (99.35%)
	60109/60121	A, n= 7	Cassava (Gomani)	-12.639611255	34.17544665	**SSA1:** 6.4, **Baf:** 0.4, **NUMTs:** 0.01	54,392	54,135 (99.53%)
	60013/60126	N, n=22	Cassava (Gomani)	-13.03129506	34.29746802	**SSA1:** 21.7, **NUMTs:** 0.17	68,562	68,224 (99.51%)
	60013/60126	A, n= 8	Cassava (Gomani)	-13.03129506	34.29746802	**Baf:** 7.84, **NUMTs:** 0.09	44,424	44,166 (99.42%)
	60127/60250	N, n= 20	Cassava (Gomani)	-13.11325369	34.32143017	**SSA1:** 5.1, **Baf:** 14.5, **NUMTs:** 0.07	34,660	32,169 (92.81%)
	60127/60250	A, n= 20	Cassava (Gomani)	-13.11325369	34.32143017	**SSA1:** 15.9, **Baf:** 3.8, **NUMTs:** 0.13, **Bac:** 0.01	62,600	62,193 (99.35%)
	60032/60263	N, n= 18	Cassava (Manyokola)	-14.11054382	33.81826233	**SSA1:** 0.63, **Baf:** 14.7, **U:** 0.08, **NUMTs:** 1.45	43,156	40,341 (93.48%)
	60032/60263	A, n= 15	Cassava (Manyokola)	-14.11054382	33.81826233	**SSA1:** 5.3, **Baf:** 5.9, **U:** 0.56, **NUMTs:** 2.6	72,172	66,784 (92.53%)
	60010/60242	N, n= 30	Cassava (Beatrice)	-12.804086285	34.21788623	**SSAS1:** 18.7, **Baf:** 10.1, **NUMTs:** 0.91, **Enc:** Y, **Eret:** Y	63,506	62,888 (99.03%)
	60037/60265	N, n= 40	Cassava (Manyokola)	-14.1593299559021	33.8348985175844	**SSA1:** 0.6, **Baf:** 38.7, **Enc:** Y, **NUMTs:** 0.16	44,808	44,228 (98.71%)
	60039/60040	N, n= 40	Cassava (Manyokola)	-13.9888087453993	33.6391004272323	**SSA1:** 38.4, **Baf:** 0.7, **Eret:** Y, **T:** 0.2, **NUMTs:** 0.52	36,530	36,207 (99.12%)
Uganda	10109/10230	N, n= 40	Cassava (TME_14)	0.93830798	33.16688609	**SSA1:** 32.8, **SSA2:** 3.4, **Eret+Enc:** Y (0.5), **NUMTs:** 2	49, 601	47,906 (96.58%)
	20018/20019	N, n= 40	Cassava (Omongole)	-0.98750407	31.42085652	**SSA1:** 32.8, **Baf:** 2.5, **Eret+Enc:** Y (4.1), **NUMTs:** 0.3	38,082	37,752 (99.13%)
	10085/10336	N, n= 40	Cassava (TME_14)	0.87832027	33.19462426	**SSA1:** 35.6, **Baf:** 0.13, **Eret+Enc:** Y (2.8), **NUMTs:** 0.96	68,176	67,185 (98.55%)

	43130	N, n= 20	Cassava (TME_14)	2.35231436	32.93393845	**SSA1:** 9.9, **SSA2:** 3.1, **Baf:** 2.4, **Eret:** Y, **NUMTs:** 1.8	7,486	6,531 (87.24%)
	43131	N, n= 40	Cow peaCowpea	2.35246646	32.93482051	**SSA1:** 8.1, **MED:** 21.8, **U:** 1.6, **Eret:** Y, **NUMTs:** 2.1	6,674	5,883 (88.15%)
	53371	N, n= 40	Sweet potato	-0.52519054	31.6441164	**SSA1:** 1, **MED:** 26.2, **IO:** 0.4, **Ug1:** 11, **NUMTs:** 0.14, **Eret:** Y (0.1)	58,646	56,426 (96.21%)
	10362	N, n= 8	Cassava (Nase_14)	0.92017081	33.21595328	**SSA1:** 5.1, **SSA2:** 1.6, **NUMTs:** 0.14	7,305	6,202 (84.90%)
	33026	N, n= 20	other crop, Malakwang Rosella-like	0.97703406	33.04814937	**SSA1:** 0.6, **MED:** 19.2, **Eret:** Y, **NUMTs:** 0.2	54,877	54,726 (99.72%)
	33042	N, n= 20	Cassava (Nase_14)	1.00622674	32.98948722	**SSA1:** 18.4, **MED:** 0.03, **Eret+Enc:** Y (1.3), **NUMTs:** 0.18	47,014	46,812 (99.57%)
Tanzania	30145/30157	N, n= 40	Cassava (busanagulwa)	-2.388214545	33.05174421	**SSA1:** 33, **Eret+Enc:** Y (6.5), **NUMTs:** 0.27	58,826	58,407 (99.29%)
	30052/30061	N, n= 40	Sweet potato in cassava field	-2.41603623925181	33.0826151071912	**SSA1:** 39.3, **Baf:** 0.4, **Enc:** Y, **NUMTs:** 0.3	62,988	62,589 (99.37%)
	40013/40014	N, n= 40	Cassava (Kiroba)	-7.15118632229587	39.1807717492072	**SSA1:** 31.5, **Baf:** 6, **NUMTs:** 1.1	71,134	68,629 (96.48%)
	40279/40288	N, n= 40	Cassava (Kiroba)	-6.58507491183717	38.8436025425142	**SSA1:** 32.4, **Baf:** 1.1, **Eret+Enc:** Y (5.8), **NUMTs:** 0.34	49,610	49,154 (99.08%)
	30359	N, n= 12	Cassava (unknown)	-2.802858945	33.05936783	**SSA1:** 9.5, **U:** 1.2, **NUMTs:** 0.1	6,931	6,219 (89.73%)
	70120	N, n= 20	Cassava (marfaransa)	-6.59705482	38.81119587	**SSA1:** 14.2, **MED:** 0.1, **Eret+Enc:** Y, **NUMTs:** 4.3	6,029	5,912 (98.06%)

DNA extraction and HTS library preparation

DNA extraction of pooled individuals was carried out using the Qiagen Blood and Tissue DNA extraction kit (cat. #69056) and followed manufacturer's instructions. Extracted gDNA from each pooled sample was quantified for DNA concentration and standardised to 0.5 ng/µl prior to PCR amplification of target mtCOI region. Two PCR reactions were carried out for each pooled sample using primers wfly-PCR-F1/R1 and wfly-PCR-F2/R2 for the first and second reaction, respectively. For each reaction, the PCR amplification of each pooled sample uses 3.5ng gDNA template in a 35µL PCR reaction volume and 30 PCR cycles (PCR conditions as described in Suppl. Material I). Amplicons from each of the pooled samples were ascertained for PCR amplification success on 1.25% 1x TAE agarose gel stained with GelRed and visualised on a UV-transilluminator. Amplicons were then purified using AMPure XP beads as instructed in the Illumina **16S Metagenomic Sequencing Library Preparation** (Part # 15044223 Rev. B) protocol and resuspended in 25 µL Qiagen EB (instead of the recommended 52.5µL). Purified pooled amplicons were individually quantified for DNA concentration using a Qubit v2.0 fluorometer (Life Technologies) and Qubit HS dsDNA assay kit and indexed by amplifying 5.0 ng of the purified amplicon in 50µL PCR volume over 11 PCR cycles (see Suppl. Material I). Indexed amplicons were purified using AMPure XP beads as described above and concentration ascertained using Qubit v2.0. Equal proportions of purified indexed amplicon (i.e., 100ng) from each pooled sample were mixed with 6x Loading Dye and run on 1.25% low melt agarose at 75 V for 2 hours. The F1/R1 combined amplicons and the F2/R2 combined amplicons were excised and purified using the Zymoclean Gel DNA Recovery Kit (cat. # D4001, D4002, D4007, D4008) following the manufacturer's protocol. Purified indexed amplicons were quantified for double stranded DNA concentration using Qubit and fragment size was estimated using TapeStation (Agilent Technologies). Amplicons were calculated for their nano-molarity and diluted to 4 nM (see Illumina 470-2016-007-B) for HTS run on MiSeq sequencer using the Illumina MiSeq Reagent Kit Version 3. A detailed protocol for the amplicon library preparation and related primer sequence information is provided as Suppl. Material I.

Reference sequences for identification of *Bemisia* mtCOI

Within the African cassava cultivation landscape context, the proportion of specific *Bemisia* and related cryptic species detected was estimated as represented by the proportion of total amplicon fragments. For this purpose, we included reference sequences from the following *Bemisia* species: SSA1 (GenBank AF344264), SSA2

(GenBank AY057173), SSA3 (GenBank AF344257), MEAM1 (GenBank KY951449), IO (GenBank KY951448), *B. afer*

(GenBank KF734668), Uganda1 (GenBank KX57085), 'African' MEAM2 (GenBank KX570778), MED (GenBank,

KY951447), SSA6 (GenBank AY903561), SSA9 (GenBank KX570854), SSA10 (GenBank KX570837), SSA13 (GenBank

KX570833) and novel sub-Saharan *Bemisia* species (i.e., MN646951; MN646952) detected by Sanger sequencing

from field-surveyed populations (Macfadyen et al., 2019; this study). We confirmed the authenticity of these

reference sequences as free from NUMTs following the methods of Kunz, Tay, Elfekih, et al. (2019). A list of

Bemisia reference sequences used in this study is provided (Suppl. Material II). The reference sequences were

used for mapping assembly in Geneious v11.1.5 using predefined parameters (Fig. 1). Intra-specific uncorrected

pair-wise distance (*p*-dist) have been demonstrated to range from 0 to 2% in *B. tabaci* and 'non-*tabaci*' cryptic

species (Kunz, Tay, Elfekih, et al., 2019); however, this was set at 3% to also enable the detection and grouping

of closely related MED cryptic species complex (e.g., ASL, see Vyskočilová et al. (2018); and the Burkina Faso

'MED' population (see Kunz, Tay, Elfekih, et al., 2019)) that is putatively a novel species, with significant host-

plant range differences from the Mediterranean region MED populations (Vyskocilova et al., 2019).

Parasitoids reference sequences

Eretmocerus and *Encarsia* parasitoid mtCOI reference sequences from the sub-Saharan region

surrounding our field survey sites were built up (MN646919, MN646949) during the course of the analysis

workflow (see Fig. 1). Due to the anticipated significant whitefly hymenopteran parasitoid species diversity in

the sub-Saharan region, we arbitrarily determine a cut-off value of 6% within the Geneious 'Map to Reference'

function. The reference Aphelinidae parasitoid partial mtCOI gene sequences used in this study are provided in

Suppl. Material II.

Fig. 1: Workflow for analysis of MiSeq generated high-throughput sequencing of mtCOI amplicon reads for
estimating proportions of *Bemisia* species in the African cassava cultivation landscape, including assessment of
hymenopteran parasitoid species. Optional steps are indicated in dashed box depending on whether amplicon
libraries were gel-purified (refer to Suppl. Material I) to remove small fragments. Quality of amplicons can also
be improved if desired, by using an Illumina sequence trimming program such as Trimmomatic (Bolger, Lohse, &
Usadel, 2014). The *de-novo* assembly step enables novel *Trialeurodes*-, parasitoid- and bacterial-related
sequences, as well as NUMTs, to be identified.

Results

Primer efficacies: The primers wfly-COI-F1/R1 and/or wfly-COI-F2/R2 were shown to be highly efficient in

successful PCR amplification of diverse representative *Bemisia* species that included Australian species

(Lipaleyrodes (*Bemisia*)), Asian, New World (NW), Mediterranean (MED), Middle East Asian Minor (MEAM1),

sub-Saharan African (SSA) species, and 'non-*tabaci*' species (i.e., *B. afer*). Sanger sequencing of PCR amplicon

from test *Bemisia* cryptic species (Table 1) partial mtCOI confirmed species identity when compared against both

the up-dated standard *B. tabaci* mtCOI reference dataset and the uncorrected pairwise nucleotide distances (*p*-

dist) (see Kunz, Tay, Elfekih, et al. (2019). While Sanger sequencing confirmed sequence integrity such as through

conservation of amino acid residues against published *B. tabaci* cryptic species HTS-derived full mtDNA genomes

(e.g., Kunz, Tay, Court, et al., 2019; Tay, Elfekih, Court, Gordon, & De Barro, 2016; Tay, Elfekih, Court, et al., 2017;

Tay, Elfekih, Polaszek, et al., 2017; Vyskočilová et al., 2018), results from HTS amplicon analyses readily detected

NUMTs such as similar to 'MEAM2' (Delatte et al., 2005; Tay, Elfekih, Court, et al., 2017), as well as partial mtCOI

genes with INDELs. Detection of co-amplified pseudogenes/NUMTs demonstrated the complexity of the *Bemisia*

genome (Chen et al., 2016; Xie et al., 2017; Xie et al., 2018). HTS-amplicon analyses also shed light on the

unexpected co-amplification of non-target whitefly genomic regions, as well as fungal, bacterial symbionts, and

parasitoids genomic regions, and highlighted the complexity of trophic interactions in *Bemisia* whiteflies (e.g.,

Rao et al., 2018; Shamimuzzaman et al., 2019; Tay, Elfekih, Polaszek, et al., 2017).

HTS-amplicon protocol and analysis workflow: Test scenarios for detection and identification of *Bemisia* cryptic

species in predetermined proportions (i.e., 20:1; 40:1; 100:1) showed that the HTS large scale sampling protocol

and the analysis workflow (Fig. 1) confidently identified the dominant as well as minor species in approximate

expected proportions (Table 2). Similarly, in the hypothetical mixed-species composition scenario that included

different proportions of mixed species (i.e., scenarios 'Mixed-II', 'Mixed-III', 'Mixed-IV'; Table 2), the HTS protocol

also successfully detected all test species while the expected proportions were affected by co-amplification

NUMTs/pseudogenes, bacterial, fungal, parasitoid, and non-specific whitefly genomic regions (Table 2). The test

scenarios therefore provided significant insights into potential weaknesses associated with our primers and the

robustness of the developed HTS-amplicon protocol. For example, across various scenarios, detected amplicon

coverage regularly fell below the expected values due to non-specific PCR co-amplification especially of nuclear

genome regions.

Field application of HTS-amplicon method: Applying the HTS amplicon protocol to field-collected samples from

Malawi, Tanzania, and Uganda showed that in the African cassava fields complex *Bemisia* species compositions

existed, although in most cases SSA1 was the dominant species (Macfadyen et al., 2019). A total of eight whitefly

species (i.e., SSA1, SSA2, IO, MED, *B. afer,* Uganda 1, two novel *Bemisia* species), and a novel *Trialeurodes* species,

were detected in different field sites. Multiple novel mtCOI haplotypes for *Eretmocerus* species (MN64915-

MN64927) and *Encarsia* species (MN646929-MN646950) were also detected (Table 3). Parasitoid sequences

were only ever detected in nymphal samples, and this was despite the active exclusion of parasitised nymphal

samples that could be visually identified prior to genomic DNA extraction. Detection of parasitoids could

represent early stages of parasitism where visual identification could be difficult. The *B. tabaci* MED species

complex (i.e., included either MED (e.g., Tay et al., 2012), MED-ASL (e.g., Vyskočilová et al., 2018), and/or MED-

Burkina Faso (e.g., Kunz, Tay, Elfekih, et al., 2019) putative species) was present as the dominant species in

Uganda from three sampling sites (field codes 33026, 43131, 53371; Table 3), and in Malawi, *B. afer* and SSA1

species dominated different field sites (Table 3). Their detection showed significant variability in species

composition within the same field sites depending on the life-stage sampled (i.e., nymphal vs. adults). Field

sample analyses also detected data as NUMTs and bacterial origins in the African populations, similar to our

hypothetical scenarios (Figs. 2, 3), thereby demonstrating the robustness of our HTS work flow and primer-

efficacy tests.

Discussion

This study successfully demonstrated the applicability, transferability and powers of the HTS platform

to investigate species compositions within an agroecological context, using the highly challenging *Bemisia*

whitefly cryptic pest species as a model system. This methodology also enabled concurrent survey of whitefly-

nymph, parasitoid diversity, thereby further contributing to our understanding of trophic interactions between

pests, beneficial insects and plant hosts across highly heterogeneous African agricultural landscapes. Our

approach transforms the molecular analysis procedures that traditionally involved low volume sampling of

biological material, to one that embraces PCR amplicon HTS analyses capable of handling high volume sampled

biological material. This achievement is underpinned by the development of an easy-to-use tailored

bioinformatic analysis workflow, and two sets of highly efficient PCR primers that have the capability to amplify

the mtCOI barcoding region of *Bemisia* hosts, their associated *Eretmocerus* and *Encarsia* parasitoids, and various

fungal and bacterial homologous gene regions.

DNA barcode amplicon HTS approaches have been reported for integrative taxonomy (e.g., Cruaud,

Rasplus, Rodriguez, & Cruaud, 2017; Sigut et al., 2017) and their potential for invasive species surveys reviewed

(Piper et al., 2019). Adoption for landscape ecological investigations including trophic interactions under field

conditions between insect pest hosts and beneficial insects is now also demonstrated to be highly effective by

our study. Determining which insect represents the major species within a diverse agriculture landscape requires

adequate sampling of the insects' populations (Kalyebi et al., 2019; Kalyebi et al., 2018; Macfadyen et al., 2019).

Conclusions based on few individuals sampled at a large number of sites are inherently problematic; there is a

need, therefore, for the development of techniques to scale up data collection and processing of samples, and

to develop associated bioinformatic workflows and analytical pipelines to provide robust findings at economic

scales. For the cassava cultivation landscape in sub-Saharan Africa, determining the dominant pest whitefly

species represents a significant challenge due to difficulties with species identification without the help of

molecular DNA markers (De Barro et al., 2011; Kunz, Tay, Elfekih, et al., 2019). Furthermore, the often-high

population sizes associated with individual fields, poor PCR primer efficacies (Elfekih, Tay, et al., 2018; Mugerwa

et al., 2018; Shatters et al., 2009; Tay, Elfekih, Court, et al., 2017) often resulting in suboptimal PCR amplification,

and the difficulty encountered in sequence quality control (Kunz, Tay, Elfekih, et al., 2019; Tay, Elfekih, Court, et

al., 2017; Vyskočilová et al., 2018), have been persistent problems in *Bemisia* molecular systematics and

diagnostics. In this study, we showed that by applying the DNA barcoding and the species 'genetic gaps' concepts

first demonstrated by Dinsdale et al. (2010; see also Kunz, Tay, Elfekih, et al., 2019), and with careful re-designing

of DNA markers for the HTS platform, it was possible to develop a highly efficient method for large-scale,

effective and economic sampling of these whitefly cryptic species to contribute to their management and control

in challenging agricultural landscape settings. The primers were also shown to be efficient in detecting novel

Encarsia and *Eretmocerus* species of the hymenopteran parasitoids in the African cassava and non-cassava fields,

thereby providing value and saving effort to understand trophic interactions between crop hosts, insect pests

and their potential biological control agents.

The traditional approach (i.e., PCR and Sanger sequencing) using individual whiteflies limits our ability

to ascertain the common pest species within a landscape because: (i) mtCOI species identification method is

laborious when specimens are to be extracted individually for Sanger sequencing, and (ii) primer efficacies are

often poor due to the large nucleotide diversity that exists between *B. tabaci* cryptic species when applying non-

hemipteran 'universal' primers (Elfekih, Tay, et al., 2018). Sampling sufficiently large quantities of *B. tabaci*

cryptic species across diverse agricultural landscape will need to rely on a standardised sampling approach

(Sseruwagi, Sserubombwe, Legg, Ndunguru, & Thresh, 2004; Macfadyen et al., 2019), followed by species

identification via molecular characterisation of the mtCOI gene. The approach developed here is especially useful

for collecting nymphal data that has a strong link to host plant use, and can be further developed to examine

parasitism rates of *Bemisia* species. However, standardized sampling techniques for nymphs are less well-

developed than for adult *B. tabaci* on cassava (which are considered easier to count and collect). In reality, the

focus of the research question should drive the sampling approach selected (be that nymphs, parasitized

nymphs, or adults); however, in all cases scaling up the numbers of individuals processed is critical. An important

distinction of sampling nymphal *cf*. adult individuals could be seen in the Malawi samples (Table 3), where our

HTS species diagnostic method detected significant differences in species compositions, especially between *B.*

afer and SSA1. Factors underpinning the detected species composition differences could involve ecological and

biological factors despite being from the same sampling sites.

While our method enables efficient ascertainment of species composition at the landscape level, this

method should not be used to determine individual maternal lineages of mtDNA haplotypes due to the high

sensitivity of detecting random PCR-introduced SNPs in individual amplicon molecules. At the landscape survey

level, ascertaining specific maternal lineages does not represent a criterion needed to enable species identity to

be determined, provided that sequence quality was adequately managed to avoid misidentification of species

status due to mtDNA pseudogenes (Elfekih, Etter, et al., 2018; Kunz, Tay, Elfekih, et al., 2019; Tay, Elfekih, Court,

et al., 2017; Vyskočilová et al., 2018). Elfekih et al. (2019) have shown that NUMTs and pseudogenes could lead

to significant misinterpretation of *Bemisia* population structure in the African cassava landscape regardless of

whether partial mtCOI or genome-wide SNPs were used (see e.g., Wosula, Chen, Fei, & Legg, 2017). A significant

limitation on our approach is the quality of the library used to interrogate the sequences generated. Sanger

sequencing-generated partial mtCOI gene sequences in the public database (e.g., NCBI; see Vyskočilová et al.

(2018) regarding MED-related pseudogenes/NUMTs; see also Kunz, Tay, Elfekih, et al. (2019) for reanalysis of *B.*

tabaci DNA database (Boykin, Savill, & De Barro., 2017)) are gradually being recognised for containing high

numbers of problematic sequences (see e.g., Nacer & do Amaral (2017); Song, Buhay, Whiting, & Crandall (2008);

Triant & Dewoody (2007)). Such readily available public DNA global databases will need to be critically reviewed

and modified, given that a large scientific community relies on these sequences to assist with species

identification.

With advances in HTS to generate genome-wide SNPs data and genome resequencing, the efficacy of

the partial mtCOI gene as a molecular marker for species diagnostics has come under scrutiny. The inability of

the mtCOI gene to effectively differentiate African *Bemisia* species would substantially reduce the efficiency of

our HTS amplicon species identification method. Conflicting genetic signatures of African *Bemisia* SSA1, SSA2 and

SSA4 species status based on partial mtCOI and from genome-wide SNP patterns have been reported (Wosula et

al., 2017). Reanalysis of the Wosula et al. (2017) mtCOI and nextRAD data by Elfekih et al. (2019) identified sub-

optimal clean-up of SNP data and inclusion of NUMTs, as main factors that underpinned species status

discrepancies between mtCOI and genome-wide SNP markers. Removal of poor quality nextRAD data, NUMTs

and pseudogenes revealed that the SSA1 and SSA2 species status were in total agreement between the mtCOI

and nuclear markers. In addition, genome-wide SNP data provided clues as to the potential presence of sub-

species status/population substructure within the SSA1 (i.e., SSA1-NW and SSA1-SE, see Elfekih et al., 2019), and

showed SSA4 to be a misidentified species due to NUMTs, with similar genome-wide SNP signature of MEAM2

from MEAM1 (Elfekih, Etter, et al., 2018). The study of Elfekih et al. (2019), therefore, provided strong support

for the validity of our DNA-barcode HTS methods for landscape species survey purposes, including for the highly

challenging African cassava whitefly complex.

Our HTS amplicon sequencing method can be adapted to understand other pest species complexes in

agricultural systems and may be especially effective for invasive insect pests with small body size and where

population sizes can build up in large numbers, such as in invasive whiteflies, thrips, mites, and aphids. We

propose, therefore, to apply the high-throughput amplicon sequencing approach to: (1) greatly increase number

of individuals that can be sampled, processed and characterised simultaneously for their 'DNA barcoding' gene

including the mtCOI, 16s rRNA, 18s rRNA and 28s rRNA genes, (2) estimate individual species proportion across

large agricultural landscapes, (3) in nymph samples, identify parasitoid species that target the agricultural pests

of interest. While the high-throughput amplicon sequencing method in a species composition study in

agroecological landscapes represents a relatively new field of research, the use of this method to identify genes

of biosecurity importance has been demonstrated for understanding the spread of pyrethroid resistance genes

in the red-legged earth mites *Halotydeus destructor* (Edwards et al., 2018). The method described by Edwards et

al. (2018) can be incorporated into our high-throughput amplicon sequencing platform for rapid species

composition identification and resistance allele frequency estimates. This would enable a genomic approach that

would simultaneously allow the identification of species composition at landscape scales, the survey of

insecticide resistance profiles (e.g., Hopkinson et al., 2019), and assessment of the implication of chemical control

strategies on diversity and density of beneficial insects such as parasitoids. Such study would be especially

informative when temporal scales are included within the study designs.

Developmental stages sampled do impact *Bemisia* species compositions and the conclusions drawn

from a survey, as demonstrated in the Malawi cassava sampling sites for adult and nymphal *Bemisia* populations.

The extent of such variability in target species composition between immature and adult life stages for many

other biological systems can also benefit from the HTS method described here. In this study, *Bemisia* nymphs

that visibly showed signs of being parasitised by hymenopteran parasitoids were actively excluded prior to

genomic processing. Despite this, *Encarsia* and *Eretmocerus* parasitoid mtCOI sequences were detected, thereby

highlighting the importance of biological control of *Bemisia* pest species by parasitoids in the sub-Saharan African

agricultural system. While we have not optimised this workflow for addressing host-parasitoid interaction

research questions, we have demonstrated the power of HTS genomic approaches to document and facilitate an

understanding of host-parasitoid interactions. Improved understanding of the ecological impact of pest species

systems from genomic perspectives will further transform our understanding of sustainable agricultural

production. Appropriate experimental designs including careful development of sampling procedures will be

needed for meaningful interpretation that takes advantage of the large volume of genomic data generated by

the HTS platform. Our study demonstrates the use of HTS amplicon methods successfully in classical field-

ecological studies and for novel cryptic species discovery. The adoption of this approach more widely has the

potential to transform global biosecurity preparedness and thereby to impact human health and food security.

Acknowledgements

Laboratory colonies of whitefly samples (*Bemisia* SSA1, SSA2, SSA3) were provided by the NRI, University

of Greenwich, UK. J. Ryu (CSIRO) assisted with sample sorting, C. Paull and A. Hulthen (CSIRO) assisted with field

surveys of African cassava whitefly samples. We thank Dr Andy Polaszek (NHM, UK) for discussion on the

Encarsisa and *Eretmocerus* parasitoids detected in the *Bemisia* hosts. This work was supported by the Natural

Resources Institute, University of Greenwich from a grant provided by the Bill & Melinda Gates foundation (Grant

Agreement OPP1058938). We would like to thank the broader African cassava whitefly project team for their

enthusiasm and support throughout this study.

Data Availability Statement

The data that support the findings of this study are available as Supplemental Material II, and from GenBank (GenBank accession numbers cited throughout). Raw data can be obtained from the corresponding author upon reasonable request.

References

Berry, S. D., Fondong, V. N., Rey, C., Rogan, D., Fauquet, C. M., & Brown, J. K. (2004). Molecular evidence for five distinct *Bemisia tabaci* (Homoptera : Aleyrodidae) geographic haplotypes associated with cassava plants in sub-Saharan Africa. *Annals of the Entomological Society of America, 97*(4), 852-859. doi:Doi 10.1603/0013-8746(2004)097[0852:Meffdb]2.0.Co;2

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics, 30*(15), 2114-2120. doi:10.1093/bioinformatics/btu170

Boykin, L. M., Kinene, T., Wainaina, J. M., Savill, A., Seal, S., Mugerwa, H., . . . Sseruwagi, P. (2018). Review and guide to a future naming system of African *Bemisia tabaci* species. *Systematic Entomology, 43*(3), 427-433. doi:10.1111/syen.12294

Boykin, L. M., Savill, A., & De Barro, P. (2017). Updated mtCOI reference dataset for the *Bemisia tabaci* species complex *F1000Res, 6*, 1835. doi:10.12688/f1000research.12858.1

Boykin, L. M., Shatters, R. G., Jr., Rosell, R. C., McKenzie, C. L., Bagnall, R. A., De Barro, P., & Frohlich, D. R. (2007). Global relationships of *Bemisia tabaci* (Hemiptera: Aleyrodidae) revealed using Bayesian analysis of mitochondrial COI DNA sequences. *Mol Phylogenet Evol, 44*(3), 1306-1319. doi:10.1016/j.ympev.2007.04.020

Chen, W. B., Hasegawa, D. K., Kaur, N., Kliot, A., Pinheiro, P. V., Luan, J. B., . . . Fei, Z. J. (2016). The draft genome of whitefly *Bemisia tabaci* MEAM1, a global crop pest, provides novel insights into virus transmission, host adaptation, and insecticide resistance. *BMC Biology, 14*. doi:ARTN 110 doi:10.1186/s12915-016-0321-y

Cruaud, P., Rasplus, J. Y., Rodriguez, L. J., & Cruaud, A. (2017). High-throughput sequencing of multiple amplicons for barcoding and integrative taxonomy. *Sci Rep, 7*, 41948. doi:10.1038/srep41948

De Barro, P. J., Liu, S. S., Boykin, L. M., & Dinsdale, A. B. (2011). *Bemisia tabaci*: a statement of species status. *Annu Rev Entomol, 56*, 1-19. doi:10.1146/annurev-ento-112408-085504

Delaquis, E., Andersen, K. F., Minato, N., Cu, T. T. L., Karssenberg, M. E., Sok, S., . . . de Haan, S. (2018). Raising the Stakes: Cassava Seed Networks at Multiple Scales in Cambodia and Vietnam. . *Frontiers in Sustainable Food Systems, 2*, 73. doi:https://doi.org/10.3389/fsufs.2018.00073

Delatte, H., Reynaud, B., Granier, M., Thornary, L., Lett, J. M., Goldbach, R., & Peterschmitt, M. (2005). A new silverleaf-inducing biotype Ms of *Bemisia tabaci* (Hemiptera : Aleyrodidae) indigenous to the islands of the south-west Indian Ocean. *Bulletin of Entomological Research, 95*(1), 29-35. doi:10.1079/Ber2004337

Dinsdale, A., Cook, L., Riginos, C., Buckley, Y. M., & De Barro, P. (2010). Refined Global Analysis of *Bemisia tabaci* (Hemiptera: Sternorrhyncha: Aleyrodoidea: Aleyrodidae) Mitochondrial Cytochrome Oxidase 1 to Identify Species Level Genetic Boundaries. *Annals of the Entomological Society of America, 103*(2), 196-208. doi:10.1603/An09061

Edwards, O. R., Walsh, T. K., Metcalfe, S., Tay, W. T., Hoffmann, A. A., Mangano, P., . . . Umina, P. A. (2018). A genomic approach to identify and monitor a novel pyrethroid resistance mutation in the redlegged earth mite, *Halotydeus destructor*. *Pestic Biochem Physiol, 144*, 83-90. doi:10.1016/j.pestbp.2017.12.002

Elfekih, S., Etter, P., Tay, W. T., Fumagalli, M., Gordon, K., Johnson, E., & De Barro, P. (2018). Genome-wide analyses of the *Bemisia tabaci* species complex reveal contrasting patterns of admixture and complex demographic histories. *PLoS ONE, 13*(1), e0190555. doi:10.1371/journal.pone.0190555

Elfekih, S., Tay, W. T., Gordon, K., Court, L. N., & De Barro, P. J. (2018). Standardized molecular diagnostic tool for
the identification of cryptic species within the *Bemisia tabaci* complex. *Pest Manag Sci, 74*(1), 170-173.
doi:10.1002/ps.4676
Elfekih, S., Tay, W. T., Polaszek, A., Gordon, K. H. J., Kunz, D., Macfadyen, S., . . . De Barro, P. J. (2019). On species
delimitation, hybridization and population structure of cassava whitefly in Africa. *Genome Bio Evol, Subitted*
doi: https://doi.org/10.1101/836072.
Gotz, M., & Winter, S. (2016). Diversity of *Bemisia tabaci* in Thailand and Vietnam and indications of species
replacement. *Journal of Asia-Pacific Entomology, 19*(2), 537-543. doi:10.1016/j.aspen.2016.04.017
Graziosi, I., Minato, N., Alvarez, E., Ngo, D. T., Hoat, T. X., Aye, T. M., . . . Wyckhuys, K. A. (2016). Emerging pests
and diseases of South-east Asian cassava: a comprehensive evaluation of geographic priorities,
management options and research needs. *Pest Manag Sci, 72*(6), 1071-1089. doi:10.1002/ps.4250
Hebert, P. D. N., Cywinska, A., Ball, S. L., & DeWaard, J. R. (2003). Biological identifications through DNA barcodes.
Proceedings of the Royal Society B-Biological Sciences, 270(1512), 313-321. doi:10.1098/rspb.2002.2218
Hopkinson, J., Pumpa, S., van Brunschot, S., Fang, G., Frese, M., Tay, W. T., & Walsh, T. (2019). Insecticide
resistance status of *Bemisia tabaci* MEAM1 (Hemiptera: Aleyrodidae) in Australia cotton production valleys.
Austral Entomology, In press.
Kalyebi, A., Macfadyen, S., Hulthen, A., Ocitti, P., Jacomb, F., Tay, W. T., . . . De Barro, P. (2019). Within-season
changes in land use impact pest abundance in smallholder African cassava production systems. *PLoS ONE,*
Submitted.
Kalyebi, A., Macfadyen, S., Parry, H., Tay, W. T., De Barro, P., & Colvin, J. (2018). African cassava whitefly, *Bemisia*
tabaci, cassava colonization preferences and control implications. *PLoS ONE, 13*(10), e0204862.
doi:10.1371/journal.pone.0204862
Kumar, N. R., Chang, J. C., Narayanan, M. B., & Ramasamy, S. (2017). Phylogeographical structure in
mitochondrial DNA of whitefly, *Bemisia tabaci* Gennadius (Hemiptera: Aleyrodidae) in southern India and
Southeast Asia. *Mitochondrial DNA A DNA Mapp Seq Anal, 28*(5), 621-631.
doi:10.3109/24701394.2016.1160075
Kunz, D., Tay, W. T., Court, L. N., Elfekih, S., Gordon, K. H. J., Evans, G. A., & De Barro, P. J. (2019). Draft
mitochondrial DNA genome of a 1920 Barbados cryptic *Bemisia tabaci* 'New World' species (Hemiptera:
Aleyrodidae). *Mitochondrial DNA Part B-Resources, 4*(1), 1183-1184. doi:10.1080/23802359.2019.1591197
Kunz, D., Tay, W. T., Elfekih, S., Gordon, K. H. J., & De Barro, P. J. (2019). Take out the rubbish - Removing NUMTs
and pseudogenes from the *Bemisia tabaci* cryptic species mtCOI database. *Sci Rep, Submitted*.
doi:https://doi.org/10.1101/724765
Lee, W., Park, J., Lee, G. S., Lee, S., & Akimoto, S. (2013). Taxonomic status of the *Bemisia tabaci* complex
(Hemiptera: Aleyrodidae) and reassessment of the number of its constituent species. *PLoS ONE, 8*(5),
e63817. doi:10.1371/journal.pone.0063817
Legg, J. P., French, R., Rogan, D., Okao-Okuja, G., & Brown, J. K. (2002). A distinct *Bemisia tabaci* (Gennadius)
(Hemiptera : Sternorrhyncha : Aleyrodidae) genotype cluster is associated with the epidemic of severe
cassava mosaic virus disease in Uganda. *Molecular Ecology, 11*(7), 1219-1229. doi:DOI 10.1046/j.1365-
294X.2002.01514.x
Legg, J. P., Jeremiah, S. C., Obiero, H. M., Maruthi, M. N., Ndyetabula, I., Okao-Okuja, G., . . . Lava Kumar, P.
(2011). Comparing the regional epidemiology of the cassava mosaic and cassava brown streak virus
pandemics in Africa. *Virus Res, 159*(2), 161-170. doi:10.1016/j.virusres.2011.04.018
Legg, J. P., Shirima, R., Tajebe, L. S., Guastella, D., Boniface, S., Jeremiah, S., . . . Rapisarda, C. (2014). Biology and
management of *Bemisia* whitefly vectors of cassava virus pandemics in Africa. *Pest Manag Sci, 70*(10), 1446-
1453. doi:10.1002/ps.3793
Macfadyen, S., Paull, C., Boykin, L. M., De Barro, P., Maruthi, M. N., Otim, M., . . . Omongo, C. A. (2018). Cassava
whitefly, *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) in East African farming landscapes: a review
of the factors determining abundance. *Bull Entomol Res, 108*(5), 565-582.
doi:10.1017/S0007485318000032
Macfadyen, S., Tay, W. T., Hulthen, A., Paull, C., Kalyebi, A., Jacomb, F., . . . De Barro, P. (2019). Landscape factors
and how they influence whitefly pests in cassava fields across East Africa. *Landscape Ecology, Submitted*.
Maruthi, M. N., Colvin, J., Thwaites, R. M., Banks, G. K., Gibson, G., & Seal, S. E. (2004). Reproductive
incompatibility and cytochrome oxidase I gene sequence variability amongst host-adapted and
geographically separate *Bemisia tabaci* populations (Hemiptera : Aleyrodidae). *Systematic Entomology,*
29(4), 560-568. doi:DOI 10.1111/j.0307-6970.2004.00272.x

Mugerwa, H., Rey, M. E., Alicai, T., Ateka, E., Atuncha, H., Ndunguru, J., & Sseruwagi, P. (2012). Genetic diversity and geographic distribution of *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) genotypes associated with cassava in East Africa. *Ecol Evol, 2*(11), 2749-2762. doi:10.1002/ece3.379

Mugerwa, H., Seal, S., Wang, H. L., Patel, M. V., Kabaalu, R., Omongo, C. A., . . . Colvin, J. (2018). African ancestry of New World, *Bemisia tabaci*-whitefly species. *Sci Rep, 8*(1), 2734. doi:10.1038/s41598-018-20956-3

Nacer, D. F., & do Amaral, F. R. (2017). Striking pseudogenization in avian phylogenetics: Numts are large and common in falcons. *Molecular Phylogenetics and Evolution, 115*, 1-6. doi:10.1016/j.ympev.2017.07.002

Piper, A. M., Batovska, J., Cogan, N. O. I., Weiss, J., Cunningham, J. P., Rodoni, B. C., & Blacket, M. J. (2019). Prospects and challenges of implementing DNA metabarcoding for high-throughput insect surveillance. *Gigascience, 8*(8). doi:10.1093/gigascience/giz092

Polston, J. E., De Barro, P., & Boykin, L. M. (2014). Transmission specificities of plant viruses with the newly identified species of the *Bemisia tabaci* species complex. *Pest Management Science, 70*(10), 1547-1552. doi:10.1002/ps.3738

Rao, R. S., Liew, T. S., Yow, Y. Y., & Ratnayeke, S. (2018). Cryptic diversity: Two morphologically similar species of invasive apple snail in Peninsular Malaysia. *PLoS ONE, 13*(5), e0196582. doi:10.1371/journal.pone.0196582

Shamimuzzaman, M., Hasegawa, D. K., Chen, W., Simmons, A. M., Fei, Z., & Ling, K. S. (2019). Genome-wide profiling of piRNAs in the whitefly *Bemisia tabaci* reveals cluster distribution and association with begomovirus transmission. *PLoS ONE, 14*(3), e0213149. doi:10.1371/journal.pone.0213149

Shatters, R. G., Jr., Powell, C. A., Boykin, L. M., Liansheng, H., & McKenzie, C. L. (2009). Improved DNA barcoding method for *Bemisia tabaci* and related Aleyrodidae: development of universal and *Bemisia tabaci* biotype-specific mitochondrial cytochrome c oxidase I polymerase chain reaction primers. *J Econ Entomol, 102*(2), 750-758. doi:10.1603/029.102.0236

Sigut, M., Kostovcik, M., Sigutova, H., Hulcr, J., Drozd, P., & Hrcek, J. (2017). Performance of DNA metabarcoding, standard barcoding, and morphological approach in the identification of host-parasitoid interactions. *PLoS ONE, 12*(12), e0187803. doi:10.1371/journal.pone.0187803

Song, H., Buhay, J. E., Whiting, M. F., & Crandall, K. A. (2008). Many species in one: DNA barcoding overestimates the number of species when nuclear mitochondrial pseudogenes are coamplified. *Proceedings of the National Academy of Sciences of the United States of America, 105*(36), 13486-13491. doi:10.1073/pnas.0803076105

Sseruwagi, P., Sserubombwe, W. S., Legg, J. P., Ndunguru, J., & Thresh, J. M. (2004). Methods of surveying the incidence and severity of cassava mosaic disease and whitefly vector populations on cassava in Africa: a review. *Virus Res, 100*(1), 129-142. doi:10.1016/j.virusres.2003.12.021

Tay, W. T., Beckett, S. J., & De Barro, P. J. (2016). Phosphine resistance in Australian *Cryptolestes* species (Coleoptera: Laemophloeidae): perspectives from mitochondrial DNA cytochrome oxidase I analysis. *Pest Manag Sci, 72*(6), 1250-1259. doi:10.1002/ps.3805

Tay, W. T., Elfekih, S., Court, L., Gordon, K. H., & De Barro, P. J. (2016). Complete mitochondrial DNA genome of *Bemisia tabaci* cryptic pest species complex Asia I (Hemiptera: Aleyrodidae). *Mitochondrial DNA Part A, 27*(2), 972-973. doi:10.3109/19401736.2014.926511

Tay, W. T., Elfekih, S., Court, L. N., Gordon, K. H. J., Delatte, H., & De Barro, P. J. (2017). The Trouble with MEAM2: Implications of Pseudogenes on Species Delimitation in the Globally Invasive *Bemisia tabaci* (Hemiptera: Aleyrodidae) Cryptic Species Complex. *Genome Biology and Evolution, 9*(10), 2732-2738. doi:10.1093/gbe/evx173

Tay, W. T., Elfekih, S., Polaszek, A., Court, L. N., Evans, G. A., Gordon, K. H. J., & De Barro, P. J. (2017). Novel molecular approach to define pest species status and tritrophic interactions from historical *Bemisia* specimens. *Scientific Reports, 7*. doi:ARTN 429 doi:10.1038/s41598-017-00528-7

Tay, W. T., Evans, G. A., Boykin, L. M., & De Barro, P. J. (2012). Will the Real *Bemisia tabaci* Please Stand Up? *PLoS ONE, 7*(11). doi:ARTN e50550 doi:10.1371/journal.pone.0050550

Tay, W. T., & Gordon, K. H. J. (2019). Going global - genomic insights into insect invasions. *Curr Opin Insect Sci, 31*, 123-130. doi:10.1016/j.cois.2018.12.002

Tay, W. T., Court, L. N., Macfayden, S., Jacomb, F., Vyskocilova, S., Colvin, J., & De Barro, J. P. (2019). A high-throughput amplicon sequencing approach for population-wide species diversity and composition survey. Supplemental Material II. *Mol Ecol Res, Submitted.*

Triant, D. A., & Dewoody, J. A. (2007). The occurrence, detection, and avoidance of mitochondrial DNA translocations in mammalian systematics and phylogeography. *Journal of Mammalogy, 88*(4), 908-920. doi:Doi 10.1644/06-Mamm-a-204r1.1

Vyskocilova, S., Seal, S., & Colvin, J. (2019). Relative polyphagy of "Mediterranean" cryptic *Bemisia tabaci* whitefly
species and global pest status implications. *Journal of Pest Science, 92*(3), 1071-1088. doi:10.1007/s10340-
019-01113-9
Vyskočilová, S., Tay, W. T., van Brunschot, S., Seal, S., & Colvin, J. (2018). An integrative approach to discovering
cryptic species within the *Bemisia tabaci* whitefly species complex. *Sci Rep, 8*(1), 10886.
doi:10.1038/s41598-018-29305-w
Wosula, E. N., Chen, W. B., Fei, Z. J., & Legg, J. P. (2017). Unravelling the Genetic Diversity among Cassava *Bemisia*
tabaci Whiteflies Using NextRAD Sequencing. *Genome Biology and Evolution, 9*(11), 2958-2973.
doi:10.1093/gbe/evx219
Xie, W., Chen, C., Yang, Z., Guo, L., Yang, X., Wang, D., . . . Zhang, Y. (2017). Genome sequencing of the
sweetpotato whitefly *Bemisia tabaci* MED/Q. *Gigascience, 6*(5), 1-7. doi:10.1093/gigascience/gix018
Xie, W., Yang, X., Chen, C., Yang, Z., Guo, L., Wang, D., . . . Zhang, Y. (2018). The invasive MED/Q *Bemisia tabaci*
genome: a tale of gene loss and gene gain. *BMC Genomics, 19*(1), 68. doi:10.1186/s12864-018-4448-9
398