A class of synaptic signaling molecules required for homeostatic potentiation also tunes homeostatic depression

Catherine J. Yeates[1,2,4], C. Andrew Frank[1,2,3]

[1] Department of Anatomy and Cell Biology, University of Iowa Carver College of Medicine, Iowa
City, IA, USA

[2] Interdisciplinary Graduate Program in Neuroscience, University of Iowa, Iowa City, IA, USA

[3] Iowa Neuroscience Institute, University of Iowa Carver College of Medicine, Iowa City, IA, USA

[4] Present Address: Department of Biology, University of Dayton, Dayton, OH, USA

* Correspondence:
C. Andrew Frank
andy-frank@uiowa.edu

Keywords: synapse, homeostasis, depression, *Drosophila melanogaster*, NMJ, neurotransmis-
sion, plasticity

Abstract

Synapses and circuits rely on homeostatic forms of regulation in order to transmit meaningful infor-
mation. The *Drosophila melanogaster* neuromuscular junction (NMJ) is a well-studied synapse that
shows robust homeostatic control of function. Most prior studies of homeostatic plasticity at the NMJ
have centered on presynaptic homeostatic potentiation (PHP). PHP happens when postsynaptic muscle
neurotransmitter receptors are impaired, triggering retrograde signaling that causes an increase in pre-
synaptic neurotransmitter release. As a result, normal levels of evoked excitation are maintained. The
counterpart to PHP at the NMJ is presynaptic homeostatic depression (PHD). Overexpression of the
Drosophila vesicular glutamate transporter (VGlut) causes an increase in the amplitude of spontaneous
events. PHD happens when the synapse responds to the challenge by decreasing quantal content during
evoked neurotransmission – again, resulting in normal levels of postsynaptic excitation.

We hypothesized that there may exist a class of molecules that affects both PHP and PHD. Impairment
of any such molecule could hurt a synapse's ability to respond to any significant homeostatic challenge.
We conducted an electrophysiology-based screen for blocks of PHD. While we did not observe a block
of PHD in the genetic conditions screened, we instead found loss-of-function conditions that led to
excess depression – i.e., a substantial deficit in evoked amplitude when combined with VGlut overex-
pression. The conditions causing this phenotype included a double heterozygous loss-of-function con-
dition for genes encoding the inositol trisphosphate receptor (IP$_3$R – *itpr*) and ryanodine receptor (*RyR*).
IP$_3$Rs and RyRs gate calcium release from intracellular stores. Pharmacological agents targeting IP$_3$R
and RyR recapitulated the genetic losses of these factors, as did lowering calcium levels from other
sources. Our data are consistent with the idea that some factors required for homeostatic potentiation
are also required for the synapse to achieve appropriate levels of homeostatic depression. Loss of such
factors may disorient compensatory plasticity signals.

1   Introduction

Animal nervous systems use forms homeostatic synaptic plasticity to maintain stable function. Over
the last 20-25 years, studies from diverse systems have revealed a wealth of information about how
forms of homeostatic synaptic plasticity are implemented (Davis, 2013; Davis and Müller, 2015;
Delvendahl and Müller, 2019; Marder and Goaillard, 2006; Pozo and Goda, 2010; Turrigiano, 2008).
In particular, the *Drosophila melanogaster* neuromuscular junction (NMJ) has uncovered many facets
of homeostatic implementation on a molecular level (Frank, 2014a; Frank et al., 2020). Much of the
NMJ homeostasis work in both Drosophila and vertebrates has focused on a form of homeostatic plas-
ticity termed presynaptic homeostatic potentiation (PHP). With PHP, manipulations that impair
postsynaptic muscle receptor function trigger an increase in presynaptic vesicle release (Cull-Candy et
al., 1980; Davis et al., 1998; Frank et al., 2006; Petersen et al., 1997; Wang et al., 2016).

Homeostatic plasticity the NMJ is a bi-directional process. First, PHP is reversible – when manipula-
tions that impair muscle receptor function are removed, the presynaptic potentiation ceases (Wang et
al., 2016; Yeates et al., 2017). Second, the Drosophila NMJ can depress quantal content in a homeo-
static manner functionally opposite to PHP: Presynaptic homeostatic depression (PHD) happens when
there is a decrease in quantal content in response to a perturbation causing an increase in quantal size.
Experimentally, one way to trigger PHD is to overexpress the *Drosophila* vesicular glutamate trans-
porter gene, *VGlut*, in motor neurons. Overexpression of the glutamate transporter leads to an increase
in the diameter of glutamatergic vesicles, an increase in quantal size across the entire distribution of
spontaneous miniature events, and very large spontaneous quantal events (Daniels, 2004). To compen-
sate for this, quantal content at the NMJ is lowered, resulting in normal evoked postsynaptic excitation
(Daniels et al., 2004).

Many genes have been shown to be necessary for PHP at the NMJ. But much less is known about PHD.
PHP and PHD result in opposite changes in quantal content, and studies suggest divergent and separa-
ble mechanisms governing these forms of homeostatic plasticity. Some genes required for homeostatic
potentiation are dispensable for homeostatic depression (Gaviño et al., 2015; Li et al., 2018; Marie et
al., 2010). Moreover, unlike homeostatic potentiation, homeostatic depression does not appear to in-
volve a change in the size of the readily releasable pool of synaptic vesicles. Rather, homeostatic de-
pression appears to involve a decrease in release probability (Gaviño et al., 2015). Finally, PHP at the
NMJ appears to be a process that is dependent the input (i.e. the type of synapse formed at the NMJ)
while PHD does not appear to be input specific (Li et al., 2018).

The degree of overlap between homeostatic depression and homeostatic potentiation is unknown. We
designed a small-scale, directed screen to test for links between these two forms of homeostatic plas-
ticity. For the screen, we targeted genes based on prior evidence that their impairment in the neuron
caused a failure of the long-term maintenance of PHP. We examined loss-of-function conditions for
these genes in a VGlut overexpression background for PHD. We did not find any cases of failed ho-
meostatic depression – the conditions we examined showed decreases in quantal content in response
to increased quantal size. However, we did find an interesting and unexpected evoked neurotransmis-
sion phenotype: a robust decrease in excitatory postsynaptic potential (EPSP) amplitude in a VGlut
overexpression background because of a profound decrease in quantal content (QC). We observed this
phenotype for a double heterozygous loss-of-function condition for the Ryanodine and IP$_3$ receptors.
In our follow-up work, pharmacology phenocopied this genetic result, and our overall findings are
consistent with the idea that the PHD system may show a heightened sensitivity to low calcium.

Prior characterizations of homeostatic depression did not report decreases in EPSP amplitude in VGlut
overexpression relative to controls. Studies at the NMJ generally suggest models in which homeostatic
compensation generally maintains evoked neurotransmission at the synapse approximately at control
levels (Davis, 2013). Our results suggest that perturbation of calcium regulation may result in a defect
in tuning homeostatic responses to maintain EPSPs at control levels.

2   **Materials and Methods**

Drosophila stocks and husbandry

Fruit fly stocks were obtained from the Bloomington Drosophila Stock Center (BDSC, Bloomington,
Indiana), Kyoto Stock Center (DGRC, Kyoto, Japan), Japan National Institute of Genetics (Mishima,
Shizuoka, Japan), Vienna Drosophila Research Center (VDRC, Vienna, Austria), or from the labs that
generated them. *w^1118* was used as a wild-type (WT) control (Hazelrigg et al., 1984). RNAi lines and
mutants used in the screen are reported in Supplemental Table S1.

Fruit flies were raised on cornmeal, molasses, and yeast medium (see BDSC website for standard rec-
ipe) in temperature-controlled conditions. Animals were reared at 25°C until they reached the wander-
ing third instar larval stage, at which point they were selected for electrophysiological recording. *UAS-
VGlut* (Daniels et al., 2004) was recombined with *OK371-GAL4* (Mahr and Aberle, 2006; Meyer and
Aberle, 2006) to drive constitutive overexpression of VGlut. The full genotype of these animals is: *w;
VGlut, OK371-Gal4/CyO-GFP*. Virgins of these flies were crossed to RNAi lines or mutants to test for
changes to homeostatic depression. *w; OK371-Gal4/+* was used as a genetic control for baseline elec-
trophysiology.

Electrophysiology and analysis

Larvae were dissected in a modified HL3 saline comprised of: NaCl (70 mM), KCl (5 mM), MgCl$_2$
(10 mM), NaHCO$_3$ (10 mM), sucrose (115 mM = 3.9%), trehalose (4.2 mM = 0.16%), HEPES (5.0
mM = 0.12%), and CaCl$_2$ (0.5 mM, except as noted).

For pharmacology, Dantrolene (R&D Systems) and Xestospongin C (Abcam) were used. Dantrolene
was mixed into saline to a final concentration of 25 μM. Larvae were cut open on the dorsal side and
allowed to incubate in the Dantrolene saline for 5 minutes. The rest of the dissection and recording was
completed in Dantrolene saline. Xestospongin C was applied in a similar manner, with the animals
allowed to incubate in 20 μM Xestospongin C saline for 5 minutes before they were recorded, also in
saline containing Xestospongin C.

Electrophysiological data were collected using an Axopatch 200B amplifier (Molecular Devices,
Sunnyvale, CA) in bridge mode, digitized using a Digidata 1440A data acquisition system (Molecular
Devices), and recorded with pCLAMP 10 acquisition software (Molecular Devices). A Master-8 pulse
stimulator (A.M.P. Instruments, Jerusalem, Israel) and an ISO-Flex isolation unit (A.M.P. Instruments)
were utilized to deliver 1 ms suprathreshold stimuli to the appropriate segmental nerve. The average
spontaneous miniature excitatory postsynaptic potential (mEPSP) amplitude per NMJ was quantified
by hand, approximately 100 individual spontaneous release events per NMJ (MiniAnalysis, Synapto-
soft, Fort Lee, NJ). Measurements from all NMJs of a given condition were then averaged. For evoked
neurotransmission, 30 excitatory postsynaptic potentials (EPSPs) were averaged to find a value for
each NMJ. These were then averaged to calculate a value for each condition. Quantal content (QC)
was calculated by the ratio of average EPSP and average mEPSP amplitudes for each individual NMJ.
An average quantal content was then calculated for each condition. EPSP variability was assessed by
measuring each of the 30 traces individually and calculating a standard deviation and range for that
NMJ. Range was defined as the maximum EPSP value minus the minimum EPSP value.

Statistical Analyses

Statistical analyses were conducted using GraphPad Prism Software. Statistical significance was as-
sessed either by Student's T-Test when one experimental data set was being directly compared to a
control data set, or one-way ANOVA with Tukey's post-hoc test when multiple data sets were being
compared. Specific *p* value ranges are noted in the Figure legends and shown in graphs as follows: * *p*
< 0.05, ** *p* < 0.01, and *** *p* < 0.001 (* and # are used in Figures if there are additional comparisons
highlighted). For some comparisons that are close to *p* < 0.05 statistical significance but do not achieve
it ($0.05 < p < 0.1$), specific values are reported on the graph itself. Calcium cooperativity data were
analyzed using a non-linear fit regression analysis on GraphPad Prism.

3   Results

A recombinant line to analyze presynaptic homeostatic depression (PHD)

Using previously published reagents, we generated a fly stock with constitutive *VGlut* transgene over-
expression. Such a stock could be used as a tool for a single-cross genetic screen. To generate the stock,
we recombined the *OK371-Gal4* motor neuron driver (Mahr and Aberle, 2006; Meyer and Aberle,
2006) with a *UAS-VGlut* transgene (Daniels et al., 2004). We placed these two genetic elements in *cis*
on *Drosophila melanogaster* Chromosome II. *OK371-Gal4* is an enhancer trap line for the *VGlut* pro-
moter itself. This ensured that GAL4-driven *UAS-VGlut* overexpression would happen in desired tis-
sues, *Drosophila* motor neurons.

We tested if the recombinant line constitutively overexpressing *UAS-VGlut* could express PHD at the
NMJ. We crossed the recombinant stock to our wild-type stock (*w[1118],* herein: WT) (Cross result,

herein: "*VGlut, OK371/+*"). By NMJ electrophysiology, we recorded from WT control, *OK371/+* con-
trol, and *w; VGlut, OK371/+*. As expected, *VGlut, OK371/+* NMJs showed an increase in spontaneous
miniature excitatory postsynaptic potential (mEPSP) amplitude compared to controls (Fig. 1A-C).
Compared to WT control NMJs, there was no significant difference in evoked postsynaptic amplitudes
for *VGlut, OK371/+* NMJs (Fig. 1D). This was because of an accurate homeostatic decrease in quantal
content (QC) (Fig 1E) – hence, successful PHD. This result matched prior studies that had used WT as
a control and a *trans OK371/UAS-VGlut* combination to induce PHD (Daniels et al., 2004; Gaviño et
al., 2015; Li et al., 2018).

Even though PHD was successful relative to WT for our test cross, we noted a small, but statistically
significant, baseline increase in the EPSP amplitude of *OK371/+* NMJs. This increase in *OK371/+*
EPSP level was present compared either to WT control or to *VGlut, OK371/+* (Fig. 1D). One possibility
is that the *OK371/+* genetic background has slightly elevated release, and the combined addition of
UAS-VGlut reveals a slight depression in evoked amplitude. Noting this potentially important differ-
ence in our driver control, we continued using the *OK371/+* heterozygous condition as a genetic back-
ground control. *OK371/+* is a closer genetic control for PHD analysis than WT.

A genetic screen suggests a role for calcium stores in homeostatic depression

We used our recombinant line to conduct a genetic screen for conditions that affect presynaptic home-
ostatic depression (PHD). We crossed the recombinant stock to *UAS-RNAi* stocks and we selected lar-
vae for recording (Materials and Methods, Fig. 2A). For the screen, we targeted a subset of genes
previously identified as in the neuron for homeostatic potentiation, or closely related genes. We tested
43 genotypes (sometimes multiple conditions for a single gene), including our homeostatic depression
condition, *VGlut, OK371/+* (Fig. 2B, C).

The aggregate results of the screen are reported here (Fig. 2B, C; raw data in Supplementary Table S1).
We recorded from 42 experimental heterozygous *mutant/+* or > *UAS-RNAi/+* conditions, in the *VGlut,*
OK371/+ genetic background. Of those 42, 12 achieved EPSPs that were numerically larger than
VGlut, OK371/+, and 22 achieved QCs that were numerically larger than *VGlut, OK371/+* (Fig. 2B,
C). Increased evoked potentials could signify failed PHP – however, none of these cases represented
statistically significant increases compared to *VGlut, OK371/+*. None were so much bigger than they
were good candidates for "failed PHD." Indeed, all of the candidates had average EPSP and QC levels
below *OK371/+* NMJ baseline recordings (Compare Figs 1D, E and Fig. 2B, C).

We noted a phenotype distinct from what we were initially seeking: two crosses yielded larvae with
striking decreases in NMJ EPSP amplitudes, more than two standard deviations below the average
EPSPs from the baseline *VGlut, OK371/+* data set (Fig. 2B). One case was knockdown of the *Survival*
motor neuron (*Smn*) gene with the *UAS-Smn[RNAi]^{JF02057}* line in the *VGlut, OK371/+* background.
This was intriguing because Drosophila *Smn* is homologous to human *SMN*. Defects in *SMN* cause
Spinal Muscular Atrophy (Lefebvre et al., 1995). Drosophila Smn has been characterized as a potential
model for Spinal Muscular Atrophy (Raimer et al., 2020; Sen et al., 2011; Spring et al., 2019). Smn
has also previously been implicated in PHP (Sen et al., 2011). However, the result for *UAS-*
Smn[RNAi]^{JF02057} was not replicated by other *Smn* knockdown or loss-of-function mutant test crosses
(Fig. 2B, C). We did not follow up on *Smn* for this study.

A second case with a striking decrease in EPSP amplitude in the screen was a double heterozygous
genetic condition in genes encoding the Drosophila Ryanodine receptor (*RyR*) and inositol 1,4,5-
trisphosphate (IP$_3$) receptor (*itpr*): *VGlut, OK371/RyR^{E4340K}; itpr^{90B}/+* (Fig. 2B, C). Ryanodine recep-
tors (RyRs) and IP$_3$ receptors (IP$_3$Rs) are localized to the endoplasmic reticulum. They mediate release
of calcium from intracellular stores (Berridge, 1984, 1987, 1998; Simkus and Stricker, 2002). The

RyR^{E4340K} mutation is a single amino acid substitution (glutamic acid to lysine) (Dockendorff et al.,
2000), and the *itpr^{90B}* mutation is null mutant generated by imprecise excision of a transposon
(Venkatesh and Hasan, 1997). We previously defined a roles for RyR, IP$_3$R, IP$_3$ signaling and upstream
components in maintaining presynaptic homeostatic potentiation (PHP) (Brusich et al., 2015; James et
al., 2019).

In parallel, we screened single mutant manipulations for both genes. Neither the *RyR^{E4340K}/+* heterozy-
gous condition, nor the *itpr^{90B}/+* heterozygous condition – nor had any other single heterozygous or
RNAi knockdown conditions in either gene – yielded as significantly depressed EPSPs in response to
PHD challenge (Fig. 2B, C). Therefore, the screen result with the double heterozygote could be due to
a genetic interaction, or it could be due to other factors in the genetic background. This preliminary
finding required further characterization.

We tested if the electrophysiological phenotype could be due to a baseline neurotransmission defect
when both genes are heterozygous. By electrophysiology, we compared NMJs from *OK371/RyR^{E4340K};*
itpr^{90B}/+ larvae as a baseline double heterozygous condition vs. NMJs from *VGlut, OK371/RyR^{E4340K};*
itpr^{90B}/+ larvae (Fig. 3A-D). Just like WT, the baseline double heterozygous condition did have a slight
decrease in EPSP amplitude compared to *OK371/+* driver control (Fig. 3A). This indicated a small,
but discernible defect in neurotransmission in animals where the IP$_3$Rs and RyRs are both impaired.
Yet the double heterozygous condition with concurrent *VGlut* gene overexpression showed a lot of
depression compared to its own genetic control – increased quantal size (Fig. 3B), but markedly de-
creased evoked amplitude (Fig. 3C) because of a large decrease in quantal content (Fig. 3D), hence
"excess" PHD.

We noted that the EPSP amplitude in individual *VGlut, OK371/RyR^{E4340K}; itpr^{90B}/+* NMJ recordings
appeared to vary markedly from stimulus to stimulus. High variability could indicate unstable neuronal
excitability or release. To check if evoked release events were indeed more variable, we completed two
additional analyses. First, we extracted the amplitude of each individual EPSP event at every NMJ
recorded. From this information, we calculated the EPSP standard deviation (S.D.) per individual NMJ.
We also calculated a range for each NMJ by subtracting the maximum EPSP of the thirty from the
minimum. We averaged these S.D. and range measures for each genotype, considering all of the indi-
vidual EPSP recordings.

For both of these EPSP variability parameters, *w; VGlut, OK371/RyR^{E4340}; itpr^{90B}/+* animals showed
statistically significant higher standard deviations and ranges compared to controls (Fig. 3E). By con-
trast, double heterozygous baseline *OK371/RyR^{E4340K}; itpr^{90B}/+* NMJs did not differ significantly from
w; OK371/+ driver control NMJs, suggesting that the variability stems from *VGlut* overexpression in
the mutant background (Fig. 3E). *w; VGlut, OK371/+* NMJs showed numerically higher variability
than *w; OK371/+*, but this was not statistically significant (Fig. 3E).

Pharmacology targeting Ryanodine and IP$_3$ receptors recapitulates loss-of-function genetics

We tested if the electrophysiological phenotypes we observed genetically could be recapitulated by
combining genetics and pharmacology. We started with the drug Dantrolene. Dantrolene is a RyR an-
tagonist (Vazquez-Martinez et al., 2003; Zhao et al., 2001). In prior work at the *Drosophila* NMJ, we
found that application of Dantrolene can abrogate the long-term maintenance of PHP (James et al.,
2019).

For our first test with Dantrolene, we used a sensitized *OK371/+; itpr^{90B}/+* genetic background. With
this background, we could pharmacologically impair RyRs while genetically impairing IP$_3$Rs. We ap-
plied 25μM of Dantrolene to: 1) *OK371/+* NMJs; 2) *VGlut, OK371/+* NMJs; 3) *OK371/+; itpr^{90B}/+*

NMJs; and 4) *VGlut, OK371/+; itpr*^{90B}/+ NMJs. The mEPSP amplitudes in both *VGlut*-overexpressing
lines were elevated compared to their respective controls (Fig. 4A). This indicated that in the presence
of Dantrolene, there was still homeostatic pressure that could induce PHD. EPSP amplitudes in both
VGlut-overexpressing lines were significantly decreased compared to their respective genetic controls
(Fig. 4B). This was because of a marked decrease in QC (Fig. 4A). Moreover, the *VGlut, OK371/+;*
itpr^{90B}/+ condition with Dantrolene had significantly decreased evoked transmission compared to
VGlut, OK371/+ with Dantrolene (Fig. 4A, B). This indicated a cumulative effect of impairing both
the IP$_3$Rs and RyRs in a PHD-challenged background.

It is possible that strong impairment of RyRs was sufficient to disorient the PHD regulation system.
We ran additional pharmaco-genetic tests with Dantrolene, this time using a second sensitized genetic
background, *OK371/RyR*^{E4340K} – with and without a *UAS-VGlut* overexpression. Again, mEPSPs were
evoked potentials became significantly larger when *VGlut* was overexpressed (Fig. 5A, left), but EPSPs
were significantly down (Fig. 5A, middle) because of a marked decrease in quantal content (Fig. 5A,
right).

Finally, we attempted the inverse pharmaco-genetic experiment from that in Figure 4. This time we
used the IP$_3$R inhibitor, Xestospongin C (Gafni et al., 1997; Wilcox et al., 1998) and the sensitized
OK371/RyR^{E4340K} genetic background. We applied 20 μM Xestospongin C, both to *OK371/RyR*^{E4340K}
NMJs and to *VGlut, OK371/RyR*^{E4340K} NMJs. The mEPSPs were numerically larger when *VGlut* was
overexpressed (Fig. 5B, left) – though this time, for the Xestospongin C dataset, the data did not achieve
statistical significance for mEPSP size ($p = 0.10$). This could indicate weak homeostatic pressure in the
presence of Xestospongin C. Nevertheless, EPSPs were significantly down (Fig. 5B, middle) because
of a marked decrease in quantal content (Fig. 5B, right).

Taking our data together, for each case where we examined a dual impairment of RyR and IP$_3$R the
EPSP amplitudes and QC were all quite low with concomitant VGlut overexpression (Figs. 3-5). These
results are consistent with excess depression.

Excess PHD in very low extracellular calcium

We wondered how impairment of channels that mediate release of calcium from intracellular stores
might cause the electrophysiological phenotypes that we observed. Our prior work suggested that these
ER calcium store channels and the signaling systems that control them are required to maintain home-
ostatic potentiation throughout life (Brusich et al., 2015; James et al., 2019). We also found a related
result: impairing Ca^{2+} store release mollified hyperexcitability phenotypes caused by gain-of-function
Ca$_V$2 amino-acid substitutions in the alpha1 subunit Cacophony. Ca$_V$2 channels mediate synaptic cal-
cium influx at the NMJ (Brusich et al., 2018). In light of these prior data, we considered two possibil-
ities for PHD. One model is that the IP$_3$R and RyR channels play a role in orienting PHD and ensuring
a proper level of depression. A different model is that calcium itself plays the important role. If this
latter idea were true, it might be the case that lowering calcium influx into the synaptic cleft would also
be sufficient to disorient the PHD system, resulting in excess depression.

As a test, we measured release over a range of low extracellular calcium concentrations (0.2-0.5 mM).
We examined six genotypes: 1) WT; 2) *w; OK371/+* ; 3) *w; VGlut, OK371/+*; 4) *w; RyR*^{E4340K}/+;
itpr^{90B}/+; 5) *w; OK371/RyR*^{E4340K} ; *itpr*^{90B}/+; and 6) *w; VGlut, OK371/RyR*^{E4340K} ; *itpr*^{90B}/+. To organ-
ize data and to calculate calcium cooperativity, we plotted quantal content as a function of calcium
concentration, with the x-y axes on a log-log scale (Fig. 6A, B). To account for different Ca^{2+} driving

forces in the different concentrations, we corrected QC for nonlinear summation in our plots and in our
subsequent analyses (NLS Corrected QC) (Martin, 1955).

Non-linear regression analyses revealed that there was not a significant difference in calcium cooper-
ativity between any of these genotypes over the range of extracellular $[Ca^{2+}]$ we tested (Fig. 6A, B).
The calculated log-log slope values of the control PHD genotypes were: WT (log-log slope = 1.810),
w; OK371/+ (log-log slope = 1.884), and *w; VGlut, OK371/+* (log-log slope = 2.117). Comparing those
three slopes with one another by nonlinear regression yielded no significant difference in slope (p =
0.91). The log-log slope values of the double heterozygous conditions were: *w; RyR^{E4340K}/+ ; itpr^{90B}/+*
(log-log slope = 1.737), *w; OK371/RyR^{E4340K}; itpr^{90B}/+* (log-log slope = 2.102), and *w; VGlut,*
OK371/RyR^{E4340K} ; itpr^{90B}/+ (log-log slope = 1.601). Comparing those slopes with one another also
yielded no significant difference (p = 0.77).

Even though there was no significant difference in calcium cooperativity of release over the range of
low $[Ca^{2+}]$ conditions examined, our data did show a very large drop in release between 0.3 and 0.2
mM $[Ca^{2+}]$ – specifically for the genotypes where PHD was induced by *UAS-VGlut* overexpression, or
for the genotypes with a double heterozygous impairment of *RyR* and *itpr*. Examining the raw data at
0.2 mM $[Ca^{2+}]$, we observed that there was significant homeostatic pressure for PHD signified by
mEPSP amplitude increases in the VGlut-overexpression background (Fig. 6C, left). Yet except for the
control NMJs, EPSP amplitudes were very much diminished (Fig. 6C, middle) because of stark drops
in QC (Fig. 6C, right).

Together, the data point to two conclusions. First, low extracellular calcium on its own appears to be a
case where the synapse experiences "excess" PHD (Fig. 6C, *VGlut, OK371/+* data). Second, double
heterozygous impairment of *RyR* and *itpr* appears to cause very low levels of baseline release in low
calcium, irrespective of PHD challenge (Fig. 6C, middle; compare with Fig. 3C). Taken together, these
data suggest that lowering presynaptic calcium by any means (impairing store release and/or impairing
influx) is sufficient to disorient the homeostatic set point for release.

Excess PHD with impaired Ca_V2 function

As a final test of the idea depressed presynaptic calcium causes excess PHD, we turned to genetics.
Drosophila Ca_V2 channels mediate synaptic calcium influx at the NMJ. We used a hypomorphic mutant
in the Ca_V2 alpha1 subunit-encoding *cacophony* gene, *cac^S*, to limit calcium influx. Ca_V2 is essential
for viability, but *cac^S* hypomorphs are viable and fertile (Kawasaki et al., 2000; Smith et al., 1998).
Prior work showed that the *cac^S* homozygous condition dampens NMJ EPSP amplitude by about 70-
80% (Frank et al., 2006); calcium imaging data suggest this is due to a ~50% decrease in Ca^{2+} influx
during evoked stimulation (Müller and Davis, 2012). Beyond this phenotype in baseline neurotrans-
mission, *cac^S* hypomorphs also block PHP expression and PHP-associated increases in presynaptic
calcium influx (Frank et al., 2006; Müller and Davis, 2012).

With a single cross, we generated hemizygous *cac^S/Y; VGlut, OK371/+* male larvae (Fig. 7A). Com-
pared to *cac^S/Y* as a baseline mutant control, *cac^S/Y; VGlut, OK371/+* NMJs have a marked increase in
mEPSP size (Fig. 7B), indicating homeostatic pressure to induce PHD (Fig. 7B). However, comparing
evoked potentials of those two conditions shows that *cac^S/Y; VGlut, OK371/+* NMJs have much
smaller EPSPs (Fig. 7C). This was because of an extreme decrease in QC (Fig. 7D) – or "excess" PHD.

4   Discussion

We began this study in search of genetic conditions that affect PHD (Fig. 2). While we did not find any
conditions that result in a block of PHD, we did find conditions that disrupt the tuning of PHD and

provide insight into the role calcium regulation plays in this form of homeostatic plasticity. When IP_3R
and RyR functions are partially impaired – either by genetics or by pharmacology – the NMJ still
executes a PHD-like process. But that process goes beyond what is appropriate for the homeostatic
pressure that is applied to the system. As a result, evoked potentials at the NMJ are much smaller than
baseline (Figs. 3-5). A similar "excess depression" phenotype is observed when extracellular $[Ca^{2+}]$ is
lowered to 0.2 mM (Fig. 6) and when the Ca_V2 alpha1 subunit gene *cacophony* harbors a hypomorphic
mutation, cac^S (Fig. 7).

This phenotype has important implications for proper homeostatic control of synapse function. Taking
our data together, we propose that presynaptic calcium plays an important role in tuning PHD to the
appropriate level. Perturbations that dampen calcium efflux from stores or perturbations that dampen
calcium influx from the extracellular environment can both disrupt PHD tuning (Fig. 8).

Known roles for calcium in tuning homeostatic plasticity

The notion that calcium plays a role in homeostatic signaling is not new. Many roles for voltage-gated
calcium channels in synaptic homeostasis are well-documented (Frank, 2014a, b). Prior to our study,
there was evidence for voltage-gated calcium channel regulation for both NMJ PHP and PHD. For
PHP, loss-of-function conditions in Ca_V2/*cacophony* can impair or block this form of homeostatic reg-
ulation (Frank et al., 2006; Frank et al., 2009; Müller and Davis, 2012; Spring et al., 2016). Calcium
imaging experiments suggest that the reason is because an increase in calcium influx through Ca_V2 is
required for the upregulation of quantal content during PHP, and mutant conditions like cac^S block this
increase (Müller and Davis, 2012). Recent studies report that Cacophony and other active zone protein
levels increase at the NMJ active zone in response to PHP homeostatic challenges (Böhme et al., 2019;
Goel et al., 2019; Gratz et al., 2019). And work from mammalian systems mirrors these findings. For
example, with mouse hippocampal cultures, TTX exposure induces a homeostatic decrease in presyn-
aptic calcium influx (Zhao et al., 2011).

The converse appears true for PHD. Calcium imaging data from two different studies has shown a
decrease in the size of calcium transients at the NMJ in response to presynaptic nerve firing in VGlut-
overexpressing animals (Gaviño et al., 2015; Li et al., 2018). The data are mixed on how these de-
creased transients might come about during PHD. Using a tagged *UAS-cacophony* cDNA transgene,
two studies verified there was a reduction in the amount GFP-tagged Cacophony alpha1 subunits in
Ca_V2 in a VGlut-overexpressing background (Gaviño et al., 2015; Gratz et al., 2019) However, one of
these same studies demonstrated that if a tagged genomic construct is used instead, that same Ca_V2
reduction is not observed (Gratz et al., 2019). Since the transgenic tagged Cacophony-GFP is the prod-
uct of a single *cac* splice isoform (Kawasaki et al., 2002; Kawasaki et al., 2004), it could be the case
that some isoforms are more dynamically trafficked at the synapse. Another possibility is that existing
active zone components are somehow modulated during PHD. Regardless of the actual mechanism,
the phenomenon appears conserved: again, with rodent hippocampal preparations, increased neuronal
activity through gabazine exposure induces a PHD-like phenomenon ultimately resulting in decreases
in calcium influx and release (Jeans et al., 2017; Zhao et al., 2011).

Similarities and differences with prior PHD studies at the NMJ

We were able to conduct a PHD screen using our recombinant stock with the *UAS-VGlut* and *OK371-*
Gal4 elements on the same chromosome. We acknowledge that this type of *cis* strain construction is
an unorthodox choice for Drosophila genetics. This is because *UAS-VGlut* is continually overexpressed

every generation. In principle, such a stock can pick up modifier mutations. The trade-off for our work
was a simplified, single-generation crossing scheme for genetic screens. As it turns out, our recombi-
nant stock with the driver and *UAS* elements in *cis* maintains consistent PHD challenge from generation
to generation, and it behaves similarly electrophysiologically to *trans OK371/VGlut* combinations used
in other studies (Daniels et al., 2004; Gaviño et al., 2015; Li et al., 2018).

There are differences between our study and the findings of other published work. Prior studies have
used WT (or *w1118*) as a control background when compared to VGlut overexpression (Daniels et al.,
2004; Gaviño et al., 2015; Li et al., 2018). This is a standard practice. Those studies reported precise
PHD when comparing WT vs. *OK371/VGlut* third instar larvae – decreased QC at *OK371/VGlut* NMJs
resulting in unchanged evoked transmission. We replicated this finding (Fig. 1). However, we also used
our Gal4 driver stock background *OK371/+* as an additional control. For that comparison, we saw a
slight depression in the evoked amplitude of *OK371, VGlut/+* NMJs (Fig. 1). One possibility is that
our recombinant stock was acting as a sensitized background to uncover conditions that might cause
excess depression.

A second difference comes from the low extracellular calcium test. A low extracellular calcium exper-
iment was previously done when VGlut overexpression was first characterized (Daniels et al., 2004).
For that study, the authors showed that QC was significantly diminished compared to wild-type NMJs
by the method of failure analysis. Taking the data of that study in aggregate, the authors concluded that
PHD was intact in a variety of conditions, including saline with very low extracellular $[Ca^{2+}]$ (0.23 mM
Ca^{2+}, 20 mM Mg^{2+}). Our study may appear to conflict with that study because we found that saline
with very low $[Ca^{2+}]$ (0.2 mM Ca^{2+}, 10mM Mg^{2+}) is conducive to "excess PHD." One possibility is
that since the original study was examining failure percentage vs. WT – and not the absolute value of
mEPSPs or EPSPs in low calcium – a finding like excess depression might not be as easily observed.
Other differences might be attributed genetic background or other differences in recording saline, like
magnesium concentration.

Finally, one other study previously examined the effects of a *cacS* mutation with concomitant VGlut
overexpression (Gaviño et al., 2015). The authors did not find the low evoked potentials that we report.
The major difference between that experiment and ours is that the prior work examined the *cacS* mu-
tation in an extracellular $[Ca^{2+}]$ (1.0 mM) that was double that of our study. The result was a Ca^{2+}
driving force that yielded robust baseline EPSPs, even in the *cacS* mutant background (Gaviño et al.,
2015). Given our results with calcium concentration (Fig. 6), a similar effect may work to resolve any
sort of tuning problem with PHD compensation that we uncovered here.

How do calcium stores tune PHD?

How exactly might calcium stores impact PHD tuning? Less is understood about this. We know that
endoplasmic reticulum (ER) can be visualized at Drosophila NMJ terminals (Summerville et al., 2016),
and recently developed imaging tools can show how nerve stimulation results in dynamic changes to
ER lumenal calcium, including at the Drosophila NMJ (de Juan-Sanz et al., 2017; Handler et al., 2019;
Oliva et al., 2020). In parallel, other groups working at the NMJ have demonstrated important roles in
baseline neurotransmission and in PHP for ER resident proteins (Genç et al., 2017; Kikuma et al.,
2017). And from our prior work at the NMJ, we know that store calcium channels and upstream sig-
naling components are important for maintaining the NMJ's capacity for PHP throughout life (Brusich
et al., 2015; James et al., 2019). We also know that disrupting these same factors can ameliorate hy-
perexcitability associated with gains of Ca_V2 function (Brusich et al., 2018). Finally, from mammalian
work it is clear that IP_3Rs, RyRs, and intracellular calcium govern a variety of forms of neuroplasticity
(Berridge, 2016), including paired pulse facilitation (Emptage et al., 2001), and modulation of voltage-
gated calcium channel activity (Catterall, 2011; Lee et al., 2000).

If PHD were simply a matter of properly functioning neurotransmission machinery, then it is not en-
tirely obvious why PHD would be so sensitive to the amount calcium available such that its tuning
point would become disoriented, either when store-operated channels were impaired or when the
amount of influx was lowered. In our study, neurotransmission has not been lowered beyond a point of
synapse failure. This means that there is still functional machinery. And PHD, per se, is not disrupted
– indeed, there is still depression. With any type of homeostatic system, there not only needs to be error
detection (large quantal size) and correction (decreased quantal content), but there also need to be
brakes applied to the system to prevent some kind of overcorrection. Our data suggest some manner of
PHD "overcorrection." In our view, this is an interesting and understudied type of phenomenon that
could be examined in other homeostatic systems as well.

So how exactly do levels of calcium (or the function of distinct types of calcium channels found at the
synapse) ultimately affect PHD correction levels? This is a difficult problem. The first step might be
to narrow the relevant tissue type(s) involved in PHD signaling. ER and store-operated channels are
relevant to the functions of many tissues. In principle, our genetic loss-of-function manipulations to
itpr and *RyR* could affect store-operated channels either in the neuron or in the muscle or in surrounding
tissues like glia. Our pharmacological manipulations using Dantrolene and Xestospongin C could also
affect multiple tissue types. Therefore, in principle, changing the levels of cytosolic calcium could
either affect local signaling in the neuron, or it could result in aberrant signaling back to the presynaptic
neuron, disorienting the homeostat.

We favor the idea that the relevant calcium signal is local in the motor neuron for two reasons. First,
from our own data, we were able to observe the excess depression (or overcorrection) phenotype either
with manipulations to store calcium or with manipulations that affect presynaptic calcium influx, in-
cluding partial loss-of-function of neuronal *cacophony*. Second, a recent study puts forth data suggest-
ing that when VGlut overexpression induces PHD, this happens exclusively because of excess presyn-
aptic glutamate release, and presynaptic depression is initiated independent of any sort of postsynaptic
response (Li et al., 2018). Such an autocrine signaling mechanism could very well reveal a role for
intracellular calcium signaling in the presynapse.

5   References

Berridge, M.J. (1984). Inositol trisphosphate and diacylglycerol as second messengers. Biochem J *220*,
345-360.

Berridge, M.J. (1987). Inositol trisphosphate and diacylglycerol: two interacting second messengers.
Annu Rev Biochem *56*, 159-193.

Berridge, M.J. (1998). Neuronal calcium signaling. Neuron *21*, 13-26.

Berridge, M.J. (2016). The Inositol Trisphosphate/Calcium Signaling Pathway in Health and Disease.
Physiol Rev *96*, 1261-1296.

Böhme, M.A., McCarthy, A.W., Grasskamp, A.T., Beuschel, C.B., Goel, P., Jusyte, M., Laber, D.,
Huang, S., Rey, U., Petzoldt, A.G.*, et al.* (2019). Rapid active zone remodeling consolidates
presynaptic potentiation. Nat Commun *10*, 1085.

Brusich, D.J., Spring, A.M., and Frank, C.A. (2015). A single-cross, RNA interference-based genetic
tool for examining the long-term maintenance of homeostatic plasticity. Frontiers in cellular
neuroscience *9*, 107.

Brusich, D.J., Spring, A.M., James, T.D., Yeates, C.J., Helms, T.H., and Frank, C.A. (2018).
Drosophila CaV2 channels harboring human migraine mutations cause synapse hyperexcitability that
can be suppressed by inhibition of a Ca2+ store release pathway. PLoS genetics *14*, e1007577.

Catterall, W.A. (2011). Voltage-gated calcium channels. Cold Spring Harbor perspectives in biology
3, a003947.

Cull-Candy, S.G., Miledi, R., Trautmann, A., and Uchitel, O.D. (1980). On the release of transmitter
at normal, myasthenia gravis and myasthenic syndrome affected human end-plates. J Physiol *299*, 621-
638.

Daniels, R.W., Collins, C.A., Gelfand, M.V., Dant, J., Brooks, E.S., Krantz, D.E., and DiAntonio, A.
(2004). Increased expression of the Drosophila vesicular glutamate transporter leads to excess
glutamate release and a compensatory decrease in quantal content. J Neurosci *24*, 10466-10474.

Davis, G.W. (2013). Homeostatic signaling and the stabilization of neural function. Neuron *80*, 718-
728.

Davis, G.W., DiAntonio, A., Petersen, S.A., and Goodman, C.S. (1998). Postsynaptic PKA controls
quantal size and reveals a retrograde signal that regulates presynaptic transmitter release in Drosophila.
Neuron *20*, 305-315.

Davis, G.W., and Müller, M. (2015). Homeostatic control of presynaptic neurotransmitter release.
Annu Rev Physiol *77*, 251-270.

de Juan-Sanz, J., Holt, G.T., Schreiter, E.R., de Juan, F., Kim, D.S., and Ryan, T.A. (2017). Axonal
Endoplasmic Reticulum Ca(2+) Content Controls Release Probability in CNS Nerve Terminals.
Neuron *93*, 867-881 e866.

Delvendahl, I., and Müller, M. (2019). Homeostatic plasticity-a presynaptic perspective. Curr Opin
Neurobiol *54*, 155-162.

Dockendorff, T.C., Robertson, S.E., Faulkner, D.L., and Jongens, T.A. (2000). Genetic
characterization of the 44D-45B region of the Drosophila melanogaster genome based on an F2 lethal
screen. Molecular & general genetics : MGG *263*, 137-143.

Emptage, N.J., Reid, C.A., and Fine, A. (2001). Calcium stores in hippocampal synaptic boutons
mediate short-term plasticity, store-operated Ca2+ entry, and spontaneous transmitter release. Neuron
29, 197-208.

Frank, C.A. (2014a). Homeostatic plasticity at the Drosophila neuromuscular junction.
Neuropharmacology *78*, 63-74.

Frank, C.A. (2014b). How voltage-gated calcium channels gate forms of homeostatic synaptic
plasticity. Frontiers in cellular neuroscience *8*, 40.

Frank, C.A., James, T.D., and Muller, M. (2020). Homeostatic control of Drosophila neuromuscular
junction function. Synapse *74*, e22133.

Frank, C.A., Kennedy, M.J., Goold, C.P., Marek, K.W., and Davis, G.W. (2006). Mechanisms
underlying the rapid induction and sustained expression of synaptic homeostasis. Neuron *52*, 663-677.

Frank, C.A., Pielage, J., and Davis, G.W. (2009). A presynaptic homeostatic signaling system
composed of the Eph receptor, ephexin, Cdc42, and CaV2.1 calcium channels. Neuron *61*, 556-569.

Gafni, J., Munsch, J.A., Lam, T.H., Catlin, M.C., Costa, L.G., Molinski, T.F., and Pessah, I.N. (1997).
Xestospongins: potent membrane permeable blockers of the inositol 1,4,5-trisphosphate receptor.
Neuron *19*, 723-733.

Gaviño, M.A., Ford, K.J., Archila, S., and Davis, G.W. (2015). Homeostatic synaptic depression is
achieved through a regulated decrease in presynaptic calcium channel abundance. eLife *4*.

Genç, Ö., Dickman, D.K., Ma, W., Tong, A., Fetter, R.D., and Davis, G.W. (2017). MCTP is an ER-
resident calcium sensor that stabilizes synaptic transmission and homeostatic plasticity. eLife *6*.

Goel, P., Dufour Bergeron, D., Bohme, M.A., Nunnelly, L., Lehmann, M., Buser, C., Walter, A.M.,
Sigrist, S.J., and Dickman, D. (2019). Homeostatic scaling of active zone scaffolds maintains global
synaptic strength. J Cell Biol *218*, 1706-1724.

Gratz, S.J., Goel, P., Bruckner, J.J., Hernandez, R.X., Khateeb, K., Macleod, G.T., Dickman, D., and
O'Connor-Giles, K.M. (2019). Endogenous Tagging Reveals Differential Regulation of Ca(2+)
Channels at Single Active Zones during Presynaptic Homeostatic Potentiation and Depression. J
Neurosci *39*, 2416-2429.

Handler, A., Graham, T.G.W., Cohn, R., Morantte, I., Siliciano, A.F., Zeng, J., Li, Y., and Ruta, V.
(2019). Distinct Dopamine Receptor Pathways Underlie the Temporal Sensitivity of Associative
Learning. Cell *178*, 60-75 e19.

Hazelrigg, T., Levis, R., and Rubin, G.M. (1984). Transformation of white locus DNA in drosophila:
dosage compensation, zeste interaction, and position effects. Cell *36*, 469-481.

James, T.D., Zwiefelhofer, D.J., and Frank, C.A. (2019). Maintenance of homeostatic plasticity at the
Drosophila neuromuscular synapse requires continuous IP3-directed signaling. eLife *8*.

Jeans, A.F., van Heusden, F.C., Al-Mubarak, B., Padamsey, Z., and Emptage, N.J. (2017). Homeostatic
Presynaptic Plasticity Is Specifically Regulated by P/Q-type Ca(2+) Channels at Mammalian
Hippocampal Synapses. Cell reports *21*, 341-350.

Kawasaki, F., Collins, S.C., and Ordway, R.W. (2002). Synaptic calcium-channel function in
Drosophila: analysis and transformation rescue of temperature-sensitive paralytic and lethal mutations
of cacophony. J Neurosci *22*, 5856-5864.

Kawasaki, F., Felling, R., and Ordway, R.W. (2000). A temperature-sensitive paralytic mutant defines
a primary synaptic calcium channel in Drosophila. J Neurosci *20*, 4885-4889.

Kawasaki, F., Zou, B., Xu, X., and Ordway, R.W. (2004). Active zone localization of presynaptic
calcium channels encoded by the cacophony locus of Drosophila. J Neurosci *24*, 282-285.

Kikuma, K., Li, X., Kim, D., Sutter, D., and Dickman, D.K. (2017). Extended Synaptotagmin Localizes
to Presynaptic ER and Promotes Neurotransmission and Synaptic Growth in Drosophila. Genetics *207*,
993-1006.

Lee, A., Scheuer, T., and Catterall, W.A. (2000). Ca2+/calmodulin-dependent facilitation and
inactivation of P/Q-type Ca2+ channels. J Neurosci *20*, 6830-6838.

Lefebvre, S., Burglen, L., Reboullet, S., Clermont, O., Burlet, P., Viollet, L., Benichou, B., Cruaud,
C., Millasseau, P., Zeviani, M.*, et al.* (1995). Identification and characterization of a spinal muscular
atrophy-determining gene. Cell *80*, 155-165.

Li, X., Goel, P., Wondolowski, J., Paluch, J., and Dickman, D. (2018). A Glutamate Homeostat
Controls the Presynaptic Inhibition of Neurotransmitter Release. Cell reports *23*, 1716-1727.

Mahr, A., and Aberle, H. (2006). The expression pattern of the Drosophila vesicular glutamate
transporter: a marker protein for motoneurons and glutamatergic centers in the brain. Gene expression
patterns : GEP *6*, 299-309.

Marder, E., and Goaillard, J.M. (2006). Variability, compensation and homeostasis in neuron and
network function. Nat Rev Neurosci *7*, 563-574.

Marie, B., Pym, E., Bergquist, S., and Davis, G.W. (2010). Synaptic homeostasis is consolidated by
the cell fate gene gooseberry, a Drosophila pax3/7 homolog. J Neurosci *30*, 8071-8082.

Martin, A.R. (1955). A further study of the statistical composition on the end-plate potential. J Physiol
130, 114-122.

Meyer, F., and Aberle, H. (2006). At the next stop sign turn right: the metalloprotease Tolloid-related
1 controls defasciculation of motor axons in Drosophila. Development *133*, 4035-4044.

Müller, M., and Davis, G.W. (2012). Transsynaptic control of presynaptic Ca(2)(+) influx achieves
homeostatic potentiation of neurotransmitter release. Curr Biol *22*, 1102-1108.

Oliva, M.K., Perez-Moreno, J.J., O'Shaughnessy, J., Wardill, T.J., and O'Kane, C.J. (2020).
Endoplasmic Reticulum Lumenal Indicators in Drosophila Reveal Effects of HSP-Related Mutations
on Endoplasmic Reticulum Calcium Dynamics. Front Neurosci *14*, 816.

Petersen, S.A., Fetter, R.D., Noordermeer, J.N., Goodman, C.S., and DiAntonio, A. (1997). Genetic
analysis of glutamate receptors in Drosophila reveals a retrograde signal regulating presynaptic
transmitter release. Neuron *19*, 1237-1248.

Pozo, K., and Goda, Y. (2010). Unraveling mechanisms of homeostatic synaptic plasticity. Neuron *66*,
337-351.

Raimer, A.C., Singh, S.S., Edula, M.R., Paris-Davila, T., Vandadi, V., Spring, A.M., and Matera, A.G.
(2020). Temperature-sensitive spinal muscular atrophy-causing point mutations lead to SMN
instability, locomotor defects and premature lethality in Drosophila. Dis Model Mech *13*.

Sen, A., Yokokura, T., Kankel, M.W., Dimlich, D.N., Manent, J., Sanyal, S., and Artavanis-Tsakonas,
S. (2011). Modeling spinal muscular atrophy in Drosophila links Smn to FGF signaling. J Cell Biol
192, 481-495.

Simkus, C.R., and Stricker, C. (2002). The contribution of intracellular calcium stores to mEPSCs
recorded in layer II neurones of rat barrel cortex. J Physiol *545*, 521-535.

Smith, L.A., Peixoto, A.A., Kramer, E.M., Villella, A., and Hall, J.C. (1998). Courtship and visual
defects of cacophony mutants reveal functional complexity of a calcium-channel alpha1 subunit in
Drosophila. Genetics *149*, 1407-1426.

Spring, A.M., Brusich, D.J., and Frank, C.A. (2016). C-terminal Src Kinase Gates Homeostatic
Synaptic Plasticity and Regulates Fasciclin II Expression at the Drosophila Neuromuscular Junction.
PLoS genetics *12*, e1005886.

Spring, A.M., Raimer, A.C., Hamilton, C.D., Schillinger, M.J., and Matera, A.G. (2019).
Comprehensive Modeling of Spinal Muscular Atrophy in Drosophila melanogaster. Frontiers in
molecular neuroscience *12*, 113.

Summerville, J.B., Faust, J.F., Fan, E., Pendin, D., Daga, A., Formella, J., Stern, M., and McNew, J.A.
(2016). The effects of ER morphology on synaptic structure and function in Drosophila melanogaster.
Journal of cell science *129*, 1635-1648.

Turrigiano, G.G. (2008). The self-tuning neuron: synaptic scaling of excitatory synapses. Cell *135*,
422-435.

Vazquez-Martinez, O., Canedo-Merino, R., Diaz-Munoz, M., and Riesgo-Escovar, J.R. (2003).
Biochemical characterization, distribution and phylogenetic analysis of Drosophila melanogaster

ryanodine and IP3 receptors, and thapsigargin-sensitive Ca2+ ATPase. Journal of cell science *116*,
2483-2494.

Venkatesh, K., and Hasan, G. (1997). Disruption of the IP3 receptor gene of Drosophila affects larval
metamorphosis and ecdysone release. Curr Biol *7*, 500-509.

Wang, X., Pinter, M.J., and Rich, M.M. (2016). Reversible Recruitment of a Homeostatic Reserve Pool
of Synaptic Vesicles Underlies Rapid Homeostatic Plasticity of Quantal Content. J Neurosci *36*, 828-
836.

Wilcox, R.A., Primrose, W.U., Nahorski, S.R., and Challiss, R.A. (1998). New developments in the
molecular pharmacology of the myo-inositol 1,4,5-trisphosphate receptor. Trends Pharmacol Sci *19*,
467-475.

Yeates, C.J., Zwiefelhofer, D.J., and Frank, C.A. (2017). The Maintenance of Synaptic Homeostasis at
the Drosophila Neuromuscular Junction Is Reversible and Sensitive to High Temperature. eNeuro *4*.

Zhao, C., Dreosti, E., and Lagnado, L. (2011). Homeostatic synaptic plasticity through changes in
presynaptic calcium influx. J Neurosci *31*, 7492-7496.

Zhao, F., Li, P., Chen, S.R., Louis, C.F., and Fruen, B.R. (2001). Dantrolene inhibition of ryanodine
receptor Ca2+ release channels. Molecular mechanism and isoform selectivity. J Biol Chem *276*,
13810-13816.

594   6   Figure Legends

Figure 1. Presynaptic homeostatic depression (PHD) works successfully with a recombinant line
of *OK371-Gal4* and *UAS-VGlut*. (A) NMJ electrophysiological data for miniature excitatory postsyn-
aptic potentials (mEPSP), excitatory postsynaptic potentials (EPSP), and quantal content (QC). Data
are normalized to WT (w^{1118}) values. *VGlut, OK371/+* NMJs have increased mEPSP but normal EPSP
because of decreased QC, indicative of successful PHD. (*** $p < 0.001$ vs. WT by one-way ANOVA
with Tukey's post-hoc). **(B)** Representative electrophysiological traces. Large traces are EPSPs; small
traces are mEPSPs. Scale bars for EPSPs (mEPSPs) are 5 mV (1 mV) and 50 ms (1000 ms). **(C)** Raw
data for mEPSPs. **(D)** Raw data for EPSPs. **(E)** Raw data for QC. For (C)-(E), bars are averages and
error bars are ± SEM. *** $p < 0.001$ vs. WT or vs. *OK371/+*; # $p < 0.05$ vs. *OK371/+*; analyses by
one-way ANOVA with Tukey's post-hoc.

Figure 2. A screen to uncover conditions with defective PHD. (A) Crossing scheme for generating
larvae for electrophysiological recording. Each animal recorded had a homeostatic challenge provided
by VGlut overexpression and a concurrent heterozygous or RNAi condition. **(B)** Data distribution for
screened conditions (x-axis = average EPSP for condition; y-axis = average QC for condition). Green
= *UAS-VGlut, OK371-Gal4/+*. Red = *UAS-VGlut, OK371-Gal4/RyRE4340K; itpr90B/+*. Purple =
UAS-VGlut, OK371-Gal4/+; UAS-Smn[RNAi]^{JF02057}/+. Dotted line: EPSP value two standard devia-
tions below *UAS-VGlut, OK371-Gal4/+* chosen as a cut off for potential follow-up hits. **(C)** Average
EPSPs for screened conditions. All conditions have a *UAS-VGlut, OK371-Gal4/+* genetic background.
">" denotes as *UAS* construct or RNAi line being driven in motor neurons by *OK371-Gal4*. "+" denotes
additional mutations present as heterozygotes. Top dotted line denotes *UAS-VGlut, OK371-Gal4/+*
average. Bottom dotted line denotes two standard deviations below *UAS-VGlut, OK371-Gal4/+* aver-
age.

Figure 3. Double heterozygous loss of the *itpr* and *RyR* genes causes excess depression. Note:
Traces and data for *OK371/+* and *VGlut, OK371/+* are repeated from Figure 1 for comparison pur-
poses. Abbreviations are as in Figure 1 as well. **(A)** NMJ electrophysiological data for mEPSP, EPSP,
and QC. Data are normalized to *OK371/+* values. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ vs.
OK371/+; ## $p < 0.01$ and ### $p < 0.001$ vs. *OK371/RyRE4340K; itpr90B/+*; analyses by one-way
ANOVA with Tukey's post-hoc. **(B)** Raw data for mEPSPs. **(C)** Raw data for EPSPs. **(D)** Raw data
for QC. For (B)-(D), bars are averages and error bars are ± SEM. * $p < 0.05$, ** $p < 0.01$, and *** $p <
0.001$ by one-way ANOVA with Tukey's post-hoc. **(E)** Representative electrophysiological traces with
S.D. and range values for EPSPs shown. The evoked amplitude S.D. and range were significantly
higher for *VGlut, OK371/RyR^{E4340K}; itpr^{90B}/+* vs. its genetic control, *OK371/RyR^{E4340K}; itpr^{90B}/+*. * p
< 0.05, ** $p < 0.01$ by one-way ANOVA with Tukey's post-hoc. Scale bars as in Figure 1.

Figure 4. Genetic impairment of *itpr* combined with pharmacological impairment of RyR causes
excess depression. (A) Raw data for mEPSPs (left); raw data for EPSPs (middle); raw data for QC
(right). All data are for the indicated NMJ genotypes with 25 µM Dantrolene; bars are averages and
error bars are ± SEM. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ by one-way ANOVA with Tukey's
post-hoc. **(B)** Representative EPSP traces. Scale bars are as in Figure 1.

Figure 5. Additional pharmaco-genetic combinations cause excess depression. (A) Raw data for
mEPSPs (left); raw data for EPSPs (middle); raw data for QC (right). All data are for the indicated
NMJ genotypes with 25 µM Dantrolene; bars are averages and error bars are ± SEM. **(B)** Data as in
(A) but with 20 µM Xestospongin C instead of Dantrolene. **(C)** Representative EPSP traces. Scale bars
are as in Figure 1. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ by Student's T-Test comparing a control
dataset (no VGlut overexpression) vs. an experimental dataset (VGlut overexpression).

Figure 6. Ca^{2+} concentration-sensitivity of PHD execution. (A) Log-log plots of recording saline
[Ca^{2+}] vs. QC corrected for non-linear summation for WT, *OK371/+*, and *VGlut, OK371/+* conditions.
Across the range of [Ca^{2+}] examined, there is no significant difference in calcium cooperativity for
these conditions (Nonlinear Regression, $p = 0.91$). **(B)** Data plotted as in (A) but this time with a double
heterozygous *RyR^{E4340K}/+; itpr^{90B}/+* genetic background. Across the range of [Ca^{2+}] examined, there
is no significant difference in calcium cooperativity for these conditions (Nonlinear Regression, $p =
0.78$). **(C)** Raw data for mEPSPs (left); raw data for EPSPs (middle); raw data for QC (right). All data
are for the indicated NMJ genotypes in 0.2 mM [Ca^{2+}]; bars are averages and error bars are ± SEM.
For mEPSPs, * $p < 0.05$ and *** $p < 0.001$ by Student's T-Test, comparing PHD-challenged genotypes
vs. unchallenged genetic controls. For EPSPs and QC, * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ vs.
OK371/+; ## $p < 0.01$; EPSP and QC analyses done across multiple genotypes by one-way ANOVA
with Tukey's post-hoc.

Figure 7. Partial impairment of Ca$_V$2/Cacophony causes excess depression. (A) Crossing scheme
for generating larvae for electrophysiological recording. Male larvae were hemizygous for the *cac^S*
hypomorphic mutation. **(B)** Raw data for mEPSPs. **(C)** Raw data for EPSPs. **(D)** Raw data for QC. For
(B)-(D), bars are averages and error bars are ± SEM. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ by
Student's T-Test comparing the control *cac^S* dataset (no *VGlut* overexpression) vs. the experimental
cac^S dataset (*VGlut* overexpression).

Figure 8. Model for how multiple calcium sources tune the process of PHD. Under baseline con-
ditions, Ca$_V$2-type calcium channels contribute to synapse function, as may RyRs and IP$_3$Rs. Under
conditions inducing PHD, synaptic vesicles are enlarged, and QC is decreased, possibly through

regulation of sources of calcium. When PHD challenge is coupled with concomitant impairment of
RyR and IP$_3$R channels, evoked potentials are significantly diminished, due to excess PHD.
669

7   Conflict of Interest

671 The authors declare that the research was conducted in the absence of any commercial or financial
672 relationships that could be construed as a potential conflict of interest.

8   Author Contributions

674 C.J.Y. and C.A.F. both did the following: designed research, performed research, analysed data, and
675 wrote and edited the paper.

9   Funding

677 Funding supporting this work includes an NSF Grant (1557792) and an NIH/NINDS Grant
678 (R01NS085164) to C.A.F. C.J.Y. was supported in part by an NIH/NINDS Predoctoral Training Grant
679 to the University of Iowa (UI) Interdisciplinary Graduate Program in Neuroscience (T32NS007421
680 – PI Daniel T. Tranel), as well as a post-comprehensive exam predoctoral summer fellowship and a
681 Ballard and Seashore Dissertation fellowship via the Graduate College at UI.

10   Acknowledgments

683 We thank members of the Frank lab for helpful comments. We thank the laboratories of Drs. Tina
684 Tootle, Fang Lin, Toshihiro Kitamoto, Pamela Geyer, and Lori Wallrath for helpful discussions. We
685 also thank Drs. Toshihiro Kitamoto, Joshua Weiner, Christopher Stipp, and Mark Stamnes for helpful
686 feedback on an earlier written version of this study.

11   Supplementary Material

688 Please see Supplementary Table S1 for raw data from the electrophysiology screen and a legend ex-
689 plaining the table.

A

w; UAS-VGlut, OK371-Gal4 ♀ X *mut* OR *UAS-gene[RNAi]* ♂

examine progeny by
electrophysiology

B

C

Screened Genotype in a *UAS-VGlut, OK371-Gal4/+* background

A

% *OK371/+* Driver Control

- ■ mEPSP
- ▨ EPSP
- □ QC

OK371/+ — *OK371/RyR^E4340K itpr^90B/+* — *VGlut, OK371/RyR^E4340K itpr^90B/+*

B

mEPSP (mV)

OK371/+ — *VGlut, OK371/+* — *OK371/ RyR^E4340K itpr^90B/+* — *VGlut, OK371/ RyR^E4340K itpr^90B/+*

*** — **

C

EPSP (mV)

OK371/+ — *VGlut, OK371/+* — *OK371/ RyR^E4340K itpr^90B/+* — *VGlut, OK371/ RyR^E4340K itpr^90B/+*

*** — ***

D

QC

OK371/+ — *VGlut, OK371/+* — *OK371/ RyR^E4340K itpr^90B/+* — *VGlut, OK371/ RyR^E4340K itpr^90B/+*

*** — ***

E

S.D. 1.22 mV
Range 5.20 mV

S.D. 1.89 mV
Range 8.01 mV

S.D. 1.46 mV
Range 5.20 mV

S.D. 2.38 mV *
Range 10.36 mV **

OK371/+ — *VGlut, OK371/+* — *OK371/RyR^E4340K itpr^90B/+* — *VGlut, OK371/RyR^E4340K itpr^90B/+*

WT

ER

Ca²⁺

RyR

IP₃R

Ca²⁺
Channel

GluRs

Evoked potentials normal

Homeostatic Depression

Evoked potentials ≈ normal

**Homeostatic Depression
+ *RyR itpr* Impairment**

Evoked potentials small