Nonameric structures of the cytoplasmic domain of FlhA and SctV in the context of the full-length protein.

Lucas Kuhlen [1,+], Steven Johnson[1], Jerry Y. Cao[1], Justin C. Deme[1,2], Susan M. Lea[1,2,*]

1. Sir William Dunn School of Pathology, South Parks Road, Oxford OX1 3RE, UK
2. Central Oxford Structural Molecular Imaging Centre, South Parks Road, Oxford OX1 3RE, UK
   + current address Imperial College, South Kensington Campus, London SW7 2AZ
   *correspondence to susan.lea@path.ox.ac.uk

Abstract

Type three secretion is the mechanism of protein secretion found in bacterial flagella and injectisomes. At its centre is the export apparatus (EA), a complex of five membrane proteins through which secretion substrates pass the inner membrane. While the complex formed by four of the EA proteins has been well characterised structurally, little is known about the structure of the membrane domain of the largest subunit, FlhA in flagella, SctV in injectisomes. Furthermore, FlhA/SctV is most often studied as a monomer and only a single structure of an SctV homologue assembled into the biologically relevant nonameric ring is available. FlhA has been shown to bind to chaperone-substrate complexes in an open state, but in the assembled ring structure SctV is in a closed state. Here, we identify FlhA and SctV homologues that can be recombinantly produced in the oligomeric state and study them using cryo-electron microscopy. The structures of the cytoplasmic domains from both FlhA and SctV are in the open state and we observe a conserved interaction between a short stretch of residues at the N-terminus of the cytoplasmic domain, known as $FlhA_L$/$SctV_L$, with a groove on the adjacent protomer's cytoplasmic domain, which stabilises the nonameric ring assembly. This represents the first structure of SctV in the open state, the first observation of the $SctV_L$ interaction with the adjacent protomer and confirms the importance of $FlhA_L$ for the stability of the FlhA nonameric ring.

Importance

Bacterial flagella are assembled from proteins secreted through a type III secretion system. A related type III secretion system is found in injectisomes, molecular syringes that bridge three membranes to secrete proteins directly from the bacterial cytoplasm into eukaryotic host cells. The major protein of the export apparatus of type III secretion is made up of a membrane and a cytoplasmic domain, which in the flagellar system can adopt an open or a closed state, is known to form a nonameric ring *in vivo*. We produced the full-length proteins from both injectisome and flagellar systems in the assembled state. The structures of the cytoplasmic domains demonstrate the conserved principle of the N-terminus of one subunit binding the membrane proximal face of the adjacent subunit to stabilise the assembled ring. Our structure of the homologue from the injectisome also demonstrates that the open state of the cytoplasmic domain is not unique to flagella.

Introduction

Type III secretion systems (T3SS) are bacterial macromolecular machines that facilitate transport of protein substrates across the bacterial cell envelope (Erhardt et al., 2010;

Wagner et al., 2018). The secretion machinery is conserved across at least two biological systems, the bacterial flagellum, which drives motility, and the virulence associated T3SS, which is an essential virulence factor for many bacterial pathogens due to the secretion of effector proteins directly into host cells (Galan, 2009). Most building blocks of the virulence associated T3SS or injectisome have a homologue in flagella and one of the most highly conserved parts of the system is a set of transmembrane proteins known collectively as the export apparatus (EA) (Deng et al., 2017).

The EA is made up of a core complex formed by the membrane proteins FliPQR (flagellar T3SS) or SctRST (virulence associated T3SS) which form a channel through which the unfolded protein substrates pass (Dietsche et al., 2016; Kuhlen et al., 2018). The regulatory subunit FlhB/SctU wraps around this core (Kuhlen et al., 2020) and the inner rod and needle assemble onto its periplasmic face (Hu et al., 2019; Torres-Vargas et al., 2019). Surprisingly, following assembly into the full T3SS, the core complex is not in the plane of the inner membrane. Instead, it is held in the plane of the periplasm by interactions with the surrounding basal body and its hydrophobic surface is covered by another membrane protein complex situated directly underneath it in the inner membrane. This protein complex is believed to be formed by the transmembrane domain of the fifth EA component, FlhA or SctV (Butan et al., 2019; Kuhlen et al., 2018). The FlhA/SctV cytoplasmic domain is known to form a nonameric ring underneath FliPQR-FlhB/SctRSTU (Abrusci et al., 2013), but the full-length FlhA/SctV complex remains poorly characterised. The entire EA is housed within the basal body of the T3SS and is thought serve as a nucleus for its assembly (Fan et al., 1997; Wagner et al., 2010).

FlhA/SctV has long been known to be the major component of the EA, but its stoichiometry was controversial for a long time (Li and Sourjik, 2011; Lilic et al., 2010) and may be dynamic *in vivo* (Li and Sourjik, 2011; Morimoto et al., 2014). Fluorescence based measurements showed that FlhA can form a large complex, but its exact stoichiometry couldn't be measured with high precision (Li and Sourjik, 2011; Morimoto et al., 2014). The crystallisation of the cytoplasmic domain of SctV from *Shigella flexneri*, $SctV_{Sf}$, as a nonameric ring matching the dimensions of a toroidal density in the cytoplasm, described by tomography, established the stoichiometry of FlhA/SctV (Abrusci et al., 2013) and the ability of the cytoplasmic domain to form a nonamer. This was later confirmed by high resolution AFM as the FlhA cytoplasmic domain also formed a nonameric ring on mica (Terahara et al., 2018) whose stability depended on the linker between cytoplasmic and membrane domains, $FlhA_L$/$SctV_L$. However, direct structural characterisation of the nonamer has so far been limited to the crystal structure of $SctV_{Sf}$, in which the crucial $SctV_L$ is not ordered, and full-length protein has been challenging to produce in an assembled state, precluding final confirmation that the full-length protein also forms a nonameric ring.

The lack of assembled ring structures of the FlhA/SctV cytoplasmic domain has also limited our understanding of the structural transitions this domain is thought to undergo in the secretion process. An open and closed state have been described, with the open state being competent for chaperone-substrate complex binding (Xing et al., 2018) while the closed state has been proposed to be responsible for binding early substrates (Inoue et al., 2019), but the open state has so far only been observed in protein that is not assembled into a ring. An interaction between $FlhA_L$/$SctV_L$ and the cytoplasmic domain of the neighbouring

subunit has been observed in crystal structures of the open state (Bange et al., 2010; Saijo-Hamano et al., 2010), but in the only structure of the assembled nonamer this interaction isn't seen. A structure of the assembled ring in the open state is needed to confirm observations that have been made with the monomeric cytoplasmic domain.

The transmembrane domain of FlhA/SctV is less well understood than the cytoplasmic domain. It is thought to contain approximately 8 transmembrane helices and conduct protons in order to use the pmf to power type III secretion (Minamino et al., 2016). Consistent with this function, the transmembrane domain contains a number of conserved charged residues essential for function (Erhardt et al., 2017). Structural studies of the membrane domain have been hampered by the difficulties in producing the full-length membrane protein in the assembled state. In addition to the predicted transmembrane helices the membrane domain also contains a short stretch of highly conserved soluble residues known as the FHIPEP domain (McMurry et al., 2004). It is known to be crucial for protein secretion, but its precise role in the secretion process remains to be elucidated (Barker et al., 2016; Erhardt et al., 2017).

We produced assembled rings of full-length FlhA/SctV from both flagellar and virulence associated systems and studied them using cryo-EM. Here, we present the structures of the nonameric cytoplasmic domain of both FlhA and SctV, the first structures of the assembled nonamer in the open state. The structures demonstrate the conserved interaction between FlhA$_L$/SctV$_L$ and the neighbouring subunit and further confirm the conserved stoichiometry of the complex.

Results

We screened a number of homologues of FlhA/SctV from many species for assembly in the *E. coli* membrane by employing GFP fusions and assaying fluorescent spot formation in the cell envelope, indicating membrane protein complex formation (Diepold et al., 2011; Li and Sourjik, 2011; Morimoto et al., 2014). FlhA from the lateral flagella of Vibrio parahaemolyticus (FlhA$_{Vp}$) and SctV from Yersinia enterocolitica (SctV$_{Ye}$) were found to assemble in *E. coli* (Fig 1a) and could be produced in large quantities at high purity (Fig 1b) using the gentle detergent LMNG. When the complex was subjected to gel filtration, a high molecular weight complex was isolated (Fig 1c).

Figure 1 *FlhA from Vibrio parahaemolyticus and SctV from Yersinia enterocolitica assemble into a large complex in the membrane of E. coli.*
a. Fluorescence images of *E. coli* cells expressing GFP tagged FlhA_{Vp} or SctV_{Ye} in the cell envelope. **b.** SDS-PAGE of purified FlhA_{Vp} and SctV_{Ye} following cleavage of GFP. **c.** Gel filtration traces of purified FlhA_{Vp} and SctV_{Ye}. The grey shaded area indicates fractions collected to make cryoEM grids.

We imaged the purified FlhA_{Vp} and SctV_{Ye} complexes using cryo-EM and analysed them using single particle analysis in Relion (Zivanov et al., 2018). Particle classification revealed that surprisingly, SctV_{Ye} had formed a dimer of nonamers and we imposed D9 symmetry for 3D refinement, resulting in a 3.7 Å reconstruction of the cytoplasmic domain double nonamer (Fig 2a). No high-resolution features of the transmembrane domain could be discerned. The SctV_{Ye} cytoplasmic domain 18-mer is made up of two nonamers dimerising via the cytoplasmic face of the cytoplasmic domain, an interaction which is not consistent with tomographic studies of the injectisome (Hu et al., 2017). The 18-mer is most likely formed by a nonphysiological association of two nonamers, perhaps due to the absence of other proteins such as YscX and YscY that are known to bind to SctV_{Ye} in the native T3SS (Diepold et al., 2012).

In contrast, FlhA_{Vp} was found to form the expected nonameric complex and we imposed C9 symmetry for particle refinement, resulting in a 3.8 Å volume of the cytoplasmic domain but little detail in the transmembrane domain (Fig 2b). Both reconstructions were of sufficient quality to allow us to build a model of the structure of the cytoplasmic domain.

Figure 2 *Cryo-EM volumes of the cytoplasmic domain of SctV and FlhA.*
a. 3D reconstruction of SctV_{Ye} at a resolution of 3.7 Å using D9 symmetry. **b.** 3D reconstruction of FlhA_{Vp} at a resolution of 3.8 Å using C9 symmetry. **c.** Model of the SctV_{Ye} linker region in the cryo-EM volume (orange) in the context of the adjacent protomer (grey).

d. Model of the FlhA$_{Vp}$ linker region in the cryo-EM volume (blue) in the context of the adjacent protomer (grey).

A striking feature of both structures is the N-terminal density of the cytoplasmic domain, formed by FlhA$_L$/SctV$_L$, which reaches out to bind to the cytoplasmic face of the adjacent protomer (Fig 2c), demonstrating this well studied interaction for the first time in the context of the full-length protein in both flagellar and injectisome T3SS. Biochemical studies have indicated that the interaction between FlhA$_L$ and the neighbouring subunit is important for assembly of the nonameric ring (Terahara et al., 2018), however, the interaction has so far only been observed in crystal structures of monomeric protein (Bange et al., 2010; Saijo-Hamano et al., 2010) and SctV$_L$ is not ordered in the crystal structure of the assembled SctV$_{Sf}$ nonamer (Abrusci et al., 2013). In both of our structures a hydrophobic residue, Trp350 in FlhA$_{Vp}$ and Phe367 in SctV$_{Ye}$, sticks into a hydrophobic groove on the neighbouring subunit. This residue in FlhA has previously been shown to be involved in ring formation of the cytoplasmic domain of FlhA (Terahara et al., 2018).

The overall dimensions of the nonameric rings are similar to that of the crystal structure of SctV$_{Sf}$ (Fig 3a), but the individual subunits are in a different state. FlhA/SctV is known to be able to adopt at least two conformations, open and closed, and FlhA$_L$/SctV$_L$ is thought to bind to the membrane proximal face of the neighbouring subunit only in the open state (Inoue et al., 2019). Comparison of our structures with FlhA from *S.* Typhimurium, FlhA$_{ST}$, in the open state and SctV$_{Sf}$, which is in the closed state shows that both of our complexes are in the open state (Fig 3b), with SctV$_{Ye}$ being the first example of an SctV homologue in the open state. The demonstration of the open state opens up the possibility that SctV binds to substrate-chaperone complexes in an analogous manner to what has been described for FlhA (Xing et al., 2018), in which chaperones bind between SD4 and SD2 and are sterically unable to access the binding site in the closed state.

A conformational change from open to closed state would be expected to lead to further changes throughout the protein. We simulated the movement of FlhA$_L$ by overlaying one subunit of FlhA$_{Vp}$ onto the SD3 or SD2 subdomain of the closed SctV$_{Sf}$ structure (Fig 3c). This demonstrates that adoption of the closed state in one subunit would affect the neighbouring subunit, potentially propagating the conformational change in a wave around the ring. An additional effect on the transmembrane domain is possible.

Figure 3 *The cryo-EM structures of SctV$_{Ye}$ and FlhA$_{Vp}$ are in the open state.*
a. Overlay of the nonameric rings of the cytoplasmic domains of SctV$_{Ye}$ and FlhA$_{Vp}$ with the crystal structure of the SctV$_{Sf}$ nonameric ring (PDB: 4A5P). **b.** Overlay of a single subunit of SctV$_{Ye}$ and FlhA$_{Vp}$ with FlhA$_{ST}$ in the open state (grey, PDB: 3a5i) and SctV$_{Sf}$ in the closed state (red, PDB: 4a5p). **c.** Overlay of FlhA$_{Vp}$ (light blue) on the SD3 domain of SctV$_{Sf}$ (red) and overlay (dark blue) on the SD1 and SD2 domains of SctV$_{Sf}$ illustrating the movement of the linker (arrow) as the protein changes from the open to the closed state.

Discussion

Full-length FlhA/SctV forms a very fragile, detergent sensitive membrane protein complex. Here, we have identified particularly stable FlhA/SctV sequences and purification conditions that have allowed preparation of the assembled ring complex for structural studies. We have determined the first nonameric structures of the cytoplasmic domain of FlhA/SctV in the open state. Multiple studies have previously implicated the FlhA linker region, FlhA$_L$, in promoting complex formation. We provide the first structural view of this interaction in the assembled state in both FlhA and SctV, showing that SctV$_L$ is the functional equivalent of FlhA$_L$ in the stabilisation of the nonameric assembly.

It has been proposed that the closed state of the FlhA/SctV cytoplasmic domain interacts with early secretion substrates, while later substrates bind to the open state (Inoue et al., 2019) which is induced by completion of the first step of filament assembly. Both of our structures are in the open state. Our structures are not affected by crystallisation artefacts and are in the context of the full-length protein, suggesting that the open state is the default state in the absence of other factors. Our structures also suggest that at least in the open state the residues known as FlhA$_L$/SctV$_L$, which have previously been described as the linker between cytoplasmic and membrane domains, do not form part of this linker but are

part of the cytoplasmic domain. Instead, the residues N-terminal to FlhA$_L$/SctV$_L$ must be responsible for linking the membrane domain to the cytoplasmic domain.

Unfortunately, we were not able to determine the structure of the membrane domain of either complex, possibly due to damage to the fragile membrane domain at the air-water interface during cryo-EM sample preparation (D'Imprima et al., 2019). However, two features are clear in 2D averages of the side view of both complexes. The cytoplasmic and membrane domains are at a set distance from each other and this distance matches that observed in tomographic reconstructions of the T3SS (Fig 4a). This suggests that the linker between the domains is not flexible, which would be inconsistent with a previously suggested model of secretion according to which the cytoplasmic domain is tethered to the membrane by a flexible linker and moves towards the membrane domain and back into the cytoplasm in a cyclical fashion (Erhardt et al., 2017).

The methods developed here may be the basis of future studies of the FlhA/SctV transmembrane domain or interactions of the cytoplasmic domain with chaperones and substrates in the context of the assembled complex.

Figure 4 *Position of the membrane domain of FlhA in the T3SS nanomachine.*
a. The volume of FlhA$_{Vp}$ is shown in red overlaid onto the tomographic reconstruction of the injectisome (EMD-8544). **b.** Selected 2D class averages of FlhA$_{Vp}$ show the high level of detail in the cytoplasmic domain and the lower information content in the presumed transmembrane, micelle-embedded, portion.

Methods

Protein purification
FlhA and SctV were produced as GFP fusion proteins in *E. coli* BL21 by expressing them from a pt12 plasmid (Dietsche et al., 2016). Cells were grown in terrific broth supplemented with rhamnose monohydrate (0.1%) and harvested after overnight growth. Cells were

resuspended in TBS (100 mM Tris, 150 mM NaCl, 1 mM EDTA) and lysed using an Emulsiflex homogeniser (Avestin). Membranes were purified by ultracentrifugation of the clarified lysate at 235,000g and solubilised in 1% LMNG using 0.1 grams of detergent per gram of membrane pellet. After gentle stirring for one hour aggregates were removed by centrifugation at 75,600g and the solubilised membrane proteins were applied to a StrepTrap column (GE healthcare) which was then washed in TBS supplemented with 0.01% LMNG. Pure protein was eluted in TBS containing 0.01% LMNG and 10 mM desthiobiotin. GFP was cleaved off by incubating with tev protease overnight. Next, the protein was concentrated and further purified by gel filtration using a superpose 6 increase (GE healthcare) in TBS containing 0.01% LMNG. The peak fraction, eluting close to 12 ml, was collected.

Cryo-EM analysis
3 µl of purified protein at a concentration between 0.8 and 1 mg/ml was deposited on a glow-discharged Quantifoil grid (2/2, 300 mesh for FlhA, 1.2/1.3, 300 mesh for SctV). Grids were blotted and plunged into liquid ethane using a Vitrobot Mark IV (FEI). The samples were imaged using a Titan Krios (FEI) equipped with a K2 detector (Gatan). Relion's implementation of MotionCor2 (Zivanov et al., 2019) was used for motion correction and SIMPLE's implementation of CTFFIND4 was used for CTF estimation (Reboul et al., 2018). Particles were picked using SIMPLE and classified in Relion3.0 (Zivanov et al., 2018). Atomic models were built in Coot (Emsley et al., 2010) and refined using phenix (Afonine et al., 2018).

Fluorescence imaging
8 µl of an overnight culture of *E. coli* BL21 expressing FlhA-GFP or SctV-GFP were applied to a glass slide and imaged using a Zeiss 880 inverted microscope equipped with a plan-apochromat 63x/1.4 NA objective and an Airyscan detector. GFP fluorescence was excited using a laser (488 nm).

Acknowledgements
We thank the staff of the Central Oxford Structural Microscopy and Imaging Centre Errin Johson and Adam Costin, Alan Wainman of the Dunn School Light Microscopy Facility and all members of the Lea group for assistance. The work was funded by grants from the Wellcome Trust 109136, 100298, 219477, the Medical Research Council (UK) S021264 and the Wolfson Foundation WL160052.

Data Availability
Coordinates and associated volumes and metadata have been deposited in the PDB and EMDB respectively. SctV EMD-11820 / PDB 7ALW, FlhA EMD-11827 / PDB 7AMY.

References
Abrusci, P., Vergara-Irigaray, M., Johnson, S., Beeby, M.D., Hendrixson, D.R., Roversi, P., Friede, M.E., Deane, J.E., Jensen, G.J., Tang, C.M.*, et al.* (2013). Architecture of the major component of the type III secretion system export apparatus. Nat Struct Mol Biol *20*, 99-104.

Afonine, P.V., Poon, B.K., Read, R.J., Sobolev, O.V., Terwilliger, T.C., Urzhumtsev, A., and Adams, P.D. (2018). Real-space refinement in PHENIX for cryo-EM and crystallography. Acta Crystallogr D Struct Biol *74*, 531-544.

Bange, G., Kummerer, N., Engel, C., Bozkurt, G., Wild, K., and Sinning, I. (2010). FlhA provides the adaptor for coordinated delivery of late flagella building blocks to the type III secretion system. Proc Natl Acad Sci U S A *107*, 11295-11300.

Barker, C.S., Inoue, T., Meshcheryakova, I.V., Kitanobo, S., and Samatey, F.A. (2016). Function of the conserved FHIPEP domain of the flagellar type III export apparatus, protein FlhA. Mol Microbiol *100*, 278-288.

Butan, C., Lara-Tejero, M., Li, W., Liu, J., and Galan, J.E. (2019). High-resolution view of the type III secretion export apparatus in situ reveals membrane remodeling and a secretion pathway. Proc Natl Acad Sci U S A *116*, 24786-24795.

D'Imprima, E., Floris, D., Joppe, M., Sanchez, R., Grininger, M., and Kuhlbrandt, W. (2019). Protein denaturation at the air-water interface and how to prevent it. Elife *8*.

Deng, W., Marshall, N.C., Rowland, J.L., McCoy, J.M., Worrall, L.J., Santos, A.S., Strynadka, N.C.J., and Finlay, B.B. (2017). Assembly, structure, function and regulation of type III secretion systems. Nat Rev Microbiol *15*, 323-337.

Diepold, A., Wiesand, U., Amstutz, M., and Cornelis, G.R. (2012). Assembly of the Yersinia injectisome: the missing pieces. Mol Microbiol *85*, 878-892.

Diepold, A., Wiesand, U., and Cornelis, G.R. (2011). The assembly of the export apparatus (YscR,S,T,U,V) of the Yersinia type III secretion apparatus occurs independently of other structural components and involves the formation of an YscV oligomer. Mol Microbiol *82*, 502-514.

Dietsche, T., Tesfazgi Mebrhatu, M., Brunner, M.J., Abrusci, P., Yan, J., Franz-Wachtel, M., Scharfe, C., Zilkenat, S., Grin, I., Galan, J.E.*, et al.* (2016). Structural and Functional Characterization of the Bacterial Type III Secretion Export Apparatus. PLoS Pathog *12*, e1006071.

Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and development of Coot. Acta Crystallogr D Biol Crystallogr *66*, 486-501.

Erhardt, M., Namba, K., and Hughes, K.T. (2010). Bacterial nanomachines: the flagellum and type III injectisome. Cold Spring Harb Perspect Biol *2*, a000299.

Erhardt, M., Wheatley, P., Kim, E.A., Hirano, T., Zhang, Y., Sarkar, M.K., Hughes, K.T., and Blair, D.F. (2017). Mechanism of type-III protein secretion: Regulation of FlhA conformation by a functionally critical charged-residue cluster. Mol Microbiol *104*, 234-249.

Fan, F., Ohnishi, K., Francis, N.R., and Macnab, R.M. (1997). The FliP and FliR proteins of Salmonella typhimurium, putative components of the type III flagellar export apparatus, are located in the flagellar basal body. Mol Microbiol *26*, 1035-1046.

Galan, J.E. (2009). Common themes in the design and function of bacterial effectors. Cell Host Microbe *5*, 571-579.

Hu, B., Lara-Tejero, M., Kong, Q., Galan, J.E., and Liu, J. (2017). In Situ Molecular Architecture of the Salmonella Type III Secretion Machine. Cell *168*, 1065-1074 e1010.

Hu, J., Worrall, L.J., Vuckovic, M., Hong, C., Deng, W., Atkinson, C.E., Brett Finlay, B., Yu, Z., and Strynadka, N.C.J. (2019). T3S injectisome needle complex structures in four distinct states reveal the basis of membrane coupling and assembly. Nat Microbiol *4*, 2010-2019.

Inoue, Y., Ogawa, Y., Kinoshita, M., Terahara, N., Shimada, M., Kodera, N., Ando, T., Namba, K., Kitao, A., Imada, K.*, et al.* (2019). Structural Insights into the Substrate Specificity Switch Mechanism of the Type III Protein Export Apparatus. Structure *27*, 965-976 e966.

Kuhlen, L., Abrusci, P., Johnson, S., Gault, J., Deme, J., Caesar, J., Dietsche, T., Mebrhatu, M.T., Ganief, T., Macek, B., *et al.* (2018). Structure of the core of the type III secretion system export apparatus. Nat Struct Mol Biol *25*, 583-590.

Kuhlen, L., Johnson, S., Zeitler, A., Baurle, S., Deme, J.C., Caesar, J.J.E., Debo, R., Fisher, J., Wagner, S., and Lea, S.M. (2020). The substrate specificity switch FlhB assembles onto the export gate to regulate type three secretion. Nat Commun *11*, 1296.

Li, H., and Sourjik, V. (2011). Assembly and stability of flagellar motor in Escherichia coli. Mol Microbiol *80*, 886-899.

Lilic, M., Quezada, C.M., and Stebbins, C.E. (2010). A conserved domain in type III secretion links the cytoplasmic domain of InvA to elements of the basal body. Acta Crystallogr D Biol Crystallogr *66*, 709-713.

McMurry, J.L., Van Arnam, J.S., Kihara, M., and Macnab, R.M. (2004). Analysis of the cytoplasmic domains of Salmonella FlhA and interactions with components of the flagellar export machinery. J Bacteriol *186*, 7586-7592.

Minamino, T., Morimoto, Y.V., Hara, N., Aldridge, P.D., and Namba, K. (2016). The Bacterial Flagellar Type III Export Gate Complex Is a Dual Fuel Engine That Can Use Both H+ and Na+ for Flagellar Protein Export. PLoS Pathog *12*, e1005495.

Morimoto, Y.V., Ito, M., Hiraoka, K.D., Che, Y.S., Bai, F., Kami-Ike, N., Namba, K., and Minamino, T. (2014). Assembly and stoichiometry of FliF and FlhA in Salmonella flagellar basal body. Mol Microbiol *91*, 1214-1226.

Reboul, C.F., Eager, M., Elmlund, D., and Elmlund, H. (2018). Single-particle cryo-EM-Improved ab initio 3D reconstruction with SIMPLE/PRIME. Protein Sci *27*, 51-61.

Saijo-Hamano, Y., Imada, K., Minamino, T., Kihara, M., Shimada, M., Kitao, A., and Namba, K. (2010). Structure of the cytoplasmic domain of FlhA and implication for flagellar type III protein export. Mol Microbiol *76*, 260-268.

Terahara, N., Inoue, Y., Kodera, N., Morimoto, Y.V., Uchihashi, T., Imada, K., Ando, T., Namba, K., and Minamino, T. (2018). Insight into structural remodeling of the FlhA ring responsible for bacterial flagellar type III protein export. Sci Adv *4*, eaao7054.

Torres-Vargas, C.E., Kronenberger, T., Roos, N., Dietsche, T., Poso, A., and Wagner, S. (2019). The inner rod of virulence-associated type III secretion systems constitutes a needle adapter of one helical turn that is deeply integrated into the system's export apparatus. Mol Microbiol *112*, 918-931.

Wagner, S., Grin, I., Malmsheimer, S., Singh, N., Torres-Vargas, C.E., and Westerhausen, S. (2018). Bacterial type III secretion systems: a complex device for the delivery of bacterial effector proteins into eukaryotic host cells. FEMS Microbiol Lett *365*.

Wagner, S., Konigsmaier, L., Lara-Tejero, M., Lefebre, M., Marlovits, T.C., and Galan, J.E. (2010). Organization and coordinated assembly of the type III secretion export apparatus. Proc Natl Acad Sci U S A *107*, 17745-17750.

Xing, Q., Shi, K., Portaliou, A., Rossi, P., Economou, A., and Kalodimos, C.G. (2018). Structures of chaperone-substrate complexes docked onto the export gate in a type III secretion system. Nat Commun *9*, 1773.

Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., and Scheres, S.H. (2018). New tools for automated high-resolution cryo-EM structure determination in RELION-3. Elife *7*.

Zivanov, J., Nakane, T., and Scheres, S.H.W. (2019). A Bayesian approach to beam-induced motion correction in cryo-EM single-particle analysis. IUCrJ *6*, 5-17.

	FlhA EMD- 11827 PDB 7AMY	YscV EMD-11820 PDB 7ALW
Data collection and processing		
Magnification	165,000	165,000
Voltage (kV)	300	300
Electron exposure (e−/Å2)	48	48
Defocus range (μm)	0.5–4	0.5–4
Pixel size (Å)	0.822	0.822
Symmetry imposed	C9	D9
Final particle images (no.)	9756	15913
Map resolution (Å)	3.75	3.7
FSC threshold	0.143	0.143
Refinement		
Initial model used	FlhA$_{ST}$ (3a5i)	SctV$_{Sf}$ (4a5p)
Model resolution (Å)	3.75	3.7
FSC threshold	0.143	0.143
Map sharpening B factor (Å^2)	-72	-58
Model composition		
Non-hydrogen atoms	24,849	50,148
Protein residues	3177	6192
Ligands	0	0
B factors (Å^2)		
Protein	64.56	57.65
R.m.s. deviations		
Bond lengths (Å)	0.0096	0.0099
Bond angles (°)	0.88	0.95
Validation		
MolProbity score	2.44	2.39
Clashscore	20.06	15.19
Poor rotamers (%)	0.66	0.98
Ramachandran plot		
Favoured (%)	85.94	82.37
Allowed (%)	13.78	17.33
Disallowed (%)	0.28	0.29

Table 1. CryoEM data collection, processing and model statistics