Massive genome decay and insertion sequence expansion drive the evolution of a novel host-restricted bacterial pathogen

Gonzalo Yebra[1,a], Andreas F Haag[2,a], Maan M Neamah[2,3,4], Bryan A Wee[1], Emily J Richardson[1], Pilar Horcajo[5], Sander Granneman[6], María Ángeles Tormo-Más[7,8], Ricardo de la Fuente[5], J Ross Fitzgerald[1,*], José R Penadés[2,7,9]*

[1]*The Roslin Institute, University of Edinburgh, Edinburgh, United Kingdom;* [2]*Institute of Infection, Immunity & Inflammation, University of Glasgow, Glasgow, United Kingdom;* [3]*Faculty of Veterinary Medicine, University of Kufa, Kufa, Iraq;* [4]*Middle Euphrates Centre for Cancer and Genetic Research, University of Kufa, Kufa, Iraq;* [5]*Facultad de Veterinaria, Universidad Complutense de Madrid, Madrid, Spain;* [6]*Centre for Synthetic and Systems Biology, University of Edinburgh, Edinburgh, United Kingdom;* [7]*Departamento de Ciencias Biomédicas, Facultad de Ciencias de la Salud, Universidad CEU Cardenal Herrera, 46113 Moncada, Valencia, Spain;* [8]*Severe Infection Group, Health Research Institute Hospital La Fe, Valencia, Spain;* [9]*MRC Centre for Molecular Bacteriology and Infection, Imperial College London, SW7 2AZ, UK*

[a] These authors contributed equally.

* Joint senior and corresponding authors.

Email addresses:

GY: Gonzalo.Yebra@ed.ac.uk; AH: Andreas.Haag@glasgow.ac.uk;

MMN: maanm.neamah@uokufa.edu.iq; BAW: B.Wee@ed.ac.uk;

EJR: e.j.richardson@bham.ac.uk; PH: phorcajo@ucm.es;

SG: Sander.Granneman@ed.ac.uk; MAT-M: tormo_man@gva.es;

R de la F: rifuente@vet.ucm.es; JRF: ross.fitzgerald@ed.ac.uk;

JRP: j.penades@imperial.ac.uk.

Abstract

Background. The emergence of new pathogens is a major threat to public and veterinary health. Changes in bacterial habitat such as those associated with a switch in host or disease tropism are often accompanied by genetic adaptation. *Staphylococcus aureus* is a multi-host bacterial species comprising strains with distinct tropisms for human and livestock species. A microaerophilic subspecies, *Staphylococcus aureus* subsp*. anaerobius*, is responsible for outbreaks of Morel's disease, a lymphadenitis in small ruminants. However, the evolutionary history of *S. aureus* subsp*. anaerobius* and its relatedness to *S. aureus* are unknown.

Results. Evolutionary genomic analyses of clinical *S. aureus* subsp*. anaerobius* isolates revealed a highly conserved clone that descended from a *S. aureus* progenitor about 1000 years ago before differentiating into distinct lineages representing African and European isolates. *S. aureus* subsp*. anaerobius* has undergone limited clonal expansion, with a restricted population size, and an evolutionary rate 10-fold slower than *S. aureus*. The transition to its current restricted ecological niche involved acquisition of a pathogenicity island encoding a ruminant host-specific effector of abscess formation, several large chromosomal re-arrangements, and the accumulation of at least 205 pseudogenes resulting in a highly fastidious metabolism. Importantly, expansion of ~87 insertion sequences (IS) located largely in intergenic regions provided distinct mechanisms for the control of expression of flanking genes, representing a novel concept of the IS regulon.

Conclusions. Our findings reveal the remarkable evolutionary trajectory of a host-restricted bacterial pathogen that resulted from extensive remodelling of the *S. aureus* genome through an array of parallel mechanisms.

Keywords: Bacterial evolution; host adaptation; host restriction; genome reduction; gene decay; insertion elements; pseudogenes; structural variation; gene expression regulation; phylodynamics.

Background

Bacteria have a remarkable capacity to adapt to new environmental niches, a characteristic that underpins their potential to become successful pathogens. Some pathogens can infect multiple host-species [1], while others become specialized and restricted to a single host. The evolutionary process of host restriction, observed across distantly-related bacterial groups, typically involves genomic events such as gene loss, gene acquisition, or chromosomal rearrangements [2-5]. The impact of such events may be most apparent after long-term associations between bacterium and host, with obligate intracellular pathogens the most extreme examples, exhibiting extremely small and compact genomes [6].

Staphylococcus aureus is a highly versatile bacterial pathogen, associated with an array of diseases in humans and livestock representing a threat to public and livestock health [7]. *S. aureus* has undergone extensive host-switching events during its evolutionary history leading to the emergence of new endemic and epidemic clones in livestock and humans [8]. In particular, the highest number of host-jump events appeared to have been between humans and ruminants in either direction [8]. A combination of gene acquisition, loss of gene function, and allelic diversification have been central to the capacity of *S. aureus* to undergo successful host-adaptation [8].

A close relative of *S. aureus*, *S. aureus* subsp. *anaerobius*, responsible for Morel's disease [9], is highly restricted to small ruminants and its clinical manifestation is confined to abscesses located at major superficial lymph nodes. Outbreaks of this disease are associated with significant economic losses and have been reported in Europe, Africa and the Middle East [10]. In contrast to *S. aureus* subsp. *aureus*, *S. aureus* subsp. *anaerobius* is microaerophilic and catalase-negative [11], but their genetic and evolutionary relatedness is poorly understood.

Here, we investigate the evolutionary history of *S. aureus* subsp. anaerobius, revealing that it descended from an ancestor resembling *S. aureus* and adapted to its current ecological niche by extensive genome diversification involving massive genome decay, gene

acquisition, and genome rearrangements. Furthermore, acquisition and expansion of

ISSau8-like insertion sequences associated with the manipulation of transcription of flanking

genes through multiple mechanisms leads us to propose the novel concept of the IS regulon.

Overall, our results reveal a remarkable example of a bacterial pathogen that has undergone

extensive genome remodelling in transition to a highly niche-restricted ecology.

Results

S. aureus **subsp.** *anaerobius* **has a highly conserved genome.**

Previously, it has been reported that *S. aureus* subsp. *anaerobius* isolates belong to a single

clonal complex as determined by MLST [10, 12]. To date only one fragmented draft genome

of *S. aureus* subsp. *anaerobius* isolated in Sudan has been publicly available, and therefore

the genome architecture has been unclear [13]. Here, we applied PacBio sequencing to the

type strain MVF7, isolated in Spain in 1981-1982 [11], to provide the first complete genome

sequence for a strain of *S. aureus* subsp. *anaerobius*. The genome consisted of a circular

chromosome of 2,755,024 bp with 2,888 coding DNA sequences (CDS), 56 tRNAs and 5

complete rRNA copies, with a GC content of 32.74% (**Fig. 1A**).

In order to further explore the genome content of *S. aureus* subsp. *anaerobius*, we obtained

genomic DNA for 39 additional strains and performed Illumina whole genome sequencing to

examine genome diversity and population genetic structure. Of these strains, 30 were

isolated from different outbreaks in Spain between 1981 and 2012, with the others from

Sudan (n=3), Italy (n=3), Poland (n=2) and Denmark (n=1) [10]. Among the isolates, we

identified 4 closely-related novel STs (see **Additional Information** in Additional File 1 for

more detail).

The combined dataset of 41 genomes had an average length of 2,685,366 bp (range:

2,604,446 to 2,758,945 bp) that contained 2,479 genes (92.7%) shared among all isolates

(core-genome) of the total 2,675 genes identified (pan-genome). On average, each strain

has a highly compact accessory genome of only 130 genes (range 107-143).

S. aureus subsp. anaerobius evolved from an ancestor that resembled S. aureus subsp. aureus over a millennium ago.

The evolutionary origin of *S. aureus* subsp. *anaerobius* and its phylogenetic relatedness to *S. aureus* subsp. *aureus* is unknown. In order to investigate its evolutionary history, we constructed a core genome alignment of *S. aureus* subsp. *anaerobius* and 807 *S. aureus* isolates that were representative of the species diversity [8] and reconstructed a maximum-likelihood phylogeny (**Fig. 2**). *S. aureus* subsp. *anaerobius* formed a distinct clade on a long branch within the *S. aureus* subsp. *aureus* diversity but was not genetically allied to any particular clonal complex. These data indicate that *S. aureus* subsp. *anaerobius* emerged from an unknown *S. aureus* lineage likely after a human to ruminant host-switch event that occurred over a millennium ago.

A core genome sequence alignment of the 41 isolates was produced with 5 recombinant regions (size range: 54 bp to 2,950 bp) excluded leaving an alignment of 2,324,010 sites with 3,443 variable sites. Phylogenetic analyses by the maximum likelihood approach revealed two distinct clades among the *S. aureus* subsp. *anaerobius* examined representing the isolates sampled in Sudan, and all others (sampled in Europe), respectively (**Fig. 3**). The average number of SNPs between pairs of *S. aureus* subsp. *anaerobius* genomes was 494 SNPs (range: 0-1,746), and the number of SNPs between the two clades was 1,697 SNPs (1,663-1,746).

Linear regression analysis provided evidence of a molecular clock-like evolution in the ML phylogeny (**Fig. S1** in Additional File 1) and time-scaled Bayesian phylogenetic analysis indicated that the model combination that best fitted the data was a strict molecular clock paired with a constant population size, though other models performed similarly (data not shown). This analysis dated the most recent common ancestor (MRCA) of *S. aureus* subsp. *anaerobius* in 943 years before present (YBP) (95% Bayesian confidence interval (BCI): 1,184-716) (**Fig. 3**). The European and Sudanese clades diverged between 235 YBP (95% BCI: 296-180) and 82 YBP (95% BCI: 105-61), respectively. The estimated

substitution rate was 4.4 (95% BCI: 3.3-5.5) $\times$ 10^{-7} substitutions/site/year (s/s/y),

approximating to 1.2 SNPs per year which is ~one order of magnitude slower than *S. aureus*

subsp*. aureus* and may reflect the relatively slow growth observed for *S. aureus*

subsp. *anaerobius* [9, 14].

S. aureus subsp. *anaerobius* has undergone large intra-chromosomal

rearrangements.

During the evolutionary genomic analysis, we discovered that *S. aureus* subsp. *anaerobius*

has undergone six large genomic translocations that ranged in size between 70 kb and

346 kb since separation from its *S. aureus* subsp. *aureus* progenitor (**Fig. 4** and **Fig. S2** in

Additional File 1). As the edges of each translocated region were flanked by identical

insertion sequence elements, we suggest that each event was the result of transposase-

mediated homologous recombination between the leading and lagging strands [15]. For 2

pairs of translocations (I, VI, and II, V, respectively) the original genomic locations were

exchanged whereas translocations III and IV were the result of excision and subsequent

insertion at a distinct genomic location (**Fig. 4**). Because all translocations occurred via an

inversion event, genes in the translocated areas conserved their original orientation and the

GC skew remained unaffected (**Fig. 4**).

PCR-based analysis of the distribution of the 6 rearrangements identified in strain MVF7

among all *S. aureus* subsp. *anaerobius* strains revealed that all 6 were conserved in the

majority of European isolates (clade II) whereas the isolates from Sudan (clade I) lacked

translocations numbers II and V, suggesting they occurred in the clade I lineage since

separation from a common ancestor (**Table S1** in Additional File 2). Although the impact of

the re-arrangements on global gene expression is unclear, large chromosomal

rearrangements in *S. aureus* subsp. *aureus* have been demonstrated to mediate transition to

less virulent strains associated with persistent infection [16, 17].

S. aureus subsp. *anaerobius* exhibits massive genome decay.

Analysis of the complete genome of *S. aureus* subsp. *anaerobius* strain MVF7 revealed the

existence of 205 pseudogenes, with a range of 201 to 210 pseudogenes per genome across

the 41 *S. aureus* subsp. *anaerobius* strains examined (see **Additional Information** in

Additional File 1). Pseudogenes originated through point mutations that caused frameshifts,

premature stop codons or alternative downstream start codons and were evenly distributed

around the genome (**Fig. 1A**). Relative to *S. aureus* subsp*. aureus*, (which contains

approximately 20-50 pseudogenes per strain [18-20]), the high frequency of pseudogenes

representing on average of 9.5% (range: 9.2-10.1%) of the total genome length represents a

remarkable example of extensive gene loss that is likely to have a major impact on bacterial

phenotype. A total of 164 pseudogenes (approximately 80% of those found in any given

isolate) were shared among all *S. aureus* subsp. *anaerobius* genomes, of which 92.1% were

caused by the same mutation in all isolates consistent with early events in the evolutionary

history of *S. aureus* subsp. *anaerobius*.

The original functions of the 205 pseudogenes present in MVF7 were classified into clusters

of orthologous groups (COGs) of proteins (see **Table S2** in Additional File 2). Of these, 82

(50%) were associated with metabolic functions, 29 (17.7%) to cellular processes and

signalling, and 10 (6.1%) to information storage and processing, and 43 (26.2%) had

unknown functions. Enrichment analyses of both KEGG pathways and GO terms (see **Table**

S3 in Additional File 2 for a full list) revealed a statistically significant enrichment of

pseudogenes involved in biosynthesis of amino acids and metabolic pathways (see

Additional Information in Additional File 1 and **Table S2** in Additional File 2). Of note, loss

of function of genes mediating resistance to oxidative killing mechanisms such as catalase

and other oxidoreductases may underpin the micro-aerophilic growth phenotype. In addition,

an array of genes with known roles in pathogenicity including capsule biosynthesis,

adherence and secreted proteins mediating interactions with the host, are predicted to be

non-functional. Taken together, the extensive loss of gene function impacting on bacterial

metabolism and pathogenicity is consistent with the highly fastidious growth requirements

and defined disease tropism of *S. aureus* subsp. *anaerobius* (**Fig. 1C**).

***S. aureus* subsp. *anaerobius* contains a novel pathogenicity island encoding a**

ruminant specific effector of abscess formation.

Our genomic analysis revealed a small accessory genome with limited strain-dependent

variation in gene content. However, 2 putative MGE were identified in all strains examined

including a 43.2 kb, novel prophage (ΦSaa1) belonging to the *Siphoviridae* family, and a

novel 13 kb *S. aureus* subsp. *anaerobius* pathogenicity island (SaaPIMVF7) (**Fig. 1B**). While

ΦSaa1 contained no putative determinants of virulence, SaaPIMVF7 encodes novel variants

of known virulence factors Staphylococcal complement inhibitor (SCIN) and the von

Willebrand factor-binding protein (vWbp) (**Fig. 1B**) [18, 21]. A phylogenetic network of all

members of the SaPI family constructed with SplitsTree v4.14.6 and the NeighborNet

algorithm [22] (**Fig. S3** in Additional File 1) revealed that SaaPIMVF7 is genetically closest

(93.1% nucleotide identity) to SaPIov2, found among isolates of *S. aureus* subsp. *aureus*

CC133 which has a tropism for small ruminants (**Fig. 1B**). Of note, SaaPIMVF7 contained 10

pseudogenes out of a total of 21 genes identified, including the integrase and primase

genes, suggesting that it can no longer be mobilized but is a stable feature of the

chromosome. Of note, both the *vwb* and *scn* genes were intact and transcribed (data not

shown), consistent with functionality (**Fig. 1C**). The vWBP has previously been

demonstrated to promote coagulation of plasma, and is involved with abscess formation

during invasive infection [23, 24]. In order to test the hypothesis that SaaPIMVF7-encoded

vWBP was functional, we cloned the *vwb* variant into the expression vector pCN51,

introduced this plasmid into a coagulase and vWbp-deficient derivative of strain RN4220

(RN4220 *coa*::*tet*M Δ*vwb*). Importantly, expression of the variant vWBP protein conferred the

capacity for coagulation of ruminant plasma (**Fig. 1B**). In summary, all strains of *S. aureus*

subsp. *anaerobius* contain a novel SaPI indicating an acquisition event that occurred over

1000 years ago prior to divergence of the 2 major subclades. Strikingly, the accumulation of

loss-of-function mutations in genes required for mobilization indicate that SaaPIMVF7 is a stable element of the genome and functional expression of vWBP with ruminant coagulation activity suggests a potential role in abscess formation, a defining characteristic of Morel's disease.

Expansion and intergenic location of insertion sequences in the genome of *S. aureus* subsp. *anaerobius.*

An additional striking feature of *S. aureus* subsp. *anaerobius* was the presence of at least 87 insertion sequences (IS) distributed around the genome in strain MVF7 (**Fig. 1A**). Each IS had 99.3% nucleotide identity with each other, and 97% identity with the previously described ISSau8 from bovine *S. aureus* strain RF122. The great majority (77 of 87) had premature stop codons that truncated the transposase gene whereas only 10 IS contained full-length intact transposase genes. The distribution of the IS elements identified in MVF7 among the rest of *S. aureus* subsp. *anaerobius* isolates was examined by mapping Illumina reads for each strain against the border between each IS and its flanking gene in the closed genome of strain MVF7 to identify reads overlapping this border. This analysis revealed that 68 out of the 87 MVF7 ISs (78.2%) were inserted at an identical location in all 41 *S. aureus* subsp. *anaerobius* isolates consistent with an ancient acquisition that occurred during the emergence of *S. aureus* subsp. *anaerobius*. An additional 7 IS elements were shared among all the European isolates, but were absent from isolates from Sudan (**Fig. 3**).

Previously, IS elements have been shown to contribute to loss of gene function or phase variation via direct insertion into CDS sequences [25]. However, all except one IS element were inserted in intergenic regions (the single exception was inserted into the *msb*A CDS), and we speculated that the non-random distribution of IS elements may influence the expression of the chromosomal genes located adjacent to the IS insertion site.

IS affects *neighbouring* gene regulation in *S. aureus* subsp*. anaerobius* by disruption of promoter region and operon structure.

Our *in silico* analysis of promoter and transcriptional terminator (TT) sequences indicated

that each IS had a functional promoter but lacked a TT between the *tnp* gene and the

downstream gene, supporting the idea that the *tnp* transcript could influence expression of

the 3'-located gene. Furthermore, 2 putative TT were identified at different sites in the IS:

TT1 is located in the 5' region of the IS genes, and is bidirectional, while TT2 is located at

the 3' end region of the IS and is unidirectional in the antisense orientation (see **Fig. 5**). The

putative role for the bidirectional TT1 would be to prevent the interference between

transcripts initiated either from the IS or from the gene located upstream to the IS. By

contrast, TT2 would prevent the impact that an antisense transcript originated from a gene

located downstream of the IS could have on the expression of the IS transcript (**Fig. 5**). This

organisation suggests a strategy that allows IS transcripts to interfere with the genes located

3' to the IS, while blocking any interference that transcripts from flanking genes could have

on the IS-derived regulatory transcript.

To validate the functionality and directionality of the TTs, we employed plasmid pCN42 [26],

in which the *bla*Z reporter is under the control of the P$_{cad}$ promoter. The conserved 5' and 3'

region of the ISs containing the bi- and unidirectional TTs, respectively, were inserted, in

both orientations, between the P$_{cad}$ promoter and the *bla*Z reporter (see **Fig. 5**), and the

expression of the reporter gene measured with and without induction of the P*cad* promotor.

The results confirm the existence, functionality and directionality of both TTs, consistent with

the hypothesis that ISs can control the expression of 3'-located chromosomal genes.

To test the idea that the intergenic IS insertions arise by selection because they influence

the expression of the neighbouring genes, we examined the impact of IS on the expression

of selected downstream genes located in the same orientation as the IS. Since *S. aureus*

subsp. *anaerobius* is not genetically tractable, we designed reporter constructs

reconstructing the genetic organisation found in *S. aureus* subsp. *anaerobius* (**Fig. S4** in

Additional File 1) and tested gene expression levels in *S. aureus* subsp. *aureus*. Specifically,

we selected four different genes (*rps*P, *dna*D, *met*C and *met*N) that were identified in

S. aureus subsp. *anaerobius* to contain an IS at distinct distances upstream of the gene start

codon, and constructed *bla*Z transcriptional reporter fusions using pCN41 [26] (**Fig. 6A-D**).

The expression of these genes in the presence or absence of upstream ISs was then tested.

Importantly, and in support of the hypothesis, absence of an IS impacted on the expression

of the neighbouring genes, confirming their regulatory role, although this effect was different

depending on the gene including increased transcription of *dna*D, *met*C or *met*N, or

decreased *rps*P transcription (**Fig. 6A-D**). The impact on downstream gene expression likely

depends on the position of the IS relative to the gene's start codon and the nature of its

integration (conservative or deleterious; see **Fig. S4** in Additional File 1). In some cases, the

IS will reduce expression by altering either the promoter or the DNA binding domain of a

putative inducer of the flanking gene, while in other cases it may promote enhanced

transcription by eliminating the binding site of a repressor. These hypotheses are currently

under investigation.

ISs affect *S. aureus* subsp. *anaerobius* gene expression by a novel anti-antisense

RNA interference mechanism.

Intriguingly, the majority of ISs (63 of 87) were located in an antisense orientation to putative

target genes. In such an orientation, the previously described effects on gene expression

would not be feasible, and we reasoned that distinct mechanisms of IS-mediated control of

neighbouring gene expression may exist. To test this hypothesis, we selected 2 examples of

S. aureus subsp. *anaerobius* genes that were located in an anti-sense orientation to the

flanking IS, including *mgr*A, encoding a global regulator of virulence [27-29], and *adh*A

encoding a zinc-dependent alcohol dehydrogenase, induced under low oxygen conditions

[30] (**Figs. S4** and **S5** in Additional File 1). Since the antisense orientation of these genes

relative to IS precludes monitoring of gene expression by transcriptional fusions, they were

cloned to express 3xFLAG tagged versions of the encoded proteins, which facilitated

measurement of expression levels by western immunoblot. To examine the impact of IS on

gene expression, expression plasmid constructs with and without IS in the antisense

orientation were introduced to *S. aureus* subsp. *aureus* strain RN4220 Δ*spa* and expression

levels compared (see scheme in **Fig. S5** in Additional File 1). Surprisingly, no differences in

the expression levels of MgrA or AdhA were observed in any of the constructs suggesting

that the presence of ISs did not actively alter expression levels of the downstream genes by

a classical anti-sense mechanism (**Figs. S5A** and **S5B**, respectively, in Additional File 1).

However, the analysis of the genetic context of *mgr*A revealed another possible mechanism

of IS-mediated control of gene expression. In the ancestral *S. aureus* subsp. *aureus*, the

gene downstream of *mgr*A is in an antisense orientation (**Fig. S4** in Additional File 1).

Furthermore, overlapping antisense transcripts for *mgr*A and its downstream gene in *S.*

aureus have been identified in a previous study [31], suggesting the possibility of antisense

interference of *mgrA* gene expression. Accordingly, we hypothesized that the presence of an

IS between the 2 genes in *S. aureus* subsp. *anaerobius* could disrupt the putative

interference (antisense) mechanism. To examine this possibility, we constructed another set

of reporters (see scheme in **Fig. 4E**) in which we used the inducible expression plasmid

pCN51 to mimic expression from the gene adjacent to *mgr*A. Since the gene located

downstream of *mgr*A belongs to a large operon that was impossible to clone into pCN51, we

used the cadmium-inducible promoter present in plasmid pCN51 as the origin of the anti-

sense mRNA for *mgr*A, mimicking the natural genetic context (**Fig. 4E**). The different

constructs were introduced into the *S. aureus* subsp. *aureus* strain RN4220 Δ*spa* and

expression from the cadmium-inducible pCN51 promoter induced. Absence of the IS

resulted in much reduced MgrA expression levels suggesting that presence of a strong

transcriptional terminator protects *mgr*A mRNA from antisense mRNA interference (**Fig. 4E**).

In support of this idea, insertion of a TT between *mgr*A and the plasmid-encoded cadmium-

inducible promoter produced the same effect as the presence of the complete IS (**Fig. 4E**). A

similar effect was identified on *sle1* encoding an N-acetylmuramyl-L-alanine amidase

involved in staphylococcal cell separation and β-lactam resistance [32] [33], whereby IS

insertion has disrupted the anti-sense regulation of Sle1 expression by a neighbouring gene

in a growth phase-dependent manner (**Fig. 4F**). Thus, IS in *S. aureus* subsp. *anaerobius* can uncouple genes from ancestral antisense regulation by acting as anti-antisense elements, a hitherto unknown regulatory role.

Overall, our data suggest that the non-random insertion of IS into *S. aureus* subsp. *anaerobius* intergenic regions has influenced the expression of neighbouring genes through multiple distinct mechanisms. The putative attenuated expression of up to 87 different chromosomal genes involved in an array of different functions (**Table S4** in Additional File 2) supports the novel concept of the *IS regulon* in which the expression of a subset of genes in a bacterial genome is controlled, not by a transcriptional regulator, but by an IS.

Discussion

While S. *aureus* is an opportunistic pathogen responsible for an array of different pathologies in different anatomical sites in humans and animals, *S. aureus* subsp. *anaerobius* is restricted to a specific infection of superficial lymph nodes in sheep and goats. Here, we provide a remarkable example of the evolutionary transition of a versatile multi-host bacterium to a fastidious highly niche-restricted endemic pathogen of small ruminants. The transition was marked by multiple distinct evolutionary processes mediating drastic changes to the genome that resulted in a complete re-modelling of bacterium-niche interactions. Previous studies have indicated a human ancestral host for *S. aureus* and endemic livestock clones are the result of host jump events that have occurred during the evolutionary history of *S. aureus* [8, 34]. Accordingly, the co-segregation of all isolates of the *S. aureus* subsp. *anaerobius* within a single monophyletic clade in the *S. aureus* phylogeny suggests a likely human to ruminant host-switch event that preceded the evolutionary transition to a highly niche-specific ecology.

Such switches to a host-restricted lifestyle [35], have occurred across the bacterial kingdom in diverse lineages such as *Mycobacterium* [36], *Shigella* [37], *Salmonella* [38], and *Burkholderia* [39], and extreme examples are represented by endosymbionts that have evolved from free-living organisms to become dependent on a single host species [2, 6].

While most known examples of host-restrictive evolution have time-scales of hundreds of

thousands up to many millions of years [6, 35], our analysis indicates that *S. aureus* subsp.

anaerobius evolved about 1000 or more years ago offering a unique insight into the relatively

early stages of evolution towards niche-restriction.

Host shifts are associated with a radical change in habitat, typically with a genetic bottleneck

that diminishes effective population size and a corresponding reduction in purifying selection

activity [2, 35]. We discovered that approximately 10% of the *S. aureus* subsp. *anaerobius*

genome is made up of pseudogenes affecting an array of metabolic and pathogenic

pathways associated with the fastidious nutritional requirements and limited virulence of *S.*

aureus subsp. *anaerobius* (**Fig. 1C**). In particular, the microaerophilic metabolism, is likely

to be due to the loss of function of catalase and other oxidoreductase genes leading to

increased sensitivity to oxidative free radicals. These data suggest adaptation to a nutrient-

rich, oxygen-limited niche, such as that provided by the lymphatic system [11]. The repair of

mildly deleterious mutations such as those resulting in loss of gene function is likely

compromised by lower levels of purifying selection leading to fixation by genetic drift, and

sufficient time has not elapsed to facilitate the deletion of genes no longer required in the

new habitat, possibly also impacted by the observed lower rate of mutation.

We identified a large number of closely-related IS elements distributed around the genome

of *S. aureus* subsp. *anaerobius* (**Fig. 1A**; **Table S2** in Additional File 2), a phenomenon that

has been observed among other bacteria evolving towards host-restriction [2, 4, 6, 39].

Ineffective purifying selection after a population bottleneck also allows insertion sequences

(IS), normally present in bacterial genomes in low numbers (<10), to expand and

disseminate around the genome in the early stages of host-restriction. Over time the IS

elements will be eventually purged, such that recently host-restricted bacteria will generally

contain more IS than ancient symbionts [2, 4]. It is well established that IS elements often

insert into coding regions resulting in gene inactivation and can also mediate chromosomal

rearrangements of genetic segments flanked by ISs via homologous recombination events

[15]. We identified five large rearrangements that occurred in the common ancestor of all

isolates and an additional event exclusive to the clade comprising European isolates.

Although the effects on bacterial phenotype resulting from the identified rearrangements is

unknown, large rearrangements can change the distance of a gene from the origin of

chromosome replication leading to altered gene copy number and expression [40, 41] and

similar rearrangements in *S. aureus* have been reported to mediate transition to reduced

virulence phenotypes such as single colony variants associated with persistent infections

[16, 17].

While IS elements often inactivate genes, it is striking that only one of 87 IS insertions in

S. aureus subsp. *anaerobius* results in gene disruption. It is established that IS elements can

also activate expression of neighbouring genes, either by an extended transcription from an

internal promoter or by the generation of a hybrid promoter [25, 42]. Uniquely, the IS-

mediated gene regulation of *S. aureus* subsp. *anaerobius* involves one established and one

novel mechanism of control. Upstream IS elements oriented in the sense orientation relative

to the flanking gene (~25% of IS) act mainly through modulation of promoter and operon

structure as previously reported [43]. However, in *S. aureus* subsp*. anaerobius*, the majority

of ISs (~75%) are located downstream in the antisense orientation. In the ancestral

S. aureus subsp. *aureus* strain, the expression of some genes is controlled by transcripts of

a downstream gene in the antisense orientation suggesting expression of the corresponding

proteins is mutually exclusive [44]. Here, we have shown that in *S. aureus*

subsp. *anaerobius,* IS inserted in an antisense orientation to flanking genes can effectively

uncouple the targeted genes from their interdependent expression control and act as anti-

antisense regulatory elements. This novel regulatory mechanism might provide a selective

advantage to *S. aureus* subsp. *anaerobius* in facilitating the simultaneous expression of

proteins required concurrently the new niche.

While some bacterial pathogens such as *Bordetella pertussis* utilise IS elements to control

the expression of their flanking genes in a strain-specific manner [43], the genomic

localisation of the majority of IS elements in *S. aureus* subsp. *anaerobius* is conserved

across the phylogeny, indicating that insertion events happened early in the evolution of

S. aureus subsp. *anaerobius*, and suggesting that the complement of genes affected may be

important for the ecology of *S. aureus* subsp. *anaerobius*. Based on this, we propose the

novel concept of the *IS regulon*, representing the set of genes that is transcriptionally

controlled by IS elements.

In conclusion, using a combination of phylodynamic, comparative genomic and molecular

biology approaches, we have dissected the relatively recent evolution of a host-restricted

bacterial pathogen underpinned by drastic changes to the genome via numerous distinct

processes. In particular, IS elements have been domesticated by *S. aureus,* contributing to

the chromosomal architecture via intra-chromosomal rearrangements, and control of gene

expression via multiple mechanisms. Taken together, our findings provide a unique and

remarkable example of the capacity of bacterial pathogens to expand into new host niches.

Methods

Bacterial strains and growth conditions.

The bacterial strains employed are detailed in **Table S5** in Additional File 2. *S. aureus* was

grown in Tryptic soy broth (TSB) or on Tryptic soy agar and *Escherichia coli* was grown in

Luria-Bertani broth (LB) or on LB agar. Antibiotic selection was used where appropriate

(erythromycin 10 µg ml^{-1} for *S. aureus* and ampicillin 100 µg ml^{-1} for *E. coli*.

Whole genome sequencing and genomic analysis.

Forty *S. aureus* subsp. *anaerobius* ovine isolates previously reported [10] were selected, of

which 31 were sampled in Spain during different outbreaks spanning 3 decades (1981-

2012). The rest of the isolates, sampled between 1996 and 2011, were from Sudan (n=3),

Italy (n=3), Poland (n=2) and Denmark (n=1). Isolates were sequenced using a MiSeq

machine (Ilumina, San Diego, CA, USA); the paired-end short reads were trimmed using

Trimmomatic v0.36 [45] and *de novo* assembled into contigs using SPAdes v3.10.0 [46] and

Velvet v1.2.10 [47]. Illumina read data is available in the European Nucleotide Archive under the study accession number PRJEB30965.

One of the Spanish isolates (MVF7, the type strain) was sequenced on the RSII platform (Pacific Biosciences, Menlo Park, CA, USA) using SMRT technology in the Centre for Genomic Research (University of Liverpool, UK). The long reads were de novo assembled into a single contig using the Hierarchical Genome Assembly Process (HGAP) method. Also included in the analysis was the only *S. aureus* subsp. *anaerobius* draft genome publicly available, isolate ST1464 (assembly GCA_000588835.1) [13]. Genes of all 41 genomes and draft assemblies were annotated using Prokka v1.12 [48], and the pan-genome was determined using Roary v3.8.2 [49] applying a 95% identity cut-off. *In silico* multilocus sequence typing (MLST) of the assemblies was performed using the MLST tool v2.8 (https://github.com/tseemann/mlst). Phage sequences and genomic islands were identified using PHASTER [50] and Island Viewer v4 [51], respectively, and manually inspected for reliability. Antimicrobial resistance genes were detected using ResFinder v3.0 [52].

Pseudogene detection.

Here, pseudogenes were defined as protein-coding sequences with homologous genes in an *S. aureus* subsp. *aureus* reference that were split or truncated (i.e. <80% of the reference gene length). For this purpose, the isolate RF122 (assembly GCA_000009005.1) was employed as a reference and a custom python script (https://github.com/GonzaloYebra/anaerobius) was developed to identify pseudogenes from the output of a Roary analysis of *S. aureus* subsp. *anaerobius* isolates compared to the *S. aureus* RF122 genome. The analyses were repeated using multiple *S. aureus* subsp. *aureus* genome sequence references, but this did not alter significantly the pseudogenes detected (data not shown). Ancestral functions of pseudogenes were predicted by assigning them to clusters of orthologous groups (COGs) using eggNOG v4.5.1 [53]. Enrichment analyses of GO (Gene Ontology) terms and KEGG (Kyoto Encyclopedia of Genes and Genomes) pathways assigned by eggNOG were performed using the R

packages topGO v2.34.0 and ClusterProfiler v3.10.0 [54]. Additionally, the Integrated

Microbial Genomes (IGM) annotation tool [55] was used to infer phenotypic metabolic

characteristics from the presence/absence of protein pathways.

Insertion sequence detection.

Transposase coding sequences were identified in the MVF7 complete genome using the

Prokka output and the ISsaga web tool hosted in the ISfinder platform [56]. Determining the

location of IS elements and other repetitive sequences by assembly of short-read sequences

is often not feasible [57]. Therefore, in order to examine the distribution of IS identified in

strain MVF7 among other isolates, Illumina reads were mapped to fragments of the MVF7

genome representing each transposase edge together with their flanking genes using BWA-

MEM [58]. In this manner, identification of reads that spanned the border between the IS and

the flanking gene indicated the presence of that IS at the same genomic location relative to

MVF7.

Phylogenetic analyses.

A core genome alignment was created by aligning the Illumina short reads and the ST1464

assembly   to   the   MVF7   whole   genome   using   Snippy   v3.1

(https://github.com/tseemann/snippy). Sites containing any gap character ('-') or unknown

nucleotide ('N') were discarded. Gubbins v2.2.0 [59] was used to detect recombinant regions

which were then discarded. A maximum likelihood (ML) tree was constructed using IQ-TREE

[60], applying the GTR nucleotide substitution model together with a gamma-distributed rate

heterogeneity across sites and 1,000 ultrafast bootstrap replicates. Temporal signal was

investigated using TempEst v1.5 [61] by means of the correlation between root-to-tip

distances and sampling dates (**Fig. S1** in Additional File 1). Bayesian phylogenetic analysis

was performed with BEAST v1.9.0 [62] using the HKY model for nucleotide substitution.

Different models were tested for the molecular clock (strict and uncorrelated lognormal

relaxed) and demographic (constant, exponential and Bayesian skygrid) models. Each of

these model combinations were run for 100 million generations, with sampling every 10,000

and discarding the initial 10% as burn-in. Runs were compared via a marginal likelihood

estimation (MLE) using path sampling and stepping stone sampling methods implemented in

BEAST. The posterior distribution of trees was summarised into a maximum clade credibility

tree.

In order to examine the relatedness of *S. aureus* subsp. *anaerobius* and *S. aureus*, another

core genome SNP tree was created, also using Snippy v3.1 and IQ-TREE (same settings

than above). It included a sample of 790 *S. aureus* genomes (corresponding to 43 different

host species and 77 clonal complexes (CCs), isolated in 50 different countries) along with 17

isolates of the most closely-related staphylococcal species (*S. schweitzeri* and *S. argenteus*)

[8]. All bioinformatic analyses were carried out using the Cloud Infrastructure for Microbial

Bioinformatics (CLIMB) facility [63].

Gene cloning.

General DNA manipulations were performed using standard procedures. PCR fragments

were amplified from genomic DNA using Kapa Hifi DNA polymerase (Kapa Biosystems).

PCR fragments were either digested using restriction endonucleases and ligated into the

respective plasmid backbone or assembled directly into a linearised plasmid backbone using

Gibson assembly (NEBuilder HiFi DNA Assembly, NEB) according to the manufacturer's

instructions. The plasmids and oligonucleotides used in this study are listed in **Tables S6**

and **S7** in Additional File 2, respectively.

Western blot analysis.

S. aureus cultures harbouring the respective pCN47 or pCN51 derivative plasmids were

diluted 1:50 from overnight cultures and grown in TSB at 37°C and 120 rpm until sample

collection. For pCN51 constructs induced with $CdCl_2$, cultures were split at early exponential

phase (OD_{540}~0.15), 5 µM $CdCl_2$ was added to half the cultures and incubation continued for

another 2 h. *S. aureus* strains carrying pCN47 reporter constructs were grown to early

exponential (OD_{540} ~0.15), mid-exponential (OD_{540} ~0.8) or stationary phase (OD_{540} ~2 after

overnight culture). A sample amount corresponding to and OD_{540} of 0.5 in 1 ml was pelleted

and stored at -20°C. The sample pellets were re-suspended in 100 µl digestion/lysis buffer

(50 mM Tris-HCl, 20 mM MgCl$_2$, 30% (w/v) raffinose) plus 1 µl of lysostaphin (12.5 µg ml^{-1})

and incubated at 37°C for 1 h. Samples for SDS-PAGE were prepared by adding 4X

NuPAGE LDS Sample Buffer (Invitrogen) and proteins denatured at 95°C for 10 min.

Samples were separated by SDS-PAGE (NuPAGE™ 4-12% Bis-Tris Protein Gels,

Invitrogen and then transferred to a PVDF transfer membrane (Thermo Scientific, 0.2 µM)

following standard procedures [64, 65]. FLAG-tagged proteins were detected using mouse

anti-FLAG-HRP antibody (Monoclonal ANTI-FLAG® M2-Peroxidase (HRP), Sigma-Aldrich)

following the manufacturer's protocol.

Enzyme assay for the quantification of β-lactamase activity in transcriptional fusion

plasmids.

Samples for pCN41 based reporters were grown to either early exponential phase (OD-

$_{540}$~0.15) or stationary phase for promoters with low activity (*met*C) (OD$_{540}$ ~2 after overnight

culture) and 1 ml of culture snap frozen. β-Lactamase assays, using nitrocefin (BD

Diagnostic systems) as substrate, were performed as previously described [66] using an

ELx808 microplate reader (BioTek) measuring absorbance at 490 nm. Promoter activity was

calculated using the following equation:

$$Promoter\ activity = \frac{dA_{490}}{dt(h)}\frac{1}{OD_{540} * DF * V}$$

Declarations

Ethics approval and consent to participate.

Not applicable.

Consent for publication.

Not applicable.

Availability of data and materials.

Illumina short-read sequence data was deposited at the European Nucleotide Archive with

study accession number PRJEB30965. The MVF7 full genome assembly was deposited at

NCBI with GenBank accession number CP062279.

Competing interests.

The authors declare that they have no competing interests.

Funding.

The study was supported by a project grant (BB/K00638X/1) and institute strategic grant

funding ISP2: BB/P013740/1 from the Biotechnology and Biological Sciences Research

Council (United Kingdom) to J.R.F.; grant MR/N02995X/1 from the Medical Research

Council (United Kingdom) to J.R.F; Wellcome Trust collaborative award 201531/Z/16/Z to

J.R.F. and J.R.P.

Authors' contributions.

JRF and JRP conceived and led the study. GY, BAW and EJR performed the genomics

analyses. AFH and MMN performed the *in vitro* analyses. SG advised on the transcriptional

analyses. PH, MAT-M and R de la F provided the isolates and their metadata. GY, AFH,

JRP and JRF wrote the first draft of the manuscript. All authors read and approved the final

manuscript.

Acknowledgments.

We wish to thank Dr. Prerna Vohra and Dr. Cosmin Chintoan-Uta (The Roslin Institute,

University of Edinburgh) for advice regarding culture of the *S. aureus* subsp. *anaerobius*

isolates under microaerophilic conditions.

References.

1.   Woolhouse ME, Taylor LH, Haydon DT. Population biology of multihost pathogens.
Science. 2001; 292(5519):1109-1112.

2.   Moran NA, Plague GR. Genomic changes following host restriction in bacteria. Curr
Opin Genet Dev. 2004; 14(6):627-633.

3.   Baumler A, Fang FC. Host specificity of bacterial pathogens. Cold Spring Harb Perspect
Med. 2013; 3(12):a010041.

4.   Bobay LM, Ochman H. The evolution of bacterial genome architecture. Front Genet.
2017; 872.

5.   Sheppard SK, Guttman DS, Fitzgerald JR. Population genomics of bacterial host
adaptation. Nat Rev Genet. 2018; 19(9):549-565.

6.   Weinert LA, Welch JJ. Why might bacterial pathogens have small genomes? Trends
Ecol Evol. 2017; 32(12):936-947.

7.   Fitzgerald JR. Livestock-associated Staphylococcus aureus: origin, evolution and public
health threat. Trends in Microbiology. 2012; 20(4):192-198.

8.   Richardson EJ, Bacigalupe R, Harrison EM, Weinert LA, Lycett S, Vrieling M, et al.
Gene exchange drives the ecological success of a multi-host bacterial pathogen. Nat
Ecol Evol. 2018.

9.   de la Fuente R, Suarez G. Respiratory deficient *Staphylococcus aureus* as the
aetiological agent of "abscess disease". Zentralbl Veterinarmed B. 1985; 32(6):397-406.

10.   de la Fuente R, Ballesteros C, Bautista V, Medina A, Orden JA, Domínguez-Bernal G, et
al. *Staphylococcus aureus* subsp*. anaerobius* isolates from different countries are clonal
in nature. Vet Microbiol. 2011; 150(1-2):198-202.

11.   De la Fuente R, Suarez G, Schleifer KH. Staphylococcus aureus subsp. anaerobius
subsp. nov., the causal agent of abscess disease of sheep. Int J Syst Bacteriol. 1985;
35(1):99-102.

12.   Szaluś-Jordanow O, Krysztopa-Grzybowska K, Czopowicz M, Moroz A, Mickiewicz M,
Lutyńska A, et al. MLST and RAPD molecular analysis of *Staphylococcus aureus* subsp*.
anaerobius* isolated from goats in Poland. Archives of Microbiology. 2018; 200(9):1407-
1410.

13.   Elbir H, Robert C, Nguyen TT, Gimenez G, El Sanousi SM, Flock JI, et al.
Staphylococcus aureus subsp*. anaerobius* strain ST1464 genome sequence. Stand
Genomic Sci. 2013; 9(1):1-13.

14.   Gibson B, Wilson DJ, Feil E, Eyre-Walker A. The distribution of bacterial doubling times
in the wild. Proc Biol Sci. 2018; 285(1880).

15.   Parkhill J, Sebaihia M, Preston A, Murphy LD, Thomson N, Harris DE, et al.
Comparative analysis of the genome sequences of *Bordetella pertussis, Bordetella
parapertussis* and *Bordetella bronchiseptica*. Nat Genet. 2003; 35(1):32-40.

16.   Cui L, Neoh H-m, Iwamoto A, Hiramatsu K. Coordinated phenotype switching with large-
scale chromosome flip-flop inversion observed in bacteria. Proceedings of the National
Academy of Sciences. 2012; 109(25):E1647-E1656.

17. Guérillot R, Kostoulias X, Donovan L, Li L, Carter GP, Hachani A, et al. Unstable
chromosome rearrangements in *Staphylococcus aureus* cause phenotype switching
associated with persistent infections. Proceedings of the National Academy of Sciences.
2019; 116(40):20135-20140.

18. Guinane CM, Ben Zakour NL, Tormo-Mas MA, Weinert LA, Lowder BV, Cartwright RA,
et al. Evolutionary genomics of *Staphylococcus aureus* reveals insights into the origin
and molecular basis of ruminant host adaptation. Genome Biol Evol. 2010; 2454-466.

19. Lerat E, Ochman H. Recognizing the pseudogenes in bacterial genomes. Nucleic Acids
Res. 2005; 33(10):3125-3132.

20. Lowder BV, Guinane CM, Ben Zakour NL, Weinert LA, Conway-Morris A, Cartwright
RA, et al. Recent human-to-poultry host jump, adaptation, and pandemic spread of
Staphylococcus aureus. Proc Natl Acad Sci USA. 2009; 106(46):19545-19550.

21. Viana D, Blanco J, Tormo-Mas MA, Selva L, Guinane CM, Baselga R, et al. Adaptation
of *Staphylococcus aureus* to ruminant and equine hosts involves SaPI-carried variants
of von Willebrand factor-binding protein. Mol Microbiol. 2010; 77(6):1583-1594.

22. Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. Mol
Biol Evol. 2006; 23(2):254-267.

23. Malachowa N, Kobayashi SD, Porter AR, Braughton KR, Scott DP, Gardner DJ, et al.
Contribution of *Staphylococcus aureus* coagulases and clumping factor A to abscess
formation in a rabbit model of skin and soft tissue infection. PLoS One. 2016;
11(6):e0158293.

24. Cheng AG, DeDent AC, Schneewind O, Missiakas D. A play in four acts:
Staphylococcus aureus abscess formation. Trends in Microbiology. 2011; 19(5):225-
232.

25. Siguier P, Gourbeyre E, Chandler M. Bacterial insertion sequences: their genomic
impact and diversity. FEMS Microbiol Rev. 2014; 38(5):865-891.

26. Charpentier E, Anton AI, Barry P, Alfonso B, Fang Y, Novick RP. Novel cassette-based
shuttle vector system for gram-positive bacteria. Appl Environ Microbiol. 2004;
70(10):6076-6085.

27. Luong TT, Lee CY. The arl locus positively regulates *Staphylococcus aureus* type 5
capsule via an mgrA-dependent pathway. Microbiology (Reading). 2006; 152(Pt
10):3123-3131.

28. Crosby HA, Schlievert PM, Merriman JA, King JM, Salgado-Pabon W, Horswill AR. The
Staphylococcus aureus global regulator MgrA modulates clumping and virulence by
controlling surface protein expression. PLoS Pathog. 2016; 12(5):e1005604.

29. Crosby HA, Tiwari N, Kwiecinski JM, Xu Z, Dykstra A, Jenul C, et al. The
Staphylococcus aureus ArlRS two-component system regulates virulence factor
expression through MgrA. Mol Microbiol. 2020; 113(1):103-122.

30. Pagels M, Fuchs S, Pane-Farre J, Kohler C, Menschner L, Hecker M, et al. Redox
sensing by a Rex-family repressor is involved in the regulation of anaerobic gene
expression in *Staphylococcus aureus*. Mol Microbiol. 2010; 76(5):1142-1161.

31. Lasa I, Toledo-Arana A, Dobin A, Villanueva M, de los Mozos IR, Vergara-Irigaray M, et
al. Genome-wide antisense transcription drives mRNA processing in bacteria. Proc Natl
Acad Sci U S A. 2011; 108(50):20172-20177.

32. Kajimura J, Fujiwara T, Yamada S, Suzawa Y, Nishida T, Oyamada Y, et al.
Identification and molecular characterization of an N-acetylmuramyl-L-alanine amidase
Sle1 involved in cell separation of Staphylococcus aureus. Mol Microbiol. 2005;
58(4):1087-1101.

33. Thalsø-Madsen I, Torrubia FR, Xu L, Petersen A, Jensen C, Frees D. The Sle1 cell wall
amidase is essential for β-lactam resistance in community-acquired methicillin-resistant
Staphylococcus aureus USA300. Antimicrob Agents Chemother. 2019; 64(1).

34. Weinert LA, Welch JJ, Suchard MA, Lemey P, Rambaut A, Fitzgerald JR. Molecular
dating of human-to-bovid host jumps by Staphylococcus aureus reveals an association
with the spread of domestication. Biol Lett. 2012; 8(5):829-832.

35. Toft C, Andersson SG. Evolutionary microbial genomics: insights into bacterial host
adaptation. Nat Rev Genet. 2010; 11(7):465-475.

36. Cole ST, Eiglmeier K, Parkhill J, James KD, Thomson NR, Wheeler PR, et al. Massive
gene decay in the leprosy bacillus. Nature. 2001; 409(6823):1007-1011.

37. Feng Y, Chen Z, Liu SL. Gene decay in *Shigella* as an incipient stage of host-
adaptation. PLoS One. 2011; 6(11):e27754.

38. Langridge GC, Fookes M, Connor TR, Feltwell T, Feasey N, Parsons BN, et al. Patterns
of genome evolution that have accompanied host adaptation in *Salmonella*. Proc Natl
Acad Sci USA. 2015; 112(3):863-868.

39. Song H, Hwang J, Yi H, Ulrich RL, Yu Y, Nierman WC, et al. The early stage of bacterial
genome-reductive evolution in the host. PLoS Pathog. 2010; 6(5):e1000922.

40. Block DH, Hussein R, Liang LW, Lim HN. Regulatory consequences of gene
translocation in bacteria. Nucleic Acids Res. 2012; 40(18):8979-8992.

41. Periwal V, Scaria V. Insights into structural variations and genome rearrangements in
prokaryotic genomes. Bioinformatics. 2015; 31(1):1-9.

42. Prentki P, Teter B, Chandler M, Galas DJ. Functional promoters created by the insertion
of transposable element IS1. J Mol Biol. 1986; 191(3):383-393.

43. Amman F, D'Halluin A, Antoine R, Huot L, Bibova I, Keidel K, et al. Primary
transcriptome analysis reveals importance of IS elements for the shaping of the
transcriptional landscape of Bordetella pertussis. RNA Biol. 2018; 15(7):967-975.

44. Toledo-Arana A, Lasa I. Advances in bacterial transcriptome understanding: From
overlapping transcription to the excludon concept. Mol Microbiol. 2020; 113(3):593-602.

45. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. 2014; 30(15):2114-2120.

46. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a
new genome assembly algorithm and its applications to single-cell sequencing. J
Comput Biol. 2012; 19(5):455-477.

47. Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de
Bruijn graphs. Genome Res. 2008; 18(5):821-829.

48. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;
30(14):2068-2069.

49. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, et al. Roary: rapid
large-scale prokaryote pan genome analysis. Bioinformatics. 2015; 31(22):3691-3693.

50. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster
version of the PHAST phage search tool. Nucleic Acids Res. 2016; 44(W1):W16-21.

51. Bertelli C, Laird MR, Williams KP, Simon Fraser University Research Computing G, Lau
BY, Hoad G, et al. IslandViewer 4: expanded prediction of genomic islands for larger-
scale datasets. Nucleic Acids Res. 2017.

52. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al.
Identification of acquired antimicrobial resistance genes. J Antimicrob Chemother. 2012;
67(11):2640-2644.

53. Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, et al. eggNOG
4.5: a hierarchical orthology framework with improved functional annotations for
eukaryotic, prokaryotic and viral sequences. Nucleic Acids Res. 2016; 44(D1):D286-
293.

54. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological
themes among gene clusters. OMICS. 2012; 16(5):284-287.

55. Markowitz VM, Chen IM, Palaniappan K, Chu K, Szeto E, Grechkin Y, et al. IMG: the
Integrated Microbial Genomes database and comparative analysis system. Nucleic
Acids Res. 2012; 40(Database issue):D115-122.

56. Varani AM, Siguier P, Gourbeyre E, Charneau V, Chandler M. ISsaga is an ensemble of
web-based methods for high throughput identification and semi-automatic annotation of
insertion sequences in prokaryotic genomes. Genome Biol. 2011; 12(3):R30.

57. Treangen TJ, Salzberg SL. Repetitive DNA and next-generation sequencing:
computational challenges and solutions. Nat Rev Genet. 2011; 13(1):36-46.

58. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 2009; 25(14):1754-1760.

59. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, et al. Rapid
phylogenetic analysis of large samples of recombinant bacterial whole genome
sequences using Gubbins. Nucleic Acids Res. 2015; 43(3):e15.

60. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol.
2015; 32(1):268-274.

61. Rambaut A, Lam TT, Max Carvalho L, Pybus OG. Exploring the temporal structure of
heterochronous sequences using TempEst (formerly Path-O-Gen). Virus Evol. 2016;
2(1):vew007.

62. Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti
and the BEAST 1.7. Mol Biol Evol. 2012; 29(8):1969-1973.

63. Connor TR, Loman NJ, Thompson S, Smith A, Southgate J, Poplawski R, et al. CLIMB
(the Cloud Infrastructure for Microbial Bioinformatics): an online resource for the medical
microbiology community. Microb Genom. 2016; 2(9):e000086.

64. Ausubel F, Brent R, Kingston R, Moore D, Seidman J, Smith J, et al: Current Protocols
in Molecular Biology. New York, NY: John Wiley & Sons; 1990.

65. Sambrook J, Fritsch EF, Maniatis T: Molecular cloning: a laboratory manual. Cold
Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 1989.

66. Ji G, Beavis R, Novick RP. Bacterial interference caused by autoinducing peptide
variants. Science. 1997; 276(5321):2027-2030.

Supplementary Information

Additional file 1 (PDF format): Additional Information; Supplementary Figures S1-S6.

Additional file 2 (XLSX format): Supplementary Tables S1-S7.

Figure Legends

Figure 1: Pseudogenes, insertion sequences and mobile genetic elements in the

***Staphylococcus aureus* subsp. *anaerobius* isolate MVF7.** (A) Circular map of the

chromosome. Rings show, from outside to inside: annotated genes in the positive strand,

those in the negative strand (with the 205 pseudogenes and 87 insertion sequences shown

in red and blue, respectively), and the mobile genetic elements found across all isolates

(gold: prophage ΦSaa1; cyan: pathogenicity island SaaPIMVF7). (B) Gene map of

SaaPIMVF7. Genes in grey are pseudogenes and genes in orange are intact by comparison

other SaPI relatives. *int:* integrase; *pri:* primase; *scn*: Staphylococcal complement inhibitor

(SCIN); *vwb*: von Willebrand factor-binding protein (vWBP). The box shows the expression

of the SaaPIMVF7-encoded *vwb*. The SaaPIMVF7 *vwb* gene was cloned into the expression

vector pCN51 under the control of a cadmium-inducible promoter, transformed into a

coagulase and vWbp-deficient derivative of strain RN4220 (RN4220 *coa*::*tet*M Δ*vwb*) and

the ability of SaaPIMVF7 vWbp to coagulate ruminant plasma was assessed. (C) Graphical

summary of the main biological functions potentially disrupted by the presence of

pseudogenes.

Figure 2: *Staphylococcus aureus* subsp. *anaerobius* represents a single clade within

the *S. aureus* phylogenetic tree. Maximum Likelihood tree constructed from a SNP

alignment of the studied *S. aureus* subsp. *anaerobius* sequences (clades in blue) and 787

sequences of different *S. aureus* subsp. *aureus* (in black). (A) Unrooted tree showing the

divergence of 17 sequences of other *Staphylococcus* (*S. schweitzeri* and *S. argenteus*)

whereas *S. aureus* subsp. *aureus* is embedded in the *S. aureus* diversity. (B) Subtree

showing the position of the *S. aureus* subsp. *anaerobius* clade (collapsed in blue) with

respect to the other *S. aureus* subsp. *aureus* clonal complexes (CC).

Figure 3: *S. aureus* subsp. *anaerobius* evolved over the last 1000 years with a stable

population size. Bayesian maximum clade credibility tree of the *Staphylococcus aureus*

subsp. *anaerobius* isolates. The x-axis is expressed in calendar years, branch colours

denote sampling country. Purple bars at each node show its most recent common ancestor

(MRCA) confidence interval. Numbers in nodes indicate the posterior probability. The

estimated dates of the MRCAs of the main lineages are indicated. The isolate that was

whole genome sequenced using PacBio technology (MVF7) is highlighted in orange. The

bottom plot represents the changes in the effective population size over time, with the

shadowed area representing the 95% credible interval, following the same timescale as the

tree.

Figure 4: *S. aureus* subsp. *anaerobius* has undergone 6 large intra-chromosomal

rearrangements. Map of the genomic translocations across the *Staphylococcus aureus*

subsp. *anaerobius* isolate MVF7 chromosome. The green blocks represent the 6 large

translocations detected in MVF7 when compared to the *Staphylococcus aureus* subsp.

aureus RF122 strain (see also **Fig. S2** in the Additional File 1). The shaded areas indicate

the original location of the translocated portions in the putative ancestral genome. The inner

ring shows the GC skew (green for positive skew, purple for negative), which is unaffected

by the translocations.

Figure 5: Characterisation of the transcriptional terminators (TTs) present in the ISs.

(A) Schematic representation of IS-associated TTs. (B) Schematic representation of and

functional assessment of the bi-directional TT1 localised in the 5' region of the IS. The bi-

directional TT1 is composed of two sub-TTs that share the same sequence but are inverted

repeats, thus generating a master bi-directional TT1. The expression of the *bla*Z reporter in

plasmid pCN42 is controlled by a cadmium-inducible promoter. The identified transcriptional

terminators were cloned between the promoter and the *bla*Z reporter gene. β-lactamase

expression was monitored either in uninduced (black bars) or Cd-induced (open bars)

cultures 180 min after induction. Data are show the means of three independent biological

replicates and error bars show the standard deviation from the mean. Statistical analysis

was performed using two-way ANOVA followed by Dunnett's post-test. Multiplicity adjusted

p-values are shown for uninduced to induced comparison and all terminator containing

reporters showed significant differences when compared to their respective empty plasmid

control. **** $p < 0.0001$, ns not significant.

Figure 6: Presence of IS influences the expression of downstream genes and genes

product through multiple mechanisms. (A-D) Reporter constructs for presence of the IS

upstream and in sense orientation of the target gene were designed in plasmid pCN41

containing a β-lactamase reporter gene. The green triangle indicates the IS position relative

to the target gene's start. All reporters were constructed either by removing the IS gene from

S. aureus subsp. *anaerobius* strain MVF84 or by introducing the IS into *S. aureus* subsp.

aureus strain RF122 to confirm the IS role on gene expression variation. In (A) the IS

removal results in a 207bp deletion to the promoter region not present in RF122. Reporter

plasmids were introduced into RN4220 as described in Methods. Data show the mean of

three biological replicates, error bars represent the mean's standard deviation. One-Way

ANOVA was performed followed by Tukey's multiple comparisons test. *$p < 0.05$, **$p < 0.01$,

***p<0.001, ****p<0.0001, *ns* not significant. (E&F) Western blot analysis of (E) *mgr*A and (F)

sle1 expression constructs to address whether the IS impacts on their expression levels by

interrupting expression of an antisense transcript. (E) *mgr*A encoding a 3xFLAG tag was

cloned in antisense to a cadmium-inducible promoter into plasmid pCN51 to mimic the

expression of an antisense transcript to *mgr*A. Constructs presented different combinations

of presence/absence of IS and/or transcriptional terminator the IS transcript or expression

from the plasmid-encoded cadmium-inducible promoter from the *mgr*A transcript. Expression

of the cadmium-inducible promoter was induced with 5 μM CdCl$_2$. (F) To assess the IS

impact in the context of a natural antisense transcript, 3xFLAG-encoding expression

constructs of *sle1* including the gene downstream of *sle1* in the ancestral genome were

cloned into the plasmid pCN47. Samples were taken during different growth phases.

Plasmids were introduced into RN4220 Δ*spa* as described in Methods. Western blots shown

are representative of at least two independent biological replicates. TT, transcriptional

terminator.

A MVF84_01231 *rps*P

B MVF84_01317c *dna*D

C MVF84_00302 *met*C

D MVF84_00796 *met*N