Massive genome decay and expansion of insertion sequences drive the evolution of a novel host-restricted bacterial pathogen

Gonzalo Yebra[1,a], Andreas F Haag[2,a], Maan M Neamah[2,3,4], Bryan A Wee[1], Emily J Richardson[1], Pilar Horcajo[5], Sander Granneman[6], María Ángeles Tormo-Más[7,8], Ricardo de la Fuente[5], J Ross Fitzgerald[1,*], José R Penadés[2,7,9,*]

[1]The Roslin Institute, University of Edinburgh, Edinburgh, United Kingdom; [2]Institute of Infection, Immunity & Inflammation, University of Glasgow, Glasgow, United Kingdom; [3]Faculty of Veterinary Medicine, University of Kufa, Kufa, Iraq; [4]Middle Euphrates Centre for Cancer and Genetic Research, University of Kufa, Kufa, Iraq; [5]Facultad de Veterinaria, Universidad Complutense de Madrid, Madrid, Spain; [6]Centre for Synthetic and Systems Biology, University of Edinburgh, Edinburgh, United Kingdom; [7]Departamento de Ciencias Biomédicas, Facultad de Ciencias de la Salud, Universidad CEU Cardenal Herrera, Moncada, Valencia, Spain; [8]Severe Infection Group, Health Research Institute Hospital La Fe, Valencia, Spain; [9]MRC Centre for Molecular Bacteriology and Infection, Imperial College London, London, United Kingdom.

[a] These authors contributed equally.

* Corresponding authors:
    JRF (ross.fitzgerald@roslin.ed.ac.uk)
    JRP (j.penades@imperial.ac.uk)

Keywords:

Bacterial evolution; host adaptation; host restriction; genome reduction; gene decay; insertion elements; pseudogenes; structural variation; gene expression regulation; phylodynamics.

Abstract

Background. The emergence of new bacterial pathogens represents a major threat to public and veterinary health. *Staphylococcus aureus* is a multi-host bacterial species comprising pathogenic clones with distinct tropisms for human and livestock species. A *S. aureus* microaerophilic subspecies, *Staphylococcus aureus* subsp*. anaerobius*, is responsible for outbreaks of a specific lymphadenitis pathology (Morel's disease) exclusively found in small ruminants. However, the evolutionary history of *S. aureus* subsp*. anaerobius* and its genetic relatedness to *S. aureus* are unknown.

Results. Evolutionary genomic analyses of clinical *S. aureus* subsp*. anaerobius* isolates sampled across 3 decades revealed this clone emerged from a *S. aureus* progenitor about 1000 years ago (95%CI: 716-1184), before differentiating into two distinct lineages representing African (emerged in 1930 [1907-1951)) and European (1777 [1716-1832]) isolates. *S. aureus* subsp*. anaerobius* has undergone limited clonal expansion, with a restricted population size, and an evolutionary rate 10-fold slower than *S. aureus*. The transition to a highly niche-specific pathogen of small ruminant lymph nodes involved acquisition of a pathogenicity island encoding an effector with ruminant host-specificity, large genomic rearrangements, and the accumulation of at least 205 pseudogenes resulting in a highly fastidious metabolism underpinning its restricted ecological niche. Importantly, acquisition and expansion of ~87 insertion sequences located in conserved intergenic regions provided distinct mechanisms for the control of expression of flanking genes, representing a novel concept of transcriptional regulon.

Conclusions. Our findings provide a remarkable example of the evolutionary trajectory of a host-restricted bacterial pathogen that resulted from extensive remodelling of the *S. aureus* genome.

Background

Bacteria are particularly able to adapt to new environmental niches, a characteristic that may promote their capacity to become highly successful pathogens. Many pathogens can infect multiple hosts [1], but occasionally strains of free-living bacteria become restricted to one particular host. The evolutionary process of host restriction, observed across distantly-related bacterial groups, typically involves one or more processes such as genome reduction, acquisition of mobile genetic elements and insertion sequences, and chromosomal rearrangements [2-4]. The magnitude of these changes may be most manifest after long-term associations between bacterium and the host, with obligate intracellular pathogens the most extreme of these cases exhibiting the smallest and most compact genomes [5].

Staphylococcus aureus is an important and versatile bacterial pathogen, capable of infecting a broad array of animals [6], and thus representing a threat for public and livestock health. Understanding the evolutionary events underpinning successful host switches may assist the design of strategies to contain the emergence of new human and animal pathogenic clones. In some cases, the genetic determinants involved in host adaptation have been identified. While some virulence factors encoded in certain mobile genetic elements play an important role in the adaptation of *S. aureus* to ruminants and poultry [7, 8], a single mutation is sufficient to allow the *S. aureus* human-to-rabbit host jump [9, 10]. These diverse strategies highlight the versatility that *S. aureus* has in adapting and infecting new hosts.

A close relative of *S. aureus*, *S. aureus* subsp. *anaerobius*, responsible for Morel's disease [11], is highly restricted to small ruminants and its clinical manifestation is confined to abscesses located near major superficial lymph nodes. Moreover, and contrary to *S. aureus* subsp. *aureus*, *S. aureus* subsp. *anaerobius* is microaerophilic and catalase-negative [12]. Importantly, outbreaks of this disease have been detected in Europe, Africa and the Middle East, causing serious economic losses [13].

Here, we investigate the evolutionary history of *S. aureus* subsp. *anaerobius*, revealing that it descended from an ancestor resembling contemporary *S. aureus* and adapted to its current

ecological niche by extensive genome diversification through a number of processes including massive genome decay, gene acquisition, and genome rearrangements. Of note, the emergence of an IS regulon, involving regulatory control of almost 90 chromosomal genes by an ISSau8-like insertion sequence, represents a novel mode of transcriptional remodelling that accompanied the transition to a new niche. Overall, our results reveal a new strategy used by *S. aureus* to adapt to a new host, highlighting the extraordinary versatility of this important pathogen.

Results

***S. aureus* subsp. *anaerobius* exhibits a highly conserved genome with limited strain-dependent variation**

The genomic architecture of *S. aureus* subsp. *anaerobius* remains largely unknown since only one fragmented draft genome was available until now, corresponding to the strain ST1464 isolated in Sudan [14]. In order to provide a fully assembled genome sequence, we applied PacBio sequencing to the type strain MVF7, isolated in Central Spain in 1981-1982 [12]. We obtained a closed genome which comprised a circular chromosome of 2,755,024bp with 32.74% GC content (**Fig. 1A**) and no plasmids. This genome contained 2,888 coding DNA sequences (CDS), 56 tRNAs and 5 complete rRNA copies.

In order to explore the population structure and the variability in gene content of *S. aureus* subsp. *anaerobius*, we performed whole genome sequencing using Illumina on a collection of 39 additional strains from different countries [13]. Of them, 30 were sampled in Spain during different outbreaks spanning 3 decades (1981-2012). The rest of the isolates, sampled between 1996 and 2011, were from Sudan (n=3), Italy (n=3), Poland (n=2) and Denmark (n=1). These newly generated assemblies were analysed together with that of the ST1464 isolate. The 39 draft assemblies presented an average length of 2,683,726bp (range: 2,612,583-2,749,895bp). The pan-genome analysis showed that the core genome (i.e. genes shared by all isolates) included 2,479 (92.7%) of the total 2,675 genes found. Of the remaining,

strain-specific genes, 66 (2.5%) were "soft core" genome (present in 95-99% of sequences), and the rest (26, 4.8%) were mostly truncated genes in assemblies with low quality. Interestingly, 15 genes were present in European isolates but absent in Sudanese ones (including 2 clusters of 4 genes each) and 5 genes vice versa.

Phylogenetic analyses reveal two distinct sub-lineages of *S. aureus* subsp. *anaerobius*

To explore the population structure of *S. aureus* subsp. *anaerobius*, we generated a core genome alignment including all our samples and the only publicly available draft assembly to make a total of 41 genomes. In this alignment we detected 5 putative recombinant regions (size range: 54-2,950bp) that were excluded from the core genome alignment. The definitive alignment was 2,324,010bp long and included 3,443 variable sites, and it was used for further phylogenetic analysis.

The phylogenetic analyses of the core genome alignment revealed two distinct lineages within *S. aureus* subsp. *anaerobius* – one including the isolates sampled in Sudan and the other one including the rest of isolates, all sampled in Europe (**Fig. 2**). The core alignment showed that the average number of SNPs between pairs of *S. aureus* subsp. *anaerobius* genomes was 494 SNPs (range: 0-1,746). This value was 231 (0-450) within the European clade and 97 (2-142) within the Sudanese clade. The average distance between these two clades was 1,697 SNPs (1,663-1,746). For reference, the average number of SNPs between the *S. aureus* subsp. *anaerobius* isolates and the *S. aureus* RF122 isolate was 27,611 (27,582-27,650).

In contrast to previous results that ascribed all *S. aureus* subsp. *anaerobius* isolates to a single, homogenous sequence type (ST1464) [13, 15], all Sudanese sequences belonged to the same ST but we found 3 other STs within European sequences (see **Additional Information** in Additional File 1 for more detail).

Linear regression analysis provided evidence of a molecular clock-like evolution in the ML phylogeny. The subsequent time-scaled Bayesian phylogenetic analysis showed that the model combination that best fitted the data was a strict molecular clock paired with a constant

population size. This analysis dated the most recent common ancestor (MRCA) of *S. aureus* subsp. *anaerobius* in the year 1069 AD (95% Bayesian confidence interval (BCI): 828-1296) (**Fig. 2**). The European and Sudanese lineages diversified in the years 1777 (95% BCI: 1716-1832) and 1930 (95% BCI: 1907-1951), respectively. The estimated evolutionary rate was 4.4 (95% BCI: 3.3-5.5) × 10^{-7} substitutions/site/year (s/s/y), which approximates to 1.2 SNPs per year. This is roughly one order of magnitude slower than *S. aureus* subsp. *aureus*. The rest of models tested in BEAST provided very similar results, with mean MRCA estimates between 1111 and 1333 AD and mean evolutionary rates between 4.2 and 5.4 × 10^{-7} s/s/y. This agrees with the slow *in vitro* growth observed for *S. aureus* subsp. *anaerobius* [11], as evolutionary rates are proportional to doubling times [16].

To ascertain the phylogenetic relationship between *S. aureus* subsp. *anaerobius* and *S. aureus* subsp. *aureus,* we constructed a core genome SNP tree with 807 whole-genome sequences representative of the *Staphylococcus* diversity [17]. In this tree (**Fig. 3**), *S. aureus* subsp. *anaerobius* formed a well-defined cluster within *S. aureus* subsp. *aureus* but without being genetically close to any clonal complex. This phylogenetic evidence reinforces the idea that this subspecies evolved and diversified from an unknown *S. aureus* lineage after a host adaptation and restriction process that started at least a millennium ago. Although there has not been any study specifically designed to estimate the *S. aureus* subsp. *aureus* emergence time, a recent article [17] which analysed a globally diverse dataset found a root date of 7,000 BC – approximately 10 times older.

Massive gene decay characterises the *S. aureus* subsp. *anaerobius* genome

The complete genome of *S. aureus* subsp. *anaerobius* strain MVF7 contained 205 pseudogenes, with an average of 204.9 pseudogenes per genome across all *S. aureus* subsp. *anaerobius* strains (range: 201-210). On average, 66.6% (range: 64.7-68.8%) of the pseudogenes were split CDSs and 33.4% (range: 31.2-35.3%) were truncated CDSs. These pseudogenes were evenly distributed across the genome (**Fig. 1A**), and originated through point mutations causing frameshifts, premature stop codons or alternative downstream start

codons. Pseudogenes represented an average of 9.5% (range: 9.2-10.1%) of the genome length across all *S. aureus* subsp. *anaerobius* genomes.

A total of 164 pseudogenes (approximately 80% of those found in any given isolate) were shared among all *S. aureus* subsp. *anaerobius* genomes. Of those pseudogenes, 92.1% were caused by the same mutation in all isolates suggesting they were an early event in its evolutionary history. The proportion of shared pseudogenes was higher within each of the two main phylogenetic clades described above: Sudanese isolates shared 183 pseudogenes (89%) and European isolates shared 179 (87%).

The functions of the 205 pseudogenes present in MVF7 were classified into clusters of orthologous groups (COGs) of proteins (for a more detailed description, see **Additional Information** in Additional File 1 and **Table S1** in Additional File 2). Whereas 43 (26.2%) had unknown or unavailable functions, half of these proteins (n=82, 50%) corresponded to metabolic functions, 29 (17.7%) to cellular processes and signalling, and 10 (6.1%) to information storage and processing. Within the pseudogenes affecting metabolic functions, most were associated with amino acids (n=29, 32.6%), carbohydrates (n=14, 15.7%), inorganic ions (n=9, 10.1%) and functions affecting energy production (n=8, 9.0%). The enrichment analyses of both KEGG pathways and GO terms (see **Table S2** in Additional File 2 for a full list) revealed a statistically significant pseudogenisation of genes involved in biosynthesis of amino acids and metabolic pathways.

Although most pseudogenes were involved in numerous metabolic pathways, others were defence mechanisms (such as catalase and other oxidoreductases, transmembrane pumps involved in antibiotic resistance and fluoride toxicity, and type I restriction enzymes), and virulence factors (such as proteins involved in capsule biosynthesis, adherence factors and exoproteins). In short, genes affected by pseudogenisation could explain the limited metabolism and pathogenicity of this bacterium.

S. aureus subsp. *anaerobius* contains novel mobile genetic elements

All *S. aureus subsp. anaerobius* sequences carried a 43.2 kb-long, novel prophage (ΦSaa1) belonging to the *Siphoviridae* family (**Fig. 1B**), which also contained many pseudogenes when compared to the closest available prophage (Φ2958PVL, accession number NC_011344). The prophage presented a novel integrase type (see **Fig. S1** in Additional File 1) and a pseudogenised variant of the virulence-associated protein E, *virE.*

We also identified one novel 13 kb Staphylococcal Pathogenicity Island (SaPI) present in all isolates, which encoded additional copies of the Staphylococcal complement inhibitor (SCIN) and the von Willebrand factor-binding protein (vWbp) (**Fig. 1B**). vWbp has previously been demonstrated to show host-specific activity, being able to coagulate ruminant plasma [7, 8]. A phylogenetic network using SplitsTree v4.14.6 and the NeighborNet algorithm [18] of SaaPIMVF7 and previously described SaPIs (**Fig. S2** in Additional File 1) showed that SaaPIMVF7 is closest to SaPIov2 (93.1% nucleotide identity), firstly described in the *S. aureus* subsp. *aureus* clonal complex CC133 (responsible for infections of small ruminants including sheep and goats) which also carries vWbp and SCIN (**Fig. 1B**). SaaPIMVF7 contained pseudogenes including the integrase and primase genes, suggesting that it is no longer mobile but stable in the chromosome. Of note, *S. aureus* subsp. *anaerobius* also has chromosomal copies of *vwb* and *scn* of which the former is pseudogenised. SaaPIMVF7-encoded *vwb* and *scn* were transcribed (**Fig. 1C**) and apparently functional in SaaPIMVF7 suggesting a key role in the pathogenicity of *S. aureus* subsp. *anaerobius.* The SaaPIMVF7-encoded *vwb* was cloned into the expression vector pCN51, and the ability of the expressed protein to coagulate ruminant plasma confirmed (**Fig. S3** in Additional File 1), verifying its role in virulence.

Insertion sequences drive *S. aureus* subsp. *anaerobius* evolution

Yet another unique feature of the *S. aureus* subsp. *anaerobius* genomes was the presence of 87 insertion sequences in MVF7, all identified as ISSau8 (IS family ISL3 described in the strain RF122) by ISsaga [19]. The average nucleotide similarity between all was 99.3%, and that to ISSau8 was 97.0%. Most of them had internal stop codons that split the transposase-coding

region in two or more fragments whereas only 10 ISs contained full-length transposase genes. ISs were distributed evenly throughout the genome (**Fig. 1A**).

Since *S. aureus* subsp. *anaerobius* has undergone massive gene decay, we initially hypothesised that ISs had caused pseudogenes, inactivating genes not required in the new niche. However, all the ISs were located in intergenic regions—except only one inserted within a gene (*msb*A), resulting into a split CDS (i.e. pseudogene). Since only ~16% of the *S. aureus* subsp. *anaerobius* genome corresponds to non-coding intergenic regions, the non-random distribution of the ISs through the genome suggested an additional role for these elements in the evolutionary adaptation of this species. We hypothesised that *S. aureus* subsp. *anaerobius* uses ISs to control the expression of the genes located downstream of the IS elements. These would represent a novel concept of regulon, in which a subset of genes would be controlled not by a transcriptional regulator but by an IS.

To test this idea, we studied the conservation of IS locations across all the *S. aureus* subsp. *anaerobius* strains. Because their high number and genetic similarity (>99% nucleotide identity) caused frequent contig breaks in the draft genomes assembled from short-reads, ISs were only reliably reconstructed in the MVF7 close genome sequence. We tested the presence of each MVF7 IS in the rest of isolates by mapping their respective Illumina reads against the border between each IS and their flanking gene, and identifying reads overlapping this boundary. This analysis revealed that 68 out of the 87 MVF7 ISs (78.2%) were inserted in the same locations (i.e. we detected reads mapping against both IS flanks) in all *S. aureus* subsp. *anaerobius* samples. We observed differences between the European and Sudanese clades: 75 (86.2%) ISs shared genomic location among all the European isolates, whereas 7 were missing in all Sudanese isolates. These data suggest that IS have been important not only for the initial host jump, but also for the origin and divergence observed between clades (**Fig. 2**). Interestingly, one of the ISs detected in all European isolates but absent in all Sudanese ones was flanking one of the translocations (number V) described below, which agrees with the fact that such translocation was not detected in the Sudanese strains.

Our prediction of promoter and transcriptional terminator (TT) sequences showed that all IS had a functional promoter but not a TT between the *tnp* gene and the downstream gene, supporting the idea that the *tnp* transcript could influence expression of the gene located 3' of the IS gene. This also reported the existence of two additional putative TTs in the IS genes: one bidirectional TT (TT1) located in the 5' region of the IS gene, and a unidirectional one (TT2, antisense orientation) located at the end (3' region) of the IS (see **Fig. 4**). The bidirectional TT1 would putatively prevent the potential interference between transcripts initiated either from the IS or from the gene located upstream to the IS. By contrast, TT2 would prevent the impact that an antisense transcript originated from a gene located downstream of the ISs could have in the expression of the IS transcript (**Fig. 4**). This organisation suggests that the ISs have established a strategy that allows their transcripts to interfere with the genes located in position 3' to the IS, while blocking any external interference that the flanking genes could have in the expression of the regulatory IS transcript.

To validate the functionality and directionality of the TTs, we used an expression construct based on the plasmid pCN42 [20], in which the *bla*Z reporter is under the control of the P$_{cad}$ promoter. The conserved 5' and 3' region of the ISs containing the bi- and unidirectional TTs, respectively, were cloned, in both orientations, between the P$_{cad}$ promoter and the *bla*Z reporter (see **Fig. 4**), and the expression of the reporter gene tested with and without induction of the P$_{cad}$ promotor. The results confirm the existence, functionality and directionality of both TTs, suggesting that the role of the ISs is to control the expression of the chromosomal genes located 3' to the ISs.

IS affect *S. aureus* subsp. *anaerobius* gene regulation by interference with promoter region and operon structure

ISs are inserted into the *S. aureus* subsp. *anaerobius* either upstream or downstream of targeted genes. To evaluate their impact on the expression of neighbouring genes, we initially analysed whether the presence of an IS controls the expression of the downstream gene located in the same orientation as the IS. Since *S. aureus* subsp. *anaerobius* is not genetically

tractable, we designed reporter constructs mimicking the genetic organisation found in *S. aureus* subsp. *anaerobius* (**Fig. S4** in Additional File 1) and tested expression levels of the selected genes or gene products in *S. aureus* subsp. *aureus*. Specifically, we selected four different genes (*rps*P, *dna*D, *met*C and *met*N) that contained an upstream IS at different distance to each gene's start codon and constructed *bla*Z transcriptional reporter fusion using pCN41 [20] (**Fig. 5A-D**). Then we tested the expression of these genes in presence or absence of the ISs. Importantly, and in support of the hypothesis, deletion of the ISs affected expression of the neighbouring genes, confirming their regulatory role, although this effect was different depending on the genes. Thus, while removal of the ISs increased transcription of *dna*D, *met*C or *met*N, it decreased *rps*P expression (**Fig. 5A-D**). Based on these results, it is tempting to speculate that the impact of the IS on downstream gene expression depends on the position of the IS relative to the gene's start codon and the nature of its integration (conservative or deleterious, see **Fig. S4** in Additional File 1). In some cases, the presence of the IS will reduce expression by altering either the promoter region or the DNA binding region of a putative inducer of the expression of the targeted gene, while in other cases it may increase transcription by eliminating the binding site of putative repressors. These ideas are now under study.

ISs affect *S. aureus* subsp. *anaerobius* gene expression through a novel mechanism of anti-antisense RNA interference

Most ISs were located downstream and in antisense orientation to putative target genes. Since the previously identified mechanism cannot be involved in the IS-mediated control of the expression of the flanking genes, we reasoned that additional strategies of gene control involving ISs should occur. To test this, we selected two genes located in an antisense orientation to the ISs and tested whether expression from the IS-contained promoter would actively influence their expression (**Figs. S4** and **S5** in Additional File 1). We selected the global regulator-encoding gene *mgr*A, which is a major effector of *S. aureus* gene regulation [21-23], and *adh*A encoding a zinc-dependent alcohol dehydrogenase, which is induced under

low oxygen environmental conditions in a RexA-dependent manner [24]. Since the expression of these flanking genes cannot be monitored using transcriptional fusions, they were cloned so they expressed 3xFLAG versions of the encoded proteins, which were used to monitor the expression of these proteins by western blot. To test the impact of the ISs, some of the plasmids contained the flanking IS (wild-type structure), while we generated as controls plasmids without the ISs (see scheme in **Fig. S5** in Additional File 1). Surprisingly, no differences in the expression levels of MgrA or AdhA were observed in any of the constructs (**Figs. S5A** and **S5B**, respectively, in Additional File 1) suggesting that the presence of ISs did not actively alter expression levels of the downstream genes by a classical anti-sense mechanism.

However, the analysis of the genetic context of *mgr*A revealed another interesting possibility involving ISs in controlling the expression of this gene. In the ancestor *S. aureus* subsp. *aureus*, the gene downstream of *mgr*A has an antisense orientation (**Fig. S4** in Additional File 1) and the transcripts from this gene could interfere with the expression of *mgr*A. This is an interesting hypothesis since overlapping antisense transcripts for *mgr*A and its downstream gene have been identified in several *S. aureus* subsp. *aureus* strains in a previous study [25]. Since as previously demonstrated the ISs have different functional transcriptional terminators (**Fig. 4**), we hypothesized that the presence of the ISs between the two contiguous genes could block this interference (antisense) mechanism. To assess whether this was the case, we constructed another set of reporters (see scheme in **Fig. 5E**) in which we used the inducible expression plasmid pCN51 to mimic expression from the gene adjacent to *mgr*A. Since the gene located downstream of *mgr*A belongs to a large operon which was unfeasible to clone completely in a plasmid, we used the cadmium-inducible promoter present in plasmid pCN51 as the origin of the anti-sense mRNA for *mgr*A, mimicking the natural genetic context. The different constructs were introduced into the *S. aureus* subsp. *aureus* strain RN4220 Δ*spa* and expression from the cadmium-inducible pCN51 promoter induced. Absence of the ISs significantly reduced MgrA expression levels (**Fig. 5E**), compared to the reporter construct

where these were present. This suggests that the IS prevents transcripts from the upstream gene through the presence of a strong transcriptional terminator and thus protects *mgr*A mRNA from antisense mRNA interference. In support of this idea, insertion of a TT between *mgr*A and the plasmid-encoded cadmium-inducible promoter produced the same effect as the presence of the complete IS (**Fig. 5E**).

To test whether this organisation affected the regulation of an IS targeted gene in its native context containing a downstream gene and its promoter in antisense orientation with known overlapping transcription [25], we selected *sle1* (aka *aaa*) encoding an N-acetylmuramyl-L-alanine amidase involved in staphylococcal cell separation [26] and essential for β-lactam resistance of *S. aureus* [27]. The gene downstream of *sle1* is expressed as a single transcript from its own promoter allowing us to construct a set of reporters similar to the constructs prepared for *mgr*A in the plasmid pCN47 [20]. Note that in this scenario all genes are expressed using they native promoter regions. Since the potential expression patterns of the gene downstream of *sle1* were unknown, we assessed the impact of the IS on the expression levels of Sle1 at different stages during bacterial growth (**Fig. 5F**). Similar to the results observed for MgrA, absence of the IS significantly reduced Sle1 expression levels albeit in a growth phase-dependent manner. Thus, certain ISs in *S. aureus* subsp. *anaerobius* uncouple genes from ancestral antisense regulation by acting as anti-antisense elements, a hitherto unprecedented regulatory role.

S. aureus subsp. anaerobius has undergone large genomic rearrangements

The widespread presence of ISs in *S. aureus* subsp. *anaerobius* caused promote genomic rearrangements by homologous recombination, a phenomenon observed in different bacterial species [28]. When compared to *S. aureus* subsp. *aureus*, the genome of *S. aureus* subsp. *anaerobius* contained at least six large genomic translocations, flanked by ISs, whose size ranged between 70 and 346 kbp (**Fig. 6** and **Fig. S6** in Additional File 1). Two pairs of translocations (I and VI, and II and V) switched their original genomic locations whereas the rest (III and IV) were produced by excision and insertion. These translocations interchanged

regions between the leading and lagging strands, but because all also involved inversion, genes in the translocated areas conserved their original orientation and the GC skew remained unaffected. Translocations did not produce any noticeable loss of genomic regions.

We designed a series of primer pairs to qualitatively test whether the same breakpoints between translocations were present in all the *S. aureus* subsp. *anaerobius* strains. The PCR results, summarised in **Table S3** in Additional File 2, showed that the genomic rearrangements observed in the MVF7 full genome were also present in the rest of European isolates, whereas the isolates from Sudan lacked the translocations numbers II and V. This agrees with the fact that one of the ISs delimiting the translocation number II was one of those present in all European isolates but absent in all Sudanese ones. It is therefore likely that this particular translocation happened after these two phylogenetic lineages diverged.

Although there were some cases of PCR pairs failing in certain European isolates, which might be due to the DNA integrity, in general this shows how some of the genomic rearrangements might have occurred after the European and African lineages separated (**Table S3** in Additional File 2).

In summary, our results demonstrate an extraordinary role of the ISs in *S. aureus* subsp*. aureus,* not just controlling the expression of many chromosomal genes, but also shaping the structure of the *S. aureus* subsp*. aureus* chromosome.

Discussion

Here, we demonstrate that *Staphylococcus aureus* subsp. *anaerobius* represents a remarkable example of the drastic genomic changes affecting a bacterial pathogen during the evolutionary process of host restriction. This process involved an outstanding variety of simultaneous evolutionary strategies such as mobile genetic element acquisition, pseudogene accumulation, large genomic rearrangements and, importantly, a novel transcriptional regulatory system orchestrated by widespread insertion elements.

This bacterium was described as a respiratory deficient variant of *S. aureus* due to its lack of aerobic growth and catalase activity [12]. While both are very closely related, their pathogenic abilities are radically different: *S. aureus* is a major pathogen responsible for acute and chronic infections both in humans and animals, while *S. aureus* subsp. *anaerobius* has only been occasionally identified when causing a specific lymphadenitis of sheep and goats. In order to reconstruct the evolutionary trajectory that transformed *S. aureus* subsp. *anaerobius* into a host- and niche-specific pathogen, we performed the first population genomics study by analysing 40 isolates sampled in different countries over the last 30 years. Our results suggest that *S. aureus* subsp. *anaerobius* evolved from a ruminant-adapted *S. aureus* strain (containing specialised mobile genetic elements) through a process of host restriction that modified drastically its genome. Such modifications accumulated gradually during restriction to a novel host of a once free-living bacterium [29], are recurring across the bacterial kingdom. They have been described in diverse lineages such as *Mycobacterium* [30], *Shigella* [31], *Salmonella* [32] and *Burkholderia* [33] (see also review in [2]).

Host shifts represent drastic, sudden changes in the environmental conditions for a bacterial population, associated with genetic bottlenecks which cause a reduction in population size and a weakened purifying (negative) selection [2, 29]. This scenario would explain the fact that one tenth of the *S. aureus* subsp. *anaerobius* genome is made up of pseudogenes, due to an incapacity to clear deleterious mutations that cause gene truncation and frameshifts, which are fixed then by genetic drift. The conservation of pseudogenes could be explained by a weakened purifying selection (which prevents purging these truncated genes no longer needed) as a consequence of a reduced population size and/or by the fact that the extreme adaptation of *S. aureus* subsp. *anaerobius* is a recent event in the evolutionary scale and that selection is limited by a reduced mutation rate. Both explanations agree with our phylodynamic analyses, which shows a relatively recent date of emergence (a thousand years ago), a restricted population size growth, and a reduced evolutionary rate.

The pseudogenisation of genes involved in many metabolic pathways, defence mechanisms, and virulence factors explains the atypical phenotype of *S. aureus* subsp. *anaerobius* when compared to *S. aureus*, including its limited pathogenicity, reduced metabolism and fastidious growth *in vitro*. In addition, its microaerophilic metabolism, caused by a lack of defence against radicals (by truncation of the catalase and others) rather than disruption of oxidative respiration, suggests that this bacterium has become confined to a nutrient-rich, oxygen-limited niche. This explanation agrees with the fact that it can only be cultured from the benign lymph node abscesses it causes [12].

An ineffective purifying selection also allows insertion sequences (IS), normally present in bacterial genomes but in low numbers (<10), to proliferate and spread across the genome in the early stages of host-restriction only to be purged later on –therefore recently adapted bacteria will harbour more ISs than more ancient symbionts [2, 4]. ISs will normally cause gene inactivation (through insertion within coding regions) and chromosomal rearrangements of genetic segments flanked by ISs.

The latter is true for *S. aureus* subsp. *anaerobius*, with five large rearrangements originated in the common ancestor of all isolates and an additional one exclusive to the European clade. Effects of genomic rearrangements are difficult to establish although some of the ones described include reduction in growth rate (again, a characteristic phenotype in *S. aureus* subsp. *anaerobius)* and colony variation [34]. In addition, large rearrangements can change the distance of a gene from the origin of chromosome replication leading to altered gene copy number and thus affecting its expression [34, 35]. During the initial analysis of the *S. aureus* subsp. *anaerobius*, and based in the high number of pseudogenes and ISs present in this species, we hypothesised that most of the pseudogenes would be consequence of the presence of an IS. A more detailed analysis revealed that this idea was wrong, and only one However, only one of the ISs in *S. aureus* subsp. *anaerobius* caused inactivation of one gene, while the rest (n=86) were inserted in intergenic regions, a non-random distribution that we hypothesised was a key feature in the evolutionary origin of *S. aureus* subsp. *anaerobius*. It

has been extensively reported that individual ISs can activate expression of the neighbouring genes, either by an extended transcription from an internal promoter or by the generation of a hybrid promoter [28, 36]. Consequently, domestication of ISs by the bacterium to control specific functions is an evolutionary process that has been unexpectedly common. An example of this is *Bordetella pertussis*, with more than 230 copies of the IS481 in each genome [37], most of which also influence the expression of their flanking genes [38]. In this case, however, the transcripts produced by the ISs are strain-specific due to the dynamic variation in IS copy number and genomic localisation [38]. By contrast, the localisation of most of the ISs present in *S. aureus* subsp. *anaerobius* is conserved in all the strains analysed, indicating they control a set of genes important in the adaptation to the new host. Thus, the insertion of most of these ISs occurred early in the *S. aureus* subsp. *anaerobius* evolution, being subsequently inherited and inactivated (note that most of the *tnp* genes were pseudogenes). This represents a new concept of IS regulon, were a set of genes is transcriptionally controlled by the presence of an ISs. Interestingly, the IS-mediated regulation involves at least two different mechanisms of control, depending on whether the genes flanking the IS are in sense or antisense orientation in relation to the IS. While upstream IS oriented in sense to the flanking gene (~25% of ISs) act mainly through modulation of promoter and operon structure, the majority of ISs (~75%) is located downstream and in antisense orientation to their flanking genes. In the ancestral *S. aureus* subsp. *aureus* strain, the expression of many of these genes is controlled through antisense mechanisms caused by transcripts of a downstream located gene in antisense orientation. This ensures that expression levels of both genes can be modulated under different environmental conditions, allowing *S. aureus* subsp. *aureus* to adapt to changing environments. By contrast, in *S. aureus* subsp. *anaerobius*, the integration of an IS in antisense orientation to flanking genes can effectively act as anti-antisense regulatory elements and uncouple the targeted genes from their native interdependent expression control. This new regulatory mechanism might provide a selective advantage to *S. aureus* subsp. *anaerobius* in its more restricted and consequently more stable host niche

enabling the simultaneous expression of both genes. This strongly suggests that expression of both proteins is required at the same time in this specialised pathogen.

Additionally, the presence of some ISs whose location was strain-dependent suggests that some ISs remain functional in order to generate more variation and/or regulation. Indeed, during its differentiation into two lineages in Africa and Europe a small proportion of ISs integrated in different genomic locations in these lineages. The absence of a few ISs in certain non-phylogenetically related isolates could be due to the limitations of the detection approach, or alternatively imply that ISs could be occasionally lost in some isolates.

Finally, our comparative genomics analysis based on newly generated whole genome sequences unveiled the previously unknown population structure of *S. aureus* subsp. *anaerobius* and revealed the existence of two distinct phylogenetic clades containing the African and European lineages, respectively, which diverged a few centuries ago. However, most of the evolutionary events that shaped the genome of this bacterium (including MGE acquisition, IS insertion, widespread pseudogenisation, and genomic rearrangements) took place before this diversification and were very likely key processes in the initial adaptation of *S. aureus* subsp. *anaerobius* to its current niche in the sheep host.

In conclusion, *S. aureus* subsp. *anaerobius* represents a remarkable example of drastic genomic changes during a relatively recent host and niche adaptation from a versatile and generalist ancestor –an evolutionary journey notably marked by accumulation of pseudogenes and the development of a new concept of regulon orchestrated by insertion elements found in conserved genome localisations.

Methods

Bacterial strains and growth conditions

The bacterial strains used in this study are detailed in **Table S4** in Additional File 2. *S. aureus* strains were grown in Tryptic soy broth (TSB) or on Tryptic soy agar plates and *Escherichia coli* strains were grown in Luria-Bertani broth (LB) or on LB agar plates. Antibiotic selection

473 was used where appropriate (erythromycin 10 µg ml^{-1} for *S. aureus* strains and ampicillin

474 100 µg ml^{-1} for *E. coli* strains).

Sample sequencing and gene content analysis

Forty *S. aureus* subsp. *anaerobius* ovine isolates previously reported [13] were analysed, of

which 31 were sampled in Spain during different outbreaks spanning 3 decades (1981-2012).

The rest of the isolates, sampled between 1996 and 2011, were from Sudan (n=3), Italy (n=3),

Poland (n=2) and Denmark (n=1).

They were sequenced using a MiSeq platform ((Ilumina, San Diego, CA, USA); the paired-end

short reads were trimmed using Trimmomatic v0.36 [39] and *de novo* assembled into contigs

using SPAdes v3.10.0 [40] and Velvet v1.2.10 [41]. Illumina read data is available in the

European Nucleotide Archive under the study accession number PRJEB30965.

One of the Spanish isolates (MVF7, the type strain) was sequenced on the RSII platform

(Pacific Biosciences, Menlo Park, CA, USA) using SMRT technology in the Centre for

Genomic Research (University of Liverpool, UK). The long reads were de novo assembled

into one, full-length contig using the Hierarchical Genome Assembly Process (HGAP) method.

The only *S. aureus* subsp. *anaerobius* draft genome publicly available, isolate ST1464

(assembly GCA_000588835.1) [14] was included in the analyses.

Genes of all 41 genomes and draft assemblies were annotated using Prokka v1.12 [42]. The

pan-genome was calculated using Roary v3.8.2 [43] applying a 95% identity cut-off. *In silico*

multilocus sequence typing (MLST) of the assemblies was performed using the MLST tool

v2.8 (https://github.com/tseemann/mlst). Phage sequences and genomic islands were

identified using PHASTER [44] and Island Viewer v4 [45], respectively, and manually

inspected for reliability. Antimicrobial resistance genes were detected using ResFinder v3.0

[46].

Pseudogene detection

Pseudogenes are gene copies that have lost their protein-coding ability. Here, pseudogenes were defined as protein-coding sequences that had homologous genes in an *S. aureus* subsp*. aureus* reference and that were split or truncated (i.e. <80% of the reference length) with respect to that reference. For this purpose, the isolate RF122 (assembly GCA_000009005.1) was used as reference and a custom python script was developed to identify pseudogenes from the output of a Roary analysis that compared the genetic content of each *S. aureus* subsp. *anaerobius* to the RF122 genome. The analyses were repeated using different *S. aureus* subsp*. aureus* genome references, but this did not alter significantly the pseudogenes detected (data not shown).

Functions of pseudogenes were investigated by assigning them to clusters of orthologous groups (COGs) using eggNOG v4.5.1 [47]. Enrichment analyses of GO (Gene Ontology) terms and KEGG (Kyoto Encyclopedia of Genes and Genomes) pathways assigned to the pseudogenes by eggNOG were performed using the R packages topGO v2.34.0 and ClusterProfiler v3.10.0 [48]. Additionally, the Integrated Microbial Genomes (IGM) annotation tool [49] was used to infer phenotypic metabolic characteristics from the presence/absence of protein pathways.

Insertion sequence detection and transcription analyses

Transposase coding sequences were identified in the MVF7 full genome using the Prokka output and the ISsaga web tool hosted in the ISfinder platform [19].

For the rest of the isolates, *de novo* assembly from Illumina short reads failed to reconstruct the highly conserved and repeated transposase genes with enough specificity, causing contig breaks in those locations. Consequently, few or no transposases could be inferred from their draft assemblies. This is a common issue when trying to resolve repetitive regions using short read data, which causes ambiguity in selecting the best location in the genome to which align such reads [50].

Therefore, the presence of ISs was tested by mapping the Illumina reads against them using BWA-MEM [51] to fragments of the MVF7 genome containing each transposase edge together with part of their flanking genes. This was done to detect Illumina reads that overlapped the border between the IS and the flanking gene as indicator of the presence of that IS in that particular genomic location.

Phylogenetic analyses

A core genome alignment was created by aligning the Illumina short reads and the ST1464 assembly to the MVF7 whole genome using Snippy v3.1 (https://github.com/tseemann/snippy). Sites containing any gap character ('-') or unknown nucleotide ('N') were discarded. Gubbins v2.2.0 [52] was used detect recombinant regions in this alignment, which were discarded.

A maximum likelihood (ML) tree was constructed using IQ-TREE [53], applying the GTR nucleotide substitution model together with a gamma-distributed rate heterogeneity across sites and 1,000 ultrafast bootstrap replicates. Temporal signal was investigated using TempEst v1.5 [54] by means of the correlation between root-to-tip distances and sampling dates.

Bayesian phylogenetic analysis was performed using BEAST v1.9.0 [55]. The HKY model for nucleotide substitution was used, whereas different models were tested for the molecular clock (strict and uncorrelated lognormal relaxed) and demographic (constant, exponential and Bayesian skygrid) models. Each of these model combinations were run for 100 million generations, with sampling every 10,000 and discarding the initial 10% as burn-in. Runs were compared via a marginal likelihood estimation (MLE) using path sampling and stepping stone sampling methods implemented in BEAST. The posterior distribution of trees was summarised into a maximum clade credibility tree.

In order to explore the relationship between *S. aureus* subsp. *anaerobius* and *S. aureus*, another core genome SNP tree was created, also using Snippy v3.1 and IQ-TREE (same

settings than above). It included a sample of 787 *S. aureus* genomes (corresponding to 43 different animal host species and 77 clonal complexes (CCs), isolated in 50 different countries) along with 17 isolates of the most closely-related staphylococcal species (*S. schweitzeri* and *S. argenteus*) [17].

Bioinformatics analyses were carried out using the Cloud Infrastructure for Microbial Bioinformatics (CLIMB) facility [56].

DNA methods

General DNA manipulations were performed using standard procedures. Cloning and assembly of reporter constructs was performed using restriction digest and ligation or Gibson assembly (NEBuilder HiFi DNA Assembly, NEB) according to the manufacturer's instructions. The plasmids and oligonucleotides used in this study are listed in **Tables S5** and **S6** in Additional File 2, respectively.

Western blot

S. aureus cultures harbouring the respective pCN47 or pCN51 derivative plasmids were diluted 1:50 from overnight cultures and grown in TSB at 37°C and 120 rpm until sample collection. For pCN51 constructs induced with $CdCl_2$, cultures were split at early exponential phase (OD_{540}~0.15), 5 µM $CdCl_2$ was added to half cultures and incubation continued for another 2 h. *S. aureus* strains carrying pCN47 reporter constructs were grown to early exponential (OD_{540} ~0.15), mid-exponential (OD_{540} ~0.8) or stationary phase (OD_{540} ~2 after overnight culture). A sample amount corresponding to and OD_{540} of 0.5 in 1 ml was pelleted and stored at -20°C. The sample pellets were re-suspended in 100 µl digestion/lysis buffer (50 mM Tris-HCl, 20 mM $MgCl_2$, 30% (w/v) raffinose) plus 1 µl of lysostaphin (12.5 µg ml^{-1}) and incubated at 37°C for 1 h. Samples for SDS-PAGE were prepared by adding 4X NuPAGE LDS Sample Buffer (Invitrogen) and proteins denatured at 95°C for 10 min. Samples were separated by SDS-PAGE (NuPAGE™ 4-12% Bis-Tris Protein Gels, Invitrogen and then transferred to a PVDF transfer membrane (Thermo Scientific, 0.2 µM) following standard procedures [57, 58]. FLAG-tagged proteins were detected using mouse anti-FLAG-HRP

antibody (Monoclonal ANTI-FLAG® M2-Peroxidase (HRP), Sigma-Aldrich) following the manufacturer's protocol.

Enzyme assay for the quantification of β-lactamase activity in transcriptional fusion plasmids

Samples for pCN41 based reporters were grown to either early exponential phase (OD$_{540}$~0.15) or stationary phase for promoters with low activity (*met*C) (OD$_{540}$ ~2 after overnight culture) and 1 ml of culture snap frozen. β-Lactamase assays, using nitrocefin (BD Diagnostic systems) as substrate, were performed as previously described [59] using an ELx808 microplate reader (BioTek) measuring absorbance at 490 nm. Promoter activity was calculated using the following equation:

$$Promoter\ activity = \frac{dA_{490}}{dt(h)}\frac{1}{OD_{540} * DF * V}$$

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Illumina short-read sequence data was deposited at the European Nucleotide Archive with study accession number PRJEB30965. The MVF7 full genome assembly was deposited at NCBI with GenBank accession number CP062279.

Competing interests

The authors declare that they have no competing interests.

Funding

The study was supported by a project grant (BB/K00638X/1) and institute strategic grant funding ISP2: BB/P013740/1 from the Biotechnology and Biological Sciences Research Council (United Kingdom) to J.R.F.; grant MR/N02995X/1 from the Medical Research Council (United Kingdom) to J.R.F; Wellcome Trust collaborative award 201531/Z/16/Z to J.R.F. and J.R.P.

Authors' contributions

JRF and JRP lead the project. GY, BAW and EJR performed the genomics analyses. AFH and MMN performed the *in vitro* analyses. SG advised on the transcriptional analyses. PH, MAT-M and R de la F provided the isolates and their metadata. GY, AFH, JRP and JRF wrote the first draft of the manuscript. All authors read and approved the final manuscript.

Acknowledgments

We wish to thank Dr. Prerna Vohra and Dr. Cosmin Chintoan-Uta (The Roslin Institute, University of Edinburgh) for their help culturing the *S. aureus* subsp. *anaerobius* isolates under microaerophilic conditions.

References

1. Woolhouse ME, Taylor LH, Haydon DT. Population biology of multihost pathogens. Science. 2001; 292(5519):1109-1112.

2. Moran NA, Plague GR. Genomic changes following host restriction in bacteria. Curr Opin Genet Dev. 2004; 14(6):627-633.

3. Baumler A, Fang FC. Host specificity of bacterial pathogens. Cold Spring Harb Perspect Med. 2013; 3(12):a010041.

4. Bobay LM, Ochman H. The evolution of bacterial genome architecture. Front Genet. 2017; 872.

5. Weinert LA, Welch JJ. Why might bacterial pathogens have small genomes? Trends Ecol Evol. 2017; 32(12):936-947.

6. Fitzgerald JR, Holden MT. Genomics of natural populations of *Staphylococcus aureus*. Annu Rev Microbiol. 2016; 70459-478.

7. Guinane CM, Ben Zakour NL, Tormo-Mas MA, Weinert LA, Lowder BV, Cartwright RA, et al. Evolutionary genomics of *Staphylococcus aureus* reveals insights into the origin and molecular basis of ruminant host adaptation. Genome Biol Evol. 2010; 2454-466.

8. Viana D, Blanco J, Tormo-Mas MA, Selva L, Guinane CM, Baselga R, et al. Adaptation of
Staphylococcus aureus to ruminant and equine hosts involves SaPI-carried variants of von
Willebrand factor-binding protein. Mol Microbiol. 2010; 77(6):1583-1594.

9. Viana D, Comos M, McAdam PR, Ward MJ, Selva L, Guinane CM, et al. A single natural nucleotide
mutation alters bacterial pathogen host tropism. Nat Genet. 2015; 47(4):361-366.

10. Wilson GJ, Seo KS, Cartwright RA, Connelley T, Chuang-Smith ON, Merriman JA, et al. A novel
core genome-encoded superantigen contributes to lethality of community-associated MRSA
necrotizing pneumonia. PLoS Pathog. 2011; 7(10):e1002271.

11. de la Fuente R, Suarez G. Respiratory deficient *Staphylococcus aureus* as the aetiological agent of
"abscess disease". Zentralbl Veterinarmed B. 1985; 32(6):397-406.

12. De la Fuente R, Suarez G, Schleifer KH. Staphylococcus aureus subsp. anaerobius subsp. nov.,
the causal agent of abscess disease of sheep. Int J Syst Bacteriol. 1985; 35(1):99-102.

13. de la Fuente R, Ballesteros C, Bautista V, Medina A, Orden JA, Domínguez-Bernal G, et al.
Staphylococcus aureus subsp*. anaerobius* isolates from different countries are clonal in nature. Vet
Microbiol. 2011; 150(1-2):198-202.

14. Elbir H, Robert C, Nguyen TT, Gimenez G, El Sanousi SM, Flock JI, et al. *Staphylococcus aureus*
subsp*. anaerobius* strain ST1464 genome sequence. Stand Genomic Sci. 2013; 9(1):1-13.

15. Elbir H, Feil EJ, Drancourt M, Roux V, El Sanousi SM, Eshag M, et al. Ovine clone ST1464: a
predominant genotype of *Staphylococcus aureus* subsp*. anaerobius* isolated from sheep in Sudan.
J Infect Dev Ctries. 2010; 4(4):235-238.

16. Gibson B, Wilson DJ, Feil E, Eyre-Walker A. The distribution of bacterial doubling times in the wild.
Proc Biol Sci. 2018; 285(1880).

17. Richardson EJ, Bacigalupe R, Harrison EM, Weinert LA, Lycett S, Vrieling M, et al. Gene exchange
drives the ecological success of a multi-host bacterial pathogen. Nat Ecol Evol. 2018.

18. Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. Mol Biol Evol.
2006; 23(2):254-267.

19. Varani AM, Siguier P, Gourbeyre E, Charneau V, Chandler M. ISsaga is an ensemble of web-based
methods for high throughput identification and semi-automatic annotation of insertion sequences in
prokaryotic genomes. Genome Biol. 2011; 12(3):R30.

20. Charpentier E, Anton AI, Barry P, Alfonso B, Fang Y, Novick RP. Novel cassette-based shuttle vector
system for gram-positive bacteria. Appl Environ Microbiol. 2004; 70(10):6076-6085.

21. Luong TT, Lee CY. The arl locus positively regulates *Staphylococcus aureus* type 5 capsule via an
mgrA-dependent pathway. Microbiology (Reading). 2006; 152(Pt 10):3123-3131.

22. Crosby HA, Schlievert PM, Merriman JA, King JM, Salgado-Pabon W, Horswill AR. The
*Staphylococcus aureu*s Global Regulator MgrA Modulates Clumping and Virulence by Controlling
Surface Protein Expression. PLoS Pathog. 2016; 12(5):e1005604.

23. Crosby HA, Tiwari N, Kwiecinski JM, Xu Z, Dykstra A, Jenul C, et al. The *Staphylococcus aureus*
ArlRS two-component system regulates virulence factor expression through MgrA. Mol Microbiol.
2020; 113(1):103-122.

24. Pagels M, Fuchs S, Pane-Farre J, Kohler C, Menschner L, Hecker M, et al. Redox sensing by a Rex-family repressor is involved in the regulation of anaerobic gene expression in *Staphylococcus aureus*. Mol Microbiol. 2010; 76(5):1142-1161.

25. Lasa I, Toledo-Arana A, Dobin A, Villanueva M, de los Mozos IR, Vergara-Irigaray M, et al. Genome-wide antisense transcription drives mRNA processing in bacteria. Proc Natl Acad Sci U S A. 2011; 108(50):20172-20177.

26. Kajimura J, Fujiwara T, Yamada S, Suzawa Y, Nishida T, Oyamada Y, et al. Identification and molecular characterization of an N-acetylmuramyl-L-alanine amidase Sle1 involved in cell separation of *Staphylococcus aureus*. Mol Microbiol. 2005; 58(4):1087-1101.

27. Thalsø-Madsen I, Torrubia FR, Xu L, Petersen A, Jensen C, Frees D. The Sle1 cell wall amidase is essential for β-lactam resistance in community-acquired methicillin-resistant *Staphylococcus aureus* USA300. Antimicrob Agents Chemother. 2019; 64(1).

28. Siguier P, Gourbeyre E, Chandler M. Bacterial insertion sequences: their genomic impact and diversity. FEMS Microbiol Rev. 2014; 38(5):865-891.

29. Toft C, Andersson SG. Evolutionary microbial genomics: insights into bacterial host adaptation. Nat Rev Genet. 2010; 11(7):465-475.

30. Cole ST, Eiglmeier K, Parkhill J, James KD, Thomson NR, Wheeler PR, et al. Massive gene decay in the leprosy bacillus. Nature. 2001; 409(6823):1007-1011.

31. Feng Y, Chen Z, Liu SL. Gene decay in *Shigella* as an incipient stage of host-adaptation. PLoS One. 2011; 6(11):e27754.

32. Langridge GC, Fookes M, Connor TR, Feltwell T, Feasey N, Parsons BN, et al. Patterns of genome evolution that have accompanied host adaptation in *Salmonella*. Proc Natl Acad Sci USA. 2015; 112(3):863-868.

33. Song H, Hwang J, Yi H, Ulrich RL, Yu Y, Nierman WC, et al. The early stage of bacterial genome-reductive evolution in the host. PLoS Pathog. 2010; 6(5):e1000922.

34. Periwal V, Scaria V. Insights into structural variations and genome rearrangements in prokaryotic genomes. Bioinformatics. 2015; 31(1):1-9.

35. Block DH, Hussein R, Liang LW, Lim HN. Regulatory consequences of gene translocation in bacteria. Nucleic Acids Res. 2012; 40(18):8979-8992.

36. Prentki P, Teter B, Chandler M, Galas DJ. Functional promoters created by the insertion of transposable element IS1. J Mol Biol. 1986; 191(3):383-393.

37. Parkhill J, Sebaihia M, Preston A, Murphy LD, Thomson N, Harris DE, et al. Comparative analysis of the genome sequences of *Bordetella pertussis, Bordetella parapertussis* and *Bordetella bronchiseptica*. Nat Genet. 2003; 35(1):32-40.

38. Amman F, D'Halluin A, Antoine R, Huot L, Bibova I, Keidel K, et al. Primary transcriptome analysis reveals importance of IS elements for the shaping of the transcriptional landscape of Bordetella pertussis. RNA Biol. 2018; 15(7):967-975.

39. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 2014; 30(15):2114-2120.

40.Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome
assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012; 19(5):455-
477.

41.Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs.
Genome Res. 2008; 18(5):821-829.

42.Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014; 30(14):2068-2069.

43.Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, et al. Roary: rapid large-scale
prokaryote pan genome analysis. Bioinformatics. 2015; 31(22):3691-3693.

44.Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster version of
the PHAST phage search tool. Nucleic Acids Res. 2016; 44(W1):W16-21.

45.Bertelli C, Laird MR, Williams KP, Simon Fraser University Research Computing G, Lau BY, Hoad
G, et al. IslandViewer 4: expanded prediction of genomic islands for larger-scale datasets. Nucleic
Acids Res. 2017.

46.Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of
acquired antimicrobial resistance genes. J Antimicrob Chemother. 2012; 67(11):2640-2644.

47.Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, et al. eggNOG 4.5: a
hierarchical orthology framework with improved functional annotations for eukaryotic, prokaryotic
and viral sequences. Nucleic Acids Res. 2016; 44(D1):D286-293.

48.Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological themes
among gene clusters. OMICS. 2012; 16(5):284-287.

49.Markowitz VM, Chen IM, Palaniappan K, Chu K, Szeto E, Grechkin Y, et al. IMG: the Integrated
Microbial Genomes database and comparative analysis system. Nucleic Acids Res. 2012;
40(Database issue):D115-122.

50.Treangen TJ, Salzberg SL. Repetitive DNA and next-generation sequencing: computational
challenges and solutions. Nat Rev Genet. 2011; 13(1):36-46.

51.Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 2009; 25(14):1754-1760.

52.Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, et al. Rapid phylogenetic
analysis of large samples of recombinant bacterial whole genome sequences using Gubbins.
Nucleic Acids Res. 2015; 43(3):e15.

53.Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic
algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015; 32(1):268-274.

54.Rambaut A, Lam TT, Max Carvalho L, Pybus OG. Exploring the temporal structure of
heterochronous sequences using TempEst (formerly Path-O-Gen). Virus Evol. 2016; 2(1):vew007.

55.Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti and the
BEAST 1.7. Mol Biol Evol. 2012; 29(8):1969-1973.

56.Connor TR, Loman NJ, Thompson S, Smith A, Southgate J, Poplawski R, et al. CLIMB (the Cloud
Infrastructure for Microbial Bioinformatics): an online resource for the medical microbiology
community. Microb Genom. 2016; 2(9):e000086.

57. Ausubel F, Brent R, Kingston R, Moore D, Seidman J, Smith J, et al: Current Protocols in Molecular Biology. New York, NY: John Wiley & Sons; 1990.

58. Sambrook J, Fritsch EF, Maniatis T: Molecular cloning: a laboratory manual. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 1989.

59. Ji G, Beavis R, Novick RP. Bacterial interference caused by autoinducing peptide variants. Science. 1997; 276(5321):2027-2030.

List of Figures and Tables

Main Figures:

Supplementary Figures (in Additional File 1)**:**

Supplementary Tables (in Additional File 2):

Figure 1. Distribution of pseudogenes, insertion sequences and mobile genetic elements in the *Staphylococcus aureus* subsp. *anaerobius* isolate MVF7. (A): Circular map of the chromosome. Rings show, from outside to inside: annotated genes in the positive strand, those in the negative strand (with the 205 pseudogenes and 87 insertion sequences shown in red and blue, respectively), and the mobile genetic elements found across all isolates (gold: prophage ΦSaa1; cyan: pathogenicity island SaaPIMVF7). (B) and (C): genes found in ΦSaa1 and SaaPIMVF7, respectively. Genes in grey are pseudogenes and genes in orange are intact (according to homology against their most similar phage (Φ2958PVL) and SaPI (SaPIov2) relatives. The histograms in blue represent the genes' expression levels (inferred from RNA-seq read coverage).

Figure 2. Bayesian maximum clade credibility tree of the *Staphylococcus aureus* subsp. *anaerobius* isolates. The x-axis is expressed in calendar years. Branch colours denote sampling country (see key). Purple bars at each node show its most recent common ancestor (MRCA) confidence interval. Numbers in nodes indicate the posterior probability. The estimated dates of the MRCAs of the main lineages are indicated. The isolate that was whole genome sequenced using PacBio technology (MVF7) is highlighted in orange. The bottom plot represents the changes in the effective population size over time, with the shadowed area representing the 95% credible interval, following the same timescale as the tree.

Figure 3. The position of *Staphylococcus aureus* subsp. *anaerobius* within the *S. aureus* diversity. Maximum Likelihood tree constructed from a SNP alignment of the studied *S. aureus* subsp. *anaerobius* sequences (clades in blue) and 787 sequences of different *S. aureus* subsp. *aureus* (in black). (A): unrooted tree showing the divergence of 17 sequences of other *Staphylococcus* (*S. schweitzeri* and *S. argenteus*) whereas *S. aureus* subsp. *aureus* is embedded in the *S. aureus* diversity. (B): a subtree showing the position of the *S. aureus* subsp. *anaerobius* clade (collapsed in blue) with respect to the other *S. aureus* subsp. *aureus* clonal complexes (CC).

A

B

Figure 4. Characterisation of the transcriptional terminators (TTs) present in the ISs. (A) Schematic representation of IS-associated TTs. (B) Schematic representation of and functional assessment of the bi-directional TT1 localised in the 5' region of the IS. The bi-directional TT1 is composed of two sub-TTs that share the same sequence but are inverted repeats, thus generating a master bi-directional TT1. The expression of the *blaZ* reporter in plasmid pCN42 is controlled by a cadmium-inducible promoter. The identified transcriptional terminators were cloned between the promoter and the *blaZ* reporter gene. β-lactamase expression was monitored either in uninduced (black bars) or Cd-induced (open bars) cultures 180 min after induction. Data show the means of three independent biological replicates and error bars show the standard deviation from the mean. Statistical analysis was performed using two-way ANOVA followed by Dunnett's post-test. Multiplicity adjusted p-values are shown for uninduced to induced comparison and all terminator containing reporters showed significant differences when compared to their respective empty plasmid control. **** p<0.0001, ns not significant.

Figure 5. Presence of IS impacts on the expression levels of downstream genes and genes product. (A-D) Reporter constructs for presence of the IS upstream and in sense orientation of the target gene were designed in plasmid pCN41 containing a β-lactamase reporter gene. The position of the IS relative to the target gene's start codon is indicated by green triangle. All reporters were constructed either by removing the IS gene from strain *S. aureus* subsp. *anaerobius* strain MVF84 or by introducing the IS into *S. aureus* subsp. *aureus* strain RF122 to confirm the causative role of the IS on any changes to gene expression. Note that in (A) the removal of the IS results in a 207 bp deletion to the promoter region not present in strain RF122. Reporter plasmids were introduced into strain RN4220 and samples taken and processed as described in Materials and Methods. Data show the mean of three independent biological replicates and error bars represent the standard deviation from the mean. Statistical analysis was performed using One-Way ANOVA followed by Tukey's multiple comparisons test. *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$, *ns* not significant. (E&F) Western blot analysis of (E) *mgr*A and (F) *sle1* expression constructs to address whether presence of the IS impacts on MgrA and Sle1 expression levels by interrupting expression of an antisense transcript. (E) The *mgr*A gene encoding a 3xFLAG tag was cloned in antisense orientation to a cadmium-inducible promoter into plasmid pCN51 to mimic the expression of an antisense transcript to *mgr*A. Constructs either contained or lacked the *S. aureus* subsp. *anaerobius* IS, or contained a transcriptional terminator (TT) separating the IS transcript or expression from the plasmid-encoded cadmium-inducible promoter from the *mgr*A transcript. Expression of the cadmium-inducible promoter was induced by the addition of 5 µM CdCl₂. (F) To assess the impact of IS in the context of a natural antisense transcript, 3xFLAG-encoding expression constructs of *sle1* including the gene downstream of *sle1* in the ancestral genome were cloned into the plasmid pCN47. Samples were taken during different growth phases as indicated. Plasmids were introduced into the *S. aureus* strain RN4220 Δ*spa* and samples for expression analysis taken and processed as described in Materials and Methods. Western blots shown are representative of results obtained in at least two independent biological replicates. TT transcriptional terminator (E-F).

Figure 6. Map of the genomic translocations across the *Staphylococcus aureus* subsp. *anaerobius* isolate MVF7 chromosome. The green blocks represent the 6 large translocations detected in MVF7 when compared to the *Staphylococcus aureus* subsp. *aureus* RF122 strain (see also **Fig. S6** in the Additional File 1). The shaded areas indicate the original location of the translocated portions in the putative ancestral genome. The inner ring shows the GC skew (green for positive skew, purple for negative), which is unaffected by the translocations.