**Large-scale transcriptomics to dissect two years of the life of a fungal phytopathogen**

**interacting with its host plant**

**Authors:**

Elise J. Gay [1], Jessica L. Soyer [1], Nicolas Lapalu [1], Juliette Linglin [1], Isabelle Fudal [1],

Corinne Da Silva [2], Patrick Wincker [2], Jean-Marc Aury [2], Corinne Cruaud [3], Anne Levrel [4],

Jocelyne Lemoine [4], Regine Delourme [4], Thierry Rouxel [1], Marie-Hélène Balesdent [1]

[1] Université Paris-Saclay, INRAE, AgroParisTech, UMR BIOGER, 78850 Thiverval-Grignon,

France.

[2] Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Université

d'Evry, Université Paris-Saclay, 91057 Evry, France

[3] Genoscope, Institut François Jacob, CEA, Université Paris-Saclay, Evry, France

[4] INRAE, Institut Agro, Univ Rennes, IGEPP, 35653, Le Rheu, France.

.

## Abstract

The fungus *Leptosphaeria maculans* has an exceptionally long and complex relationship

with its host plant, *Brassica napus*, during which it switches between different lifestyles,

including asymptomatic, biotrophic, necrotrophic, and saprotrophic stages. The fungus is also

exemplary of "two-speed" genome organisms in which gene-rich and repeat-rich regions

alternate. Except for a few stages of plant infection under controlled conditions, nothing is

known about the genes mobilized by the fungus throughout its life cycle, which may last

several years in the field. We show here that about 9% of the genes of this fungus are highly

expressed during its interactions with its host plant. These genes are distributed into eight

well-defined expression clusters, corresponding to specific infection lifestyles or to tissue-

specific genes. All expression clusters are enriched in effector genes, and one cluster is

specific to the saprophytic lifestyle on plant residues. One cluster, including genes known to be involved in the first phase of asymptomatic fungal growth in leaves, is re-used at each asymptomatic growth stage, regardless of the type of organ infected. The expression of the genes of this cluster is repeatedly turned on and off during infection. Whatever their expression profile, the genes of these clusters are located in regions enriched in heterochromatin, either constitutive or facultative. These findings provide support for the hypothesis that fungal genes involved in niche adaptation are located in heterochromatic regions of the genome, conferring an extreme plasticity of expression. This work opens up new avenues for plant disease control, by identifying stage-specific effectors that could be used as targets for the identification of novel durable disease resistance genes, or for the in-depth analysis of chromatin remodeling during plant infection, which could be manipulated to interfere with the global expression of effector genes at crucial stages of plant infection.

## Author Summary

Fungi are extremely important organisms in the global ecosystem. Some are damaging plant pathogens that threaten global food security. A knowledge of their biology and pathogenic cycle is vital for the design of environmentally-friendly control strategies. Unfortunately, many parts of their life cycle remain unknown, due to the complexity of their life-cycles and technical limitations. Here, we use a rapeseed pathogen, *Leptosphaeria maculans*, which has a particularly complex life-cycle, to show that large-scale RNA-Seq analyses of fungal gene expression can decipher all stages of the fungal cycle over two years of interaction with living or dead hosts, in laboratory and agricultural conditions. We found that the fungus uses about 9% of the genes of its genome specifically during interactions with the plant, and observed waves of extremely tight, complex regulation during the colonization

49  of specific tissues and specific parts of the life-cycle. Our findings highlight the importance of

50  genes encoding effectors, small secreted proteins manipulating the host. This work opens up

51  new avenues for plant disease control through the identification of stage-specific effectors

52  leading to the discovery of novel durable disease resistance genes, or the analysis of

53  epigenetic regulation, which could be manipulated to interfere with effector gene expression.

## Introduction

Fungi play a vital role in global ecosystems. They have highly complex life-cycles that

often remain poorly understood, due to the multiplicity of fungal species and their behavior.

Plant-associated fungi (endophytes, symbionts and phytopathogens) contribute to this

complexity and diversity of behavior, as they may have beneficial, neutral or detrimental

interactions with their hosts; some are of paramount importance in favoring the growth of

plants or their adaptation to the environment, whereas others are highly damaging pathogens

of cultivated plants [1].

Many fungal pathogens of annual crops display contrasting infection strategies (e.g. the

biotrophs *Blumeria graminis* and *Ustilago maydis,* the hemibiotroph *Colletotrichum*

*higginsianum* or the necrotroph *Botrytis cinerea*), with reproducible short cycles of

colonization/sporulation, lasting from hours to weeks, on plant leaves. For such species, high-

throughput RNA sequencing has made it possible to describe the stage-specific expression of

genes during the few days or weeks over which the infection process occurs, under laboratory

conditions. Such studies have identified concerted waves of pathogenicity gene expression

during fungal infection [2-3], sets of genes involved in the switch from asymptomatic to

necrotrophic growth in the hemibiotrophic fungus *Zymoseptoria tritici* [4,5], or four waves of

expression for genes encoding putative effectors in *C. higginsianum* [6]. Short life-cycles, of

no more than a few weeks on the host, are compatible with the description of biological and

phytopathogenic features in experiments performed in controlled conditions.

In many other models, including symbionts and endophytes, the fungi spend much

longer interacting with their host, and make use of different pathogenic/symbiotic

developmental programs to adapt to different lifestyles during host colonization (this is the

case in *Leptosphaeria maculans*, the fungus studied here), different host tissues (e.g. *U.*

*maydis*; [7]), or different plant species (e.g. rust species with complex pathogenic cycles

including two alternate host species; [8]). For these species, RNA-Seq approaches also seem

to be the approach of choice for dissecting the entire fungal life-cycle and describing the

biology of the fungus, together with the sets of genes involved and their regulation. However,

the use of RNA-Seq to describe the fungal life-cycle is challenging and technically

demanding. Even for fungal species that can be grown in axenic media and for which

miniaturized pathogenicity assays have been developed, parts of the pathogenicity cycle

remain inaccessible and cannot be reproduced in laboratory conditions. It is, therefore,

essential to sample and monitor isolates from the wild. The second challenge is developing a

sufficiently comprehensive *a priori* knowledge of all the relevant stages of the fungal life-

cycle, and a means of sampling all of these stages for analysis. The third challenge is the

generation of sufficient amounts of high-quality samples (often corresponding to minute

amounts of fungal organisms in large amounts of plant material, or samples obtained blind,

without knowing whether the fungus is present in the plant at the time of sampling) for RNA-

Seq, to ensure the generation of enough reads for appropriate statistical analyses. The fourth

challenge is coping with complex field samples, containing not only the phytopathogen and

the host plant, but also many other fungal or bacterial species, necessitating a careful

evaluation of the specificity of RNA-Seq mapping before analysis. The final challenge is

coping with field samples subject to natural infection in variable environmental conditions,

potentially resulting in a high level of heterogeneity in terms of the presence of the fungus and

its developmental stage within the tissues, and attaining a biologically relevant continuity

when monthly samples are obtained over a whole growing season.

*L. maculans*, a pathogen of rapeseed (*Brassica napus*), has an exceptionally complex

life cycle compared to most other fungal phytopathogens attacking annual plants. It can infect

different plant tissues, undergo multiple switches from biotrophy to necrotrophy during a

lengthy lifespan in plant tissues, and can also live as a saprophyte on plant residues. The

epidemiological cycle of *L. maculans* is well described [9,10]. It begins with the

hemibiotrophic colonization of young leaves by spores generated by sexual reproduction

(ascospores) in early autumn (October-November) in Europe. The fungus first colonizes the

leaf tissues as a biotroph, for a few days or weeks, depending on the climatic conditions,

without causing any symptoms. It then induces the development of necrotrophic leaf lesions

in which its asexual spores (conidia) are produced. The fungus then migrates, without causing

symptoms, from the petiole to the stem, where it lives in the plant tissues, as an endophyte, for

several months. Finally, at the end of the growing season (May to July in Europe), it switches

back to necrotrophic behavior, inducing the formation of a damaging stem canker that may

result in plant lodging. Having completed all these stages of infection on living plant tissues,

*L. maculans* then switches to a saprotrophic lifestyle, living on crop residues for up to three

years. It develops structures for sexual reproduction to create the new inoculum (ascospores)

for subsequent seasons on these residues. Due to its length and complexity, only a limited part

of the life-cycle of *L. maculans* is amenable to laboratory experiments. Most of the RNA-seq-

based time-course experiments published to date were performed during the first two weeks

of infection, on cotyledons in controlled conditions [11-13], or in one set of stem infection

conditions in controlled conditions [14].

In many models analyzing primary leaf infection, RNA-Seq approaches have

highlighted the importance of genes encoding small secreted proteins (SSPs), acting as

effectors, which often compromise plant defense responses [15]. In *L. maculans*, SSP genes

are strongly expressed during cotyledon infection in controlled conditions [11-13], whereas

another set of putative effectors and a toxin encoded by a secondary metabolite gene cluster,

sirodesmin PL, are recruited during stem infection [14,16]. To date, 20 effector genes have

been characterized in *L. maculans*, corresponding to the most documented class of

pathogenicity genes in this species. These genes include nine cloned "early" effector genes

(*AvrLm* genes) [17-25] located in AT-rich regions of the fungal genome enriched in

transposable elements (TEs) [26] and encoding typical SSPs with few, if any, matches to

sequences in databases, frequently enriched in cysteine residues. These genes are expressed

and act during asymptomatic cotyledon infection. Eleven "late" putative effector genes, with

molecular features similar to those of "early" effectors, were then shown to be expressed

during stem colonization [14]. Unlike "early" effectors, these genes are located in gene-rich

regions of the genome.

In addition to effector genes, the possible role in virulence of dozens of genes of

different functional categories has been studied in *L. maculans*. These genes encode proteins

involved in the biosynthesis of two secondary metabolites: abscisic acid (ABA) [27] and the

toxin sirodesmin [16], and genes disrupted or silenced in *L. maculans,* 11 of which have been

implicated in fungal virulence [28-36]. Again, all these studies were performed in controlled

conditions, generally with miniaturized pathogenicity tests on cotyledons, and nothing is

known about the involvement of the candidate genes, or of other pathogenicity genes, in other

parts of the fungal life cycle. 'Late' effectors have been identified only in controlled

conditions, and their involvement in the lengthy process of stem colonization in the field

requires further investigation.

*L. maculans* has a well-defined bipartite genome consisting of gene-rich, GC-

equilibrated regions and large AT-rich regions enriched in transposable elements (TE) but

depleted of genes. The AT-rich regions account for 34.8% of the genome and are enriched in

H3K9me3 (trimethylation of the lysine 9 residue of histone H3) marks, typical of constitutive

heterochromatin in other species, whereas the GC-rich regions are enriched in H3K4me2

(dimethylation of the lysine 4 residue of histone H3) marks, typical of euchromatin [37-38].

In addition, species-specific genes and genes silenced during axenic growth display an

enrichment in H3K9me3 and H3K27me3 (trimethylation of the lysine 27 residue of histone

H3; [38]) heterochromatin modifications. The chromatin-based regulation of the expression of

'early' effector genes has been experimentally validated for a few *AvrLm* genes in this model

[37].

We hypothesized here that, with the use of careful experimental and sampling

procedures, it should be possible to use a large-scale transcriptomic approach to decipher the

whole of the complex life cycle of *L. maculans* in interaction with its host, in both controlled

and field conditions, and over time scales ranging from weeks (controlled condition

experiments) to months or years (follow-up of plant infection on leaves and stems for a whole

growing season, survival of *L. maculans* on residues remaining on the soil for one year). Our

objective was, thus, to use RNA-Seq data to describe the complete fungal life-cycle and

pathogenicity cycle and to identify the genes and functions involved in each stage. We found

that about 9% of the genes of the fungus were expressed specifically during host-fungus

interactions. A clustering of all the genes involved in the host-fungus interaction led to the

detection of eight major waves of gene expression that could be consistently related to the

consecutive stages of the *L. maculans* life-cycle, to tissue-specific expression, or

corresponded to genes recycled for all similar lifestyles at different stages of the fungal life-

cycle. The regulation of expression was strongly associated with the hosting of genes of the

eight waves in heterochromatin landscapes, and heterochromatin dynamics is thought to be

instrumental in allowing the recycling of expression for genes from the same wave during all

stages of fungal biotrophy.

## Results

**Biological and RNA samples representing all aspects of the life-cycle of the fungus in**

**interaction with its host**

The RNA-Seq experiment was designed to generate samples corresponding to all stages

of the interaction of *L. maculans* with its host plant, either alive (**Fig 1E-C**) or dead (stem

residues after harvest; **Fig 1D**), in controlled conditions or in field experiments under natural

inoculum pressure, over periods of time ranging from a few days to months or years (**Fig 1**;

For a detailed description of biological samples, see the M&M section). As a reference, we

included a series of samples grown in axenic conditions promoting a particular fungal

developmental stage (mycelium growth, pycnidium differentiation, pseudothecium

differentiation, resting conidia, conidial germination) (**Fig 1E**), to distinguish between these

fundamental biological processes and those specific to interaction with the plant. In total, 102

biological samples, corresponding to 37 different sets of conditions, were subjected to RNA-

Seq. Several quality control checks and parameter optimization were required before this

heterogeneous dataset could be subjected to statistical analyses. *In fine,* 32 sets of conditions

were retained for statistical analyses, whereas five sets of conditions were excluded because

too few fungal reads were available (**S1 Text**, **S1** and **S2 Tables**).

**Fig 1. Conditions and samples used for the RNA-Seq study of the *Leptosphaeria maculans* life cycle.**
Time-course studies are represented by an arrow and the sampling time points are indicated. (A) RNA was extracted from cotyledons infected with ascospores one or two days post inoculation (DPI). Another time-course study was conducted by inoculation with conidia, with RNA extraction at six time points, from 2 to 15 DPI. Young leaves from naturally infected fields were sampled 2 or 2.5 months post sowing (MPS). (B) Petiole colonization was achieved on petioles inoculated in controlled conditions, with RNA extraction at 7 and 14 DPI. (C) Stem colonization samples were obtained in controlled conditions, at 14 and 28 DPI. Stem base samples from naturally infected fields were collected and used for RNA extraction every two to three months over an entire growing season. (D) Leftover stem residues were sampled every two to three months for one year after harvest (dates expressed in months post harvest, MPH). (E) We used 10 sets of *in vitro* growth conditions, based on two different media (liquid or solid) and corresponding to different physiological states (mycelium, sporulation, germinating and resting conidia, *in vitro* crossing conditions). The samples highlighted in bold did not provide enough fungal RNA-Seq reads for statistical analysis.

**Quality control analyses highlight the consistency of the RNA-Seq dataset**

Clustering based on pairwise Pearson's correlation analyses of the 32 sets of conditions

identified four groups of clustered conditions (**Fig 2**). In each of these four groups, principal

component analysis (PCA) showed a low level of variation between biological replicates,

except for a few field samples and *in vitro* crosses (**S2 Text**, **S1A-D Fig**). The four clusters

(**Fig 2**) corresponded to: (i) the early infection stage on cotyledons and leaves and

colonization of the petiole (group 2), (ii) the late infection stage on stems (group 1), (iii) the

saprophytic lifestyle (group 3), in controlled conditions or in the field, and (iv) all the *in vitro*

growth conditions promoting the differentiation of pseudothecia and/or pycnidia (group 4).

Within each group, lifestyle provided a second level of sample discrimination, with

samples from biotrophic or endophytic stages distinguished from necrotrophic-stage samples

(**Fig 2**). For example, within group 2, samples displaying necrotrophic behavior on petioles 14

days post infection (DPI) were more closely related to the necrotrophic samples collected

from cotyledons 12 and 15 DPI than to the corresponding biotrophic samples (petioles 7 DPI)

(**Fig 2**). Similarly, in group 1, the residues collected immediately after harvest grouped with

the last samples collected from the stem base, even though the residues did not originate from

the same growing season, or from the same rapeseed variety, suggesting that the transition

from stem necrosis to saprophytic life takes place after harvest (**Fig 2**).

Some axenic growth conditions were also grouped with some *in planta* conditions.

Within group 2, germinating conidia grouped with the earliest samples collected from

cotyledons, consistent with the use of conidia to inoculate plant tissues in controlled

conditions (**Fig 2**). Samples grown in axenic culture on solid V8 medium under conditions

promoting mycelial growth over sporulation clustered within group 3, indicating that mycelial

growth is a major component of the saprophytic lifestyle in the wild. By contrast, mycelium

grown in static liquid medium grouped with all conditions in which the fungus colonized stem

tissues, possibly reflecting growth under low-oxygen conditions in plant vessels or

intercellular spaces.

**Fig 2. Hierarchical clustering of Leptosphaeria maculans gene expression in 32 conditions representing the fungal life cycle**

Each gene with an FPKM count > 2 in at least one set of conditions was retained for the correlation analysis. A pairwise Pearson's correlation analysis was performed between the 32 sets of conditions on Log2(FPKM+1) values. The resulting Pearson's correlation matrix was subjected to hierarchical clustering (represented by the tree) with the ward.D2 method. The Pearson's correlation results are shown on a color scale. Four groups of clustered samples were identified (GP1 to GP4). The experimental conditions are indicated as IPF, in planta field conditions; IPC, in planta controlled conditions; IV, in vitro conditions and RF, residues in field conditions. The infected plant tissues collected or the conditions under which the in vitro samples were obtained are detailed, together with the corresponding time points for sampling (DPI, days post-inoculation; MPS, months post-sowing; MPH, months post-harvest).

**Analytical strategy for identifying genes involved in pathogen-plant interactions**

For the identification of fungal genes specifically involved in at least one stage of the

pathogen-plant interaction, we used the 10 sets of *in vitro* growth conditions as 10 sets of

control conditions, to exclude all genes involved in basic fungal metabolism or life traits other

than those associated with plant infection (**Fig 3**). Using very stringent criteria (LogFC > 4; *p*

value < 0.01), we focused on genes specifically overexpressed *in planta* relative to *in vitro*

control conditions. We analyzed the expression patterns of these genes, their functional

annotation, their genomic localization, and their location with respect to histone modifications

during axenic growth. We then focused on genes encoding SSPs from the new repertoire

generated in this study (**S3 Text**). Finally, the expression profiles of co-expressed SSP gene

were used as a template for the identification of waves of genes displaying a high level of co-

regulation (R-squared > 0.80) an exemplifying specific pathogenic strategies (**Fig 3**).

**Fig 3. Pipeline used to describe *Leptosphaeria maculans* genes overexpressed in at least one**
**condition *in planta***
We found that 1,207 of the 13,047 annotated genes were overexpressed in at least one of the 22 sets of
conditions *in planta* relative to the 10 sets of *in vitro* conditions. These genes included 272 predicted
to encode small secreted proteins (SSPs). We analyzed the set of 1,207 genes to identify the major
expression profiles by gene expression clustering. We defined eight expression clusters (clusters 1 to
8). For each cluster, we analyzed expression patterns, functional enrichment, and the genomic
localization of the genes. For the specific detection of highly coregulated waves of SSPs, we
performed a linear regression analysis of the mean expression value for each SSP subset present in
each cluster against the expression of all 13,047 genes. This analysis identified 95 SSP genes and 185
non-SSP genes as highly coregulated genes.

**Genes overexpressed during interaction with the plant define eight biologically relevant**

**expression clusters**

Relative to axenic growth conditions, 1207 genes were found to be overexpressed in at

least one set of conditions *in planta*; thus, 9.2% of *L. maculans* genes are specifically

mobilized during the interaction of the fungus with its host, alive or dead. Eight expression

profiles were identified, referred to hereafter as clusters 1 to 8, defining a complex landscape

of expression profiles (**Fig 4**). We analyzed the enrichment of each cluster in GO terms,

focusing on CAZymes (Carbohydrate-active enzymes) associated with degradation activities,

to identify the functions and pathways associated with each expression profile (**S2-S5 Fig**).

**Fig 4. Heatmap obtained by clustering the 1,207 *Leptosphaeria maculans* genes upregulated *in planta***

For each set of conditions, we calculated the mean FPKM value of the replicates, which we then subjected to $Log_2$ transformation and scaling. The self-organizing map method was used on the $Log_2(FPKM+1)$ values, to define eight expression clusters (clusters 1 to 8) for the 1,207 genes. The level of expression ranged from 4 (yellow) to -4 (green). Genes located in AT-rich regions (AT) or encoding small secreted proteins (SSPs) are indicated by red and black lines, respectively, on the left. The conditions are described in the legend to the *x* axis of the heatmap. Three sample features are described (i) the experimental conditions: IPF, *in planta* field conditions; IPC, *in planta* controlled conditions; RF, residues in field conditions, (ii) the type of plant tissue sampled and (iii) the sampling time points (DPI, days post inoculation; MPS, months post sowing; MPH, months post harvest. The ten sets of *in vitro* conditions are ordered as follows: conidia (a. non-germinating, b. germinating); incompatible mating conditions (c. JN2XJN2 7days (d), d. JN2XJN2 20d, e. JN2XJN2 35d); compatible mating conditions (f. JN2XNZT4 7d, g. JN2XNZT4 20d, h. JN2XNZT4 35d); mycelial growth (i. in liquid medium, j. on solid medium).

Clusters 1, 2 and 3 included genes expressed sequentially during cotyledon infection and were representative of three consecutive stages occurring during the two-week period of cotyledon colonization by *L. maculans*. Cluster 1 encompassed 97 genes highly expressed at the first stages of cotyledon infection, when the conidia germinate and the hyphae penetrate the wounded plant tissues (2-5 DPI). It was enriched in GO terms corresponding to carbohydrate metabolic processes ($p = 1\mathrm{x}10^{-2}$) and catalytic activities ($p = 1\mathrm{x}10^{-3}$), including peptidase and hydrolase activities (all $p \leq 1\mathrm{x}10^{-2}$). An enrichment in cutinase activity ($p = 1\mathrm{x}10^{-2}$), specific to this cluster, was also observed (**S2-S3, S5 Figs**).

The 148 genes of cluster 2 displayed peaks of expression during all stages of asymptomatic growth within plant tissues (cotyledon colonization 5-9 DPI, first stage of petiole colonization, first two dates of stem colonization in the field). The genes of this cluster were annotated with few GO terms or functions, but were strongly enriched in SSP genes ($p = 2\mathrm{x}10^{-16}$; **S3 Fig**). Cluster 3 encompassed 87 genes highly expressed during the necrotrophic stage on both cotyledons (9-15 DPI) and petioles, but not during the necrotrophic stem canker stage in the field. This cluster was enriched in carboxylic ester hydrolase activities ($p = 3\mathrm{x}10^{-3}$) suggesting a role in degradation activities specifically targeting the plant cell wall.

Cluster 4 encompassed 238 genes highly expressed during the shift from biotrophy to necrotrophy in cotyledons (7-9 DPI) and in stem bases infected in the field (7-8 months post-sowing, 'MPS'). This cluster displayed the strongest enrichment in catalytic processes ($p = 2\mathrm{x}10^{-10}$) and hydrolase activities ($p = 1\mathrm{x}10^{-12}$), and a specific enrichment in proteolysis processes ($p = 2\mathrm{x}10^{-2}$; **S3 Fig**). It also displayed the largest number of CAZymes (63 genes; **S5 Fig**) targeting different substrates, mostly plant cell-wall components, such as pectin, arabinose and galactan (**S5 Fig**).

Two clusters (cluster 5 and 6) displayed an increase in expression late in stem infection,

but differed in terms of the timing of expression and the functions involved. Cluster 5

contained 128 genes highly expressed during the asymptomatic colonization of stems in the

field (at 7-8 MPS) and on residues immediately after harvest. It was enriched in catalytic

activities ($p = 1\times10^{-3}$), mostly linked to saccharide degradation, suggesting involvement in

activities relating to the uptake of sugar resources (carbon-oxygen lyase activity acting on

polysaccharides; hydrolase activity; cellulose binding; **S3 Fig**). Cluster 6 contained 141 genes

highly expressed during colonization of the stem base in the field, with maximal expression at

10 MPS, corresponding to the time at which stem canker develops. This cluster was enriched

in catalytic activities ($p = 2\times10^{-4}$) but was also linked to cofactor binding ($p = 3\times10^{-6}$),

monooxygenase ($p = 1\times10^{-2}$), and oxidoreduction activities ($p = 5\times10^{-5}$), involved in

detoxification processes (**S3 and S4 Figs**). The higher proportion of xylanase genes in cluster

6 than in cluster 5 (**S5 Fig**) suggests a change in the mode of nutrition during the necrotrophic

stage on stems.

Finally, clusters 7 and 8 were specific to saprophytic behavior on stem residues. The

198 genes in cluster 7 showed displayed an increase in expression from the last sampling on

necrotic stems until the last sampling on stem residues one year after harvest. This cluster was

enriched in genes relating to mono-oxygenase activities ($p = 1\times10^{-4}$) and contained the largest

number of genes encoding LMPOs (lytic polysaccharide monooxygenases), cellulases and

xylanases. Cluster 8 encompassed 170 genes specifically expressed during all stages

occurring on senescent or dead plant tissue, with the highest levels of expression during

saprophytic growth on residues. It was also enriched in oxidoreductase processes ($p = 1\times10^{-3}$),

in response to oxidative stresses and oxidant detoxification associated with an enrichment in

catalase ($p = 1\times10^{-3}$) and peroxidase ($p = 3\times10^{-3}$).

The different clusters of gene expression thus can be designated as follows: "Penetration

and establishment" (cluster 1), "Biotrophy" (cluster 2), "Cotyledon and petiole necrotrophy"

(cluster 3), "Biotrophy to necrotrophy transition" (cluster 4), "Stem biotrophy" (cluster 5),

"Stem necrotrophy" (cluster 6), "Stem canker and saprophytism" (cluster 7) and

"Saprophytism" (cluster 8).

**Contribution of candidate pathogenicity genes to fungal life in interaction with the plant**

**1- Genes involved in fungal pathogenicity in other fungal species.** Several genes

previously shown to be involved in fungal pathogenicity were found in one of the eight

expression clusters, and were mostly associated with biotrophic stages of the interaction. One

RALF (rapid alkalinization factor) gene known to favor fungal infection [39] was identified in

cluster 1 (Lmb_jn3_05329). Two of the six salicylate hydroxylase genes, known to

compromise plant salicylic acid signaling in other models [40], were found in clusters 1

(Lmb_jn3_12582) and 2 (Lmb_jn3_11892). Three of the four LysM-domain containing genes

of the genome, known to protect chitin from degradation by plant enzymes or to scavenge

chitin oligomers [41], were found in clusters 2 (two genes, Lmb_jn3_00461,

Lmb_jn3_10047) and 6 (one gene, Lmb_jn3_04300). Three isochorismatase genes

(Lmb_jn3_11641, Lmb_jn3_10045, Lmb_jn3_06552), interfering with plant salicylate and

jasmonate defense signaling in other phytopathogens, were present in clusters 1, 4 and 6,

respectively. Finally, one of the two genes encoding NEP (necrosis and ethylene-inducing

protein), frequently associated with necrotrophic behavior [42], was found to be

overexpressed *in planta*, and belonged to cluster 4 (Lmb_jn3_10035).

**2- Secondary metabolite gene clusters.** The genome of *L. maculans* contains 22 genes

predicted by SMURF tools to encode secondary metabolite enzymes (SM) and annotated as

"polyketide synthase" (PKS, 11) "non-ribosomal peptide synthetase" (NRPS, 5), or "others"

(dimethylallyltryptophan synthase, 1; NRPS-like, 4; PKS-like, 1). Seven of the 1207 genes in

the eight expression clusters encoded SM enzymes from the PKS or NRPS family. Four of

these genes were highly expressed during necrotrophic stages of the life-cycle within the stem

(cluster 6), whereas the other three were associated with clusters 2, 4 and 5 (**S3 Table**).

Among the SM genes analyzed in previous studies [27,43-44], the PKS gene involved

in abscisic acid (ABA) synthesis was linked to biotrophic behavior (cluster 2), whereas genes

involved in the synthesis of the toxins sirodesmin PL and phomenoic acid were associated

with stem necrotrophy, together with four other PKS genes and one NRPS gene (cluster 6)

(**S3 Table**). Secondary metabolite enzymes encode an SM backbone that may be modified by

tailoring enzymes, forming complex secondary metabolite gene clusters (SMGC). Here, only

three of the seven SM enzyme genes overexpressed *in planta* showed co-regulation with all or

part of surrounding genes of their SMGC: (i) the PKS responsible for ABA production (all

genes of the ABA SMGC being included in cluster 2), (ii) the NRPS responsible for

sirodesmin PL production (16 of the 25 genes of the SMGC included in cluster 6), and (iii) an

unknown NRPS-like gene, co-expressed with nine other surrounding genes also included in

cluster 6 (**S6 Fig**).

**3- Genes involved in *L. maculans* pathogenicity.**  Thirty-six of the 52 genes previously

identified as candidate pathogenicity genes in *L. maculans* (**S4 Table**) did not belong to the

SM or avirulence effector gene (see below) categories. Fifteen of these genes were identified

within a single cluster (**S4 Table**): cluster 1 (2 genes), cluster 2 (5 genes), cluster 3 (1 gene),

cluster 4 (4 genes), cluster 5 (2 genes) or cluster 7 (1 gene). Nine had previously been selected

as putative pathogenicity genes on the basis of their overexpression during infection, for

inactivation with CRISP-cas9, but the inactivation of these genes individually resulted in no

visible pathogenicity defect at the cotyledon stage [45] (**S4 Table**). Only one gene, encoding a

3-ketoacyl-thiolase (cluster 1), was shown to be involved in pathogenicity, but the mutant also

displayed impaired *in vitro* growth and germination [28] (**S4 Table**). The other five genes

have not been validated by functional approaches and are, thus, still considered to be

candidate pathogenicity-related genes. Four of these genes (three GH genes and one ABC

transporter) were included in the late expression clusters 4, 5 and 7, suggesting that they may

be *bona fide* pathogenicity genes (**S4 Table**).

By contrast, another 21 putative pathogenicity genes did not belong to any of the

expression clusters. Nineteen had previously been analyzed in functional studies (knockout or

silenced mutants), and 10 were shown to induce a loss or reduction of pathogenicity following

inactivation (**S4 Table**). However, the inactivation of six of these genes also induced defects

in growth, sporulation or germination [30,33,36,45-46], suggesting a general effect on

different stages of the fungal life-cycle with no specific overexpression during interactions

with the plant.

**4- Validated or candidate SSP effectors of *L. maculans*.** SSPs have been identified in *L.*

*maculans* and were classified as "early" effectors (including all known avirulence proteins,

inducing an avirulence phenotype when matching the corresponding resistance genes) or

"late" candidate effectors by Gervais *et al*. [14]. The role of avirulence proteins as effectors

suppressing plant defense responses or increasing/decreasing the size of cotyledon lesions has

been investigated in a few cases following comparisons of near-isogenic *L. maculans* isolates,

gene silencing or complementation (**S4 Table**, [21-22,47-48]). The most recently published

version of the genome assembly of *L. maculans* isolate JN3 ([49]; **S4 Text**) included a

number of major changes. We therefore produced a new repertoire of genes encoding putative

SSPs, encompassing 1,070 genes (8.2% of the entire *L. maculans* gene set) (**S3 Text**, **S7 Fig,**

**S5 and S6 Tables**).

All expression clusters displayed a significant enrichment in genes encoding SSPs (**S3**

**Fig**). We found that 272 of the 1207 genes upregulated during the *L. maculans* interaction

with *B. napus* (22.5%) encoded SSPs. This enrichment was limited in clusters 1, 3, 6, 7 and 8,

which contained only 14-17% of SSP-encoding genes ($0.013 < p < 3.10^{-5}$), but three other

clusters displayed a much higher level of enrichment in genes encoding SSPs: cluster 4

(23.1%, $p = 7x10^{-16}$), cluster 5 (31.2%, $p = 2x10^{-16}$) and cluster 2 (45.3%, $p = 2x10^{-16}$).

Interestingly, both clusters 2 and 5 are associated with biotrophic behavior of the fungus. All

the currently known *AvrLm* genes belonged to the "Biotrophy" cluster (cluster 2) (**S4 Table**).

This cluster displayed sequential rounds of overexpression/repression, with three peaks of

overexpression occurring during the three biotrophic stages (5-9 DPI on cotyledons, leaf

infection in the field; 7 DPI on petioles; 6-7 MPS in field plant stems) (**Fig 5; S8 Fig**).

The ten "late" effectors, expressed during stem colonization in controlled conditions

[14] were scattered over four different clusters (3, 4, 5, 6), along with a series of other genes

encoding SSPs (cluster 3, 14 SSPs; cluster 4, 55 SSPs; cluster 5, 40 SSPs and cluster 6, 22

SSPs) (**Fig 5**). The "late" effectors in clusters 5 and 6 were specific to stem colonization (and

the first stages of saprotrophy on residues), but those in clusters 3 and 4 were also expressed

during cotyledon infection. These "late" effector candidates were associated with either

biotrophic behavior on stems (cluster 5) or necrotrophic behavior on cotyledons and stems

(clusters 3 and 6).

Clusters 1, 7 and 8, were also found to be enriched in genes encoding SSPs (*p* value <

$5x10^{-3}$). They contained 16, 33 and 24 SSP genes, respectively. Candidate effectors may,

therefore, be mobilized very early in colonization (cluster 1) and during saprophytic life

(clusters 7 and 8) (**S8 Fig**).

**Fig 5. Expression of known and putative effectors of *Leptosphaeria maculans* within the eight expression clusters**

The scaled Log$_2$(FPKM +1) expression values for the nine avirulence effector (*AvrLm*) genes, and the eight candidate late effector genes shown to be upregulated in at least one set of conditions *in planta* are represented and grouped according to cluster assignment. The mean scaled Log$_2$(FPKM +1) values of all the small secreted protein (SSP) genes in each cluster are indicated (black bold curve). The total number of SSP genes in each cluster is indicated. The conditions are described in the legend to the *x* axis. Three sample features are described: (i) the experimental conditions: IPF, *in planta* field conditions; IPC, *in planta* controlled conditions; RF, residues in field conditions. (ii) the type of plant tissue sampled and (iii) the sampling time points (DPI, days post inoculation; MPS, months post sowing; MPH, months post harvest.

Additional RNA-Seq samples (ascospores germinating on unwounded cotyledons 24 and 48 hours post inoculation; stem base samples 2, 3 and 5 MPS) were available. They provided too few *L. maculans* reads for statistical analyses, but these samples allowed us to confirm that genes belonging to cluster 2, and "early" effector genes in particular, were among the genes most strongly expressed both during ascospore germination on cotyledons and very early stages of stem colonization in winter (**S5 Text**, **S7 Table**).

**Histone methylation associated with gene regulation**

All cloned avirulence genes of *L. maculans* identified to date are associated with H3K9me3. Furthermore, H3K9me3 and H3K27me3 domains are significantly enriched in genes encoding effectors, regardless of their expression [38]. During axenic culture, 7,373 *L. maculans* genes are associated with H3K4me2, 104 with H3K9me3, 2,020 with H3K27me3 and 101 with the two heterochromatin histone modifications [38]. We combined the genome-wide histone maps previously generated *in vitro* with our analysis of gene expression during infection, to determine whether (i) all genes, (ii) only (putative) effector genes, or (iii) only certain subsets of genes upregulated *in planta* were associated with a particular chromatin landscape. H3K4me2-domains encompass 56% of *L. maculans* genes, but only 20% of the genes upregulated *in planta* were associated with H3K4me2 ($\chi^2$ test=662; $p < 2 \times 10^{-16}$). By contrast, the genes upregulated *in planta* were significantly enriched in genes located in an

H3K27me3 domain, as 44% of the upregulated genes were located in such a domain ($\chi^2$ test =

749, $p < 2x10^{-16}$). Strikingly, only 205 genes were located in H3K9me3 or H3K27me3

domains *in vitro*, but 85 of these genes were upregulated *in planta*. Thus, regardless of the

wave of expression concerned, the genes upregulated *in planta* were enriched in genes

associated with heterochromatin *in vitro*.Only cluster 2 was significantly enriched in genes

associated with the H3K9me3 histone modification (**Fig 6**). Interestingly, all clusters of genes

upregulated *in planta* relative to axenic culture were enriched in H3K27me3-associated genes

*in vitro,* whereas these clusters contained significantly fewer genes located in H3K4me2-

domains *in vitro* than the rest of the genome (**Fig 6**). The integration of RNA-seq and ChIP-

seq data revealed that the genes upregulated *in planta* were not randomly located in the

genome of *L. maculans*, with H3K4me2-domains significantly depleted of genes upregulated

during rapeseed infection, and both types of heterochromatin domains significantly enriched

in such genes.

**Fig 6. Association of *Leptosphaeria maculans* genes upregulated *in planta* with the histone modification marks.** The genome-wide histone map was generated by ChIP-seq analysis during axenic growth [38]. The proportion of genes associated with any of the histone modifications H3K4me2, H3K9me3 and H3K27me3 throughout the entire genome was used as a reference set for detecting the over- or under-representation of specific methylation marks in the eight gene clusters overexpressed *in planta*, in chi-squared tests (***: $p < 0.001$, **: $p < 0.01$). K4, H3K4me2; K9, H3K9me3; K27, H3K27me3.

**The co-expression of effector genes defines waves of co-regulated gene expression.**

Almost all the expression clusters were enriched in genes encoding SSPs. However within each cluster, the genes encoding SSPs could have different expression profiles, due to the clustering method (**S8 Fig**). We aimed here to increase the robustness of the detection of SSP expression waves. We defined eight SSP reference expression profiles by calculating the mean levels of expression for SSP genes within the eight clusters. We selected genes displaying significant co-regulation (SSP and non-SSP genes), by analyzing the linear regression between the expression patterns of the entire set of 13,047 genes and that of each SSP reference expression profile. Genes significantly co-expressed with the reference

expression wave tested ($R^2 > 0.80$, $p < 0.05$) were included in the new SSP expression waves

and were considered to be "highly coregulated genes".

For clusters 1, 3, and 8, no highly coregulated genes could be identified (**Fig. 7**). For the

other five clusters, waves of highly coregulated genes were identified, each containing five to

39 highly coregulated SSP genes (including 90 SSP genes from the 178 previously assigned

to these clusters), together with 185 highly coregulated genes encoding other types of proteins

(**Fig 7**). Two waves of expression, corresponding to clusters 5 and 6, displayed coregulation

of a limited number of SSP genes (seven and five, respectively, i.e. 17% of the SSP genes of

these clusters), together with 20 and 25 genes encoding other proteins, respectively. However,

only 15 (cluster 5) and 13 (cluster 6) of these genes belonged to the clusters (**Fig 7**),

suggesting that additional coregulated genes are either expressed in axenic conditions, or do

not display specific regulation during plant infection. For clusters 2, 4 and 7, a higher

proportion of highly coregulated SSP genes was found, with 58% (exact confidence interval

[45-70%]), 34% ([22-48%]) and 63% ([45-79%]) of the SSP genes of the clusters retained in

the waves of highly coregulated genes, respectively (**Fig 7**). Thus, clusters 2 and 7 displayed

the highest proportion of highly coregulated SSP genes, suggesting that the regulation of

expression is much tighter in these two clusters than in the others. The highly coregulated

genes in the "Biotrophy" wave (contained in cluster 2) included 39 SSPs, 23 (59%) of which

displayed no sequence matches to the NCBI nr protein database (**S8 Table**) and could be

considered *L. maculans*–specific effectors. These genes also included other genes mentioned

above, such as the genes of the ABA cluster, providing addition support for their role in the

biotrophic parts of the interaction. The highly coregulated genes in the "Stem canker and

necrotrophy" wave included most of the SSP genes of the cluster, but only two of these genes

(9.5%) were specific to *L. maculans*. This wave also included a large number of genes (24)

encoding CAZymes.

**Fig 7. Identification of highly coregulated *Leptosphaeria maculans* genes based on mean SSP gene expression *in planta***

The mean scaled $Log_2$(FPKM +1) values for small secreted protein (SSP) genes in each cluster (cluster 1 to 8) was used to create the eight reference expression profiles (black bold curve). The 22 sets of biological conditions used to average SSP gene expression are described in the legend to the *x* axis. Three sample features are described: (i) the experimental conditions: IPF, *in planta* field conditions; IPC, *in planta* controlled conditions; RF, residues in field conditions (ii) the type of plant tissue sampled and (iii) the sampling time points (DPI, days post inoculation; MPS, months post sowing; MPH, months post harvest). A linear regression analysis was performed to determine the correlation between the scaled $Log_2$(FPKM+1) value for the whole set of genes in the 22 sets of conditions and these eight reference expression profiles. A gene was considered to be highly correlated if its expression fitted the reference expression profile with an R-squared value $> 0.80$ and a corrected *p*-value $< 0.05$. The expression values of the resulting highly correlated gene subsets (SSP in blue and non-SSP in yellow) are plotted. The Venn diagrams show the number of genes (S: SSP, NS: Non-SSP) present in each cluster (green circles) and the number of genes identified as highly correlated with the corresponding expression profile (red circles). The asterisks indicate the significance of the SSP enrichment in the initial cluster (***: $p < 0.001$, **: $p < 0.01$).

## Discussion

In this paper, we used large-scale transcriptomic analyses to decipher the complex life cycle of a phytopathogenic fungus, to identify genes paramount for its interactions with the host plant and to shed light on the underlying regulatory mechanisms. In particular, given that many parts of the *L. maculans* life cycle remain poorly described, we focused on how the fungus copes with its different lifestyles, their timing and the transitions between them, and the regulation of gene expression throughout the long pathogenic/saprophytic life of the fungus. We generated RNA-Seq data for biological samples corresponding to the main relevant stages of infection in controlled conditions or in field conditions, and at different time scales (from days to years). Using this large dataset, we then focused on sets of genes mobilized exclusively when the fungus interacts with the plant. We used 10 different sets of axenic growth conditions promoting different aspects of the fungal life-cycle (e.g. vegetative growth, asexual sporulation, sexual reproduction) as basic expression features for the identification of pathogenicity genes *per se*, contrasting with the common practice of using only a few axenic conditions as controls [7,13-14] or performing pairwise comparisons of the infection time course [11]. We dealt with the variable number of fungal reads between

samples and the diverse *in vitro* conditions, by applying very stringent conditions to

determine whether a gene was specifically overexpressed (LogFC>4, $p < 0.01$). With these

criteria, we showed that 1,207 genes (less than 10% of the genes of the fungal genome) were

overexpressed during interactions with the living or dead plant relative to axenic conditions.

These genes may be considered the maximal set of genes mobilized by the fungus to infect

the plant without involvement in basal biological processes. These 1,207 upregulated genes

were consistently grouped into eight expression profiles, some of which displayed strong

coregulation of a number of the genes within the cluster. The sequential expression profiles

are consistent with the associated pathogenicity functions, confirming the relevance of this

RNA-Seq study performed over a large timescale.

**Transcriptome-based description of a complex fungal lifecycle**

Our RNA-Seq data highlighted and strengthened our knowledge of the biology of the

fungus all along its life-cycle, including for "obscure parts" of this cycle, as follows (Fig. 8) :

(i) Ascospores, which are difficult to generate for pathogenicity tests in controlled

conditions, are known to be the main source of infection in the field, germinating within a few

hours to produce a hypha that penetrates leaf tissues via stomata or wounds [50]. We found

that germinating hyphae very rapidly (2 DPI) produced "early" effectors (including known

AvrLm effectors), detected only 5-7 DPI as a massive peak of expression when inoculations

were performed with conidia ([11,13-14], this study). This result is consistent with

microscopy-assisted RNA-Seq analyses in other models, which have suggested that a few

hyphal tips produce early effectors to counter the initial defense reactions of the plant upon

contact [4,51]. This suggests that, like other plant-associated fungi, *L. maculans* manipulates

the plant as soon as it enters the tissues, to initiate its own growth within plant tissues, without

causing symptoms.

(ii) Following the infection of leaves or cotyledons and primary leaf spots development,

the hyphae grow within the intercellular spaces within tissues in advance of the leaf spots

towards and within the petiole and then enter the stem tissues [52-54]. Our RNA-Seq data

highlight the kinetics of this process and the lifestyle transitions occurring during cotyledon

infection, with three successive waves of gene expression related to different stages of

infection (penetration, biotrophy, and necrotrophy). Consistent with the biological data, the

pattern of gene expression during petiole colonization was similar to that in cotyledons, with a

symptomless stage during which the same set of genes was mobilized in cotyledons or

petioles (**Fig 8**).

(iii) We also found that isolates infecting leaves in the field had a pattern of gene

expression similar to that of a reference isolate used to inoculate cotyledons in controlled

conditions at 7 and 9 DPI, demonstrating that infections reproduced with routine protocols in

controlled conditions can accurately mimic some stages of field infection.

(iv) The monitoring of stem samples for a full year in field conditions provided new

information about the dynamics of stem colonization and necrosis in the field. We found that

the fungus was present very early in the stem base tissues, with fungal reads detectable as

early as December, as little as one to two months after the initial leaf infection. However,

levels of fungal transcriptomic activity remained very low between December and February,

and it was only at the end of winter (from March on) that the fungus displayed a quantifiable,

steady increase in transcriptomic activity.

(v) Fungal life within stem tissues, over a period of eight months, was characterized by

complex patterns of gene expression, with some of the genes concerned expressed exclusively

during stem colonization, highlighting the lengthy transition from biotrophy to necrotrophy

leading to stem canker (**Fig 8**). The gene expression patterns observed for stem infection at

six and seven months post sowing (March-April) were very similar to those observed for

biotrophic stem infection in controlled conditions at 14 and 28 DPI, again demonstrating that

controlled conditions adequately reproduce at least one stage of stem colonization in the field.

(vi) The fungus has been reported to survive on residues for between two and four

years, depending on the climatic conditions [50]. However, we currently know nothing about

how *L. maculans* survives as a saprobe on residues. A recent metabarcoding study showed

that *L. maculans* was the predominant fungal species on rapeseed residues, on which many

other fungal species, including other phytopathogens were also present, highlighting complex

interplay between the species [55]. Our RNA-Seq analysis of residues left on the soil for one

year confirmed these findings and systematically revealed transcriptomic activity of *L.*

*maculans* on the residues, with a mean of 22% of the total RNA-Seq reads corresponding to

*L. maculans* throughout the year, but with seasonal variations.

**Fig 8. Overview of the eight expression profiles detected during the infection cycle of _Leptosphaeria maculans_**

Each curve represents the mean expression value for the genes making up the eight clusters described in this study. The eight profiles are divided into three different plots, depending on the timing of their major expression peaks: (A) early infection and colonization of cotyledons, (B) late colonization of stems, (C) development on crop residues. The characteristics associated with each sample are indicated: mode of nutrition (B: biotrophy, N: necrotrophy, S: saprotrophy); the experimental conditions: (i) IPF, _in planta_ field conditions; IPC, _in planta_ controlled conditions; RF, residues in field conditions. (ii) the type of plant tissue sampled and (iii) the sampling time points (DPI, days post inoculation; MPS, months post sowing; MPH, months post harvest. We did not include field-infected leaves and the stem infected in controlled conditions, to prevent redundancy and generate a simplified model. The major gene functions identified in each gene cluster are shown on the right.

**Expression profiles and characteristics of genes specifically involved in interaction with the plant**

We observed successive or partly overlapping expression profiles, highlighting various life traits or feeding strategies specific to a particular lifestyle or colonized tissue (cluster 1, cluster 2, cluster 4, cluster 8), or with a combination of lifestyle/tissue specificities (cluster 3, cluster 5, cluster 6, cluster 7) (**Fig 8**). For example, necrotrophic behavior is tissue-specific, with genes expressed during cotyledon or petiole necrosis (cluster 3) or in stem tissues at the end of the growing season (cluster 6; **Fig 8**). By contrast, cluster 2 is involved in biotrophic behavior, whatever the tissue colonized.

The set of 1,207 genes specifically expressed during interaction with the plant is strongly enriched in candidate effectors, with enrichment in effectors observed for all expression clusters. As shown for cotyledon infection in other studies [13], and expanded here to other biotrophic or necrotrophic stages of plant infection, biotrophic behavior was associated with enrichment in very few, if any GO terms, whereas biotrophy-necrotrophy transitions were strongly enriched in genes involved in catalytic processes and CAZymes. Strict necrotrophic behavior within stems is associated with an overrepresentation of PKS-encoding genes (including the sirodesmin biosynthesis gene cluster) relative to other necrotrophic clusters.

The fungus expresses a set of 148 genes in all biotrophic/endophytic/asymptomatic stages of colonization (cluster 2). Interestingly, we found that the expression of these genes was up- and down-regulated on multiple occasions during the interaction with the plant. The recycling of the same set of genes for a similar lifestyle highlights the key role played by this set of genes in establishing symptomless growth within the plant, with at least one set of stem-specific genes (cluster 5) also associated with biotrophic behavior, but only during stem colonization. Cluster 2 (and, to a lesser extent, cluster 5) was found to be strongly enriched in

genes encoding candidate effectors, which are generally thought to act by compromising plant

immunity [15]. One of these candidate effectors, AvrLm10A, has been shown to compromise

leaf symptom development [23], and two others, *AvrLm4-7* and *AvrLm1,* interfere with SA

and ET signaling [47-48]. In addition, cluster 2 includes genes reported in various models to

be involved in the suppression of plant defense responses (salicylate hydroxylase or the ABA

cluster; [27,56-57]) or the scavenging of chitin oligomers, such as the LysM genes [41]. This

strongly suggests that cluster 2 plays an important role in hiding the fungus from the plant

surveillance machinery, or in compromising or suppressing plant defense responses to various

extents.

Our findings suggest that the previous classification of effectors as "early" or "late" [14]

is not relevant. "Early" effector genes were found to be expressed on several occasions during

symptomless cotyledon infection and colonization, the symptomless colonization of petioles

and during at least three months of symptomless growth in the tissues of the stem base.

Similarly, the "late" effector genes identified by Gervais *et al*. [14] and thought to be specific

to stem colonization according to experiments in controlled condition, were actually found in

four different expression clusters, only one of which was specific to biotrophic behavior in

stem tissues (cluster 5, **Fig 8**).

Importantly, cluster 2 contains all the genes encoding avirulence proteins identified to

date in *L. maculans*. During leaf or cotyledon infection, these avirulence proteins can be

recognized by the products of the major resistance genes (*Rlm* genes) of the plant, resulting in

complete resistance to avirulent isolates. The re-use of these same genes at other stages of

plant colonization strongly suggests that recognition can also take place during the systemic

growth of the fungus within petioles or stems, resulting in efficient resistance at these stages

of colonization, provided that the *Rlm* genes are constitutively expressed in all plant tissues.

In addition to the nine already cloned avirulence genes, cluster 2 contains 58 additional SSP

genes (39 of which are highly coregulated with all *AvrLm* genes), which are candidates of

choice for the screening of genetic resources for novel resistance genes.

We also found that effector gene expression was not restricted to interaction with the

living plant. Genomic studies on fungal saprotrophs focus on CAZymes and class-II

peroxidases [58]. We show here that *L. maculans* recruits a typical cocktail of CAZymes,

including the lytic polysaccharide mono-oxygenases (LMPO) involved in wood decay, but

also peroxidases, cytochrome P450 and stress response genes, during one year of life on crop

residues. In addition, cluster 7 encompasses 21 SSP genes specifically expressed during

saprophytic growth on stem residues (**Fig 8**). It has been suggested that effector genes arise

from genes used by saprobes to suppress ecological competitors [59]. The steady expression

of a series of coregulated candidate effector genes during many months of life on residues,

regardless of the changes in the mycoflora present on the residues over time [55], may

indicate a need for the fungus to produce a minimal set of molecules involved in ecological

competition.

**Heterochromatin-based regulation of expression as a key player in the lifetime**

**coregulation of genes involved in pathogenicity**

One of the key findings of our work is that all clusters of genes overexpressed during

interaction with the host are significantly associated with localization within a

heterochromatin domain during axenic growth. Evidence is accumulating, for many other

plant-associated fungi, regardless of their lifestyles and modes of nutrition, that the genes

involved in fungus-plant interaction are often present in regions of heterochromatin within the

genome (e.g. in [60-62]). Genes highly expressed during infection are often associated with

heterochromatin during axenic culture (e.g. in *Z. tritici* or *L. maculans* [4,37,63]) and histone

modifications (H3K9me3 and / or H3K27me3) have been shown to play a major role in

regulating secondary metabolite gene clusters or effector genes (e.g. in *F. graminearum*, *F.*

*fujikuroi*, *Z. tritici*, *Epichloe festucae* [64-68]). In the endophyte *E. festucae* and the pathogen

*Z. tritici*, the derepression of putative effector genes or secondary metabolite gene clusters

upon infection is associated with an underlying dynamic of histone modifications, which has

been experimentally validated [64,67]. These data suggest that there is a generic chromatin-

based control of plant infection mechanisms. We previously showed that, in *L. maculans,*

heterochromatin domains (either H3K9me3 or H3K27me3) display significant enrichment in

putative effector genes, and that all known avirulence genes are associated with H3K9me3

during axenic growth [38]. We found here that the genes included in the eight clusters

specifically overexpressed during interaction with the plant relative to axenic growth were

also enriched in H3K9me3 and/or H3K27me3 histone modifications *in vitro*. One striking

finding of our study is that genes from the various waves of expression are scattered

throughout the genome, but are significantly associated with heterochromatin, facilitating the

temporal regulation of sets of genes involved in the same process. Seven of the eight

expression clusters identified in the lifecycle of *L. maculans* contained genes displaying

significant enrichment in H3K27me3 heterochromatin *in vitro*. This histone modification is an

important regulator of development and response to several biotic and abiotic stresses, in

various fungal or plant models (e.g. [69-70]). By contrast, genes associated with biotrophic or

biotrophic/asymptomatic life within the plants (cluster 2, **Fig 8**) displayed enrichment in

H3K9me3 *in vitro*. The genes in this cluster are located in TE-rich regions of the genome and

were found to be more enriched in heterochromatin modifications than the genes of the other

clusters. They included the largest number of candidate effector genes, and the largest number

of highly coregulated genes (65 genes). In *L. maculans*, the TE-rich regions are comprised of

heterochromatin, leading to the silencing of associated genes; loss of the repressive histone

modification H3K9me3 is a prerequisite to induce the expression of these genes [37]. The

unique association between genes involved in the symptomless spread of the fungus and

H3K9me3 suggests that this histone modification is involved in a regulatory mechanism

important for the concealment of the fungus from the plants defenses. Conversely, this cluster

also contains all the known avirulence genes, providing additional support for the role of

H3K9me3 as the primary regulator of expression for genes at the forefront of the battle with

the plant, and as a target of choice for the establishment of effector-triggered immunity during

co-evolution of the plant and the pathogen.

The genes associated with H3K9me3 in cluster 2 appear to be more strictly coregulated

than those in other clusters, in which genes are associated with H3K27me3. Our findings

suggest that chromatin remodeling during interaction with the plant is highly dynamic, rather

than occurring only once when the fungus infects the plant, as might have been anticipated

from analyses of species with a short lifecycle, or from the few days of interaction amenable

to laboratory experiments. This combined analysis of transcriptomic and epigenomic data thus

supports the notion that the finely tuned temporal progress of fungal infections is underpinned

by a highly dynamic phenomenon based on the successive opening and closing of different

genomic regions.

## 759　Materials and methods

### 760　Plant and fungal materials

Isolate JN2 (v23.1.2) was used in all experiments. Unless otherwise stated, the fungus

was cultured on V8-agar medium, and conidia for inoculation tests were produced as

previously described [71]. Isolate Nz-T4, which is sexually compatible with JN2, was also

used in *in vitro* crossing experiments. Both isolates were described by Balesdent *et al*. [72].

Three winter *B. napus* cultivars were used, depending on the experiment: 'Darmor-*bzh*',

the reference sequenced cultivar [73], 'Darmor' and 'Bristol'. 'Darmor-*bzh*' is a dwarf

isogenic line resulting from the introduction of the dwarf *bzh* gene into 'Darmor' [74].

'Darmor-*bzh*' and 'Darmor' are characterized by a high level of quantitative resistance in the

field, and 'Bristol' is moderately susceptible [75].

**Production of biological material for RNA sequencing**

*In vitro* **culture.** The fungus was grown under different *in vitro* conditions, to mimic as many

physiological stages as possible. All replicates used for RNA sequencing were true

independent biological replicates. The following physiological conditions were reproduced:

mycelial growth on V8-agar medium or in liquid Fries medium, pycnidiospore production on

V8-agar medium, pycnidiospore germination in liquid Fries medium, and sexual mating *in*

*vitro* at three time points (7, 20 and 35 days). All samples were immediately frozen in liquid

nitrogen after recovery and stored at -80°C until RNA extraction. Detailed protocols for the

production of samples in each set of conditions are provided in the S6 Text.

**Cotyledon inoculation and sampling.** The cotyledons of 10-day-old plants of cv Darmor-*bzh*

were inoculated with pycnidiospore suspensions, as previously described ([76]; S6 Text).

Control plants were mock-inoculated with sterile deionized water. Samples for RNA-Seq

were recovered at 0 (with or without wound), 2, 5, 7, 9, 12, 15 and 17 DPI. At each time

point, eight cotyledons from eight different plants were randomly selected. The plant tissues

around the inoculation site were cut with a 10 mm-diameter disposable hole-punch and the 16

corresponding samples were pooled together in a Falcon tube, immediately frozen in liquid

nitrogen and stored at -80°C until extraction. Two replicates were recovered at each time

point and the whole experiment was repeated once.

**Petiole inoculation and sampling.** The petioles of cvs Darmor-*bzh* and Bristol were

inoculated as described by Dutreux *et al*. [49]. Plants were grown in individual pots and the

petioles of the second and third true leaves were inoculated with 8 μL of $10^7$ spores.mL$^{-1}$

pycnidiospore suspensions. Petioles were sampled at 0 (mock), 7 and 14 DPI. At each date,

the two inoculated petioles of four plants were cut at their base and placed in Petri dishes.

Each petiole was cut 1 cm away from the point of inoculation, and the eight upper fragments

obtained were pooled in a 15 mL Falcon tube. The contents of this tube constituted one

sample of the upper part of the petiole. All samples were immediately frozen in liquid

nitrogen and stored at -80°C until extraction. Two independent replicates were recovered at

each time point and the whole experiment was repeated.

**Petiole inoculation and stem sampling.** Plants from the three varieties were sown in

individual pots and grown for three weeks before inoculation (S6 Text). When the plants

reached the three-leaf stage, the petiole of the second leaf was cut 0.5 cm from the stem. We

applied 10 μL of pycnidiospores ($10^7$ spores mL$^{-1}$) or sterile water (for mock inoculation) to

the petiole section. The plants were kept in the dark, under high humidity, for 36 h, and were

then transferred to a growth chamber (S6 Text). Three biological replicates were performed.

Each replicate consisted of three blocks of five plants per variety for RNA-Seq sampling. At

14 and 28 DPI, a 0.5 cm-long section of stem was excised from each plant with a razor blade.

The stem section was harvested at the level at which the inoculated petiole inserted into the

stem, at 14 DPI, and 0.5 cm below this level at 28 DPI. The tissues harvested from each block

at each time point (five individual plants) were pooled as a single sample. Harvested tissues

were immediately frozen in liquid nitrogen and stored at −80°C.

**Crop residue sampling.** In 2012-2013, the susceptible cultivar Alpaga was sown at Grignon,

France, where it was subject to natural infection. Disease incidence, assessed in the fall,

revealed high levels of infection, and pre-harvest assessments revealed high levels of disease

severity, variable between plants (S6 Text). After harvest, the stem bases of the plants were

removed from the field and kept outside until use, either to re-inforce the natural inoculum for

the 2013-2014 field assay (see below), or for RNA extraction. For RNA extraction, only

plants presenting necrosis on more than 50% of the stem area were used, and stem residues

were collected at seven time points after harvest. For one sampling date, two stem residues were pooled to constitute a single sample, and four samples (four replicates) were used per date for RNA sequencing.

**Stem bases from field sampling.** A field experiment was set up at Grignon, France in the fall of 2013, with two cultivars, Darmor-*bzh* and Bristol, with natural, but reinforced inoculum (S6 Text). Stem base tissues (1 cm above the collar of the plants) were collected from both cultivars in the field, at six time points, for RNA extraction (11/20/2013; 12/18/2013; 02/13/2014; 03/13/2014; 04/04/2014; 05/14/2014, 07/08/2014, i.e. one week before harvest (S6 Text)). For each cultivar and time point, we collected three samples (each containing three pooled stem bases) for RNA sequencing. Stem canker severity was assessed by G2 scoring, beginning at the April sampling (S6 Text). Only samples from the susceptible cultivar Bristol collected from March to July were kept for RNA-Seq statistical analyses.

**Young leaves from field sampling.** A field experiment was set up at Grignon, France in the fall of 2017. It included three plots of cv Darmor, subjected to natural, but reinforced inoculum (S6 Text). When typical leaf lesions due to *L. maculans* were visible on more than 60% of the plants (16/11/2017), leaves with leaf lesions were harvested. Five disks of leaf tissue with typical leaf lesions were pooled in a 15 mL Falcon tube (S6 Text). All samples were immediately frozen in liquid nitrogen and stored at -80°C until extraction. Pooled disks from one plot constitutes one replicate, and a similar sampling was performed two weeks later.

**Infection of cotyledons with ascospores.** Stem residues bearing mature pseudothecia were placed on a metallic grid over seven-day-old plants (S6 Text). The stem residues were left over the plants for 24 hours, in the dark, under saturation, to promote ascospore ejection. The plants were then transferred to a growth chamber. After 24 h and 48 h, six cotyledons per pot were randomly cut, pooled in a single 15 mL tube, frozen in liquid nitrogen and stored at -

842    80°C. Cotyledons from four selected neighboring pots (i.e., a total of 24 cotyledons; S6 Text)

843    were pooled to produce a single sample of the early stage of cotyledon infection with

844    ascospores. Two samples were obtained for each time point and the experiment was repeated

845    once.

**RNA extraction and sequencing**

Cotyledons, leaf disks, and freeze-dried cultures of fungal spores or mycelia were

ground with a Retsch MM300 mixer mill in Eppendorf tubes, with one tungsten carbide bead

per tube, for 45 s, at 30 oscillations per second. Petioles were ground with a pestle and mortar.

Stem bases from the field and stem residues were ground with a Retsch MM300 mixer mill,

with zirconium oxide jars and beads, for 40 s, at 30 oscillations per second. Inoculated stem

samples were ground in liquid nitrogen, in the presence of steel marbles, with a TissueLyser II

(Qiagen).

RNA was extracted with Trizol reagent (Invitrogen, Cergy Pontoise, France), as

previously described [18]. RNA-Seq libraries were prepared as previously described [49].

Each library was sequenced with 101 bp paired-end read chemistry, on a HiSeq2000 Illumina

sequencer.

**Annotation of genomic data**

**Comparisons of gene annotations between the two published version of the *L. maculans***

**strain JN3 genome.** The reference genome assembly of *L. maculans* isolate v23.1.3 (aka.

JN3) was generated in 2007, by Sanger sequencing [26]. Dutreux *et al.* [49] subsequently

generated an improved genome assembly, to achieve assembly at the chromosome scale. We

compared this new version of the genome sequence with that published in 2011, using BLAT

suite tools [77] (S4 Text).

**Functional annotations.** Functional annotations were predicted with BLAST2GO [78].

Genes encoding CAZymes were annotated manually, together with *de novo* prediction with

dbCAN tools v3 [79]. PFam domains were annotated with PFamScan software version 1.6

[80]. For discrimination between proteins with and without annotated functions, the

BLAST2GO results and PFam annotations were used, but with the terms "hypothetical

protein", "predicted protein" and "uncharacterized protein" discarded. PKS and NRPS genes

were predicted with the SMURF online tool, using default parameters [81].

**Annotation of genes encoding SSPs.** We generated a new repertoire of genes encoding SSPs,

by first creating a secretome of predicted secreted proteins with three filters. The predicted

secretome contained all the proteins with no more than one transmembrane domain predicted

by TMHMM version 2.0. [82], and either a signal peptide predicted by SignalP version 4.1

[83], or an extracellular localization predicted by TargetP version 1.1 [84]. The final SSP

repertoire was created by applying a size cutoff of 300 amino acids to the predicted

secretome. In parallel, EffectorP version 1.0 [85] was applied to the whole predicted

secretome and detected four proteins of more than 300 amino acids in length, which were

added to the SSP repertoire.

**Detection of AT-rich regions.** AT-rich and GC-equilibrated regions were annotated with

OcculterCut [86] and post-processed to avoid the detection of regions that were too small.

OcculterCut uses a predefined window of 1 kb to analyze AT/GC content, and this can lead to

the detection of small GC-rich-regions within AT-rich regions in the case of isolated genes.

We overcame this problem by defining reliable detected regions (10 kb for AT and 6 kb for

GC), which we used as anchors for the merging of regions (S6 Text).

**Analyses on genomic data**

**Genomic localization and enrichment in epigenetic marks.** For genes in the eight clusters, we performed Chi² tests to detect enrichment in the associated epigenetic marks (H3K4me2, H3K9me3 or H3K29me3), recovered from Soyer *et al*. [38], to compare the distribution of marks in the whole set of genes. Enrichments were considered significant for $p$ values $< 0.05$; all analyses were performed in R.

**Gene Ontology enrichment analysis.** The Bingo plugin of Cytoscape software v3.5.1 [87] was used for Gene Ontology (GO) enrichment analysis. For each gene, the GO identifier was generated with BLAST2GO software. Significant enrichments in "Molecular Function", "Biological Process" and "Cellular Component" were detected with a hypergeometric test, by comparing the proportion of genes identified for a given GO function in the subset of interest to that for the whole set of protein-coding genes in *L. maculans*.

**BLAST.** A BLAST alignment was generated with non-redundant protein sequences from the GenPept, Swissprot, PIR, PDF, PDB, and NCBI RefSeq databases. Only the 10 best BLAST hits were selected, with the max_target_seqs: 10 and max_hsps: 10 options.

**Analyses on transcriptomic data**

**Mapping.** Raw RNA-Seq reads were mapped with STAR software version 020201 [88]. As *Leptosphaeria biglobosa* isolates could be present in the field samples, we assessed the mapping specificity of the final parameters for the genome of *L. maculans* (S1 Text, **S1 Table**). We chose a number of two base-pair mismatches for a one-step process of mapping onto the concatenated genome of the two species for the final analysis. Based on the maximum intron size within the genome, we allowed for an intron size of 10 000 bp. The other parameters used were as follows: outFilterMultimapNmax: 100; SeedSearchStartLmax: 12; alignSJoverhangMin: 15; alignIntronMin: 10. We then selected the correctly paired reads

in BAM files with Samtools v1.6 [89]. Finally, FeaturesCounts version v1.5.1 [90] was used

to quantify gene expression for the uniquely mapped and paired reads.

**Sample correlation analysis.** Pairwise Pearson's correlation analysis was performed on all

samples, with the Log2(FPKM +1)-transformed expression values for the 32 selected sets of

conditions. The resulting correlation matrix was clustered with the Ward.D2 method. PCA

was performed with the FactoRmineR R package [91] on the scaled Log2(FPKM +1)

expression values for the four groups identified in the correlation matrix.

**Detection of differentially expressed genes.** Differential expression analyses were

performed with the EdgeR package v3.20.9 [92]. Genes with > 30 reads in at least one

condition were kept for statistical analysis. Samples were normalized by the TMM method.

We fitted a negative binomial generalized linear model to the data with the glmFIt function.

We then used the glmLRT function to compare gene expression in each of the 22 *in planta*

conditions with that in the 10 *in vitro* conditions, using appropriate coefficients and contrasts

(-1/10 for the 10 *in vitro* conditions and +1 for the tested conditions). We also used

coefficients to mask the effects of replicate on cotyledon, petiole and stem samples from

controlled conditions. Only genes overexpressed relative to the10 *in vitro* conditions with

LogFC > 4 and a *p*-value < 0.01 were selected.

**Clustering** For the clustering step, genes with a FPKM > 2 in at least one condition were

Log2(FPKM +1)-transformed and scaled. The whole 1207 gene set overexpressed in at least

one set of conditions *in planta* was extracted from this transformed count table and the self-

organizing map method of the Kohonen R package v3.0.8 [93] was used to classify those

genes into eight clusters, with 200 iterations.

**Detection of co-expressed genes.** For each cluster, the mean scaled Log2(FPKM +1) value

for SSP gene expression was calculated and used as a reference for an analysis of linear

regression against the scaled Log2(FPKM +1) expression of all genes. Genes with an R-

squared value > 0.80 and an adjusted $p$-value < 0.05 were considered to be highly correlated.

## Acknowledgments

We would like to thank Christophe Montagnier and the experimental unit team, INRAE,

France, for the setting up and management of the experimental rapeseed fields; Guillem

Rigaill from the Institute of Plant Sciences Paris-Saclay, France, for helpful comments and

advice on statistics, and Antoine Gravot, (INRAE IGEPP, France), for helpful discussions and

review of the manuscript. For preliminary analyses and genomic work, we would like to thank

Fabien Dutreux (CEA, Genoscope, Evry, France). This work was supported in part by

Promosol (METAPHOR project), and France Génomique (Leptolife project). The "Effectors

and Pathogenesis of *L. maculans*" group benefits from the support of Saclay Plant Sciences-

SPS (ANR-17-EUR-0007).

## Author contribution

Conceptualization, RD, TR, MHB; Data curation, EJG, NL, CDA, JMA, CC; Formal

analysis EJG, JLS, NL, CDA, JMA; Funding acquisition, PW, TR, MHB; investigation, EJG,

NL, JL, CC, AL, JL, MHB; methodology, EJG, JL; Project administration, TR; resources,

EJG, NL, CDA, PW, JMA, CC, MHB; supervision, PW, JMA, TR, MHB; validation, JLS, IF;

TR; visualization; EJG, JLS; writing draft, EJG, JLS, NL, TR, MHB; writing review, EJG,

JLS, NL, IF, CDA, PW, JMA, CC, RD, TR, MHB.

**References**

1. Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, et al.
Emerging fungal threats to animal, plant and ecosystem health. Nature. 2012;484(7393):
186–194.

2. Dobon A, Bunting DCE, Cabrera-Quio LE, Uauy C, Saunders DGO. The host-pathogen
interaction between wheat and yellow rust induces temporally coordinated waves of gene
expression. BMC Genomics. 2016;17: 380.

3. Dong Y, Li Y, Zhao M, Jing M, Liu X, Liu M, et al. Global genome and transcriptome
analyses of *Magnaporthe oryzae* epidemic isolate 98-06 uncover novel effectors and
pathogenicity-related genes, revealing gene gain and lose dynamics in genome evolution.
PLoS Pathog. 2015;11(4): e1004801.

4. Haueisen J, Möller M, Eschenbrenner CJ, Grandaubert J, Seybold H, Adamiak H, et al.
Highly flexible infection programs in a specialized wheat pathogen. Ecol Evol. 2019;9:
275–294.

5. Palma-Guerrero J, Torriani SFF, Zala M, Carter D, Courbot M, Rudd JJ, et al.
Comparative transcriptomic analyses of *Zymoseptoria tritici* strains show complex
lifestyle transitions and intraspecific variability in transcription profiles. Mol Plant Pathol.
2016;17: 845–59.

6. Kleemann J, Rincon-Rivera LJ, Takahara H, Neumann U, van Themaat EVL, van der
Does HC, et al. Sequential delivery of host-induced virulence effectors by appressoria and
intracellular hyphae of the phytopathogen *Colletotrichum higginsianum*. PLoS Pathog.
2012;8(4): e1002643.

7. Lanver D, Müller AN, Happel P, Schweizer G, Haas FB, Franitza M, et al. The biotrophic
development of *Ustilago maydis* studied by RNA-seq Analysis. Plant Cell. 2018;30: 300–
323.

8. Lorrain C, Marchal C, Hacquard S, Delaruelle C, Pétrowski J, Petre B, et al. The rust
fungus *Melampsora larici-populina* expresses a conserved genetic program and distinct
sets of secreted protein genes during infection of its two host plants, larch and poplar. Mol
Plant Microbe Intercact. 2018;31: 695-706

9. Rouxel T, Balesdent MH. The stem canker (blackleg) fungus, *Leptosphaeria maculans*,
enters the genomic era. Mol Plant Pathol. 2005;6: 225–241.

10. West JS, Balesdent M-H, Rouxel T, Narcy JP, Huang Y-J, Roux J, et al. Colonization of
winter oilseed rape tissues by A/Tox+ and B/Tox0 *Leptosphaeria maculans* (phoma stem
canker) in France and England. Plant Pathol. 2002;51: 311–321.

11. Haddadi P, Ma L, Wang H, Borhan MH. Genome-wide transcriptomic analyses provide
insights into the lifestyle transition and effector repertoire of *Leptosphaeria maculans*
during the colonization of *Brassica napus* seedlings. Mol Plant Pathol. 2016;17: 1196–
210.

12. Lowe RGT, Cassin A, Grandaubert J, Clark BL, Van de Wouw AP, Rouxel T, et al.
Genomes and transcriptomes of partners in plant-fungal-interactions between canola
(*Brassica napus*) and two *Leptosphaeria* species. PLoS ONE. 2014;9(7): e103098.

13. Sonah H, Zhang X, Deshmukh RK, Borhan MH, Fernando WGD, Bélanger RR.
Comparative transcriptomic analysis of virulence factors in *Leptosphaeria maculans*
during compatible and incompatible interactions with canola. Front Plant Sci. 2016;7:
1784.

14. Gervais J, Plissonneau C, Linglin J, Meyer M, Labadie K, Cruaud C, et al. Different
waves of effector genes with contrasted genomic location are expressed by *Leptosphaeria*
*maculans* during cotyledon and stem colonization of oilseed rape. Mol Plant Pathol.
2017;18: 1113–1126.

15. Lo Presti LL, Lanver D, Schweizer G, Tanaka S, Liang L, Tollot M, et al. Fungal
effectors and plant susceptibility. Annu Rev Plant Biol. 2015;66: 513–145.

16. Elliott CE, Gardiner DM, Thomas G, Cozijnsen A, Van de Wouw AP, Howlett BJ.
Production of the toxin sirodesmin PL by *Leptosphaeria maculans* during infection of
*Brassica napus*. Mol Plant Pathol. 2007;8: 791–802.

17. Balesdent M-H, Fudal I, Ollivier B, Bally P, Grandaubert J, Eber F, et al. The dispensable
chromosome of *Leptosphaeria maculans* shelters an effector gene conferring avirulence
towards *Brassica rapa*. New Phytol. 2013;198: 887–898.

18. Fudal I, Ross S, Gout L, Blaise F, Kuhn ML, Eckert MR, et al. Heterochromatin-like
regions as ecological niches for avirulence genes in the *Leptosphaeria maculans* genome:
Map-Based Cloning of *AvrLm6*. Mol Plant Microbe Interact. 2007;20: 459–470.

19. Ghanbarnia K, Ma L, Larkan NJ, Haddadi P, Fernando WGD, Borhan MH.
*Leptosphaeria maculans* AvrLm9: a new player in the game of hide and seek with
AvrLm4-7. Mol Plant Pathol. 2018;19: 1754–64.

20. Ghanbarnia K, Fudal I, Larkan NJ, Links MG, Balesdent M-H, Profotova B, et al. Rapid identification of the *Leptosphaeria maculans* avirulence gene *AvrLm2* using an intraspecific comparative genomics approach. Mol Plant Pathol. 2015;16: 699–709.

21. Gout L, Fudal I, Kuhn M-L, Blaise F, Eckert M, Cattolico L, et al. Lost in the middle of nowhere: the *AvrLm1* avirulence gene of the Dothideomycete *Leptosphaeria maculans*. Mol Microbiol. 2006;60: 67–80.

22. Parlange F, Daverdin G, Fudal I, Kuhn M-L, Balesdent M-H, Blaise F, et al. *Leptosphaeria maculans* avirulence gene *AvrLm4-7* confers a dual recognition specificity by the *Rlm4* and *Rlm7* resistance genes of oilseed rape, and circumvents *Rlm4*-mediated recognition through a single amino acid change. Mol Microbiol. 2009;71: 851–863.

23. Petit☐Houdenot Y, Degrave A, Meyer M, Blaise F, Ollivier B, Marais C-L, et al. A two genes-for-one gene interaction between *Leptosphaeria maculans* and *Brassica napus*. New Phytol. 2019;223: 397–411.

24. Plissonneau C, Daverdin G, Ollivier B, Blaise F, Degrave A, Fudal I, et al. A game of hide and seek between avirulence genes *AvrLm4-7* and *AvrLm3* in *Leptosphaeria maculans*. New Phytol. 2016;209: 1613–1624.

25. Van de Wouw AP, Lowe RGT, Elliott CE, Dubois DJ, Howlett BJ. An avirulence gene, *AvrLmJ1*, from the blackleg fungus, *Leptosphaeria maculans*, confers avirulence to *Brassica juncea* cultivars. Mol Plant Pathol. 2014;15: 523–530.

26. Rouxel T, Grandaubert J, Hane JK, Hoede C, van de Wouw AP, Couloux A, et al. Effector diversification within compartments of the *Leptosphaeria maculans* genome affected by Repeat-Induced Point mutations. Nature Commun. 2011;2: 202.

27. Darma R, Lutz A, Elliott CE, Idnurm A. Identification of a gene cluster for the synthesis of the plant hormone abscisic acid in the plant pathogen *Leptosphaeria maculans*. Fungal Genet Biol. 2019;130: 62–71.

28. Elliott CE, Howlett BJ. Overexpression of a 3-Ketoacyl-CoA thiolase in *Leptosphaeria maculans* causes reduced pathogenicity on *Brassica napus*. Mol Plant Microbe Interact. 2006;19: 588–596.

29. Elliott CE, Harjono, Howlett BJ. Mutation of a gene in the fungus *Leptosphaeria maculans* allows increased frequency of penetration of stomatal apertures of arabidopsis thaliana. Mol Plant. 2008;1: 471–481.

30. Feng J, Zhang H, Strelkov SE, Hwang S-F. The *LmSNF1* Gene is required for pathogenicity in the canola blackleg pathogen *Leptosphaeria maculans*. PLoS ONE. 2014;9: e92503.

31. Idnurm A, Howlett BJ. Analysis of loss of pathogenicity mutants reveals that repeat-induced point mutations can occur in the Dothideomycete *Leptosphaeria maculans*. Fungal Genet and Biol. 2003;39: 31–37.

32. Remy E, Meyer M, Blaise F, Chabirand M, Wolff N, Balesdent M-H, et al. The *Lmpma1* gene of *Leptosphaeria maculans* encodes a plasma membrane H+-ATPase isoform essential for pathogenicity towards oilseed rape. Fungal Genet Biol. 2008;45: 1122–1134.

33. Remy E, Meyer M, Blaise F, Simon UK, Kuhn D, Chabirand M, et al. The *Lmgpi15* gene, encoding a component of the glycosylphosphatidylinositol anchor biosynthesis pathway, is required for morphogenesis and pathogenicity in *Leptosphaeria maculans*. New Phytol. 2008;179: 1105–1120.

34. Remy E, Meyer M, Blaise F, Simon UK, Kuhn D, Balesdent MH, et al. A key enzyme of the Leloir pathway is involved in pathogenicity of *Leptosphaeria maculans* toward oilseed rape. Mol Plant Microbe Interact. 2009;22: 725–736.

35. Soyer JL, Hamiot A, Ollivier B, Balesdent M-H, Rouxel T, Fudal I. The APSES transcription factor LmStuA is required for sporulation, pathogenic development and effector gene expression in *Leptosphaeria maculans*. Mol Plant Pathol. 2015;16: 1000–1005.

36. Van de Wouw AP, Pettolino FA, Howlett BJ, Elliott CE. Mutations to LmIFRD affect cell wall integrity, development and pathogenicity of the ascomycete *Leptosphaeria maculans*. Fungal Genet Biol. 2009;46:695–706.

37. Soyer JL, Ghalid ME, Glaser N, Ollivier B, Linglin J, Grandaubert J, et al. Epigenetic control of effector gene expression in the plant pathogenic fungus *Leptosphaeria maculans*. PLoS Genetics. 2014;10: e1004227.

38. Soyer JL, Clairet C, Gay EJ, Lapalu N, Rouxel T, Stukenbrock EH, et al. Genome-wide mapping of histone modifications in two species of *Leptosphaeria maculans* showing contrasting genomic organization and host specialization. bioRxiv. 2020 May 10;2020.05.08.084566.

39. Murphy E, De Smet I. Understanding the RALF family: a tale of many species. Trends Plant Sci. 2014;19:664–671.

40. Rabe F, Ajami□Rashidi Z, Doehlemann G, Kahmann R, Djamei A. Degradation of the plant defence hormone salicylic acid by the biotrophic fungus *Ustilago maydis*. Mol Microbiol. 2013;89: 179–188.

41. De Jonge R, Thomma BPHJ. Fungal LysM effectors: extinguishers of host immunity? Trends Microbiol. 2009;17:151–157.

42. Gijzen M, Nürnberger T. Nep1-like proteins from plant pathogens: recruitment and
diversification of the NPP1 domain across taxa. Phytochemistry. 2006;67 :1800–1807.

43. Elliott CE, Callahan DL, Schwenk D, Nett M, Hoffmeister D, Howlett BJ. A gene cluster
responsible for biosynthesis of phomenoic acid in the plant pathogenic fungus,
*Leptosphaeria maculans*. Fungal Genet and Biol. 2013;53: 50–58.

44. Gardiner DM, Jarvis RS, Howlett BJ. The ABC transporter gene in the sirodesmin
biosynthetic gene cluster of *Leptosphaeria maculans* is not essential for sirodesmin
production but facilitates self-protection. Fungal Genet Biol. 2005;42: 257–263.

45. Urquhart AS, Idnurm A. Limitations of transcriptome-based prediction of pathogenicity
genes in the plant pathogen *Leptosphaeria maculans*. FEMS Microbiol Lett. 2019;366:
fnz080.

46. Idnurm A, Howlett BJ. Isocitrate lyase is essential for pathogenicity of the fungus
*Leptosphaeria maculans* to canola (*Brassica napus*). Eukaryot Cell. 2002 10;1: 719–724.

47. Ma L, Djavaheri M, Wang H, Larkan NJ, Haddadi P, Beynon E, et al. *Leptosphaeria*
*maculans* effector protein avrlm1 modulates plant immunity by enhancing MAP kinase 9
phosphorylation. iScience. 2018;3: 177–191.

48. Nováková M, Šašek V, Trdá L, Krutinová H, Mongin T, Valentová O, et al.
*Leptosphaeria maculans* effector AvrLm4-7 affects salicylic acid (SA) and ethylene (ET)
signalling and hydrogen peroxide ($H_2O_2$) accumulation in *Brassica napus*. Mol Plant
Pathol. 2016;17: 818–831.

49. Dutreux F, Da Silva C, d'Agata L, Couloux A, Gay EJ, Istace B, et al. De novo assembly
and annotation of three *Leptosphaeria genomes* using Oxford Nanopore MinION
sequencing. Scientific Data. 2018;5: 180235.

50. West JS, Kharbanda PD, Barbetti MJ, Fitt BDL. Epidemiology and management of
*Leptosphaeria maculans* (phoma stem canker) on oilseed rape in Australia, Canada and
Europe. Plant Pathol. 2001;50: 10–27.

51. Valent B, Khang CH. Recent advances in rice blast effector research. Curr Opin Plant
Biol. 2010;13: 434–441.

52. Hammond KE, Lewis BG. Variation in stem infections caused by aggressive and non-
aggressive isolates of *Leptosphaeria maculans* on *Brassica napus* var. *oleifera*. Plant
Pathol. 1987;36: 53–65.

53. Hammond KE, Lewis BG. The establishment of systemic infection in leaves of oilseed
rape by *Leptosphaeria maculans*. Plant Pathol. 1987;36: 135–147.

54. Johnson RD, Lewis BG. Variation in host range, systemic infection and epidemiology of
*Leptosphaeria maculans*. Plant Pathol. 1994;43: 269–277.

55. Kerdraon L, Balesdent M-H, Barret M, Laval V, Suffert F. Crop residues in wheat-oilseed
rape rotation system: a pivotal, shifting platform for microbial meetings. Microb Ecol.
2019;77: 931–945.

56. Chanclud E, Morel J-B. Plant hormones: a fungal point of view. Mol Plant Pathol.
2016;17: 1289–1297.

57. Qi G, Chen J, Chang M, Chen H, Hall K, Korin J, et al. Pandemonium breaks out:
disruption of salicylic acid-mediated defense by plant pathogens. Mol Plant. 2018;11:
1427–1439.

58. Ohm RA, Riley R, Salamov A, Min B, Choi I-G, Grigoriev IV. Genomics of wood-
degrading fungi. Fungal Genet Biol. 2014;72: 82–90.

59. Tyler BM, Rouxel T. Effectors of fungi and oomycetes: their virulence and avirulence
functions and translocation from pathogen to host cells. ; In Molecular Plant Immunity.
2013; G. Sessa, Ed., Wiley Blackwell Press, Chap. 7, pp. 123-167.

60. Cheeseman K, Weitzman JB. Host–parasite interactions: an intimate epigenetic
relationship. Cell Microbiol. 2015;17: 1121–1132.

61. Gómez-Díaz E, Jordà M, Peinado MA, Rivero A. Epigenetics of host-pathogen
interactions: the road ahead and the road behind. PLoS Pathog. 2012 Nov 29 [cited 2020
Apr 27];8(11). Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3510240/

62. Soyer JL, Rouxel T, Fudal I. Chromatin-based control of effector gene expression in
plant-associated fungi. Curr Opin Plant Biol. 2015;26: 51–56.

63. Fouché S, Badet T, Oggenfuss U, Plissonneau C, Francisco CS, Croll D. Stress-driven
transposable element de-repression dynamics and virulence evolution in a fungal
pathogen. Mol Biol Evol. 2020;37: 221–239.

64. Chujo T, Scott B. Histone H3K9 and H3K27 methylation regulates fungal alkaloid
biosynthesis in a fungal endophyte-plant symbiosis. Mol Microbiol. 2014;92: 413–434.

65. Connolly LR, Smith KM, Freitag M. The *Fusarium graminearum* histone H3 K27
methyltransferase KMT6 regulates development and expression of secondary metabolite
gene clusters. PLoS Genet [Internet]. 2013 Oct 31 [cited 2020 Apr 27];9(10). Available
from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3814326/

66. Gacek A, Strauss J. The chromatin code of fungal secondary metabolite gene clusters.
Appl Microbiol Biotechnol. 2012;95:1389–1404.

67. Soyer JL, Grandaubert J, Haueisen J, Schotanus K, Stukenbrock EH. *In planta* chromatin
immunoprecipitation in *Zymoseptoria tritici* reveals chromatin-based regulation of
putative effector gene expression. bioRxiv. 2019 Feb 8;544627.

68. Studt L, Rösler SM, Burkhardt I, Arndt B, Freitag M, Humpf H-U, et al. Knock-down of
the methyltransferase Kmt6 relieves H3K27me3 and results in induction of cryptic and
otherwise silent secondary metabolite gene clusters in *Fusarium fujikuroi*. Environ
Microbiol. 2016;18: 4037–4054.

69. Basenko EY, Sasaki T, Ji L, Prybol CJ, Burckhardt RM, Schmitz RJ, et al. Genome-wide
redistribution of H3K27me3 is linked to genotoxic stress and defective growth. PNAS.
2015;112: E6339–6348.

70. Wang G, Köhler C. Epigenetic processes in flowering plant reproduction. J Exp Bot.
2017;68: 797–807.

71. Ansan-Melayah D, Balesdent MH, Buée M, Rouxel T. Genetic characterization of
*AvrLm1*, the first avirulence gene of *Leptosphaeria maculans*. Phytopathology. 1995;85:
1525–1529.

72. Balesdent MH, Attard A, Kühn ML, Rouxel T. New avirulence genes in the
phytopathogenic fungus *Leptosphaeria maculans*. Phytopathology. 2002;92:1122–1133.

73. Chalhoub B, Denoeud F, Liu S, Parkin IAP, Tang H, Wang X, et al. Early allopolyploid
evolution in the post-neolithic *Brassica napus* oilseed genome. Science. 2014;345(6199):
950–953.

74. Foisset N, Delourme R, Barret P, Renard M. Molecular tagging of the dwarf BREIZH
(Bzh) gene in *Brassica napus*. Theor Appl Genet. 1995;91: 756–761.

75. Pilet ML, Delourme R, Foisset N, Renard M. Identification of loci contributing to
quantitative field resistance to blackleg disease, causal agent *Leptosphaeria maculans*
(Desm.) Ces. et de Not., in Winter rapeseed (*Brassica napus*). Theor Appl Genet.
1998;96: 23–30.

76. Gall C, Balesdent M-H, Desthieux I, Robin P, Rouxel T. Polymorphism of Tox$^0$
*Leptosphaeria maculans* isolates as revealed by soluble protein and isozyme
electrophoresis. Mycol Res. 1995;99: 221–229.

77. Kent WJ. BLAT-The BLAST-Like Alignment Tool. Genome Res. 2002;12: 656–664.

78. Conesa A, Götz S. Blast2GO: A comprehensive suite for functional analysis in plant
genomics. Int J Plant Genomics [Internet]. 2008 [cited 2019 Nov 12];2008. Available
from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2375974/

79. Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. dbCAN: a web resource for automated
carbohydrate-active enzyme annotation. Nucleic Acids Res. 2012;40(Web Server
issue):W445–451.

80. Mistry J, Bateman A, Finn RD. Predicting active site residue annotations in the Pfam
database. BMC Bioinformatics. 2007 Aug 9;8(1): 298.

81. Khaldi N, Seifuddin FT, Turner G, Haft D, Nierman WC, Wolfe KH, et al. SMURF:
genomic mapping of fungal secondary metabolite clusters. Fungal Genet Biol.
2010;47:736–741.

82. Möller S, Croning MDR, Apweiler R. Evaluation of methods for the prediction of
membrane spanning regions. Bioinformatics. 2001;17: 646–653.

83. Nielsen H. Predicting Secretory Proteins with SignalP. Methods Mol Biol. 2017;1611:
59-73. doi: 10.1007/978-1-4939-7015-5_6.

84. Almagro Armenteros JJ, Salvatore M, Emanuelsson O, Winther O, von Heijne G,
Elofsson A, et al. Detecting sequence signals in targeting peptides using deep learning.
Life Sci Alliance. 2019;2(5): 01900429.

85. Sperschneider J, Gardiner DM, Dodds PN, Tini F, Covarelli L, Singh KB, et al.
EffectorP: predicting fungal effector proteins from secretomes using machine learning.
New Phytol. 2016;210: 743–761.

86. Testa AC, Oliver RP, Hane JK. OcculterCut: A comprehensive survey of AT-rich regions
in fungal genomes. Genome Biol Evol. 2016;8: 2044–2064.

87. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: A
software environment for integrated models of biomolecular interaction networks.
Genome Res. 2003;13: 2498–504.

88. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29: 15–21.

89. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence
alignment/map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.

90. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014 Apr 1;30(7):923–30.

91. Lê S, Josse J, Husson F. FactoMineR□: An *R* package for multivariate analysis. J Stat
Soft [Internet]. 2008 [cited 2019 Nov 12];25(1). Available from:
http://www.jstatsoft.org/v25/i01/

92. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. 2010;26: 139–140.

93. Wehrens R, Buydens LMC. Self- and super-organizing maps in R: The kohonen Package.
J Stat Soft. 2007;21:1–19.

94. Gardiner DM, Howlett BJ. Negative selection using thymidine kinase increases the
efficiency of recovery of transformants with targeted genes in the filamentous fungus
*Leptosphaeria maculans*. Curr Genet. 2004;45: 249–255.

95. Sexton AC, Paulsen M, Woestemeyer J, Howlett BJ. Cloning, characterization and
chromosomal location of three genes encoding host-cell-wall-degrading enzymes in
*Leptosphaeria maculans*, a fungal pathogen of *Brassica* spp. Gene. 2000;248: 89–97.

96. Taylor JL, Condie J. Characterization of ABC transporters from the fungal
phytopathogen *Leptosphaeria maculans*. In Biology of Plant-Microbe Interactions (P. J.
G. M. de Wit, T. Bisseling and W. J.Stiekema, Eds.), 2000 Vol. 2, pp. 71–76. Int. Soc.
Plant–Microbe Interact, St. Paul, MN.

97. Idnurm A, Warnecke DC, Heinz E, Howlett BJ. Characterisation of neutral trehalase and
UDP-glucose:sterol glucosyltransferase genes from the plant pathogenic fungus
*Leptosphaeria maculans*. Physiol Mol Plant Pathol. 2003 ;62: 305–313.

98. Idnurm A, Taylor JL, Pedras MSC, Howlett BJ. Small scale functional genomics of the
blackleg fungus, *Leptosphaeria maculans*: analysis of a 38 kb region. Australas Plant
Pathol. 2003;32: 511–519.

99. Sexton AC, Howlett BJ. Characterisation of a cyanide hydratase gene in the
phytopathogenic fungus *Leptosphaeria maculans*. Mol Gen Genet. 2000;263: 463–470.

100. Urquhart AS, Idnurm A. Sit4-associated protein is required for pathogenicity of
*Leptosphaeria maculans* on *Brassica napus*. Curr Microbiol. 2017;74: 1438–1446.

101. Wilson LM, Idnurm A, Howlett BJ. Characterization of a gene (sp1) encoding a secreted
protein from *Leptosphaeria maculans*, the blackleg pathogen of *Brassica napus*. Mol
Plant Pathol. 2002;3: 487–493.

102. Vincent D, Balesdent M-H, Gibon J, Claverol S, Lapaillerie D, Lomenech A-M, et al.
Hunting down fungal secretomes using liquid-phase IEF prior to high resolution 2-DE.
Electrophoresis. 2009;30: 4118–4136.

103. Balesdent MH, Attard A, Ansan-Melayah D, Delourme R, Renard M, Rouxel T. Genetic
control and host range of avirulence toward *Brassica napus* cultivars Quinta and Jet Neuf
in *Leptosphaeria maculans*. Phytopathology. 2001;91:70–76.