In Situ Optical Quantification of Extracellular Electron Transfer from *Shewanella oneidensis* using Plasmonic Metal Oxide Nanocrystals

Austin J. Graham*[1,2], Stephen L. Gibbs*[1], Camila A. Saez Cabezas*[1,2], Yongdan Wang[1], Allison M. Green[1,2], Delia J. Milliron^[1,2], Benjamin K. Keitz^[1,2]

[1]McKetta Department of Chemical Engineering, University of Texas at Austin, Austin, TX 78712

[2]Center for Dynamics and Control of Materials, University of Texas at Austin, Austin, TX 78712

*These authors contributed equally.

^To whom correspondence should be addressed: milliron@che.utexas.edu; keitz@utexas.edu

Keywords: extracellular electron transfer, biosensors, colloidal semiconductor nanocrystals, doped metal oxides, synthetic biology, localized surface plasmon resonance

Abstract

Extracellular electron transfer (EET) is a critical form of microbial metabolism that enables respiration on a variety of inorganic substrates, including metal oxides. For this reason, engineering EET processes has garnered significant interest for applications ranging from bioelectronics to materials synthesis. These applications require a strong understanding of electron flux from EET-relevant microbes. However, quantifying current generated by electroactive bacteria has been predominately limited to biofilms formed on electrodes, which require long incubation times, electrode colonization, and convolute contributions to EET from planktonic cells. To address this, we developed a platform for quantifying time-resolved EET flux from cell suspensions using aqueous dispersions of plasmonic tin-doped indium oxide nanocrystals. Tracking the change in optical extinction during electron transfer and fitting the optical response to a free electron model enabled quantification of current generation and electron transfer rate constants from planktonic *Shewanella oneidensis* cultures. Using this method, we differentiated between starved and actively respiring *S. oneidensis*, and between cells of varying genotype using an EET knockout strain. In addition, we quantified current production ranging from 0.12 – 0.68 fA • cell^{-1} from *S. oneidensis* cells engineered to differentially express a key EET gene using an inducible genetic circuit. Overall, our results validate the utility of colloidally stable plasmonic metal oxide nanocrystals as quantitative biosensors in native biological environments and contribute to a fundamental understanding of planktonic *S. oneidensis* electrophysiology using simple *in situ* spectroscopy.

Introduction

Extracellular electron transfer (EET) is a microbial respiratory process by which electron flux from carbon metabolism is transported across the cell membrane and directed onto inorganic substrates (1). This capability makes electrogenic bacteria useful organisms for microbial fuel

cells (2, 3), bioelectronics (4–6), and material syntheses (7, 8), and has led to significant interest

in controlling their electron flux using synthetic biology (9, 10). Improving EET-dependent

technologies such as controlling electron transfer dynamics (11), implementing genetic logic (12),

and orchestrating electron transfer dynamics between proteins (13) necessitates a stronger

quantitative understanding of bacterial electrophysiology and simple methods to measure current

generation. EET is usually quantified from biofilms formed on electrodes in microbial fuel cells or

bioelectrochemical systems (1, 14–16). EET from single cells has also been measured in such

systems, providing sensitive quantitative measurements and highlighting variability in EET

capabilities between individual cells (17–19). However, these measurements often require long

incubation times and electrode colonization (~days), and the contribution of suspended (*i.e.*,

planktonic) microbes is generally unclear and not controlled in electrode-based platforms (20).

EET from planktonic microbes is typically quantified using soluble substrates such as ferric citrate

(12, 21, 22), but such molecules may be passively or actively transported across the cell

membrane (23), convoluting reduction pathways. In addition, reduction of soluble substrates like

ferric citrate does not necessarily translate to electricity generation (24). Furthermore,

implementation of synthetic gene programs such as genetic logic operations often require cell

turnover and growth, which are convoluted in biofilms. Overall, there is a need for reliable *in situ*

biosensors of EET that are both compatible with planktonic culture and more accurately capture

electron transfer dynamics to insoluble substrates.

Plasmonic nanoparticles address this need because they are insoluble but maintain colloidal

stability in a variety of solvents. An intrinsically high concentration of conductive electrons results

in a strong absorption at the localized surface plasmon resonance (LSPR) peak frequency, ω_{LSPR},

and enables participation in redox reactions with molecules near the surface many times over.

The LSPR peak position and intensity depend on the particle's conduction electron concentration,

n_e, such that tracking the optical response can report on the number of redox reactions occurring

at the nanoparticle surface (25–27). Thus, optically tracking the plasmonic response can monitor

electron transfer similar to tracking current from a traditional film electrode. Doped metal oxide

nanocrystals are particularly promising candidates because they sustain plasmonic resonance

but have n_e ~1-3 orders of magnitude lower than Au and Ag (28, 29), making their optical

response more sensitive to single electron transfer events (25, 26). Further, the surface of doped

metal oxide nanocrystals can be post-synthetically functionalized with various ligands to

customize solvent compatibility to ensure colloidal stability and tune the interactions with other

species in solution (30–32). Indeed, the advantageous properties of doped metal oxide

nanocrystals have already enabled their use in spectroscopic measurement of EET (33–36).

However, these methods did not directly quantify the relationship between the spectroscopic

response and the electron transfer rate (*i.e.*, current generation). Building on prior work, we

recently developed a model for fitting the plasmonic optical response of doped metal oxide

nanocrystal dispersions which extracts the number of free electrons in each nanocrystal, enabling

in situ tracking of biological electron transfer events using spectroscopy (37).

Leveraging our ability to quantify the change in optical response with the aqueous biocompatibility

of colloidal nanocrystal dispersions, we aimed to quantify *in situ* electron flux from planktonic

microbes in culture suspensions. We chose *Shewanella oneidensis* as a model electroactive

bacterium that is attractive for its respiratory plasticity and genetic tractability. Using non-

destructive optical measurements to track the LSPR peak in real-time, we successfully quantified

kinetic EET rate constants from *S. oneidensis* of varying metabolic activity and genotype. We also

quantitatively differentiated electron flux from an engineered *S. oneidensis* strain with tuned EET

gene expression levels. Our method enabled quantification of *S. oneidensis* population currents

from planktonic culture, with the average cell exhibiting $0.05 - 2.8$ fA $\cdot$ cell^{-1} depending on

metabolic activity, genotype, and gene expression level. Overall, our results indicate that

plasmonic semiconductor nanocrystals are a reliable sensing platform for probing metabolic

activity in cellular environments and contribute to a quantitative understanding of EET.

Results & Discussion

Preparation of Sn-doped indium oxide (ITO) nanocrystal aqueous dispersions.

ITO nanocrystals (5.78 ± 0.64 nm radius, 9.4 at% Sn) were synthesized using established colloidal

methods and a slow injection approach (Figure 1a, Materials and Methods) (38). Nanocrystal size

and Sn doping concentration were measured with small-angle X-ray scattering (SAXS) (Figure

S1) and inductively coupled plasma-optical emission spectroscopy, respectively. To disperse ITO

nanocrystals in aqueous solvents, the hydrophobic organic ligands bound to the surface of the

as-synthesized nanocrystals were replaced by a hydrophilic polymer. Specifically, we chose

poly(acrylic acid) grafted with methoxy-terminated poly(ethylene oxide) (PAA-mPEO$_4$). This

custom-designed and biocompatible random co-polymer has previously been used to

functionalize different types of metal oxides (39, 40) and been shown to be suitable for facilitating

electrochemical redox reactions in thin films of polymer-wrapped ITO nanocrystals (41).

Figure 1. Preparation of polymer-wrapped oxidized nanocrystals for EET biosensing. (a) Scanning
transmission electron microscopy image of as-synthesized ITO nanocrystals. Scale bar = 40 nm. (b)
Dynamic light scattering of charge-stabilized, polymer-wrapped, and oxidized polymer-wrapped ITO
nanocrystals. The ITO nanocrystals remain colloidally stable and do not exhibit signs of significant
aggregation after each processing step (inset) since the hydrodynamic diameter measurements of the three
samples are in close agreement. (c) Quantification of polymer adsorption on the ITO nanocrystal surface
using thermogravimetric analysis. The mass loss (18%) around 300 °C corresponds to the decomposition
of PAA-mPEO$_4$.

The functionalization of ITO nanocrystals with PAA-mPEO$_4$ was achieved in a two-step process.

First, the native stabilizing ligands were chemically removed by exposing the ITO nanocrystals to

nitrosonium tetrafluoroborate, which resulted in bare, charge-stabilized ITO nanocrystals

dispersible in N,N-dimethylformamide (30). Then, charge-stabilized ITO nanocrystals were mixed

with PAA-mPEO$_4$ and subsequently transferred into Milli-Q water. In a previous study, we found

that colloidal stability in aqueous media is due to the adsorption of PAA-mPEO$_4$ on ITO and is

likely facilitated by the coordination of carboxylate species at the nanocrystal surface (41). Fourier

transform infrared spectroscopy of nanocrystal dispersions at each processing step confirmed

that the native ligands were effectively removed and replaced by PAA-mPEO$_4$ (Figure S2). Finally,

polymer-wrapped ITO nanocrystals were chemically oxidized upon the addition of ammonium

cerium (IV) nitrate to promote EET from *S. oneidensis.* As expected, oxidation of the nanocrystal

dispersion caused a red-shift and decrease in intensity of the LSPR peak (Figure S3a). After

oxidation, the polymer-wrapped ITO nanocrystals were dispersed in a deuterated medium

(*Shewanella* Basal Medium, SBM, Table S1). Deuterated buffer avoided saturating the

spectrophotometer detector, as strong water absorption in the near-infrared (NIR) (42) (6000-

3000 cm^{-1}) overlaps significantly with the ITO LSPR peak (Note S1).

Avoiding the formation of large ITO nanocrystal aggregates in the dispersion before or during EET

is key for interpretation and analysis of optical measurements. Therefore, we monitored the

hydrodynamic diameter of the nanocrystal dispersion after each processing step (*i.e.,* ligand

removal, polymer wrapping, and oxidation post-polymer wrapping) using dynamic light scattering to probe the stability of the colloid (Figure 1b). The hydrodynamic diameters of the charge-stabilized and polymer-wrapped samples were centered around ~10 nm, which is in good agreement with the diameter obtained from SAXS. The slight increase in the hydrodynamic diameter of the polymer-wrapped and oxidized sample (D_h ~15 nm) could suggest interaction between the colloid and charged species during oxidation, but we did not detect signs of significant aggregation. This dispersion was colloidally stable for at least five months after oxidation and transfer into deuterated SBM (Figure S3b). We hypothesize that the colloidal stability of our aqueous dispersions of polymer-wrapped ITO (PAAPEO-ITO) is due to sufficient PAA-mPEO$_4$ adsorption on the nanocrystal surface (18% by mass, Figure 1c), which we quantified by thermogravimetric analysis, and thus effective steric stabilization.

S. oneidensis remains viable in the presence of PAAPEO-ITO.

All bacterial strains and plasmids used in this study are listed in Table S3. We first ensured that polymer-wrapped nanocrystals were not cytotoxic by quantifying cell viability in their presence (43). Aerobically pregrown cells were diluted to OD$_{600}$ = 0.2 in a mixture of ~1 mg/mL (~0.2 µM) PAAPEO-ITO and incubated for 2 h at 30 ℃. Unbound Live/Dead stain was washed away and cells were imaged using green (live) and red (dead) fluorescence channels. Under these conditions, *S. oneidensis* MR-1 (wild-type) remained predominately alive, with an 86 ± 2% viable population (Figure S4). To assess cell health for longer experiments, colony counting was used to quantify the viable population after 8 and 72 h incubation with PAAPEO-ITO and 20 mM lactate at room temperature in deuterated SBM with casamino acids. As expected, there were minimal effects on the viable population (Figure S5). Minor decreases in viable population were likely due to the lack of an available electron acceptor aside from ITO, which was only present at 0.3 µM (compared to standard growth conditions in 40 mM fumarate). In contrast to fumarate, a nanocrystal can accept more than one electron, but not enough to account for this concentration difference. These results indicate that cell viability is not substantially impacted by exposure to the polymer-wrapped nanocrystals.

Optical extinction of PAAPEO-ITO tracks respiratory electron flux from *S. oneidensis.*

Once cell viability was established, we measured the *in situ* optical response of the PAAPEO-ITO LSPR during electron transfer from *S. oneidensis* MR-1. Aerobically pregrown cells were washed

and resuspended in 1.5 mg/mL PAAPEO-ITO in deuterated SBM with casamino acids and 20

mM lactate as the carbon source. The solution was immediately transferred to a cuvette, sealed,

and loaded into a spectrophotometer in transmission mode to monitor the LSPR peak over time.

The time-resolved increase in optical density and blue shift of the LSPR peak indicate continuous

electron transfer to the nanocrystals (Figure 2a). These results confirm that *S. oneidensis* is able

to reduce PAAPEO-ITO and the kinetics of this process are resolvable using NIR spectroscopy.

Given the aerobic preparation of the cell culture and sample, they also suggest that dissolved

oxygen is first consumed through aerobic respiration, followed by activation of anaerobic EET

pathways (*i.e.*, reduction of PAAPEO-ITO), as we have previously observed for molecular electron

acceptors (44, 45). We then measured EET from a *S. oneidensis* MR-1 population deprived of a

carbon source (*e.g.*, lactate), which should inhibit respiration and prevent generation of electron

flux. The change in optical extinction was significantly diminished in these samples (Figure 2b).

Some minimal reduction was observed, which can likely be attributed to excess metabolism from

pregrowth (7), or to non-specific reducing capacity from cell stress and lysis. Indeed, a reaction

containing lysed *S. oneidensis* slightly reduced PAAPEO-ITO even without oxygen removal,

indicating that cell death could lead to a small amount of nonspecific reduction (Figure S6).

Similarly, an *E. coli* sample demonstrated minimal PAAPEO-ITO reduction, as would be expected

from this EET-deficient microbe (Figure 3c). Together, these measurements confirm that

regulating metabolic flux manifests in an observable difference to the optical response of

PAAPEO-ITO, validating this platform as an effective *in situ* biosensor that is specific to EET.

192
193
194 **Figure 2. EET from metabolically active *S. oneidensis* leads to time-resolved increased optical**
195 **extinction of PAAPEO-ITO.** Experimental extinction spectra collected each hour from *t* = 0 to 8 h and
196 associated fits for PAAPEO-ITO inoculated with (a) *S. oneidensis* MR-1 that was provided 20 mM lactate
197 as a carbon source, (b) *S. oneidensis* MR-1 not provided a carbon source, or (c) *E. coli* MG1655 provided
198 with 20 mM lactate as a carbon source.
199

Quantitative spectral analysis enables quantification of EET from *S. oneidensis* with varying metabolic activity and genotype.

Following qualitative observations of the LSPR peak over time, we fit the peaks to a free electron model that extracts the number of conductive electrons in a nanoparticle. Using a deuterated buffer allowed for the collection of clean spectra and nearly the full plasmon peak for the ITO nanocrystals could be resolved. ITO nanocrystals were previously incubated with *S. oneidensis*, but only to quantify total electrons transferred from wild-type *S. oneidensis*, and only in non-deuterated buffer (35). Therein the water absorption obscured the signal below ~5500 cm^{-1}, meaning the high energy tail of the LSPR peak and the ultraviolet band gap absorption were used to quantify electron transfer. While that work was an important step towards quantifying EET using ITO nanocrystals, our ability to resolve nearly the full plasmonic response allows for a simpler, more reliable quantification using our recently developed model, the heterogeneous ensemble Drude approximation (HEDA). The HEDA model extracts intrinsic material properties, such as electron concentration, from the plasmonic response of a dilute dispersion of nanocrystals (37). By accounting for ensemble heterogeneities in nanocrystal size and nanocrystal electron concentration, it enables more accurate acquisition of material properties.

Immediately upon inoculating the cells with oxidized PAAPEO-ITO, we began collecting time-resolved spectra, continuing for up to 48 hours. We fit spectra using the HEDA model to quantify the number of electrons within the average nanocrystal in an ensemble at each timepoint (Note S2). Since we quantified both the concentration of nanocrystals and cells, we could calculate the cumulative electrons transferred per cell over time. This quantification revealed two consistent electron transfer regimes across samples – a variable regime during the first ~1 h, where electron transfer was nonlinear (Figure 3a, inset; Figure S7a), followed by a steady-state electron transfer regime. In previous studies, aerobic *S. oneidensis* at OD$_{600}$ = 0.2 took between 10 min to 1 h to fully consume dissolved oxygen and turn on anaerobic EET pathways, depending on the electron acceptor and reaction conditions (44, 45). Therefore, we attribute this initial variable domain to a combination of oxygen removal via aerobic respiration and cell stress as they adjust to a new electron acceptor and undergo a metabolic shift.

Beyond 8 h of reaction in high EET samples (*e.g.*, wild-type *S. oneidensis* with lactate), we noticed effects of nanocrystal saturation and precipitation (Figure 3a, Figure S7b). Thus, we defined the steady-state regime as between 1 and 8 h, and used this domain to determine electron transfer

kinetics. Notably, OD_{600} = 0.2 is approximately the saturation density of anaerobic *S. oneidensis*

in SBM with 20 mM lactate. Upon adjusting to their new anaerobic environment, the cells should

divert minimal metabolic flux toward growth and instead direct it toward energy production via

EET (46). This, combined with the lack of cell growth between inoculation and 8 h (Figure S5),

validates the appearance of steady-state electron transfer in our system. As such, the linear

increase in electrons transferred per cell justified using linear regressions to obtain kinetic rate

constants and compute population-averaged current generation.

242
Figure 3. HEDA fitting of PAAPEO-ITO LSPR response during EET from *S. oneidensis* enables
quantification of steady-state electron transfer kinetics. (a) EET from *S. oneidensis* to PAAPEO-ITO
exhibits three domains. (Inset) The early domain (≤ 1 h) shows variability due to aerobic respiration,
introduction of a deuterated solvent, and cell stress. This regime includes a lag phase (0–20 min) followed
by a sharp increase in electron transfer (20–60 min). Steady-state electron transfer (1–8 h) precedes a later
domain (≥ 8 h) exhibiting effects of nanocrystal saturation, precipitation, and cell death. (b) The Mtr pathway
in *S. oneidensis*. In the Δ*mtrC*Δ*omcA*Δ*mtrF* knockout (right), three critical extracellular cytochrome genes
are removed from the genome, hindering EET capabilities. (c) Steady-state EET (1–8 h) where dashed
lines represent linear regression of the electrons transferred over time and k_{EET} is the slope of the regression
(Table 1). Data points shown are the mean ± SD of *n* = 3 fits of biological replicates.
253

We first determined steady-state electron transfer rates to PAAPEO-ITO for *S. oneidensis* MR-1

with and without lactate as a carbon source, which qualitatively showed a large difference in

optical response (Figure 2). When 20 mM lactate was present as a carbon source, enabling

continuous respiration and generation of EET flux, the rate of electron transfer was k_{EET} = 2.0 ±

0.2 x 10^7 electrons • cell^{-1} • h^{-1}, or 0.89 ± 0.07 fA • cell^{-1}, which is approximately 5-fold greater than

when the cells were starved (Table 1). To further assess background electron transfer and

dynamic range, we used an EET-deficient strain of *S. oneidensis*. *S. oneidensis*

Δ*mtrC*Δ*omcA*Δ*mtrF* is a triple-knockout of three key extracellular cytochromes responsible for

direct electron transfer to a variety of substrates (Figure 3b), and as a result, exhibits significantly

hindered electron transfer activity toward both electrodes and soluble acceptors (11, 14). As

expected, this strain showed significantly attenuated electron transfer to PAAPEO-ITO (Figure

3c). Additionally, there was an order-of-magnitude decrease in the electron transfer rate constant
compared to the wild-type strain (Table 1). It should be noted that statistical error associated with
accumulated electron transfer is dominated by variation in *S. oneidensis* EET rates among
biological replicates, not from error associated with spectra collection or fitting (Figure S8). The
collected spectra are highly resolved, meaning the signal is stable over the collection time period
of ~1 minute and the fitting procedure consistently matches experimental data and converges to
the same solution even for a range of randomly generated initial guesses (Table S6).
272
273 **Table 1. Metrics obtained from HEDA fitting during EET from *S. oneidensis*.** Data shown are results
274 of the steady-state regime, between 1 and 8 h, and are the mean ± SD of *n* = 3 HEDA model fits of biological
275 replicates.
276

Sample	e⁻ Transferred • Cell⁻¹	k_{EET} (e⁻ • cell⁻¹ • h⁻¹)	Current • Cell⁻¹ (fA)
S. oneidensis MR-1 (+ 20 mM lactate)	$(1.3 \pm 0.2) \times 10^8$	$(2.0 \pm 0.2) \times 10^7$	0.89 ± 0.07
S. oneidensis MR-1 (− 20 mM lactate)	$(2.5 \pm 3.1) \times 10^7$	$(4.8 \pm 2.4) \times 10^6$	0.21 ± 0.11
S. oneidensis Δ*mtrC*Δ*omcA*Δ*mtrF* (+ 20 mM lactate)	$(0.9 \pm 1.4) \times 10^7$	$(1.2 \pm 1.2) \times 10^6$	0.05 ± 0.05

277
278
Although the physical dimensions of the nanocrystals (5.78 ± 0.64 nm radius, compared to cell
length on the ~1 μm scale) should prevent biofilm formation, we sought to ensure that EET was
occurring from planktonic cultures. Therefore, we tested a biofilm-deficient knockout (*S.*
oneidensis S2933, termed Δlysis-operon) (47). This strain did not exhibit significantly different
current generation than wild-type *S. oneidensis*, indicating that electron transfer is occurring from
planktonic cells (Figure 4). We repeated the wild-type sample as this experiment was performed
with a newly synthesized batch of nanocrystals compared to Figure 3 (Note S3).
286
287

Figure 4. Planktonic cells are responsible for EET to PAAPEO-ITO. Extinction spectra for PAAPEO-ITO dispersions incubated with (a) *S. oneidensis* MR-1 and (b) and *S. oneidensis* S2933 (Δlysis-operon, a biofilm-deficient knockout) (47). (c) Steady-state EET (1–8 h) where dashed lines represent linear regression of the electrons transferred over time and k_{EET} is the slope of the regression (Table 1). For MR-1, k_{EET} = (2.4 ± 0.2) x 10^7 e^- • $cell^{-1}$ • h^{-1}; for S2933, k_{EET} = (2.1 ± 0.4) x 10^7 e^- • $cell^{-1}$ • h^{-1}. Data points shown are the mean ± SD of n = 3 HEDA model fits of biological replicates.

All fit results are reported in Tables S4-7. Overall, current quantified from the wild-type strain with 20 mM lactate is comparable to results from other methods that have measured electron flux from *S. oneidensis*. These studies found current generation from either *S. oneidensis* populations (~2 fA • $cell^{-1}$) (48) or from single cells trapped on electrodes (~100 fA • $cell^{-1}$), although such studies typically exclude non- and low-performing single cells, which would change the population average (18, 19). In addition, the necessity of deuterium in our buffer would lead to kinetic isotope effects on proton pumping and therefore electron transfer kinetics. Previous studies have determined that *S. oneidensis* can recover from this and reach deuterium efflux equilibrium (49); however, this may contribute to slightly lower overall current generation measured in our system compared to others (Note S1). Overall, these results contribute to a quantitative understanding of EET from planktonic cell populations and suggest that simple spectroscopic techniques can be used as a screening tool to elucidate the electroactive capabilities of various species or strains.

EET output from engineered *S. oneidensis* with variable *mtrC* expression can be quantitatively differentiated using PAAPEO-ITO.

Synthetic biologists commonly use genetic circuits to control protein output, and emergent applications of EET will likely employ similar strategies. However, unlike fluorescence or secreted protein, it is difficult to directly quantify the relationship between EET gene expression and resultant electron flux, especially in planktonic cultures. Toward this goal, we used an *S. oneidensis* strain that tailors expression of a critical EET gene, *mtrC*, in response to inducing

molecule concentration. Genetically engineered *S. oneidensis* Δ*mtrC*Δ*omcA*Δ*mtrF* + *mtrC*
proportionally expresses *mtrC* in the presence of isopropyl ß-D-1-thiogalactopyranoside (IPTG)
as an inducer (12, 45). Cultures of *S. oneidensis* Δ*mtrC*Δ*omcA*Δ*mtrF* + *mtrC* were pregrown
anaerobically in varying concentrations of IPTG (ranging from 0 to 1000 μM), leading to differing
expression levels of *mtrC* and resultant EET flux. Cultures were then washed and normalized to
OD_{600} = 0.2 in a reaction mixture containing PAAPEO-ITO, 20 mM lactate, 25 μg/mL kanamycin
for plasmid maintenance, and the corresponding concentration of inducing molecule. As outlined
above, spectra were collected through 8 hours and fit with the HEDA model (Figure 5a–d). Fit
results are reported in Table S8. As expected, the resultant electron transfer rate during steady-
state was a function of IPTG concentration (Figure 5e), increasing about 5.5-fold from no
induction/leaky expression (0 μM IPTG) to full induction (1000 μM IPTG). The consistent linear
electron transfer rate across all engineered strains after 1 h again supports steady-state electron
transfer directed toward energy production. A wild-type (MR-1), empty vector control grown in
1000 μM IPTG also showed a high electron transfer rate, even higher than the wild-type without
a plasmid, which could be attributed to the presence of antibiotic, physiological changes from
plasmid maintenance, and/or an anaerobic pregrowth. In this sample, saturation effects were
observed at 4 hours, so the rate constant was derived from only the preceding timepoints during
steady-state electron transfer (Figures S9–10). It is notable that even the fully induced *mtrC*
complement did not completely recapture the wild-type EET phenotype, exhibiting ~4.5-fold less
activity compared to the empty vector control. This may highlight the importance of other missing
EET components in this strain, specifically *omcA* and *mtrF*, in binding and electron transfer to
metal oxide substrates (50, 51). Tiered electron flux based on varying gene expression was also
quantified as current output, ranging from 0.12 ± 0.04 (uninduced) to 0.68 ± 0.08 (fully induced)
fA • cell^{-1} at steady-state (Table 2). The rate constants were fit with an activating Hill Function
gene expression model as a guide to the eye; as expected, increasing concentrations of IPTG
correspond to increasing values of k_{EET} (Figure 5f, Note S4). Together, these results exemplify
the quantitative capabilities of our method for measuring electron flux after varying expression
levels of a single EET gene. This deepens our understanding of the relationship between *mtrC*
expression and EET output to heterogenous acceptors in planktonic cultures, and may be used
to predict electron flux based on *mtrC* expression for a desired application.
348

Figure 5. HEDA fitting of PAAPEO-ITO LSPR response during EET from engineered *S. oneidensis*
reveals gradients in population current output as a result of differential gene expression. *In situ*
extinction spectra and HEDA fits for PAAPEO-ITO during EET from (a–d) *S. oneidensis ΔmtrCΔomcAΔmtrF*
+ *mtrC* with (a) 1000, (b) 100, (c) 10, and (d) 0 µM inducing molecule (IPTG). (e) Steady-state EET (1–8 h)
where dashed lines represent linear regression of the electrons transferred over time and k_{EET} is the slope
of the regression (Table 2). Data points shown are the mean ± SD of *n* = 3 HEDA model fits of biological
replicates. (f) Rate constants exhibit an activating Hill Function-type response to increasing IPTG
concentration. Data points shown are the mean ± SD of *n* = 3 independent linear regressions of HEDA
model fits of biological replicates.

Table 2. Metrics obtained from HEDA fitting during EET from engineered *S. oneidensis* MR-1 +
***empty* and *ΔmtrCΔomcAΔmtrF* + *mtrC*.** Data shown are results of the steady-state regime, between 1
and 8 h for all samples except for MR-1 + *empty*, which was between 1 and 3 h. Data shown are the mean
± SD of *n* = 3 HEDA model fits of biological replicates.

Sample	e⁻ Transferred • Cell⁻¹	k_{EET} (e⁻ • cell⁻¹ • h⁻¹)	Current • Cell⁻¹ (fA)
S. oneidensis MR-1 + *empty* (1000 µM IPTG)	$(3.2 \pm 1.0) \times 10^8$	$(6.2 \pm 1.3) \times 10^7$	2.76 ± 0.60
ΔmtrCΔomcAΔmtrF + *mtrC* (1000 µM IPTG)	$(1.0 \pm 0.3) \times 10^8$	$(1.5 \pm 0.2) \times 10^7$	0.68 ± 0.08
ΔmtrCΔomcAΔmtrF + *mtrC* (100 µM IPTG)	$(8.3 \pm 1.9) \times 10^7$	$(1.2 \pm 0.1) \times 10^7$	0.53 ± 0.05
ΔmtrCΔomcAΔmtrF + *mtrC* (10 µM IPTG)	$(4.1 \pm 1.3) \times 10^7$	$(5.6 \pm 1.1) \times 10^6$	0.25 ± 0.05
ΔmtrCΔomcAΔmtrF + *mtrC* (0 µM IPTG)	$(1.9 \pm 1.3) \times 10^7$	$(2.6 \pm 1.0) \times 10^6$	0.12 ± 0.04

Conclusions

Together, our results demonstrate that the sensing capabilities of plasmonic ITO nanocrystals in microbial suspensions can address outstanding challenges in quantifying biological electron transfer. Using our platform, we successfully measured current generation on insoluble substrates from planktonic cultures, enabling differentiation of *S. oneidensis* cells of varying metabolic activity and genotype. In addition, we quantified changes in current generation by cells engineered to express varying levels of a critical EET gene, *mtrC*. This will enable fundamental studies of biological electron transfer metrics and resolution of EET gene expression dynamics. For example, PAAPEO-ITO biosensors can explore, in real time, how cells change electron flux in response to their environment, including the presence of a new electron acceptor or an inducing molecule. This non-destructive, *in situ* spectroscopic measurement could also be integrated with high-throughput technologies such as well plates and flow cytometers to screen genetic mutants or new bacterial strains based on electron transfer rate, in addition to metrics like growth rate and fluorescent protein expression.

Our recent progress in both surface functionalization and spectroelectrochemical modeling led to PAAPEO-ITO as an accurate EET sensing platform; however, further advances in each of these areas will enable even more applications. For example, continuing to develop novel nanocrystal surface chemistry, such as functionalizing with bioactive polymers, could expand our capability to target cellular machinery. In addition, standardizing the optical signatures of ITO nanocrystals with known surface potentials could allow both electron tracking and mapping of potential energy landscapes across the cell. Overall, our design highlights concurrent progress in nanocrystal

390   biosensors and synthetic biology for EET, enabling more effective evaluation of critical biological

391   processes.

Materials and Methods

394
Chemicals and reagents: Deuterium oxide (D_2O, Sigma-Aldrich, 99.9 atom % D), sodium DL-lactate
($NaC_3H_5O_3$, TCI, 60% in water), sodium fumarate ($Na_2C_4H_2O_4$, VWR, 98%), HEPES buffer solution
($C_8H_{18}N_2O_4S$, VWR, 1 M in water, pH = 7.3), potassium phosphate dibasic (K_2HPO_4, Sigma-Aldrich),
potassium phosphate monobasic (KH_2PO_4, Sigma-Aldrich), sodium chloride (NaCl, VWR), ammonium
sulfate (($NH_4)_2SO_4$, Fisher Scientific), magnesium (II) sulfate heptahydrate ($MgSO_4 \cdot 7H_2O$, VWR), trace
mineral supplement (ATCC), casamino acids (VWR), silicone oil (Alfa Aesar), N,N-dimethylformamide
(DMF, Sigma-Aldrich, ≥99.8%), nitrosonium tetrafluoroborate ($NOBF_4$, Sigma-Aldrich, 95%), ammonium
cerium (IV) nitrate (Sigma-Aldrich, ≥98.5%), toluene (Fisher Scientific, ≥99.5%), tin (IV) acetate (Sigma
Aldrich), indium (III) acetate (STREM 99.99%), oleic acid (Sigma Aldrich, technical grade, 90%), oleyl
alcohol Sigma Aldrich, technical grade, 85%), hexane (Fisher Scientific, various methylpentanes 4.2%,
≥98.5%) isopropyl ß-D-1-thiogalactopyranoside (IPTG, Teknova), kanamycin sulfate ($C_{18}H_{38}N_4O_{15}S$,
Growcells), nail polish (Electron Microscopy Sciences), BacLight Live/Dead Stain (Invitrogen) were all used
as purchased unless otherwise noted. All media components were autoclaved or sterilized using 0.2 μm
PES filters.

409
Nanocrystal synthesis: Tin-doped indium oxide nanocrystals were synthesized using a slow-injection
approach pioneered by the Hutchison group (38, 52). Indium acetate and tin acetate in a 9:1 molar ratio
were dissolved in oleic acid at an overall concentration of 0.5 mmol metal per mL oleic acid. The solution
was heated to 150°C and kept at that temperature for at least two hours to convert the metal-acetate
complexes into the metal-oleate precursor. Using a syringe pump, the precursor was injected into a hot
bath (290 °C) of oleyl alcohol at a rate of 0.3 mL/min. For this work, 4.5 mL of precursor was injected. The
nanocrystals were collected from the reaction mixture and purified by flocculating with ethanol, centrifuging
at 9000 rpm, and dispersing in hexane three consecutive times.

418
Nanocrystal ligand stripping: $NOBF_4$ was used to chemically remove the native organic ligands bound
to the nanocrystal surface following a previously established method (30). Briefly, 60 mg of $NOBF_4$ were
added to a two-phase mixture of DMF (2 mL) and ITO nanocrystal dispersion in hexane (~50 mg/mL, 2
mL). The mixture was sonicated for 30 min to promote ligand removal and therefore the phase transfer of
nanocrystals from hexane to DMF. After discarding the hexane layer, charge-stabilized ITO nanocrystals
dispersed in DMF were purified by performing seven cycles of flocculation with toluene, centrifugation at
7500 rpm for 5 min, and redispersion in DMF.

426
Nanocrystal polymer wrapping: PAA-mPEO$_4$ random copolymer was synthesized (39, 40) and ITO
nanocrystals were polymer-wrapped with PAA-mPEO$_4$ following previously established methods (41).
Briefly, a PAA-mPEO$_4$ solution was prepared by dissolving 10 mg of polymer in 700 μL of DMF. Then, 300
μL of ITO nanocrystal dispersed in DMF (~90 mg/mL) were added dropwise to the PAA-mPEO$_4$ solution
under gentle stirring. This nanocrystal-polymer mixture was stirred overnight and subsequently added
dropwise into 19 mL of Milli-Q water while stirring. The aqueous mixture was stirred at 600 rpm for 48 hours.
Polymer-wrapped ITO nanocrystals were recovered and purified by performing three cycles of spin dialysis
(centrifugation at 4000 rpm for 10 min per cycle) using 50 kDa Millipore Amicon Ultra centrifugal tubes. The
final dispersion in Milli-Q water was filtered through a 0.45 μm PVDF syringe filter (Acrodisc, Pall).

436
Oxidation of polymer-wrapped nanocrystals: Polymer-wrapped ITO nanocrystals dispersed in Milli-Q
water were chemically oxidized following a previously established method (35). Briefly, 120 μL of polymer-
wrapped ITO (~7 mg/mL) were mixed with 25 μL of an aqueous solution of ammonium cerium nitrate (25
mM). Instead of scaling up the volume of the reaction, the reagents were combined in multiple separate
vials to achieve the desired final mass of oxidized and polymer-wrapped nanocrystals. The mixture was left
to stand without stirring for 15 min and then purified by performing three cycles of spin dialysis
(centrifugation at 4000 rpm for 10 min per cycle) using 50 kDa Millipore Amicon Ultra centrifugal tubes. The

resulting oxidized and polymer-wrapped ITO nanocrystals were then dispersed in deuterated SBM + CAAs
(Table S1) by performing a solvent exchange through three additional spin dialysis cycles with excess
deuterated buffer instead of Milli-Q water (final concentration: 1.5 mg/mL).

447
Nanocrystal Characterization: a. Electron microscopy. ITO nanocrystals were imaged in bright-field
scanning transmission electron microscopy mode at a 30 kV accelerating voltage on a Hitachi S5500
SEM/STEM instrument. The sample was drop-cast on Type-A ultrathin carbon copper TEM grids (Ted
Pella, 01822, 400 mesh) from dilute ITO dispersions in hexane. **b. Small-angle X-ray scattering (SAXS).**
The size of as-synthesized ITO nanocrystals was determined by SAXS. A dilute dispersion (~ 1 mg/mL),
enclosed in a sealed glass capillary (Charles-Supper Company, Boron Rich, 1.5 mm diameter, 0.01 mm
wall thickness), was measured in transmission configuration at a sample-detector distance of 1 m on a
SAXSLAB Ganesha instrument using Cu Kα radiation. Details of SAXS data analysis are included in the
Supplementary Information. **c. Dynamic light scattering.** The hydrodynamic diameters of ligand-stripped,
polymer-wrapped, and oxidized ITO nanocrystals were measured on a Malvern Zetasizer Nano ZS. Dilute
ITO nanocrystals dispersions (~1 mg/mL) were enclosed in disposable plastic micro cuvettes (ZEN0040,
Malvern). **d. Thermogravimetric analysis.** PAA-mPEO$_4$ adsorption on the ITO nanocrystal surface was
quantified using a Mettler Toledo TGA 2. In a typical experiment, 100 µL of polymer-wrapped ITO
nanocrystals dispersed in Milli-Q water (~10 mg/mL) was added to a disposable aluminum crucible and
dried under vacuum at room temperature for 24 hours. The sample was measured in dry synthetic air from
30 ºC to 600 ºC at a 5 ºC/min ramp rate. **e. Fourier transform infrared spectroscopy.** Ligand-capped,
ligand-stripped, and polymer-wrapped ITO nanocrystals were measured in transmission geometry with a
Bruker Vertex 70 spectrometer. Samples were drop-cast from dilute nanocrystal dispersion (~1 mg/mL) on
undoped, double side polished silicon substrates (Virginia Semiconductor Inc.) **f. UV/Vis/NIR**
Spectroscopy. The cuvettes were placed in an Agilent Cary series ultraviolet-visible-near infrared
spectrophotometer in transmission mode for spectra collection. Spectra were collected from 3032-10000
cm^{-1} for a collection time of ~1 min per spectrum. **g. Inductively coupled plasma-optical emission**
spectroscopy (ICP-OES). To quantify the volume fraction of nanocrystals in dispersion, as well as the
doping percentage of tin atoms in the indium oxide lattice, we took a known volume of the dispersed
nanocrystals, dried them into a pellet and dissolved them in aqua regia for two days. We then diluted the
dissolved ions to 2% v/v acid and loaded the dilute acid into the Varian 720-ES ICP-OES to quantify the tin
and indium concentrations. We calibrated the instrument for tin and indium using standards purchased from
Sigma Aldrich (TraceCERT® 1000mg In/L nitric acid and 1000mg Sn/L nitric acid).

476
Bacterial strains and culture: Bacterial strains and plasmids are listed in Table S3. Cultures were
prepared as follows: bacterial stocks stored in 22.5% glycerol at -80 ºC were streaked onto LB agar plates
(for wild-type and knockout strains) or LB agar with 25 µg/mL kanamycin (for plasmid-harboring strains)
and grown overnight at 30 ºC for *Shewanella*, 37 ºC for *E. coli*. Single colonies were isolated and inoculated
into *Shewanella* Basal Medium (SBM) supplemented with 100 mM HEPES, 0.05% trace mineral
supplement (Table S2), 0.05% casamino acids, and 20 mM sodium lactate (2.85 µL of 60% w/w sodium
lactate per 1 mL culture) as the electron donor. Aerobic cultures were pregrown in 15 mL culture tubes at
30 ºC and 250 rpm shaking. Anaerobic, plasmid-harboring cultures were pregrown using the same
procedure outlined above, but in degassed growth medium in a humidified anaerobic chamber,
supplemented with 40 mM sodium fumarate (40 µL/mL of a 1 M stock) as the electron acceptor, 25 µg/mL
kanamycin, and varying concentrations of IPTG. Descriptions of the *mtrC* circuit and pCD24r1 plasmid, as
well as the empty vector circuit and the pCD8 plasmid, have been published previously (12). Cultures were
washed 3x after pregrowth using SBM supplemented with 0.5% casamino acids (degassed for anaerobic
cultures). Before the final centrifugation, OD$_{600}$ was measured using a NanoDrop 2000C spectrophotometer
and the culture volume normalized such that the final OD$_{600}$ in the reaction with PAAPEO-ITO would be
0.2.

493
Viability assessments: All microscopy was performed using a Nikon Ti2 Eclipse inverted epifluorescence
microscope. Cells assessed for viability by microscopy were incubated with PAAPEO-ITO for 2 h at 30 ºC.
The cells were centrifuged, and the nanocrystals were washed away using 0.85% NaCl solution. The cell
suspensions were then incubated in the dark in the BacLight Live/Dead stain mix (1.5 µL/mL Syto9, 2.5
µL/mL propidium iodide in 0.85% NaCl solution) for 20 minutes. Stained cells were then washed 3x in 0.85%
NaCl solution to remove unbound dye. Cell suspensions were loaded onto glass microscope slides and a

no. 1 coverslip placed on top. Nail polish was used to seal the coverslip to the glass slide to prevent
evaporative losses. Fluorescence for each stain (green for Syto9, red for propidium iodide) was measured
using GFP and Texas Red excitation/emission filter cubes on a Nikon Ti2 Eclipse, as outlined previously[21].
Live and dead populations were quantified using thresholding and masking in Fiji 1.0. For colony counting,
cell samples before and after incubation with PAAPEO-ITO were diluted by factors of 10^5-10^7 in LB medium,
and 100 µL of these dilutions was plated onto LB-agar. The viable population (CFU/mL) for each sample
was quantified by averaging $\frac{\text{CFU}}{\text{mL}} = \frac{\text{\# of colonies} \cdot \text{dilution}}{(100\ \mu\text{L} / 1000\ \mu\text{L} \cdot \text{mL}^{-1})}$ for each replicate.

507
Electron transfer experiments and spectroscopy measurements:
Bacteria and PAAPEO-ITO nanocrystals were mixed in a disposable centrifuge tube and immediately
transferred to a Spectrocell near-IR quartz cuvette and the cuvette was sealed for the duration of the
experiment. The cuvette was loaded into an Agilent Cary UV/vis/NIR spectrophotometer for spectra
collection. On average, the first spectrum was taken ~3 min after mixing. Thereafter, a spectrum was
collected every 10 minutes for the first hour and then every hour until 8 hours. Three cuvettes were used
to carry out triplicate experiments simultaneously, two with pathlengths of 1 mm and the third with
pathlength 0.5 mm. All spectra were backgrounded to a cuvette filled solely with deuterated SBM buffer,
without PAAPEO-ITO or bacteria.

Author Contributions: A.J.G., S.L.G., C.A.S.C., D.J.M., and B.K.K. conceived the project; A.J.G., S.L.G.,
C.A.S.C., Y.W., and A.M.G. performed experiments; S.L.G., C.A.S.C., and Y.W. performed nanocrystal
synthesis and characterization; A.J.G. designed biological experiments and performed microbial culture;
A.J.G., S.L.G., C.A.S.C., D.J.M., and B.K.K. analyzed the results; A.J.G., S.L.G., C.A.S.C., D.J.M., and
B.K.K. wrote the manuscript with input from all authors.

Acknowledgements: *S. oneidensis ΔmtrCΔomcAΔmtrF* was a generous gift from Prof. Jeffrey Gralnick (U.
Minnesota), and *E. coli* MG1655 and *S. oneidensis* S2933 were generous gifts from Prof. Lydia Contreras
(UT Austin). Christopher Dundas is thanked for his contribution of the engineered *S. oneidensis* strain.
A.J.G. and S.L.G. were supported through National Science Foundation Graduate Research Fellowships
(Program Award No. DGE-1610403). Portions of this work, including polymer synthesis and
characterization by Swagat Sahu and Brett A. Helms, were carried out as a user project at the Molecular
Foundry, which is supported by the Office of Science, Office of Basic Energy Sciences of the U.S.
Department of Energy under contract no. DE-AC02-05CH11231. This research was primarily supported by
the National Science Foundation through the Center for Dynamics and Control of Materials: an NSF
Materials Research Science and Engineering Center under DMR-1720595. Additional research support
was provided by the Welch Foundation (Grants F-1929 and F-1848), the National Institute of General
Medical Sciences of the National Institutes of Health under Award Number R35GM133640 (B.K.K.), and
an NSF CAREER Award (1944334, B.K.K.). Additional research support was provided by the Welch
Foundation (Grants F-1929 and F-1848) and by National Institute of General Medical Sciences of the
National Institutes of Health under Award Number R35GM133640 (B.K.K.). The authors acknowledge use
of shared research facilities supported in part by the Texas Materials Institute including the SAXS
instrument acquired under an NSF MRI grant (CBET-1624659), the Center for Dynamics and Control of
Materials: an NSF MRSEC (DMR-1720595), and the NSF National Nanotechnology Coordinated
Infrastructure (ECCS-1542159). We gratefully acknowledge the use of facilities within the core microscopy
lab of the Institute for Cellular and Molecular Biology, University of Texas at Austin.

ORCID:
Austin J. Graham: 0000-0001-8924-181X
Stephen L. Gibbs: 0000-0003-2533-0957
Camila A. Saez Cabezas: 0000-0002-8734-0096
Allison M. Green: 0000-0002-7734-4202
Delia J. Milliron: 0000-0002-8737-451X
Benjamin K. Keitz: 0000-0003-3314-0053

References

1. L. Shi, *et al.*, Extracellular electron transfer mechanisms between microorganisms and minerals. *Nat. Rev. Microbiol.* **14**, 651–662 (2016).

2. J. M. Flynn, D. E. Ross, K. A. Hunt, D. R. Bond, J. A. Gralnick, Enabling Unbalanced Fermentations by Using Engineered Electrode-Interfaced Bacteria. *mBio* **1**, e00190-10-e00190-17 (2010).

3. D. R. Bond, "Electrodes as electron acceptors, and the bacteria who love them" in *Geomicrobiology: Molecular and Environmental Perspective*, L. Barton, M. Mandl, A. Loy, Eds. (Springer, 2010), pp. 385–399.

4. D. E. Ross, J. M. Flynn, D. B. Baron, J. A. Gralnick, D. R. Bond, Towards Electrosynthesis in *Shewanella*: Energetics of Reversing the Mtr Pathway for Reductive Metabolism. *PLoS ONE* **6**, e16649-9 (2011).

5. Y. Gao, M. Mohammadifar, S. Choi, From Microbial Fuel Cells to Biobatteries: Moving toward On-Demand Micropower Generation for Small-Scale Single-Use Applications. *Adv. Mater. Technologies* **4**, 1900079 (2019).

6. X. Fang, S. Kalathil, G. Divitini, Q. Wang, E. Reisner, A three-dimensional hybrid electrode with electroactive microbes for efficient electrogenesis and chemical synthesis. *Proc. Natl. Acad. Sci. U.S.A.* **9**, 5074–5080 (2020).

7. G. Fan, C. M. Dundas, A. J. Graham, N. A. Lynd, B. K. Keitz, *Shewanella oneidensis* as a living electrode for controlled radical polymerization. *Proc. Natl. Acad. Sci. U.S.A.* **4**, 4559–4564 (2018).

8. C. M. Dundas, A. J. Graham, D. K. Romanovicz, B. K. Keitz, Extracellular Electron Transfer by *Shewanella oneidensis* Controls Palladium Nanoparticle Phenotype. *ACS Synth. Biol.* **7**, 2726–2736 (2018).

9. N. Schuergers, C. Werlang, C. M. Ajo-Franklin, A. A. Boghossian, A synthetic biology approach to engineering living photovoltaics. *Energ. Environ. Sci.* **10**, 1102–1115 (2017).

10. B. Tian, *et al.*, Roadmap on semiconductor–cell biointerfaces. *Phys Biol* **15**, 031002 (2018).

11. D. Coursolle, J. A. Gralnick, Modularity of the Mtr respiratory pathway of *Shewanella oneidensis* strain MR-1. *Mol. Microbiol.* **55**, 995–1008 (2010).

12. C. M. Dundas, D. J. F. Walker, B. K. Keitz, Tuning Extracellular Electron Transfer by *Shewanella oneidensis* Using Transcriptional Logic Gates. *ACS Synth Biol* (2020) https:/doi.org/10.1021/acssynbio.9b00517.

13. J. T. Atkinson, *et al.*, Metalloprotein switches that display chemical- dependent electron transfer in cells. *Nat. Chem. Biol.*, 1–10 (2018).

14. D. Coursolle, D. B. Baron, D. R. Bond, J. A. Gralnick, The Mtr Respiratory Pathway Is Essential for Reducing Flavins and Electrodes in *Shewanella oneidensis*. *J. of Bacteriol.* **192**, 467–474 (2009).

15. A. R. Rowe, *et al.*, Tracking Electron Uptake from a Cathode into *Shewanella* Cells: Implications for Energy Acquisition from Solid-Substrate Electron Donors. *mBio* **9**, e02203-17–19 (2018).

16. Q. Wang, A.-A. D. Jones, J. A. Gralnick, L. Lin, C. R. Buie, Microfluidic dielectrophoresis illuminates
the relationship between microbial cell envelope polarizability and electrochemical activity. *Sci. Adv.* **5**,
eaat5664 (2019).

17. H. Liu, G. J. Newton, R. Nakamura, K. Hashimoto, S. Nakanishi, Electrochemical Characterization of
a Single Electricity-Producing Bacterial Cell of *Shewanella* by Using Optical Tweezers. *Angew. Chem. Int.*
49, 6596–6599 (2010).

18. B. J. Gross, M. Y. El-Naggar, A combined electrochemical and optical trapping platform for measuring
single cell respiration rates at electrode interfaces. *Rev. Sci. Instrum.* **86**, 064301–9 (2015).

19. S. Pirbadian, M. S. Chavez, M. Y. El-Naggar, Spatiotemporal mapping of bacterial membrane
potential responses to extracellular electron transfer. *Proc. Natl. Acad. Sci. U.S.A.* 202000802 (2020).

20. G. Méhes, *et al.*, Organic Microbial Electrochemical Transistor Monitoring Extracellular Electron
Transfer. *Adv. Sci.*, 2000641 (2020).

21. H. H. Hau, J. A. Gralnick, Ecology and Biotechnology of the Genus *Shewanella*. *Ann. Rev. of*
Microbiol. **61**, 237–258 (2007).

22. E. D. Brutinel, J. A. Gralnick, Shuttling happens: soluble flavin mediators of extracellular electron
transfer in *Shewanella*. *Appl. Microbiol. Biot.* **93**, 41–48 (2012).

23. P. Chandrangsu, C. Rensing, J. D. Helmann, Metal homeostasis and resistance in bacteria. *Nat. Rev.*
Microbiol. **15**, 338–350 (2017).

24. H. Richter, M. Lanthier, K. P. Nevin, D. R. Lovley, Lack of Electricity Production by *Pelobacter*
carbinolicus Indicates that the Capacity for Fe(III) Oxide Reduction Does Not Necessarily Confer Electron
Transfer Ability to Fuel Cell Anodes. *Appl. Environ. Microb.* **73**, 5347–5353 (2007).

25. A. M. Schimpf, C. E. Gunthardt, J. D. Rinehart, J. M. Mayer, D. R. Gamelin, Controlling Carrier
Densities in Photochemically Reduced Colloidal ZnO Nanocrystals: Size Dependence and Role of the
Hole Quencher. *J. Am. Chem. Soc.* **135**, 16569–16577 (2013).

26. A. M. Schimpf, S. D. Lounis, E. L. Runnerstrom, D. J. Milliron, D. R. Gamelin, Redox Chemistries and
Plasmon Energies of Photodoped In_2O_3 and Sn-Doped In_2O_3 (ITO) Nanocrystals. *J. Am. Chem. Soc.* **137**,
518–524 (2015).

27. A. Agrawal, *et al.*, Rationalizing the Impact of Surface Depletion on Electrochemical Modulation of
Plasmon Resonance Absorption in Metal Oxide Nanocrystals. *ACS Photon.* **5**, 2044–2050 (2018).

28. A. Agrawal, *et al.*, Localized Surface Plasmon Resonance in Semiconductor Nanocrystals. *Chem.*
Rev. **118**, 3121–3207 (2018).

29. R. S. Lakes, Negative-Poisson's-Ratio Materials: Auxetic Solids. *Ann. Rev. Mater. Res.* **47**, 63–81
(2017).

30. A. Dong, *et al.*, A Generalized Ligand-Exchange Strategy Enabling Sequential Surface
Functionalization of Colloidal Nanocrystals. *J. Am. Chem. Soc.* **133**, 998–1006 (2011).

31. J. D. Roo, *et al.*, Unravelling the Surface Chemistry of Metal Oxide Nanocrystals, the Role of Acids
and Bases. *J. Am. Chem. Soc.* **136**, 9650–9657 (2014).

32. R. M. Dragoman, *et al.*, Surface-Engineered Cationic Nanocrystals Stable in Biological Buffers and
High Ionic Strength Solutions. *Chem. Mater.* **29**, 9416–9428 (2017).

33. S.-J. Yuan, *et al.*, A Photometric High-Throughput Method for Identification of Electrochemically
Active Bacteria Using a WO_3 Nanocluster Probe. *Sci. Rep.* **3**, 1315 (2013).

34. L. Su, *et al.*, Modifying Cytochrome c Maturation Can Increase the Bioelectronic Performance of
Engineered Escherichia coli. *ACS Synth. Biol.* **9**, 115–124 (2019).

35. R. J. Mendelsberg, *et al.*, Dispersible Plasmonic Doped Metal Oxide Nanocrystal Sensors that
Optically Track Redox Reactions in Aqueous Media with Single-Electron Sensitivity. *Adv. Opt. Mater.* **3**,
1293–1300 (2015).

36. Y. Cao, *et al.*, A Synthetic Plasmid Toolkit for *Shewanella oneidensis* MR-1. *Front. Microbiol.* **10**, 410
(2019).

37. S. L. Gibbs, *et al.*, Intrinsic Optical and Electronic Properties from Quantitative Analysis of Plasmonic
Semiconductor Nanocrystal Ensemble Optical Extinction (2020). https://arxiv.org/abs/1812.10142v2.

38. A. W. J. and J. E. Hutchison, Continuous Growth of Metal Oxide Nanocrystals: Enhanced Control of
Nanocrystal Size and Radial Dopant Distribution. *ACS Nano* **10**, 6942–6951 (2016).

39. J. T. Duong, *et al.*, Efficient polymer passivation of ligand-stripped nanocrystal surfaces. *J. Polym. Sci.*
A Polym. Chem. **50**, 3719–3727 (2012).

40. M. J. Bailey, R. van der Weegen, P. J. Klemm, S. L. Baker, B. A. Helms, Stealth Rare Earth Oxide
Nanodiscs for Magnetic Resonance Imaging. *Adv. Healthcare Mater.* **1**, 437–442 (2012).

41. A. Maho, *et al.* Aqueous Processing and Spray Deposition of Polymer-Wrapped Tin-Doped Indium
Oxide Nanocrystals as Electrochromic Thin Films. *Chem. Mater.*
(2020). doi:10.1021/acs.chemmater.0c02399

42. K. F. Palmer, D. Williams, Optical properties of water in the near infrared. *J. Opt. Soc. Am.* **64**, 1107
(1974).

43. Y. Zhang, *et al.*, Influence of spatial distribution of cationic functional groups at nanoparticle surfaces
on bacterial viability and membrane interactions. *J. Am. Chem. Soc.* **142**, 10814–10823 (2020).

44. G. Fan, A. J. Graham, J. Kolli, N. A. Lynd, B. K. Keitz, Aerobic radical polymerization mediated by
microbial metabolism. *Nat. Chem.* **12**, 638–646 (2020).

45. A. J. Graham, *et al.*, Genetic Control of Radical Cross-linking in a Semisynthetic Hydrogel. *ACS*
Biomater. Sci. Eng. **6**, 1375–1386 (2020).

46. K. A. Hunt, J. M. Flynn, B. Naranjo, I. D. Shikhare, J. A. Gralnick, Substrate-Level Phosphorylation is
the Primary Source of Energy Conservation during Anaerobic Respiration of *Shewanella oneidensis*
Strain MR-1. *J. of Bacteriol.* **192**, 3345–3351 (2010).

47. L. Binnenkade, L. Teichmann, K. M. Thormann, Iron Triggers λSo Prophage Induction and Release of
Extracellular DNA in *Shewanella oneidensis* MR-1 Biofilms. *Appl. Environ. Microb.* **80**, 5304–5316 (2014).

48. F. Zhang, J.-H. Wu, H.-Q. Yu, Probing Microbial Extracellular Respiration Ability Using Riboflavin.
Anal. Chem. **92**, 10606–10612 (2020).

49. A. Okamoto, Y. Tokunou, S. Kalathil, K. Hashimoto, Proton Transport in the Outer-Membrane
Flavocytochrome Complex Limits the Rate of Extracellular Electron Transport. *Angew. Chem.* **129**, 9210–
9214 (2017).

50. T. Fukushima, *et al.*, The Molecular Basis for Binding of an Electron Transfer Protein to a Metal Oxide
Surface. *J. Am. Chem. Soc.* **139**, 12647–12654 (2017).

51. X. Jing, *et al.* Outer membrane c-type cytochromes OmcA and MtrC play distinct roles in enhancing
the attachment of *Shewanella oneidensis* MR-1 cells to geothite. *Appl. Environ. Microbiol.* (2020).
doi:10.1128/AEM.01941-20.

52. D. Ito, S. Yokoyama, T. Zaikova, K. Masuko, J. E. Hutchison, Synthesis of Ligand-Stabilized Metal
Oxide Nanocrystals and Epitaxial Core/Shell Nanocrystals via a Lower-Temperature Esterification
Process. *ACS Nano* **8**, 64–75 (2014).

a

b

c

a

Extinction (OD)

S. oneidensis MR-1: + lactate
— pre-steady state EET
— steady state EET
— post-steady state EET

t=48 hr
t=0

(inset) t=8 hr, t=0

Wavenumber (cm⁻¹)

b

S. oneidensis MR-1:

S. oneidensis Δ*mtrC*Δ*omcA*Δ*mtrF*:

MtrC
OmcA   OmcA
MtrB
MtrA
CymA
e⁻

c

Electrons Transferred • Cell⁻¹

S. oneidensis MR-1:
+ lactate
– lactate

Δ*mtrC*Δ*omcA*Δ*mtrF*:
+ lactate

Time (h)

a *S. oneidensis* ΔmtrCΔomcAΔmtrF + mtrC (1000 μM IPTG)

b *S. oneidensis* ΔmtrCΔomcAΔmtrF + mtrC (100 μM IPTG)

c *S. oneidensis* ΔmtrCΔomcAΔmtrF + mtrC (10 μM IPTG)

d *S. oneidensis* ΔmtrCΔomcAΔmtrF + mtrC (0 μM IPTG)

e
S. oneidensis MR-1:
+ empty (1000 μM IPTG)

ΔmtrCΔomcAΔmtrF:
+ mtrC (1000 μM IPTG)
+ mtrC (100 μM IPTG)
+ mtrC (10 μM IPTG)
+ mtrC (0 μM IPTG)

f *S. oneidensis* ΔmtrCΔomcAΔmtrF + mtrC