1   **Evolution of Honey Resistance in Experimental Populations of Bacteria Depends on the Type of**

2   **Honey, and Has no Major Side Effects for Antibiotic Susceptibility**

4   <u>Running title</u>: Honey resistance evolution in *E. coli*

6   <u>Authors</u>: Anna M. Bischofberger[1], Katia R. Pfrunder Cardozo[1], Michael Baumgartner[1], Alex R. Hall[1]

8   <u>Contact information</u>: [1]Institute of Integrative Biology, ETH Zurich, Switzerland

9   anna.bischofberger@env.ethz.ch

10   katia.pfrunder@env.ethz.ch

11   michael.baumgartner@env.ethz.ch

12   alex.hall@env.ethz.ch

<u>Abstract</u>:

With rising antibiotic resistance, alternative treatments for communicable diseases are increasingly

relevant. One possible alternative for some types of infections is honey, used in wound care since before

2000 BCE and more recently in licensed, medical-grade products. However, it is unclear whether

medical application of honey results in the evolution of bacterial honey resistance, and whether this has

collateral effects on other bacterial traits such as antibiotic resistance. Here, we used single-step

screening assays and serial transfer at increasing concentrations to isolate honey-resistant mutants of

Escherichia coli. We only detected bacteria with consistently increased resistance to the honey they

evolved in with two of the four tested honey products, and the observed increases were small (maximum

two-fold increase in IC_{90}). Genomic sequencing and experiments with single-gene knockouts showed a

key mechanism by which bacteria increased their honey resistance was by mutating genes involved in

detoxifying methylglyoxal, which contributes to the antibacterial activity of *Leptospermum* honeys.

Crucially, we found no evidence that honey adaptation conferred cross-resistance or collateral

sensitivity against nine antibiotics from six different classes. These results reveal constraints on

bacterial adaptation to different types of honey, improving our ability to predict downstream

consequences of wider honey application in medicine.

Keywords:

honey, *Escherichia coli*, anti-bacterial agents, microbial drug resistance, bacteria, communicable

diseases, leptospermum

Acknowledgments:

A.B. thanks Richard Allen for help with statistical analysis. A.H. acknowledges Swiss National Science

Foundation project 31003A_165803.

Article type:　　Original research article

Introduction

Antimicrobial resistance is one of the biggest challenges facing global public health (WHO, 2018). To preserve the effectiveness of antibiotics and to treat infections caused by resistant bacteria, alternative approaches are required that can be used instead of antibiotics or after they have failed. One possible alternative currently being investigated for some applications is honey (Descottes, 2009; Knipping *et al.*, 2012; Vandamme *et al.*, 2013). Produced by the honey bee, *Apis mellifera*, honey has a long history in human medicine (Breasted, 1948) and has remained a staple treatment in traditional medicine. More recently, medically certified honeys and honey-containing products have been licensed in various part of the world, primarily for topical application, such as in wound healing (Molan and Betts, 2004; Cooke *et al.*, 2015), and treatment of otorhinolaryngological diseases (Werner and Laccourreye, 2011). The idea is appealing: honey is generally cheap and non-harmful to patients (Dunford and Hanano, 2004; Knottenbelt, 2014), inhibits bacterial growth (Basualdo *et al.*, 2007; Carter *et al.*, 2016) and can promote wound closure and healing (Molan, 1999). If honey could be used instead of antibiotics for some applications, this could contribute to managing antibiotic resistance. However, several open questions remain about whether bacteria exposed to inhibitory concentrations of honey evolve resistance to it, which genes or pathways are involved, and whether any such evolutionary responses have downstream effects on other properties relevant for treatment, in particular antibiotic resistance.

Despite open questions about how bacteria evolve honey resistance, the physiological causes of honey's antibacterial activity have been investigated in various species (Kwakman *et al.*, 2008, 2011; Badet and Quero, 2011; Camplin and Maddocks, 2014). Multiple mechanisms have been implicated, including high sugar/low water content, acidity, hydrogen peroxide and non-peroxide molecules such as methylglyoxal (White *et al.*, 1963; Allen *et al.*, 1991; Molan, 1992; Mavric *et al.*, 2008). At the phenotypic level, various effects of honey have been reported across different species, including changes in cell-wall integrity and cell shape (Henriques *et al.*, 2011; Brudzynski and Sjaarda, 2014; Wasfi *et al.*, 2016), quorum sensing (Truchado *et al.*, 2009; Lee *et al.*, 2011; Wang *et al.*, 2012), iron acquisition (Kronda *et al.*, 2013; Ankley *et al.*, 2020) and biofilm formation (Merckoll *et al.*, 2009; Badet and Quero, 2011; Halstead *et al.*, 2016). Perhaps unsurprisingly given this complex picture of inhibitory mechanisms, research to date on the evolution of honey resistance in bacteria has reached no

clear consensus. For example, Cooper *et al.* (2010) reported no stable increase in resistance after

exposure of various species to medical-grade *Leptospermum* honey *in vitro*, while Camplin and

Maddocks (2014) and Lu *et al* (2019) detected increased resistance in *Pseudomonas aeruginosa* cells

exposed to honey in biofilms *in vitro*. Thus, it remains unclear how rapidly bacteria exposed to honey

can evolve reduced susceptibility to it. Another important question is whether adaptation to honey has

side effects for antibiotic susceptibility. This is important for assessing the risk that honey application

could contribute to the spread of antibiotic resistance, affecting the success of antibiotic treatments used

in combination or later against the same bacteria. There is some evidence that honey resistance affects

susceptibility to rifampicin and imipenem in *P. aeruginosa* (Camplin and Maddocks, 2014). To

understand the general picture of how honey resistance affects antibiotic susceptibility, we need to test

a wider range of bacteria and antibiotics, and characterise the genetic pathways by which bacteria

become resistant.

We chose to study *E. coli* because it is a common pathogen in humans and animals (Dowd *et al.*,

2008; Suojala *et al.*, 2013), frequently associated with surface wounds that might be suitable for honey

treatment. Antibacterial resistance has increased in *E. coli* (Tadesse *et al.*, 2012; O'Neill, 2014), and

many of the resistance mechanisms found in *E. coli* can also be found in other species (Philippon *et al.*,

1989; Paulsen *et al.*, 1996), suggesting it as a good model for studying resistance evolution. To find out

how bacteria respond to honey exposure phenotypically and genotypically, we experimentally evolved

E. coli in the presence of four different honeys (two medical-grade honeys, two commercially available

honeys) by gradually exposing bacteria to increased honey concentrations during serial passage. We

then measured honey susceptibility of evolved bacteria from this experiment, as well as population

growth in the absence of honey. To identify genes involved in honey resistance, we used whole-genome

sequencing of these evolved isolates, as well as single-gene knockout variants. We also used a second

screen for honey-resistant mutants, by exposing many replicate populations to high honey

concentrations in a single-step (selective plating of overnight cultures). Lastly, we tested for collateral

effects of honey adaptation on antibiotic resistance by determining the phenotypic resistance of honey-

adapted isolates to antibiotics of different classes. Our results show that, even upon gradually increasing

exposure, large changes in honey resistance in *E. coli* populations growing *in vitro* are rare. However,

for some honey products we identified mechanisms driving moderate increases in honey resistance, and

we find no indication for cross-resistance between honey and antibiotics.

<u>Materials and Methods</u>

Organisms and Growth Conditions

We used *E. coli* K-12 MG1655 as parental strain for the evolution experiment and isolation of single-

step mutants. We used *E. coli* K-12 BW25113 and single-gene knockout variants derived from it (Keio

Knockout Collection (Baba *et al.*, 2006)) for knockout experiments. We stored all isolates in 25%

glycerol at -80°C. We performed routine culturing in lysogeny broth (LB) at 37°C with shaking at

180rpm.

Honeys and Antibiotics

The different honey products we used are listed in Table 1. We purchased SurgihoneyRO™ in 20g and

50g tubes from H&R Healthcare, Hull, UK; Medihoney™ Medical Honey in 20g tubes from Puras AG,

Bern, Switzerland, and in 50g tubes from The Honeydoctor, The Littledart Company Ltd, Tiverton,

UK; *MGO*™ Manuka Honey MGO™550+ in 250g jars from nu3 GmbH, Berlin, Germany; and

Fairtrade Liquid Blossom Honey in 550g jars from Coop, Basel, Switzerland. Honeys were stored in a

cool, dark place and, in the case of medical-grade honeys, opened tubes were only used as long as

recommended by the manufacturer. Non-medical-grade honeys were used for a maximum of six months

after opening. After plating honey samples on agar plates, we observed colony-forming units with the

commercial honey and Manuka honey. This was no longer the case after filtering honey solutions with

a Filtropur S 0.45 filter (Sarstedt, Germany) (Wasfi *et al.*, 2016). Accordingly, for all four honeys,

honey-containing growth media were prepared immediately before the start of each experiment by

diluting honey in LB (pre-heated to 55°C) and filter sterilizing. We purchased amoxicillin (product

number A8523), chloramphenicol (product number 23275), ciprofloxacin (product number 17850),

gentamicin (product number 48760), kanamycin (product number 60615), neomycin trisulfate salt

hydrate (product number N1876), polymyxin B (product number 5291), tobramycin (product number

T4014), and trimethoprim (product number 92131) from Sigma-Aldrich (Merck KGaA, Germany). We

prepared antibiotic stock solutions at the outset of the experiments and filter sterilized (Filtropur S 0.2

(Sarstedt, Germany)) and stored them according to the manufacturers' instructions (stock solutions:

amoxicillin 25mg/mL in sterile distilled water (dH$_2$O); chloramphenicol 50mg/mL in 70% ethanol;

ciprofloxacin 20mg/mL in 0.1M HCl; gentamicin 50mg/mL in dH$_2$O; kanamycin 40mg/mL in dH$_2$O;

neomycin 40mg/mL in dH$_2$O; polymyxin B 20mg/mL in dH$_2$O; tobramycin 40mg/mL in dH$_2$O;

trimethoprim 25mg/mL in DMSO).

Measuring Susceptibility of Isolates to Different Honeys

We used the 90% inhibitory concentration (IC$_{90}$) of each antibacterial compound as an indicator of

resistance. We defined the IC$_{90}$ as the lowest concentration tested above which bacterial growth did not

exceed 10% of growth of the same isolate in the absence of antibacterials. We also tried an alternative

definition, taking the IC$_{90}$ as the lowest concentration tested where growth did not exceed 10% of the

same isolate's growth in the absence of antibacterials (even if there was a score above this cut-off at

one of the higher concentrations, which we observed in a minority of dose-response curves); this did

not alter our overall conclusions. For each assay, we transferred independent LB-overnight cultures

(cultured in flat-based 96-well microplates (Sarstedt, Germany)) into microplates filled with various

honey concentrations and plain LB as a control, using a pin replicator (1/100 dilution, 2μL in 200μL).

We used slightly different concentration ranges in different sets of assays (as described below), which

were conducted with independent controls on different days. We incubated microplates overnight at

37°C and quantified bacterial growth by measuring optical density at 600nm (OD$_{600}$) with a microplate

reader (Infinite® 200 PRO, Tecan Trading AG, Switzerland) at the beginning and end of the experiment

(0h and 24h). We corrected OD scores for the optical density of the media. In each assay, we made 3-5

replicates of all tested strains in all tested compounds.

Experimental Evolution (Fig. S1A)

We serially passaged multiple selection lines of *E. coli* K-12 MG1655 in filtered solutions of each of

the four honey products and in the absence of honey for 22 days, transferring daily (4 honey treatments

+ 1 control treatment = 5 evolution environments). In summary: at each transfer, each selection line

was inoculated into multiple wells containing various honey concentrations. After overnight incubation,

we transferred from the well with the highest concentration supporting viable growth (Fig. S1A). In

more detail: to begin the experiment, we inoculated six selection lines in each evolution environment,

each with an independent colony (5 evolution environments x 6 colonies = 30 selection lines). In this

first step, we cultured each selection line over night in 5mL of LB at 37°C with shaking at 180rpm.

Then, for every selection line, we inoculated seven microplate wells filled with: 200μL of

unsupplemented LB ("rescue well"), LB supplemented with one of five concentrations of the respective

honey product, or honey stock solution (30 or 50% (w/v)). We incubated microplates overnight at 37°C

and quantified bacterial growth by measuring OD_{600} after 0h and 24h. For every selection line, we

transferred 5μL from one of the seven inoculated wells to seven wells in a new microplate. Each time,

we transferred from the well at the highest honey concentration where ΔOD_{600} (OD_{600} 24h – OD_{600} 0h)

> 0.1 (an arbitrary cut-off we took as an indication of viable growth). In cases where ΔOD_{600} in all

honey-supplemented wells was < 0.1, we used the rescue-well culture to inoculate the fresh microplate.

To gradually expose selection lines to higher concentrations of honey, we adjusted the range of

concentrations tested over time, according to the performance of individual selection lines

(concentrations and OD scores over time are given in Fig. S2. In the control treatment (no honey), we

used a single well for each selection line at each transfer. We did this for 22 days, freezing microplates

from days 3, 6, 9, 12, 15, 18, 21 and 22.

At the end of the experiment, we isolated a single colony isolate at random from each selection line.

We streaked out a sample from the well with the highest honey concentration where ΔOD_{600} > 0.1 onto

LB agar. After overnight incubation, we picked one colony per selection line, grew it overnight in 5mL

LB and stored it at -80°C. We used colony PCR (primer sequences: forward: 5′-AGA CGA CCA ATA

GCC GCT TT-3′; reverse: 5′-TTG ATG TTC CGC TGA CGT CT-3′) to ensure that all colony isolates were

E. coli K-12 MG1655. For five selection lines (Medihoney_01, Commercial_02, Commercial_03,

Commercial_04, Commercial_05), we found no colonies when streaking out samples at day 22. We

initially screened for honey-resistant phenotypes of colony isolates from the other selection lines by

culturing each colony isolate in honey-supplemented medium at a concentration in which the parental

strain was not able to grow. This led us to exclude five isolates that did not grow at honey concentrations

above that of the parental strain (Surgihoney_01, Surgihoney_02, Surgihoney_03, Surgihoney_06,

Commercial_01). We then proceeded with the remaining 14 serially-passaged putative honey-resistant

mutants, plus six control isolates serially passaged in LB, sequencing their genomes and quantifying

their honey susceptibility as described below/above respectively.

To test whether isolates from different evolution environments had different honey-resistance

profiles, we used a linear-mixed effects model (lmer function in R's lmerTest package (R version: 4.0;

package version: 3.1-2)), with evolution environment and assay compound as fixed effects and

genotype (isolate) as a random effect. We compared models with and without the interaction between

fixed effects, using the anova function of the stats package to test significance. When looking at

individual assay compounds, we used evolution environment as a fixed effect and genotype as a random

effect. We used maximum likelihood estimation ($REML = F$). Wild-type data was excluded from this

analysis; we tested for differences in IC_{90} between evolved isolates and the wild type separately, with

t-tests (*p*-values adjusted for multiple testing using the Holm-Bonferroni method).

Genetic Analysis of Serially-Passaged Mutants

We sequenced the 14 serially-passaged, putative resistant mutants and the six LB-adapted control

isolates. We used the QIAGEN Genomic-tip 20/G (Cat. No. 10223, Qiagen, the Netherlands) according

to the manufacturer's instructions for genomic DNA (gDNA) extraction. Overnight cultures inoculated

with single colonies were centrifuged at 5000xg at room temperature for 10min. After removal of the

supernatant, we stored cell pellets at -20°C until further processing. We quantified the obtained gDNA

using the Quant-iTTM dsDNA BR (Broad Range) Assay Kit (Thermo Fisher Scientific, USA) in the

QubitTM Fluorometer (Thermo Fisher Scientific). We used a Nanodrop (Thermo Fisher Scientific) to

control the purity of gDNA (ratios A_{260}/A_{280}). We sequenced at the Functional Genomic Center, Zurich,

Switzerland, using the Illumina Hiseq 4000 platform after library preparation with the Nextera XT DNA

Library Prep kit (Illumina, USA).

We trimmed and quality-filtered all sequences with trimmomatic (Bolger *et al.*, 2014). We mapped

the sequence of the ancestral strain of our resistance isolation experiment against the reference sequence

of *E. coli* K-12 MG1655 (NCBI accession number: U00096) using breseq 0.33.1 (Deatherage and

Barrick, 2014). We used gdtools implemented in breseq to integrate the identified mutations into the

reference genome. For variant calling, we mapped all sequences of the serially-passaged and single-

step putative resistant mutants against the refined reference genome using breseq. The sequencing data

has been deposited in the European Nucleotide Archive under the study accession number PRJEB35347

(https://www.ebi.ac.uk/ena).

Experiments with Single-Gene Knockout Variants

After identifying genes that potentially contribute to honey adaptation (Table S1), we tested for further

evidence of the role of these genes in honey resistance using single-gene knockout variants from the

Keio Knockout Collection (Baba *et al.*, 2006) for *E. coli* K-12. We tested for a change in the resistance

phenotype of these knockout variants relative to the ancestral strain of the knockout collection, *E. coli*

K-12 BW25113, using the same resistance phenotyping assay as described above. When choosing genes

to investigate, we concentrated on those (1) which were affected by independent mutations in at least

two selection lines, (2) where mutations were not detected in isolates from the control treatment, (3)

which were not annotated as "pseudogene", "intergenic" nor "non-coding", and (4) for which there is

an available knockout variant in the Keio Collection (limited to non-essential genes).

Measuring Population Growth of Serially-Passaged Mutants in the Absence of Honey

We tested whether our serially passaged isolates showed altered population growth in the absence of

honey relative to the ancestral strain. To do this, we grew four independent cultures (each inoculated

from a different colony) of each serially-passaged isolate (including control-evolved isolates, $n = 20$)

and the ancestral isolate, each in 150μL LB in a microplate in a randomized layout. After overnight

incubation, we used a pin replicator to inoculate a fresh microplate (all wells filled with 150μL LB).

We then measured OD_{600} every 15min for 24h (shaking before each measurement). We used R's nls

and SSlogis functions in the stats package to estimate the growth rate and yield for each culture, with

yield corresponding to the *Asym* parameter of the models (= asymptote) and growth rate corresponding

to the inverse of the *scal* parameter (= numeric scale parameter on input axis). We tested for a difference

in growth parameters between each evolved isolate and the parental strain with a *t*-test, adjusting *p*-

values using sequential Bonferroni correction. We also tested for an average difference in growth rate

or yield among evolved isolates from different evolution environments. We did this using a linear

mixed-effects model, with evolution environment as a fixed effect and genotype as random effect, and

excluding data from the wild type. We used maximum likelihood estimation (*REML = F*) with the lmer

function in the lmerTest package.

Measuring Susceptibility of Serially-Passaged Mutants to Antibiotics

We measured the phenotypic resistance (IC_{90}) of the 20 serially-passaged isolates (14 putative resistant

mutants, six LB-adapted control isolates) and of the parental strain (*E. coli* K-12 MG1655) for nine

antibiotics representing six different classes: amoxicillin (penicillin), ciprofloxacin (fluoroquinolone),

chloramphenicol (chloramphenicol), gentamicin (aminoglycoside), kanamycin (aminoglycoside),

neomycin (aminoglycoside), polymyxin B (polymyxin), tobramycin (aminoglycoside), and

trimethoprim (dihydrofolate reductase inhibitor). We decided to test several aminoglycoside drugs

(gentamicin, kanamycin, neomycin, tobramycin) because previous studies have found that bacteria

exposed to honey reduce the expression of genes involved in the TCA cycle (Lee *et al.*, 2011; Jenkins

et al., 2014), while others report a link between aminoglycoside susceptibility and defects or down-

regulated gene expression in the bacterial respiratory chain, including the TCA cycle (Magnet and

Blanchard, 2005; Chittezham Thomas *et al.*, 2013; Shan *et al.*, 2015; Su *et al.*, 2018; Zhou *et al.*, 2019).

We conducted the assays using a similar protocol as described above for honey. In brief, we first

incubated independent replicate populations of each isolate in a randomized layout in microplates

overnight. From these microplates we inoculated assay plates filled with LB supplemented with

antibiotics at various concentrations in two-fold broth dilution series, measuring bacterial growth by

the change in OD_{600} (0h, 24h) as described above.

Single-Step Isolation of Honey-Resistant Mutants (Fig. S1B)

As a second screen for mutants of *E. coli* K12 MG1655 with increased honey-resistance, we plated

aliquots of multiple independent overnight cultures, grown in the absence of antibiotics, on agar

supplemented with each honey product.

For each honey type, we first grew 54 independent overnight cultures (250μL per culture in LB in a

96-well microplate; Fig. S1B). We then plated each culture onto a separate agar plate (prepared in 6-

well culture plates filled with 4mL honey-supplemented agar per well (Sarstedt, Germany)), plating

100μL of 18 separate cultures at each of three concentrations per honey product. We chose

concentrations 1.25-, 1.5-, 2- or 3-times higher than previously determined IC_{90}s of the wild-type strain

in liquid. We then incubated the plates overnight at 37°C before checking for resistant mutants. For

SurgihoneyRO™, Medihoney™ and Manuka honey, we picked six putative honey-resistant colonies,

each from a separate well. When isolating colonies, we prioritized those from wells with higher

concentrations of honey. For commercial honey, we observed no viable colonies on honey-

supplemented agar despite four attempts on different days (total of 216 cultures). We cultured the

selected putative resistant colonies overnight in 5mL LB and suspended them in 25% glycerol for

storage at -80°C. We then tested these 18 single-step putative honey-resistant mutants for phenotypic

resistance as described above. We also used a second test to see if putative honey resistance phenotypes

were robust, by streaking out frozen stocks of each colony isolate on honey-supplemented agar at a

concentration inhibitory to the parental strain. Only three of 18 single-step putative honey-resistant

mutants were able to form colonies under these conditions (Medihoney_10, Surgihoney_09,

Surgihoney_10).

283
284   **<u>Results</u>**
285

Experimental Evolution of Manuka Honey-Resistant Bacteria by Serial Passaging

During serial passage at gradually increasing honey concentrations (Fig. S1A), most selection lines

reached higher population densities at concentrations that were initially inhibitory to the ancestral strain

(Fig. S2 & S3). Of 14 putative honey-resistant mutants and six control isolates, each isolated from a

different selection line after 22 days, several showed increased resistance compared to the ancestral

strain against one or more honey products (Fig. 1 & S4). The average change in resistance varied among

evolved isolates passaged with different honey compounds (effect of evolution environment: $F_{4, 20}$ =

1.799, p > 0.1). These differences also depended on which honey compound was used in the assay

(evolution environment × assay environment interaction: $\chi^2(12)$ = 234.99, p < 0.001). We observed the

largest change in resistance for mutants selected with Manuka honey or Medihoney™ when they were

assayed with Manuka honey (mean change relative to the ancestor of 2.13-fold (s.d. = 0.22) and 2.07-

fold, respectively). Manuka honey resistance of Manuka honey- and Medihoney™-evolved isolates was

also significantly higher than for control-evolved isolates from the honey-free LB-medium treatment

($t_{9.7}$ = 7.73, p < 0.001 and $t_{8.9}$ = 6.77, p < 0.001, respectively). On average, Manuka honey-evolved and

Medihoney™-evolved isolates also had moderately increased resistance to Medihoney™ (mean change

relative to the ancestor of 1.15-fold (s.d. = ±0.16) and 1.17-fold (s.d. = ±0.1), respectively), but this was

not significantly different from that of control-evolved isolates on average (Manuka: t_2 = -0.30, p >

0.05; Medihoney™: t_2 = 0, p > 0.05). Some other individual isolates had consistently increased Manuka

honey or SurgihoneyRO™ resistance (all replicates for a given isolate higher than all replicates for the

ancestor; Fig. 1 & S4), but these changes were small compared to those for Manuka honey- and

Medihoney™-evolved isolates tested with Manuka honey. Thus, after experimental evolution we

observed the strongest evidence of honey adaptation with Manuka honey.

Honey Adaptation Is Linked to Mutations in *nemAR* and *clpP*

Genomic sequencing of our serially passaged isolates revealed changes at several loci (Fig. 2, Table

S1). Some loci were mutated multiple times independently in honey-evolved colony isolates, but not in

control-evolved isolates (Table S1, Fig. 2), indicating a possible role in adaptation to honey. Two such

genes were *clpP* (serine protease, mutated in two Manuka honey-evolved isolates) and *nemR* (DNA-

binding transcriptional repressor, mutated in four Manuka honey-evolved isolates and three

Medihoney™-evolved isolates; Table S1, Fig. 2). The intergenic region between *ydhL* and *nemR* was

mutated in the remaining two Manuka honey-adapted isolates and the remaining two Medihoney™-

adapted isolates (Table S1, Fig. 2). An additional gene for which we found mutations in multiple isolates

serially passaged in Manuka honey is *yafS* (methyltransferase; Table S1, Fig. 2), located next to *gloB*,

the gene encoding glyoxalase II, an enzyme involved in detoxification of reactive aldehydes (Ozyamak

et al., 2010). In control-evolved isolates, we found parallel mutations in *fimE*, a recombinase

responsible for on-to-off switching of type 1 fimbriae expression, and *flhD*, one of two transcriptional

activators of the *E. coli* flagellar regulon but also involved in cell division (Klemm, 1986; Iino *et al.*,

1988; Liu and Matsumura, 1994; Prüß and Matsumura, 1996), consistent with past work with LB-

adapted isolates (Knöppel *et al.*, 2018). In summary, we found parallelisms in both control-evolved and

honey-evolved isolates, and in particular every Manuka honey- and Medihoney™-evolved isolate had

a mutation in or affecting *nemR*. As we discuss below, this may reflect the role of *nemR* in

methylglyoxal degradation (Ozyamak *et al.*, 2013), an active component of *Leptospermum* honeys.

Single-Gene Knockouts Support a Role for *nemR* in Honey Resistance

We used single-gene knockout variants of several genes that were mutated in serially-passaged isolates

to test for further evidence that they play a role in honey resistance (Fig. 3 & S5). The knockout variant

Δ*nemR* had increased Manuka honey resistance (all replicates higher than all replicates of the wild-type

strain). This variant also showed increased average resistance to Medihoney™, although this was not

consistent across all replicate assays (Fig. 3 & S5). This is consistent with the above finding that all

isolates serially passaged in Manuka honey or Medihoney™ had mutations in or close to *nemR*, and all

were more resistant to Manuka honey and Medihoney™ than the ancestral strain. In Δ*clpP*, the

knockout of the other gene for which we observed independent mutations in multiple honey-evolved

isolates, we observed a similar pattern: increased resistance to Manuka honey (in all replicate assays)

and higher average Medihoney™ resistance (two out of three replicates higher than wild-type). This is

consistent with the increased resistance for these honeys we observed with the two Manuka honey-

selected isolates with mutations in *clpP* (Fig. 2 & S4). Two other knockout variants, Δ*nemA* and Δ*eda*,

had consistently altered resistance to Manuka honey, with Δ*eda* having a higher susceptibility than the

ancestral strain. In summary, independent deletion of two genes, *nemR* and *clpP,* which were also

directly or indirectly affected by mutations in several of our serially-passaged isolates, conferred

increased resistance to Manuka honey and Medihoney™.

Impaired Growth in the Absence of Honey for Honey-Evolved Compared With Control-Evolved Isolates

Most serially passaged isolates (both honey-evolved and control-evolved) had increased population growth rates relative to the ancestral strain in the absence of honey (Fig. 6). However, control-evolved isolates had higher growth rates on average compared to all four types of honey-evolved isolates (linear mixed-effects model: Manuka honey: t_{20} = -4.355, p < 0.001; Medihoney™: t_{20} = -5.538, p < 0.001; SurgihoneyRO™: t_{20} = -4.078, p < 0.001; commercial honey: t_{20} = -4.671, p < 0.001). By contrast, we found little effect of serial passaging on growth yield in evolved isolates compared to the ancestral strain (Fig. S6). Comparing the growth yield in different evolution environments, we found no significant difference between isolates serially passaged in LB and honey-evolved isolates (linear mixed-effects model: all p > 0.05; Fig. S6). In summary, for most experimentally evolved isolates we found a positive effect of serial passage on growth rate, but honey-adapted isolates had a lower growth rate on average compared to isolates serially passaged in the absence of honey.

No Evidence for Cross-Resistance Between Honey and Antibiotics

We determined the phenotypic resistance of our serially-passaged isolates and the ancestral strain to nine antibiotics of six different classes (Fig. 5; Fig. S7). We found only a single case out of 180 combinations (20 serially-passaged isolates × 9 antibiotics) where antibiotic resistance was consistently altered compared to the ancestral strain (higher/lower IC_{90} for all mutant replicates compared to all replicates of ancestor). This was for one Manuka honey-evolved isolate, Manuka_05, tested with polymyxin B. For seven of the other eight antibiotics, the largest difference shown by individual replicates relative to the median of the ancestral replicates was two-fold or less. For the remaining antibiotic, kanamycin, some individual replicates had differences of four-fold compared to the ancestor. The overall lack of major differences between evolved isolates and the ancestral strain was supported by Analysis of Variance (testing each evolved-isolate-versus-wild-type combination separately, with genotype (evolved vs wild type) and assay compound (antibiotic) as factors, including the interaction term and with *p*-values adjusted using the Holm-Bonferroni method); this indicated a single significant

effect of genotype, for isolate Manuka_06). Thus, we found no evidence for appreciable changes in

sensitivity to antibiotics in serially-passaged honey-adapted isolates.

Single-Step Screening Supports Similar Variation of Resistance Evolution Among Honey

Products

We tested whether our finding that resistance evolution was more pervasive with Manuka honey than

the other honey compounds also holds true when bacteria are exposed to inhibitory honey

concentrations suddenly, rather than gradually. We did this by plating aliquots from multiple

independent, replicate overnight cultures directly onto honey-supplemented agar, and picking the

resulting colony isolates that showed a putative resistance phenotype. As for our serially passaged

isolates above, we observed the largest increases in resistance for colony isolates from the Manuka-

honey treatment, when assayed with Manuka honey (Fig. S8). For SurgihoneyRO™ and Medihoney™,

we observed colonies on some of our plates, but after picking six colonies for each compound and

testing their phenotypic resistance, none of them had consistently altered resistance to the honey they

were selected with. When we streaked out frozen stocks of these 12 putative honey-resistant colony

isolate on honey-supplemented agar at a concentration inhibitory to the parental strain, only 3 out of 12

isolates formed colonies (Medihoney_10, Surgihoney_09, Surgihoney_10). Thus, most putative

SurgihoneyRO™- and Medihoney™-resistant mutants showed unstable phenotypic resistance across

multiple rounds of isolation, culturing and restreaking (we discuss this further in the context of earlier,

similar observations below). For commercial honey, we obtained no viable colonies despite repeating

the assay an additional three times on different days (216 aliquots in total for commercial honey). In

summary, we isolated single-step resistant mutants with Manuka honey, but found no colony isolates

with robust resistance phenotypes to SurgihoneyRO™, Medihoney™ or commercial honey, consistent

with our finding in serial passage that resistance evolved most readily against Manuka honey.

DISCUSSION

We found *E. coli* does not easily develop large increases in honey resistance upon serial passage at

gradually increasing concentrations, or upon plating of many replicate populations at inhibitory

concentrations. However, we identified a set of genes important in adaptation to honey, which conferred

moderate increases in resistance to Manuka honey. The known physiological role of this genetic

pathway helps to explain why these genetic changes, and associated increases in phenotypic resistance,

were specific to *Leptospermum* honeys (detoxification of methylglyoxal, which contributes to

antibacterial activity of these honeys). This indicates the likelihood of honey resistance upon medical

application may depend critically on the type or combination of honey products used. Furthermore, we

found that honey adaptation *in vitro* had only minimal side-effects for antibiotic resistance.

The first important implication of our results is that *E. coli* does not readily become resistant to

honey, which is promising in the context of expanding medical use of honey. Previous studies also had

difficulty in isolating mutants with stable honey resistance, and/or found that honey resistance

phenotypes revert quickly in the absence of honey (Cooper *et al.*, 2010). This is consistent with our

observation that several putative resistant mutants from our single-step screen had unstable phenotypes,

and some serially passaged populations that attained the ability to grow at increased honey

concentrations subsequently lost this ability upon transfer to fresh medium (referred to by (Abdel-Azim

et al., 2019) as *second transfer crash* (STC)). One possible explanation is that non-genetic changes such

as persister formation or other morphological changes enable temporary population survival or growth

in otherwise inhibitory concentrations. Consistent with honey inducing physiological responses that are

not necessarily heritable, Brudzynski and Sjaarda (2014) observed morphological changes in *E. coli*

cultures exposed to honey, including changes in cell shape / filamentation.

The second key implication of our results is that, to our knowledge, this is the first report of a specific

genetic mechanism linked to decreased honey susceptibility, namely changes affecting the *nemAR*

operon, and alternatively via changes affecting *clpP*. In support, *nemR* and its operon have previously

been described as being involved in physiological processes that we can expect to be beneficial during

honey exposure. Manuka honey and Medihoney™ are both *Leptospermum* honeys, whose high non-

peroxide antimicrobial activity is attributed to their relatively high content of methylglyoxal (Allen *et*

al., 1991; Mavric *et al.*, 2008), a small amount of which bacteria produce intracellularly (Tötemeyer *et*

al., 1998). Ozyamak *et al.* (2013) linked the *nemAR* operon to methylglyoxal detoxification. The *nemAR*

operon is located upstream from the glyoxalase system (Ozyamak *et al.*, 2013) which consists of GlxI

and GlxII (glyoxalase I and II, encoded by *gloA* and *gloB* respectively), the two most important enzymes

in the detoxification pathway of methylglyoxal (Clugston *et al.*, 1998; MacLean *et al.*, 1998;

Mannervik, 2008). Deletion of *nemR*, the repressor gene, results in increased transcription both of *nemA*

and *gloA* (Ozyamak *et al.*, 2013). Thus, the role we find for *nemAR* in manuka honey resistance is

consistent with its known role in methylglyoal detoxification. Similarly, *clpP*, another gene we found

independently mutated in multiple Manuka honey-adapted isolates, was shown by Jenkins *et al.* (2014)

to be overexpressed in *Leptospermum* honey-exposed methicillin-resistant *S. aureus*, supporting a role

for this gene in honey resistance in multiple species.

That methylglyoxal detoxification plays a key role in increased honey resistance is further supported

by the variation we observed among honey compounds. Robust resistant mutants emerged in our single-

step screen only with a *Leptospermum* honey (Manuka) with a high concentration of methylglyoxal,

but not with the other three compounds. During serial passage we saw a similar trend: the most

consistent increases in resistance were with the two *Leptospermum* honey products. By contrast,

resistance of putative single-step resistant mutants isolated from SurgihoneyRO™-supplemented agar

was not stable, and we could not isolate any spontaneous mutants for the commercial honey we used.

This commercial honey is a blend of honeys with unknown flower sources from different South-

American countries. Drawing a parallel to antibiotics, where combinations of antibiotics can make it

harder for bacteria to evolve resistance (Baym *et al.*, 2016), we speculate that resistance evolution

against this honey was rare in our experiments because it is a form of combination treatment. This raises

the possibility that effective application of honey in treatment may benefit from combining multiple

types of honey products with different mechanisms of action. We note that despite variation among

honey products, all isolates in our experiment were inhibited by honey concentrations comparable to

those found in medical-grade honey products (63–100% in products recommended/licensed for medical

wound care, http://www.medihoney.de/index.html). We hypothesize that the multi-faceted nature of

honey's antibacterial activity (Wang *et al.*, 2012; Nolan *et al.*, 2019) contributes to the difficulty of

bacteria evolving resistance to such high concentrations.

The third major implication of our results is that honey adaptation did not come with collateral

effects on antibiotic resistance. This is a second aspect that is promising in the context of wider medical

application of honey, and contrasts starkly with the frequent occurrence of cross-resistances between

antibiotics (Szybalski and Bryson, 1952; Gutmann *et al.*, 1985) and between antibiotics and some other

antimicrobial agents (Loughlin *et al.*, 2002; Braoudaki and Hilton, 2004; Baker-Austin *et al.*, 2006;

Allen *et al.*, 2017; Bischofberger *et al.*, 2020). We do not exclude that more significant changes in

antibiotic resistance might be observed in mutants or strains with larger changes in honey resistance

than the moderate increases we observed here, if such strains exist. A key avenue for future work is

therefore to determine whether any genetic variation in natural populations associated with variable

honey resistance is independent of variation in antibiotic resistance, as in our experiment. Similarly,

because honey has shown effective inhibition of a wide range of pathogens (Carter *et al.*, 2016; Hillitt

et al., 2017; Yabes *et al.*, 2017), and the principal genes involved in honey resistance in our experiment

are conserved in multiple pathogenic species (e.g., *nemR* or analogues are present in *Klebsiella*

pneumonia, Acinetobacter baumannii and *Salmonella typhimurium*), our work identifies candidate loci

that may be involved in resistance in other species. This is another relevant area for future research in

the context of medical honey application.

Our results also provided indirect evidence of growth costs associated with honey adaptation.

Growth in the absence of selecting antibacterials is widely considered an important parameter in the

long-term spread of resistance (Andersson and Hughes, 2010). The larger increase in growth rate

observed in our control-evolved isolates compared to honey-adapted isolates suggests either (i) that

honey-adaptation constrained adaptation to other aspects of the environment, preventing acquisition of

other beneficial mutations, or (ii) that honey-resistance mutations combined with other types of

beneficial mutations conferred a smaller net increase in growth rate in honey-free growth medium

compared to other beneficial mutations alone. Our sequence data are more consistent with the former:

control-evolved populations had parallel mutations in flagella and fimbriae genes (*flhD* and *fimE*),

consistent with past work with LB-adapted *E. coli* (Knöppel *et al.*, 2018). These mutations were much

less common in honey-adapted colony isolates (Fig. 2). This could be due, for example, to epistatic

interactions between the different types of adaptive mutations (Scanlan *et al.*, 2015). A full

understanding of the growth costs and benefits associated with individual honey-resistance alleles and

their interactions with other types of beneficial mutations is beyond the scope of this paper, and was

not our aim here, but would make an interesting avenue for future work.

In conclusion, honey resistance in our experiment only evolved with a subset of the compounds we

tested, and only to a moderate degree. This is promising in the context of medical application of honey.

A further positive aspect is that we found no evidence of strong downstream effects on antibiotic

susceptibility in isolates adapted to honey *in vitro* via chromosomal mutation. Finally, we identified

putative genetic mechanisms involved in honey adaptation, via changes affecting genes involved in

detoxifying methylglyoxal, making this mechanism most relevant for *Leptospermum* honeys such as

Manuka honey.

Tables

Table 1: Honey products used in this study.

Product name	Product description	Initial IC$_{90}$ for *E. coli* K-12 MG1655	Manufacturer
SurgihoneyRO™	medical honey; sterile, bio-engineered; undiluted; poly-origin (unspecified)	5%(w/v)	H&R Healthcare, Hull, UK
Medihoney™ Medical Honey	medical honey; sterile, undiluted; *Leptospermum* honey (methylglyoxal content unspecified)	16%(w/v)	sorbion austria, Zwölfaxing, Austria
MGO™ Manuka Honey MGO™550+	consumer product; natural, undiluted; *Leptospermum* honey (methylglyoxal content of 550 mg/kg)	10%(w/v)	Manuka Health New Zealand Ltd, Te Awamutu, New Zealand
Commercial honey (Fairtrade Liquid Blossom Honey)	consumer product; natural, undiluted; poly-origin (Chili, Guatemala, (Mexico))	10%(w/v)	Coop, Basel, Switzerland

Supplementary Table 1. Mutations involved in adaptation to serial passaging in different honey

products and in LB. The table contains all mutations registered in serially-passaged isolates (14

isolates adapted to four different honeys, six control isolates) using the Illumina Hiseq 4000 platform.

Gene description according to breseq and the reference sequence of the ancestral isolate, E. coli K-12

MG1655. (*see .xslx document "tableS1_Bischofbergeretal.xslx"*)

Figures

Figure 1.

Figure 1. Susceptibility of serially-passaged, putative resistant mutants with four different honey

compounds. Each cell gives the median IC_{90} for a given isolate (the ancestral strain K-12 MG1655, in

the top row, or a putative resistant mutant serially passaged with one of the four honey compounds,

labelled according to compound and replicate selection line) assayed with one of the four honey

compounds (columns). Each value is the median of four independent replicates; combinations where

all replicates of a putative resistant mutant were higher than all replicates of the ancestor in the same

treatment are indicated with a "+". Individual replicates for each strain are shown in Fig. S4.

Figure 2.

Figure 2. Genetic changes in 20 serially-passaged isolates of *E. coli* K-12 MG1655. The outermost

ring represents the ancestral isolate with coloured tiles for all genes in which we found mutations in

serially-passaged isolates. Each inner circle represents one serially-passaged isolate: in turquoise:

control treatment; in grey: four different honey treatments. Genetic mutations are indicated with

coloured tiles: red: mutations in one gene; yellow: intergenetic mutations; blue: deletions >1000bp.

Figure 3.

Figure 3. Honey susceptibility of single-gene deletion variants. Each cell shows the median IC_{90}

towards four different honeys (columns) for the ancestral strain *E. coli* K-12 BW25113 (top row) or

one of 28 single-gene knockout variants (other rows). Each cell is the median value of three independent

replicates; combinations where all replicates of a putative resistant mutant were higher/lower than all

replicates of the ancestor in the same treatment are indicated with a "+"/"-". Individual replicates for

each strain are shown in Fig. S5.

Figure 4.

Figure 4. Growth rate of experimentally evolved isolates in the absence of honey. Population

growth rate (*y*-axis) is shown for the wild type (MG1655), six mutants serially passaged in honey-free

LB (LB_01-LB_06), and 14 mutants serially passaged in different honeys (labelled according to honey

and replicate; *x*-axis). Each black line gives the median of four replicates (replicates shown as dots).

Figure 5.

Figure 5. Resistance of serially-passaged, putative honey-resistant mutants against nine different

antibiotics, relative to the wild type. The matrix shows the median IC_{90} towards nine antibiotics of 20

serially-passaged isolates relative to the median IC_{90} of the parental strain, *E. coli* K-12 MG1655.

Blue/dark seagreen signify susceptibility, yellow/light green resistance relative to the parental strain;

combinations where all replicates of a given isolate were lower than all replicates of the ancestor in the

same treatment are indicated with a "-". Individual replicates for each strain are shown in Fig. S7.

 Supplementary Figures

A

use o/n culture to inoculate:
- 6 wells with increasing C_{honey}
- 1 well with control

determine transfer culture (O) with OD measurements

pick independent colony

incubate o/n at 37°C

incubate o/n at 37°C

repeat 22 times

inoculate new microplate with transfer culture (O); C_{honey} adjusted

after 22d: plate transfer cultures (O) on honey-supplemented agar plates; isolate resistant colonies

B

pick independent colony

incubate 2h at 37°C

use culture to inoculate 54 LB-filled wells (1:10 dilution)

incubate o/n at 37°C

plate cultures on honey-supplemented agar plates; isolate resistant colonies

 Supplementary Figure 1.

Supplementary Figure 1. Schematic protocol of experimental resistance evolution. (A) Serial passage: For each replicate selection line, an independent overnight culture was used to inoculate a range of honey concentrations and honey-free medium (LB). The highest honey concentration that showed viable growth ((OD_{600} 24h – OD_{600} 0h) > 0.1, transfer culture (outlined in red)) was used to inoculate wells filled with various concentrations in a new microplate. Control treatment was performed by daily passaging in honey-free medium (LB). This was repeated daily for 22 days. (B) Single-step screening: for each compound we made multiple overnight cultures before plating on honey-supplemented agar and screening for resistant mutants.

Supplementary Figure 2.

Supplementary Figure 2. Population densities at different time points and honey concentrations

for 24 selection lines passaged with different honeys for 22 days. Each row of panels (A-D) shows

population density, estimated by OD_{600} (*y*-axis) of six selection lines passaged with one of the four

honey compounds. The *x*-axis shows the concentration of the respective honey in % (w/v), and each

line gives the observed population densities across all tested concentrations on a given day (different

days/lines have different colours, going from blue to green to yellow to red over earlier/later time

points).

Supplementary Figure 3.

Supplementary Figure 3. Highest honey concentration each selection line was able to grow at

during 22d of serial passage in honey-supplemented medium. For each honey compound, we

passaged six selection lines by daily transfer into increasing concentrations of honey. Each point

represents the highest concentration for which an individual selection line showed $\Delta OD_{600} > 0.1$ after

24h incubation at any time during the experiment (points are jittered along the axes). The red line

represents the initial IC_{90} of the wild type strain, *E. coli* K-12 MG1655, for each honey.

Supplementary Figure 4.

Supplementary Figure 4. Resistance of serially-passaged putative resistant mutants to four

different honeys. Each of the four panels shows the IC_{90} of 14 honey-adapted (and six control-adapted)

isolates and of the parental strain *E. coli* K-12 MG1655 to four honey products after 24h incubation.

The black lines represent the median of four independent replicates per serially-passaged isolate,

replicates are shown as dots (jittered along the *y*-axis). The red horizontal line represents the median

IC_{90} of four independent replicates of the parental strain.

Supplementary Figure 5.

Supplementary Figure 5. Honey resistance of single-gene deletion variants. Each panel shows the

IC_{90} of 28 single-gene knockout variants from the Keio Knockout Collection and of the parental strain

E. coli K-12 BW25113 after 24h incubation. Each black line is the median of three replicates per

knockout variant; replicates are shown as dots (jittered along the *y*-axis). The dashed red line represents

the median IC_{90} of three replicates of the wild-type strain, *E. coli* K-12 BW25133.

Supplementary Figure 6.

Supplementary Figure 6. Growth yield of serially passaged, putative honey-resistant isolates in

honey-free growth medium (LB). Growth yield (*y*-axis) is shown for the wild type (MG1655), six

isolates serially passaged in LB (LB_01-LB_06), and 14 isolates serially passaged in different honeys

(labelled according to honey and replicate) (*x*-axis). Each black line represents the median of four

replicates (points).

Supplementary Figure 7.

Supplementary Figure 7. No evidence for cross-resistance between four honeys and nine

antibiotics. Each panel shows the IC_{90} of 20 isolates that were serially-passaged in honey or LB relative

to the IC_{90} of the wild type (*E. coli* K-12 MG1655; individual replicates for the wild type are also shown

relative to the median for the wild type, which is the red horizontal line in each panel) towards one

antibiotic. Each black line is the median of four independent replicates per isolates; replicates are shown

as dots (jittered along the *y*-axis).

Supplementary Figure 8.

Supplementary Figure 8. IC_{90} of 18 single-step putative resistant mutants to four different honeys.

Each of the four panels shows the IC_{90} values of 18 single-step mutants and of the parental strain *E. coli*

K-12 MG1655 to four different honeys. The black lines represent the median of five independent

replicates per single-step putative resistant mutant, replicates are shown as dots (jittered along the *y*-

axis). The red horizontal line represents the median IC_{90} of five independent replicates of the wild-type

strain.

Acknowledgements

See title page.

Data Archiving Statement

The data that support the findings of this study will be made openly available at Dryad Digital

Repository: to be completed after manuscript is accepted for publication.

References

Abdel-Azim, S.G., Abdel-Azim, A.G., Piasecki, B.P., and Abdel-Azim, G.A. (2019) Characterization of the Gain and Loss of Resistance to Antibiotics versus Tolerance to Honey as an Antimutagenic and Antimicrobial Medium in Extended-Time Serial Transfer Experiments. *Pharmacognosy Res* **11**: 147–54.

Allen, K.L., Molan, P.C., and Reid, G.M. (1991) A Survey of the Antibacterial Activity of Some New Zealand Honeys. *J Pharm Pharmacol* **43**: 817–822.

Allen, R.C., Pfrunder-Cardozo, K.R., Meinel, D., Egli, A., and Hall, A.R. (2017) Associations among antibiotic and phage resistance phenotypes in natural and clinical Escherichia coli isolates. *MBio* **8**: e01341-17.

Andersson, D.I. and Hughes, D. (2010) Antibiotic resistance and its cost: Is it possible to reverse resistance? *Nat Rev Microbiol* **8**: 260–271.

Ankley, L.M., Monteiro, M.P., Camp, K.M., O'Quinn, R., and Castillo, A.R. (2020) Manuka honey chelates iron and impacts iron regulation in key bacterial pathogens. *J Appl Microbiol* **128**: 1015–1024.

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., et al. (2006) Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: The Keio collection. *Mol Syst Biol* **2**:.

Badet, C. and Quero, F. (2011) The in vitro effect of manuka honeys on growth and adherence of oral bacteria. *Anaerobe* **17**: 19–22.

Baker-Austin, C., Wright, M.S., Stepanauskas, R., and McArthur, J. V. (2006) Co-selection of antibiotic and metal resistance. *Trends Microbiol* **14**: 176–182.

Basualdo, C., Sgroy, V., Finola, M.S., and Marioli, J.M. (2007) Comparison of the antibacterial activity of honey from different provenance against bacteria usually isolated from skin wounds. *Vet Microbiol* **124**: 375–381.

Baym, M., Stone, L.K., and Kishony, R. (2016) Multidrug evolutionary strategies to reverse antibiotic resistance. *Science (80-)* **351**:.

Bischofberger, A.M., Baumgartner, M., Pfrunder-Cardozo, K.R., Allen, R.C., and Hall, A.R. (2020) Associations between sensitivity to antibiotics, disinfectants and heavy metals in natural, clinical and laboratory isolates of Escherichia coli. *Environ Microbiol* **00**: 1–16.

Bolger, A.M., Lohse, M., and Usadel, B. (2014) Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.

Braoudaki, M. and Hilton, A.C. (2004) Low level of cross-resistance between triclosan and antibiotics in Escherichia coli K-12 and E. coli O55 compared to E. coli O157. *FEMS Microbiol Lett* **235**: 305–309.

Breasted, J.H. (1948) The Edwin Smith Surgical Papyrus, Breasted,J.H. (ed) Chicago: The University of Chicago Press.

Brudzynski, K. and Sjaarda, C. (2014) Antibacterial compounds of canadian honeys target bacterial cell wall inducing phenotype changes, growth inhibition and cell lysis that resemble action of β-lactam antibiotics. *PLoS One* **9**: 106967.

Camplin, A.L. and Maddocks, S.E. (2014) Manuka honey treatment of biofilms of Pseudomonas aeruginosa results in the emergence of isolates with increased honey resistance. *Ann Clin Microbiol Antimicrob* **13**: 19.

Carter, D.A., Blair, S.E., Cokcetin, N.N., Bouzo, D., Brooks, P., Schothauer, R., and Harry, E.J. (2016) Therapeutic manuka honey: No longer so alternative. *Front Microbiol* **7**: 1–11.

Chittezham Thomas, V., Kinkead, L.C., Janssen, A., Schaeffer, C.R., Woods, K.M., Lindgren, J.K., et

al. (2013) A dysfunctional tricarboxylic acid cycle enhances fitness of Staphylococcus epidermidis during β-lactam stress. *MBio* **4**: e00437-13.

Clugston, S.L., Barnard, J.F.J., Kinach, R., Miedema, D., Ruman, R., Daub, E., and Honek, J.F. (1998) Overproduction and characterization of a dimeric non-zinc glyoxalase I from Escherichia coli: Evidence for optimal activation by nickel ions. *Biochemistry* **37**: 8754–8763.

Cooke, J., Dryden, M., Patton, T., Brennan, J., and Barrett, J. (2015) The antimicrobial activity of prototype modified honeys that generate reactive oxygen species (ROS) hydrogen peroxide. *BMC Res Notes* **8**: 20.

Cooper, R.A., Jenkins, L., Henriques, A.F.M., Duggan, R.S., and Burton, N.F. (2010) Absence of bacterial resistance to medical-grade manuka honey. *Eur J Clin Microbiol Infect Dis* **29**: 1237–1241.

Deatherage, D.E. and Barrick, J.E. (2014) Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using breseq. *Methods Mol Biol* **1151**: 165–188.

Descottes, B. (2009) Cicatrisation par le miel, l'expérience de 25 années. *Phytothérapie* **7**: 112–116.

Dowd, S.E., Sun, Y., Secor, P.R., Rhoads, D.D., Wolcott, B.M., James, G.A., and Wolcott, R.D. (2008) Survey of bacterial diversity in chronic wounds using Pyrosequencing, DGGE, and full ribosome shotgun sequencing. *BMC Microbiol* **8**: 43.

Dunford, C.E. and Hanano, R. (2004) Acceptability to patients of a honey dressing for non-healing venous leg ulcers. *J Wound Care* **13**: 193–197.

Gutmann, L., Williamson, R., Moreau, N., Kitzis, M.D., Collatz, E., Acar, J.F., and Goldstein, F.W. (1985) Cross-resistance to nalidixic acid, trimethoprim, and chloramphenicol associated with alterations in outer membrane proteins of klebsiella, enterobacter, and serratia. *J Infect Dis* **151**: 501–507.

Halstead, F.D., Webber, M.A., Rauf, M., Burt, R., Dryden, M., and Oppenheim, B.A. (2016) In vitro activity of an engineered honey, medical-grade honeys, and antimicrobial wound dressings against biofilm-producing clinical bacterial isolates. *J Wound Care* **25**: 93–102.

Henriques, A.F., Jenkins, R.E., Burton, N.F., and Cooper, R.A. (2011) The effect of manuka honey on the structure of Pseudomonas aeruginosa. *Eur J Clin Microbiol Infect Dis* **30**: 167–171.

Hillitt, K.L., Jenkins, R.E., Spiller, O.B., and Beeton, M.L. (2017) Antimicrobial activity of Manuka honey against antibiotic-resistant strains of the cell wall-free bacteria Ureaplasma parvum and Ureaplasma urealyticum. *Lett Appl Microbiol* **64**: 198–202.

Iino, T., Komeda, Y., Kutsukake, K., Macnab, R.M., Matsumura, P., Parkinson, J.S., et al. (1988) New unified nomenclature for the flagellar genes of Escherichia coli and Salmonella typhimurium. *Microbiol Rev* **52**: 533–535.

Jenkins, R., Burton, N., and Cooper, R. (2014) Proteomic and genomic analysis of methicillin-resistant staphylococcus aureus (MRSA) exposed to manuka honey in vitro demonstrated down-regulation of virulence markers. *J Antimicrob Chemother* **69**: 603–615.

Klemm, P. (1986) Two regulatory fim genes, fimB and fimE, control the phase variation of type 1 fimbriae in Escherichia coli. *EMBO J* **5**: 1389–1393.

Knipping, S., Grünewald, B., and Hirt, R. (2012) Erste Erfahrungen mit medizinischem Honig in der Wundbehandlung im Kopf-Hals-Bereich. *HNO* **60**: 830–836.

Knöppel, A., Knopp, M., Albrecht, L.M., Lundin, E., Lustig, U., Näsvall, J., and Andersson, D.I. (2018) Genetic adaptation to growth under laboratory conditions in Escherichia coli and Salmonella enterica. *Front Microbiol* **9**: 1–16.

Knottenbelt, D.C. (2014) Honey in wound management: Myth, mystery, magic or marvel? *Vet J* **199**: 5–6.

Kronda, J.M., Cooper, R.A., and Maddocks, S.E. (2013) Manuka honey inhibits siderophore
production in Pseudomonas aeruginosa. *J Appl Microbiol* **115**: 86–90.

Kwakman, P.H.S., Van den Akker, J.P.C., Güçlü, A., Aslami, H., Binnekade, J.M., de Boer, L., et al.
(2008) Medical-Grade Honey Kills Antibiotic-Resistant Bacteria In Vitro and Eradicates Skin
Colonization. *Clin Infect Dis* **46**: 1677–1682.

Kwakman, P.H.S., te Velde, A.A., de Boer, L., Vandenbroucke-Grauls, C.M.J.E., and Zaat, S.A.J.
(2011) Two major medicinal honeys have different mechanisms of bactericidal activity. *PLoS*
One **6**: e17709.

Lee, J.H., Park, J.H., Kim, J.A., Neupane, G.P., Cho, M.H., Lee, C.S., and Lee, J. (2011) Low
concentrations of honey reduce biofilm formation, quorum sensing, and virulence in Escherichia
coli O157:H7. *Biofouling* **27**: 1095–1104.

Liu, X. and Matsumura, P. (1994) The FlhD/FlhC complex, a transcriptional activator of the
Escherichia coli flagellar class II operons. *J Bacteriol* **176**: 7345–7351.

Loughlin, M.F., Jones, M. V., and Lambert, P.A. (2002) Pseudomonas aeruginosa cells adapted to
benzalkonium chloride show resistance to other membrane-active agents but not to clinically
relevant antibiotics. *J Antimicrob Chemother* **49**: 631–639.

Lu, J., Cokcetin, N.N., Burke, C.M., Turnbull, L., Liu, M., Carter, D.A., et al. (2019) Honey can
inhibit and eliminate biofilms produced by Pseudomonas aeruginosa. *Sci Rep* **9**: 1–13.

MacLean, M.J., Ness, L.S., Ferguson, G.P., and Booth, I.R. (1998) The role of glyoxalase I in the
detoxification of methylglyoxal and in the activation of the KefB K+ efflux system in
Escherichia coli. *Mol Microbiol* **27**: 563–571.

Magnet, S. and Blanchard, J.S. (2005) Molecular insights into aminoglycoside action and resistance.
Chem Rev **105**: 477–497.

Mannervik, B. (2008) Molecular enzymology of the glyoxalase system. *Drug Metabol Drug Interact*
23: 13–27.

Mavric, E., Wittmann, S., Barth, G., and Henle, T. (2008) Identification and quantification of
methylglyoxal as the dominant antibacterial constituent of Manuka (Leptospermum scoparium)
honeys from New Zealand. *Mol Nutr Food Res* **52**: 483–489.

Merckoll, P., Jonassen, T.Ø., Vad, M.E., Jeansson, S.L., and Melby, K.K. (2009) Bacteria, biofilm
and honey: A study of the effects of honey on "planktonic" and biofilm-embedded chronic
wound bacteria. *Scand J Infect Dis* **41**: 341–347.

Molan, P.C. (1992) THE ANTIBACTERIAL ACTIVITY OF HONEY. 1. The nature of the
antibacterial activity. *Bee World1* **73**: 5–28.

Molan, P.C. (1999) Why honey is effective as a medicine. I. Its use in modern medicine. *Bee World*
80: 80–92.

Molan, P.C. and Betts, J.A. (2004) Clinical usage of honey as a wound dressing: an update. *J Wound*
Care **13**: 353–356.

Nolan, V.C., Harrison, J., and Cox, J.A.G. (2019) Dissecting the antimicrobial composition of honey.
Antibiotics **8**: 1–16.

O'Neill, J. (2014) Antimicrobial Resistance: Tackling a crisis for the health and wealth of nations The
Review on Antimicrobial Resistance Chaired.

Ozyamak, E., De Almeida, C., De Moura, A.P.S., Miller, S., and Booth, I.R. (2013) Integrated stress
response of Escherichia coli to methylglyoxal: Transcriptional readthrough from the nemRA
operon enhances protection through increased expression of glyoxalase I. *Mol Microbiol* **88**:
936–950.

Ozyamak, E., Black, S.S., Walker, C.A., MacLean, M.J., Bartlett, W., Miller, S., and Booth, I.R.

(2010) The critical role of S-lactoylglutathione formation during methylglyoxal detoxification in
Escherichia coli. *Mol Microbiol* **78**: 1577–1590.

Paulsen, I.T., Brown, M.H., and Skurray, R.A. (1996) Proton-dependent multidrug efflux systems.
Microbiol Rev **60**: 575–608.

Philippon, A., Labia, R., and Jacoby, G. (1989) MINIREVIEW Extended-Spectrum beta-Lactamases.
Antimicrob Agents Chemother **33**: 1131–1136.

Prüß, B.M. and Matsumura, P. (1996) A regulator of the flagellar regulon of Escherichia coli, flhD,
also affects cell division. *J Bacteriol* **178**: 668–674.

Scanlan, P.D., Hall, A.R., Blackshields, G., Friman, V.P., Davis, M.R., Goldberg, J.B., and Buckling,
A. (2015) Coevolution with bacteriophages drives genome-wide host evolution and constrains
the acquisition of abiotic-beneficial mutations. *Mol Biol Evol* **32**: 1425–1435.

Shan, Y., Lazinski, D., Rowe, S., Camilli, A., and Lewis, K. (2015) Genetic Basis of Persister
Tolerance to Aminoglycosides in Escherichia coli. *MBio* **6**: e00078-15.

Su, Y. bin, Peng, B., Li, H., Cheng, Z. xue, Zhang, T. tuo, Zhu, J. xin, et al. (2018) Pyruvate cycle
increases aminoglycoside efficacy and provides respiratory energy in bacteria. *Proc Natl Acad*
Sci U S A **115**: E1578–E1587.

Suojala, L., Kaartinen, L., and Pyörälä, S. (2013) Treatment for bovine Escherichia coli mastitis - an
evidence-based approach. *J Vet Pharmacol Ther* **36**: 521–531.

Szybalski, W. and Bryson, V. (1952) Genetic studies on microbial cross resistance to toxic agents:
Cross resistance of Escherichia coli to fifteen antibiotics. *J Bacteriol* **64**: 489–499.

Tadesse, D.A., Zhao, S., Tong, E., Ayers, S., Singh, A., Bartholomew, M.J., and McDermott, P.F.
(2012) Antimicrobial drug resistance in Escherichia coli from humans and food animals, United
States, 1950-2002. *Emerg Infect Dis* **18**: 741–749.

Tötemeyer, S., Booth, N.A., Nichols, W.W., Dunbar, B., and Booth, I.R. (1998) From famine to feast:
The role of methylglyoxal production in Escherichia coli. *Mol Microbiol* **27**: 553–562.

Truchado, P., López-Gálvez, F., Gil, M.I., Tomás-Barberán, F.A., and Allende, A. (2009) Quorum
sensing inhibitory and antimicrobial activities of honeys and the relationship with individual
phenolics. *Food Chem* **115**: 1337–1344.

Vandamme, L., Heyneman, A., Hoeksema, H., Verbelen, J., and Monstrey, S. (2013) Honey in
modern wound care: A systematic review. *Burns* **39**: 1514–1525.

Wang, R., Starkey, M., Hazan, R., and Rahme, L.G. (2012) Honey's ability to counter bacterial
infections arises from both bactericidal compounds and QS inhibition. *Front Microbiol* **3**: 1–8.

Wasfi, R., Elkhatib, W.F., and Khairalla, A.S. (2016) Effects of selected egyptian honeys on the
cellular ultrastructure and the gene expression profile of Escherichia coli. *PLoS One* **11**:
e0150984.

Werner, A. and Laccourreye, O. (2011) Honey in otorhinolaryngology: When, why and how? *Eur*
Ann Otorhinolaryngol Head Neck Dis **128**: 133–137.

White, J.W., Subers, M.H., and Schepartz, A.I. (1963) The identification of inhibine, the antibacterial
factor in honey, as hydrogen peroxide and its origin in a honey glucose-oxidase system. *BBA -*
Biochim Biophys Acta **73**: 57–70.

WHO (2018) Fact Sheet Antibacterial Resistance.

Yabes, J.M., White, B.K., Murray, C.K., Sanchez, C.J., Mende, K., Beckius, M.L., et al. (2017) In
Vitro activity of Manuka Honey and polyhexamethylene biguanide on filamentous fungi and
toxicity to human cell lines. *Med Mycol* **55**: 334–343.

Zhou, S., Zhuang, Y., Zhu, X., Yao, F., Li, Haiyan, Li, Huifang, et al. (2019) YhjX regulates the
growth of Escherichia coli in the presence of a subinhibitory concentration of gentamicin and

mediates the adaptive resistance to gentamicin. *Front Microbiol* **10**: 1–10.

788