Title: Genetic variation in heat tolerance of the coral *Platygyra daedalea* offers the potential for
adaptation to ocean warming.

Running head: Genetics of corals' heat tolerance

Holland Elder[1]*

Virginia Weis[1]

Jose Montalvo-Proano[2,3]

Veronique J.L Mocellin[2]

Andrew H. Baird[3]

Eli Meyer[1, 4]

Line K. Bay[2, 4]

1. Oregon State University, Corvallis, OR, USA. 97331
2. Australian Institute of Marine Science, 1526 Cape Cleveland Road, Cape Cleveland
4810, Queensland, Australia
AIMS@JCU
3. ARC Centre of Excellence for Coral Reef Studies, James Cook University, 1 James Cook
Dr, Douglas QLD 4814, Australia
4. These authors contributed equally

*corresponding author

760-622-9116

hollandelder@gmail.com

Keywords: coral reefs; resilience, heritability, genomic markers, allele frequency change

Paper type: Primary research article.

Global Change Biology

Abstract

32   Reef-building corals are foundational species in coral reef ecosystems and are threatened

by many stressors including rising ocean temperatures. In 2015/16 and 2016/17, corals around

the world experienced consecutive bleaching events and most coral populations are projected to

experience temperatures above their current bleaching thresholds annually by 2050. Adaptation

to higher temperatures is therefore necessary if corals are to persist in a warming future. While

many aspects of heat stress have been well studied, few data are available for predicting the

capacity for adaptive cross-generational responses in corals. To address this knowledge gap, we

quantified the heritability and genetic variation associated with heat tolerance in *Platygyra*

daedalea from the Great Barrier Reef (GBR). We tracked the survival of replicate quantitative

genetic crosses (or families) of coral larvae from six parents in a heat stress selection experiment.

We also identified allelic shifts in heat-selected survivors *versus* paired, non-selected controls of

the same coral crosses. We estimated narrow sense heritability to be 0.66 and detected a total of

1,069 single nucleotide polymorphisms (SNPs) associated with heat tolerance. An overlap of 148

unique SNPs shared between experimental crosses indicates that specific genomic regions are

responsible for heat tolerance of *P. daedalea* and some of these SNPs fall in coding regions.

These findings suggest that this *P. daedalea* population has the genetic prerequisites for

adaptation to increasing temperatures. This study also provides knowledge for the development

of high throughput genomic tools to screen for variation within and across populations to harness

or enhance adaptation through assisted gene flow and assisted migration.

Introduction

Increasing ocean temperatures brought about by human induced climate change have increased the frequency and severity of coral bleaching events worldwide to the point that it is recognized as a primary threat to coral reefs (Hutchings et al., 2019). Bleaching is the process of loss of dinoflagellate symbionts (Family Symbiodiniaceae) from host corals (Lajeunesse et al., 2018). Severe heat stress and mass bleaching caused by marine heat waves can result in rapid and widespread coral mortality (Glynn, 1993). Global and regional bleaching events have resulted in a substantial loss of coral cover worldwide over the last few decades (Hughes et al., 2018). Over the last five years alone, 30% of reefs in the Great Barrier Reef (GBR) showed decreases in coral cover (Hughes et al., 2019; Hughes et al., 2018; Long Term Reef Monitoring Program, 2019). For coral reefs to persist in an ocean that will continue to warm for the foreseeable future, corals will need to rapidly become more tolerant or resilient to higher temperatures. Consequently, there is now an urgent need to understand adaptive and evolutionary capacity of corals to this threat.

Mechanisms of thermal and bleaching tolerance are diverse. For example, the presence of low background densities of heat tolerant symbiont species can aid in recovery from heat stress (Bay et al., 2016; Berkelmans & van Oppen, 2006; Manzello et al., 2019; Oliver & Palumbi, 2011; Silverstein, Cunning, & Baker, 2014). In addition, plastic genomic responses including, epigenetic alterations and changes in gene expression can result in acclimatization in corals (Barshis et al., 2013; Dixon et al., 2018; Kenkel et al., 2013; Liew et al., 2020; Oliver & Palumbi, 2011; Thomas et al., 2019). Many studies have documented genetic variation in coral host heat tolerance that supports the potential for adaptation (Dixon et al., 2015; Howells et al.,

2016; Kenkel et al., 2013; Meyer et al., 2009; Woolsey et al., 2014). However, there are few

datasets that predict adaptation and evolution in corals. Working knowledge of the genomic

regions that influence heat tolerance, the extent to which coral host genetics contributes to heat

tolerance, and its cross generational persistence in a population will be essential to predict corals'

responses into the future. This knowledge will also inform conventional management approaches

as well as restoration programs using assisted gene flow and assisted migration to enhance the

tolerance of local populations (van Oppen et al., 2015).

Adaptation requires genetic variation underpinning fitness-related phenotypes. If

variation in a fitness-related phenotype has a genetic basis and is heritable and not otherwise

constrained, then selection can act to further propagate it within populations (Charmantier &

Garant, 2005). Narrow sense heritability describes the total phenotypic variance in a trait that can

be attributed to parental or additive genetics and is used to determine the genetic contribution to

traits such as heat tolerance (Falconer & Mackay, 1996; Lynch & Walsh, 1998). Estimates of

narrow sense heritability are essential for calculating selection differentials and in order to use

the breeder's equation to model the persistence of a trait in a population (Falconer & Mackay,

1996; Lynch & Walsh, 1998). Specifically, estimates of narrow sense heat heritability of heat

tolerance in corals will allow us to calculate potential rates of adaptation under various selection

scenarios.

Current estimates of heritability suggest that the genetic variation to support adaptation of

increased heat tolerance is present in coral populations in many different regions. For example, a

heritability value of 0.89 was found for bleaching resistance in *Orbicella faveolata* in the

Caribbean (Dziedzic et al. 2019) and values between 0.48 and 0.74 for heat tolerance for

Platygyra daedalea in the Persian Gulf (Kirk et al., 2018). Heritability was similarly high (0.87)

for heat tolerance in the larvae of the coral *A. millepora* from the GBR (Dixon et al., 2015).

However, heritability can vary substantially between species, between populations of

conspecifics, and is influenced by the environment in which it is measured (Falconer & Mackay,

1996; Lynch & Walsh, 1998), therefore a heritability estimate from the Persian Gulf should not

be used to estimate selection for heat tolerance or potential for adaptation in a population on the

GBR (Dixon et al., 2015). To understand the heritability of heat tolerance, selection for heat

tolerance, and the potential for adaptation in GBR populations, we need more heritability

estimates of heat tolerance from populations on the GBR.

Platygyra daedalea is a good candidate for estimates of heritability because it is widely

distributed in the Indo-Pacific and occurs in the much warmer Persian Gulf. Heritability was

recently estimated for a population of *P. daedalea* in the Persian Gulf, which provides an

opportunity to compare of heritability between populations from warmer and relatively cooler

environments, which could help reveal different genetic mechanisms of heat tolerance of this

species between the two environments (Kirk et al., 2017). The Gulf regularly experiences

temperatures of 33 - 34˚C in the summer and the bleaching threshold for *P. daedalea* is between

35 and 36˚C (Howells et al., 2016; Riegl et al., 2011). Indeed, the bleaching threshold

temperature for the coral assemblages is 3 - 7˚C higher in the Gulf when compared to other parts

of the world (29 - 32˚C), including the GBR (Berkelmans, 2002; Coles & Riegl, 2013; Hoegh-

Guldberg, 1999). This makes the comparison of heritability and genomic heat tolerance of these

two regions important for understanding how increased thermal tolerance evolves.

The genetic architecture of heat tolerance can also influence the response to selection and

the ability to develop markers of tolerance for conservation management as well as new

restoration approaches like assisted gene flow and assisted migration. Understanding the genetic

architecture of heat tolerance involves identifying coding and non-coding regions in the genome

that influence thermal tolerance. For example, in plants, allele frequency analysis has identified

markers associated with traits such as yield, drought tolerance and heat tolerance (Singh et al.,

2017; Venuprasad et al., 2009; Vikram et al., 2012; M. Wang et al., 2019). These markers can be

used to identify individuals in wild populations with increased heat tolerance. Numerous studies

have found differential gene expression associated with thermal tolerance in corals e.g. (Barshis

et al., 2013; Kenkel et al., 2013; Ruiz-jones & Palumbi, 2017; Traylor-knowles et al., 2017).

Fewer have identified genomic markers in coding and non-coding genes that are associated with

a thermally tolerant phenotype (Bay & Palumbi, 2014; Dixon et al., 2015; Dziedzic et al., 2019;

Jin et al., 2016). While these studies represent progress in identifying markers which connect the

genotype to the phenotype of heat tolerance, they also represent only two species from the Indo-

Pacific and one from the Caribbean.

To understand the potential for the GBR coral *Platygyra daedalea* to persist in a warming

environment through selection and genetic adaptation, we estimated heritability and identified

genomic regions associated with heat tolerance. Estimating heritability of heat tolerance allowed

us to quantify the host genetic contribution to this trait. We identified markers associated with

heat tolerant phenotypes by conducting a heat selection experiment and then assessing changes

in allele frequency in the survivors. Sequencing our heat-selected pools of bi-parental crosses

allowed us to identify markers associated with heat tolerance and examine how widespread these

markers were across families in both coding and non-coding genomic regions. This study

addresses fundamental questions regarding the adaptive potential of corals to heat stress and

provides evidence of their ability to survive and persist in a changing environment.

Methods

Experimental crosses and larval culture

To study the genetic basis and heritability of heat tolerance, we produced a series of crosses which are larval pools produced with sperm and eggs from two different parental colonies. For these experiments, we used 6 colony fragments of *Platygyra daedalea* collected from Esk Reef, an inshore reef in the central region of the Great Barrier Reef (18˚ 46'21.95" S, 146˚ 31' 19.33" E) with Australian Institute of Marine Science (AIMS) permit G12/35236.1. The corals were maintained in the National Sea Simulator at AIMS at 27˚C in flow-through sea water aquaria with 25% reduced ambient sunlight. In November 2016 (the 5th night after the full moon, at 18:30) the six parental colonies were isolated in ~60L bins with 0.04-μm filtered sea water (FSW). Between 18:30-19:00, the colonies spawned and gamete bundles were collected. After collection, gamete bundles were washed with FSW. Sperm and eggs were then separated using 60 μm mesh sieves. Eggs were thrice rinsed to eliminate residual sperm. We combined gametes from pairs of colonies according to a partial diallele design in Figure 1 to produce 10 families (hereafter termed crosses). Each cross was maintained at a density of 0.5 to 1.0 planula larvae per ml in separate 12L culture vessels with flow-through FSW at 27˚C and gentle aeration.

Heat stress experiments

To quantify the variation in coral heat tolerance, we measured larval survival during controlled heat stress exposure for each cross following Dixon et al. (2015). Larvae from each cross were incubated in one of two temperatures: ambient reef temperatures (27˚C) or an acute heat stress (35.5˚C). Within the 60L flow through FSW tanks, larvae were contained in

individual mesh bottomed 20ml wells (to allow for water circulation). We stocked 20 larvae

from each cross into six replicate wells and randomly distributed each cross among floating six-

well plates to control for position effects (six wells per cross per treatment in total). To reach

35.5˚C in the heat stress groups the temperature was raised from 27˚C by 1˚C per hour. Water

temperatures were maintained thereafter by computer-controlled heaters heating the water

incoming to the system. To develop a high-resolution time series of larval mortality we

monitored larval survival with manual counts of each well every 12 hours for the next ten days.

All larval survival data is available at https://github.com/hollandelder/Platygyradaedalea

_HeritabiltyandMarkersofThermalTolerance.

Statistical analysis of heat tolerance

We investigated the effect of heat stress using a Kaplan Meier Survival Analysis and a

Cox Proportional Hazards (CPH) model that compared individual survival at 35.5˚C in the

context of cross and temperature (Kaplan and Meier 1958 and Cox 1972). This analysis allowed

us to determine the strength of genetic effects between crosses on the survival of larvae at high

temperatures. For this analysis, we used the Survival package in R (Therneau 2015). Our model

fit individual survival, allowing slopes to vary for temperature, cross, and the interaction of

temperature and cross. The code for the Kaplan Meier survival plots and CPH model can be

referenced at https://github.com/hollandelder/Platygyradaedalea_HeritabiltyandMarkersof

ThermalTolerance. We were also able to obtain Kaplan Meier estimates for the probability of

survival given hour of exposure to heat stress, providing statistics for comparing crosses.

We used these data to estimate narrow-sense heritability of variation in heat tolerance

(Falconer & Mackay, 1996; Wilson et al., 2010). For this analysis, we used a model for assessing

heritability using restricted maximum likelihood (REML) implemented in the software package

Wombat (Meyer 2007). We followed the same procedure we have previously described for

analysis of genetic variation in coral larvae (Dixon et al., 2015; Meyer et al., 2009). This analysis

uses known half-sibling relationships among larval biparental crosses to partition variance into

genetic and non-genetic components from which narrow-sense heritability (h^2) can be calculated

(Lynch & Walsh, 1998). We estimated heritability of survival from the time-course survival

described above, at the experiment endpoint, and from the cumulative probability of survival.

Genetic analysis of selection for heat stress tolerance

To identify genetic markers under selection during acute heat exposure, we conducted a

controlled stress experiment to eliminate the most heat sensitive genotypes in each pool. To

provide sufficient larvae for these experiments, we stocked larvae from each cross into replicated

flow-through culture containers in each of two treatments (500 larvae per container). We

maintained four replicate containers per cross at ambient reef temperatures (27˚C) throughout the

experiment and exposed another four containers to a heat stress treatment. For this treatment, we

gradually increased temperatures by 1˚C per hour to a maximum temperature of 35.5˚C then

maintained that temperature thereafter. Temperatures were maintained as described above by

incubating the flow-through culture containers in larger 60L flow-through tanks. We sampled

100 - 150 larvae from each replicate after 53 h, and preserved larvae in 100% ethanol for

subsequent genetic analysis.

We prepared genotyping libraries from genomic DNA from pooled larval samples. We

extracted genomic DNA using the Omega Bio-tek E.Z.N.A. Tissue Kit (Omega Bio-tek,

Norcross, GA). We genotyped thousands of single nucleotide polymorphisms (SNPs) in each

sample using 2bRAD, a sequencing-based approach for SNP genotyping (Wang et al. 2012) that

has been previously applied in corals (Dixon et al., 2015; Howells et al., 2016). We labeled each

library with a sample-specific pair of barcodes and combined libraries in equimolar ratios based

on qPCR using SensiFAST SYBR Hi-ROX master mix (Bioline), then sequenced the combined

pools on four lanes of 50bp single end reads (Illumina HiSeq2500) at Ramaciotti Centre for

Genomics at the University of New South Wales (Sydney, Australia).

Prior to statistical analysis, we processed the raw data to exclude low-quality reads. We

removed any reads that had ten or more bases that we considered low quality (Q-score > 30). We

removed any reads having twelve bases matching Illumina adaptors. This quality filtering

process produced a set of high-quality reads for genotyping analysis. All filtering scripts used are

available at (https://github.com/Eli-Meyer/sequence_processing).

To enable comparisons between this genomic data set from a GBR *P. daedalea*

population with the unusually heat-tolerant *P. daedalea* population from the Persian Gulf (

Howells et al., 2016), we mapped our reads against a reference we previously developed from

larvae of Persian Gulf *P. daedalea* (SRA accession: SRP066627). We aligned high quality reads

against this reference and filtered alignments to determine allele frequencies at each position as

described previously (Howells et al., 2016; Wang et al., 2012).

After alignment to the reference, we further filtered sequence data to remove loci with

weak or ambiguous matches, those with low coverage, those that occurred in too few samples,

and samples with too many missing loci using the following process. First, we removed all loci

with ambiguous matches identified by mapping to two or more regions in the reference. Then we

removed weak matches or those that did not span at least 32 of a 36 base sequence and matched

fewer than 30 of the bases within that span of sequence. We then filtered loci coverage with an

initial coverage filter requiring that there needed to be at least 10x coverage of a tag. We also

removed monomorphic loci as well as those not found in 25% of samples. We removed samples

with < 30,000 loci. We required that each locus be genotyped in at least seven out of eight

replicates for each cross in order to compare alleles at that locus across replicates in each

treatment. We also removed tags with more than two SNPs, because they are likely to come from

repetitive regions. Finally, we filtered so there would be one SNP per tag to control for non-

independence and linkage. All 2bRAD sequence filtering was conducted using scripts available

at (https://github.com/Eli-Meyer/2brad_utilities).

To compare allele frequencies between control larvae and the survivors of heat stress, we

compared replicate samples from each treatment within each cross. Since these samples

represented pools of large numbers of individuals (>100) rather than single diploid individuals,

we interpreted nucleotide frequencies in each sample directly as allele frequencies in the pool,

following the same general approach previously described as Pool-Seq (Kofler et al., 2011). We

applied a stringent coverage threshold of 80x to minimize sampling error that is likely to distort

estimates of allele frequencies at lower coverage. To test for differences in allele frequencies

between control larvae and the those surviving heat stress, we used logistic regression to analyze

the number of observations of each allele in each sample, conducting generalized linear models

in R (glm with argument "cross=binomial(logit)"). This allowed us to determine the effects of

treatment on allele frequencies at each locus, identifying loci associated with heat tolerance in

each cross. We examined the presence and absence of SNPs identified as associated with heat

tolerance from the allele frequency analysis between all combinations of two and three crosses.

To investigate the functional implications of genetic associations, we mapped each

2bRAD tag to the *P. daedalea* transcriptome (Kirk et al., 2017). To find the gene function, we

searched for the UniProt match for the annotation of each matched transcript. We required a

match of 34 base pairs of the 36 base pair tag so that matches would be unambiguous. This was

necessary, because of the short length of the restriction fragments analyzed here. It is likely that

if these standards were relaxed, more of the differentially enriched genomic regions would map

to functional regions but could also introduce false positives.

Results

Survival in heat stress experiments

A Cox proportional hazards (CPH) model revealed significant effects of heat x cross (p-

value < 0.0001). Endpoint survival probabilities ranged more among crosses at 35.5˚C (0.405 to

0.750) than at 27˚C (0.861 to 0.975) (Fig. 3). Cross DB had the greatest survival probability at

35.5˚C 0.75 (+/- 0.04), while cross CA had the lowest survival probability 0.41 (+/- 0.04) (Fig.3).

Coral E provided sperm in both EC and EF crosses which are both in the top four survival

probabilities 0.68 (+/- 0.04) and 0.64 (+/- 0.04) respectively. Coral C provided eggs in crosses

DC and BC and provided sperm in cross CA, and this coral is associated with the bottom three

survival probabilities 0.48 (+/- 0.04), 0.44 (+/- 0.05), and 0.40 (+/- 0.04). These analyses

demonstrate substantial variation in heat tolerance among larval crosses which can be attributed

to genetic background.

Quantitative genetic analysis of variation in heat tolerance

Variation in survival during heat stress was highly heritable, with additive genetic

variation explaining 66% of variation in survival (i.e. $h^2 = 0.66$; Table 1). We also estimated h^2

using the survival probabilities from the Kaplan Meier analysis, and recovered a similar estimate

($h^2 = 0.64$). Both metrics reflect the total mortality over the entire time course.

Genomic analysis of markers under selection during heat stress

Genomic analysis of heat stressed versus control larval pools found that experimental

treatment altered allele frequencies at hundreds of loci. Logistic regression of multilocus SNP

genotypes from surviving heat stress versus control larval pools (Fig. 4) uncovered tens-to-

hundreds of markers in each cross that were significantly affected by heat stress (Table 2). The

number of alleles significantly associated with survival under heat stress differed widely among

crosses. This variation was not correlated with the number of markers genotyped in each sample

($P = 0.47$, $R^2 = 0.067$), genetic distance between the parents ($P = 0.46$, $R^2 = 0.071$), or heat

tolerance of each cross ($P = 0.49$, $R^2 = 0.063$) (Table 2). The variation was also not associated

with a particular parent. For example, parents C and D each produced crosses with few markers

under selection (BC and DB), but the combination of these parents in cross DC had the most:

366 SNPs associated with survival under heat stress (Table 2). This comparison suggests the

large number of alleles changing in frequency in cross DC results from a combination of parental

alleles at many loci, producing larvae with multiple deleterious alleles.

To evaluate the generalizability of genetic associations we compared these significant

loci among crosses. A proportion of markers were consistently associated with heat stress across

multiple crosses; 2 - 17 markers identified in each cross were shared with one or more crosses,

and 2 - 5 of these were identified in all three (Figure 5). Altogether, 148 unique markers were

present in at least 2 crosses (Table 3). The same two markers were found in seven crosses, a

single marker overlapped in six crosses, and the same four markers overlapped in five crosses

(Table 3). However, most markers associated with heat tolerance in one cross were not

associated with heat tolerance in unrelated crosses. For example, cross DC shared no parents

with crosses EF or FA and 82-92% of the markers identified in each cross were not associated

with heat stress in the other two.

Altogether, 22 markers could be unambiguously assigned to a position in the

transcriptome and five SNPs were located in annotated transcripts (Table 4). Reproducible

associations across different genetic backgrounds were also apparent for four transcripts without

matches in UniProt. Furthermore, five loci in transcripts were associated with heat tolerance in

more than one family (Table 4).

Discussion

Genetic background affects heat tolerance

Our survival model shows that heat tolerance is significantly affected by genetic

background. Survival probabilities varied significantly between crosses indicating that genetics

contributed to variation in heat tolerance. Coral parent E was associated with crosses that had

high survival probabilities and Coral parent C was associated with crosses that had lower

survival probabilities. We conclude that parental genetics contribute to heat tolerance of

offspring and consideration of parental genetics will be important in coral and conservation as

well as assisted gene flow and assisted migration.

Heritability and modeling heat tolerance

Quantifying the amount of genetic variance that contributes to a phenotype is important

for evaluating adaptive potential of a population (Falconer & Mackay, 1996). Our results add to

only three published narrow sense heritability estimates for heat tolerance in corals, all of which

find this trait is highly heritable (Dixon et al., 2015; Dziedzic et al., 2019; Kirk et al., 2018). Our

study is the only estimate of heritability for a sub-massive and relatively heat tolerant coral

species from the GBR. Our result for *P. daedalea* likewise indicates that heritability in larval

survival after heat stress was substantial ($h^2 = 0.66$) and suggests that the *P. daedalea*

populations have substantial standing genetic variation which contributes to heat tolerance and

that this population has an innate capacity to adapt to some level of ocean warming.

Heritability estimates from different populations can give us valuable information about

selection in a given geographic region. Heritability estimates are dependent on the environment

and the standing genetic variation in the population for which the estimate is made (Falconer &

Mackay, 1996). If there are low levels of genetic variation contributing to a trait and all parents

express the same phenotype, the heritability estimate will be lower (Falconer & Mackay, 1996).

A novel comparison is the contrast of the heritability estimate (0.48) found by Kirk and

colleagues for *P. daedalea* in the Persian Gulf (Kirk et al., 2018). The Persian Gulf *P. daedalea*

has lived in a very shallow and warm sea for thousands of years (Purkis & Riegl, 2012), which

may mean that a degree of selection for heat tolerance has already taken place resulting in less

genetic diversity in the population (Falconer & Mackay, 1996). The lower heritability value is

evidence to support this and the higher degree of heritability for heat tolerance in the central

GBR *P. daedalea* population suggests that until very recently, heat stress has not been the

predominant selective pressure on this population.

Responses to selection depend on genetic variation in the trait under selection (Falconer

& Mackay, 1996; Lynch & Walsh, 1998) and our study contributes to building the necessary

quantitative genetic data to model heat tolerance under various selection regimes. In its simplest

form, this can be expressed in the breeder's equation, which describes the response to selection

as the product of the selection differential (S) and heritability of the trait under selection. To our

knowledge, selection differentials have not been empirically estimated in corals, but considering

the widespread reductions in coral populations following coral bleaching (Bellwood et al., 2004;

Bruno & Selig, 2007; De'ath et al., 2012; Hoegh-Guldberg et al., 2007; Hughes et al., 2017,

2018) they are likely high for sensitive species. For truncation selection, where all individuals

with trait values above a threshold contribute offspring to the next generation, selection

differentials are equal to the difference in mean trait values between the selected population and

the original population (Falconer & Mackay, 1996; Lynch & Walsh, 1998) expressed as standard

deviations of the population mean. If the selection differential equals one standard deviation, all

parents with traits one standard deviation above the population mean contribute to the next

generation. Based on the distribution of cumulative survival probabilities in heat stress from our

experiment (0.57 ± 0.18) and the heritability of variation in heat tolerance we estimated from

these data (0.66), the Breeder's Equation predicts a response to selection of 0.12. We would

expect that if larvae of the next generation after selection were subjected to the same heat stress

treatment, the average mortality would be 0.48 (0.59 - 0.12), a 20% reduction in overall

mortality. If instead of the high h^2 estimated in our study (0.66), we modeled the responses to

selection using a low h^2 value of 0.1, we would predict only a 3% reduction in mortality in the

following generation. While the estimates above are underpinned by too many assumptions to

predict responses in natural populations of corals, the comparison highlights how empirical

estimations of narrow-sense heritability can be used to predict adaptive responses. The relevance

of these estimates for population responses depends on the extent to which larval survival during

heat stress is directly under selection in natural populations, or alternatively the effectiveness of

larval heat tolerance as a proxy for heat tolerance in adult colonies.

Allele frequency analysis

Physiological and fitness traits can be determined by single or few loci of large effects,

but commonly they are underpinned by many loci of small effect. Our genomic analysis

identified tens-to-hundreds of genetic markers associated with heat tolerance in each cross, based

on changes in allele frequencies between heat selected and control pools. Comparing between

crosses allowed us to investigate the generality of these patterns among different genetic

backgrounds. This comparison revealed a number of markers with consistent changes in allele

frequencies in multiple crosses (Table 3, 4, and Supplemental Table 1). However, many of the

associations detected in each cross were not shared in comparisons of multiple crosses. The lack

of overlap for many markers likely results from the underlying distribution of those alleles in our

parental population. In isolated populations the loci contributing to a trait are predicted to be

fewer and they are of larger effect size (Savolainen et al., 2013). However, in populations

experiencing immigration, the locus effect size and the number of loci that contribute to a trait

become more variable (Savolainen et al., 2013). Corals can migrate in their larval phase before

metamorphosis into an adult, so it is entirely possible that individuals originating from disparate

reefs were among those chosen for our parental population. The observation that many markers

associated with heat stress were specific to individual crosses suggests that many alleles of small

effect contribute to variation in heat tolerance and that many different combinations of these

small effect alleles can affect heat tolerance (Bay & Palumbi, 2014; Kirk et al., 2018). These

differences may also result in part from epistatic interactions where the effects of variation at

these loci depend on the genetic background. Few data on either frequencies of heat tolerance

alleles or epistatic effects are available for any coral species. However, the repeated observation

of some of these associations across different genetic backgrounds provides strong evidence that

these markers are linked to genetic factors contributing to variation in heat tolerance.

Future studies can use markers identified in this and other studies. Starting with genetic

markers associated with habitat variation in previous studies (Capper et al., 2015), we can test

these associations through a combination of laboratory experiments and extensive population

surveys of allele frequencies (Jin et al., 2016). These studies revealed that the frequencies of

alleles associated with enhanced tolerance for environmental stress range from 0.48 in some

populations to 0.92 in others (Jin et al., 2016). Further studies are needed to extend these surveys

across species, markers, and geographic regions to develop an ecosystem-scale perspective on

the potential for coral adaptive responses to ocean warming.

SNPs associated with heat tolerance occurring in coding genes

In our study, we found a subset of regions in coding genomic regions associated with heat

tolerance, five of which were observed in multiple crosses (Tables 3 and 4). These SNPS were

found in genes associated with the extra-cellular matrix and metabolism. There were also SNPs

in four unannotated transcripts. These SNP-containing genes found in this study are candidates

for further study of their role in this trait and could serve as heat tolerance markers.

Three SNPs were associated with coding genes that play roles in the extracellular matrix:

(1) GPI_Mannosyltransferase_2, which functions in the assembly of

glycosylphosphatidylinositol (GPI) anchors that attach proteins to cell membranes (Kang et al.

2005), (2) cad96ca, an E type cadherin, that initiates cell-cell adhesion (Lodish et al. 2013), and

(3) metalloendopeptidase, a metal-dependent protease that restructures the extracellular matrix

(Lodish et al. 2013) (Table 4). A SNP was also found in Serine/threonine-protein kinase (PAK3)

a member of cell regulatory pathways influencing regulation of cytoskeletal formation, cell

migration, and the cell cycle (Deleris et al. 2011). Three coral crosses had a SNP in the Krebs

cycle enzyme isocitrate dehydrogenase (IDH) (Garret & Grisham, 2013). This enzyme has been

identified in gene expression studies as important to the coral thermal stress response (Kenkel,

Meyer, et al., 2013; Polato et al., 2010). IDH participates in handling reactive oxygen species

and is one of the 44 proteins identified as the minimal stress proteome (Kultz, 2005). The SNP in

IDH associated with heat tolerance further strengthens its proposed role in the thermal stress

response in corals.

Many studies have pointed out the abundance of unannotated genes in the genomes of

early-diverging animals and the absence of any pathways to understand their function. The four

SNPs that we found which were unannotated in the transcriptome show overlap between crosses

highlighting the potential importance of understanding the role these unknown genes in thermal

tolerance. These four unannotated transcripts warrant further functional investigations, such as

full sequencing, domain analysis, and expression in functional assays. Knowing the function of

these genes will be important for having a complete picture of the alleles controlling thermal

tolerance.

Comparisons of populations of *P. daedalea* in more typical reef temperature conditions

(this study) and unusually warm conditions (Kirk et al., 2018) will be important to understanding

the genomic basis of heat tolerance. Two SNPs from the present study were also significantly

associated with heat tolerance in the Persian Gulf population and thus represent excellent

candidates for further study. This could include screens for natural abundance among thermal

habitats and potential validation for gene function through knock-down from CRISPR-Cas9

induced mutagenesis (Cleves et al., 2018). This unexpected lack of shared markers is further

evidence that many alleles of small effect contribute to heat tolerance and that they vary in their

distribution across populations.

Conclusions

Our findings demonstrate that heat tolerance is highly heritable (i.e., > 60%) in the coral

Platygyra daedalea from the GBR demonstrating high potential for adaptation. The heat stress

time course (Table 1) supports the conclusion that genetic variation explains a substantial

fraction of the variation in heat tolerance among our crosses. Our analysis identified hundreds of

markers associated with heat tolerance, including markers that were repeatedly associated with

heat tolerance across different genetic backgrounds. This supports the conclusion that heat

tolerance is a complex polygenic trait determined by many alleles of small affect that are likely

to be species and population specific. Despite this complexity, we found markers in multiple

crosses that may be important for heat tolerance in *P. daedalea*. These markers associated with

heat tolerance in our study of *P. daedalea* from typical coral reef conditions in the GBR will

provide a valuable comparison with future genomic studies of heat tolerant populations from the

Persian Gulf. The four unannotated transcripts found in this study warrant further functional

investigations to find their role in the animal and examine how they might function in heat

tolerance. Our findings build on the growing body of evidence that natural populations of corals

harbor substantial genetic variation in heat tolerance that can support adaptive responses to

selection and identify genetic markers for heat tolerance in a globally distributed coral species.

These heat tolerance markers are essential to any type of reseeding or assisted gene flow and

assisted migration programs, which may be necessary to maintaining reefs in the future as we

work to reduce CO_2 emissions and mitigate the effects of climate change.

Acknowledgements

We thank the staff of the National Sea Simulator at the Australian Institute of Marine Science for

their assistance in the collection and husbandry of the corals used in these experiments. We also

thank Ida Bjornsbo for her tireless assistance in counting coral larvae at all hours of the day and

night. This research was conducted under Great Barrier Reef Marine Park Authority permit

(G11/34671.1) and supported with funding from the National Science Foundation Graduate

Research Fellowship to Holland Elder, National Science Foundation Graduate Research

Opportunities Worldwide to Holland Elder and internal grants from AIMS to Line K. Bay.

Authors' contributions

E.M., H.E., and L.B. conceived the experiments. H.E., V.M., J.M., and L.B. conducted

experiments, L.B. and A.B. provided advice and logistical support through the project, H.E.

conducted genetic and statistical analysis and wrote the manuscript. L.B. and E.M. helped with

genetic and statistical analysis and L.B., V.W. and A. B. helped write the manuscript.

References

Barshis, D. J., Ladner, J. T., Oliver, T. a, Seneca, F. O., Traylor-Knowles, N., & Palumbi, S. R.

    (2013). Genomic basis for coral resilience to climate change. *Proceedings of the National*

    Academy of Sciences of the United States of America, *110*(4), 1387–1392.

    https://doi.org/10.1073/pnas.1210224110

Bay, L. K., Doyle, J., Logan, M., & Berkelmans, R. (2016). Recovery from bleaching is

    mediated by threshold densities of background thermo-tolerant symbiont types in a reef-

    building coral. *Royal Society Open Science*, *3*(160322).

Bay, R. a, & Palumbi, S. R. (2014). Report Multilocus Adaptation Associated with Heat

    Resistance in Reef-Building Corals. *Current Biology*, 1–5.

    https://doi.org/10.1016/j.cub.2014.10.044

Bellwood, D. R., Hughes, T. P., Folke, C., & Nystro, M. (2004). Confronting the coral reef

    crisis. *Nature*, *429*(June), 827–833.

Berkelmans, R. (2002). Time-integrated thermal bleaching thresholds of reefs and their variation

    on the Great Barrier Reef. *Marine Ecology Progress Series*, *229*, 73–82.

Berkelmans, R., & van Oppen, M. J. H. (2006). The role of zooxanthellae in the thermal

    tolerance of corals: a "nugget of hope" for coral reefs in an era of climate change.

    Proceedings. Biological Sciences / The Royal Society, *273*(1599), 2305–2312.

    https://doi.org/10.1098/rspb.2006.3567

Bruno, J. F., & Selig, E. R. (2007). Regional Decline of Coral Cover in the Indo-Pacific : Timing

    , Extent , and Subregional Comparisons. *PloS One*, *8*.

    https://doi.org/10.1371/journal.pone.0000711

Capper, R. L., Jin, Y. K., Lundgren, P. B., Peplow, L. M., Matz, M. V, & Oppen, M. J. H. Van.

(2015). Quantitative high resolution melting : two methods to determine SNP allele

frequencies from pooled samples. *BMC Genetics*, *16*(62), 1–13.

https://doi.org/10.1186/s12863-015-0222-z

Charmantier, A., & Garant, D. (2005). Environmental quality and evolutionary potential : lessons

from wild populations. *Proceedings of the Royal Society*, *272*, 1415–1425.

https://doi.org/10.1098/rspb.2005.3117

Cleves, P. A., Strader, M. E., Bay, L. K., Pringle, J. R., & Matz, M. V. (2018). CRISPR/Cas9-

mediated genome editing in a reef building coral. *PNAS*, *115*(20), 5235–5240.

https://doi.org/10.1073/pnas.1722151115

Coles, S. L., & Riegl, B. M. (2013). Thermal tolerances of reef corals in the Gulf : A review of

the potential for increasing coral survival and adaptation to climate change through assisted

translocation. *Marine Pollution Bulletin*, *72*(2), 323–332.

https://doi.org/10.1016/j.marpolbul.2012.09.006

Cunning, R., Vaughan, N., Gillette, P., Capo, T. R., Mate, J. L., & Baker, A. C. (2015). Dynamic

regulation of partner abundance mediates response of reef coral symbioses to environmental

change. *Ecology*, *96*(5), 1411–1420.

Cunning, R, Silverstein, R. N., & Baker, A. C. (2015). Investigating the causes and consequences

of symbiont shuffling in a multi-partner reef coral symbiosis under environmental change.

Proc. R. Soc. B, *282*(20141725).

Cunning, Ross, Silverstein, R. N., & Baker, A. C. (2018). Symbiont shuffling linked to

differential photochemical dynamics of Symbiodinium in three Caribbean reef corals. *Coral*

Reefs, *37*(1), 145–152. https://doi.org/10.1007/s00338-017-1640-3

De'ath, G., Fabricius, K. E., Sweatman, H., & Puotinen, M. (2012). The 27 – year decline of

coral cover on the Great Barrier Reef and its causes. *PNAS*, *109*(44), 17995–17999.

https://doi.org/10.1073/pnas.1208909109

Dixon, G. B., Davies, S. W., Aglyamova, G. V., Meyer, E., Bay, L. K., & Matz, M. V. (2015).

Genomic determinants of coral heat tolerance across latitudes. *Science*, *348*(6242), 1460–

1462. https://doi.org/10.1126/science.1261224

Dixon, G., Davies, S., Aglyamova, G., Meyer, E., Bay, L. K., & Matz, M. V. (2015). Genomic

Determinants of coral heat tolerance across latitudes. *Science*, *348*(6242), 1460–1462.

Dixon, G., Liao, Y., Bay, L. K., & Matz, M. V. (2018). Role of gene body methylation in

acclimatization and adaptation in a basal metazoan. *PNAS*, *115*(52), 13342–13346.

https://doi.org/10.1073/pnas.1813749115

Dziedzic, K. E., Elder, H., & Meyer, E. (2019). Heritable variation in bleaching responses and its

functional genomic basis in reef-building corals (Orbicella faveolata). *Molecular Ecology*,

1–16.

Falconer, D. S., & Mackay, T. F. C. (1996). *Introduction to Quantitative Genetics*.

Garret, R. H., & Grisham, C. M. (2013). *Biochemistry*.

Glynn, P. W. (1993). Coral reef bleaching: ecological perspectives. *Coral Reefs*, *12*, 1–17.

https://doi.org/10.1007/BF00303779

Hoegh-Guldberg, O. (1999). Climate change, coral bleaching and the future of the world's coral

reefs. *Marine and Freshwater Research*, *50*(8), 839–866. https://doi.org/10.1071/MF99078

Hoegh-Guldberg, O., Harvell, C. D., Sale, P. F., Edwards, A. J., Caldeira, K., Knowlton, N.,

Eakin, C. M., Muthiga, N., Bradbury, R. H., Dubi, A., Hatziolos, M. E., Mumby, P. J.,

Hooten, A., Steneck, R. S., Greenfield, P., Gomez, E., & Iglesias-Prieto, R. (2007). Coral

Reefs Under Rapid Climate Change and Ocean Acidification. *Science*, *318*(5857), 1737–

1742.

Howells, E. J., Abrego, D., Meyer, E., Kirk, N., & Burt, J. A. (2016). Host adaptation and

unexpected symbiont partners enable reef-building corals to tolerate extreme temperatures.

Global Change Biology, 1–13. https://doi.org/10.1111/gcb.13250

Hughes, T., Kerry, J., Baird, A., Connolly, S., Chase, T., Dietzel, A., Hill, T., Hoey, A.,

Hoogenboom, M., Jacobson, M., Kerswell, A., Madin, J., Mieog, A., Paley, A., Pratchett,

M., Torda, G., & Woods, R. (2019). Global warming impairs stock - recruitment dynamics

of corals. *Nature*, *568*, 387–390. https://doi.org/10.1038/s41586-019-1081-y

Hughes, T. P., Kerry, J. T., Álvarez-noriega, M., Álvarez-romero, J. G., Anderson, K. D., Baird,

A. H., Babcock, R. C., Beger, M., Bellwood, D. R., Berkelmans, R., Bridge, T. C., Butler, I.

R., Byrne, M., Cantin, N. E., Comeau, S., Connolly, S. R., Cumming, G. S., Dalton, S. J.,

Diaz-Pulido, G., … Wilson, S. K. (2017). Global warming and recurrent mass bleaching of

corals. *Nature*, *543*, 373–377. https://doi.org/10.1038/nature21707

Hughes, T. P., Kerry, J. T., Baird, A. H., Connolly, S. R., Dietzel, A., Eakin, C. M., Heron, S. F.,

Hoey, A. S., Hoogenboom, M. O., Liu, G., Mcwilliam, M. J., Pears, R. J., Pratchett, M. S.,

Skirving, W. J., Stella, J. S., & Torda, G. (2018). Global warming transforms coral reef

assemblages. *Nature*, *556*, 492–496.

Hutchings, P., Kingsford, M., & Hoegh-Guldberg, O. (2019). *The Great Barrier Reef: Biology,*

Environment, and Management. Csiro Publishing.

Jin, Y. K., Lundgren, P., Lutz, A., Raina, J., Howells, E. J., Paley, A. S., Willis, B. L., & Oppen,

M. J. H. Van. (2016). *Genetic markers for antioxidant capacity in a reef-building coral*.

May.

Kenkel, C. D., Goodbody-Gringley, G., Caillaud, D., Davies, S. W., Bartels, E., & Matz, M. V.

(2013). Evidence for a host role in thermotolerance divergence between populations of the

mustard hill coral (Porites astreoides) from different reef environments. *Molecular Ecology*,

22(16), 4335–4348. https://doi.org/10.1111/mec.12391

Kenkel, C. D., Meyer, E., & Matz, M. V. (2013). Gene expression under chronic heat stress in

populations of the mustard hill coral (Porites astreoides) from different thermal

environments. *Molecular Ecology*, *22*(16), 4322–4334. https://doi.org/10.1111/mec.12390

Kirk, N., Howells, E., Abrego, D., Burt, J., & Meyer, E. (2018). Genomic and transcriptomic

signals of thermal tolerance in heat-tolerant corals (Platygyra daedalea) of the

Arabian/Persian Gulf. *Molecular Ecology*, *27*, 5180–5194.

Kofler, R., Orozco-terWengel, P., de Maio, N., Pandey, R. V., Nolte, V., Futschik, A., Kosiol,

C., & Schlötterer, C. (2011). Popoolation: A toolbox for population genetic analysis of next

generation sequencing data from pooled individuals. *PLoS ONE*, *6*(1).

https://doi.org/10.1371/journal.pone.0015925

Kultz, D. (2005). Molecular And Evolutionary Basis of the Cellular Stress Response. *Annual*

Review Physiology, *67*, 225–257.

https://doi.org/10.1146/annurev.physiol.67.040403.103635

Lajeunesse, T. C., Parkinson, J. E., Gabrielson, P. W., Jeong, H. J., Reimer, J. D., Voolstra, C.

R., Santos, S. R., Lajeunesse, T. C., Parkinson, J. E., Gabrielson, P. W., Jeong, H. J., &

Reimer, J. D. (2018). Systematic Revision of Symbiodiniaceae Highlights the Antiquity and

Diversity of Coral Endosymbionts Article Systematic Revision of Symbiodiniaceae

Highlights the Antiquity and Diversity of Coral Endosymbionts. *Current Biology*, *28*(16),

2570-2580.e6. https://doi.org/10.1016/j.cub.2018.07.008

Liew, Y. J., Howells, E. J., Wang, X., Michell, C. T., Burt, J. A., Idaghdour, Y., & Aranda, M.

    (2020). Intergenerational epigenetic inheritance in reef-building corals. *Nature Climate*

    Change, *10*, 254–259. https://doi.org/10.1038/s41558-019-0687-2

Long Term Reef Monitoring Program. (2019). *Annual Report on Reef Condition: Mixed bill of*

    health for the Great Barrier Reef. https://www.aims.gov.au/reef-monitoring/gbr-condition-

    summary-2018-2019

Lynch, M., & Walsh, B. (1998). *Genetics and Analysis of Quantitative Traits*.

Manzello, D. P., Jankulak, M., Matz, M. V, Enochs, I. C., Valentino, L., Carlton, R. D.,

    Kolodziej, G., Serrano, X., & Towle, E. K. (2019). Role of host genetics and heat - tolerant

    algal symbionts in sustaining populations of the endangered coral Orbicella faveolata in the

    Florida Keys with ocean warming. *Global Change Biology*, *25*, 1016–1031.

    https://doi.org/10.1111/gcb.14545

Marshall, P. A., & Baird, A. H. (2000). Bleaching of corals on the Great Barrier Reef:

    differential susceptibilities among taxa. *Coral Reefs*, *19*(2), 155–163.

    https://doi.org/10.1007/s003380000086

Meyer, E., Davies, S., Wang, S., Willis, B., Abrego, D., Juenger, T., & Matz, M. (2009). Genetic

    variation in responses to a settlement cue and elevated temperature in the reef-building coral

    Acropora millepora. *Marine Ecology Progress Series*, *392*, 81–92.

    https://doi.org/10.3354/meps08208

Mieog, J. C., Van Oppen, J. H., Berkelmans, R., Stam, W. T., & Olsen, J. L. (2009).

    Quantification of algal endosymbionts (Symbiodinium) in coral tissue using real-time

    PCR. *Molecular Ecology Resources*, *9*, 74–82. https://doi.org/10.1111/j.1755-

    0998.2008.02222.x

Núnez-Pons, L., Bertocci, I., & Baghdasarian, G. (2017). Symbiont dynamics during thermal

acclimation using cnidarian-dinoflagellate model holobionts. *Marine Environmental*

Research, *130*, 303–314. https://doi.org/10.1016/j.marenvres.2017.08.005

Oliver, T. a., & Palumbi, S. R. (2011). Do fluctuating temperature environments elevate coral

thermal tolerance? *Coral Reefs*, *30*(2), 429–440. https://doi.org/10.1007/s00338-011-0721-y

Oliver, T. A., & Palumbi, S. R. (2011). Many corals host thermally resistant symbionts in high-

temperature habitat. *Coral Reefs*, *30*, 241–250. https://doi.org/10.1007/s00338-010-0696-0

Polato, N. R., Voolstra, C. R., Schnetzer, J., DeSalvo, M. K., Randall, C. J., Szmant, A. M.,

Medina, M., & Baums, I. B. (2010). Location-specific responses to thermal stress in larvae

of the reef-building coral Montastraea faveolata. *PloS One*, *5*(6), e11221.

https://doi.org/10.1371/journal.pone.0011221

Purkis, S. J., & Riegl, B. M. (2012). Geomorphology and Reef Building in the SE Gulf. In *Coral*

Reefs of the Gulf: Adaptation to Climatic Extremes in the World's Hottest Sea (pp. 33–50).

https://doi.org/10.1007/978-94-007-3008-3

Riegl, B., Purkis, S., Al-cibahy, A., Abdel-moati, M., & Hoegh-Guldberg, O. (2011). Present

Limits to Heat-Adaptability in Corals and Population-Level Responses to Climate

Extremes. *PloS One*, *6*(9). https://doi.org/10.1371/journal.pone.0024802

Ruiz-jones, L. J., & Palumbi, S. R. (2017). Tidal heat pulses on a reef trigger a fine-tuned

transcriptional response in corals to maintain homeostasis. *Science Advances*, *3*(e1601298),

1–11.

Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local adaptation.

Nature Reviews Genetics, *14*(November), 807–820. https://doi.org/10.1038/nrg3522

Silverstein, R. N., Cunning, R., & Baker, A. C. (2014). Change in algal symbiont communities

after bleaching, not prior heat exposure, increases heat tolerance of reef corals. *Global*

Change Biology, 236–249. https://doi.org/10.1111/gcb.12706

Singh, D., Singh, C. K., Sewak, R., Tomar, S., & Pal, M. (2017). Genetics and Molecular

Mapping of Heat Tolerance for Seedling Survival and Pod Set in Lentil. *Crop Science*, *57*,

3059–3067. https://doi.org/10.2135/cropsci2017.05.0284

Thomas, L., López, E. H., Morikawa, M. K., & Palumbi, S. R. (2019). Transcriptomic resilience

, symbiont shuffling , and vulnerability to recurrent bleaching in reef - building corals.

Molecular Ecology, *28*, 3371–3382. https://doi.org/10.1111/mec.15143

Traylor-knowles, N., Rose, N. H., & Palumbi, S. R. (2017). The cell specificity of gene

expression in the response to heat stress in corals. *Journal of Experimental Biology*, *220*,

1837–1845. https://doi.org/10.1242/jeb.155275

Traylor-knowles, N., Rose, N. H., Sheets, E. A., & Palumbi, S. R. (2017). Early Transcriptional

Responses during Heat Stress in the Coral Acropora hyacinthus. *Biological Bulletin*, *232*,

91–100.

van Oppen, M. J. H., Oliver, J. K., Putnam, H. M., & Gates, R. D. (2015). Building coral reef

resilience through assisted evolution. *Proceedings of the National Academy of Sciences*,

112(8), 2307–2313. https://doi.org/10.1073/pnas.1422301112

Venuprasad, R., Dalid, C. O., Del Valle, M., Zhoa, D., Espiritu, M., Sta Cruz, M. T., Amante,

M., Kumar, A., & Atlin, G. N. (2009). Identification and characterization of large-effect

quantitative trait loci for grain yield under lowland drought stress in rice using bulk-

segregant analysis. *Theoretical Applied Genetics*, *120*, 177–190.

https://doi.org/10.1007/s00122-009-1168-1

Vikram, P., Swamy, B. P. M., Dixit, S., Ahmed, H., Cruz, M. T. S., Singh, A. K., Ye, G., &

Kumar, A. (2012). Field Crops Research Bulk segregant analysis : " An effective approach

for mapping consistent-effect drought grain yield QTLs in rice ." *Field Crops Research*,

134, 185–192. https://doi.org/10.1016/j.fcr.2012.05.012

Wang, M., Liu, W., Jiang, B., & Peng, Q. (2019). Genetic Analysis and Related Gene Primary

Mapping of Heat Stress Tolerance in Cucumber Using Bulked Segregant Analysis.

Horticulture Science, *54*(3), 423–428. https://doi.org/10.21273/HORTSCI13734-18

Wang, S., Meyer, E., McKay, J., & Matz, M. (2012). 2b-RAD: a simple and flexible method for

genome-wide genotyping. *Nature Methods*, *9*(August), 808–810.

http://www.nature.com/nmeth/journal/v9/n8/abs/nmeth.2023.html

Wilson, A. J., Clements, M. N., Morrissey, M. M., Re, D., Postma, E., Walling, C. A., Kruuk, L.

E. B., & Nussey, D. H. (2010). An ecologist ' s guide to the animal model. *Journal of*

Animal Ecology, *79*, 13–26. https://doi.org/10.1111/j.1365-2656.2009.01639.x

Woolsey, E. S., Keith, S. a., Byrne, M., Schmidt-Roach, S., & Baird, a. H. (2014). Latitudinal

variation in thermal tolerance thresholds of early life stages of corals. *Coral Reefs*, *34*, 471–

478. https://doi.org/10.1007/s00338-014-1253-z

Tables

Table 1. Estimates of narrow-sense heritability of thermal tolerance obtained from different survival metrics. Censored and uncensored KM estimates refer to cumulative probability of survival estimated with or without censoring individuals surviving at the experimental endpoint.

Data	Survival	h^2
Survival at 240hr	59%	0.66
KM Estimate	59%	0.64

Table 2. The number of loci at which allele frequencies were significantly altered in survivors of thermal stress in each cross.

Cross	Significant markers	Markers genotyped	Parental distance	Survival (10 d)
AB	76	15,217	0.247	0.637
AF	34	19,399	0.245	0.551
BA	95	11,616	0.247	0.742
BC	69	22,820	0.257	0.442
CA	75	23,173	0.246	0.406
DB	44	23,999	0.256	0.758
DC	336	17,548	0.246	0.483
EC	77	16,734	0.262	0.683
EF	124	13,904	0.257	0.654
FA	139	16,691	0.245	0.639

Table 3. Markers identified in four or more crosses.

Marker	SNP Location	Marker Found in Crosses	Number Crosses
denovoLocus34739	1, 21	AB, AF, BC, CA, DC, EF, FA	7
denovoLocus44191	16, 17	AB, AF, BA, CA, DB, EF, FA	7
denovoLocus9359	1	AB, BC, CA, DC, EF, FA	6
denovoLocus121909	11	AB, BA, CA, EC, FA	5
denovoLocus1291	4, 6	AB, BA, BC, DB, EF	5
denovoLocus22543	1, 25	BA, CA, EC, EF, FA	5
denovoLocus93322	3, 25	AB, AF, DC, EC, EF	5
denovoLocus13062	26, 36	CA, EC, EF, FA	4
denovoLocus18723	1	BA, DB, DC, EC	4
denovoLocus29413	1	BC, DC, EF, FA	4
denovoLocus36225	1, 4	AB, AF, EC, EF	4
denovoLocus51123	4	AB, AF, DC, EF	4

Table 4. SNP markers affected by heat stress that map to transcripts. Listed in order of the number of crosses the SNP marker was found.

Markers	Crosses	Transcripts	Descriptions (UniProt match)
denovoLocus32940	AF, DC, EF	c31022_g1_i1	Not annotated (no match)
denovoLocus14433	BA, DC	c40970_g1_i1	Not annotated (no match)
denovoLocus24976	DC, EF	c17319_g1_i1	Not annotated (no match)
denovoLocus55443	DB, DC	c19185_g1_i1	Not annotated (no match)
denovoLocus31374	FA	c12147_g2_i1	GPI mannosyltransferase 2 (U3DAR7)
denovoLocus2995	EC	c8644_g1_i2	Putative cad96ca fragment (L7MF99)
denovoLocus4290	AB	c36043_g1_i1	Metalloendopeptidase (A7RVF7)
denovoLocus7677	DC	c16415_g1_i3	S/T-protein kinase PAK 3 (T2MHJ0)
denovoLocus22915	AF, BA, CA	c15457_g1_i1	Isocitrate dehydrogenase 1 (E9C6G6)

696   **Figures**

699   **Fig. 1. Crossing design and heat stress experiments.** (**A**) Six *Platygyra daedalea* colonies

700   were sampled from Esk Reef on the GBR. (**B**) Crossing design used to produce ten crosses

701   among those parental colonies. (**C**) Representative micrograph (10X) of planula larvae used for

702   these experiments (53 hours post-fertilization).

Fig. 3. Heat tolerance differs among crosses. Kaplan Meier plots of cumulative probability of

survival for each family in control 27.5°C (blue) and elevated 35.5°C (red) temperature. Panels

are arranged in order of heat tolerance, based on survival probability at 240 hr. Shaded regions

represent 95% confidence intervals.

Fig. 4. Heat stress alters allele frequencies in surviving larvae. Panels show correlations

between the average allele frequencies in survivors of replicate heat stress treatments relative to

the corresponding controls. Markers with significantly altered allele frequencies (logistic

regression, FDR < 0.05) are highlighted in red.

Fig. 5. Venn diagrams depicting the overlap of genetic markers in sets of three crosses

Many markers affected by heat stress were specific to each cross, but a subset of markers showed

consistent changes in allele frequency during heat stress across multiple crosses. Examples

shown include (**a**) crosses EF, AF, and AB; and (**b**) crosses DC, EF, and FA. The Venn diagram

(**a**) shows shared markers between the three crosses that had the most shared markers and (**b**)

shows shared markers between three crosses with only one shared parent. These comparisons

demonstrate independent associations between heat stress and genotype at these loci across

different genetic backgrounds.