Defects in plant immunity modulate the rates and patterns of RNA virus evolution

Rebeca Navarro[1,&], Silvia Ambrós[1,&], Anamarija Butković[1], José L. Carrasco[1],

Rubén González[1], Fernando Martínez[1], Beilei Wu[1,$], and Santiago F. Elena[1,2,*]

[1]Instituto de Biología Integrativa de Sistemas (CSIC - Universitat de València), Paterna,

46182 València, Spain.

[2]The Santa Fe Institute, Santa Fe NM87501, USA.

[*]Correspondence, e-mail: santiago.elena@csic.es.

[&]Equal contribution.

[$]Permanent address: Institute of Plant Protection, Chinese Academy of Agricultural

Sciences, Haidian, 100193 Beijing, China.

Keywords: experimental evolution; generalism; plant immunity; plant-virus interactions;

response to infection; specialization; virus evolution

Abstract

It is assumed that host genetic variability for susceptibility to infection necessarily conditions virus evolution. Differences in host susceptibility can either drive the virus to diversify into strains that track different defense alleles (*e.g.*, antigenic diversity) or to infect only the most susceptible genotypes. To clarify these processes and their effect on virulence, we have studied how variability in host defense responses determine the evolutionary fate of viruses. To accomplish this, we performed evolution experiments with *Turnip mosaic potyvirus* in *Arabidopsis thaliana* mutants. Mutant plants had disruptions in infection-response signaling pathways or in genes whose products are essential for potyvirus infection. Genotypes were classified into five phenogroups according to their response to infection. In order to disentangle how host susceptibility affects virus adaptation, independent viral lineages were evolved in selected plant genotypes. Evaluating disease-related traits of the evolved lineages, we found that evolution proceeded faster in the most resistant hosts than in the most permissive ones, as expected for adaptation to a harsh environment. By sequencing the genomes of the evolved viral lineages, we found that the multifunctional protein VPg turned out to be the target of selection in most host genotypes. When all evolved viral lineages were tested for fitness in all plant genotypes used in the experiments, we found compelling evidences that generalist viruses were selected by the most restrictive plant genotypes, while permissive genotypes selected for specialist viruses. Overall, this work illustrates how different host defense signaling pathways constrain not only disease-related traits but virus evolution.

Introduction

The spectrum of disease severity can be attributed to heterogeneity in virus virulence or in host factors; the two are not necessarily independent explanations and they must actually complement and/or interact with each other resulting in an arms-race coevolutionary process. A problem faced by viruses is that host populations consist of individuals that had different degrees of susceptibility to infection (Schmid-Hempel and Koella 1994; Pfenning 2001; Sallinen et al. 2020). Therefore, adaptive changes improving viral fitness in one host genotype may be selected against, or be neutral, in an alternative one. Genetic variability in susceptibility of hosts and infectiousness of viruses have been well studied in animals and plants (*e.g.*, Schmid-Hempel and Koella 1994; Altizer 2006; Hughes and Boomsma 2006; Brown and Tellier 2011; Anttila et al. 2015; Parrat et al. 2016; González et al. 2019; Sallinen et al. 2020). The interaction between host and parasite genotypes has been explained in the light of two different theoretical models that represent the two extremes on a continuum of possibilities. At the one extreme, the so-called gene-for-gene (GFG) model, in which a virus genotype exists that can infect all host genotypes and a universally susceptible host genotype should also exist (Flor 1956). Resistance occurs when a host "resistance" gene is matched by at least one virus' "avirulence" gene. Polymorphism in infectivity and resistance can be maintained only if virulence pays a cost. At the other extreme, the matching-alleles (MA) model is based on self- versus non-self-recognition systems in invertebrates. Infection is not possible unless the virus possesses all alleles that match those of the host (Frank 1993). In this case, polymorphism in infectivity and resistance are maintained by negative frequency-dependent selection.

Variability within host populations arises from nonrandom spatial distributions of genotypes: social groups of animals are more closely related to each other than to other members of the population, and plant crops are generally cultivated as genetically homogeneous plots. These situations facilitate virus transmission among genetically similar host genotypes. Spatial structure and local migration predict evolution of less aggressive exploitation in horizontally transmitted parasites (reviewed in Parrat et al. 2016). For example, Boots and Mealor (2007) observed that *Plodia interpunctella* granulosis virus evolved in spatially structured lepidopteran host populations became less virulent than the one maintained in well-mixed host populations. Berngruber et al. (2015) showed that a latent λ bacteriophage won competitions against a virulent one in a spatially structured host populations but lost in well-mixed populations. Similar results were

observed in laboratory evolution experiments with *Arabidopsis thaliana* and tobacco etch virus (TEV): a negative association between the permissiveness to infection of different plant accessions and TEV virulence was observed (Hillung et al. 2014). Finally, host population structure also promotes coexistence of hosts and parasites reducing the pathogens dispersal (Brockhurst et al. 2006).

Theoretically, in the absence of host heterogeneity, parasites must evolve toward a host exploitation strategy that maximizes transmission with low virulence (Haraguchi and Sasaki 2000; Regoes et al. 2000; Rodríguez and Torres-Sorando 2001; Ganusov et al. 2002; Gandon 2004; Lively 2010; Moreno-Gámez et al. 2013). Chabas et al. (2018) experimentally showed that evolutionary emergence is more likely when the host population contains intermediate levels of resistant hosts, confirming this prediction using a set of phages and bacterial hosts with alterations in the CRISPR/Cas-mediated immunity.

Viral infection of plants is a complex system in which the virus parasitizes the host and utilizes all its cellular resources to replicate and systemically spread. In response, plants have evolved intricated signaling mechanisms that limit the spread of the virus, resulting in resistance (Zhou and Zhang 2020). A variety of factors contribute to plant resistance to viral infections. Broadly speaking, these factors can be classified into *basal* if they are pre-existing and limit within-cell propagation and cell-to-cell spread, and *inducible* if they are only activated upon infection and inhibit systemic virus movement and replication. Basal mechanisms include susceptibility (S) genes that involve alleles of cellular proteins that do not interact properly with viral factors; *e.g.*, translation initiation factors required by the virus for successful exploitation of the cell's protein synthesis machinery, heat shock proteins that assist the formation of multiprotein complexes, or DNA binding phosphatases (Carr et al. 2010; Mäkinen 2019). In contrast, inducible mechanisms include genes whose expression results in a broad-scale change in plant physiology via diverse signal transduction pathways, particularly those regulated by the hormones salicylic acid (SA), jasmonic acid (JA) and ethylene (ET) (Soosaar et al. 2005; Carr et al. 2010). These changes include local cell apoptosis (*e.g.*, hypersensitive responses - HR; Loebenstein 2009), the upregulation of nonspecific responses against many different types of pathogens throughout the entire plant (systemic acquired resistance - SAR- and induced systemic resistance - ISR) (Kachroo et al. 2006; Carr et al. 2010), and the activation of the RNA-silencing-based resistance, which seems to play a role both in basal and inducible mechanisms (Voinnet 2001; Carr et al. 2010). Early host

responses following virus detection include changes in ion fluxes (mainly Ca^{++}), activation of signaling pathways, major alterations of transcriptomic profiles, generation of reactive oxygen species (ROS) and production of nitric oxide (NO) (Soosaar et al. 2005). These immediate changes are followed by HR and the recruitment of SA and JA/ET signaling pathways. The SA-mediated defense signaling pathway results in SAR, while the JA/ET-mediated defense signaling pathway results in ISR, the latter being specifically involved in interactions between plants and beneficial microbes. Indeed, it appears that ISR is not effective against infection by most viruses (Ton et al. 2002; Loebenstein 2009; Pieterse et al. 2009). Both SAR and ISR pathways converge into two master regulators, the *ENHANCED DISEASE SUSCEPTIBILITY 1* (*EDS1*) and the *PHYTOALEXIN DEFICIENT 4* (*PAD4*) genes; *EDS1* and *PAD4* repress ISR and promote SAR. Although the SA and JA/ET pathways have been classically viewed as mutually antagonistic, several studies have revealed positive and negative crosstalk between them (van Wees et al. 2000; Pieterse et al. 2012) as well as with the RNA-silencing pathway (Soosaar et al. 2005; Carr et al. 2010; Yang et al. 2020). This crosstalk, which has the network topological structure of an incoherent feed-forward loop, creates robustness and tunability in the plant immune network (Mine et al. 2017).

An open question is how the intricacies of the underlying defense regulatory pathways of *A. thaliana*, the existence of multiple defense responses and their potential crosstalk determines the evolutionary fate of viruses. For instance, would two independent defense pathways select for specialist viruses adapted to counteract each one thus conforming a fitness tradeoff? Would a tradeoff in the investment of plant resources between pathways (*e.g.*, SAR *vs* ISR) result in a diversification of virus adaptive strategies? Would viruses targeting crosstalk points across pathways evolve as generalists? Here we will use turnip mosaic virus (TuMV; species *Turnip mosaic potyvirus*, genus *Potyvirus*, family *Potyviridae*) - *A. thaliana* experimental pathosystem to explore these questions. After exploring the variability in phenotypic responses to TuMV infection of a collection of *A. thaliana* mutants in basal and inducible resistance mechanisms, we choose nine mutants which cover the entire spectrum in phenotypic responses. Then, we performed evolution experiments on each plant mutant genotype and track the evolution of several disease-related traits. At the end of the evolution experiment, we evaluated the effect of host genotypes in the rates of virus evolution and the contribution of historical contingency, selection and stochasticity into the outcome of evolution. Next, we explored whether the different TuMV lineages evolved as specialist

146 or generalists depending on the disrupted defense mechanism of their local hosts. Finally,

147 we sought to identify the molecular changes experienced in the genome of the different

148 viral lineages and to explore the possible adaptive value of a few convergent mutations.

Results

Classification of *A. thaliana* genotypes according to their phenotypic response to

TuMV infection

The 22 *A. thaliana* mutant genotypes used in this study are shown in table 1, including

information about the affected signaling pathways or cellular processes as well as the

expected phenotype of infection relative to wildtype (WT) plants based on the description

of these mutants. Genotypes were classified according to their phenotypic response to

TuMV infection based in five different disease-related traits measured 18 days post-

inoculation (dpi). The disease-related traits measured were change in dry weight (ΔDW),

disease progression (area under the disease progress step curve or *AUDPS*), infectivity

(*I*), symptoms severity (*SS*), and viral load (*VL*). Figure 1 shows the nearest-neighbor

clustering of genotypes according to their multi-trait phenotypic response to TuMV

infection (supplementary figure S1, Supplementary Material online); we found five

significant phenogroups (hereafter named as G1 to G5). For all four members of G1 the

response was consistent with an enhanced resistance response to TuMV infection: no

significant ΔDW whilst a significant and consistent reduction in *AUDPS*, *I*, *SS*, and *VL*

relative to infected WT plants. Three members of G1 had enhanced SAR response (table

1) and the fourth one (*i4g2*) is a well-known *S* gene involved in plant resistance to

potyviruses (Nicaise et al. 2007; Gallois et al. 2010). The only member of G2 was the

strong apoptosis-inducer *p58^{IPK}*, which shows no significant changes in *AUDPS*, *I*, *SS*,

and *VL* relative to infected WT plants but a significant reduction in ΔDW. Infected *p58^{IPK}*

plants were heavier than the noninfected controls which points to a case of increased

tolerance to infection. The only member of G3 was *dbp2*, that specifically increased

resistance against another potyvirus (Castelló et al. 2011). This mutant showed a very

interesting response to TuMV infection: ΔDW was significantly increased while *AUDPS*,

I and *SS* were significantly reduced; no effect in *VL* was observed. Thus, *dbp2* represents

an intermediate response to TuMV infection between the highly resistant G1 and those

genotypes in G4. G4 represents a sort of hotchpotch formed by genotypes that do not

show a clear phenotypic difference from WT. Remarkably, though, with the exception of *cpr5-2*, *dcl2 dcl4*, and *dip2*, all show significant reductions in *VL*. Finally, *jin1* was the only member of G5, which shows a significant increase in all measured traits except *VL*, consistent with and enhanced susceptibility to TuMV infection. This is a surprising finding since *jin1* has been described as inducing a constitutive expression of the SA-dependent defenses that made plants extremely resistant to the infection of biotrophic bacterial pathogens (Laurie-Berry et al. 2006). To further characterize the progression of infection, we selected a subset of representatives from groups G1 (*eds8-1*) and G4 (*cpr5-2*, *dcl2 dcl4*, *hsp90-1*, and WT) and evaluated daily *I*, *SS* and *VL* (supplementary figure S2, Supplementary Material online). The *cpr5-2* plants started showing symptoms faster, though only 80% of plants showed symptoms after 15 dpi. Only 20% of *eds8-1* plants showed symptoms of infection, which were delayed. Infected *hsp90-1* and *dcl2 dcl4* plants were undistinguishable from the infected WT ones. Regarding *VL*, all genotypes accumulated less virus than WT plants, amongst them *cpr5-2* accumulated the least amount of virus (supplementary figure S2, Supplementary Material online).

Representatives from the five groups were selected for the subsequent evolution experiment (shadowed rows in table 1): *eds8-1* and *i4g2* from G1 representing the hardest selection regimes (more resistant genotypes); *p58^{IPK}* from G2; *dbp2* from G3; *cpr5-2*, *dcl2 dcl4*, *hsp90-1*, *npr1-1*, and WT from G4; and *jin1* from G5 representing the softest selective regime (less resistant genotype).

Experimental evolution in representative *A. thaliana* genotypes

First, we sought to evaluate the dynamics of evolution for *AUDPS* (figure 2), *I* (figure 3) and *VL* (figure 4). Data in these figures were fitted all together into the multivariate analysis of covariance (MANCOVA) model described in equation 1 of the Materials and Methods section. Notice that lineage *cpr5-2*/L2 showed a quite different evolution pattern from the other four lineages evolved in *cpr5-2*. It never increased in either of the three traits and was lost after passage five (figures 2 and 3). Therefore, we ended up with 46 evolved lineages (two in the WT, four in *cpr5-2* and five in each of the other eight plant genotypes).

Table 2 shows the results of this MANCOVA. Despite a considerable amount of noise in the time series, we found some significant results that can be summarized as follows. First, a net significant and large ($\eta_P^2 = 0.735$) effect associated to the evolutionary passages (*t*) has been observed, indicating that the three phenotypic traits

related to TuMV infection evolved during the experiment. Second, plant genotypes (G)

had a highly significant and large ($\eta_P^2 = 0.547$) effect on the phenotypic traits, suggesting

that TuMV evolutionary dynamics were strongly influenced by the local host genotype

in which it was being passaged. More interestingly, this effect was strongly dependent

on the passage number ($t \times G$; $\eta_P^2 = 0.463$), thus suggesting that the slope of the relationship

between phenotypic traits and time was influenced by the local host genotype. This effect

will be further evaluated in the next section. Third, independent lineages evolved in the

same host genotype show a strong degree of evolutionary parallelism, as indicated by

nonsignificant differences among viral lineages within local host genotype (factors $L(G)$

and $t \times L(G)$). However, this last conclusion should be taken with certain caution since the

power of the two tests is relatively low ($1 - \beta < 0.800$) and hence we might be wrongly

failing to reject the null hypothesis (type II error).

A relevant question in evolutionary biology is the extent in which ancestral

differences determine the fate of evolution. In other words, the relative contribution of

adaptation, chance and historical contingency to the evolution or organismal fitness.

Following the logic exposed by Travisano et al. (1995), we can imagine two situations.

First, ancestral statistical differences among phenotypes could be preserved along

evolution despite a net increase in the mean trait values (due to selection) and differences

among replicated lineages (due to chance; $i.e.$, mutation and drift). In this situation, we

should expect a nonzero slope in a regression of the evolved phenotypic values against

the ancestral ones. The closer the slope to a value of one, the more importance of ancestral

differences. Second, if initial trait variation among ancestral genotypes was eliminated

from the evolved populations because the combined effect of adaptation and chance, we

should expect a regression slope smaller than one. The less effect of ancestral differences,

the flatter the slope, being zero in the extreme case where ancestral differences are

completely erased. To evaluate the role of the ancestral differences in $AUDPS$, I and VL

of TuMV across the nine different plant genotypes, we evaluated the magnitude of their

change at the end of the evolution experiment. Figure 5 shows the plots of evolved vs

ancestral values. For illustrative purposes, the solid diagonal lines represent the null

hypothesis of absolute preservation of ancestral differences. In the case of $AUDPS$ (figure

5A), a significant regression exists between evolved and ancestral values ($R = 0.491$, $F_{1,44}$

$= 14.000$, $P < 0.001$), with slope 0.284 ± 0.076 (± 1 SD). Since the slope is still

significantly different from zero, yet clearly flatter than the diagonal ($t_{44} = 9.422$, $P <$

0.001), we conclude that ancestral differences have been mostly removed by the combined action of selection and chance, yet not completely erased. In the case of I (figure 5B), the regression was not significant ($R = 0.094$, $F_{1,45} = 0.405$, $P = 0.528$), thus suggesting that for this trait ancestral differences among $A.$ $thaliana$ genotypes have been fully erased by adaptation and chance. Finally, in the case of VL (figure 5C; WT not included in this analysis), the regression of evolved values on the ancestral ones was significant ($R = 0.417$, $F_{1,43} = 8.822$, $P = 0.005$), suggesting again that ancestral phenotypic differences have been largely removed by selection and chance yet some still persist (regression slope -0.142 ± 0.048 significantly different from one: $t_{42} = 23.903$, $P < 0.001$).

Rates of phenotypic evolution for disease-related traits

As mentioned in the previous section, we have observed a significant effect of $A.$ $thaliana$ genotypes in the temporal evolution of the three phenotypic traits. To further explore this effect, we estimated the rates of phenotypic evolution of $AUDPS$ and I using the ARIMA(1,0,0) model described by equation 2 in Materials and Methods. VL was not included in this analysis since the number of data points (three) in the time series was the same as the number of parameters to be estimated, thus likely incurring in overfitting problems.

Figure 6 shows the estimated rates of evolution (ν) for host genotypes ordered from the most resistant (G1) to the most permissive (G5) ones. Very interestingly, a significant negative regression coefficient exists for both traits [$AUDPS$: $-1.313 \cdot 10^{-2} \pm 0.383 \cdot 10^{-2}$ ($R = 0.459$, $F_{1,44} = 11.724$, $P = 0.001$) and I: $-2.576 \cdot 10^{-3} \pm 0.790 \cdot 10^{-3}$ ($R = 0.4413$, $F_{1,44} = 10.621$, $P = 0.002$)], suggesting that evolution always proceeded faster in the most restrictive hosts and slower in the most permissive ones.

Dissecting the role of specific defenses in TuMV evolution

The rates of TuMV phenotypic evolution estimated in the previous section were further analyzed in the context of the mutated defense signaling pathways or recessive resistances. Rates of evolution ν_{AUDPS} and ν_I were fitted to the MANOVA model shown in equation 3 of the Materials and Methods section. Figure 7 graphically summarizes the results of these analyses. First, we sought for differences in rates of evolution between permissive (soft selection) and restrictive (hard selection) hosts (figure 7A). The

multivariate analysis showed that lineages evolved in the more restrictive hosts evolved

significantly faster than those evolved in the more permissive hosts ($\Lambda = 0.780$, $F_{2,43} =$

6.080, $P = 0.005$). This effect was mostly driven by differences in the rate of infectivity

evolution v_I, as shown by the corresponding univariate analyses, being on average 3.14

times faster in the most restrictive hosts. This observation confirms the results shown in

the previous section.

Next, we evaluated whether differences exist between the rates of phenotypic

evolution for viruses evolved in host genotypes carrying mutations affecting basal or

inducible resistances (figure 7B; WT data for this analysis excluded). In this case, the

multivariate analysis found a small, yet significant effect ($\Lambda = 0.790$, $F_{2,34} = 4.507$, $P =$

0.018), though none of the two univariate analyses had enough power to detect it (indeed,

changes went in opposite directions: 1.02-fold larger for v_{AUDPS} in the plants with

mutations in basal resistances but 1.31 times larger for v_I in the plants with mutations in

the inducible resistances).

Figure 7C shows the comparation of rates of TuMV phenotypic evolution for

lineages evolved in plant genotype mutants for the SA-dependent signaling defense

pathways. In this case we have three categories compared to WT: mutations inducing a

down-expression of the pathway, mutations with no effect on this pathway and mutations

resulting in overexpression. Again, the MANOVA found highly significant differences

among the three categories ($\Lambda = 0490$, $F_{4,84} = 8.996$, $P < 0.001$). The univariate analyses

showed that most of the differences were due to a 2.38 times faster evolution of I in the

host genotypes over-expressing the SAR defenses, than in those with a WT-like or even

down-expression of the SA signaling which results in a reduced SAR.

SA- and JA/ET-dependent pathways have been considered as antagonistic defenses

(see Introduction), although more recent evidence points toward an active crosstalk

between both (van Wees et al. 2000; Pieterse et al. 2012; Mine et al. 2017). Under the

antagonistic hypothesis, we should expect the opposite trend among lineages evolved in

plants with affected JA/ET signaling than the one observed for the lineages evolved in

plants with affected SA signaling (figure 7C). Figure 7D shows the data of evolution

rates classified according to whether host genotypes had mutations affecting the JA/ET

defense signaling pathway that resulted ISR. The MANOVA found a significant

difference between the two categories ($\Lambda = 0.766$, $F_{2,43} = 6.580$, $P = 0.003$), entirely

driven by differences in v_I, as confirmed by the univariate analyses. Indeed, as expected,

the rates of evolution for infectivity were 2-fold faster for TuMV lineages evolved in the ISR deficient hosts, in agreement with the SAR/ISR antagonistic hypothesis.

Finally, we tested the effect of knocking down the host's RNA-silencing pathway in the rates of TuMV evolution (figure 7E). In this case, the multivariate analysis also found highly significant differences between rates of viral evolution in plants with a fully functional RNA-silencing pathway and the silencing-deficient *dcl2 dcl4* mutant ($\Lambda = 0.850$, $F_{2,43} = 3.801$, $P = 0.030$). The average ν_I values were 21-fold larger in plants expressing a fully functional RNA-silencing pathway than in *dcl2 dcl4* plants unable of siRNA production.

In conclusion from this section, we found that regardless of the defense signaling pathway, viruses evolved in permissive plants always evolve slower than their counterparts evolved in more restrictive plant genotypes.

Permissiveness of *A. thaliana* genotypes to infection drives the evolution of TuMV lineages into specialists or generalists

To analyze the specificity of adaptation of each evolved TuMV lineage, we performed a full cross-infection experiment. In this experiment, the 44 lineages evolved in the mutant genotypes (excluding the two WT-evolved lineages) were inoculated into ten plants of the nine *A. thaliana* mutant genotypes used in the evolution experiments (excluding WT plants). The presence of symptoms in each plant was recorded daily for up to 12 dpi. This dataset was analyzed using two different approaches, first an ANOVA-like that evaluates the effect of host genotypes, viral lineages and their interactions (Schmid-Hempel 2011); and second, the inference and characterization of nestedness, modularity and specialization of a bipartite infection network (Weitz et al. 2013). For the first approach we fitted the presence/absence of symptoms data to the logistic regression model shown in equation 4 of the Materials and Methods section. The results are shown in table 3. A test host genotype (*TG*) factor describes the effect of differences among host genotypes, irrespective of the infecting viral lineage; and thus, characterizes whether some host genotypes are more susceptible than others to infection. As shown in table 3, *TG* has a highly significant effect, both by itself and in the interaction with *t*. The magnitude of the net effect was very large ($\eta_P^2 = 0.569$), while its interaction with *t* was rather small in magnitude ($\eta_P^2 = 0.010$). On average, the most susceptible host genotypes were *jin1*, *dcl2 dcl4* and *cpr5-2* ($\bar{f} = 1.000$), whereas the most resistant host genotypes

were $p58^{IPK}$ ($\bar{f}$ = 0.459 ±0.013) and *eds8-1* ($\bar{f}$= 0.233 ±0.013).   Similarly, a highly

significant local host genotype (*LG*) effect exists by itself (of large magnitude η_P^2 = 0.288)

as well as in combination with *t* (though the magnitude of the interaction was rather small

in magnitude η_P^2 = 0.005), which means that the observation depends on the viral lineage

independently of the host genotype.   On average, the lineages evolved in *eds8-1* ($\bar{f}$ =

0.997 ±23.710) and *dbp2* ($\bar{f}$ = 0.994 ±26.108) induced symptoms faster; while in contrast,

lineages evolved in *i4g2* ($\bar{f}$ = 0.788 ±279.090) showed symptoms much slower.   Finally,

and most interestingly, a highly significant interaction term *TG×LG* (as well as in

combination with *t*) indicates that the outcome depends on the particular combinations of

host genotypes and TuMV lineages.   Indeed, the magnitude of the interaction effect was

large (η_P^2 = 0.314), though its dependence with dpi was not so relevant (η_P^2 = 0.048).   For

example, in the most susceptible genotypes, where *cpr5-2* perfectly illustrates this case:

lineages evolved in *cpr5-2*, *dbp2*, *dcl2 dcl4*, *eds8-1*, and $p58^{IPK}$ all show $\bar{f}$ = 1.000,

whereas lineages evolved in *i4g2* and *jin1* show much lower $\bar{f}$ values (0.674 ±0.028 and

0.849 ±0.032, respectively).   In the opposite extreme, *e.g.*, *eds8-1*-evolved lineages show

$\bar{f}$ = 0.575 ±0.050 while *jin1*-evolved ones show $\bar{f}$ = 0.124 ±0.030; while all other lineages

show intermediate $\bar{f}$ values.

For the second analytical approach, we used the full cross-infection dataset to

estimate the corresponding *AUDPS* values, thus creating an infection matrix of 44 rows

[*L*(*LG*)] by nine columns (*TG*).   This matrix was transformed into a new binary matrix in

which a value of 1 meant that the *AUDPS* for a given TuMV lineage, in a particular *TG,*

was greater or equal than the one observed for its *LG* and 0 otherwise.   Figure 8A shows

the packed infection matrix with rows and columns organized to better highlight its

nestedness and modularity (Weitz et al. 2013).   The matrix could be also represented as

a bipartite infection network (figure 8B) that provides the same information but in a more

graphical way.   The infection network was significantly nested (T = 14.706, P < 0.001),

with the most permissive genotypes *jin1* and *dcl2 dcl4* being susceptible to most viral

lineages (universally susceptible hosts), while the most resistant genotype *eds8-1* and

$p58^{IPK}$ were only successfully infected by some lineages that were precisely evolved in

these two genotypes.   Likewise, the most generalist viral lineages were *eds8-1*/L4,

$p58^{IPK}$/L3 and $p58^{IPK}$/L4 that infected the nine mutant host genotypes (figure 8) with

equal efficiency (universally virulent viruses).   On another hand, the most specialized

viral lineages that were able to successfully infect only their local hosts, were all surviving

lineages evolved in *cpr5-2*, four lineages evolved in *dcl2 dcl4* and four lineages evolved
in *jin1* (figure 8). Therefore, we conclude that local host's resistance to infection
positively correlates with the host range of the evolved lineages: more resistant host
genotypes selected for very generalist viruses while less resistant host genotypes selected
for very specialized viruses, in agreement with a GFG infection model (Weitz et al. 2013).

Next, we computed Blüthgen et al. (2006) specialization indexes for the packed
matrix (figure 8). Firstly, we evaluated the standardized species-level measure of partner
diversity *d'*. *d'* ranged from zero for the most generalist TuMV evolved lineages (*eds8-*
1/L4, *p58^{IPK}*/L3 and *p58^{IPK}*/L4) to one for the most specialist ones (the four surviving
cpr5-2-evolved lineages, *dbp2*/L2, *dcl2 dcl4*/L2 - L5 and *jin1*/L1 - L4) (last column in
figure 8A). Secondly, we computed the network-level specialization index H'_2, obtaining
a value of $H'_2 = 0$ (uncorrected $H_2 = 5.024$, $H'_{min} = 3.534$, $H'_{max} = 5.024$; $P < 0.001$),
which means that, overall, the binary infection network shows a degree of specialization.
Most of the lineages were able of infect one or a few host genotypes, a result mainly
driven by those lineages evolved in the less resistant host genotypes.

Finally, we sought to explore the modularity of the infection network. Indeed, from
an ecological standpoint, a significant network-level degree of specialization means that
some TuMV lineages do not interact well with some of the *A. thaliana* genotypes, thus
creating the possibility of modularity. In this context, a module will refer to an aggregated
set of viral lineages and their hosts characterized by more interactions within the module
than between modules (Newman 2006; Dormann and Strauss 2014). To this end, we
computed the *Q* modularity index, which ranges from zero, when the community has no
more links within modules than expected by chance, to a maximum value of one. We
found a small yet significant modularity ($Q = 0.234$, $P = 0.047$) in the infection matrix
(figure 8A). What are the implications of modularity in TuMV pathogenesis? Modularity
is expected for a MA infection model (Weitz et al. 2013). More similar host genotypes,
e.g., those with mutations affecting the same signaling pathway, may be selecting for viral
lineages with similar properties, thus being able of infect a subset of plant genotypes with
equal efficiency. An alternative hypothesis that can explain modularity may be parallel
evolution of lineages evolved into the same host genotype; the most representative cases
of this possibility are the lineages evolved in *cpr5-2*, *dcl2 dcl4* and *jin1*. To distinguish
between these two hypotheses, we computed a reduced matrix averaging the observed
AUDPS among lineages evolved in the same host genotype and transforming this nine-
by-nine infection matrix into its corresponding packed binary matrix, as described above.

If the reduced matrix still shows significant modularity, the first hypothesis will hold on,

but if modularity vanishes by averaging lineages evolved in the same host genotype, then

the alternative hypothesis will be more parsimonious. The reduced matrix is still

significantly nested ($T = 13.064$, $P < 0.001$), but modularity was not significant anymore

($Q = 0.286$, $P = 0.137$). This supports the hypothesis that the observed modularity was

driven by convergent phenotypic evolution of lineages evolved into the same host

genotype rather than by the overlap of selective pressures by genotypes carrying

mutations in the same defense mechanism.

Genomic differences/similarities among TuMV lineages and VPg as a potential

target of selection

Finally, we have explored the molecular changes experienced by the TuMV evolved

lineages. Full genomic consensus sequences were determined for 45 evolved lineages.

A total of 119 mutational events have been observed, affecting 73 nucleotide positions

(supplementary figure S3A and table S1, Supplementary Material online). According to

the type of nucleotide substitution involved, 98 were transitions and 21 transversions;

regarding their effect on the protein sequence, 30 were synonymous and 89

nonsynonymous. Interestingly, some mutations, including 13 nonsynonymous and four

synonymous, have been observed multiple times in independent lineages. As discussed

in the Materials and Methods section, treating each lineage as an observation and each

host genotype as a subpopulation, the average nucleotide diversity within host genotypes,

referred only to the 73 polymorphic sites, was $\pi_S = 0.067 \pm 0.007$. On the other hand, the

nucleotide diversity for the entire sample was $\pi_T = 0.073 \pm 0.008$. Hence, the estimated

interhost genotypes nucleotide diversity was $\delta_{ST} = 0.006 \pm 0.003$, which results in a

coefficient of nucleotide differentiation (Nei 1982) of $N_{ST} = 0.081 \pm 0.039$, a value

significantly greater than zero ($z = 2.077$, $P = 0.019$). Thus, we conclude that minor yet

significant genetic differentiation has been generated among viral lineages replicating in

different host genotypes. To assess whether selection played a role in virus genetic

differentiation among *A. thaliana* genotypes, we performed a Tajima's D test (Tajima

1989) and found that it was significantly negative ($D = -2.496$, $P = 0.006$).

Next, we sought to characterize the distribution of mutational events along the nine

non-overlapping cistrons (supplementary figure S3B, Supplementary Material online).

The frequency of mutations per cistron, relative to the length of the corresponding cistron,

was fitted to the logistic regression model shown in equation 5 of the Materials and

Methods. Highly significant differences exist ($\chi^2 = 143.206$, 9 d.f., $P < 0.001$), yet

entirely due to the ~11-fold larger mutation frequency observed in the *VPg* cistron relative

to the rest of the genome (supplementary figure S3B, Supplementary Material online).

Notice that all mutations observed in *VPg* are nonsynonymous and that all lineages except

eds8-1/L3 carry at least one mutation in this cistron (supplementary figure S3A and table

S1, Supplementary Material online).

Convergent nonsynonymous mutations are, *a priori*, good candidates for adaptive

mutations. Mutation CP/V148I appears in two lineages evolved in *cpr5-2*

(supplementary figure S3A and supplementary table S1, Supplementary Material online),

mutation CP/S70N appear in lineage *dcl2 dcl4*/L1 and three lineages evolved in *jin1*

(supplementary figure S3A and table S1, Supplementary Material online), and two

different mutations affecting CP amino acid 112, CP/D112G (lineage *npr1-1*/L4) and

CP/D112A (lineage *i4g2*/L5), that result in a similar replacement of side chains. In

addition, 10 nonsynonymous mutations in *VPg* are shared by several lineages

(supplementary figure S3A and table S1, Supplementary Material online). Out of these

10 cases, three seem particularly promising candidates. They all are located in a narrow

region of VPg (residues 113, 115 and 118) (supplementary figure S3A and table S1,

Supplementary Material online). Firstly, mutations G6237A, G6237C, A6238G, and

U6239A all affect the same codon, resulting in amino acid replacements D113N, D113H,

D113G, and D113E, respectively. The expected functional effect of these mutations was

evaluated using the algorithms implemented in SNAP2 (Hecht, Bromberg and Rost

2015). D113E is predicted to be functionally neutral (score = –76, accuracy = 87%),

while D113G (32, 66%), D113H (43, 71%) and D113N (30, 66%) are predicted to have

a functional effect. Among these three amino acid replacements, D113G, which shows

an intermediate value of the functional effect score, is particularly interesting, involving

a very strong change in the side radical from a long negatively charged one to a small

nonpolar one. This replacement has been observed in lineages *cpr5-2*/L3, *cpr5-2*/L4,

cpr5-2/L5, *npr1-1*/L1, *npr1-1*/L5, *hsp90-1*/L2, and WT/P. All four mutant hosts where

these lineages evolved belong to the phenogroup G4 (figure 1). Secondly, mutations

A6243C, A6243G and A6244G also affect the same codon, resulting in amino acid

replacements N115H (–42, 72%), N115E (–66, 82%) and N115S (–30, 61%). All three

are predicted to be functionally neutral. Thirdly, mutations U6252 and G6253A affect

the same codon and result in amino acid replacements R118C (−10, 53%) and R118H

(56, 78%), respectively. The most frequent among these two replacements is R118H,

which represents a conservative change among large positively charged side radicals, yet

has a strong expected functional effect. R118H has been observed in four out of five

TuMV lineages evolved in *jin1* (phenotypic group G5 in figure 1). D113G thus represents

a case of convergent evolution non-specific of the local host genotype (*i.e.*, a candidate

for a generalist mutation) while R118H represents a mutation highly specific of the local

host genotype (*i.e.*, a specialist mutation).

To further characterize the possible adaptive value of mutations VPg/D113G and

VPg/R118H, we created by site-directed mutagenesis the two mutated versions of *VPg*

and cloned them into the ancestral p35STunos infectious clone (Chen et al. 2003).

Viruses were recovered from these clones and three disease-related phenotypic traits (*e.g.*,

AUDPS, *I* and *SS*) evaluated both in the WT plants as well as in their corresponding local

hosts (*i.e.*, *cpr5-2* for mutant VPg/D113G and *jin1* for mutant VPg/R118H). Figure 9

summarizes the results of these experiments. Regarding VPg/D113G (figure 9A - C), the

results strongly depend on the combination of phenotypic trait and plant genotype: it

shows a significant negative effect in both *AUDPS* (−14.2%; $z = 1.687$, $P = 0.046$) and *I*

(−23.9%; $z = 6.344$, $P < 0.001$) but a positive one in *SS* (16.7%; $z = 2.597$, $P = 0.005$)

when evaluated in WT plants. However, it shows a largely negative yet not significant

effect in *cpr5-2* plants for *AUDPS* (−60.8%; $z = 1.055$, $P = 0.146$) and significant negative

effects both for *I* (−60.1%; $z = 10.614$, $P < 0.001$) and *SS* (−46.7%; $z = 3.228$, $P = 0.001$).

These negative effects on the local host genotype *cpr5-2* do not support a possible

beneficial effect of this mutation by itself. Regarding the possible adaptive effect of the

amino acid replacement VPg/R118H (figure 9D - F), the results are also dependent on the

combination of phenotypic trait and host plant genotype. When tested in WT plants, the

mutation has significant negative effects both for *AUDPS* (−11.4%; $z = 2.29$, $P = 0.013$)

and *I* (−9.0%; $z = 2.749$, $P = 0.003$) but a significantly positive effect for *SS* (43.6%; $z =$

7.881, $P < 0.001$). In contrast, in the local host *jin1* the effect was positive in all three

traits; significant for *AUDPS* (9.2%; $z = 1.683$, $P = 0.046$) and *SS* (68.2%; $z = 12.447$, P

< 0.001) but not for *I* (0.3%; $z = 1.342$, $P = 0.090$). These results support the conclusion

that the amino acid replacement VPg/R118H has an overall beneficial effect on the local

host genotype *jin1* in which it was selected.

Discussion

Interaction between TuMV and different plant defenses

In this study we have explored the effect that mutations in different host disease signaling pathways or in recessive resistance *S* genes have in the outcome of virus evolution. Our experimental pathosystem consisted of two well studied components: the model plant *A. thaliana* and one virus with high prevalence in natural populations on this host, TuMV (Pagán et al. 2010). Among the disease signaling pathways, we have studied genes involved in the SA, JA/ET and RNA-silencing; among the *S* genes, we have included heat shock proteins, transcription factors and components of the translation machinery. In a preliminary set of experiments, we compared phenotypic responses of different plant mutant genotypes to TuMV infection. Our results were, in some cases, at odds with the expected phenotypes for *R* genes, the most unexpected one being the result for *jin1*. *JASMONATE INSENSITIVE 1* (*JIN1*) is a negative regulator of SA-mediated defense responses, henceforth the mutant *jin1* has a constitutive expression of SAR (Laurie-Berry et al. 2006). Surprisingly, it turns out to be the most sensitive genotype to TuMV infection, showing enhanced symptoms of disease (phenogroup G5). This may reflect the fact that the function of genes involved in disease signaling have been mostly defined in terms of plant interactions with biotrophic and necrotrophic bacteria and fungi but rarely in response to virus infections. Mutant genotypes only affecting components of the ISR pathway (*e.g.*, *coi1-4*) behaved as WT plants in response to TuMV infection, thus confirming previous reports that ISR was inefficient against viral infections (Ton et al. 2002; Loebenstein 2009; Pieterse et al. 2009). However, mutant *eds8-1*, which avoids ISR but enhances SAR (Love et al. 2007) turned out to be among the most resistant genotypes to infection (phenogroup G1). In agreement with our findings of a variable response to SA-signaling, Singh et al. (2004) discussed examples of viruses in which SA-dependent responses to viral infection were strongly dependent on the virus used in the experiments. Replication of viruses such as alfalfa mosaic virus, potato virus X or turnip vein clearing virus was inhibited by treating plants with SA, while cucumber mosaic virus appeared unaffected by the treatment and accumulated to normal levels.

The RNA-silencing pathway is considered as the main plant defense against viral pathogens (Voinnet 2001), with *DICER LIKE 2* (*DCL2*) and *DICER LIKE 4* (*DCL4*) encoding for the two dicer enzymes responsible for generating the 22- and 21-nucleotides long antiviral siRNAs. Therefore, the double mutant *dcl2 dcl4* was expected to show a

highly sensitive response to TuMV infection. However, infected plants were, overall,

hardly distinguishable from the WT plants in their response to infection (supplementary

figures S2 and S3, Supplementary Material online). A possible explanation would be the

strong suppressor activity of HC-Pro in the WT plants that effectively counteracts the

defense mechanisms (Kaschau and Carrington 2001). Indeed, it has been shown that the

outcome of the interplay between plant RNA-silencing response and potyviruses is

mostly driven by the efficiency of the viral HC-Pro suppression activity (Lin et al. 2007;

Torres-Barceló et al. 2008).

The response to infection of genotypes carrying mutations in S genes was, in

general, consistent with the *a priori* expectation. Mutant *i4g2* belonging to phenogroup

G1, that is defective for the eIF4(iso)E factor, shows a strong resistance to TuMV

infection (Nicaise et al. 2007; Charron et al. 2008). Likewise, in agreement with previous

descriptions, mutant *dbp2* belonging to phenogroup G3, shows enhanced resistance to

potyvirus infection (Castelló et al. 2011). Finally, mutant *hsp90-1* turns out to show a

response to TuMV infection equivalent to the WT plants, likely due to its functional

redundancy with other heat-shock proteins present in the cell that may be coopted by the

virus to assist in their expected functions (Verchot 2012).

An interesting finding was the effect of the knocked down expression of gene

$P58^{IPK}$ on TuMV replication. In mammals P58[IPK], a tetratricopeptide repeat (TPR)-

containing protein, is recruited by viruses (*e.g.*, influenza A virus) to inhibit interferon

activation and cell death mediated by the dsRNA-activated protein kinase (PKR), thus

favoring viral spread (Goodman et al. 2011). In our experiments, $p58^{IPK}$ mutant plants

showed enhanced tolerance to infection by showing weaker symptoms than WT plants

despite the same level of virus accumulation (supplementary figure S1, Supplementary

Material online). In sharp contrast, Bilgin et al. (2003) found that infection of $p58^{IPK}$-

silenced *Nicotiana benthamiana* and *A. thaliana* plants infected with tobacco mosaic

virus led to death. Given the limited information available from other plant-virus

pathosystems, additional experiments would be necessary to shed light into the potential

antiviral role in plants of this gene.

An important conclusion of our study is that the generally accepted model of plant

defense signaling pathways, mainly based in experiments done with bacteria and fungi,

may not describe well the interaction between plants and viruses. By contrast, recessive

S gene-based resistance seems to better explain differences in susceptibility to infection

and viral accumulation, thus being a more promising target for future development of
resistant plants.
580
Evolution of specialist and generalist viral strategies depends on the host genotype
The GFG and MA models of host-virus interaction modes represent the two ends of a
continuum of possible outcomes (Agrawal and Lively 2002). The most relevant
difference between both models regards the expected genetic heterogeneity in both host
and virus populations. With a pure GFG interaction, the susceptible host types are
expected to disappear and the resistant types to dominate the population. *Vice versa*, the
most virulent virus allele would become fixed in the population at the cost of mild alleles.
However, constitutive activation of defenses is known to be costly for *A. thaliana* [*e.g.*,
SA-related defense responses pay a fitness cost in absence of pathogens (Traw et al.
2007)] and high virulence usually comes with a cost in terms of pathogen's transmission
(Acevedo et al. 2019). Hence, a pure GFG strategy seems unlikely to be achieved. By
contrast, with a pure MA interaction, negative frequency-dependent selection emerges,
such that rare *A. thaliana* resistance alleles have advantage and, as a result, a genetic
polymorphism shall be maintained (Schmid-Hempel 2011). Natural populations of *A.*
thaliana contain considerable amount of genetic variability for tolerance (Pagán et al.
2008) and for immunity-related genes (Todesco et al. 2010, van de Weyer et al. 2019),
thus suggesting that an arms race between pathogens and plants is still ongoing but it is
unclear whether it may result in pure GFG or MA interactions or lie somewhere in
between. Evolution experiments with such different pathosystems as TuMV-*A. thaliana*
(González et al. 2019), *Octosporea bayeri-Daphnia magna* (Altermatt and Ebert 2008)
and *Serratia marcescens-Caenorhabditis elegans* (Gibson et al. 2020; White et al. 2020)
have produced congruent results: parasites exposed to heterogeneous host populations
evolved significantly lower virulence than parasites exposed to homogeneous host
populations. However, a significant difference exists among pathosystems: while viruses
exposed to genetically heterogenous host populations evolved as no-cost generalists,
evolution of generalism in more complex parasites was constrained by a fitness tradeoff,
as expected for the jack-of-all trades hypothesis (Bedhomme et al. 2015).
Our results, as well as those by Hillung et al. (2014) and González et al. (2019)
have shown the evolution of significantly nested bipartite infection networks, a finding
compatible with the existence of a combination of specialist and generalist viruses and of
more permissive and resistant host genotypes. Indeed, these studies have also shown that

more permissive hosts selected for more specialized viruses while more resistant hosts

selected for more generalist viruses, here matching the predictions of the GFG model.

Our observation of small yet significant modularity in the infection network was easily

explained by convergent evolution of TuMV lineages evolved in the same host genotype.

However, it has been recently shown in a long-term survey of the prevalence of different

plant viruses in different hosts and habitats that nestedness and modularity in host-

pathogen infection networks is possible due to the spatially patched distribution of

habitats and temporal successions of plant species (Valverde et al. 2020): small spatial

scales create modularity that coexist with global nestedness. This pattern may change

spatially and temporally but remains stable over long evolutionary timescales.

Role of natural selection

We found evidence of significant genetic differentiation among TuMV lineages evolved

in different plant genotypes. To test whether these differences were driven by selection

we performed a Tajima's D test (Tajima 1989). The resulting negative D value was

significant, which is compatible with the action of purifying selection, the presence of

slightly deleterious mutations segregating in the populations or fast population

expansions (Yang 2006). How to distinguish between these explanations? In

independent fast expanding populations, many new mutations may be generated and

rising in frequency in each population, thus being observed as singletons, mutations

present in only one of the many coexisting genomes in each evolving TuMV lineage.

Singletons inflate the number of segregating sites and thus cause $D < 0$. Indeed, this is

the case here: 56 out of the 73 observed variable sites are singletons, thus the observed

pattern of molecular diversity among lineages evolved in the same host genotype and in

different host genotypes is likely to be due to the fast expansions of viral populations.

However, we have found additional evidence supporting the action of positive

selection: the existence of a number of convergent nonsynonymous mutations arising in

independent lineages. Some of these nonsynonymous mutations evolved in the same host

genotype but others arise in different host genotypes (supplementary table S1 and figure

S3, Supplementary Material online). Yet without a clear association with the particular

signaling pathway, mutated S gene (table 1) or the phenogroup they belong to (figure 1).

Interestingly, most of these convergent mutations happened in the *VPg* cistron, which

also turns out to be the most variable one. VPg plays many essential roles in genome

transcription (it is linked to the 5'-end of the viral genome and provides the hydroxyl

group that primes the synthesis of the complementary strains by the viral RdRp),

translation [directly interacts with the eukaryotic initiation factors eIF(iso)4E and

eIF(iso)4G] and interacts with all other viral proteins (Bosque et al. 2014) and some of

the host cell proteins (Martínez et al. 2016; Martínez et al. 2020). Indeed, in previous

evolution experiments with potyviruses, *VPg* has also been shown to be an important

target of selection. For example, Agudelo-Romero et al. (2008) found that a single amino

acid replacement in VPg was enough to largely increase TEV *I*, *SS*, *VL* and in *A. thaliana*.

Similarly, Gallois et al. (2010) found that *A. thaliana* plants with knock-out mutations in

the *eIF(iso)4E*, *eIF(iso)4G1* and *eIF(iso)4G2* genes were resistant to TuMV infection.

Two mutations in the VPg (E116Q and N163Y) were enough to overcome this resistance

and return to the original infection phenotype, though yeast-two hybrid assays showed

that none of these mutations affected the binding of VPg with eIF(iso)4E (Gallois et al.

2010). As a final example, one of the most extensively used resistance genes against

potato virus Y in commercial pepper cultivars is *pvr²*, which has many different alleles

(Nicaise et al. 2007; Charron et al. 2008). The *pvr²* locus encodes for the eIF4E factor

which, as mentioned above, physically interacts with VPg. Interestingly, all the

resistance-breaking viral isolates found so far contain mutations in the *VPg* cistron

(Duprat et al. 2002; Moury et al. 2004; Ayme et al. 2006).

One of the two mutations identified by Gallois et al. (2010), VPg/E116Q, affects

the same protein domain as mutations D113G and R118H identified and characterized

here. The mechanism by which these two mutations may confer a selective advantage to

TuMV lineages cannot be inferred from our studies. We found VPg/D113G in several

lineages evolved in different host genotypes, while VPg/R118H was only found in the

jin1 lineage. Our assays failed to find a beneficial effect of VPg/D113G but confirmed

the host-specific beneficial effect of VPg/R118H in *jin1* plants.

Unfortunately, we have not been able to exhaustively test all observed mutations

and, hence, several others may still be potential candidates for adaptive mutations. More

specifically, we have not tested possible epistatic interactions among mutations fixed in

the same genome. For example, three lineages carry more than one nonsynonymous

mutation in *VPg* (*dbp2*/L1 D113N and N115E, *eds8-1*/L1 N115H and E116G, and *eds8-*

1/L4 H33Y, D113N and K121E) (supplementary table S1, Supplementary Material

online). In addition, 25 lineages carry additional nonsynonymous mutations in at least

one other cistron besides *VPg* (supplementary table S1, Supplementary Material online).

Two lineages have up to four additional nonsynonymous mutations: (*i*) *cpr5-2*/L5 that

has mutations in proteins P3/T318M-T326S, CI/V148I and VPg/D113G and (*ii*) *dcl2*

dcl4/L1 that has mutations in proteins P1/V54A, CI/I378V, VPg/R118C, and CP/S70N.

Therefore, plenty of opportunities for epistatic effects both within and among cistrons

exist, and will be evaluated in future experiments.

Methods
686

Plants, virus and growth conditions
687

A collection of 21 different *A. thaliana* mutants of the Col-0 accession were used for this

study (table 1). In all experiments described below, plants were all maintained in a BSL2

climatic chamber under a photoperiod of 8 h light (LED tubes at PAR 90 - 100 μmol/m^2/s)

at 24 ºC and 16 h dark at 20 ºC and 40% relative humidity.

Prior to the inoculation experiments, we created a large stock of TuMV infectious

saps. Saps were obtained from TuMV-infected *N. benthamiana* Domin plants inoculated

with the infectious plasmid p35STunos that contains a cDNA of TuMV genome

(GeneBank accession AF530055.2) under the control of the cauliflower mosaic virus 35S

promoter and the *nos* terminator (Chen et al. 2003) as described elsewhere (González et

al. 2019; Corrêa et al. 2020). This TuMV sequence variant corresponds to the YC5 isolate

from calla lily (*Zantesdeschia* sp.) (Chen et al. 2003). After the plants showed symptoms

of infection, they were pooled and frozen with liquid N$_2$. This frozen plant tissue was

homogenized into a fine powder using a Mixer Mill MM400 (Retsch GmbH, Haan,

Germany). For inoculations, the 0.01 g of powder was diluted in 1 mL inoculation buffer

(50 mM phosphate buffer pH 7.0, 3% PEG6000, 10% Carborundum) and 5 μL of the

inoculum was gently rubbed into the plant leaves. Plants were all inoculated when they

reached the growth stage 3.5 in the Boyes et al. (2001) scale. This synchronization

ensures that all hosts were at the same phenological stage when inoculated.

Phenotyping infected plants
707

Five different disease-related traits were measured for each infected plant for 18 dpi. (*i*)

Change in dry weight (ΔDW) of the aerial part of infected plants, with a precision of 10

mg, relative to the corresponding noninfected controls (supplementary figure S1A;

Supplementary Material online). (*ii*) Severity of symptoms (*SS*) evaluated in a semi-

quantitative discrete scale (figure 1B in Corrêa et al. 2020) ranging from zero for

asymptomatic infections to four for plants showing a generalized necrosis and wilting

(supplementary figure S1B, Supplementary Material online). (*iii*) The area under the

disease progress stairs (*AUDPS*), that summarizes the speed at which the disease

progresses in a group of plants (supplementary figure S1C, Supplementary Material

online) and takes values in the range (0, *N*) where *N* is the total number of plants included

in the assay (Simko and Piepho 2012). (*iv*) Infectivity measured as the number of

symptomatic plants out of the number of inoculated plants at 18 dpi (*I*; supplementary

figure S1D, Supplementary Material online). And (*v*) viral load (*VL*) measured by

absolute RT-qPCR as the number of viral genomes per ng of total RNA in the plant as

described below (supplementary figure S1E, Supplementary Material online).

The five traits were not all independent but showed some significant pairwise

positive correlations: *AUDPS* with *I* ($r = 0.772$, 19 df, $P < 0.001$), *SS* ($r = 0.807$, 19 df, P

< 0.001) and *VL* ($r = 0.472$, 19 df, $P = 0.031$), and *I* with *SS* ($r = 0.746$, 19 df, $P < 0.001$).

Basically, ΔDW is orthogonal with the other four traits while the other four traits showed

some degree of association. Interesting is the case of *VL*, which was only (weakly)

correlated to *AUDPS*.

Experimental evolution

Five TuMV lineages were evolved during 12 consecutive serial passages in each one of

the nine selected mutant genotypes. In addition, two lineages were also evolved in WT

plants. To begin the evolution experiment, ten 21-days old *A. thaliana* plants per lineage

and genotype were inoculated as described above with the virus stock previously created

from infected tissue of *N. benthamiana* plants inoculated with an TuMV isolate YC5

infectious clone. The next passages were made by harvesting the symptomatic plants at

14 dpi, preparing the infectious sap as described above and inoculating it to a new healthy

population of ten plants. These saps were 1/10 diluted with inoculation buffer and used

to inoculate the next batch of plants.

Total RNA extractions

Tissue from each pool of infected plants per lineage, genotype and serial passage was

collected, frozen with liquid N_2 and preserved at -80 ºC until it was homogenized into

fine powder using a Mixer Mill MM400. Next, an aliquot of approximately 100 mg of

grounded tissue was used for total RNA (RNAt) extraction. The RNAt was extracted

with the Agilent Plant RNA isolation Mini kit (Agilent Technologies, Santa Clara CA,

USA). Aliquots of RNAt per each sample were separated and their concentration adjusted

at approximately 50 ng/μL to estimate viral accumulation by RT-qPCR (see below).

Quantification of *VL*

VL of each plant sample per lineage, genotype and selected passage was quantified by

absolute real-time quantitative RT-PCR (RT-qPCR) using standard curves and the

primers TuMV F117 forward (5'-CAATACGTGCGAGAGAAGCACAC-3') and F118

reverse (5'-TAACCCCTTAACGCCAAGTAAG-3') that amplify a 173 nucleotides

fragment from the *CP* cistron of TuMV genome, as previously described (Corrêa et al.

2020). Briefly, standard curves were constructed using ten serial dilutions of the TuMV

genome, that was synthesized by *in vitro* transcription as detailed previously (Cervera et

al. 2018), in RNAt extract obtained from healthy *A. thaliana* plants used as control in the

experiments. Amplification reactions were run in a 20 μL volume using the GoTaq 1-

Step RT-qPCR System (Promega, Madison WI, USA) and the recommended

manufacturer's instructions as described in Cervera et al. (2018) in an ABI StepOne Plus

Real-time PCR System (Applied Biosystems, Foster City CA, USA). The cycling

conditions consisted in: an RT phase of 5 min at 42 ºC and 10 min at 95 ºC followed by

a PCR stage consisting in 40 cycles of 5 s at 95 ºC and 34 s at 60 ºC; and the final melt

curve profile that consisted in 15 s at 95 ºC, 1 min at 60ºC and 15 s at 95 ºC. Negative

controls consisted of healthy RNAt plant extract (mock-inoculated non infected control)

and water. For each sample, three technical replicates were quantified. The results were

analysed using the StepOne software 2.2.2 (Applied Biosystems).

TuMV genome amplifications and sequencing

Evolved viral genomes of the passage 12 from each lineage and genotype were amplified

by high-fidelity RT-PCR using the AccuScript Hi-Fi (Agilent Technologies) reverse

transcriptase and Phusion DNA polymerase (Thermo Scientific, Waltham MA, USA)

following the manufacturer's instructions. Each complete TuMV genome was amplified

into three overlapping amplicons of 3114 (5' fragment R1), 3697 (central region R2) and

3287 nucleotides (3' fragment R3) using three primer sets. For RT reactions an aliquot

of the corresponding RNAt (150-300 ng) was mixed with 0.25 μM of the 1R-P3 (5'-

CGAGTAGTATCTTATAGCACAGCGCTCCGACC-3'),   2R-NIa   (5'-

TGTCTGGAATCGGTAGCAAATGTAGCTGAGTTGTG-3') or 3R-polyAR (5'-

TTTTTTTTTTTTTTTTTTTTGTCCCTTGCATCATATCAAATG-3') primer to

synthesize the R1, R2 or R3 cDNA fragment, respectively, that were denatured 5 min at

65 ºC and cooled on ice. Then a mix containing AccuScript Hi-Fi 1× Buffer, 1 mM of

dNTPs, 8 mM of DTT, 4U of Ribolock RNase inhibitor (Thermo Scientific) and 0.5 μL

of AccuScript Hi-Fi (Agilent Technologies) was added up to a 10 μL volume. RT

reactions consisted of 90 min at 42 ºC to synthesize the cDNA followed by an incubation

of 5 min at 70 ºC to inactivate the enzyme. PCR reactions were performed in a 50 μL

volume containing a mix of 1× Phusion Buffer, 0.4 μM of dNTPs, 0.2 μM of each primer,

0.5-1 μL of DMSO, 2U of Phusion DNA polymerase (Thermo Scientific) and 1 μL of the

corresponding RT reaction. R1 fragment was amplified using the primer set 1F-5UTR

(5'-GCAAACGCAGACCTTTCGAAGCACTCAAGC-3') and 1R-P3 and the following

PCR conditions: an initial denaturation of 30 s at 98 ºC, 3 cycles of 10 s at 98 ºC, 20 s at

67 ºC and 2 min at 72 ºC, 3 cycles of 10 s at 98 ºC, 20 s at 65 ºC and 2 min at 72 ºC, and

32 cycles of 10 s at 98 ºC, 20 s at 63 ºC and 2 min at 72 ºC, followed by a final extension

step of 5 min at 72 ºC. Fragments R2 and R3 were amplified with primer set 2F-P3 (5'-

TGGGAGCTTGCGGATGGTGGATACACAATTC-3') and 2R-NIa or 3F-NIa (5'-

CTCGTTATATGGAGTCGGTTTCGGACCACTCATAT-3')   and   3R-polyAR,

respectively and a PCR with the same denaturation and extension steps than the fragment

R1 but different amplification steps: a stage consisting of 15 cycles of 10 s at 98 ºC, 20 s

at 67 ºC and 2 min at 72 ºC followed by 23 cycles of 10 s at 98 ºC, 20 s at 65 ºC and 2

min at 72 ºC for fragment R2. For fragment R3 15 cycles of 10 s at 98 ºC, 20 s at 67 ºC

and 2 min at 72 ºC followed by 23 cycles of 10 s at 98 ºC, 20 s at 65 ºC and 2 min at 72

ºC were used. PCR products were purified with the MSB Spin PCRapace Kit (Stratec

Molecular, Coronado CA, USA) and then Sanger-sequenced. Full-length consensus viral

sequences were obtained assembling the sequences of the three amplified products by

using the Genious R9.0.2 program.

Construction of TuMV *VPg* mutants

Mutants in the *VPg* cistron were synthesized by long inverse site-directed PCR

mutagenesis using the QuickChange II XL Kit (Stratagene, San Diego CA, USA)

following the manufacturer's instructions and using p35STunos as template. The

mutagenic primers were designed following the manufacturer's recommendations.

Primer set cpr5-2F (5'-GGAGGATGAGTTGGGTCCAAATGAAATACGTGT-3') and

cpr5-2R (5'- ACACGTATTTCATTTGGACCCAACTCATCCTCC-3') was used to

generate the A6238G (D113G) mutant while primer set jin1F (5'-

GGATCCAAATGAAATACATGTGAATAAGACAATTC-3') and jin1R (5'-

GAATTGTCTTATTCACATGTATTTCATTTGGATCC-3') to obtain the G6253A

(R118H) mutant. PCR protocol consisted of: a denaturation step of 2 min at 95 ºC

followed by 20 cycles of 20 s at 95 ºC, 10 s at 60 ºC and 7 min at 68 ºC, and a final

extension of 5 min at 68 ºC. After *Dpn*I digestion and transformation of electrocompetent

Escherichia coli DH5α, the presence of the desired mutations in the infectious clone and

the absence of undesired nucleotide changes was confirmed by sequencing.

Bioassays of the TuMV *VPg* mutants

The infectivity, viability and pathogenicity of both VPg TuMV mutants was confirmed

by performing three independent bioassays. In each experiment, three batches of 24 *A.*

thaliana plants (three-weeks old) of genotypes WT, *cpr5-2* and *jin1* were inoculated with

the WT TuMV plasmid clone used as reference and with plasmids of each of the TuMV

VPg mutants. Another two batches of 24 plants from *cpr5-2* or *jin1* genotypes were

inoculated with the WT TuMV plasmid and the corresponding plasmid of the TuMV VPg

mutant, the VPg/D113G mutant in the case of the *cpr5-2* genotype and that of

VPg/R118H for the *jin1* genotype. *A. thaliana* plants were inoculated by abrasion of

three leaves applying equal amounts of the plasmid inoculum, a total of approximately 7

μg per plant distributed in 3 μL per leaf. Plasmids were previously purified by Midiprep

using the NucleoBond Xtra Midi Kit (Macherey-Nagel, Düren, Germany) and

resuspended in distilled water. Number of infected plants and symptom intensity were

collected every day until 15 dpi.

Statistical analyses

All statistical analyses described hereafter were performed with SPSS version 27 software

(IBM, Armonk, NY), unless otherwise indicated.

AUDPS (figure 2), *I* (figure 3) and *VL* (figure 4) data were fitted all together to a

multivariate analysis of covariance (MANCOVA) model with plant genotype (*G*) as main

factor, independent evolution lineages (*L*) were nested within *G* and passage (*t*) was

introduced in the model as a covariable. The full model equation thus reads:

$\mathbf{P}_{ijk}(t) \sim \Pi + t + G_i + L(G)_{ij} + (t \times G)_i + [t \times L(G)]_{ij} + \boldsymbol{\varepsilon}_{ijk},$ (Eq. 1)

846 where $\mathbf{P}_{ijk}(t) = (AUDPS_{ijk}, I_{ijk}, VL_{ijk})^{\mathrm{T}}$ is the vector of phenotypic traits observed at time t,

847 for an individual infected plant k of evolutionary lineage j of genotype i, $\mathbf{\Pi}$ represents the

848 vector of phenotypic grand mean values and $\mathbf{\varepsilon}_{ijk}$ stands for the vector of errors assumed

849 to be Gaussian distributed at every t. The significance of each factor, the covariable and

850 their interactions was evaluated using the Wilks' Λ method. The magnitude of the effects

851 was evaluated using the η_P^2 statistic (proportion of total variability in the traits vector

852 attributable to each factor in the model; conventionally, values of $\eta_P^2 \geq 0.15$ are considered

853 as large effects).

854  Rates of phenotypic evolution for $AUDPS$ and I were estimated by fitting the time

855 series data to a first-order autoregressive integrated moving-average, ARIMA(1,0,0),

856 model (Elena and Sanjuán, 2005; González et al. 2019). The model equation fitted has

857 the form:

858 $Y_t - \rho_1 Y_{t-1} = Y_0 + \nu_Y t + \varepsilon_t,$        (Eq. 2)

859 where Y_k represents the variable being analyzed at passage k, ρ_1 measures the degree of

860 self-similarity in the time-series data (correlation between values at passages t and $t - 1$),

861 ε_t represents the sampling error at passage t, and ν_Y represents the linear dependency of

862 variable Y with passage number, that is, the rate of phenotypic evolution. The rates of

863 phenotypic evolution were further analyzed in the context of the mutated defense

864 signaling pathway genes or recessive resistances. Rates of phenotypic evolution were

865 fitted to a multivariate analysis of variance (MANOVA) in which the plant genotype,

866 serving as host for the evolutionary lineages (G), was nested within the factor being

867 analyzed (*i.e.*, the type of selection –hard *vs* soft-, the mode of resistance –signaling

868 pathways *vs* S-, and whether mutations affected the SA-dependent, the JA/ET-dependent

869 signaling or the RNA-silencing pathways; table 1). The full model equation now thus

870 reads:

871 $\mathbf{V}_{ijk} \sim \mathbf{\Phi} + X_{ij} + X(G)_{ij} + \mathbf{\varepsilon}_{ijk},$      (Eq. 3)

872 where $\mathbf{V}_{ijk} = (\nu_{AUDPS}, \nu_I)^{\mathrm{T}}$ is the vector of rates of phenotypic evolution observed for the

873 k replicate lineage evolved in plant genotype j within the factor X_i, $\mathbf{\Phi}$ represents the vector

874 of grand mean values and $\mathbf{\varepsilon}_{ijk}$ the vector of Gaussian errors.

876 **Analysis of infection network**

The first statistical approach consisted in fitting a logistic regression model to the

presence/absence of symptoms data using GLM techniques with a Binomial probability

distribution and a probit link function. The model equation reads as follows:

$\text{probit}[f_{ijkl}(t)] \sim \phi + t + TG_i + t \times TG_i + LG_j + t \times LG_j + L(LG)_{jk} + t \times L(LG)_{jk} +$

$(TG \times LG)_{ij} + t \times (TG \times LG)_{ij} + [TG \times L(LG)]_{ijk} + t \times [TG \times L(LG)]_{ijk} + \varepsilon_{ijkl},$

(Eq. 4)

where $f_{ijkl}(t)$ is the frequency of symptomatic plants t dpi in the test host genotype TG, for

the viral lineage L evolved in the local host genotype LG. TG and LG were considered

as orthogonal factors, L was nested within LG and dpi (t) was treated as a covariable. The

model includes all main factors, their corresponding nested and factorial interactions as

well as their interactions with the covariable. ϕ represents the grand mean of the probit-

transformed f values, and ε the error term assumed to be Binomial. The significance of

the different factors was evaluated using likelihood-ratio tests (LRT).

For the second statistical approach, the binary infection matrix was analyzed using

tools borrowed from the field of network biology to explore whether they show random

associations between viral lineages and host genotypes, one-to-one associations,

nestedness indicative of a GFG type of interaction, or modularity (Weitz et al. 2013). The

statistical properties of the infection matrix were evaluated using the R package

"bipartite" version 2.15 (Dormann and Strauss 2008) in R version 4.0.0 (R Core Team

2020) under RStudio 1.2.1335. Four different summary statistics were evaluated: T

nestedness (Bascompte et al. 2003), Q modularity (Newman 2006) and the d' species-

level (or Kullback-Leibler divergence) and H_2' network-level (or two-dimensional

normalized Shannon entropy) specialization indexes (Blüthgen et al. 2006). H_2' ranges

between zero and one for extreme generalists and specialists, respectively. Statistical

significance of these statistics was evaluated using Bascompte et al. (2003) null model.

The statistics of molecular evolution

Treating each lineage as an observation and each host genotype as a subpopulation, we

evaluated the average nucleotide diversity within host genotypes, π_S, the nucleotide

diversity for the entire sample, π_T, the interhost genotypes nucleotide diversity, δ_{ST}, and

the estimate of the proportion of interhost genotypes nucleotide diversity, known as

coefficient of nucleotide differentiation (Nei 1982), $N_{ST} = \delta_{ST}/\pi_T$. Standard deviations of

estimates were inferred from 1000 bootstrap samples. All these computations were done

using MEGA 11 (Tamura et al. 2021) and the lowest-BIC nucleotide substitution model

Kimura 2-parameters (Kimura 1980). Tajima's *D* test of selection (Tajima 1989) and its

statistical significance were evaluated using DnaSP6 (Rozas et al. 2017).

The frequency of mutations (*m*) per cistron (*C*), relative to the length of the

corresponding cistron, was fitted to the following logistic regression model using GLM

techniques with a Binomial probability distribution and a probit link function:

$\text{probit}(m_i) \sim \mu + C_i + \varepsilon_i$ (Eq. 5)

where μ is the average genomic mutation frequency and *i* refers to the 10 cistrons in the

main ORF.

The functional effect of mutations found in the *VPg* cistron were evaluated *in silico*

using the SNAP2 algorithm (Hecht et al. 2015). The algorithm provides a functionality

score and an accuracy index for each possible mutation affecting a coding sequence.

Negative score values mean functional neutrality while positive values should be taken

as indications of functional effects; the larger the values, the stronger the effect.

Supplementary Material

927 Supplementary data are available *at Molecular Biology and Evolution* online.

Acknowledgments

931 We thank Francisca de la Iglesia and Paula Agudo for excellent technical assistance and

932 the rest of the EvolSysVir lab members for fruitful discussions. This work was supported

933 by grants BFU2015-65037-P and PID2019-103998GB-I00 (Agencia Estatal de

934 Investigación - FEDER) and PROMETEU2019/012 (Generalitat Valenciana) to S.F.E.

References

Acevedo MA, Dillemuth FP, Flick AJ, Faldyn MJ, Elderd BD. 2019. Virulence-driven

trade-offs in disease transmission: a meta-analysis. *Evolution* 73:636-647.

Agrawal A, Lively CM. 2002. Infection genetics: gene-for-gene versus matching-alleles

models and all points in between. *Evol Ecol Res*. 4:79-90.

Agudelo-Romero P, Carbonell P, Pérez-Amador MA, Elena SF. 2008. Virus adaptation
by manipulation of host's gene expression. *PLoS ONE* 3:e2397.

Altermatt F, Ebert D. 2008. Genetic diversity of Daphnia magna populations enhances
resistance to parasites. *Ecol Lett*. 11:918-928.

Altizer S, Kaitala V, Laakso J, Ruokolainen L. 2006. Seasonality and the dynamics of
infectious diseases. *Ecol Lett*. 9:467-484.

Anttila J, Kaitala V, Laakso J, Ruokolainen J. 2015. Environmental variation generates
environmental opportunist pathogen outbreaks. *PLoS ONE* 10:e0145511.

Ayme V, Souche S, Caranta C, Jacquemond M, Chadoeuf J, Palloix A, Moury B. 2006.
Different mutations in the genome-linked protein VPg of potato virus Y confer
virulence on the *pvr2³* resistance in pepper. *Mol Plant Microbe Interact.* **19**: 557-
563.

Bascompte J, Jordano P, Melián CJ, Olesen JM. 2003. The nested assembly of plant-
animal mutualistic networks. *Proc Natl Acad Sci USA.* 100:9389-9387.

Bedhomme S, Hillung J, Elena SF. 2015. Emerging viruses: why are not jacks of all
trades? *Curr Opin Virol.* 10:1-6.

Berngruber T, Lion S, Gandon S. 2015. Spatial structure, transmission modes and the
evolution of viral exploitation strategies. *PLoS Pathog.* 11:e1004810.

Bilgin DD, Liu Y, Schiff M, Dinesh-Kumar SP. 2003. *P58^{IPK}*, a plant ortholog of double-
stranded RNA-dependent protein kinase PKR inhibitor, functions in viral
pathogenesis. *Dev Cell.* 4:651-661.

Blüthgen N, Menzel F, Blüthgen N. 2006. Measuring specialization in species interaction
networks. *BMC Ecol*. 6:9.

Boots M, Mealor M. 2007. Local interactions select for lower pathogen infectivity.
Science 315:1284-1286.

Bosque G, Folch-Fortuny A, Picó J, Ferrer A, Elena SF. 2014. Topology analysis and
visualization of *Potyvirus* protein-protein interaction network. *BMC Syst Biol*. 8:129.

Bouché N, Lauressergues D, Gasciolli V, Vaucheret J. 2006. An antagonistic function for
Arabidopsis DCL2 in development and a new function for DCL4 in generating viral
siRNAs. *EMBO J.* 25:3347-3356.

Boyes DC, Zayed AM, Ascenzi R, McCaskill MJ, Hoffman NE, Davis KR, Görlach J.
2001. Growth stage-based phenotypic analysis of Arabidopsis: a model for high
throughput functional genomics in plants. *Plant Cell* 13:1499-1510.

Brockhurst M, Buckling A, Rainey P. 2006. Spatial heterogeneity and the stability of
host-parasite coexistence. *J Evol Biol*. 19:374-379.

Brown JKM, Tellier A. 2011. Plant-parasite coevolution: bridging the gap between
genetics and ecology. *Annu Rev Phytopathol*. 49:345-367.

Cao H, Bowling SA, Gordon AS, Dong X. 1994. Characterization of an Arabidopsis
mutant that is nonresponsive to inducers of systemic acquired resistance. *Plant Cell*
6:1583-1592.

Carr JP, Lewsey MG, Palukaitis P. 2010. Signaling in induced resistance. *Adv Virus Res*.
76:57-121.

Castelló MJ, Carrasco JL, Navarrete-Gómez M, Daniel J, Granot D, Vera P. 2011. A plant
small polypeptide is a novel component of DNA-binding protein phosphatase 1-
mediated resistance to plum pox virus in Arabidopsis. *Plant Physiol*. 157:2206-2215.

Cervera H, Ambrós S, Bernet GP, Rodrigo G, Elena SF. 2018. Viral fitness correlates
with the magnitude and direction of the perturbation induced in the host's
transcriptome: the *Tobacco etch potyvirus*-tobacco case study. *Mol Biol Evol*.
35:1599-1615.

Chabas H, Lion S, Nicot A, Meaden S, van Houte, S, Moineu S, Wahl LM, Westra ER,
Gandon S. 2018. Evolutionary emergence of infectious diseases in heterogeneous
host populations. *PLoS Biol*. 16:e2006738.

Charron C, Nicolaï, M, Gallois JL, Robaglia C, Moury B, Palloix A, Caranta, C. 2008.
Natural variation and functional analyses provide evidence for coevolution between
plant eIF4E and potyviral VPg. *Plant J*. 54:56-68.

Chen CC, Chao CH, Chen CC, Yeh SD, Tsai HT, Chang CA. 2003. Identification of
turnip mosaic virus isolates causing yellow stripe and spot on calla lily. *Plant Dis*.
87:901-905.

Corrêa RL, Sanz-Carbonell A, Kogej Z, Müller SY, Ambrós S, López-Gomollón S,
Gómez G, Baulcombe DC, Elena SF. 2020. Viral fitness determines the magnitude
of transcriptomic and epigenomic reprogramming of defense responses in plants. *Mol*
Biol Evol. 37:1866-1881.

Cui H, Qiu J. Zhou Y, Bhandari DD, Zhao C, Bautor J, Parker JE. 2018. Antagonism of
transcription factor MYC2 by EDS1/PAD4 complexes bolsters salicylic acid defense
in Arabidopsis effector-triggered immunity. *Mol Plant* 11:1053-1066.

Dormann CF, Gruber B, Fruend J. 2008. Introducing the bipartite package: analyzing
ecological networks. *R News* 8:8-11.

Dormann CF, Strauss R. 2014. A method for detecting modules in quantitative bipartite
networks. *Meth Ecol Evol*. 5:9-98.

Duprat A, Caranta C, Revers F, Menand B, Browning KS, Robaglia C. 2002. The
Arabidopsis eukaryotic initiation factor (iso)4E is dispensable for plant growth but
required for susceptibility to potyviruses. *Plant J*. 32:927-934.

Elena SF, Sanjuán R. 2005. RNA viruses as complex adaptive systems. *BioSystems*
81:31-41.

Flor HH. 1956. The complementary genetic systems in flax and flax rush. *Adv Genet*.
8:29-54.

Frank SA. 1993. Specificity versus detectable polymorphism in host-parasite genetics.
Proc R Soc B. 254:191-197.

Gallois JL, Charron C, Sánchez F, Pagny G, Houvenaghel MC, Moretti A, Ponz F, Revers
F, Caranta C, German-Retana, S. 2010. Single amino acid changes in the turnip
mosaic virus viral genome-linked protein (VPg) confer virulence towards
Arabidopsis thaliana mutants knocked out for eukaryotic initiation factors
eIF(iso)4E and eIF(iso)4G. *J Gen Virol*. 91:288-293.

Gandon S. 2004. Evolution of multihost parasites. *Evolution* 58:455-469.

Ganusov VV, Bergstrom CT, Antia R. 2002. Within-host population dynamics and the
evolution of microparasites in a heterogeneous host population. *Evolution* 56:213-
223.

Gibson AK, Baffoe-Bonnie H, Penley MJ, Lin J, Owens R, Khalid A, Morran LT. 2020.
The evolution of parasite host range in heterogeneous host populations. *J Evol Biol*.
33:773-782.

González R, Butković A, Elena SF. 2019. Role of host genetic diversity for susceptibility-
to-infection in the evolution of virulence of a plant virus. *Virus Evol*. 5:vez024.

Goodman AG, Tanner BCW, Chang ST, Esteban M, Katze MG. 2011. Virus infection
rapidly activates the P58[IPK] pathway, delaying peak kinase activation to enhance viral
replication. *Virology* 417:27-36.

Gupta V, Willits MG, Glazebrook J. 2000. *Arabidopsis thaliana EDS4* contributes to
salicylic acid (SA)-dependent expression of defense responses: evidence for
inhibition of jasmonic acid signaling by SA. *Mol Plant Microbe Interact*. 13:503-
511.

Haraguchi Y, Sasaki A. 2000. The evolution of parasite virulence and transmission rate
in a spatially structured population. *J Theor Biol*. 203:85-96.

Hecht M, Bromberg Y, Rost B. 2015. Better prediction of functional effects for sequence
variants. *BMC Genomics* 16:S1.

Hillung J, Cuevas JM, Valverde S, Elena SF. 2014. Experimental evolution of an
emerging plant virus in host genotypes that differ in their susceptibility to infection.
Evolution 68:2467-2480.

Hughes WHO, Boomsma JJ. 2006. Does genetic diversity hinder parasite evolution in
social insect colonies? *J Evol Biol*. 19:132-143.

Kachroo P, Chandra-Shekara AC, Klessig DF. 2006. Plant signal transduction and
defense against viral pathogens. *Adv Virus Res*. 66:161-191.

Kaschau K, Carrington JC. 2001. Long-distance movement and replication maintenance
functions correlate with silencing suppression activity of potyviral HC-Pro. *Virology*
285:71-81.

Kimura M. 1980. A simple method for estimating evolutionary rate of base substitutions
through comparative studies of nucleotide sequences. *J Mol Evol*. 16:111-120.

Lalić J, Agudelo-Romero P, Carrasco P, Elena SF. 2010. Adaptation of *Tobacco etch*
potyvirus to a susceptible ecotype of *Arabidopsis thaliana* capacitates it for systemic
infection of resistant ecotypes. *Phil Trans R Soc B*. 65:1997-2008.

Laurie-Berry N, Joardar V, Street IH, Kunkel BN. 2006. The *Arabidopsis thaliana*
JASMONATE INSENSITIVE 1 gene is required for suppression of salicylic acid-
dependent defenses during infection by *Pseudomonas syringae*. *Mol Plant Microbe*
Interact. 19:789-800.

Lin SS, Wu HW, Jan FJ, Hou RF, Yeh SD. 2007. Modifications of the helper component-
protease of zucchini yellow mosaic virus for generation of attenuated mutants for
cross protection against severe infection. *Phytopathology* 97:287-296.

Lively CM. 2010. The effect of host genetic diversity on disease spread. *Am Nat*.
175:e149-e152.

Loebenstein G. 2009. Local lesions and induced resistance. *Adv Virus Res*. 75:73-117.

Love AJ, Laval V, Geri C, Laird J, Tomos AD, Hooks MA, Milner JJ. 2007. Components
of Arabidopsis defense- and ethylene-signaling pathways regulate susceptibility to
cauliflower mosaic virus by restricting long-distance movement. *Mol Plant Microbe*
Interact. 20:659-670.

Mäkinen K. 2019. Plant susceptibility genes as a source for potyvirus resistance. *Ann*
Appl Biol. 173:122-129.

Martínez F, Rodrigo G, Aragonés V, Ruiz M, Lodewijk I, Fernández U, Elena SF, Daròs
JA. 2016. Interaction network of *Tobacco etch potyvirus* NIa protein with the host
proteome during infection. *BMC Genomics* 17:1.

Martínez F, Toft C, Hillung J, Carrasco JL, Giménez-Santamarina S, Yenush L, Rodrigo
G, Elena SF. 2020. A comprehensive physical interaction map between the *Turnip*
mosaic potyvirus and *Arabidopsis thaliana* proteomes. *Res. Square*
doi:10.21203/rs.3.rs-149993/v2.

Mine A, Nobori T, Salazar-Rodon MC, Winkelmüller TM, Anver S, Becker D, Tsuda K.
2017. An incoherent feed-forward loop mediates robustness and tunability in a plant
immune network. *EMBO Rep*. 18:464-476.

Moreno-Gámez S, Stephan W, Tellier L. 2013. Effect of disease prevalence and spatial
heterogeneity on polymorphism maintenance in host-parasite interactions. *Plant*
Pathol. 62:133-141.

Moury B, Morel C, Johansen E, Guilbaud L, Souche S, Ayme V, Caranta C, Palloix A,
Jacquemond M. 2004. Mutations in potato virus Y genome-linked protein determine
virulence towards recessive resistances in *Capsicum annuum* and *Lycopersicon*
hirsutum. *Mol Plant Microbe Interact*. 17:322-329.

Nawrath C, Heck, S, Parinthawong N, Métraux JP. 2002. *EDS5*, an essential component
of salicylic acid-dependent signaling for disease resistance in Arabidopsis, is a
member of the MATE transporter family. *Plant Cell* 14:275-286.

Nawrath C, Métraux JP. 1999. Salicylic acid induction-deficient mutants of Arabidopsis
express PR-2 and PR-5 and accumulate high levels of camalexin after pathogen
inoculation. *Plant Cell* 11:1393-1404.

Nei M. 1982. Evolution of human races at the gene level. In *Human Genetics, part A: the*
Unfolding Genome (ed. B. Bonné-Tamir). Alan R. Liss, New York.

Newman MEJ. 2006. Modularity and community structure in networks. *Proc Natl Acad*
Sci USA. 103:8577-8582.

Nicaise V, Gallois JL, Chafiai F, Allen LM, Schurdi-Levraud V, Browning KS,
Candresse T, Caranta C, Le Gall O, German-Retana S. 2007. Coordinated and
selective recruitment of eIF4E and eIF4G factors for potyvirus infection in
Arabidopsis thaliana. *FEBS Lett*. 581:1041-1046.

Pagán I, Alonso-Blanco C, García-Arenal F. 008. Host responses in life-history traits and
tolerance to virus infection in *Arabidopsis thaliana*. *PLoS Pathog*. 4:e1000124.

Pagán I, Fraile A, Fernández-Fueyo E, Montes N, Alonso-Blanco C, García-Arenal F.

2010. *Arabidopsis thaliana* as a model for the study of plant-virus co-evolution. *Phil*

Trans R Soc B 365:1983-1995.

Parrat SR, Numminen E, Laine AL. 2016. Infectious disease dynamics in heterogeneous

landscapes. *Annu Rev Ecol Evol Syst*. 47:283-306.

Pfenning K. 2001. Evolution of pathogen virulence: the role of variation in host

phenotype. *Proc R Soc B*. 268:755-760.

Pieterse CMJ, León-Reyes A, van der Ent S, van Wees SC. 2009. Networking by small-

molecule hormones in plant immunity. *Nat Chem Biol*. 5:308-316.

Pieterse CMJ, van der Does D, Zamioudis C, León-Reyes A, van Wees SCM. 2012.

Hormonal modulation of plant immunity. *Annu Rev Cell Dev Biol*. 28:489-521.

Regoes RR, Nowak MA, Bonhoeffer S. 2000. Evolution of virulence in a heterogeneous

host population. *Evolution* 54:64-71.

Rodríguez D, Torres-Sorando L. 2001. Models of infectious diseases in spatially

heterogeneous environments. Bull Math Biol. 63:547-571.

Rozas J, Ferrer-Mata A, Sánchez-del Barrio JC, Guirao-Rico S, Librado P, Ramos-Osins

SE, Sánchez-Gracia A. 2017. DnaSP 6: DNA sequence polymorphism analysis of

large data sets. *Mol Biol Evol*. 34:3299-3302.

Sallinen S, Norberg A, Susi H, Laine, A.L. 2020. Intraspecific host variation plays a key

role in virus community assembly. *Nat Commun*. 11:5610.

Schmid-Hempel P. 2011. *Evolutionary Parasitology*. Oxford University Press, Oxford:

UK, pp. 259-272.

Schmid-Hempel P, Koella JC. 1994. Variability and its implications for host-parasite

interactions. *Parasitol Today* 10:98-102.

Simko I, Piepho HP. 2012. The area under the disease progress stair: calculation,

advantage, and application. *Phytopathology* 102:381-389.

Singh DP, Moore CA, Gilliland A, Carr JP. 2004. Activation of multiple antiviral defence

mechanisms by salicylic acid. *Mol Plant Pathol*. 5:57-63.

Sofer L, Cabanillas D, Gayral M, Téplier R, Pouzoulet J, Ducousso M, Dufin L, Bréhélin

C, Ziegler-Graff V, Brault V, Revers Frédéric. 2017. Identification of host factors

potentially involved in *RTM*-mediated resistance during potyvirus long distance

movement. *Arch Virol*. 162:1855-1865.

Soosaar JLM, Burch-Smith TM, Dinesh-Kumar SP. 2005. Mechanisms of plant

resistance to viruses. *Nat Rev Microbiol*. 3:789-799.

Tajima F. 1989. Statistical methods for testing the neutral mutation hypothesis by DNA
polymorphism. *Genetics* 123:585-595.

Tamura K, Stecher G, Kumar S. 2021. Molecular evolutionary genetics analysis version
11. *Mol Biol Evol* 38: 3022-3027.

Thines B, Katsir L, Melotto M, Niu Y, Mandaokar A, Liu G, Nomura K, He SY, Howe
GA, Browse J. 2007. JAZ repressor proteins are targets of the SCF(COI1) complex
during jasmonate signalling. *Nature* 448:661-665.

Todesco M, Balasubramanian S, Hu TT, Traw MB, Horton M, Epple P, Kuhns C,
Sureshkumar S, Schwartz C, Lanz C, Laitinen RAE, Huang Y, Chory J, Lipka V,
Borevitz JO, Dangl JL, Bergelson J, Nordborg M, Weigel D. 2010. Natural allelic
variation underlying a major fitness trade-off in *Arabidopsis thaliana*. *Nature*
465:632-636.

Ton J, van Pelt JA, van Loon LC, Pieterse CMJ. 2002. Differential effectiveness of
salicylate-dependent and jasmonate/ethylene-dependent induced resistance in
Arabidopsis. *Mol Plant Microbe Interact*. 15:27-34.

Torres-Barceló C, Martín S, Daròs JA, Elena SF. 2008. From hypo- to hypersuppression:
effect of amino acid substitutions on the RNA-silencing suppressor activity of the
tobacco etch potyvirus HC-Pro. *Genetics* 180:1039-1049.

Travisano M, Mongold, JA, Bennett, AF, Lenski RE. 1995. Experimental test of the roles
of adaptation, chance, and history in evolution. *Science* 267:87-90.

Traw MB, Kniskern JM, Bergelson J. 2007. SAR increases fitness of *Arabidopsis*
thaliana in the presence of natural bacterial pathogens. *Evolution* 61:2444-2449.

Valverde S, Vidiella B, Montáñez R, Fraile A, Sacristán S, García-Arenal F. 2020.
Coexistence of nestedness and modularity in host-pathogen infection networks. *Nat*
Ecol Evol. 4:568-577.

Van de Weyer AL, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, Witek K, Jones JDG,
Dangl JL, Weigel D, Bemm F. 2019. A species-wide inventory of NLR genes and
alleles in *Arabidopsis thaliana*. *Cell* 178:1260-1272.

Van Wees SCM, de Swart EAM, van Pelt JA, van Loon, LC, Pieterse CMJ. 2000.
Enhancement of induced disease resistance by simultaneous activation of salicylate-
and jasmonate-dependent defense pathways in *Arabidopsis thaliana*. *Proc Natl Acad*
Sci USA. 97:8711-8716.

Verchot J. 2012. Cellular chaperones and folding enzymes are vital contributors to
membrane bound replication and movement complexes during plant RNA virus
infection. *Front Plant Sci*. 3:271.

Voinnet O. 2001. RNA silencing as a plant immune system against viruses. *Trends Genet*.
17:449-459.

Weitz JS, Poisot T, Meyer JR, Flores CO, Valverde S, Sullivan MB, Hochberg ME. 2013.
Phage-bacteria infection networks. *Trends Microbiol*. 21:82-91.

White PS, Choi A, Pandey R, Menezes A, Penley M, Gibson AK, de Roode J, Morran L.
2020. Host heterogeneity mitigates virulence evolution. *Biol Lett.* 16:20190744.

Yang Z. 2006. *Computational Molecular Evolution*. Oxford University Press. Oxford:
UK, pp 262-264.

Yang Z, Huang Y, Yang J, Yao S, Zhao K, Wang D, Qin Q, Bian Z, Li Y, Lan Y, Zhou
T, Wang H, Liu C, Wang W, Qi Y, Xu Z, Li Y. 2020. Jasmonate signaling enhances
RNA silencing and antiviral defense in rice. *Cell Host Microbe* 28:89-103.

Yates A, Antia R, Regoes RR. 2006. How do pathogen evolution and host heterogeneity
interact in disease emergence? *Proc R Soc B*. 273: 3075-3083.

Zhou JM, Zhang Y. 2020. Plant immunity: danger perception and signaling. *Cell*
181:978-989.

Figures and Tables

FIG. 1. Nearest-neighbor clustering, based on the squared Euclidean distance, of the 21 mutant genotypes of *A. thaliana* according to their phenotypic similarity in response to TuMV infection. Genotypes selected as hosts for the evolution experiments are highlighted in yellow.

FIG. 2. Evolution of disease progression (*AUDPS*) along the serial passages of experimental evolution on each different host genotype. Different symbols and lines represent the independent evolutionary lineages. Panels are arranged from the most resistant genotype (*eds8-1*) to the most sensitive one (*jin1*) according to the groups defined in figure 1.

FIG. 3. Evolution of infectivity (*I*) along the serial passages of experimental evolution on each different host genotype. Different symbols and lines represent the independent evolutionary lineages. Panels are arranged from the most resistant genotype (*eds8-1*) to the most sensitive one (*jin1*) according to the groups defined in figure 1.

FIG. 4. Evolution of viral load (*VL*) along the serial passages of experimental evolution on each different host genotype. Different symbols and lines represent the independent evolutionary lineages. Panels are arranged from the most resistant genotype (*eds8-1*) to the most sensitive one (*jin1*) according to the groups defined in figure 1. TuMV *VL* was not evaluated for WT plants along the evolution experiment.

1202

FIG. 5. Test of the contribution of historical contingency to the observed pattern of adaptation of TuMV to the different *A. thaliana* genotypes. (A) Evolved *vs* ancestral values for *AUDPS*, (B) for *I*, and (C) for *VL*. The diagonal lines represent the null hypothesis of historical differences being fully preserved despite adaptation. The solid lines represent the linear regression of the data (dashed lines represent the 95% CI for the regression lines).

1203

FIG. 6. Estimated rates of evolution for *AUDPS* (A) and *I* (B) obtained from the fitting of an ARIMA(1,0,0) model (equation 2) to the data shown in figure 2 and figure 3, respectively. Different symbols represent different independent lineages evolved on the corresponding host genotype. Solid lines represent the linear regression of the data; dashed lines the 95% confidence intervals for the regression lines. In both cases, *A. thaliana* genotypes are ranked from the less to the more permissive to TuMV infection according to figure 1.

FIG. 7. The rates of evolution for *AUDPS* and *I* differ among possible mechanisms of resistance to infection. Rates of evolution are expressed in log-scale to better visualize the slowest rates of *I* evolution. Data were fitted to the MANOVA model defined by equation. 3. Error bars represent ±1 SEM.

1206

FIG. 8. Analysis of the full cross-infection matrix. (A) Packed matrix that highlights its nested structure, compatible with a gene-for-gene infection model. Black squares represent cases in which the *AUDPS* values were equal or greater than those observed for the corresponding lineage in its local host. Last column shows the species-level specialization index *d'* (*d'* = 0 means most generalist pathogen, *d'* = 1 means most specialist pathogen). (B) Bipartite host genotype - viral lineage infection network. The size of the nodes is proportional to their degree. The phenotypic groups defined in figure 1 are indicated above the host genotype.

1207

Fig. 9. Phenotypic disease-related traits (*AUDPS*, *I* and *SS*) of TuMV VPg/D113G relative to WT VPg. (A) – (C) Evaluated in the WT *A. thaliana* and the *cpr5-2* mutant genotype. (D) – (F) Evaluated in the WT *A. thaliana* and the *jin1* mutant genotype. Error bars represent ±1 SEM.

1208

Table 1. Different arabidopsis genotypes used in this study. Highlighted in gray are those used for the evolution experiments.

Genotype	Gene name	Affected pathway	Expected phenotype relative to wildtype plants	Reference
Wildtype				
coi1-4	CORONATINE INSENSITIVE 1 (AT2G39940)	Repression of JA-responsive genes	No ISR, no effect on virus	Thines et al. (2007)
cpr5-2	CONSTITUTIVE EXPRESSOR OF PR GENES 5 (AT5G64930)	Membrane protein, negative regulator of pathogen-dependent SA signaling	More resistant, constitutive SAR	Love et al. (2007)
dbp2	DNA-BINDING PROTEIN PHOSPHATASE 2	Transcriptional regulation of gene expression in potyvirus-infected plants	More resistant to potyvirus infection	Castelló et al. (2011)
dcl2	DICER-LIKE 2 (AT3G03300)	Partial loss of RNA-silencing	No effect, siRNAs produced by DCL4	Bouché et al. (2006)
dcl4	DICER-LIKE 4 (AT5G20320)	Partial loss of RNA-silencing	No effect, siRNA produced by DCL2	Bouché et al. (2006)
dcl2 dcl4	Double mutant dcl2 dcl4	Complete loss of RNA-silencing	More susceptible, no siRNA production	Bouché et al. (2006)
dip2	DBP-INTERACTING PROTEIN 2 (AT5G03210)	Transcriptional regulation of gene expression in potyvirus-infected plants	More susceptible	Castelló et al. (2011)
eds4-1	ENHANCED DISEASE SUSCEPTIBILITY 4 (AT5G51200)	Loss of SA-dependent signaling	More susceptible, no SAR	Gupta et al. (2000)
eds5-1	ENHANCED DISEASE SUSCEPTIBILITY 5 (AT4G39030)	Lipase-like protein, positive regulator of pathogen-dependent SA signaling	More susceptible, no SAR	Nawrath et al. (2002)

eds8-1	*ENHANCED DISEASE SUSCEPTIBILITY 8*	Reduced expression of plant defensin genes, reduced ISR	More resistant, enhanced SAR	Love *et al.* (2007)
ein2-1	*ETHYLENE INSENSITIVE 2* (AT5G03280)	MAPK, ET signaling intermediate, negative regulator SA-dependent signaling	More resistant, enhanced SAR	Love *et al.* (2007)
etr1-1	*ETHYLENE RESPONSE 1* (AT1G66340)	ET receptor, negative regulator SA-dependent signaling	More resistant, enhanced SAR	Love *et al.* (2007)
hsp90-1	*HEAT SHOCK PROTEIN 1* (AT5G52640)	Recessive r gene, required for membrane-bound replication complexes; protein folding	More resistant, missing component for viral replication	Verchot (2012)
i4g1	*EUKARYOTIC TRANSLATION INITIATION FACTOR (ISO) 4G 1* (AT3G60240)	Recessive *r* gene, initiation of viral RNA translation	More resistant, missing component for viral gene expression	Nicaise *et al.* (2007)
i4g2	*EUKARYOTIC TRANSLATION INITIATION FACTOR (ISO) 4G 2*	Recessive *r* gene, initiation of viral RNA translation	More resistant, missing component for viral gene expression	Nicaise *et al.* (2007)
jin1	*JASMONATE INSENSITIVE 1* (AT1G32640)	Loss of JA signaling; negative regulator of SA-dependent signaling	More resistant, enhanced SAR	Laurie-Berry *et al.* (2006)
npr1-1	*NONEXPRESSER OF PR GENES 1* (AT1G64280)	Ankyrin-repeat protein required for *PR-1* activation	More susceptible, no SAR, no ISR	Cao *et al.* (1994)
$p58^{IPK}$	*HOMOLOG OF MAMMALIAN P58^{IPK}* (AT5G03160)	Constitutive activation of PKR	More resistant, strong apoptosis-mediated HR	Bilgin *et al.* (2003)

pad4-1	*PHYTOALEXIN DEFICIENT 4* (AT3G52430)	Lipase-like protein, positive regulator pathogen-dependent SA signaling	More susceptible, no SAR	Cui *et al*. (2018)
sid2-1	*SA INDUCTION DEFICIENT 2* (AT1G74710)	Isochorismate synthase, required for SA bioshynthesis	More susceptible, no SAR	Nawrath and Métraux (1999)

ET - ethylene; HR – hypersensitive response; ISR - induced systemic resistance; JA - jasmonic acid; MAPK – mitogen-activated protein kinase; PKR - protein kinase RNA-activated; SA - salicylic acid; SAR - systemic acquired resistance

1209

Table 2. Results of the multivariate analysis of variance (MANCOVA) for the three phenotypic traits evaluated along the course of experimental evolution. The different factors are defined in equation 1.

Source of variation	Wilk's Λ	F	df_1, df_2	P	η_P^2	$1 - \beta$
Π	0.004	3377.616	3, 42	< 0.001	0.996	1.000
t	0.265	38.799	3, 42	< 0.001	0.735	1.000
G	0.093	6.467	24, 122.414	< 0.001	0.547	1.000
$t \times G$	0.155	4.608	24, 122.414	< 0.001	0.463	1.000
$L(G)$	0.529	0.278	108, 126.692	1.000	0.191	0.373
$t \times L(G)$	0.466	0.351	105, 126.677	1.000	0.225	0.487

η_P^2: magnitude of the effect; $1 - \beta$: power of the test.

1210

1211

Table 3. Results of the probit regression testing for the specificity of adaptation. The different model factors are defined in equation. 4.

Source of variation	LRT	df	P	η_P^2	$1 - \beta$
ϕ	21224.493	1	< 0.001	0.959	1.000
t	25995.056	1	< 0.001	0.815	1.000
TG	295.160	8	< 0.001	0.569	1.000
$t \times TG$	568.212	8	< 0.001	0.010	0.981
LG	20.545	8	0.008	0.288	0.661
$t \times LG$	30.200	8	< 0.001	0.005	0.804
$L(LG)$	207.257	35	< 0.001	0.385	1.000
$t \times L(LG)$	235.073	35	< 0.001	0.016	0.984
$TG \times LG$	258.888	64	< 0.001	0.314	1.000
$t \times TG \times LG$	365.512	64	< 0.001	0.018	0.961
$TG \times L(LG)$	788.661	280	< 0.001	0.048	0.999
$t \times TG \times L(LG)$	1033.649	280	< 0.001	0.042	0.993

LRT: likelihood ratio test distributed as a χ^2; η_P^2: magnitude of the effect; $1 - \beta$: power of the test.

1212