Defects in plant immunity modulate the rates and patterns of RNA virus evolution

Rebeca Navarro[1], Silvia Ambrós[1], Fernando Martínez[1], Beilei Wu[1,$], José L. Carrasco[1], and Santiago F. Elena[1,2,*]

[1]Instituto de Biología Integrativa de Sistemas (CSIC – Universitat de València), Paterna, 46182 València, Spain.

[2]The Santa Fe Institute, Santa Fe NM87501, USA.

[*]Correspondence, e-mail: santiago.elena@csic.es.

[$]Permanent address: Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Haidian, 100193 Beijing, China.

Keywords: experimental evolution; generalism; plant immunity; plant-virus interactions; response to infection; specialization; virus evolution

Abstract

It is assumed that host genetic variability for susceptibility to infection will necessarily condition the evolution of viruses, either by driving them to diversification into strains that track the different host defense alleles (*e.g.*, antigenic diversity), or by canalization to infect only the most susceptible genotypes. Associated to these processes, virulence may or may not increase. To tackle these questions, we performed evolution experiments with turnip mosaic virus (TuMV) in *Arabidopsis thaliana* genotypes that differ in mutations in genes involved in resistance pathways and in genes whose products are essential for potyviruses infection. Plant genotypes classified into five groups according to their degree of resistance and intensity of symptoms. We found that evolution proceeded faster in the most resistant hosts than in the most permissive ones, as expected for adaptation to a harsh environment. The multifunctional viral protein VPg turned out to be the target of selection in most host genotypes. When all evolved TuMV lineages were tested for fitness in all plant genotypes used in the experiments, we found that the infection matrix was significantly nested, suggesting the evolution of generalist viruses selected by the most restrictive mutant genotypes. At the other side, a modular pattern, driven by convergent evolution of lineages evolved in the same host genotype, was also observed.

The spectrum of disease severity can be attributed to heterogeneity in virus' virulence or in host's factors; the two are not necessarily independent explanations and they must actually complement and/or interact each other. A problem faced by viruses is that host populations consist of individuals that had different degrees of susceptibility to infection (Schmid-Hempel & Koella 1994; Pfenning 2001). Therefore, adaptive changes improving viral fitness in one host genotype may be selected against, or be neutral, in an alternative one. Genetic variability in susceptibility of hosts and infectiousness of viruses have been well studied in animals and plants (*e.g.*, Schmid-Hempel & Koella 1994; Altizer 2006; Hughes & Boomsma 2006; Brown & Tellier 2011; Anttila *et al*. 2015; Parrat *et al*. 2016; González *et al*. 2019). The interaction between host and parasite genotypes has been explained in the light of two different theoretical models that represent the two extremes in a continuum of possibilities. At the one extreme, the so-called gene-for-gene (GFG) model, in which a virus genotype exists that can infect all host genotypes and a universally susceptible host genotype should also exist (Flor 1956). Resistance occurs when a host "resistance" gene is matched by at least one virus "avirulence" gene. Polymorphism in infectivity and resistance can be maintained only if virulence pays a cost. At the other extreme, the matching-alleles (MA) model is based on self- versus non-self-recognition systems in invertebrates. Infection is not possible unless the virus possesses all alleles that match those of the host (Frank 1993). In this case, polymorphism in infectivity and resistance are maintained by negative frequency-dependent selection.

Viral infection of plants is a complex system in which the virus parasitizes the host and utilizes all its cellular resources to replicate and systemically spread. In response, plants have evolved intricated signaling mechanisms (Zhou & Zhang 2020) that limit the spread of the virus, resulting in resistance. A variety of factors contribute to plant

resistance to viral infections. Broadly speaking, these factors can be classified into *basal*
if they are pre-existing and limit within-cell propagation and cell-to-cell spread, and
inducible if they are only activated upon infection and inhibit systemic virus replication
and movement. Basal mechanisms include susceptibility (*S*) genes that involve alleles of
cellular proteins that do not interact properly with viral factors, specially translation
initiation factors, required by the virus for successful exploitation of the cell's protein
synthesis machinery, heat shock proteins, that assist them in the formation of multiple
multiprotein complexes, or DNA binding phosphatases (Truniger & Aranda 2009; Carr
et al. 2010; Mäkinen 2019). In contrast, inducible mechanisms include most of resistance
(*R*) genes whose expression result in a broad-scale change in plant physiology via diverse
signal transduction pathways, particularly those regulated by the hormones salicylic acid
(SA), jasmonic acid (JA) and ethylene (ET) (Soosaar *et al.* 2005; Carr *et al.* 2010). These
changes include local cell apoptosis (*e.g.*, hypersensitive responses - HR; Loebenstein
2009), the upregulation of nonspecific responses against many different types of
pathogens throughout the entire plant (systemic acquired resistance – SAR- and induced
systemic resistance – ISR) (Kachroo *et al.* 2006; Carr *et al.* 2010), and the activation of
the RNA-silencing-based resistance, which seems to play a role both in basal and
inducible mechanisms (Voinnet 2001; Carr *et al.* 2010). Early host responses following
virus detection include changes in ion fluxes (mainly Ca^{++}), activation of signaling
pathways, major alterations of transcriptomic profiles, generation of reactive oxygen
species (ROS) and production of nitric oxide (NO) (Soosaar *et al.* 2005). These
immediate changes are followed by HR and the recruitment of SA and JA/ET signaling
pathways. The SA-mediated defense signaling pathway results in SAR, while the JA/ET-
mediated defense signaling pathway results in ISR, the latter being specifically involved
in symbiotic interactions between plants and beneficial microbes. Indeed, it appears that

ISR is not effective against infection by most viruses (Ton *et al.* 2002; Loebenstein 2009;

Pieterse *et al.* 2009). Interestingly, both SAR and ISR pathways converge into two master

regulators, the *ENHANCED DISEASE SUSCEPTIBILITY 1* (*EDS1*) and the

PHYTOALEXIN DEFICIENT 4 (*PAD4*) genes; *EDS1* and *PAD4* repress ISR and promote

SAR. Although the SA and JA/ET pathways have been classically viewed as mutually

antagonistic, several studies have revealed positive and negative crosstalk between them

(van Wees *et al.* 2000; Pieterse *et al.* 2012) as well as between them and the RNA-

silencing pathway (Soosaar *et al.* 2005; Carr *et al.* 2010; Yang *et al.* 2020). This

crosstalk, which has the network topological structure of an incoherent feed-forward loop,

creates robustness and tunability in the plant immune network (Mine *et al.* 2017).

A particularly well study example of *S* genes in the context of genetic variability

for resistance and virus adaptation in *Arabidopsis thaliana* (L.) Heynh are the

RESTRICTION TO TEV MOVEMENT (*RTM*) genes, composed by at least five different

loci (Sofer *et al.* 2017) that encode for a variety of functions needed by tobacco etch virus

(TEV; genus *Potyvirus*, family *Potyviridae*), and other potyviruses, to induce a systemic

infection of the plant. The mechanism of resistance in this case is well understood:

dominant alleles in all five loci result in resistance, whereas homozygous deletions in at

least one of the loci result in increased susceptibility to infection and different degrees of

systemic infection. By means of experimental evolution, it has been shown that TEV can

readily adapt to different plant ecotypes. In particular, serial passages in susceptible plant

genotypes *rtm1*/*rtm1* result in a viral isolate (TEV-*At*17) capable of infecting otherwise

fully resistant genotypes (Agudelo-Romero *et al.* 2008; Lalić *et al.* 2010). Additional

passages of TEV-*At*17 in plants with different *RTM* configurations resulted in increased

local adaptation and, more interestingly, in the evolution of a GFG-like interaction

mechanism: the most susceptible plant genotype selected for less pathogenic and highly

specialized viruses while the most resistant plant genotype selected for more virulent and

generalist TEV isolates (Hillung *et al.* 2015). Later on, these results were confirmed with

turnip mosaic virus (TuMV; genus *Potyvirus*, family *Potyviridae*) in the context of fully

monomorphic or maximally polymorphic experimental *A. thaliana* populations

(González *et al.* 2019).

However, these previous studies have barely paid any attention to the intricacies of

the underlying defense regulatory pathways of *A. thaliana*, the existence of multiple

defense responses (*e.g.*, SAR, ISR, RNA-silencing, and the diversity of *S* genes) and the

potential crosstalk among them. Therefore, an open question is how this intricacy

determines the evolutionary fate of viruses. For instance, would two independent defense

pathways select for specialist viruses adapted to counteract each one thus conforming a

fitness tradeoff? Would a tradeoff in the investment of plant resources between pathways

(*e.g.*, SAR *vs* ISR) result in a diversification of virus adaptive strategies? Would viruses

target crosstalk points across pathways thus evolving as generalists? Here we will use

the TuMV-*A. thaliana* experimental pathosystem to explore this sort of questions. After

exploring the variability in phenotypic responses to TuMV infection of a collection of *A.*

thaliana mutants in different *S* and *R* genes, we choose nine mutants which represent the

entire diversity in phenotypic responses, including both basal and inducible resistance

mechanisms. Then, we performed evolution experiments on each plant mutant genotype

and track the evolution of several disease-related traits. At the end of the evolution

experiment, we evaluated the effect of host genotypes in the rates of virus evolution and

the contribution of contingency, selection and stochasticity into the outcome of evolution.

Next, we explored whether the different TuMV lineages evolved as specialist or

generalists depending on the mutated defense mechanism of their local hosts. Finally,

we sought to identify the molecular changes experienced in the genome of the different

viral lineages and explored the possible adaptive value of a few convergent mutations.

Results

Classification of *A. thaliana* genotypes according to their phenotypic response to

TuMV infection

The 21 *A. thaliana* mutant genotypes used in the study are shown in Table 1, including

information about the affected signaling pathways or cellular processes as well as the

expected phenotype of infection relative to wild-type (WT) plants based on the

description of these mutants. Genotypes were classified according to their phenotypic

response to TuMV infection based in five different disease-related traits measured 18

days post-inoculation (dpi). Fig. 1 shows the neighbor-joining clustering of genotypes

according to their multi-trait phenotypic response to TuMV infection (data in Fig. S1);

we found five significant groups (hereafter named as G1 to G5). For all four members of

G1 the response was highly homogenous and consistent with an enhanced resistance

response to TuMV infection: no significant change in ΔDW whilst a significant and

consistent reduction in *SS*, *AUDPS*, *I*, and *VL* relative to infected WT plants. Three

members of G1 had enhanced SAR response (Table 1) and the fourth one (*i4g2*) is a well-

known *S* gene involved in plant resistance to potyvirus (Nicaise *et al.* 2007; Gallois *et al.*

2010). The only member of G2 was the strong apoptosis-inducer mutant *p58^{IPK}*, which

shows no significant changes in *SS*, *AUDPS*, *I*, and *VL* relative to infected WT plants but

a significant reduction in ΔDW, indeed, infected plants were heavier than the noninfected

controls, thus being this a case of increased tolerance to infection. The only member of

G3 was mutant *dbp2*, also described as specifically increasing resistance against another

potyvirus (Castelló *et al.* 2011), which shows a very interesting response to TuMV

infection: ΔDW was significantly increased while *SS*, *AUDPS* and *I* were significantly

reduced; no effect in *VL* was observed. Thus, *dbp2* represents an intermediate response

to TuMV infection between the highly resistant G1 and those genotypes in G4. G4

represents a sort of hotchpotch formed by genotypes that at most differ from WT plants

in two traits, but without a clear pattern. Remarkably, though, with the exception of *cpr5-*

2, *dcl2 dcl4*, and *dip2*, all show significant reductions in *VL*. Therefore, members from

G4 would be considered similar to WT in their phenotypic response to TuMV infection.

Finally, *jin1* was the only member of G5, which shows a significant increase in all

measured traits except *VL*, consistent with and enhanced susceptibility to TuMV

infection. This is a surprising finding since *jin1* has been described as inducing a

constitutive expression of the SA-dependent defenses that made plants extremely

resistant to the infection of biotrophic bacterial pathogens (Laurie-Berry *et al.* 2006). To

further characterize the progression of infection, we selected a subset of representatives

from groups G1 (*eds8-1*) and G4 (*cpr5-2*, *dcl2 dcl4* and *hsp90-1*) and evaluated daily *I*,

SS and *VL* (Fig. S2). The *cpr5-2* plants started showing symptoms faster, though only

80% of plants showed symptoms after 15 dpi. Only 20% of plants of genotype *eds8-1*

showed symptoms of infection, which were delayed on time. Infected *hsp90-1* and *dcl2*

dcl4 plants were undistinguishable from the WT infected ones. Regarding *VL*, all

genotypes accumulated less virus than WT plants, with *cpr5-2* being the one that did so

the less (Fig. S2)

Representatives from the five groups were selected for the subsequent evolution

experiment, covering the entire spectrum of responses to TuMV infection and

mechanisms of resistance/susceptibility (shadowed rows in Table 1): *eds8-1* and *i4g2*

from G1 representing the hardest selection regimes (more resistant genotypes); *p58*[IPK]

from G2; *dbp2* from G3; *cpr5-2*, *dcl2 dcl4*, *hsp90-1*, and *npr1-1* from G4; and *jin1* from

G5 representing the softest selective regime (less resistant genotype).

Experimental evolution in the representative *A. thaliana* genotypes

First, we sought to evaluate the dynamics of evolution for *AUDPS* (Fig. 2), *I* (Fig. 3) and

VL (Fig. 4). Data in these figures were fitted all together to the multivariate analysis of

covariance (MANCOVA) model described in Eq. 1 of the Methods section. Notice that

lineage *cpr5-2*/L2 showed a quite different evolution pattern from the other four lineages

evolved in *cpr5-2*. It never increased in either of the three traits and was lost after passage

five (Fig. 2, Fig. 3). Therefore, we ended up with 44 evolved lineages.

Table 2 shows the results of this MANCOVA. Despite a considerable amount of

noise in the time series, we found some significant results that can be summarized as

follows. First, a net significant and large ($\eta_P^2 = 0.735$) effect associated to the

evolutionary passages has been observed, indicating that the three phenotypic traits

related to TuMV infection evolved during the experiment. Second, plant genotypes had

a highly significant and large ($\eta_P^2 = 0.547$) effect on the phenotypic traits, suggesting that

TuMV evolutionary dynamics were strongly influenced by the local host genotype in

which it was being passaged. More interestingly, this effect associated to host plant

genotypes was strongly dependent on the passage number ($\eta_P^2 = 0.463$), thus suggesting

that the slope of the relationship between phenotypic traits and time was influenced by

the local host genotype. This effect will be further evaluated in the next section. Third,

independent lineages show a strong degree of evolutionary parallelism, as indicated by

nonsignificant differences among viral lineages evolved in the same local host genotype

(factors $L(G)$ and $t \times L(G)$). However, this last conclusion should be taken with certain

level of caution since the power of the two tests is relatively low $(1 - \beta < 0.800)$ and

hence we might be wrongly failing to reject the null hypothesis (type II error).

A relevant question in evolutionary biology is the extent in which ancestral

differences determine the fate of evolution. In other words, the relative contribution of

adaptation, chance and historical contingency to the evolution or organismal fitness.

Following the logic exposed by Travisano *et al.* (1995), we can imagine two situations.

First, ancestral statistical differences among phenotypes could be preserved along

evolution despite a net increase in the mean trait values (due to selection) and differences

among replicated lineages (due to chance; *i.e.*, mutation and drift). In this situation, we

should expect a nonzero slope in a regression of the evolved phenotypic values against

the ancestral ones. The closer the slope to a value of one, the more importance of ancestral

differences. Second, if initial trait variation among ancestral genotypes was eliminated

from the evolved populations because the combined effect of adaptation and chance, we

should expect a regression slope smaller than one. The less effect of ancestral differences,

the flatter the slope, being zero in the extreme case of ancestral differences being

completely erased. To evaluate the role of the ancestral differences in *AUDPS*, *I* and *VL*

of TuMV across the nine different plant genotypes, we evaluated the magnitude of their

change at the end of the evolution experiment. Fig. 5 shows the plots of evolved *vs*

ancestral values. For illustrative purposes, the diagonals represent the null hypothesis of

absolute preservation of ancestral differences. In the case of *AUDPS* (Fig. 5A), a

significant regression exists between evolved and ancestral values ($R = 0.510$, $F_{1,42} =$

14.732, $P < 0.001$), with a slope of 0.296 ±0.077 (±1 SD). Since the slope is still

significantly different from zero, yet clearly flatter than the diagonal ($t_{42} = 9.112$, $P <$

0.001), we conclude that ancestral differences have been mostly removed by the

combined action of selection and chance yet not completely erased. In the case of *I* (Fig.

5B) the regression was not significant ($R = 0.087$, $F_{1,43} = 0.031$, $P = 0.568$), thus suggesting that for this trait ancestral differences among *A. thaliana* genotypes have been fully erased by adaptation and chance. Finally, in the case of *VL* (Fig. 5C), the regression of evolved values on ancestral ones was significant ($R = 0.417$, $F_{1,43} = 8.822$, $P = 0.005$), suggesting again that ancestral phenotypic differences have been largely removed by selection and chance yet some still persist (regression slope -0.142 ± 0.048 significantly different from one: $t_{42} = 23.903$, $P < 0.001$).

Rates of phenotypic evolution for disease-related traits

As mentioned in the previous section, we have observed a significant effect of *A. thaliana* genotypes in the temporal evolution of the three phenotypic traits. To further explore this effect, we estimated the rates of phenotypic evolution of *AUDPS* and *I* using the ARIMA(1,0,0) model described in the Methods section (Eq. 2). *VL* was not included in this analysis since the number of data points (three) in the time series was the same than the number of parameters to be estimated, thus likely incurring in overfitting problems.

Fig. 6 shows the estimated rates of evolution, β, for host genotypes ordered from the most resistant (G1) to the most permissive (G5) ones. Very interestingly, a significant negative regression coefficient exists for both traits [*AUDPS*: $-1.460 \cdot 10^{-2} \pm 0.412 \cdot 10^{-2}$ ($R = 0.480$, $F_{1,42} = 12.561$, $P = 0.001$) and *I*: $-3.161 \cdot 10^{-3} \pm 0.909 \cdot 10^{-3}$ ($R = 0.473$, $F_{1,42} = 12.105$, $P = 0.001$)], suggesting that evolution always proceeded faster in the most restrictive hosts and slower in the most permissive ones.

Dissecting the role of specific defense mechanisms in TuMV evolution

The rates of TuMV phenotypic evolution estimated in the previous section were further analyzed in the context of the mutated defense signaling pathway encoded by *R* genes or

recessive resistances encoded by *S* genes. Rates of evolution β_{AUDPS} and β_I were fitted to

the MANOVA model shown in Eq. 3 of the Methods section. Fig. 7 graphically

summarizes the results of these analyses. First, we sought for differences in rates of

evolution between permissive (soft selection) and restrictive (hard selection) hosts (Fig.

7A). The multivariate analysis shows that lineages evolved in the more restrictive hosts

evolved significantly faster than those evolved in the more permissive hosts ($\Lambda = 0.327$,

$F_{2,34} = 34.923$, $P < 0.001$), with the effect being mostly driven by differences in the rate

of infectivity evolution β_I, as shown by the corresponding univariate analyses, being on

average 3.14 times faster in the most restrictive hosts. This observation confirms the

results shown in the previous section.

Next, we evaluated whether differences exist between the rates of phenotypic

evolution for viruses evolving in host genotypes carrying mutations affecting resistance

(*R*; *i.e.*, the SAR, ISR and RNA-silencing pathways) or susceptibility (*S*; *dbp2*, *dip2*,

hsp90-1, *i4g2*, and *p58^{IPK}*) genes (Fig. 7B). In this case, the multivariate analysis found

a small yet significant effect ($\Lambda = 0.790$, $F_{2,34} = 4.507$, $P = 0.018$), though none of the

two univariate analyses had enough power to detect it (indeed, changes went in opposite

directions: 1.02-fold larger for β_{AUDPS} in the *S* plants but 1.31 times larger for β_I in the *R*

plants).

Fig. 7C shows the comparison of rates of TuMV phenotypic evolution for lineages

evolving in plant genotypes mutant for the SA-dependent signaling defense pathways. In

this case we have three categories: mutations inducing a down-expression of the pathway,

mutations with no effect on this pathway and mutations resulting in overexpression.

Again, the MANOVA found highly significant differences among the three categories (Λ

$= 0.213$, $F_{4,68} = 19.861$, $P < 0.001$). The univariate analyses show that most of the

differences were due to a faster evolution of *I*, being 2.38 times faster in the host

genotypes over-expressing the SAR defenses than in those with a WT or even down-

expression of the SA signaling.

SA- and JA/ET-dependent pathways are considered as antagonistic defenses,

although more recent evidences point towards an active crosstalk between both (van Wees

et al. 2000; Pieterse *et al.* 2012; Mine *et al.* 2017). Under the antagonistic hypothesis,

we should expect the opposite trend than observed for the lineages evolved in plants over-

expressing SAR defenses (Fig. 7C). Fig. 7D shows the evolution rates data classified

according to whether host genotypes had mutations affecting the JA/ET defense (ISR)

signaling pathway. The MANOVA found a significant difference between the two

categories ($\Lambda = 0.433$, $F_{2,34} = 22.258$, $P < 0.001$), entirely driven by differences in β_I, as

confirmed by the univariate analyses. Indeed, as expected, the rates of evolution for

infectivity were 2-fold faster for TuMV lineages evolved in the ISR deficient hosts, in

agreement with the SAR/ISR antagonistic hypothesis.

Finally, we tested the effect of knocking down the host's RNA-silencing pathway

in the rates of TuMV evolution (Fig. 7E). In this case, the multivariate analysis also found

highly significant differences between rates of viral evolution in plants with a fully

functional RNA-silencing pathway and *dcl2 dcl4* plants ($\Lambda = 0.490$, $F_{2,34} = 17.729$, $P <$

0.001), with average β_I values being 21-fold larger in plants expressing a fully functional

RNA-silencing pathway than in *dcl2 dcl4* plants unable of producing siRNAs.

In conclusion from this section, we found that regardless the defense signaling

pathway, viruses evolving in permissive plants always evolve slower than their

counterparts evolving in more restrictive plant genotypes.

Permissiveness of *A. thaliana* genotypes to infection drive the evolution of specialist

or generalist TuMV lineages

To analyze the specificity of adaptation of each evolved TuMV lineage, we performed a

full cross-infection experiment in which all the 44 evolved lineages were inoculated into

ten plants of the nine *A. thaliana* genotypes used in the evolution experiments. The

presence of symptoms on each plant was evaluated daily for up to 12 dpi. These data

were analyzed using two different approaches, first an ANOVA-like that evaluates the

effect of host genotypes, viral lineages and their interactions (Schmid-Hempel, 2011);

and second, the inference and characterization of nestedness, modularity and

specialization of an infection network (Weitz *et al*. 2013). For the first approach we fitted

the presence/absence of symptoms data to the logistic regression model shown in Eq. 4

of the Methods section. The results are shown in Table 3. Despite its apparent

complexity, interpretation is straightforward. A *TG* factor describes the effect of

differences among host genotypes, irrespective of the infecting viral lineage; it thus

characterizes whether some host genotypes are more susceptible than other to infection.

As shown in Table 3, a *TG* has a highly significant effect, both by itself and in the

interaction with *t*. The magnitude of the net effect was very large ($\eta_P^2 = 0.569$), while its

interaction with *t* was rather small in magnitude ($\eta_P^2 = 0.010$). On average, the most

susceptible host genotypes were *jin1*, *dcl2 dcl4* and *cpr5-2* ($\bar{f} = 1.000$), whereas the most

resistant host genotypes were *p58^{IPK}* ($\bar{f} = 0.459 \pm 0.013$) and *eds8-1* ($\bar{f} = 0.233 \pm 0.013$).

Similarly, a highly significant *LG* effect exists by itself (of large magnitude $\eta_P^2 = 0.288$)

as well as in combination with *t* (though the magnitude of the interaction was rather small

in magnitude $\eta_P^2 = 0.005$), which means that the observation depends on the viral lineage

independently of the host genotype. On average, the lineages evolved in *eds8-1* ($\bar{f} =$

0.997 ± 23.710) and *dbp2* ($\bar{f} = 0.994 \pm 26.108$) were those inducing symptoms faster; by

contrast, lineages evolved in *i4g2* ($\bar{f} = 0.788 \pm 279.090$) showed symptoms much slower.

Finally, and most interesting, a highly significant interaction term *TG×LG* (as well as in

combination with *t*) indicates that the outcome depends on the particular combinations of

host genotypes and TuMV lineages. Indeed, the magnitude of the interaction effect was

large ($\eta_P^2 = 0.314$), though its dependence with dpi was not so relevant ($\eta_P^2 = 0.048$). For

example, in the extreme of the most susceptible genotypes, the case of *cpr5-2* perfectly

illustrates this case: lineages evolved in *cpr5-2*, *dbp2*, *dcl2 dcl4*, *eds8-1*, and *p58^{IPK}* all

show $\bar{f} = 1.000$, whereas lineages evolved in *i4g2* and *jin1* show much lower $\bar{f}$ values

(0.674 ±0.028 and 0.849 ±0.032, respectively). In the opposite extreme, *e.g.* in *eds8-1*

lineages evolved in *eds8-1* show $\bar{f} = 0.575$ ±0.050 while lineages evolved in *jin1* show $\bar{f}$

= 0.124 ±0.030; all other lineages showing intermediate values.

For the second analytical approach, we used the above infection dataset to estimate

the corresponding *AUDPS* values, thus creating an infection matrix of 44 rows [*L*(*LG*)]

by nine columns (*TG*). This matrix was transformed into a new binary matrix in which a

value of one meant that the *AUDPS* for a given TuMV lineage in a particular *TG* was

greater or equal than observed for its *LG* and zero otherwise. Fig. 8A shows the packed

infection matrix with rows and columns organized to better highlight is nestedness and

modularity (Weitz *et al*. 2013). The matrix could we used to build a binary infection

network (Fig. 8B) that provides the same information in a more graphical way. The

infection network was significantly nested (*T* = 14.706, *P* < 0.001), with the most

permissive genotypes *jin1* and *dcl2 dcl4* being susceptible to most viral lineages

(universally susceptible hosts), while the most resistant genotype *eds8-1* and *p58^{IPK}* were

only successfully infected by some lineages that were precisely evolved in these two

genotypes. Likewise, the most generalist viral lineages were *eds8-1*/L4, *p58^{IPK}*/L3 and

p58^{IPK}/L4 that infected with equal efficiency all nine host genotypes (universally virulent

viruses), while the most specialized viral lineages, able only to successfully infect their

local hosts, were all surviving lineages evolved in *cpr5-2*, four lineages evolved in *dcl2*

dcl4 and four lineages evolved in *jin1* (Fig. 8). Therefore, we conclude that local host's

resistance to infection positively correlates with the host range of the evolved lineages:

more resistant host genotypes selected for very generalist viruses while less resistant host

genotypes selected for very specialized viruses, in agreement with a gene-for-gene

infection model (Weitz *et al.* 2013).

Finally, we computed Blüthgen *et al.* (2006) specialization indexes for the packed

matrix shown in Fig. 8. Firstly, we evaluated the standardized species-level measure of

partner diversity *d'*. *d'* ranged from zero for the most generalist TuMV lineages (*eds8-*

1/L4, *p58^{IPK}*/L3 and *p58^{IPK}*/L4) to one for the most specialist ones (the four surviving

cpr5-2-evolved lineages, *dbp2*/L2, *dcl2 dcl4*/L2 - L5 and *jin1*/L1 - L4) (last column in

Fig. 8A). Secondly, we computed the network-level specialization index H'_2, obtaining a

value of $H'_2 = 0$ (uncorrected $H_2 = 5.024$, $H'_{min} = 3.534$, $H'_{max} = 5.024$; $P < 0.001$), which

means that, overall, the binary infection network shows a degree of specialization, with

most lineages being able of infecting one or few host genotype, a result mainly driven by

those TuMV lineages evolved in the less resistant host genotypes.

Finally, we sought to explore if the infection network in Fig. 8B also shows

evidences of modularity. Indeed, from an ecological standpoint, a significant network-

level degree of specialization means that some TuMV lineages do not interact well with

some of the *A. thaliana* genotypes, thus creating the possibility of modularity. In this

context, a module will refer to an aggregated set of viral lineages characterized by more

interactions within the module than between modules (Newman 2006; Dormann &

Strauss 2014). To this end, we computed the *Q* modularity index, which ranges from

zero, when the community has no more links within modules than expected by chance,

to a maximum value of one. We found a small yet significant modularity ($Q = 0.234$, *P*

= 0.047) in the infection matrix (Fig. 8A). What are the implications of modularity in

TuMV pathogenesis? Modularity is expected for a MA infection model (Weitz *et al*.

2013). More similar host genotypes, *e.g.*, those with mutations affecting the same

signaling pathway, may be selecting for viral lineages with similar properties, thus being

able of infecting with equal efficiency the subset of plant genotypes. A less appealing

alternative hypothesis to explain the observed modularity may be parallel evolution of

lineages evolved into the same host genotype; the most representative cases of this

possibility being the lineages evolved in *cpr5-2*, *dcl2 dcl4* and *jin1*. To distinguish

between these two hypotheses, we computed a reduced matrix averaging the observed

AUDPS among TuMV lineages evolved in the same host genotype and transforming this

nine by nine infection matrix into a packed binary matrix as described above. If the

reduced matrix still shows significant modularity, the first hypothesis will hold, while if

modularity disappears by averaging lineages evolved in the same host genotype, then the

alternative hypothesis will be more parsimonious. The reduced matrix is still

significantly nested ($T = 13.064$, $P < 0.001$), but modularity was not significant anymore

($Q = 0.286$, $P = 0.137$), thus supporting the alternative hypothesis that the observed

modularity was driven by convergent evolution of lineages evolved into the same host

genotype rather than by the overlap of selective pressures by genotypes carrying

mutations in the same defense mechanism.

Genomic differences/similarities among TuMV lineages and VPg as a potential

target of selection

Finally, we have explored the molecular changes experienced by the TuMV evolved

lineages. Full genomic consensus sequences were determined for all 44 evolved lineages.

A total of 114 mutational events have been observed, affecting 71 nucleotide position

(Supplementary Fig. S3A and Table S1). According to the type of nucleotide substitution

involved, 93 were transitions and 21 transversions; regarding their effect on the protein

sequence, 30 were synonymous and 84 nonsynonymous. Interestingly, some mutations,

including 13 nonsynonymous and four synonymous, have been observed multiple times

in independent lineages. As discussed in the Methods section, treating each lineage as an

observation and each host genotype as a subpopulation, the average nucleotide diversity

within host genotypes, referred only to the 71 polymorphic sites, is $\pi_S = 0.067 \pm 0.007$.

On the other hand, the nucleotide diversity for the entire sample is $\pi_T = 0.074 \pm 0.007$.

Consequently, the estimate of interhost genotypes nucleotide diversity is $\delta_{ST} = 0.007$

± 0.003, and the estimated coefficient of nucleotide differentiation (Nei 1982), $N_{ST} = 0.090$

± 0.033, a value significantly greater than zero ($z = 2.727$, $P = 0.003$). Thus, we conclude

that minor yet significant genetic differentiation has been generated among viral lineages

replicating in different host genotypes. To assess whether selection played a role in

genetic differentiation among *A. thaliana* genotypes, we performed a Tajima's *D* test

(Tajima 1989) and found that it was significantly negative ($D = -2.487$, $P = 0.006$).

Next, we sought to characterize the distribution of mutational events along the nine

non-overlapping cistrons (Fig. S3B). The frequency of mutations per cistron, relative to

the length of the corresponding cistron, was fitted to the logistic regression model shown

in Eq. 5 of the Methods. Highly significant differences exist ($\chi^2 = 127.545$, 9 d.f., $P <$

0.001), yet entirely due to the ~16-fold larger mutation frequency observed in the *VPg*

cistron relative to the rest of the genome (Fig. S3B). Notice that all mutations observed

in *VPg* are nonsynonymous and that all lineages except *eds8-1*/L3 carry at least one

mutation in this cistron (Fig. S3A and Table S1).

Convergent nonsynonymous mutations are, *a priori*, good candidates for adaptive

mutations. Mutation CP/V148I appears in two lineages evolved in *cpr5-2* (Fig. S3A and

Table S1), mutation CP/S70N appear in lineage *dcl2 dcl4*/L1 and three lineages evolved

in *jin1* (Fig. S3A and Table S1), and two different mutations affecting CP amino acid 112, CP/D112G (lineage *npr1-1*/L4) and CP/112A (lineage *i4g2*/L5), that result in a similar replacement of side chains. In addition, seven nonsynonymous mutations in *VPg* are shared by several lineages (Fig. S3A and Table S1). Out of these seven cases, three seem particularly promising candidates. They all are located in a narrow region of VPg (residues 113, 115 and 118) (Fig. S3A and Table S1). Firstly, mutations G6237A, G6237C, A6238G, and U6239A all affect the same codon, resulting in amino acid replacements D113N, D113H, D113G, and D113E, respectively. The expected functional effect of these mutations was evaluated using the algorithms implemented in SNAP2 (Hecht *et al*. 2015). D113E is predicted to be functionally neutral (score = −76, accuracy = 87%), while D113G (32, 66%), D113H (43, 71%) and D113N (30, 66%) are predicted to have a functional effect. Among these three amino acid replacements, D113G, which shows an intermediate value of the functional effect score, is particularly interesting, involving a very strong change in the side radical from a long negatively charged one to a small nonpolar one. This replacement has been observed in lineages *cpr5-2*/L3, *cpr5-2*/L4, *cpr5-2*/L5, *npr1-1*/L1, *npr1-1*/L5, and *hsp90-1*/L2. All three hosts for these lineages belong to the phenotypic group G4 in Fig. 1. Secondly, mutations A6243C, A6243G and A6244G also affect the same codon, resulting in amino acid replacements N115H (−42, 72%), N115E (−66, 82%) and N115S (−30, 61%). All three are predicted to be functionally neutral. Thirdly, mutations U6252 and G6253A affect the same codon and result in amino acid replacements R118C (−10, 53%) and R118H (56, 78%), respectively. The most frequent among these two replacements is R118H, which represents a conservative change among large positively charged side radicals, yet has a strong expected functional effect. R118H has been observed in four out of five TuMV lineages evolved in the *A. thaliana* mutant *jin1* (phenotypic group G5 in Fig. 1).

D113G thus represents a case of convergent evolution non-specific of the local host

genotype (*i.e.*, a candidate for a generalist mutation) while R118H represents a mutation

highly specific of the local host genotype (*i.e.*, a specialist mutation).

To further characterize the possible adaptive value of mutations VPg/D113G and

VPg/R118H, we created by site-directed mutagenesis the two mutated versions of *VPg*

and cloned them into the ancestral p35STunos infectious clone (Chen *et al.* 2003).

Viruses were recovered from these clones and three disease-related phenotypic traits (*e.g.*,

AUDPS, *I* and *SS*) evaluated both in the WT plants as well as in their corresponding local

hosts (*i.e.*, *cpr5-2* for mutant VPg/D113G and *jin1* for mutant VPg/R118H). Fig. 9

summarizes the results of these experiments. Regarding VPg/D113G (Fig. 9A - C), the

results strongly depend on the combination of phenotypic trait and plant genotype: it

shows a significant negative effect in both *AUDPS* (-14.2%; $z = 1.687$, $P = 0.046$) and *I*

(-23.9%; $z = 6.344$, $P < 0.001$) but a positive one in *SS* (16.7%; $z = 2.597$, $P = 0.005$)

when evaluated in WT plants. However, it shows a largely negative yet not significant

effect in *cpr5-2* plants for *AUDPS* (-60.8%; $z = 1.055$, $P = 0.146$) and significant negative

effects both for *I* (-60.1%; $z = 10.614$, $P < 0.001$) and *SS* (-46.7%; $z = 3.228$, $P = 0.001$).

These negative effects on the local host genotype *cpr5-2* do not support a possible

beneficial effect of this mutation by itself. Regarding the possible adaptive effect of the

amino acid replacement VPg/R118H (Fig. 9D - F), the results are also dependent on the

combination of phenotypic trait and host plant genotype. When tested in WT plants, the

mutation has significant negative effects both for *AUDPS* (-11.4%; $z = 2.29$, $P = 0.013$)

and *I* (-9.0%; $z = 2.749$, $P = 0.003$) but a significantly positive effect for *SS* (43.6%; $z =$

7.881, $P < 0.001$). By contrast, in the local host *jin1* the effect was positive in all three

traits; significant for *AUDPS* (9.2%; $z = 1.683$, $P = 0.046$) and *SS* (68.2%; $z = 12.447$, P

< 0.001) but not for *I* (0.3%; $z = 1.342$, $P = 0.090$). These results support the conclusion

that the amino acid replacement VPg/R118H has an overall beneficial effect on the local

host genotype *jin1* in which it was selected.

Discussion

Interaction between TuMV and the different defense pathways

In this study we have explored the effect that mutations in host's different disease

signaling pathways (*R* genes) or in recessive resistance genes (*S*) have in the outcome of

virus evolution. Our experimental pathosystem consisted in two well studied

components: the model plant *A. thaliana* and one virus with high prevalence in natural

populations on this host (Pagán *et al.* 2010), TuMV. Among the *R* genes, we have

included in our study genes involved in the SA, JA/ET and RNA-silencing pathways,

among the *S* genes, we have included heat shock proteins, transcription factors and

components of the translation machinery. In a preliminary set of experiments, we

stablished the similarities and differences in phenotypic responses of different plant

mutant genotypes to TuMV infection. Our results were, in some cases, at odds with the

expected phenotypes for *R* genes, being the results for *jin1* the most unexpected ones.

JASMONATE INSENSITIVE 1 (*JIN1*) is a negative regulator of SA-mediated defense

responses, henceforth the mutant *jin1* has a constitutive expression of SAR (Laurie-Berry

et al. 2006). Surprisingly, it turns out to be the most sensitive genotype to TuMV

infection, showing enhanced symptoms of disease (phenotypic group G5). This may

reflect the fact that the function of *R* genes, especially those involved in SAR and ISR,

have been mostly defined in terms of plant interactions with biotrophic and necrotrophic

bacteria and fungi but rarely in response to virus infections. Mutant genotypes only

affecting components of the ISR pathway (*e.g.*, *coi1-4*) behaved as wild-type plants in

response to TuMV infection, thus confirming previous reports that ISR was inefficient

against viral infections (Ton et al. 2002; Loebenstein 2009; Pieterse *et al*. 2009).

However, mutant *eds8-1*, which avoids ISR but enhances SAR (Love *et al*. 2007) turned

out to be among the most resistant genotypes to infection (phenotypic group G1). In

agreement with our findings of a variable response to SA-signaling, Singh *et al*. (2004)

discussed examples of viruses in which SA-dependent responses to viral infection were

strongly dependent on the virus used in the experiments: while viruses such as alfalfa

mosaic virus, potato virus X or turnip vein clearing virus replication was inhibited by

treating plants with SA, cucumber mosaic virus was apparently unaffected by the

treatment and accumulated to normal levels.

The RNA-silencing pathway is considered as the main plant defense against viral

pathogens (Voinnet 2001), with *DICER LIKE 2* (*DCL2*) and *DICER LIKE 4* (*DCL4*)

encoding for the two dicer enzymes responsible for generating the 22- and 21-nucleotides

long antiviral siRNAs. Therefore, the double mutant *dcl2 dcl4* was expected to show a

highly sensitive response to TuMV infection. However, infected plants were, overall,

hardly distinguishable from the WT plants in their response to infection (Fig. S2 and Fig.

S3). A possible explanation would be the strong suppressor activity of HC-Pro in the WT

plants that effectively counteracts the defense mechanism (Kaschau & Carrington 2001).

Indeed, it has been shown that the outcome of the interplay between plant RNA-silencing

response and potyvirus is mostly driven by the efficiency of the viral HC-Pro suppression

activity (Li *et al*. 2007; Torres-Barceló *et al*. 2008).

The response to infection of genotypes carrying mutations in *S* genes was, in

general, consistent with the *a priori* expectation. Mutant *i4g2*, that is defective for the

eIF4(iso)E factor, shows a strong resistance to TuMV infection (Nicaise *et al*. 2007;

Charron *et al*. 2008), belonging to phenotypic group G1. Likewise, in agreement with

previous descriptions, mutant *dbp2* shows enhanced resistance to potyvirus infection

(Castelló *et al*. 2011), belonging to phenotypic group G3. Finally, mutant *hsp90-1* turns

out to show a response to TuMV infection equivalent to the WT plants, likely due to its

functional redundancy with other heat-shock proteins present in the cell that may be

coopted by the virus to assist in their expected functions (Verchot 2012).

An interesting finding was the effect on TuMV replication of knocking down the

expression of gen *P58IPK*. In mammals P58[IPK], a tetratricopeptide repeat (TPR)-

containing protein, is recruited by viruses (*e.g.*, influenza A virus) to inhibit interferon

activation and cell death mediated by the dsRNA-activated protein kinase (PKR), thus

favoring viral spread (Goodman *et al*. 2011). In our experiments, *p58IPK* mutant plants

show enhanced tolerance to infection, showing weaker symptoms than WT plants despite

accumulating viruses at the same level (Fig. S1). In sharp contrast, Bilgin *et al*. (2003)

found that infection of *p58IPK*-silenced *Nicotiana benthamiana* and *A. thaliana* plants

infected with tobacco mosaic virus lead to death. Given the limited information available

from other plant-virus pathosystems, additional experiments would be necessary to shed

light into the potential antiviral role in plants of this gene.

One side, yet important, conclusion of our study is that the generally accepted

model of plant defense signaling pathways, mainly based in experiments done with

bacteria and fungi, may not well describe the interaction between plants and viruses. By

contrast, recessive resistances based on *S* genes seem to better explain differences in

susceptibility to infection and viral accumulation, thus being more promising targets for

future development of resistant plants.

Evolution of specialist and generalist viral strategies depend on the host genotype

The GFG and MA models of host-virus interaction modes represent the two ends of a

continuum of possible outcomes (Agrawal & Lively 2002). The most relevant difference

between both models regards the expected genetic heterogeneity in both host and virus

populations. With a pure GFG interaction the susceptible host types are expected to

disappear and the resistant types will dominate the population. *Vice versa*, the most

virulent virus allele would drive to fixation at the cost of losing mild alleles. However,

constitutive activation of defenses is known to be costly for *A. thaliana* [*e.g.*, SA-related

defense responses pay a fitness cost in absence of pathogens (Traw *et al.* 2007)] and high

virulence usually comes with a cost in terms of pathogen's transmission (reviewed in

Acevedo *et al.* 2019). Hence, a pure GFG strategy seems unlikely to be achieved. By

contrast, with a pure MA interaction, negative frequency-dependent selection emerges,

such that rare *A. thaliana* resistance alleles have advantage and, as a result, a genetic

polymorphism shall be maintained (Schmid-Hempel 2011). Natural populations of *A.*

thaliana contain considerable amount of genetic variability for tolerance (Pagán *et al.*

2008) and for immunity-related genes (Todesco *et al.* 2010, van de Weyer *et al.* 2019),

thus suggesting that an arms race between pathogens and plants should be going on yet it

is unclear whether it may result in pure GFG or MA interactions or lie somewhere in

between. Evolution experiments with such different pathosystems as TuMV/*A. thaliana*

(González *et al.* 2019), *Octosporea bayeri*/*Daphnia magna* (Altermatt & Ebert 2008) and

Serratia marcescens/*Caenorhabditis elegans* (Gibson et al. 2020; White *et al.* 2020) have

produced congruent results in one aspect: parasites exposed to heterogeneous host

populations evolved significantly lower virulence than parasites exposed to homogeneous

host populations. However, a significant difference exists among pathosystems: while

viruses exposed to genetically heterogenous host populations evolved as no-cost

generalists, evolution of generalism in more complex parasites was constrained by a

fitness tradeoff, as expected for the jack-of-all trades hypothesis (Bedhomme *et al*. 2015).

Our results, as well as those by Hillung *et al*. (2014) and González *et al*. (2019)

have shown the evolution of significantly nested binary infection networks, a finding

compatible with the existence of a combination of specialist and generalist viruses and of

more permissive and resistant host genotypes. Indeed, these studies have also shown that

more permissive hosts selected for more specialized viruses while more resistant hosts

selected for more generalist viruses, here matching the predictions of the GFG model.

Our observation of small yet significant modularity in the infection network was easily

explained by convergent evolution of TuMV lineages evolved in the same host genotype.

However, it has been recently shown in a long-term survey of the prevalence of different

plant viruses in different hosts and habitats that nestedness and modularity in host-

pathogen infection networks is possible due to the spatially patched distribution of

habitats and temporal successions of plant species (Valverde *et al*. 2020): small spatial

scales create modularity that coexist with global nestedness. This pattern may change

spatially and temporally but remains stable over long evolutionary timescales.

Role of natural selection

We found evidences of significant genetic differentiation among TuMV lineages evolved

in different plant genotypes. To test whether these differences were driven by selection

we performed a Tajima's D test (Tajima 1989). The resulting D value was significative

and negative, which is compatible with the action of purifying selection, the presence of

slightly deleterious mutations segregating in the populations or fast population

expansions (Yang 2006). How to distinguish between these explanations? In

independent fast expanding populations, many new mutations may be generated and

rising in frequency in each population, thus being observed as singletons, mutations

present in only one of the many coexisting genomes in each evolving TuMV lineage.

Singletons inflate the number of segregating sites and thus cause $D < 0$. Indeed, this is

the case here: 54 out of the 71 observed variable sites are singletons, thus the observed

pattern of molecular diversity among lineages evolved in the same host genotype and in

different host genotypes is likely to be due to the fast expansions of viral populations.

However, we have found additional evidence supporting the action of positive

selection: the existence of a number of convergent nonsynonymous mutations arising in

independent lineages, some evolved in the same host genotype but some others rising in

different host genotypes (Table S1 and Fig. S3) yet without a clear association with the

particular signaling pathway or *S* gene being mutated (Table 1) nor the phenotypic cluster

they belong to (Fig. 1). Interestingly, most of these convergent mutations happened in

the *VPg* cistron, which turns out to be also the most variable one. VPg plays many

essential roles in genome transcription (it is linked to the 5'-end of the viral genome and

provides the hydroxyl group that primes the synthesis of the complementary strains by

the viral RdRp), translation (directly interacts with the eukaryotic initiation factors

eIF(iso)4E and eIF(iso)4G) and interacts with all other viral proteins (Bosque *et al*. 2014)

and some other cell proteins (Martínez *et al*. 2016). Indeed, in previous evolution

experiments with potyviruses, *VPg* has also been shown to be an important target of

selection. For example, Agudelo-Romero *et al*. (2008) found that a single amino acid

replacement in VPg was enough to largely increase TEV *I*, *VL* and *SS* in *A. thaliana*.

Similarly, Gallois *et al*. (2010) found that *A. thaliana* plants with knock-out mutations in

the *eIF(iso)4E*, *eIF(iso)4G1* and *eIF(iso)4G2* genes were resistant to TuMV infection.

Two mutations in the VPg (E116Q and N163Y) were enough to overcome this resistance

and return to the original infection phenotype, though yeast-two hybrid assays showed

that none of these mutations affected the binding of VPg with eIF(iso)4E. As a final

example, one of the most extensively used resistance genes in pepper commercial

cultivars against potato virus Y (PVY; genus *Potyvirus*, family *Potyviridae*) is the *pvr2*,

which has many different alleles (Nicaise *et al*. 2007; Charron *et al*. 2008). The *pvr2*

locus encodes for the eIF4E factor which, as mentioned above, physically interacts with

VPg. Interestingly, all the resistance-breaking viral isolates found so far contain

mutations in the *VPg* cistron (Duprat *et al*. 2002; Moury *et al*. 2004; Ayme *et al*. 2006)

Interestingly, one of the two mutations identified by Gallois *et al*. (2010),

VPg/E116Q, affects the same protein domain that the two mutations we have identified

and characterized (D113G and R118H). The mechanism why these two mutations may

confer a selective advantage to TuMV lineages cannot be inferred from our studies. We

found VPg/D113G in several lineages evolved in different host genotypes, while

VPg/R118H was only found in the *jin1* lineages. Our assays failed to find a beneficial

effect of VPg/D113G but confirmed the host-specific beneficial effect of VPg/R118H in

jin1 plants.

Unfortunately, due to economical constrains, we have not been able to exhaustively

testing all observed mutations and, hence, several others may still be potential candidates

for adaptive mutations. More specifically, we have not tested possible epistatic

interactions among mutations fixed in the same genome. Three lineages carry more than

one nonsynonymous mutation in *VPg* (*dbp2*/L1 D113N and N115E, *eds8-1*/L1 N115H

and E116G, and *eds8-1*/L4 H33Y, D113N and K121E) (Table S1). In addition, 25

lineages carry additional nonsynonymous mutations in at least one other cistron besides

VPg (Table S1). Two lineages have up to three additional nonsynonymous mutations:

cpr5-2/L5 that has P3/T318M-T326S CI/V148I and VPg/D113G and *dcl2 dcl4*/L1 that

has P1/V54A CI/I378V VPg/R118C and CP/S70N. Therefore, plenty of opportunities

for epistatic effects both within and among cistrons exist that will be evaluated in future

experiments.

Methods

Plants, virus and growth conditions

A collection of 21 different *A. thaliana* mutants of the Col-0 accession were used for this

study (Table 1). In all experiments described below, plants were all maintained in a BSL-

2 climatic chamber under a photoperiod of 8 h light (LED tubes at PAR 90 - 100

μmol/m^2/s) at 25 ºC and 16 h dark at 20 ºC.

Prior to the inoculation experiments, we created a large stock of TuMV infectious

saps. Saps were obtained from TuMV-infected *N. benthamiana* Domin plants inoculated

with the infectious plasmid p35STunos that contains a cDNA of TuMV genome

(GeneBank accession AF530055.2) under the control of the cauliflower mosaic virus 35S

promoter and the *nos* terminator (Chen *et al.* 2003) as described elsewhere (González *et*

al. 2019; Corrêa *et al.* 2020). This TuMV sequence variant corresponds to the YC5

isolate from calla lily (*Zantesdeschia* sp) (Chen *et al.* 2003). After plants showed

symptoms of infection, they were pooled frozen with liquid N$_2$ and homogenized into a

fine powder using a Mixer Mill MM400 (Retsch GmbH, Haan, Germany). For

inoculations, the necessary amount of power was diluted in inoculation buffer (50 mM

phosphate buffer pH 7.0, 3% PEG6000, 10% Carborundum) and 5 μL of the inoculum

was gently rubbed into the plant leaves. Plants were all inoculated when reached growth

stage 3.5 in the Boyes' scale (Boyes 2001). This synchronization ensures that they all

were at the same phenological state when inoculated.

Phenotyping infected plants

The following five different traits were measured for each infected plant 18 dpi. (*i*) Change in dry weight (ΔDW) of the aerial part of infected plants, with a precision of 10 mg, relative to the corresponding noninfected controls (Fig. S1A). (*ii*) Severity of symptoms (*SS*) was evaluated in a semi-quantitative discrete scale (Fig 1B in Corrêa *et al*. 2020) ranging from zero for asymptomatic infections to four for plants showing a generalized necrosis and wilting (Fig. S1B). (*iii*) The area under the disease progress stairs (*AUDPS*) that summarizes the speed at which the disease severity progresses in a group of plants (Fig. S1C) and takes values in the range zero and the total number of plants included in the assay (Simko & Piepho 2012). (*iv*) Infectivity measured as the number of symptomatic plants out of the number of inoculated plants 18 dpi (*I*; Fig. S1D). And (*v*) viral load (*VL*) measured by absolute RT-qPCR as the number of viral genomes per ng of total RNA in the plant as described below (Fig. S1E).

The five traits were not all independent but showed some significant pairwise positive correlations: *AUDPS* with *I* ($r = 0.772$, 19 df, $P < 0.001$), *SS* ($r = 0.807$, 19 df, $P < 0.001$) and *VL* ($r = 0.472$, 19 df, $P = 0.031$), and *I* with *SS* ($r = 0.746$, 19 df, $P < 0.001$). Basically, ΔDW is orthogonal with the other four traits while the other four traits show some degree of association. Interesting is the case of *VL*, which is only (weakly) correlated to *AUDPS*.

Experimental evolution

Five TuMV lineages were evolved during 12 consecutive serial passages on each one of the nine selected mutant genotypes. To begin the evolution experiment, 10 21-days old *A. thaliana* plants per lineage, three leaves per plant, were inoculated as described above with the sap from *N. benthamiana*. Fourteen dpi, symptomatic *A. thaliana* plants were

harvested and used to prepare infectious saps as described above. These saps were 1/10

diluted in inoculation buffer and used to inoculate the next batch of plants.

Total RNA extractions

Tissue from each pool of infected plants per lineage, ecotype and serial passage was

collected, frozen with liquid N_2 and preserved at -80 ºC until it was homogenized into

fine powder using a Mixer Mill MM400. Next, an aliquot of approximately 100 mg of

grounded tissue per sample for which viral load was quantified, was used for total RNA

(RNAt) extraction performed with the Agilent Plant RNA isolation Mini kit (Agilent

Technologies, Santa Clara CA, USA). Aliquots of RNAt per each sample were separated

and their concentration adjusted at 50 ng/µL to estimate viral accumulation by RT-qPCR

(see below).

Quantification of *VL*

VL of each plant sample per lineage, ecotype and selected passage was quantified by

absolute real-time quantitative RT-PCR (RT-qPCR) using standard curves and the

primers TuMV F117 forward (5'-CAATACGTGCGAGAGAAGCACAC-3') and F118

reverse (5'-TAACCCCTTAACGCCAAGTAAG-3') that amplify a 173 nucleotides

fragment from the *CP* cistron of TuMV genome, as previously described (Corrêa *et al*.

2020). Briefly, standard curves were constructed using ten serial dilutions of the TuMV

genome, that was synthesized by *in vitro* transcription as detailed previously (Cervera *et*

al. 2018), in RNAt extract obtained from healthy *A. thaliana* plants used as control in the

experiments. Amplification reactions were run in a 20 µL volume using the GoTaq 1-

Step RT-qPCR System (Promega, Madison WI, USA) and the recommended

manufacturer's instructions as described (Cervera *et al*. 2018) in an ABI StepOne Plus

Real-time PCR System (Applied Biosystems, Foster City CA, USA). The cycling

conditions consisted in: an RT phase of 5 min at 42 ºC and 10 min at 9 5ºC followed by

a PCR stage consisting in 40 cycles of 5 s at 95 ºC and 34 s at 60 ºC; and the final melt

curve profile analysis that consisted in 15 s at 95 ºC, 1 min at 60ºC and 15 s at 95 ºC.

Negative controls consisted in healthy RNAt plant extract (non infected control) and

water. Quantitative reactions were run as three technical replicates per sample and results

were analysed using the StepOne software 2.2.2 (Applied Biosystems).

TuMV genome amplifications

Evolved viral genomes of the passage 12 from each lineage and ecotype were amplified

by high-fidelity RT-PCR by using the AccuScript Hi-Fi (Agilent Technologies) reverse

transcriptase and Phusion DNA polymerase (Thermo Scientific, Waltham MA, USA)

following the manufacturer's instructions. Each complete TuMV genome was amplified

into three overlapping amplicons of 3114 (5' fragment R1), 3697 (central region R2) and

3287 nucleotides (3' fragment R3) using three primer sets. For RT reactions an aliquot

of the corresponding RNAt (150-300 ng) was mix with 0.25 μM of the 1R-P3 (5'-

CGAGTAGTATCTTATAGCACAGCGCTCCGACC-3'), 2R-NIa (5'-

TGTCTGGAATCGGTAGCAAATGTAGCTGAGTTGTG-3') or 3R-polyAR (5'-

TTTTTTTTTTTTTTTTTTTTGTCCCTTGCATCATATCAAATG-3') primer to

synthesize the R1, R2 or R3 cDNA fragment, respectively, that were denatured 5 min at

65 ºC and cooled on ice. Then it was added a mix containing AccuScript Hi-Fi 1× Buffer,

1 mM of dNTPs, 8 mM of DTT, 4U of Ribolock RNase inhibitor (Thermo Scientific) and

0.5 μL of AccuScript Hi-Fi (Agilent Technologies) in a 10 μL volume. RT reactions

consisted in a step of 90 min at 42 ºC to synthesize the cDNA followed by an incubation

of 5 min at 70 ºC to inactivate the enzyme. PCR reactions were performed in a 50 μL

volume containing a mix of 1× Phusion Buffer, 0.4 μM of dNTPs, 0.2 μM of each primer,

0.5-1 μL of DMSO, 2U of Phusion DNA polymerase (Thermo Scientific) and 1 μL of the

corresponding RT reaction. R1 fragment was amplified using the primer set 1F-5UTR

(5'-GCAAACGCAGACCTTTCGAAGCACTCAAGC-3') and 1R-P3 and the following

PCR conditions: an initial denaturation of 30 s at 98 ºC, 3 cycles of 10 s at 98 ºC, 20 s at

67 ºC and 2 min at 72 ºC, 3 cycles of 10 s a 98 ºC, 20 s at 65 ºC and 2 min at 72 ºC, and

32 cycles of 10 s at 98 ºC, 20 s at 63 ºC and 2 min at 72 ºC, followed by a final extension

step of 5 min at 72 ºC. Fragments R2 and R3 were amplified with primer set 2F-P3 (5'-

TGGGAGCTTGCGGATGGTGGATACACAATTC-3') and 2R-NIa or 3F-NIa (5'-

CTCGTTATATGGAGTCGGTTTCGGACCACTCATAT-3')   and   3R-polyAR,

respectively and a PCR with the same denaturation and extension steps than the fragment

R1 but different amplification steps: an stage consisting in 15 cycles of 10 s at 98 ºC, 20

s at 67 ºC and 2 min at 72 ºC followed by 23 cycles of 10 s at 98 ºC, 20 s at 65 ºC and 2

min at 72 ºC for fragment R2, while 15 cycles of 10 s at 98 ºC, 20 s at 67 ºC and 2 min at

72 ºC followed by 23 cycles of 10 s at 98 ºC, 20 s at 65 ºC and 2 min at 72 ºC for fragment

R3. PCR products were purified with the MSB Spin PCRapace Kit (Stratec Molecular,

Coronado CA, USA) and then Sanger-sequenced. Full-length consensus viral sequences

were obtained assembling the sequences of the three amplified products by using the

Genious R9.0.2 program.

Construction of TuMV *VPg* mutants

Mutants in the *VPg* cistron were synthesized by long inverse site-directed PCR

mutagenesis using the QuickChange II XL Kit (Stratagene, San Diego CA, USA)

following the manufacturer's instructions using p35STunos as template. Also, mutagenic

primers were designed following the manufacturer's recommendations. Primer set cpr5-

2F (5'-GGAGGATGAGTTGGGTCCAAATGAAATACGTGT-3') and cpr5-2R (5'-

ACACGTATTTCATTTGGACCCAACTCATCCTCC-3') was used to generate the

A6238G (D113G) mutant while primer set jin1F (5'-

GGATCCAAATGAAATACATGTGAATAAGACAATTC-3') and jin1R (5'-

GAATTGTCTTATTCACATGTATTTCATTTGGATCC-3') to obtain the G6253A

(R118H) mutant. PCR protocol consisted in: a denaturation step of 2 min at 95 ℃

followed by 20 cycles of 20 s at 95 ℃, 10 s at 60 ℃ and 7 min at 68 ℃ and a final

extension of 5 min at 68 ℃. After *Dpn*I digestion and transformation of electrocompetent

Escherichia coli DH5α, the presence of the desired mutations in the infectious clone and

the absence of undesired nucleotide changes was confirmed by sequencing.

Bioassays of the TuMV *VPg* mutants

The infectivity, viability and pathogenicity of both VPg TuMV mutants was confirmed

by performing three independent bioassays. In each experiment, three batches of 24 *A.*

thaliana plants (three-weeks old) of genotypes wild-type, *cpr5-2* and *jin1* were inoculated

with the wild-type TuMV plasmid clone used as reference and with plasmids of each of

the TuMV VPg mutants, while another two batches of 24 plants from *cpr5-2* or *jin1*

genotypes were inoculated with the WT TuMV plasmid and the corresponding plasmid

of the TuMV VPg mutant, the VPg/D113G mutant in the case of the *cpr5-2* ecotype and

that of VPg/R118H for the *jin1* genotype. *A. thaliana* were inoculated by abrasion of

three leaves applying equal amounts of the plasmid inoculum, a total of approximately 7

μg per plant distributed in 3 μL per leaf. Plasmids were previously purified by Midiprep

using the NucleoBond Xtra Midi Kit (Macherey-Nagel, Düren, Germany) and

resuspended in distilled water. Number of infected plants and symptom intensity were

collected every day until 15 dpi.

Statistical analyses

All statistical analyses described hereafter were performed with SPSS version 26 software (IBM, Armonk, NY), unless otherwise indicated.

AUDPS (Fig. 2), *I* (Fig. 3) and *VL* (Fig. 4) data were fitted all together to a multivariate analysis of covariance (MANCOVA) model in which plant genotype (*G*) was the main factor, the independent evolution lineages (*L*) were nested within *G* and passage (*t*) was introduced in the model as a covariable. The full model equation thus reads:

$$\mathbf{P}_{ijk}(t) \sim \mathbf{\mu} + t + G_i + L(G)_{ij} + (t \times G)_i + [t \times L(G)]_{ij} + \mathbf{\varepsilon}_{ijk}, \qquad \text{(Eq. 1)}$$

where $\mathbf{P}_{ijk}(t) = (AUDPS_{ijk}, I_{ijk}, VL_{ijk})^{\mathrm{T}}$ is the vector of phenotypic traits observed at time *t*, for an individual infected plant *k* of evolutionary lineage *j* of genotype *i*, $\mathbf{\mu}$ represents the vector of phenotypic grand mean values and $\mathbf{\varepsilon}_{ijk}$ stands for the vector of errors assumed to be Gaussian distributed at every *t*. The significance of each factor, the covariable and their interactions was evaluated using the Wilks' Λ method. The magnitude of the effects was evaluated using the η_P^2 statistic (proportion of total variability in the traits vector attributable to each factor in the model; conventionally, values of $\eta_P^2 \geq 0.15$ are considered as large effects).

Rates of phenotypic evolution for *AUDPS* and *I* were estimated by fitting the time series data to a first-order autoregressive integrated moving-average, ARIMA(1,0,0), model (Elena and Sanjuán, 2005; González *et al.* 2019). The model equation fitted has the form:

$$Y_t - \rho_1 Y_{t-1} = Y_0 + \beta_Y t + \varepsilon_t, \qquad \text{(Eq. 2)}$$

where Y_k represents the variable being analyzed at passage *k*, ρ_1 measures the degree of self-similarity in the time-series data (correlation between values at passages *t* and *t* − 1),

ε_t represents the sampling error at passage t, and β_Y represents the linear dependency of

variable Y with passage number, that is, the rate of phenotypic evolution. The rates of

phenotypic evolution were further analyzed in the context of the mutated defense

signaling pathway encoded by R genes or recessive resistances encoded by S genes. Rates

of phenotypic evolution were fitted to a multivariate analysis of variance (MANOVA) in

which the plant genotype serving as host for the evolutionary lineages (G) was nested

within the factor being analyzed (*i.e.*, the type of selection –hard *vs* soft-, the mode of

resistance –R *vs* S-, and whether mutations affected the SA-dependent, the JA/ET-

dependent signaling or the RNA-silencing pathways; Table 1). The full model equation

now thus reads:

$\mathbf{B}_{ijk} \sim \boldsymbol{\mu} + X_{ij} + X(G)_{ij} + \boldsymbol{\varepsilon}_{ijk},$ (Eq. 3)

where $\mathbf{B}_{ijk} = (\beta_{AUDPS}, \beta_I)^{\mathrm{T}}$ is the vector of rates of phenotypic evolution observed for the

k replicate lineage evolved in plant genotype j within the factor X_i, as above, $\boldsymbol{\mu}$ represents

the vector of grand mean values and $\boldsymbol{\varepsilon}_{ijk}$ the vector of Gaussian errors.

Analysis of infection the network

The first statistical approach consisted in fitting a logistic regression model to the

presence/absence of symptoms data using GLM techniques with a Binomial probability

distribution and a probit link function. The model equation reads as follows:

$\mathrm{probit}\left[f_{ijkl}(t)\right] \sim \phi + t + TG_i + t \times TG_i + LG_j + t \times LG_j + L(LG)_{jk} + t \times L(LG)_{jk} +$

$(TG \times LG)_{ij} + t \times (TG \times LG)_{ij} + [TG \times L(LG)]_{ijk} + t \times [TG \times L(LG)]_{ijk} + \varepsilon_{ijkl},$

(Eq. 4)

where $f_{ijkl}(t)$ is the frequency of symptomatic plants t dpi in the test host genotype TG, for

the viral lineage L evolved in the local host genotype LG. TG and LG were considered

as orthogonal factors, L was nested within LG and dpi (t) was treated as a covariable. The

model includes all main factors, their corresponding nested and factorial interactions as

well as their interactions with the covariable. ϕ represents the grand mean of the probit-

transformed f values, and ε the error term assumed to be Binomial. The significance of

the different factors was evaluated using likelihood-ratio tests (LRT).

For the second statistical approach, the binary infection matrix was analyzed using

tools borrowed from the field of network biology to explore whether they show random

associations between viral lineages and host genotypes, one-to-one associations,

nestedness indicative of a GFG type of interaction, or modularity (Weitz et al. 2013). The

statistical properties of the infection matrix were evaluated using the R package

"bipartite" version 2.15 (Dormann et al. 2008) in R version 4.0.0 (R Core Team 2020)

under RStudio 1.2.1335. Four different summary statistics were evaluated: T nestedness

(Bascompte et al. 2003), Q modularity (Newman 2006) and the d' species-level (or

Kullback-Leibler divergence) and H_2' network-level (or two-dimensional normalized

Shannon entropy) specialization indexes (Blüthgen et al. 2006). H_2' ranges between zero

and one for extreme generalists and specialists, respectively. Statistical significance of

these statistics was evaluated using Bascompte et al. (2003) null model.

The statistics of molecular evolution

Treating each lineage as an observation and each host genotype as a subpopulation, we

evaluated the average nucleotide diversity within host genotypes, π_S, the nucleotide

diversity for the entire sample, π_T, the interhost genotypes nucleotide diversity, δ_{ST}, and

the estimate of the proportion of interhost genotypes nucleotide diversity, known as

coefficient of nucleotide differentiation (Nei 1982), $N_{ST} = \delta_{ST}/\pi_T$. Standard deviations of

estimates were inferred from 1000 bootstrap samples. All these computations were done

using MEGA X (Kumar et al. 2018) and the lowest-BIC nucleotide substitution model

Kimura 2-parameters (Kimura 1980). Tajima's *D* test of selection (Tajima 1989) and its

statistical significance were evaluated using DnaSP6 (Rozas *et al*. 2017).

The frequency of mutations (*m*) per cistron (*C*), relative to the length of the

corresponding cistron, was fitted to the following logistic regression model using GLM

techniques with a Binomial probability distribution and a probit link function:

$\text{probit}(m_i) \sim \mu + C_i + \varepsilon_i$ (Eq. 5)

where μ is the average genomic mutation frequency and *i* refers to the 10 cistrons in the

main ORF.

The functional effect of mutations found in the *VPg* cistron were evaluated *in silico*

using the SNAP2 algorithm (Hecht *et al*. 2015). The algorithm provides a functionality

score and an accuracy index for each possible mutation affecting a coding sequence.

Negative score values mean functional neutrality while positive values should be taken

as indications of functional effects; the larger the values, the stronger the effect.

References

Acevedo MA, Dillemuth FP, Flick AJ, Faldyn MJ, Elderd BD. 2019. Virulence-driven

trade-offs in disease transmission: a meta-analysis. *Evolution* **73**: 636-647.

Agrawal A, Lively CM. 2002. Infection genetics: gene-for-gene versus matching-alleles

models and all points in between. *Evol. Ecol. Res*. **4**: 79-90.

Agudelo-Romero P, Carbonell P, Pérez-Amador MA, Elena SF. 2008. Virus adaptation

by manipulation of host's gene experession. *PLoS ONE* **3**: e2397.

Altermatt F, Ebert D. 2008. Genetic diversity of Daphnia magna populations enhances

resistance to parasites. *Ecol. Lett*. **11**: 918-928.

Altizer S, Kaitala V, Laakso J, Ruokolainen L. 2006. Seasonality and the dynamics of

infectious diseases. *Ecol. Lett*. **9**: 467-484.

Anttila J, 2015. Environmental variation generates environmental opportunist pathogen

outbreaks. *PLoS ONE* **10**: e0145511.

Ayme V, Souche S, Caranta C, Jacquemond M, Chadoeuf J, Palloix A, Moury B (2006)

Different mutations in the genome-linked protein VPg of *Potato virus Y* confer

virulence on the *pvr2³* resistance in pepper. *Mol. Plant Microbe Interact.* **19**: 557-

563.

Bascompte J, Jordano P, Melián CJ, Olesen JM. 2003. The nested assembly of plant-

animal mutualistic networks. *Proc. Natl. Acad. Sci. USA* **100**: 9389-9387.

Bedhomme S, Hillung J, Elena SF. 2015. Emerging viruses: why are not jacks of all

trades? *Curr Opin Virol* **10**: 1-6.

Bilgin DD, Liu Y, Schiff M, Dinesh-Kumar SP. 2003. *P58^{IPK}*, a plant ortholog of double-

stranded RNA-dependent protein kinase PKR inhibitor, functions in viral

pathogenesis. *Dev. Cell* **4**: 651-661.

Blüthgen N, Menzel F, Blüthgen N. 2006. Measuring specialization in species interaction

networks. *BMC Ecol.* **6**: 9.

Bosque G, Folch-Fortuny A, Picó J, Ferrer A, Elena SF. 2014. Topology analysis and

visualization of *Potyvirus* protein-protein interaction network. *BMC Syst. Biol.* **8**:

129.

Bouché N, Lauressergues D, Gasciolli V, Vaucheret J. 2006. An antagonistic function for

Arabidopsis DCL2 in development and a new function for DCL4 in generating viral

siRNAs. *EMBO J.* **25**: 3347-3356.

Boyes DC. 2001. Growth stage-based phenotypic analysis of *Arabidopsis*: a model for

high throughput functional genomics in plants. *Plant Cell* **13**: 1499-1510.

Brown JKM, Tellier A. 2011. Plant-parasite coevolution: bridging the gap between

genetics and ecology. *Annu. Rev. Phytopathol.* **49**: 345-367.

Cao H, Bowling SA, Gordon AS, Dong X. 1994. Characterization of an *Arabidopsis*

mutant that is nonresponsive to inducers of systemic acquired resistance. *Plant Cell*

6: 1583-1592.

Carr JP, Lewsey MG, Palukaitis P. 2010. Signaling in induced resistance. *Adv. Virus Res*.

76: 57-121.

Castelló MJ, Carrasco JL, Navarrete-Gómez M, Daniel J, Granot D, Vera P. 2011. A plant

small polypeptide is a novel component of DNA-binding protein phosphatase 1-

mediated resistance to *Plum pox virus* in *Arabidopsis*. *Plant Physiol*. **157**: 2206-

2215.

Cervera H, Ambrós S, Bernet GP, Rodrigo G, Elena SF. 2018. Viral fitness correlates

with the magnitude and direction of the perturbation induced in the host's

transcriptome: the tobacco etch potyvirus-tobacco case study. *Mol. Biol. Evol*. **35**:

1599-1615.

Charron C, Nicolaï, M, Gallois JL, Robaglia C, Moury B, Palloix A, Caranta, C. 2008.

Natural variation and functional analyses provide evidence for coevolution between

plant eIF4E and potyviral VPg. *Plant J*. **54**: 56-68.

Chen CC, Chao CH, Chen CC, Yeh SD, Tsai HT, Chang CA. 2003. Identification of

Turnip mosaic virus isolates causing yellow stripe and spot on calla lily. *Plant Dis*.

87: 901-905.

Corrêa RL, Sanz-Carbonell A, Kogej Z, Müller SY, Ambrós S, López-Mogollón S,

Gómez G, Baulcombe DC, Elena SF. 2020. Viral fitness determines the magnitude

of transcriptomic and epigenomic reprogramming of defense responses in plants.

Mol. Biol. Evol. **37**: 1866-1881.

Cui H, Qiu J. Zhou Y, Bhandari DD, Zhao C, Bautor J, Parker JE. 2018. Antagonism of

transcription factor MYC2 by EDS1/PAD4 complexes bolsters salicylic acid defense

in *Arabidopsis* effector-triggered immunity. *Mol. Plant* **11**: 1053-1066.

Dormann CF, Gruber B, Fruend J. 2008. Introducing the bipartite package: analyzing

ecological networks. *R News* **8**: 8-11.

Dormann CF, Strauss R. 2014. A method for detecting modules in quantitative bipartite

networks. *Meth. Ecol. Evol*. **5**: 9-98.

Duprat A, Caranta C, Revers F, Menand B, Browning KS, Robaglia C. 2002. The

Arabidopsis eukaryotic initiation factor (iso)4E is dispensable for plant growth but

required for susceptibility to potyviruses. *Plant J*. **32**: 927-934.

Elena SF, Sanjuán R. 2005. RNA viruses as complex adaptive systems. *BioSystems* **81**:

31-41.

Flor HH. 1956. The complementary genetic systems in flax and flax rush. *Adv. Genet*. **8**:

29-54.

Frank SA. 1993. Specificity versus detectable polymorphism in host-parasite genetics.

Proc. R. Soc. B **254**: 191-197.

Gallois JL, Charron C, Sánchez F, Pagny G, Houvenaghel MC, Moretti A, Ponz F, Revers

F, Caranta C, German-Retana, S. 2010. Single amino acid changes in the turnip

mosaic virus viral genome-linked protein (VPg) confer virulence towards

Arabidopsis thaliana mutants knocked out for eukaryotic initiation factors

eIF(iso)4E and eIF(iso)4G. *J. Gen. Virol*. **91**: 288-293.

Gibson AK, Baffoe-Bonnie H, Penley MJ, Lin J, Owens R, Khalid A, Morran LT. 2020.

The evolution of parasite host range in heterogeneous host populations. *J. Evol. Biol.*

33: 773-782.

González R, Butković A, Elena SF. 2019. Role of host genetic diversity for susceptibility-

to-infection in the evolution of virulence of a plant virus. *Virus Evol.* **5**: vez024.

Goodman AG, Tanner BCW, Chang ST, Esteban M, Katze MG. 2011. Virus infection

rapidly activates the P58^{IPK} pathway, delaying peak kinase activation to enhance viral

replication. *Virology* **417**: 27-36.

Gupta V, Willits MG, Glazebrook J. 2000. *Arabidopsis thaliana EDS4* contributes to

salicylic acid (SA)-dependent expression of defense responses: evidence for

inhibition of jasmonic acid signaling by SA. *Mol. Plant Microbe Interact.* **13**: 503-

511.

Hecht M, Bromberg Y, Rost B. 2015. Better prediction of functional effects for sequence

variants. *BMC Genomics* **16**: S1.

Hillung J, Cuevas JM, Valverde S, Elena SF. 2014. Experimental evolution of an

emerging plant virus in host genotypes that differ in their susceptibility to infection.

Evolution **68**: 2467-2480.

Hughes WHO, Boomsma JJ. 2006. Does genetic diversity hinder parasite evolution in

social insect colonies? *J. Evol. Biol.* **19**: 132-143.

Kachroo P, Chandra-Shekara AC, Klessig DF. 2006. Plant signal transduction and

defense against viral pathogens. *Adv. Virus Res*. **66**: 161-191.

Kaschau K, Carrington JC. 2001. Long-distance movement and replication maintenance

functions correlate with silencing suppression activity of potyviral HC-Pro. *Virology*

285: 71-81.

Kimura M. 1980. A simple method for estimating evolutionary rate of base substitutions

through comparative studies of nucleotide sequences. *J. Mol. Evol.* **16**: 111-120.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: molecular evolutionary

genetics analysis across computing platforms. *Mol. Biol. Evol.* **35**: 1547-1549.

Lalić J, Agudelo-Romero P, Carrasco P, Elena SF. 2010. Adaptation of tobacco etch

Potyvirus to a susceptible ecotype of *Arabidopsis thaliana* capacitates it for systemic

infection of resistant ecotypes. *Phil. Trans. R. Soc. B* **65**: 1997-2008.

Laurie-Berry N, Joardar V, Street IH, Kunkel BN. 2006. The *Arabidopsis thaliana*

JASMONATE INSENSITIVE 1 gene is required for suppression of salicylic acid-

dependent defenses during infection by *Pseudomonas syringae*. *Mol. Plant Microbe*

Interact. **19**: 789-800.

Lin SS, Wu HW, Jan FJ, Hou RF, Yeh SD. 2007. Modifications of the helper component-

protease of *Zucchini yellow mosaic virus* for generation of attenuated mutants for

cross protection against severe infection. *Phytopathology* **97**: 287-296.

Loebenstein G. 2009. Local lesions and induced resistance. *Adv. Virus Res*. **75**: 73-117.

Love AJ, Laval V, Geri C, Laird J, Tomos AD, Hooks MA, Milner JJ. 2007. Components

of *Arabidopsis* defense- and ethylene-signaling pathways regulate susceptibility to

Cauliflower mosaic virus by restricting long-distance movement. *Mol. Plant Microbe*

Interact. **20**: 659-670.

Mäkinen K. 2019. Plant susceptibility genes as a source for potyvirus resistance. *Ann.*

Appl. Biol. **173**: 122-129.

Martínez F, Rodrigo G, Aragonés V, Ruiz M, Lodewijk I, Fernández U, Elena SF, Daròs

JA. 2016. Interaction network of tobacco etch potyvirus NIa protein with the host

proteome during infection. *BMC Genomics* **17**, 1

Mine A, Nobori T, Salazar-Rodon MC, Winkelmüller TM, Anver S, Becker D, Tsuda K.

2017. An incoherent feed-forward loop mediates robustness and tunability in a plant

immune network. *EMBO Rep*. **18**: 464-476.

Moury B, Morel C, Johansen E, Guilbaud L, Souche S, Ayme V, Caranta C, Palloix A,

Jacquemond M. 2004. Mutations in Potato virus Y genome-linked protein determine

virulence towards recessive resistances in *Capsicum annuum* and *Lycopersicon*

hirsutum. *Mol. Plant Microbe Interact*. **17**: 322-329.

Nawrath C, Heck, S, Parinthawong N, Métraux JP. 2002. *EDS5*, an essential component

of salicylic acid-dependent signaling for disease resistance in *Arabidopsis*, is a

member of the MATE transporter family. *Plant Cell* **14**: 275-286.

Nawrath C, Métraux JP. 1999. Salicylic acid induction-deficient mutants of *Arabidopsis*

express PR-2 and PR-5 and accumulate high levels of camalexin after pathogen

inoculation. *Plant Cell* **11**: 1393-1404.

Nei M. 1982. Evolution of human races at the gene level. In *Human Genetics, part A: the*

Unfolding Genome (ed. B. Bonné-Tamir). Alan R. Liss, New York.

Newman MEJ. 2006. Modularity and community structure in networks. *Proc. Natl. Acad.*

Sci. USA **103**: 8577-8582.

Nicaise V, Gallois JL, Chafiai F, Allen LM, Schurdi-Levraud V, Browning KS,

Candresse T, Caranta C, Le Gall O, German-Retana S. 2007. Coordinated and

selective recruitment of eIF4E and eIF4G factors for potyvirus infection in

Arabidopsis thaliana. *FEBS Lett*. **581**: 1041-1046.

Pagán I, Alonso-Blanco C, García-Arenal F. 008. Host responses in life-history traits and

tolerance to virus infection in *Arabidopsis thaliana*. *PLoS Pathog*. **4**: e1000124.

Pagán I, Fraile A, Fernández-Fueyo E, Montes N, Alonso-Blanco C, García-Arenal F.

2010. *Arabidopsis thaliana* as a model for the study of plant-virus co-evolution. *Phil.*

Trans. R. Soc. B **365**: 1983-1995.

Parrat SR, Numminen E, Laine AL. 2016. Infectious disease dynamics in heterogeneous

landscapes. *Annu. Rev. Ecol. Evol. Syst*. **47**: 283-306.

Pfenning K. 2001. Evolution of pathogen virulence: the role of variation in host

phenotype. *Proc. R. Soc. B* **268**: 755-760.

Pieterse CMJ, León-Reyes A, van der Ent S, van Wees SC. 2009. Networking by small-
molecule hormones in plant immunity. *Nat. Chem. Biol.* **5**: 308-316.

Pieterse CMJ, van der Does D, Zamioudis C, León-Reyes A, van Wees SCM. 2012.
Hormonal modulation of plant immunity. *Annu. Rev. Cell Dev. Biol.* **28**: 489-521.

Rozas J, Ferrer-Mata A, Sánchez-del Barrio JC, Guirao-Rico S, Librado P, Ramos-Osins
SE, Sánchez-Gracia A. 2017. DnaSP 6: DNA sequence polymorphism analysis of
large data sets. *Mol. Biol. Evol.* **34**: 3299-3302.

Schmid-Hempel P. 2011. *Evolutionary Parasitology*. Oxford University Press, Oxford:
UK, pp. 259-272.

Schmid-Hempel P, Koella JC. 1994. Variability and its implications for host-parasite
interactions. *Parasitol. Today* **10**: 98-102.

Simko I, Piepho HP. 2012. The area under the disease progress stair: calculation,
advantage, and application. *Phytopathology* **102**: 381-389.

Singh DP, Moore CA, Gilliland A, Carr JP. 2004. Activation of multiple antiviral defence
mechanisms by salicylic acid. *Mol. Plant Pathol.* **5**: 57-63.

Sofer L, Cabanillas D, Gayral M, Téplier R, Pouzoulet J, Ducousso M, Dufin L, Bréhélin
C, Ziegler-Graff V, Brault V, Revers Frédéric. 2017. Identification of host factors
potentially involved in *RTM*-mediated resistance during potyvirus long distance
movement. *Arch. Virol.* **162**: 1855-1865.

Soosaar JLM, Burch-Smith TM, Dinesh-Kumar SP. 2005. Mechanisms of plant
resistance to viruses. *Nat. Rev. Microbiol.* **3**: 789-799.

Tajima F. 1989. Statistical methods for testing the neutral mutation hypothesis by DNA
polymorphism. *Genetics* **123**: 585-595.

Thines B, Katsir L, Melotto M, Niu Y, Mandaokar A, Liu G, Nomura K, He SY, Howe

GA, Browse J. 2007. JAZ repressor proteins are targets of the SCF(COI1) complex

during jasmonate signalling. *Nature* **448**: 661-665.

Todesco M, Balasubramanian S, Hu TT, Traw MB, Horton M, Epple P, Kuhns C,

Sureshkumar S, Schwartz C, Lanz C, Laitinen RAE, Huang Y, Chory J, Lipka V,

Borevitz JO, Dangl JL, Bergelson J, Nordborg M, Weigel D. 2010. Natural allelic

variation underlying a major fitness trade-off in *Arabidopsis thaliana*. *Nature* **465**:

632-636.

Ton J, van Pelt JA, van Loon LC, Pieterse CMJ. 2002. Differential effectiveness of

salicylate-dependent and jasmonate/ethylene-dependent induced resistance in

Arabidopsis. *Mol. Plant Microbe Interact*. **15**: 27-34.

Torres-Barceló C, Martín S, Daròs JA, Elena SF. 2008. From hypo- to hypersuppression:

effect of amino acid substitutions on the RNA-silencing suppressor activity of the

tobacco etch potyvirus HC-Pro. *Genetics* **180**: 1039-1049.

Travisano M, Mongold, JA, Bennett, AF, Lenski RE. 1995. Experimental test of the roles

of adaptation, chance, and history in evolution. *Science* **267**: 87-90.

Traw MB, Kniskern JM, Bergelson J. 2007. SAR increases fitness of *Arabidopsis*

thaliana in the presence of natural bacterial pathogens. *Evolution* **61**: 2444-2449.

Truniger V, Aranda MA. 2009. Recessive resistance to plant viruses. *Adv. Virus Res*. **75**:

119-159.

Valverde S, Vidiella B, Montáñez R, Fraile A, Sacristán S, García-Arenal F. 2020.

Coexistence of nestedness and modularity in host-pathogen infection networks. *Nat.*

Ecol. Evol. **4**: 568-577.

Van de Weyer AL, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, Witek K, Jones JDG,

Dangl JL, Weigel D, Bemm F. 2019. A species-wide inventory of NLR genes and

alleles in *Arabidopsis thaliana*. *Cell* **178**: 1260-1272.

Van Wees SCM, de Swart EAM, van Pelt JA, van Loon, LC, Pieterse CMJ. 2000.

Enhancement of induced disease resistance by simultaneous activation of salicylate-

and jasmonate-dependent defense pathways in *Arabidopsis thaliana*. *Proc. Natl.*

Acad. Sci. USA **97**: 8711-8716.

Verchot J. 2012. Cellular chaperones and folding enzymes are vital contributors to

membrane bound replication and movement complexes during plant RNA virus

infection. *Front. Plant Sci*. **3**, 271.

Voinnet O. 2001. RNA silencing as a plant immune system against viruses. *Trends Genet*.

17: 449-459.

Weitz JS, Poisot T, Meyer JR, Flores CO, Valverde S, Sullivan MB, Hochberg ME. 2013.

Phage-bacteria infection networks. *Trends Microbiol*. **21**: 82-91.

White PS, Choi A, Pandey R, Menezes A, Penley M, Gibson AK, de Roode J, Morran L.

2020. Host heterogeneity mitigates virulence evolution. *Biol. Lett* **16**: 20190744.

Yang Z. 2006. *Computational Molecular Evolution*. Oxford University Press. Oxford:

UK, pp 262-264.

Yang Z, Huang Y, Yang J, Yao S, Zhao K, Wang D, Qin Q, Bian Z, Li Y, Lan Y, Zhou

T, Wang H, Liu C, Wang W, Qi Y, Xu Z, Li Y. 2020. Jasmonate signaling enhances

RNA silencing and antiviral defense in rice. *Cell Host Microbe* **28**: 89-103.

Zhou JM, Zhang Y. 2020. Plant immunity: danger perception and signaling. *Cell* **181**:

978-989.

1124

1125   **Data availability**

1126 Raw data will be available upon request.

1127

Acknowledgments

1129 We thank Francisca de la Iglesia and Paula Agudo for excellent technical assistance and

1130 the rest of the EvolSysVir lab members for fruitful discussions. This work was supported

1131 by grant PID2019-103998GB-I00 (Agencia Estatal de Investigación - FEDER) and

1132 PROMETEU2019/012 (Generalitat Valenciana) to S.F.E.

1133

Author contributions

1135 S.F.E. conceived the study, analyzed the data and wrote the paper. R.N., S.A., F.M.,

1136 B.W., and J.L.C. performed all the experiments.

1137

Competing interests

1139 The authors declare no competing interests.

1140

Figures and Tables

Fig. 1. Neighbor-joining clustering of the 21 mutant genotypes of Arabidopsis according to their phenotypic similarity in response to TuMV infection. Genotypes selected as hosts for the evolution experiments are highlighted in yellow.

Fig. 2.　Evolution of disease progression (*AUDPS*) along the serial passages of experimental evolution on each different host genotype.　Different symbols and lines represent the independent evolutionary lineages.　Panels are arranged from the most resistant genotype (*eds8-1*) to the most sensitive one (*jin1*) according to the groups defined in Fig. 1.

1144

Fig. 3. Evolution of infectivity (*I*) along the serial passages of experimental evolution on each different host genotype. Different symbols and lines represent the independent evolutionary lineages. Panels are arranged from the most resistant genotype (*eds8-1*) to the most sensitive one (*jin1*) according to the groups defined in Fig. 1.

1145

Fig. 4. Evolution of viral load (*VL*) along the serial passages of experimental evolution on each different host genotype. Different symbols and lines represent the independent evolutionary lineages. Panels are arranged from the most resistant genotype (*eds8-1*) to the most sensitive one (*jin1*) according to the groups defined in Fig. 1.

1146

Fig. 5. Test of the contribution of historical contingency to the observed pattern of adaptation of TuMV to the different Arabidopsis genotypes. (A) Evolved *vs* ancestral values for *AUDPS*, (B) for *I*, and (C) for *VL*. The diagonal lines represent the null hypothesis of historical differences being fully preserved despite adaptation. The solid lines represent the linear regression of the data (dashed lines represent the 95% CI for the regression lines).

Fig. 6. Estimated rates of evolution for *AUDPS* (A) and *I* (B) obtained from the fitting of an ARIMA(1,0,0) model (Eq. 2) to the data shown in Fig. 2 and Fig. 3, respectively. Different symbols represent different independent lineages evolved on the corresponding host genotype. Solid lines represent the linear regression of the data; dashed lines the 95% confidence intervals for the regression lines. In both cases, Arabidopsis genotypes are ranked from the less to the more permissive to TuMV infection according to Fig. 1.

Fig. 7. The rates of evolution for *AUDPS* and *I* differ among possible mechanisms of resistance to infection. Rates of evolution are expressed in log-scale to better visualize the slowest rates of *I* evolution. Data were fitted to the MANOVA model defined by Eq. 3. Error bars represent ±1 SEM.

1149

Fig. 8. Analysis of the full cross-infection matrix. (A) Packed matrix that highlights the its nested structure, compatible with a gene-for-gene infection model. Black squares represent cases in which the *AUDPS* values were equal or greater than those observed for the corresponding lineage in its local host. Last column shows the species-level specialization index *d'* (*d'* = 0 means most generalist pathogen, *d'* = 1 means most specialist pathogen). (B) Bipartite host genotype - viral lineage infection network. The size of the nodes is proportional to their degree. The phenotypic groups defined in Fig. 1 are indicated above the host genotype.

1150

Fig. 9. Phenotypic disease-related traits (*AUDPS*, *I* and *SS*) of TuMV VPg/D113G relative to TuMV wild-type VPg. (A) – (C) Evaluated in the wild-type *Arabidopsis* and the *cpr5-2* mutant genotype. (D) – (F) Evaluated in the wild-type *Arabidopsis* and the *jin1* mutant genotype. Error bars represent ±1 SEM.

1151

Table 1. Different Arabidopsis mutant genotypes used in this study. Highlighted in gray those used for the evolution experiments.

Genotype	Gene name	Affected pathway	Expected phenotype relative to wild-type plants	Reference
coi1-4	*CORONATINE INSENSITIVE 1* (AT2G39940)	Repression of JA-responsive genes	No ISR, no effect on virus	Thines *et al.* (2007)
cpr5-2	*CONSTITUTIVE EXPRESSOR OF PR GENES 5* (AT5G64930)	Membrane protein, negative regulator of pathogen-dependent SA signaling	More resistant, constitutive SAR	Love *et al.* (2007)
dbp2	*DNA-BINDING PROTEIN PHOSPHATASE 2*	Transcriptional regulation of gene expression in potyvirus-infected plants	More resistant to potyvirus infection	Castelló *et al.* (2011)
dcl2	*DICER-LIKE 2* (AT3G03300)	Partial loss of RNA-silencing	No effect, siRNAs produced by DCL4	Bouché *et al.* (2006)
dcl4	*DICER-LIKE 4* (AT5G20320)	Partial loss of RNA-silencing	No effect, siRNA produced by DCL2	Bouché *et al.* (2006)
dcl2 dcl4	Double mutant *dcl2 dcl4*	Complete loss of RNA-silencing	More susceptible, no siRNA production	Bouché *et al.* (2006)
dip2	*DBP-INTERACTING PROTEIN 2* (AT5G03210)	Transcriptional regulation of gene expression in potyvirus-infected plants	More susceptible	Castelló *et al.* (2011)
eds4-1	*ENHANCED DISEASE SUSCEPTIBILITY 4* (AT5G51200)	Loss of SA-dependent signaling	More susceptible, no SAR	Gupta *et al.* (2000)
eds5-1	*ENHANCED DISEASE SUSCEPTIBILITY 5* (AT4G39030)	Lipase-like protein, positive regulator of pathogen-dependent SA signaling	More susceptible, no SAR	Nawrath *et al.* (2002)
eds8-1	*ENHANCED DISEASE SUSCEPTIBILITY 8*	Reduced expression of plant defensin genes, reduced ISR	More resistant, enhanced SAR	Love *et al.* (2007)
ein2-1	*ETHYLENE INSENSITIVE 2* (AT5G03280)	MAPK, ET signaling intermediate, negative regulator SA-dependent signaling	More resistant, enhanced SAR	Love *et al.* (2007)
etr1-1	*ETHYLENE RESPONSE 1* (AT1G66340)	ET receptor, negative regulator SA-dependent signaling	More resistant, enhanced SAR	Love *et al.* (2007)

hsp90-1	*HEAT SHOCK PROTEIN 1* (AT5G52640)	Recessive r gene, required for membrane-bound replication complexes; protein folding	More resistant, missing component for viral replication	Verchot (2012)
i4g1	*EUKARYOTIC TRANSLATION INITIATION FACTOR (ISO) 4G 1* (AT3G60240)	Recessive *r* gene, initiation of viral RNA translation	More resistant, missing component for viral gene expression	Nicaise *et al.* (2007)
i4g2	*EUKARYOTIC TRANSLATION INITIATION FACTOR (ISO) 4G 2*	Recessive *r* gene, initiation of viral RNA translation	More resistant, missing component for viral gene expression	Nicaise *et al.* (2007)
jin1	*JASMONATE INSENSITIVE 1* (AT1G32640)	Loss of JA signaling; negative regulator of SA-dependent signaling	More resistant, enhanced SAR	Laurie-Berry *et al.* (2006)
npr1-1	*NONEXPRESSER OF PR GENES 1* (AT1G64280)	Ankyrin-repeat protein required for *PR-1* activation	More susceptible, no SAR, no ISR	Cao *et al.* (1994)
p58^{IPK}	*HOMOLOG OF MAMMALIAN P58^{IPK}* (AT5G03160)	Constitutive activation of PKR	More resistant, strong apoptosis-mediated HR	Bilgin *et al.* (2003)
pad4-1	*PHYTOALEXIN DEFICIENT 4* (AT3G52430)	Lipase-like protein, positive regulator pathogen-dependent SA signaling	More susceptible, no SAR	Cui *et al.* (2018)
sid2-1	*SA INDUCTION DEFICIENT 2* (AT1G74710)	Isochorismate synthase, required for SA bioshynthesis	More susceptible, no SAR	Nawrath and Métraux (1999)

ET - ethylene; HR – hypersensitive response; ISR - induced systemic resistance; JA - jasmonic acid; MAPK – mitogen-activated protein kinase; PKR - protein kinase RNA-activated; SA - salicylic acid; SAR - systemic acquired resistance

1152

Table 2. Results of the multivariate analysis of variance (MANCOVA) for the three phenotypic traits evaluated along the course of experimental evolution. The different factors are defined in Eq. 1.

Source of variation	Wilk's Λ	F	df_1, df_2	P	η_P^2	$1 - \beta$
μ	0.004	3377.616	3, 42	< 0.001	0.996	1.000
t	0.265	38.799	3, 42	< 0.001	0.735	1.000
G	0.093	6.467	24, 122.414	< 0.001	0.547	1.000
$t{\times}G$	0.155	4.608	24, 122.414	< 0.001	0.463	1.000
$L(G)$	0.529	0.278	108, 126.692	1.000	0.191	0.373
$t{\times}L(G)$	0.466	0.351	105, 126.677	1.000	0.225	0.487

η_P^2: magnitude of the effect; $1 - \beta$: power of the test.

1153

1154

Table 3. Results of the probit regression testing for the specificity of adaptation. The different model factors are defined in Eq. 4.

Source of variation	LRT	df	P	η_P^2	$1 - \beta$
ϕ	21224.493	1	< 0.001	0.959	1.000
t	25995.056	1	< 0.001	0.815	1.000
TG	295.160	8	< 0.001	0.569	1.000
$t{\times}TG$	568.212	8	< 0.001	0.010	0.981
LG	20.545	8	0.008	0.288	0.661
$t{\times}LG$	30.200	8	< 0.001	0.005	0.804
$L(LG)$	207.257	35	< 0.001	0.385	1.000
$t{\times}L(LG)$	235.073	35	< 0.001	0.016	0.984
$TG{\times}LG$	258.888	64	< 0.001	0.314	1.000
$t{\times}TG{\times}LG$	365.512	64	< 0.001	0.018	0.961
$TG{\times}L(LG)$	788.661	280	< 0.001	0.048	0.999
$t{\times}TG{\times}L(LG)$	1033.649	280	< 0.001	0.042	0.993

LRT: likelihood ratio test distributed as a χ^2; η_P^2: magnitude of the effect; $1 - \beta$: power of the test.

1155

Additional Information

Fig. S1. Disease-related traits evaluated for the 21 Arabidopsis genotypes listed in Table 1. (A) Change in dry weight (ΔDW; genetic component of variance $\sigma_G^2 = 93.09\%$, one-way ANOVA $F_{20,84} = 71.820$, $P < 0.001$). (B) Severity of symptoms (SS; $\sigma_G^2 = 75.88\%$, $F_{20,84} = 17.643$, $P < 0.001$). (C) Area under the disease progress stairs ($AUDPS$; $\sigma_G^2 = 99.96\%$, $F_{20,399} = 2626.754$, $P < 0.001$). (D) Infectivity (I; $\sigma_G^2 = 78.91\%$, $F_{20,399} = 4.563$, $P < 0.001$). (E) Viral load (VL; $\sigma_G^2 = 78.99\%$, $F_{20,84} = 22.113$, $P < 0.001$). Error bars represent ±1 SEM, except for the *AUDPS*, which are 95% confidence intervals based in 1000 bootstrap pseudo-replicates.

1159

Fig. S2. Daily variation of (A) infectivity (*I*), (B) symptoms severity (*SS*) and (C) viral load (*VL*) for genotypes representative of groups G1 (*eds8-1* more resistant than wild-type to TuMV infection) and G4 (*cpr5-2*, *dcl2 dcl4*, *hsp90-1* similar to wild-type in response to infection) in Fig. 1. Error bars represent ±1 SEM.

1160

Fig. S3. Distribution of observed mutations per TuMV cistron. (A) Schematic representation of the distribution of synonymous (green boxes) and nonsynonymous (red boxes) distribution along TuMV genome. (B) Frequency of mutations observed on each cistron. Error bars represent ±1 SEM.

1161

Table S1. Nucleotide and amino acid changes found in the consensus sequences of the TuMV evolved viruses. SNP positions referred to the complete genome. Amino acid changes positions are referred to the mature peptide.

ORF	Lineages	Mutation	Amino acid change	Amino acid group change	Syn:Nosyn	Ts:Tv
P1	$p58^{IPK}$/L4	A233G	synonymous			
	dcl2 dcl4/L1	U289C	V45A	conservative nonpolar by nonpolar		
	eds8-1/L3	G298U	G75V	conservative nonpolar by nonpolar		
	$p58^{IPK}$/L1	A490G	E121G	negatively charged by nonpolar	3:5	7:1
	npr1-1/L3	C633U	P169S	nonpolar by nonpolar, strong structural effect		
	eds8-1/L3	U692C	synonymous			
	i4g2/L3	C806U	synonymous			
	hsp90-1/L4	G1087A	G320E	nonpolar by negatively charged		
HC-Pro	dbp2/L4	G1273A	S20N	small polar by larger polar		
	cpr5-2/L4	A1430G	synonymous			
	dbp2/L4	G1512A	D100N	negatively charged by polar		
	dcl2 dcl4/L1	U1520C	synonymous		5:4	9:0
	jin1/L4	C1760U	synonymous			
	jin1/L5					
	npr1-1/L3	C1916U	synonymous			

	Mutant/Line	Nucleotide	AA change	Description		
	i4g2/L4	G2316A	V367I	conservative small nonpolar by longer nonpolar		
	dbp2/L2	A2493G	T326S	small polar by nonpolar		
P3	*dbp2*/L3	C3205U	A206V	conservative nonpolar by nonpolar		
	jin1/L4 *jin1*/L5	U3269C	synonymous		2:3	3:1
	cpr5-2/L5	C3541U	T318M	small polar by long nonpolar		
	cpr5-2/L5	A3564U	T326S	conservative polar by polar		
6K1	cpr5-2/L5	U3686C	synonymous		2:0	2:0
	p58^{IPK}/L5	U3743C	synonymous			
	dcl2 dcl4/L5	G4032A	A72T	small nonpolar by polar		
	dcl2 dcl4/L5	G4148A	synonymous			
	cpr5-2/L4 *cpr5-2*/L5	G3269A	V148I	conservative small nonpolar by larger nonpolar		
CI	*npr1-1*/L1	A4357G	N186S	positively charged by small polar	6:13	15:4
	npr1-1/L3	U4409G	I200M	conservative nonpolar by nonpolar		
	i4g2/L5	G4448A	synonymous			
	jin1/L1	C4559U	synonymous			
	jin1/L2	A4674G	I291V	conservative nonpolar by nonpolar		
	dcl2 dcl4/L1	A4941G	I378V	conservative nonpolar by nonpolar		

	jin1/L1	C4945A	P379Q	nonpolar by polar, strong structural effect		
	cpr5-2/L4	G5183A	synonymous			
	cpr5-2/L5					
	i4g2/L3	G5193U	A416S	nonpolar by polar		
	npr1-1/L1	G5213A	synonymous			
	i4g2/L1	U5372C	synonymous			
	i4g2/L4					
	i4g2/L1	C5193G	A532G	conservative nonpolar by nonpolar		
	p58^{IPK}/L2	U5519C	synonymous			
6K2	*jin1*/L3	A5900G	synonymous		1:0	1:0
VPg	*eds8-1*/L4	C5977U	H33Y	positively charged by polar		
	eds8-1/L5					
	hsp90-1/L1	C6219U	L107F	conservative nonpolar by nonpolar		
	dcl2 dcl4/L4				0:47	45:2
	dbp2/L1					
	dbp2/L5	G6237A	D113N	negatively charged by polar		
	i4g2/L2					
	i4g2/L5					

dcl2 dcl4/L2			
dcl2 dcl4/L3			
eds8-1/L4			
p58[IPK]/L4			
npr1-1/L2			
eds8-1/L2	G6237C	D113H	negatively charged by positively charge
cpr5-2/L3			
cpr5-2/L4			
cpr5-2/L5	A6238G	D113G	negatively charged by nonpolar
npr1-1/L1			
npr1-1/L5			
hsp90-1/L2			
npr1-1/L4	U6239A	D113E	conservative negatively charged by negatively charged
i4g2/L3	A6243C	N115H	polar by positively charged
eds8-1/L3			
dbp2/L1			
dcl2 dcl4/L5	A6243G	N115E	polar by negatively charged
p58[IPK]/L1			
dbp2/L2	A6244G	N115S	conservative polar by polar
dbp2/L3			

dbp2/L4			
jin1/L3			
eds8-1/L1			
hsp90-1/L5	A6247G	E116G	negatively charged by small nonpolar
p58^{IPK}/L2			
i4g2/L1			
i4g2/L4	U6252C	R118C	positively charged by nonpolar
dcl2 dcl4/L1			
p58^{IPK}/L3			
jin1/L1			
jin1/L2	G6253A	R118H	conservative positively charged by positively charged
jin1/L4			
jin1/L5			
hsp90-1/L3			
hsp90-1/L4			
cpr5-2/L1	A6261G	K121E	positively charged by negatively charged
npr1-1/L3			
eds8-1/L4			
p58^{IPK}/L5			

NIa-Pro	*npr1-1*/L4	G6950A	synonymous			
	dcl2 dcl4/L3	A7002U	I176F	nonpolar by nonpolar	2:2	3:1
	dcl2 dcl4/L3	G7029A	D187N	negatively charged by polar		
	jin1/L3	C7181U	synonymous			
NIb	*i4g2*/L5	A7394G	synonymous			
	eds8-1/L3	C8271U	synonymous			
	eds8-1/L5					
	npr1-1/L5	U8302C	L366P	nonpolar by nonpolar, strong structural effect	5:3	7:1
	npr1-1/L5	C8308U*	T368I	polar by nonpolar		
	dbp2/L3	C8337G	L378V	conservative nonpolar by nonpolar		
	i4g2/L1	U8345C	synonymous			
	dbp2/L3	G8360A	synonymous			
CP	*hsp90-1*/L2	G8792A	synonymous			
	jin1/L3					
	jin1/L4	G8965A	S70N	conservative polar by polar		
	jin1/L5				1:10	1:10
	dcl2 dcl4/L1					
	npr1-1/L4	A9091G	D112G	negatively charged by small nonpolar		
	i4g2/L5	A9091C	D112A	negatively charged by small nonpolar		

				1162
cpr5-2/L4	G9224A	M156I	conservative nonpolar by nonpolar	
dcl2 dcl4/L2	A9244G	E163G		
dbp2/L5	A9252G	I166V	negatively charged by small nonpolar	
hsp90-1/L1	C9541U	A262V	conservative nonpolar by nonpolar	

[*]Polymorphic positions; Syn:Nosyn: number of synonymous *vs* nonsynonymous mutations; Ts:Tv: number of transitions *vs* transversions. The two highlighted nonsynonymous mutations D113G and R118H in VPg have been used in further experiments.