1

Isoform-specific roles of the *Drosophila* filamin-type protein Jitterbug (Jbug) during development

SeYeon Chung[1, ¶, *], Thao Phuong Le[1, ¶], Vishakha Vishwakarma[1], Yim Ling Cheng[2] and Deborah J. Andrew[2]

[1] Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA

[2] Department of Cell Biology, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

[¶] These authors contributed equally to this work.

[*]Author for correspondence: seyeonchung@lsu.edu

Running title: Roles of Jitterbug in *Drosophila* development

Key workds: *Drosophila*, filamin, *jitterbug*, actin, thoracic bristles, epidermal denticles, tubular epithelial organs, trans-splicing, Crb

ABSTRACT

Filamins are highly conserved actin-crosslinking proteins that regulate organization of the actin cytoskeleton. As key components of versatile signaling scaffolds, filamins are implicated in developmental anomalies and cancer. Multiple isoforms of filamins exist, raising the possibility of distinct functions for each isoform during development and in disease. Here, we provide an initial characterization of *jitterbug* (*jbug*), which encodes one of the two filamin-type proteins in *Drosophila*. We generate Jbug antiserum that recognizes all of the spliced forms and reveals differential expression of different Jbug isoforms during development, and a significant maternal contribution of Jbug protein. To reveal the function of Jbug isoforms, we create new genetic tools, including a null allele that deletes all isoforms, hypomorphic alleles that affect only a subset, and UAS lines for Gal4-driven expression of the major isoforms. Using these tools, we demonstrate that Jbug is required for viability and that specific isoforms are required in the formation of actin-rich protrusions including thoracic bristles in adults and ventral denticles in the embryo. We also show that specific isoforms of Jbug show differential localization within epithelia and that maternal and zygotic loss of *jbug* disrupts Crumbs (Crb) localization in several epithelial cell types.

Introduction

Actin-crosslinking proteins organize actin filaments into higher-order structures such as orthogonal actin arrays and parallel actin bundles. Filamins are highly conserved large cytoplasmic proteins that crosslink actin filaments into dynamic three-dimensional structures (Nakamura *et al.* 2011; Razinia *et al.* 2012). Filamins consist of actin-binding Calponin-homology (CH) domains at their N-terminus, followed by a long C-terminal rod-like domain of filamin-type immunoglobulin (IG-FLMN) repeats. Dimerization of filamins through the last C-terminal IG-FLMN repeat allows the formation of a flexible V-shaped structure that holds two actin filaments at large angles to create either a loose three-dimensional actin network (Nakamura *et al.* 2011; Razinia *et al.* 2012) or parallel bundles of actin filaments (Sokol and Cooley 1999; Gay *et al.* 2011). In addition to binding to actin, filamins interact with transmembrane receptors, adhesion molecules and even transcription factors, and are thus involved in multiple cell functions, including motility, maintenance of cell shape, and differentiation (Nakamura *et al.* 2011; Razinia *et al.* 2012). Mutations in filamins are associated with a wide range of congenital anomalies and have been shown to both promote and inhibit metastasis and cancer growth (Razinia *et al.* 2012; Savoy and Ghosh 2013; Shao *et al.* 2016; Sasaki *et al.* 2019), emphasizing the multiple critical roles of filamins in development and disease.

Studies in *Drosophila* have helped refine our understanding of the roles of filamins in vivo. *Drosophila* has two orthologs of human filamin A: Cheerio (Cher) and Jitterbug (Jbug). Cher shares the organization of the protein and 46% identity and 61% similarity in amino acid sequence with human filamin A. Jbug, with 23% identity and 36% similarity in amino acid sequence with human filamin A, has some distinct features, including an actin-binding domain consisting of three, instead of two, CH domains. Several studies have revealed roles for Cher in organizing the F-actin cytoskeleton in multiple developmental contexts and diseases, including formation of ovarian germline ring canals (Li *et al.* 1999; Sokol and Cooley 1999; Huelsmann *et al.* 2016), follicle cell migration (Sokol and Cooley 2003), tissue morphogenesis during cellularization (Krueger *et al.* 2019), and *Drosophila* tumorigenesis (Külshammer and Uhlirova 2013). Compared to the extensive studies on Cher, our understanding of the roles of Jbug is more limited. Originally identified as a gene that causes bang-sensitive seizure – the phenotype that gives the gene its *jitterbug* name (Song and Tanouye 2006), *jbug* is required in photoreceptor cells for axon targeting (Oliva *et al.* 2015) and in tendon cells to maintain their shape at the muscle-tendon junction (Olguín *et al.* 2011; Manieu *et al.* 2018).

Multiple isoforms of filamin proteins exist in both *Drosophila* and mammals (Sokol and Cooley
1999; Browne *et al.* 2000; Van der Flier and Sonnenberg 2001; Gorlin *et al.* 1990; Wang *et al.*
2007). Originally, only two Cher isoforms were reported: a large isoform (FLN240) that contains
both actin-binding CH domains and IG-FLMN repeats and a smaller one (FLN90) that contains
only IG-FLMN repeats (Sokol and Cooley 1999). Recent annotations indicate that *cher* encodes
six longer isoforms (>240 kDa), which include two actin-binding CH domains and 18-22 IG-FLMN
repeats, and four shorter isoforms (90-100 kDa), which include only eight IG-FLMN repeats
without any CH domains (Flybase; www.flybase.org; Thurmond *et al.* 2019). Importantly, a recent
study in *Drosophila* larval neuromuscular junctions revealed a role for the shorter Cher isoform
(FLN90) as a postsynaptic scaffold in *Drosophila* larval neuromuscular junctions during synapse
formation (Lee and Schwarz 2016), suggesting distinct roles for different isoforms. Like *cher*, *jbug*
also encodes multiple isoforms with a range of protein sizes. However, the roles for *jbug* have
been revealed mostly using RNA interference (RNAi) constructs that target most of the existing
splice forms (Olguín *et al.* 2011; Manieu *et al.* 2018), preventing the characterization of isoform-
specific roles for Jbug.

Here, we report on the creation of several new genetic tools for parsing out the roles of the
different Jbug isoforms, including a null allele that deletes all isoforms and hypomorphic alleles
that affect only a few. We also create UAS constructs for expressing the major Jbug isoforms. We
demonstrate that different isoforms are differentially expressed during development and localize
to different domains in embryonic epithelia. We show isoform-specific roles of Jbug in formation
of epidermal denticles during embryogenesis and thoracic bristles in adults. Finally, we also report
that removal of both maternal and tissue-specific supplies of *jbug* disrupts tissue morphology and
reduces Crb accumulation in the subapical domain of epithelial cells.
94

Materials and Methods

Fly stocks and husbandry

Fly lines used in our experiments were: *Oregon R*; *jbug[20]*, *jbug[30]*, *jbug[133]*, UAS-Jbug-RC, UAS-

Jbug-RI, UAS-Jbug-RF (this work); UAS-Jbug-RL (Olguín *et al.* 2011); *fkh-Gal4* (Henderson and

Andrew 2000); *69B-Gal4* (Rorth 1996); *da-Gal4* (Wodarz *et al.* 1995); *matα-Gal4* (Häcker and

Perrimon 1998), *Df(2R)59AB*, *Df(2R)Exel6079*, *TRiP.JF01166* (Bloomington Stock Center);

GD8664, *GD13033* (Vienna *Drosophila* RNAi Center). All crosses were performed at 25°C.

Generation of *jbug* mutant alleles

jbug mutants were generated by homologous recombination (Gong and Golic 2003). For *jbug[20]*

null mutants, genomic fragments upstream and downstream of the *jbug* ORF were amplified by

PCR using the following primers (restriction sites in bold and linkers in italic):

in-fusion_L-22737893, 5'-*CTAGTCTAGGGCGCGCC*GATGAGTTGTGGCTTGAGCA-3'

in-fusion_R-22742442, 5'-*TAGGGGATCACGTACG*GCCAAACCAATGCTGAAGAT-3'

NotI_L-22717574, 5'-AT**GCGGCCGC**TAGAGGGGAGAGCAAGTGGA-3'

Acc65I_R-22721186, 5'-ATA**GGTACC**GTGTGAGCTTCGGGATCAAT-3'

For *jbug[30]* and *jbug[133]* hypomorphic alleles, genomic fragments upstream and downstream of an

exon common to all *jbug* splice forms were amplified using the following primers (restriction sites

in bold):

AscI_L-22727168, 5'-AT**GGCGCGCC**AAAGCCGTTGAAGAAAGCAA-3'

BsiWI_R-22731840, 5'-ATA**CGTACG**CTCGATGCACTTTGTCTCCA-3'

NotI_L-22717574, 5'-AT**GCGGCCGC**TAGAGGGGAGAGCAAGTGGA-3'

Acc65I_R-22721186, 5'-ATA**GGTACC**GTGTGAGCTTCGGGATCAAT-3'

PCR fragments were cloned into pW25, which carries *white[+]*, the recognition site for I-SceI

endonuclease, and FRT sites. The constructs were injected into *w[1118]* embryos by Rainbow

Transgenic Flies, Inc. Transformants were crossed to flies carrying hs-I-SceI and hs-Flp, and

progeny were heat shocked (37°C) for 1 hour 48–72 hours after egg laying (AEL). Recombination

and insertion were confirmed by genomic PCR and reverse-transcriptase PCR (RT-PCR) followed

by sequencing.

The following primers were used for RT-PCR for *jbug*[20] shown in Supplemental Figure 2

(restriction sites in bold and linkers in italic):

jbug 5-1, 5'-GGAGAACAAGTACCGGGTGA-3'

jbug 5-2, 5'-*GGATCCTCGAGCATATG*TCCTTCCACGTTCACCTCGA-3' (same as Jbug-Ab-3' for

making Jbug antibody)

jbug 3-1, 5'-AT**GCGGCCGC**TAGAGGGGAGAGCAAGTGGA-3' (same as NotI_L-22717574 used

for amplifying a homology arm downstream of the *jbug* ORF)

jbug 3-2, 5'-*CGCGCGGCAGCCATATG*ATGTCCTCACCCGGCCTAAC-3' (same as Jbug-Ab-5'

for making Jbug antibody)

Generation of UAS-Jbug-RC, -RI and -RF transgenic flies

An ORF for *jbug-RC* (1230 bp) was PCR-amplified using the existing cDNA RE40504 as a

template. A full length cDNA for *jbug-RF* (9430 bp) was created by In-Fusion cloning of three

fragments. The 5' fragment was prepared by enzyme digestion of RE40504 (cut with ApaI and

BamHI). The middle fragment (1189 bp) was amplified by reverse transcriptase-PCR (RT-PCR),

and the ApaI and BglII sites were added to the 5' and the 3' end of the fragment, respectively.

The 3' fragment was amplified by PCR using the existing cDNA GH03118 from the BglII site to

the end of the ORF, and the BamHI site was added to the 3' end. The ligated full cDNA was

confirmed by sequencing. A full cDNA for *jbug-RI* (5989 bp) was created by PCR amplification

using the *jbug-RF* cDNA as a template and confirmed by sequencing. *jbug-RC*, *jbug-RI* and *jbug-

RF* ORFs were PCR-amplified and cloned into the pUAST vector (Brand and Perrimon, 1993) to

create UAS lines. The DNA constructs were injected into w^{1118} embryos by Rainbow Transgenics,

Inc. (UAS-Jbug-RC) and GenetiVision (UAS-Jbug-RI and UAS-Jbug-RF).

Generation of Jbug antibody

A PCR fragment of 1212 bp common to all *jbug* splice forms was cloned into the pET15b vector.

The fragment corresponds to almost the entire open reading frame for the Jbug-PC isoform

except for the last six amino acids and contains four IG-FLMN repeats. Primers used for In-Fusion

cloning were (linkers in italic):

Jbug-Ab-5', 5'- *CGCGCGGCAGCCATATG*ATGTCCTCACCCGGCCTAAC-3'

Jbug-Ab-3', 5'- *GGATCCTCGAGCATATG*TCCTTCCACGTTCACCTCGA-3'

The construct was transformed into BL21-DE3 cells for protein induction with IPTG. Recombinant
protein was purified using Ni-NTA agarose (Qiagen). Rabbit polyclonal antibodies were generated
by Covance, Inc. and used at a dilution of 1:500 for immunohistochemistry and 1:5,000 for western
blot.
160

Whole-mount in situ hybridization
In situ hybridization was performed as described previously (Lehmann and Tautz 1994). Probes
specific for a subset of *jbug* splice forms are shown in Figure 1A. DNA fragments for probes 1
and 3 were PCR-amplified using wild-type genomic DNA, cloned into the pCR2.1 vector and
sequenced to verify. Probe 2 was made from cDNA GH03013, linearized at the KpnI site to create
a probe specific to all jbug splice forms. Antisense digoxigenin-labeled RNA probes were
generated using T7 (probes 1 and 3) or SP6 (probe 2) polymerase on linearized templates. The
molecular regions corresponding to the RNA probes are as follows:

Probe 1 (specific for *jbug-RF/M/N* and *jbug-RL*): 22734199 – 22731666
Probe 2 (detects all *jbug* transcripts): 22722586-22722906, 22723936-22724162, 22724531-
22725474
Probe 3 (specific for *jbug-RF/M/N* and *jbug-RI*): 22721775 – 22720075
173

Western blot
Protein extracts from embryos 0-2 and 2-16 hours AEL, 1st, 2nd, 3rd instar larvae, pupae (2 days
after puparium formation (APF), and 2-day old male and female adults were used. For embryos
homozygous for *jbug^{20}*, *Df(2R)59AB* and *Df(2R)Exel6079* and for 1st instar larvae homozygous
for *jbug^{20}* and transheterozygous for *jbug^{30}*/*Df(2R)Exel6079*, a balancer chromosome that
contains the twi-Gal4 UAS-GFP construct was used to distinguish desired genotypes from
heterozygous siblings. Rabbit α-Jbug (this work; 1:5,000), mouse α-Actin (RRID:AB_11004139;
Invitrogen;   1:2,000),   and   HRP-labelled   secondary   antibodies   (RRID:AB_430833;
RRID:AB_430834; Promega; 1:2,000) were used. Protein bands were detected using Pierce ECL
Western Blotting Substrate.
184

185 **Scanning electron microscopy**
186 Wild type (*Oregon R*) and *jbug^{30}* homozygous male flies were coated with gold palladium and
187 examined and photographed in a LEO/Zeiss Field-emission SEM.
188

Antibody staining and confocal microscopy

Embryos were collected on grape juice-agar plates and processed for immunofluorescence using

standard procedures. Briefly, embryos were dechorionated in 50% bleach, fixed in 1:1

heptane:formaldehyde for 40 min and devitellinized with 80% EtOH, then stained with primary

and secondary antibodies in PBSTB (1X PBS, 0.1% Triton X-100, 0.2% BSA). For Jbug and

phalloidin staining, embryos were hand-devitellinized. Antibodies used include: mouse α-Crb

(RRID:AB_528181; DSHB (Cq4); 1:10), mouse α-α-Spec (RRID:AB_528473; DSHB (3A9); 1:2) ,

rabbit α-Neurexin IV (Baumgartner *et al.* 1996; 1:2,000), rabbit α-CrebA (RRID:AB_10805295;

1:5,000), guinea pig α-Sage (Fox *et al.* 2013; 1:100), chicken α-GFP (RRID:AB_2534023;

Invitrogen; 1:500), rat α-E-Cad (RRID:AB_528120; DSHB (DCAD2); 1:50), rabbit α-Jbug (this

work; 1:500), rabbit α-β-Gal (RRID:AB_221539; Invitrogen; 1:500). Alexa fluor 488, 568, 647-

labelled secondary antibodies were used at 1:500 (RRID:AB_142924; RRID:AB_141778;

RRID:AB_2535766; RRID:AB_143157; RRID:AB_2535812; Molecular Probes). Alexa fluor-568-

labeled phalloidin was used for F-actin labelling (RRID:AB_2632953; Molecular Probes; 1:250).

All images were taken with a Leica SP8 confocal microscope using either a 63x, 1.4 NA or a 40x,

1.3 NA oil objective and the LAS X software (Leica).

Quantification of denticle precursors in the ventral epidermis

Embryos stained for phalloidin and E-Cad were imaged with a Leica SP8 confocal microscope. A

maximum intensity projection was generated using two focal planes (0.3 μm apart) in the region

of adherens junctions. Denticle belts in abdominal segments 4-6 in stage 16 embryos were

selected for quantification. Six columns of cells from anterior to posterior of each belt were

analyzed. In each column, denticle precursors of four cells closest to the ventral midline were

counted. For each genotype, five to eight embryos were quantified. Unpaired Student's t-test with

Welch's correction was used for statistical analysis.

Data availability

All data and methods required to confirm the conclusions of this work are within the article, figures,

and supplemental materials.

Results

Different splice forms of *jbug* are differentially expressed during development

Ten annotated splice forms of *jbug*, which encode six different protein isoforms, are reported in Flybase (www.flybase.org; Thurmond *et al.* 2019). All isoforms contain multiple repeats of filamin-type immunoglobulin (IG-FMLN) domains (Figure 1, A and B), which form a rod-like structure in filamin. Two Jbug isoforms consist exclusively of these repeats; Jbug-PC/G/H/J/K (each splice form encodes the same 42.4 kDa protein; hereafter referred to as Jbug-PC) has four IG-FMLN repeats and Jbug-PI (185.7 kDa) has sixteen (Figure 1B). On the other hand, Jbug-PL (175.6 kDa) and Jbug-PF/M/N, the three longest Jbug isoforms (>300 kDa), contain three tandem calponin-homology (CH) actin-binding domains in their N-termini as well as multiple IG-FMNLs in their C-termini (seven and nineteen IG-FMLNs for Jbug-PL and Jbug-PF/M/N, respectively) (Figure 1B).

High-throughput expression data available in Flybase (www.flybase.org; Thurmond *et al.* 2019) showed high expression of *jbug* splice forms throughout all developmental stages, except during 0-10 hours of embryogenesis when there is a minimal expression of *jbug-RF/M/N* and *jbug-RL* (Figure 1C). BDGP Expression Data showed that, in addition to the strong ectodermal expression at earlier stages, *jbug* is also upregulated in epithelial tubular organs, including the trachea primordia and the salivary gland (https://insitu.fruitfly.org/cgi-bin/ex/insitu.pl). Our previous work confirmed this expression and revealed that *jbug* tracheal expression requires Trachealess, a major transcription factor in the trachea (Chung *et al.* 2011).

in situ hybridization using probes specific for different *jbug* splice forms further revealed tissue-specific expression of specific *jbug* splice forms (Supplemental Figure 1). A 5' probe (Figure 1A; probe1) that should detect *jbug-RL* (the form that includes the CH actin-binding domains) and the longest splice forms *jbug-RF/M/N* revealed no expression until embryonic stage 12, when expression in the clypeolabrum (CL) was first detected at low levels (Supplemental Figure 1, left column). This CL expression persisted through embryogenesis with later additional expression in another group of anterior cells that, together with the CL-positive cells, forms the mouth and low level expression in posterior cells that form the posterior spiracles. A 3' probe (Figure 1A; probe 3) that detects *jbug-RI* (the form that includes sixteen IG-FLMN repeats without the CH actin-binding domains) and *jbug-RF/M/N* reveals broad ectodermal expression in early embryos, with tissue-specific expression apparent by stage 11 in the CL, salivary gland, and tracheal pits (tp), with some low level expression in the foregut and subset of midline cells (Supplemental Figure 1; right column). By stage 12, the CL and tp staining are no longer detectable and expression is now also observed in epidermal stripes. By late stages, expression is also observed in the region of

the posterior spiracles. A probe that detects all *jbug* splice forms (Figure 1A, probe 2) revealed a

pattern that was the sum of the expression patterns observed with the more 5' and 3' isoform-

specific probes. Given the very limited overlap in staining patterns with the 5' (probe 1) and 3'

(probe 3) probes, this finding suggests that *jbug-RL* and *jbug-RI* are the major mRNA splice forms

made in embryos and that *jbug-RL* is much more restricted in its expression than *jbug-RI*, both in

terms of how early the transcripts are detected and how broadly expression is observed. If *jbug*

RF/M/N transcripts are produced in embryos, the expression must be limited to the late posterior

spiracles, the only site where all three probes detected signals. Likewise, if *jbug-RC* is made in

embryos, it must completely overlap the expression patterns of *jbug-RL* and *jbug-RI*.

To analyze protein expression levels of Jbug during *Drosophila* development, we generated Jbug

antiserum that recognizes a protein domain encoded by all ten *jbug* splice forms (Figure 1A). This

domain includes the four IG-FLMN repeats present in all Jbug isoforms (Figure 1B). Our Western

blot analysis using wild-type protein extracts revealed that Jbug proteins are detected at high

levels throughout all developmental stages (Figure 1D). Strong Jbug signals in protein extracts

from 0-2 hour embryos (Figure 1D) also suggest a maternally provided pool of Jbug. The Jbug

antiserum detected bands in size ranges predicted from the major splice forms (42.4 kDa, 175.6

kDa, 185.7 kDa, >300 kDa). Importantly, the different isoforms were differentially expressed

during development (Figure 1D). Jbug-PC (42.4 KDa) was abundant only in pupae and Jbug-PL

(175.6 kDa) was most abundant in adults (Figure 1D). Bands corresponding in size to Jbug-PI

(185.7 kDa) and Jbug-PF/M/N (>300 kDa) were detected at all stages with slight differences in

protein levels. The PI band had the highest intensity at almost all stages, with the exception of

PC in pupae, where the levels of PI and PC were similar (Figure 1D). Jbug-PF/M/N isoforms differ

only slightly from one another in regions outside the CH and IG-FLMN domains. Due to their huge

sizes and a very slight difference in molecular weight among Jbug-PF/M/N, we were not able to

distinguish exactly which of the three isoforms is expressed at any specific stage. The large size

of these isoforms potentially also diminishes membrane transfer, potentially leading to relatively

weak signals. Alternatively, the relatively weak signals could indicate that only very low levels of

the longest splice forms are made. Notably, several intermediate-sized bands that did not match

the predicted size of annotated isoforms were also detected at different levels throughout

development (Figure 1D). The existence of these bands suggests either that more Jbug isoforms

exist than have been annotated or that proteolyzed and/or phosphorylated fragments are being

detected (also see Figure 1E; most of these extra bands are absent in *jbug* null mutant larval

extracts). Indeed, roles for proteolytic forms of Filamin A (Bedolla *et al.* 2009; Savoy and Ghosh

2013; Yue *et al.* 2013) as well as phosphorylated Filamin A (Ohta and Hartwig 1996; Tirupula *et*

al. 2015) have been reported from studies of the mammalian proteins. Overall, our data indicate

that different splice forms of *jbug* are differentially expressed during *Drosophila* development.

288   Null and hypomorphic *jbug* alleles have been generated

To investigate roles for *jbug* in development, we generated mutant alleles for *jbug* using

homologous recombination (Gong and Golic 2003). Two donor DNA constructs were used to

target different genomic regions. One construct targets the genomic region encompassing the

translation start sites for all splice forms and extends through the exons shared by all splice forms;

the other targets only the region containing the exons shared by all splice forms. With the

construct that targets the larger region, we obtained a null mutant, *jbug^{20}* (Figure 1A). PCR

analysis confirmed that the entire targeted region was replaced with the mini-*white^{+}* (mini-*w^{+}*)

gene (data not shown). Reverse transcriptase-PCR (RT-PCR) using mRNA extracted from

homozygous *jbug^{20}* embryos further confirmed that no *jbug* transcripts are produced in *jbug^{20}*

embryos (Supplemental Figure 2, A-C). Both *jbug^{20}* homozygous animals and transheterozygotes

for *jbug^{20}* over either of the two deficiency lines that delete all or part of the *jbug* locus (*59AB* and

Exel6079; Figure 1A) died during the first instar larval stage (Figure 1H; Table 1). Western blot

analysis using Jbug antiserum showed that whereas a significant amount of the Jbug proteins

were detected in *jbug^{20}* embryos (Figure 1, E and H), almost all protein bands were absent by the

first instar larval stage, with the exception of two low molecular weight bands that are likely to be

from nonspecific cross-reactivity (Figure 1E). Similarly, the same Jbug isoforms were detected

both in *59AB* and *Exel6079* homozygous embryos as observed in WT (Figure 1F). These data

suggest that Jbug proteins from maternally provided *jbug* mRNA or protein persist through

embryogenesis. Since only very low level *jbug* mRNA is detected in 0-2 hour wild-type embryos,

based on RNA-Seq (Figure 1C, Flybase; www.flybase.org) and in situ hybridization (Fly-FISH;

http://fly-fish.ccbr.utoronto.ca; BDGP; https://insitu.fruitfly.org/cgi-bin/ex/insitu.pl; this study), it is

the Jbug protein and not *jbug* mRNA, that is maternally provided.

With the DNA construct that aimed to remove the genomic region common for all splice forms of

jbug, we unexpectedly obtained two hypomorphic alleles, *jbug^{30}* and *jbug^{133}*, that affect only a

subset of splice forms. Incomplete recombination occurred in these two alleles, resulting in either

a part of (for *jbug^{30}*) or the entire DNA construct (for *jbug^{133}*) being inserted into the *jbug* locus

(Figure 1, A' and A''). In *jbug^{30}*, the more 3' homology arm in the donor DNA recombined correctly,

but the other homology arm was inserted into the genome along with the mini-*w+* gene, rather

than being recombined (Figure 1A'). In *jbug*[133], homologous recombination failed to occur; instead,

the entire donor construct was inserted into the 3' UTR of the splice forms that have longer C-

termini (*jbug-RI* and *jbug-RF/M/N*) (Figure 1A''). These recombination events disrupted *jbug-RI*

and *jbug-RF/M/N* in both alleles. Western blot analysis revealed that the Jbug-PF/M/N isoforms

(>300 kDa) were either absent or decreased to undetectable levels in *jbug*[30] mutants at all stages

(Figure 1E). Jbug-PI (185.7 kDa) was also absent in *jbug*[30], and instead, a smaller protein (~90

kDa) was detected in *jbug*[30] mutant animals at all stages (Figure 1, E and H). This additional band

was not detected in wild-type or *jbug*[133] protein extracts at any stage, suggesting that insertion of

the DNA construct generates a truncated form of Jbug-PI in *jbug*[30]. In *jbug*[133], both Jbug-PF/M/N

and Jbug-PI protein bands were slightly decreased in intensity (Figure 1, E and H), suggesting

that insertion of the construct in the 3' UTR region either affects mRNA stability or inhibits

translation of these isoforms. Consistent with the molecular data, we did not detect any significant

changes in the level of Jbug-PC (42.4 kDa) and Jbug-PL (175.6 kDa) in *jbug*[30] and *jbug*[133] protein

extracts (Figure 1, E and H). Taken together, we generated new alleles for *jbug*, including a null

mutant and two hypomorphic alleles that disrupt a subset of Jbug isoforms, specifically Jbug-PI

and Jbug-PF/M/N.

jbug hypomorphs show semi-lethality and defects in bristle development on the thorax

Many homozygous flies in *jbug*[30] and *jbug*[133] died during pupation, leaving a number of dead

pupae that never eclosed. In both alleles, homozygous adult flies that survived were fertile and

homozygous lines could be maintained. Interestingly, adult flies homozygous for *jbug*[30] or *jbug*[133]

produced short, bent and rough-ended bristles on the thorax (Figure 2A), suggesting a role for

Jbug in bristle development, perhaps in crosslinking actin bundles that form the bristles.

Consistent with our findings that levels of Jbug-PF/M/N and Jbug-PI were more significantly

reduced in *jbug*[30] than in *jbug*[133] (Figure 1, E and H), *jbug*[30] flies showed a higher expressivity of

bristle defects than *jbug*[133]. Whereas almost every thoracic bristle had defects in *jbug*[30]

homozygous adults, only a few bristles showed distinguishable defects in *jbug*[133] homozygotes

(Figure 2A). Phalloidin staining in *jbug*[30] pupae showed changes in F-actin staining consistent

with the bent bristle phenotype (Figure 2B), suggesting that the bristle defects arise during

pupation.

To analyze the bristle defects in more detail, we performed scanning electron microscopy (SEM)

in wild-type and *jbug*[30] homozygous mutant adults. Whereas wild-type animals showed nicely

elongated and tapered bristles that suggested evenly aligned actin bundles (Figure 2C, a and a'),

jbug³⁰ mutant flies had bristles that were short, curved (Figure 2C, c), bent (Figure 2C, d) and

rough-ended (Figure 2C, e). Even when the bristles looked relatively normal, they were often

shorter than wild type and mildly twisted (Figure 2C; compare b and b' to f and f').

Transheterozygotes of *jbug^{30}* or *jbug^{133}* over the *jbug^{20}* null allele or the *59AB* deficiency line that

deletes the entire *jbug* gene (Figure 1A) showed both semi-lethality and the same bristle defects

as observed in homozygous *jbug^{30}* and *jbug^{133}* adults (Table 1). These data suggest that both

phenotypes are due to loss of *jbug* function. Consistent with our data that *jbug^{30}* is a more severe

hypomorphic allele than *jbug^{133}* (Figure 1, E and H), fewer transheterozygous *jbug^{30}*/*jbug^{20}* or

jbug^{30}/*59AB* flies survived to the adult stage compared to transheterozygous *jbug^{133}*/*jbug^{20}* or

jbug^{133}/*59AB* flies (Table 1). Taken together, these data suggest that Jbug-PF/M/N and/or PI are

important for viability and actin bundle arrangement within bristles.

We also performed complementation tests using the deficiency line *Exel6079*, which deletes

through the genomic region encoding the N-terminal exons of transcripts of *jbug-RL* and *jbug-*

RF/M/N but leaves the coding region for *jbug-RC* and *jbug-RI* intact (Figure 1A). Interestingly,

transheterozygous *jbug^{30}*/*Exel6079* (and *jbug^{133}*/*Exel6079*) flies showed neither semi-lethality nor

bristle defects (Table 1). To test which isoforms are present in *jbug^{30}*/*Exel6079* flies, we performed

Western blot using extracts from *jbug^{30}*/*Exel6079* transheterozygous 1st instar larvae obtained

from *jbug^{30}* homozygous mothers and found that the truncated PI (~90 Kd) band present in *jbug^{30}*

is no longer detected; only a full-length PI band is observed in the transheterozygotes. Also

surprisingly, the large MW Jbug-PF/M/N bands that were absent/reduced in *jbug^{30}* homozygotes

were easily detected in *jbug^{30}*/*Exel6079* larvae (Figure 1G). This finding was unexpected because

Jbug-PF/M/N should be absent in transheterozygous *jbug^{30}*/*Exel6079* flies; *jbug^{30}* and *Exel6079*

disrupt the 3' and the 5' ends of *jbug-RF/M/N*, respectively. Since *jbug^{30}*/*Exel6079* larvae were

obtained from *jbug^{30}* homozygous mothers, Jbug-PI or Jbug-PF/M/N cannot be provided

maternally. This result is challenging to reconcile, unless there is some form of transvection or

transplicing that would generate full length transcripts coming from the 5' regions of *jbug^{30}* and 3'

regions of *Exel6079*. Such events could also explain the recovery of the full-length *jbug-RI*

transcript and corresponding loss of the shortened form of Jbug-PI (Figure 1G).

Expression of Jbug-RF rescues the bristle defects and semi-lethality of *jbug* hypomorphs

To begin to parse out the functions of the different Jbug isoforms, we created three new UAS

lines encoding Jbug-PC (42 kDa protein), Jbug-PI (185.6 kDa) and Jbug-PF (>300 kDa protein).

We also obtained a UAS construct for Jbug-PL (175.6 kDa protein) expression from the Mlodzik

lab (Olguín *et al.* 2011). We first tested whether expression of any specific *jbug* splice form(s)
could rescue the lethality of *jbug²⁰* null mutants. Each UAS construct was expressed in *jbug²⁰*
mutant background using *daughterless-Gal4* (*da-Gal4; Wodarz et al., 1995*), which is expressed
in all cells, or *69B-Gal4* (Rorth 1996), which is expressed in ectodermal cells. None of the UAS
constructs rescued the lethality of *jbug²⁰*; no larvae survived past the 1st instar larval stage as in
jbug²⁰. Co-overexpression of *jbug-RL* and *jbug-RI* together using da-Gal4 in the *jbug²⁰* mutant
background also did not rescue the lethality. These data suggest that multiple Jbug isoforms are
required for viability.

We next tested whether expression of specific Jbug isoforms could rescue the bristle defects in
jbug³⁰, the severe hypomorphic allele. Consistent with the data that Jbug-PC is not affected in
jbug³⁰ mutants, overexpression of *jbug-RC* with *da-Gal4* or with *69B-Gal4* in an otherwise wild-
type background did not result in any overt defects in bristles (Figure 2D). Overexpression of
Jbug-PI in an otherwise wild-type background also did not cause any defects in bristles (Figure
2E). On the other hand, overexpression of *jbug-RF* in otherwise wild-type animals resulted in
shorter and thicker bristles (Figure 2F), suggesting that proper levels of Jbug-PF/M/N are critical
for normal bristle development. Phalloidin staining in *fkh>Jbug-RF* pupae showed the bent bristle
phenotype and irregular F-actin alignment, similar to observed in *jbug³⁰* pupal bristles (Figure 2B).
Importantly, whereas expression of *jbug-RC* (Figure 2G) or *jbug-RI* (Figure 2H) in *jbug³⁰* mutants
did not rescue the bristle phenotypes, expression of *jbug-RF* in *jbug³⁰* mutants using *da-Gal4* or
69B-Gal4 nearly completely rescued the bristle phenotypes (Figure 2I). Moreover, *jbug-RF*
expression in *jbug³⁰* mutants also rescued semi-lethality. 86% (27/32) of pupae emerged as adult
flies when *jbug-RF* was expressed in *jbug³⁰* mutants, whereas only 26% (9/35) of *jbug³⁰*
homozygous pupae emerged as adult flies. These data suggest that the bristle defects and semi-
lethality observed in *jbug³⁰* are due to loss of the longest (>300 kDa) Jbug isoforms.
407

Jbug hypomorphs show a delay in actin-based prehair formation and mild defects in wing
hair orientation
Besides thoracic bristle defects, *jbug³⁰* and *jbug¹³³* homozygous adults also showed a mild swirling
pattern of wing hairs (Supplemental Figure 3B and data not shown). Phalloidin staining of actin-
rich prehairs at 32 hours after puparium formation (APF) revealed that hair formation is delayed
in *jbug³⁰* pupal wings (Supplemental Figure 3, D and E). A swirling pattern of wing hairs and
delayed prehair formation are often observed in mutants for planar cell polarity (PCP) genes
(Adler 2012). *jbug³⁰* mutant eyes, however, did not have any defects in ommatidial rotation

characteristic of loss of key PCP genes (Supplemental Figure 3, F and G), suggesting that Jbug

is not involved in ommatidial rotation. Overall, these data suggest that proper levels for Jbug-

PF/M/N and/or Jbug-PI are required for the timely maturation of the actin bundles that form the

prehairs. In support of this, we find that the wing hair polarity defects are fully rescued in

jbug30/Exel6079 flies (Supplemental Figure 3C).

Jbug proteins localize in a dynamic pattern in the embryonic epithelium

We next analyzed the subcellular localization of Jbug proteins during embryogenesis in wild type

and *jbug* mutant alleles. During early development, Jbug was detected as small punctate

structures in the cytoplasm of all cells (Figure 3, A and B). During stages 11-13, Jbug was

detected at a high level both near the apical surface and at and near the adherens junctions (AJs)

in all epidermal cells (Figure 3, C and M). AJ Jbug signals became weaker over time, and by

stage 15, only mesh-like signals on the apical surface remained (Figure 3, D and M). In *jbug[20]*

mutant embryos, AJ signals were absent, but signals on the apical surface were still detected

(Figure 3, E and M). At stage 15 and later, apical surface signals were significantly decreased

compared to wild-type levels (Figure 3, F and M). Since the *jbug[30]* and *jbug[133]* homozygotes that

survived to adult stages were fertile, we analyzed Jbug signals in *jbug[30]* or *jbug[133]* embryos from

homozygous mothers. Interestingly, in *jbug[30]* embryos from *jbug[30]* homozygous mothers, Jbug

signals were both very weak and dispersed in epidermal cells throughout the stages and largely

absent in AJs (Figure 3, G, H and M), suggesting that Jbug-PF/M/N and Jbug-PI might be major

isoforms that show strong embryonic expression and AJ localization. Expression of Jbug-RF in

the *jbug[30]* mutant background restored strong signals at AJs (Figure 3K) and on the apical surface

(Figure 3L) at stage 13 and 15, respectively. Consistent with the minor changes in Jbug protein

levels in *jbug[133]*, Jbug staining was mostly normal, albeit slightly reduced, in the epidermal cells

in *jbug[133]* embryos from homozygous *jbug[133]* mothers (Figure 3, I, J and M).

Jbug is required for denticle formation in the epidermis

Jbug antibody staining in the ventral regions of stage 14 embryos revealed intense signals on the

apical surface that presaged the actin-rich denticles that begin to form during stage 15 (Figure 4,

B and C). This prompted us to ask if denticle formation was normal in *jbug* mutants. In wild-type

embryos, six rows of epidermal cells in the anterior portion of each abdominal segment normally

produce one or more denticles, forming a trapezoidal belt (Bejsovec 2013; Figure 4, C and D).

Embryos homozygous for *jbug[20]* did not show any overt morphological defects, likely due to strong

maternal contribution, but they exhibited defects in denticle formation (Figure 5B). Compared to

wild type (Figure 5A), many denticles in *jbug[20]* embryos were small and premature (Figure 5B).

Also, the overall number of denticles were fewer in *jbug[20]* embryos compared to wild type. We

counted the number of denticles in each cell from a grid of 24 cells (Figure 5K) for at least five

individuals of each genotype. *jbug[20]* embryos had fewer denticles per cell than wild-type embryos

in several rows of cells (C1, C3, C4 and C6; Figure 5, L and M; Supplemental Figure 4, A and B).

Denticles in *jbug[30]* embryos from a homozygous stock were relatively normal in their size and

morphology, but these embryos had significantly fewer denticles compared to wild type (Fgure

5C). Irrespective of which row of cells (C1-C6) was examined, *jbug[30]* embryos had, on average,

fewer denticles per cell than wild-type embryos (Figure 5, L and M; Supplemental Figure 4, A and

B). These data suggest important roles of Jbug in actin bundle arrangement for denticle formation.

We next tested whether expression of *jbug-RF* or *jbug-RI* rescues the denticle defects in *jbug[20]*

or *jbug[30]* embryos. Expression of *jbug-RF* using the *da-Gal4* driver in the otherwise wild-type

condition did not affect the number of denticles nor the size (Figure 5, F, L and M). Expression of

jbug-RF using the *da-Gal4* driver in *jbug[20]* mutants largely restored denticle size (compare Figure

5B and 5D), suggesting a role of the Jbug isoforms with actin-binding domains in formation of

mature denticles. However, Jbug-RF expression in *jbug[20]* minimally rescued the decreased

denticle number phenotype; it slightly increased the number of denticles in one row (C6), but not

significantly in other rows (C1-C5) (Figure 5, D, L and M; Supplemental Figure 4, A and B).

Similarly, expression of *jbug-RF* using the *da-Gal4* driver in *jbug[30]* mutants slightly increased the

number of denticles in one row (C1), but not in other rows (C2-C6) (Figure 5, E, L and M;

Supplemental Figure 4, A and B). On the other hand, overexpression of *jbug-RI* using the *da-Gal4*

driver in the otherwise wild-type condition significantly increased the number of denticles in

several rows (C2-C4; Supplemental Figure 4, A and B). In all rows, Jbug-RI-overexpressing

embryos have more cells with more than three denticles per cell than wild type (Figure 5L;

Supplemental Figure 4, A and B). Moreover, expression of *jbug-RI* using *da-Gal4* in *jbug[30]*

mutants significantly rescued the decreased denticle number phenotype in several rows (Figure

5, G, L and M; Supplemental Figure 4, A and B). These data suggest a critical role of Jbug-PI in

formation of the actin-rich denticles in the embryo.

We further tested whether the maternal pool of *jbug* is required for denticle formation. Knocking

down *jbug-RL* and *jbug-RF/M/N* by RNAi (*GD13033*; Figure 1A) using *da-Gal4* also showed

similar, albeit weaker, defects to those in *jbug[20]* embryos: a decrease in the size and numbers of

denticles per cell in several rows (Figure 5, I, L and M; Supplemental Figure 4, A and B),

suggesting roles for the Jbug isoforms with actin-binding domains in denticle formation. Maternal

knockdown of *jbug* using the *matα-Gal4* driver (Häcker and Perrimon 1998) and two strong RNAi

lines (*GD8664* and *GD13033*; each targets a subset of *jbug* splice forms; Figure 1A) resulted in

very few embryos, suggesting an essential role for Jbug in oogenesis. Maternal knockdown of

jbug using the weaker RNAi line (*JF01166* that targets all splice forms; Figure 1A) resulted in

mostly normal overall embryonic morphology, except for slightly wavy epidermis (Figure 5J), and

these embryos showed defects in denticle formation. As observed in *jbug*[20] and *jbug*[30] mutants or

da-Gal4>jbug RNAi GD13033 embryos, some cells failed to form denticles; others formed short,

premature denticles with weak phalloidin signals (Figure 5, J, L and M). Quantification showed a

decrease in the average number of denticles per cell in most columns (except for C2) of each

denticle belt with *matα>jbug RNAi* (Figure 5, L and M; Supplemental Figure 4, A and B). Taken

together, our data suggest that both maternally and zygotically provided Jbug proteins are

required for denticle formation.

Jbug is upregulated in embryonic epithelial tubes

Consistent with mRNA expression (BDGP; https://insitu.fruitfly.org/cgi-bin/ex/insitu.pl; Fly-FISH;

http://fly-fish.ccbr.utoronto.ca; Supplemental Figure 1), antibody staining in wild-type embryos

revealed accumulation of Jbug protein in the trachea and the salivary gland (Figure 6, A and E).

Interestingly, Jbug was abundant in both the cytoplasm and in a distinct apical domain of both

tracheal and salivary gland cells, which appeared to be just under the apical surface (Figure 6, A

and E), as opposed to following the AJ pattern of E-Cad. This apical staining in the salivary gland

was prominent up to stage 14; at later stages, Jbug signals were observed mostly in the cytoplasm

and detected in the apical domain only as a thin layer along the lumen (compare Figure 6E to

Figure 7A).

Jbug signals in tubular epithelial organs were significantly decreased in *jbug*[20] embryos, even with

regards to surrounding tissues (Figure 6, B and F), suggesting that zygotic *jbug* expression is

required for upregulation of Jbug in these tissues. In *jbug*[20], the apical staining in the trachea was

absent and in the salivary gland was more punctate. In *jbug*[30] embryos (from *jbug*[30] parents),

apical staining of Jbug in the trachea and salivary glands was also absent; instead, uniform Jbug

signals were detected in the cytoplasm (Figure 6, C and G). Consistent with slight changes in

Jbug protein levels in *jbug*[133], Jbug signals were mostly normal, albeit slightly reduced, in the

trachea and salivary glands in *jbug*[133] embryos from *jbug*[133] parents (Figure 6, D and H). These

data suggest that it is either the Jbug-PF/M/N and/or Jbug-PI that are enriched in the apical

domain in tubular epithelial organs in the embryo.

Despite the absence or mislocalization of Jbug in the apical domain of trachea and salivary glands,

both organs developed normally without overt morphological defects in *jbug*[20] and *jbug*[30] embryos

(Figure 6, B, C, F and G). Crumbs (Crb), the apical determinant protein, E-Cadherin (E-Cad), a

key adherens junction protein, Discs large (Dlg), a septate junction component, and α-Spectrin

(α-Spec), the lateral membrane-associated protein, also localized properly in salivary glands in

jbug[20] or *jbug*[30] embryos (Supplemental Figure 5, A-I), suggesting that the overall epithelial

polarity is intact in salivary glands in these mutants. Also, phalloidin signals in salivary glands in

jbug[20] and *jbug*[30] embryos were comparable to wild type (Supplemental Figure 5, D-F), suggesting

that F-actin is largely intact in salivary glands in *jbug* mutants.

To further test where each of the jbug isoforms localizes and to test the effect of Jbug

overexpression in tubular epithelial tissues, we overexpressed the different Jbug isoforms (Jbug-

PC, PI, PL and PF) in the salivary gland using the salivary gland-specific *fork head-Gal4* driver

(*fkh-Gal4*; Henderson and Andrew 2000). All isoforms primarily localized in the cytoplasm when

overexpressed (Figure 7, D-G). Interestingly, however, both untagged and C-terminal GFP-

tagged Jbug-PC were detected both in the cytoplasm and in nuclei (Figure 7, B and C). These

data suggest that this smallest isoform – which is primarily expressed in pupae (Figure 1E) –

might function in the nucleus as well. Similar nuclear localization has been observed with human

Filamin A. Although full-length filamin A is predominantly cytoplasmic, a C-terminal fragment of

Filamin A colocalizes with the androgen receptor to the nucleus to act as a co-transcription factor

(Loy *et al.* 2003). Overexpression of each isoform resulted in an expanded and irregular luminal

domain (Figure 7, B-G). Interestingly, Crb was mislocalized and dispersed along the entire apical

surface in Jbug-overexpressing salivary glands (Figure 7, B-G) rather than being enriched in a

sub-apical domain just apical to the AJs as in control salivary glands (Fig 7A, inset in Crb column;

Wodarz *et al.* 1995; Tepass 1996; Chung and Andrew 2014). These data indicate that

overexpression of multiple Jbug isoforms can disrupt the sub-apical domain enrichment of Crb.

Phalloidin and other junctional and polarity markers, including E-Cad, Nrx-IV (a septate junction

marker) as well as α-Spec, localized properly in Jbug-overexpressing salivary glands

(Supplemental Figure 6), suggesting that the overall epithelial polarity is not disrupted by Jbug

overexpression. Taken together, these data suggest that appropriate levels of Jbug isoforms are

important for proper Crb localization and apical organization of epithelial cells during

embryogenesis.

Jbug is required for maintaining Crb levels in embryonic epithelial cells

Maternal knockdown (*matα>JF01166*) or zygotic loss of all *jbug* splice forms (*jbug[20]*), or specific

maternal and zygotic loss of the *jbug-RF/M/N* and *jbug-RI* (*jbug[30]*) did not show overt

morphological defects in the embryo, except for mild wavy epidermis (Figures 5, B and J) and

denticle formation defects (Figure 5). We therefore knocked down both maternal and zygotic pools

of all *jbug* splice forms using RNAi. Compared to normal morphology in control embryos (Figure

8, A and E) or in embryos knocked down for *jbug* zygotically (using *da-Gal4*; Figure 8, B and F)

or maternally (using *matα*-Gal4; Figure 8, C and G), knockdown of *jbug* both maternally and

zygotically using *matα-Gal4* and *da-Gal4* (*jbug* MZ knockdown) resulted in a severely disrupted

morphology in all embryos produced (Figure 8, D and H). The embryos were often so disrupted

that their stages could not be ascertained. Whereas strong Jbug signals were still detected in

epithelial cells in embryos knocked down for *jbug* maternally or zygotically (Figure 8, F'' and G'';

compare to control in Figure 8E''), Jbug signals were absent in the apical surface in many

epidermal cells in *jbug* MZ knockdown embryos (Figure 8H''). Importantly, Crb signals were also

absent in those cells (Figure 8H'''). These data suggest a critical role for Jbug in regulating Crb

levels and localization in epithelial cells.

We also knocked down *jbug* in the salivary gland in embryos also maternally knocked down for

jbug, using *matα-Gal4* and *fkh-Gal4*. No overt morphological defects were observed in salivary

glands in these embryos at stage 11, when the salivary gland begins to invaginate to form a tube

(Supplemental Figure 7). However, compared to continuous Crb signals in control or in salivary

glands knocked down for *jbug* maternally or zygotically only, huge gaps in Crb signals were

detected in salivary glands in *jbug* MZ knockdown embryos (Supplemental Figure 7, A''''-D''''). At

later stages, knockdown of *jbug* resulted in a slightly irregular salivary gland lumen (Figure 8, I-

M). Importantly, compared to the continuous sub-apical localization of Crb in control or SG cells

with zygotic knockdown for *jbug*, the salivary glands knocked down for *jbug* maternally showed

discontinuous Crb signals (Figure 8I'''). Salivary glands knocked down for *jbug* both maternally

and zygotically showed reduced Crb signals with wider gaps (Figure 8L''') or mislocalization of

Crb signals to the apical surface (Figure 8M'''). Overall, these data reveal that Jbug is required

for proper levels and distribution of Crb in epithelia.

Discussion

Filamins play important roles in actin reorganization in many developmental processes and disease contexts. In this study we show that different isoforms of Jbug, a *Drosophila* filamin-type protein, are differentially expressed and localized during development. Using new genetic tools that we generated, we reveal essential roles for different Jbug isoforms in viability, epithelial morphogenesis and formation of actin-rich structures.

Roles of different Jbug isoforms in various developmental stages in *Drosophila*

Our developmental western analysis (Figure 1), along with developmental RNA-Seq data by Flybase (www.flybase.org; FB2021_02), reveals differential expression of different Jbug isoforms throughout development. Although transcripts of isoforms containing actin-binding domains (*jbug-RF/M/N* and *jbug-RL*) are not expressed until several hours of embryogenesis, proteins for these isoforms are detected at early hours of embryogenesis, suggesting Jbug proteins are provided maternally. Our studies reveal roles of different Jbug isoforms at various developmental processes, including oogenesis, epithelial morphogenesis, and viability at larval and pupal stages.

Severe defects in egg laying upon maternal knockdown of *jbug* using two strong RNAi lines that target a different subset of isoforms (*GD13033* and *GD8664*) suggest critical roles for Jbug isoforms in oogenesis. *GD13033* targets Jbug-PF/M/N and Jbug-PL, suggesting roles for the actin-binding domain-containing isoforms in oogenesis. Homozygous *jbug*[30] flies can be maintained as a homozygous stock depite the apparent absence of Jbug-PF/M/N and truncated Jbug-PI, suggesting that these Jbug isoforms do not play an essential role in oogenesis. Since Jbug-PC is only made in pupae, the main isoform for oogenesis is likely to be Jbug-PL (Figure 9).

Our study also reveals a critical role of Jbug in epithelial morphogenesis during embryogenesis. Embryos knocked down for *jbug* both maternally and zygotically show severe morphological defects with disrupted Crb localization (Figure 8). The precise mechanism underlying Crb regulation by Jbug remains to be discovered. As filamin binds to many interacting proteins via IG-FLMN repeats, it will be interesting to test whether Jbug regulates Crb by directly interacting with it. It is also possible that Crb reduction is an indirect effect of disrupted actin cytoskeleton by loss of Jbug. Dynamic localization of Jbug isoforms to the apical surface during embryogenesis (Figures 3 and 4) suggests potential roles for different Jbug isoforms in remodeling the actin cytoskeleton and contributing to the mechanical stability of the plasma membrane and the cell cortex. Interestingly, homozygous *jbug*[30] embryos from homozygous mothers, where only Jbug-PC and Jbug-PL are intact, show no overt defects during embryogenesis except for defects in

denticle formation at later stages (Figure 5). As Jbug-PC is not expressed during embryogenesis

(Figure 1), these data indicate that Jbug-PL is largely sufficient for epithelial morphogenesis

during embryogenesis (Figure 9).

Jbug is also required for viability. The $jbug^{20}$ null allele lacking all Jbug isoforms is 1st instar larval

lethal. 1st instar larvae of homozygous $jbug^{20}$ or transheterozygous animals of $jbug^{20}$ and a

deficiency line appear normal in their overall size, morphology and motility, but they do not molt

into the 2nd instar. Based on the role of Jbug in regulating Crb localization during embryogenesis

(Figures 7 and 8), the lethality of $jbug^{20}$ homozygous larvae might be linked to disruption of Crb

localization and/or epithelial organization during the larval stage. As overexpressing a single

isoform or Jbug-RL and Jbug-RI together failed to rescue the lethality of $jbug^{20}$, multiple Jbug

isoforms may be required for viability during early stages of development.

Our $jbug$ hypomorphic alleles reveal roles for Jbug-PF/M/N and/or Jbug-PI in organismal viability

late in development (Table 1). Most flies homozygous for $jbug^{30}$ or $jbug^{133}$, or transheterozygous

flies for these $jbug$ alleles and a deficiency completely removing $jbug$ die during late pupation.

Jbug-PF expression in the $jbug^{30}$ mutant background rescued this semi-lethality, suggesting that

the longest splice forms (PF/M/N) are required at these late stages. Interestingly, anatomy RNA-

Seq by Flybase (www.flybase.org; FB2021_02) shows increased $jbug$ expression in the fat body

at pupal stage P8 (2 days old; moderate levels of expression) compared to the white prepupae

stage (very low expression), opening up the possibility for a role for Jbug in the fat body. The fat

body of $Drosophila$ is equivalent to vertebrate adipose tissue and liver in its storage and major

metabolic functions, including immune response and sensing nutritional conditions. It remains to

be revealed whether the lethality caused by the loss (or significant reduction) of Jbug-PF/M/N is

related to defects in morphogenesis, metabolic functions, or both.

Jbug plays a role in formation of actin-rich protrusions

Our results identify roles for specific Jbug isoforms in formation of two prominent actin-rich

protrusions in $Drosophila$, bristles on the thorax and denticles in the ventral epidermis of the

embryo (Figures 2 and 5). Jbug may regulate assembly of actin bundles by interacting with actin-

regulating proteins, such as capping protein and the Arp2/3 complex components. Mutants for

capping protein or the Arp2/3 complex components show bristle defects reminiscent of the $jbug$

mutant phenotypes (Hudson and Cooley 2002; Frank et $al.$ 2006). Several studies in other

systems also suggest that the Arp2/3 complex synergizes with filamins. In filamin-deficient

melanoma cells, the Arp2/3 complex is still abundant but unable to stabilize the plasma

membranes or support locomotion (Cunningham *et al.* 1992). Also, branched actin filaments

made with Arp2/3 do not have the closing loop structures necessary to form three-dimensional

structures (Mullins *et al.* 1998; Blanchoin *et al.* 2000), unless the filaments are assembled in the

presence of filamin (Niederman *et al.* 1983; Wolosewick and Condeelis 1986). Other good

candidates that might interact with Jbug to organize the actin cytoskeleton include the Rho family

small GTPases and their regulators. The Rho family small GTPases, including Rho, Rac, and

Cdc42 are known to physically interact with Filamin (Ohta *et al.* 1999; Jeon *et al.* 2008; Del Valle-

Pérez *et al.* 2010), as are regulators of Rho GTPases, including guanine nucleotide exchange

factors (Bellanger *et al.* 2000; Pi *et al.* 2002), GTPase activating proteins (Ohta *et al.* 2006;

Mammoto *et al.* 2007; Nakamura *et al.* 2009) and Rho kinase (Ueda *et al.* 2003). It will be

interesting to test whether Jbug works with any of these proteins to regulate bristle and/or denticle

formation.

Defects in actin-rich protrusions in *jbug*^{30} are observed in the presence of Jbug-PL, which contains

actin-binding domains and short IG-FLMN repeats, suggesting that this isoform is not sufficient to

form normal bristles and denticles in *Drosophila*. Indeed, bristle defects are rescued by expression

of Jbug-PF, one of the longest isoforms that contain long IG-FLMN repeats following the actin-

binding domains (Figure 2). These data suggest an important role for protein interactions through

the long IG-FLMN repeats in forming actin-rich protrusions. Our interpretation is consistent with a

role of filamins in acting as a molecular scaffold. More than 90 filamin-binding partners have been

identified, including intracellular signaling molecules, receptors, ion channels, transcription factors,

and cytoskeletal and adhesion proteins (Razinia *et al.* 2012).

Surprisingly, defects in denticle numbers in *jbug*^{30} embryos are rescued by expression of Jbug-

PI, an isoform that contains long IG-FLMN repeats without the actin-binding domains, but only

minimally by expression of Jbug-PF (Figure 5). These data suggest a major role of Jbug-PI in

denticle formation (Figure 9). Although Jbug is upregulated in the anterior side of each segment

in the embryo, where denticle belts form, Jbug is not enriched in the denticles themselves (Figure

4). Interestingly, another *Drosophila* filamin – Cher – is enriched in denticles, colocalizing with

actin protrusions (Dilks and DiNardo 2010). The different subcellular localization of Jbug and Cher

in denticle forming cells suggests distinct roles for the two *Drosophila* filamins. Cher likely acts as

a component of denticles that organizes actin filaments. A simple model for Jbug function is that

actin-binding domains might act to stabilize actin filaments and that protein-protein interactions

through long IG-FLMN repeats might recruit other key components for denticle formation to the

apical plasma membrane where actin-rich protrusions arise. Still, Jbug-RF/M/N and/or Jbug-RL

appear to contribute to denticle formation. Although overexpression of *jbug-RF* in *jbug*[20] or *jbug*[30]
mutant embryos does not fully rescue the decrease of denticle numbers (Figure 5), zygotic
knockdown of *jbug-RF/M/N* and *jbug-RL* using the RNAi line specifically targeting them (*GD13033*)
results in premature/small denticle formation and also decreases the number of denticles in some
rows (Figure 5), suggesting roles for the actin-binding domain-containing isoforms in denticle
formation (Figure 9).

Jbug is likely to affect PCP indirectly

jbug[30] and *jbug*[133] alleles show a delay of actin-rich prehair formation in the pupal wing and a mild
swirling pattern of adult wing hairs (Supplemental Figure 3). Both phenotypes are often shown in
mutants of PCP genes (Adler, 2002). *jbug* mutants, however, do not show any defects in
ommatidial rotation, suggesting that Jbug is less likely to directly affect the core PCP pathway.
Rather, the phenotypes in pupal/adult wing hairs are likely to be caused by defects in formation
of actin protrusions, similar to what we observe in thoracic bristles (Figure 2) and denticles (Figure
5) in *jbug* mutants. Consistent with this idea, Jbug has been proposed to modulate the mechanical
properties of actin filaments in tendon cells to regulate planar orientation of thoracic bristles;
alterations in the ability to adjust tensile forces upon *jbug* knockdown disrupt the planar orientation
of bristles (Olguin et al., 2011).

Roles of Jbug in formation of tubular organs

Several studies suggest roles for filamin A in tubular epithelial morphogenesis and diseases in
tubular organs. During branching morphogenesis of mammary glands, the levels of filamin A
expression and the extent of the integrin-filamin interaction modulate collagen remodeling to
optimize matrix stiffness to support tubulogenesis (Gehler *et al.* 2009). Filamin A is also an
important regulator for Polycystin-2, a key molecule in autosomal polycystic kidney disease
(Wang *et al.* 2015). Our data also suggests a role for Jbug in apical domain reorganization in
tubular epithelial cells (Figures 7 and 8). Jbug shows strong apical enrichment in the early
embryonic trachea and the salivary gland (Figure 6) and both Jbug loss and overexpression cause
defective tube morphology with disrupted Crb localization along the apical surface of salivary
gland cells (Figures 7 and 8). These data support the idea that appropriate levels of Jbug are
important for apical organization during tubular organ development. Loss of apical Jbug
enrichment, however, does not cause overt defects in tubular organ formation in *jbug* null or
hypomorphic alleles (Figure 6). In these alleles, abundant cytoplasmic Jbug signals still remain,
and the levels or distribution of Crb was not noticeably changed (Supplemental Figure 5).

Overexpressed Jbug also mainly localizes to the cytoplasm in salivary gland cells, with only slight

apical enrichment (Figure 7). These data suggest that apical enrichment may not be necessary

for Jbug function in the formation of tubular epithelia.

Transvection or trans-splicing might occur in the *jbug* locus

The surprising rescue that we have observed of viability, bristle morphology and wing hair

orientation of *jbug^{30}* and to a lesser extent with *jbug^{133}* with *Exel6079*, a deficiency that deletes

all *jbug* exons upstream of the transcription start sites of *RC* and *RI* (Table 1; Supplemental Figure

3), suggests epigenetic regulation. One possible mechanism is transvection, an epigenetic

phenomenon that results from interactions between alleles on homologous chromosomes in

tissues where the chromosomes are paired (King *et al.* 2019), typically with enhancers of one

allele on one chromosome affecting expression of the other allele on the homologous

chromosome. Here, however, we have to propose transvection via template-switching, wherein

transcripts from the *jbug^{30}* allele would switch templates at some point outside the *Exel6079*

deficiency but before the large insertion in *jbug^{30}*, for example. Another possibility is trans-splicing,

a gene regulatory mechanism that joins exons from two separate transcripts to produce chimeric

mRNA, and has been detected in most eukaryotes (Horiuchi and Aigaki 2006; Lasda and

Blumenthal 2011). In *Drosophila*, trans-splicing has been demonstrated to be an essential

process for two genes, *longitudinals lacking* (*lola*) and *modifier of mdg4* (*mod(mdg4)*). In the *lola*

locus, the 5'-end of *lola* transcripts contain five exons that splice into 20 variants of 3' exons to

generate 20 protein isoforms (Ohsako *et al.* 2003). In the *mod(mdg4)* locus, two mutant alleles of

the gene, each located in one of the two homologous chromosomes, restore the wild-type function

of the gene (Mongelard *et al.* 2002), much like is observed here. The presence of Jbug-PF/M/N

isoforms (and the rescue of PI size) in transheterozygous animals for *jbug* hypomophs and

Exel6079 (Figure 1G) supports the idea that trans-splicing might occur, where the 5' exons of

jbug-RF/M/N transripts on *jbug^{30}* or *jbug^{133}* are spliced to 3' exons of *jbug* transcripts on the

Exel6079 chromosome. Recent studies have revealed conserved sequences in the *mod(mdg4)*

intron that promote trans-splicing in this gene (Gao *et al.* 2015; Tikhonov *et al.* 2018). Identification

of the sequences that promote trans-splicing in the *jbug* locus will help better understand this

surprising gene regulatory mechanism.

Table 1. Complementation test

	Number of Progeny	
	CyO	Non-CyO
jbug^{20}/CyO x *jbug^{20}/CyO*	>200	0
jbug^{20}/CyO x *Df(2R)59AB/CyO*	>200	0
jbug^{20}/CyO x *Df(2R)Exel6079/CyO*	>200	0
jbug^{30}/CyO x *jbug^{20}/CyO*	176	12*
jbug^{30} (F) x *jbug^{20}/CFL* (M)	144	50*
jbug^{30}/CyO x *Df(2R)59AB/CyO*	231	41*
jbug^{30}/CyO x *Df(2R)Exel6079/CyO*	98	91
jbug^{133}/CyO x *jbug^{20}/CyO*	141	62*
jbug^{133} (F) x *jbug^{20}/CFL* (M)	64	70*
Jbug^{133}/CyO x *Df(2R)59AB/CyO*	153	47*
jbug^{133}/CyO x *Df(2R)Exel6079/CyO*	140	100
* Transheterozygous flies show the same bristle defects as *jbug^{30}* or *jbug^{133}* homozygotes (See Figure 2 for phenotypes in homozygotes).		

Figure legends

Figure 1. Differential expression of Jbug isoforms throughout *Drosophila* development and generation of null and hypomorphic alleles of *jbug*. (A) Ten annotated splice forms of *jbug* are reported by FlyBase. Deletion regions for two deficiency lines in the *jbug* locus and the *jbug*[20] null allele are indicated. Cyan bars, three RNAi lines for *jbug* that were tested in this study. Magenta bar, a common exon for all isoforms that was used to generate an antibody against Jbug. Light green bars, regions used for making antisense probes for in situ hybridization. (A', A'') Molecular alterations in two hypomorphic alleles of *jbug*. Blue and red boxes represent homology arms that were used for homologous recombination. Genomic sequence locations are shown for the homology arms. Hypomorphic alleles were generated by incomplete homologous recombination, with a partial (*jbug*[30], A') or a complete (*jbug*[133], A'') insertion of the donor DNA into the *jbug* isoforms that have a longer C terminus (*RF/M/N* and *RI*). (B) Six different Jbug protein isoforms are made from the different annotated splice forms. (C) Developmental RNA-Seq data by Flybase (www.flybase.org; FB2021_02). Note that only minimal expression of *jbug-RF/M/N* and *jbug-RL* is detected until 10 hours of embryogenesis (red arrow). (D-F) Western blot using protein extracts from wild type and *jbug* mutants at different stages. Bands that correspond to the predicted molecular weights of several isoforms are marked with dots and arrowheads with different colors. Arrowheads of the same colors are used to indicate the major isoforms in B. (D) Developmental western blot using protein extracts from embryos (0-2 and 2-16 hours AEL at 25°C), 1st, 2nd and 3rd instar larvae, pupae (2 days APF) and male (M) and female (F) adults (2 days post-eclosion). (E) Western blots using protein extracts from different *jbug* mutants. Asterisks, two small bands that are still observed in *jbug*[20] larvae, which are likely from nonspecific cross-reactivity. (F) In both *59AB* and *Exel6079* homozygous embryos, the same protein bands are detected as in wild type. Note that two faint but distinct bands are visible for >300 KDa, suggesting that Jbug-PM (306.1 kDa) and the other two bigger isoforms, Jbug-PF (318.9 kDa) and Jbug-PN (322.7 kDa), are separated in this gel. (J) In *jbug*[30]/*Exel6079* transheterozygous larvae, Jbug-PF/M/N, Jbug-PI and Jbug-PL bands are all detected. (H) Protein levels of Jbug isoforms in each *jbug* mutant. Each mutant's viability is also shown.

Figure 1

774
775

Figure 2. *jbug³⁰* and *jbug¹³³* show bristle defects on the adult thorax, which are rescued by

expression of Jbug-RF. (A) Compared to wild type, *jbug³⁰* and *jbug¹³³* adult flies show distinct

thoracic bristle defects, including short bristles with rough ends and bent bristles (arrowheads).

jbug¹³³ flies have milder defects than *jbug³⁰*. (B) Phalloidin staining (red) shows the bent bristle

phenotype in *jbug³⁰* and *da>Jbug-RF* pupae (arrows) compared to the straight wild-type bristle.

(C) SEM images for bristles in wild-type and *jbug³⁰* male flies. Bristles in the red and yellow boxes

on the flies are shown in higher magnification in a-f. Compared to the nicely elongated and tapered

wild-type bristle (a), bristles in *jbug³⁰* mutants are short and distorted (c), bent (d), and rough-

ended (e). Even when the bristle looks relatively normal, it is often shorter than the wild type and

slightly twisted (compare b and f). (a'-f') Higher magnification of a-f is shown. (D-F) Whereas

overexpression of Jbug-RC (D) or Jbug-RI (E) using *69B-Gal4* or *da-Gal4* in an otherwise wild-

type animal does not cause overt defects in thoracic bristles, overexpression of Jbug-RF causes

short thick bristles (F; red arrowheads). (G-I) Although expression of Jbug-RC (G) or Jbug-RI (H)

does not rescue the bristle defects in *jbug³⁰* (red arrowheads), Jbug-RF expression almost

completely rescues the bristle defects in *jbug³⁰*, leaving only a couple of short bristles (I; cyan

arrowheads).

Figure 2

A

| WT | jbug³⁰ | jbug¹³³ | B |

C

WT   100μm

jbug³⁰   100μm

| | WT | jbug³⁰ |

| a | a' | c | c' | d | d' | e | e' |
| b | b' | f | f' |

D   *da>Jbug-RC*   *69B>Jbug-RC*

E   *da>Jbug-RI*   *69B>Jbug-RI*

F   *da>Jbug-RF*   *69B>Jbug-RF*

G   *jbug³⁰ ; da>Jbug-RC*   *jbug³⁰ ; 69B>Jbug-RC*

H   *jbug³⁰ ; da>Jbug-RI*   *jbug³⁰ ; 69B>Jbug-RI*

I   *jbug³⁰ ; da>Jbug-RF*   *jbug³⁰ ; 69B>Jbug-RF*

793
794

**Figure 3. Analysis of Jbug protein localization in the embryonic epidermis in different *jbug*
alleles.** (A, B) Confocal images for early embryos stained for Jbug (green) and phalloidin
(magenta). Jbug localizes as small apical punctate structures. (C-L) Confocal images for
epidermis in stage 13 (C, E, G, I, K) and stage 15 (D, F, H, J, L) embryos stained for Jbug (green)
and E-Cad (magenta). (C, D) In wild type, whereas high Jbug signals are observed both on the
apical surface (arrowheads) and at adherens junctions (arrows) at stage 13 (C), Jbug mostly
localizes more uniformly across the apical surface (arrowheads) at stage 15 (D). (E, F) In *jbug*[20]
mutants, junctional Jbug signals are absent, and signals on the apical surface are weak
(arrowheads in E). (G, H) In *jbug*[30] mutants, overall Jbug signals are very weak and diffused
throughout the stages. (I, J) In *jbug*[133] embryos, signals at both the apical surface (arrowheads)
and junctions (arrows) are present but are slightly weaker than wild type. (K, L) Jbug-RF
overexpression in ectodermal cells driven by *69B-Gal4* in the *jbug*[30] mutant background (K)
results in upregulation of Jbug signals in the entire apical domain with junctional enrichment at
stage 13 (arrows). Overexpression of Jbug-RF in all cells by *da-Gal4* in the *jbug*[30] mutant
background (L) causes upregulation of Jbug signals in the apical domain at stage 15 (arrowheads).
(M) A summary of Jbug protein localization in the embryonic epidermis in wild type and *jbug*
mutant alleles.

Figure 3

	WT	jbug²⁰	jbug³⁰ (homozygous stock)	jbug¹³³ (homozygous stock)
St. 13	Apical surface & Junctions	Apical surface (weak)	Weak and diffuse	Apical surface & Junctions
St. 15	Apical surface	Apical surface (weak)	Weak and diffuse	Apical surface

Figure 4. Jbug protein is upregulated in the anterior part of each segment just prior to the
formation of actin-rich denticles in the same region. (A-D) Epidermis in stages 13 (A), 14 (B),
15 (C) and 16 (D) embryos stained for Jbug (green), phalloidin (red), and E-Cad (magenta). (A)
At stage 13, Jbug is mostly uniform across epidermis. (B) At stage 14, actin begins to be
upregulated in the anterior side of each segment. Jbug also shows accumulation in the apical
domain of epidermal cells in the anterior side of the segment. (C-D) At stages 15 (C) and 16 (D),
actin-rich denticles form in the anterior side of each segment. Jbug still shows higher levels in
those regions. A and P represent the anterior and the posterior side, respectively.
822

Figure 4

Figure 5. Jbug regulates the formation of denticle precursors in the ventral epidermis of
the embryo. (A-G) Stage 16 embryos stained for E-Cad (magenta; cell boundaries) and phalloidin
(green; denticles). Ventral epidermis in wild type (A), *jbug*[20] (B), *jbug*[30] (C), *jbug*[20] ; *da-Gal4>Jbug-*
RF (D), *jbug*[30] ; *da-Gal4>Jbug-RF* (E), *da-Gal4>Jbug-RF* (F), *jbug*[30] ; *da-Gal4>Jbug-RI* (G), *da-*
Gal4>Jbug-RI (H), *da-Gal4>jbug RNAi* (*GD13033*) (I) and *matα-Gal4>jbug RNAi* (*JF01166*) (J) is
shown. (A) Epidermal cells at the denticle belt in the abdominal segment 5 (yellow box) are shown
at higher magnification. Dotted yellow line, ventral midline. (B-J) Epidermal cells of the same
region as in A are shown. For all the embryos, anterior is to the left. (B, J) *jbug*[20] mutants (B) and
matα-Gal4>jbug RNAi embryos (J) show mild defects in the embryo morphology, such as wavy
epidermis (areas with dotted lines; cells are out of focus). Occasionally, cells either fail to form
denticles (arrow in J) or form very short denticles (arrowheads in B, I and J). (K) A cartoon for
quantification of the denticle numbers. Four cells that are closest to the ventral midline were
chosen in each of the six columns of cells in denticle belts in abdominal segments 4, 5 and 6
(column 1, most anterior). (L) Percentage of cells that have different numbers of denticles in C1
and C6. See Supplemental Figure 4 for quantification for C2-C5. (M) Quantification of the average
number of denticles per cell in C1 and C6. Student's t-test with Welch's correction is used for
statistics calculation. (O) A summary of denticle defects in *jbug* mutant alleles and RNAi
knockdown and rescue results with Jbug-RF and -RI.
841
842

Figure 5

843
844

Figure 6. Jbug protein is apically enriched in the embryonic trachea and the salivary gland,

and this enrichment is absent in *jbug²⁰* and *jbug³⁰* mutants. (A-D) Confocal images of stage

14 embryonic trachea stained for Jbug (green) and E-Cad (magenta). (A) In wild type, weak Jbug

signals are observed in all tracheal cells, with enrichment in the apical domain (arrowheads). (B)

In *jbug²⁰* embryos, Jbug signals in the trachea are mostly absent (arrows). (C) In *jbug³⁰* mutant

embryos, Jbug signals are diffused in the cytoplasm in tracheal cells (asterisks), with no apical

enrichment (arrows; compare to E-Cad signals (arrowheads) in the apical domain). (D) In *jbug¹³³*,

apical enrichment of Jbug is still present (arrowheads). (E-H) Stage 14 salivary glands stained for

Jbug (green) and E-Cad (magenta). (E) Similar to tracheal signals, Jbug is enriched in the apical

domain of wild-type salivary gland cells (arrowheads) with weak signals in the cytoplasm. (F) In

jbug²⁰, overall Jbug signals are significantly decreased in the salivary gland and apical enrichment

is mostly gone (arrow; compare to E-Cad signals (arrowhead) in the apical domain). (G) In *jbug³⁰*,

Jbug is not apically enriched in the salivary gland (arrow; compare to E-Cad signals (arrowhead)

in the apical domain) but diffused in the cytoplasm. (H) In *jbug¹³³*, apical enrichment of Jbug is still

present (arrowheads). (I) A summary of Jbug protein localization in the trachea and the salivary

gland in wild type and *jbug* mutant alleles.

Figure 7. Jbug overexpression in the salivary gland causes mislocalization of Crb along

the apical surface and an expanded and irregular apical domain. (A-D) stage 16 salivary

glands stained for Crb (green), Jbug (magenta), GFP (magenta in C) and salivary gland nuclear

markers CrebA (A-C and F) and Sage (D and E). Salivary glands overexpressing Tmem-GFP

(control, A), Jbug-RC (B), Jbug-RC-GFP (C), Jbug-RI (D), Jbug-RL (E) and Jbug-RF (F, G) are

shown. Note that Jbug mainly localizes to the cytoplasm and only shows weak apical signals in

stage 16 salivary glands (red arrow in A; compare to Figure 4E). Jbug-RC (B) and Jbug-RC-GFP

(C) localize both to salivary gland nuclei and the cytoplasm when overexpressed. Jbug

overexpression causes an expanded and irregular salivary gland lumen (B-G). Irregular salivary

gland lumen is worsened when stronger UAS lines are used (compare B and C; also compare F

and G). Compared to subapical Crb signals along AJs in control salivary gland cells (inset in A;

yellow arrows), Crb signals are dispersed along the entire apical domain in Jbug-overexpressing

salivary glands (insets in B-G). (H) Quantification of the irregular apical membrane morphology in

salivary glands overexpressing different Jbug isoforms. Strong UAS lines shown in B and G were

used to overexpress Jbug-RC and Jbug-RF. For mild irregularity, salivary glands with the uneven

apical domain were counted. For severe irregularity, salivary glands with bulging and constriction

phenotypes were counted.

Figure 7

Figure 8. Jbug is required for proper Crb levels in epithelial cells during embryogenesis.

(A-H) Confocal images of stage 14/15 embryos of control (*da-Gal4/+*; A and E), zygotic (*da>jbug*

RNAi; B and F), maternal (*matα>jbug RNAi*; C and G), and both maternal and zygotic knockdown

of *jbug* (*matα, da>jbug RNAi*; D and H). Anterior is to the left. Ventral (A, D, E), dorsal (F) and

lateral (B, C, G and H) views are shown. (A-D) Embryos stained for E-Cad (magenta). (E-H)

Embryos stained for Crb (green) and Sage (a SG marker; magenta). (E'-H''') Higher magnification

of the boxed regions in E-H. Crb (green) and Jbug (magenta) are shown. Compared to strong

apical surface Jbug signals in epidermal cells in control (E'') and embryos knocked down for *jbug*

only zygotically (F'') or maternally (G''), Jbug signals are absent in the apical surface in many

epidermal cells in *jbug* M/Z knockdown embryos (yellow arrowheads in H' and H''). In those cells,

Crb is also absent (yellow arrows in H'''). Compare to neighboring cells with significant apical

surface Jbug signals (cyan arrowheads in H' and H'') and relatively normal Crb localization (cyan

arrows in H'''). Asterisks, residual or nonspecific Jbug signals. (I-M) Confocal images of stage 14

salivary glands stained for Crb (green) and Jbug (magenta). White dotted lines, salivary gland

boundaries. Control (*fkh-Gal4/+*; I), zygotic (*fkh>jbug RNAi*; J), maternal (*matα>jbug RNAi*; K),

and both maternal and zygotic knockdown of *jbug* (*matα, fkh>jbug RNAi*; two representative

images are shown in L and M). (I'-M'') Jbug (I'-M') and Crb (I''-M'') only. Jbug signals are reduced

in *jbug* knockdown (cyan arrowheads in J'-M'), with a more significant reduction in maternal (K')

and maternal/zygotic knockdown (K'-M'). (I'''-M''') Higher magnification of the boxed regions in I'-

M''. Maternal and maternal/zygotic knockdown of *jbug* results in gaps (yellow arrowheads in K'''

and M''') or mislocalized Crb signals to the apical surface (red arrowheads in L''').

Figure 8

905   **Figure 9. Isoform-specific roles and features of Jbug.**

Jbug isoforms	Roles/specific features during development
Jbug-PC/G/H/J/K	Pupal stage-specific, Potential role in the nucleus
Jbug-PI	Embryonic denticle formation (numbers)
Jbug-PL	Oogenesis, Epithelial morphogenesis during embryogenesis
Jbug-PF/M/N	Thoracic bristle formation, Embryonic denticle formation (shape/size)

Calponin-homology domain   Filamin-type immunoglobulin domain

907 **Supplemental Figure 1. In situ hybridization.** In situ hybridization of wild-type embryos using
908 probes specific for different *jbug* splice forms as shown in Figure 1A. CL, clypeolabrum. SG,
909 salivary gland. tp, tracheal pits. PS, posterior spiracles.
910

Supplemental Figure 2. Reverse transcriptase-PCR (RT-PCR) reveals a loss of *jbug* mRNA

in *jbug*[20] null mutants. (A) Genomic structures for the jbug locus. Blue and red arrows represent

primers used for RT-PCR. (B and C) RT-PCR using cDNAs made from wild-type (OR) and *jbug*[20]

homozygous embryos. Whereas RT-PCR using wild-type cDNA produces a PCR product of

expected sizes (left lanes in B and C), no bands are detected in *jbug*[20] (right lanes in B and C).

Supplemental Figure 3. Mutations in *jbug* cause mild planar cell polarity defects in adult

wing hair orientation. (A-C) Adult wings. Compared to the largely distal orientation of wing hairs

in the wild-type wing (A), wings in *jbug^{30}* mutants (B) show mild swirling patterns of wing hairs. (C)

jbug^{30}/Exel6079 flies do not show defects in the wing hair orientation. White boxed regions are

shown in higher magnification in the bottom. (D, E) Wild-type (D) and *jbug^{30}* mutant (E) pupal

wings (32 hours APF) stained for phalloidin. Prehair formation is delayed in the pupal wing in

jbug^{30}. (F, G) Adult ommatidia of wild type (F) and *jbug^{30}* (G) near the dorsal/ventral boundary,

the equator. Schematic drawings are shown in the panels below the actual images. Green and

magenta shapes indicate the orientation of ommatidia normally found in the dorsal and ventral

hemisphere of the eye, respectively. *jbug^{30}* flies do not show planar cell polarity defects in the

orientation of ommatidia.

Supplemental Figure 3

 Supplemental Figure 4. Jbug regulates the formation of denticle precursors in the ventral

epidermis of the embryo. (A) Percentage of cells that have different numbers of denticles in

rows C2-C5. (B) Quantification of the average number of denticles per cell in C2-C5. Student's t-

test with Welch's correction is used for statistics calculation. The same four cells as shown in

Figure 5K were chosen for quantification for each genotype.

Supplemental Figure 5. Epithelial polarity markers and F-actin and are not affected in

salivary glands in *jbug* mutants. (A-I) Confocal images of stage 16 salivary glands. (A-C) Wild-

type (A), *jbug*[20] (B) and *jbug*[30] (C) salivary glands stained for Crb (green) and E-Cad (magenta).

(D-F) Wild-type (E), *jbug*[20] (F) and *jbug*[30] (G) salivary glands stained for phalloidin (green) and

Dlg (magenta). (G-I) Wild-type (G), *jbug*[20] (H) and *jbug*[30] (I) salivary glands stained for α-Spec.

Insets, higher magnification of the boxed region.

Supplemental Figure 6. F-actin, epithelial polarity markers and junction markers are not
significantly altered in salivary glands overexpressing different Jbug isoforms. (A-J)
Confocal images of stage 16 control (*fkh-Gal4/+*, A and F) and salivary glands overexpressing
Jbug-RC (B and G), Jbug-RI (C and H), Jbug-RL (D and I) and Jbug-RF (E and J). (A-E) Salivary
gland stained for phalloidin (green), α-Spec (magenta) and a salivary gland nuclear marker CrebA
(blue). (F-J) Salivary glands stained for Nrx-IV (green) and E-Cad (magenta). Insets, higher
magnification of the boxed region.
951

Supplemental Figure 6

Supplemental Figure 7. (A-D) Confocal images of stage 11 embryos stained for Crb (green),

and Sage (blue). Control (*fkh-Gal4/+*; A), zygotic (*fkh>jbug RNAi*; B), maternal (*matα>jbug RNAi*;

C), and both maternal and zygotic knockdown of *jbug* (*matα, fkh>jbug RNAi*; D). (A'-D''') Higher

magnification of the salivary glands in A-D. Crb (green), Jbug (magenta) and Sage (blue) signals

are shown. (A''-D''') Crb (A''-D'') and Jbug (A'''-D''') only. Jbug signals are slightly reduced in *jbug*

knockdown, with a more significant reduction in maternal (C''') and maternal/zygotic knockdown

(D'''). (A''''-D'''') Higher magnification of the boxed region in A''-D''. Maternal/zygotic knockdown

of *jbug* results in gaps (yellow arrowheads) and mislocalized Crb signals to the apical surface (red

arrowheads).

Acknowledgments

We thank the members of the Chung and the Andrew laboratories for their comments and suggestions. We thank M. Mlodzik and the Bloomington stock center for fly stocks and the Developmental Studies Hybridoma Bank for antibodies. We thank Flybase for the gene information. This work is supported by start-up fund from Louisiana State University to S.C. and NIH R01 DE013899 to D.J.A.

References

Adler P. N., 2012 The frizzled/stan pathway and planar cell polarity in the Drosophila wing. Curr.

Top. Dev. Biol. 101: 1–31. https://doi.org/10.1016/b978-0-12-394592-1.00001-6

Baumgartner S., J. T. Littleton, K. Broadie, M. A. Bhat, R. Harbecke, *et al.*, 1996 A Drosophila

neurexin is required for septate junction and blood-nerve barrier formation and function.

Cell 87: 1059–1068. https://doi.org/10.1016/S0092-8674(00)81800-0

Bedolla R. G., Y. Wang, A. Asuncion, K. Chamie, S. Siddiqui, *et al.*, 2009 Nuclear versus

cytoplasmic localization of filamin a in prostate cancer: Immunohistochemical correlation

with metastases. Clin. Cancer Res. 15: 788–796. https://doi.org/10.1158/1078-0432.CCR-

08-1402

Bejsovec A., 2013 Wingless/Wnt signaling in Drosophila: The pattern and the pathway. Mol.

Reprod. Dev. 80: 882–894. https://doi.org/10.1002/mrd.22228

Bellanger J. M., C. Astier, C. Sardet, Y. Ohta, T. P. Stossel, *et al.*, 2000 The Rac1- and RhoG-

specific GEF domain of trio targets filamin to remodel cytoskeletal actin. Nat. Cell Biol. 2:

888–892. https://doi.org/10.1038/35046533

Blanchoin L., K. J. Amann, H. N. Higgs, J. Marchand, D. A. Kaiser, *et al.*, 2000 Direct

observation of dendritic actin ®lamentnetworksnucleatedbyArp2/3 complex and

WASP/Scar proteins. Nature 404: 1007–1011. https://doi.org/10.1038/35010008

Browne K. A., R. W. Johnstone, D. A. Jans, and J. A. Trapani, 2000 Filamin (280-kDa actin-

binding protein) is a caspase substrate and is also cleaved directly by the cytotoxic T

lymphocyte protease granzyme B during apoptosis. J. Biol. Chem. 275: 39262–39266.

https://doi.org/10.1074/jbc.C000622200

Chung S. Y., C. Chavez, and D. J. Andrew, 2011 Tracheales s (Trh) regulates all tracheal genes

during Drosophila embryogenesis. Dev. Biol. 360: 160–172.

https://doi.org/10.1016/j.ydbio.2011.09.014

Chung S., and D. J. Andrew, 2014 Cadherin 99C regulates apical expansion and cell

rearrangement during epithelial tube elongation. Development 141: 1950–60.

https://doi.org/10.1242/dev.104166

Cunningham C. C., J. B. Gorlin, D. J. Kwiatkowski, J. H. Hartwig, A. Janmey, *et al.*, 1992 Actin-

Binding Protein Requirement for Cortical Stability and Efficient Locomotion Published by :

American Association for the Advancement of Science Stable URL :

http://www.jstor.org/stable/2876160. Science (80-.). 255: 325–327.

https://doi.org/10.1126/science.1549777

Dilks S. A., and S. DiNardo, 2010 Non-cell-autonomous control of denticle diversity in the

Drosophila embryo. Development 137: 1395–1404. https://doi.org/10.1242/dev.045450

Flier A. Van der, and A. Sonnenberg, 2001 Structural and functional aspects of filamins. Biochim. Biophys. Acta - Mol. Cell Res. 1538: 99–117. https://doi.org/10.1016/S0167-4889(01)00072-6

Fox R. M., A. Vaishnavi, R. Maruyama, and D. J. Andrew, 2013 Organ-specific gene expression: The bHLH protein Sage provides tissue specificity to Drosophila FoxA. Dev. 140: 2160–2171. https://doi.org/10.1242/dev.092924

Frank D. J., R. Hopmann, M. Lenartowska, and K. G. Miller, 2006 Capping Protein and the Arp2/3 Complex Regulate Nonbundle Actin Filament Assembly to Indirectly Control Actin Bundle Positioning during Drosophila melanogaster Bristle Development. Mol. Biol. Cell 17: 3930–3939. https://doi.org/10.1091/mbc.E06-06-0500

Gao J. L., Y. J. Fan, X. Y. Wang, Y. Zhang, J. Pu, *et al.*, 2015 A conserved intronic U1 snRNP-binding sequence promotes trans-splicing in Drosophila. Genes Dev. 29: 760–771. https://doi.org/10.1101/gad.258863.115

Gay O., B. Gilquin, F. Nakamura, Z. A. Jenkins, R. McCartney, *et al.*, 2011 RefilinB (FAM101B) targets FilaminA to organize perinuclear actin networks and regulates nuclear shape. Proc. Natl. Acad. Sci. U. S. A. 108: 11464–11469. https://doi.org/10.1073/pnas.1104211108

Gehler S., M. Baldassarre, Y. Lad, J. L. Leight, M. A. Wozniak, *et al.*, 2009 Filamin A–$_N$L1 Integrin Complex Tunes Epithelial Cell Response to Matrix Tension. Mol. Biol. Cell 20: 3224–3238. https://doi.org/10.1091/mbc.E08

Gong W. J., and K. G. Golic, 2003 Ends-out, or replacement, gene targeting in Drosophila. Proc. Natl. Acad. Sci. U. S. A. 100: 2556–2561. https://doi.org/10.1073/pnas.0535280100

Gorlin J. B., R. Yamin, S. Egan, M. Stewart, T. P. Stossel, *et al.*, 1990 Human endothelial actin-binding protein (ABP-280, nonmuscle filamin): A molecular leaf spring. J. Cell Biol. 111: 1089–1105. https://doi.org/10.1083/jcb.111.3.1089

Häcker U., and N. Perrimon, 1998 DRhoGEF2 encodes a member of the Dbl family of oncogenes and controls cell shape changes during gastrulation in Drosophila. Genes Dev. 12: 274–284. https://doi.org/10.1101/gad.12.2.274

Henderson K. D., and D. J. Andrew, 2000 Regulation and function of Scr, exd, and hth in the Drosophila salivary gland. Dev. Biol. 217: 362–374. https://doi.org/10.1006/dbio.1999.9560

Horiuchi T., and T. Aigaki, 2006 Alternative trans-splicing: a novel mode of pre-mRNA processing. Biol. Cell 98: 135–140. https://doi.org/10.1042/bc20050002

Hudson A. M., and L. Cooley, 2002 A subset of dynamic actin rearrangements in Drosophila requires the Arp2/3 complex. J. Cell Biol. 156: 677–687.

https://doi.org/10.1083/jcb.200109065

Huelsmann S., N. Rintanen, R. Sethi, N. H. Brown, and J. Ylänne, 2016 Evidence for the

mechanosensor function of filamin in tissue development. Sci. Rep. 6: 2–9.

https://doi.org/10.1038/srep32798

Jeon Y. J., J. S. Choi, J. Y. Lee, K. R. Yu, S. H. Ka, *et al.*, 2008 Filamin B Serves as a Molecular

Scaffold for Type I Interferon-induced c-Jun NH2-terminal Kinase Signaling Pathway. Mol.

Biol. Cell 19: 5116–5130. https://doi.org/10.1091/mbc.e08-06-0576

King T. D., J. E. Johnson, and J. R. Bateman, 2019 Position Effects In fl uence Transvection in.

213: 1289–1299. https://doi.org/https://doi.org/10.1534/genetics.119.302583

Krueger D., T. Quinkler, S. A. Mortensen, C. Sachse, and S. de Renzis, 2019 Cross-linker–

mediated regulation of actin network organization controls tissue morphogenesis. J. Cell

Biol. 218: 2743–2761. https://doi.org/10.1083/JCB.201811127

Külshammer E., and M. Uhlirova, 2013 The actin cross-linker filamin/cheerio mediates tumor

malignancy downstream of JNK signaling. J. Cell Sci. 126: 927–938.

https://doi.org/10.1242/jcs.114462

Lasda E. L., and T. Blumenthal, 2011 Trans-splicing. Wiley Interdiscip. Rev. RNA 2: 417–434.

https://doi.org/10.1002/wrna.71

Lee G. Y., and T. L. Schwarz, 2016 Filamin, a synaptic organizer in Drosophila, determines

glutamate receptor composition and membrane growth. Elife 5: 1–29.

https://doi.org/10.7554/eLife.19991

Lehmann R., and D. Tautz, 1994 In Situ Hybridization to RNA. Methods Cell Biol. 44: 575–598.

https://doi.org/doi: 10.1016/s0091-679x(08)60933-4

Li M., M. Serr, K. Edwards, S. Ludmann, D. Yamamoto, *et al.*, 1999 during Drosophila

Oogenesis. Cell 146: 1061–1073.

Loy C. J., K. S. Sim, and E. L. Yong, 2003 Filamin-A fragment localizes to the nucleus to

regulate androgen receptor and coactivator functions. Proc. Natl. Acad. Sci. U. S. A. 100:

4562–4567. https://doi.org/10.1073/pnas.0736237100

Mammoto A., S. Huang, and D. E. Ingber, 2007 Filamin links cell shape and cytoskeletal

structure to Rho regulation by controlling accumulation of p190RhoGAP in lipid rafts. J. Cell

Sci. 120: 456–467. https://doi.org/10.1242/jcs.03353

Manieu C., G. H. Olivares, F. Vega-Macaya, M. Valdivia, and P. Olguín, 2018 Jitterbug/Filamin

and Myosin-II form a complex in tendon cells required to maintain epithelial shape and

polarity during musculoskeletal system development. Mech. Dev. 154: 309–314.

https://doi.org/10.1016/j.mod.2018.09.002

Mongelard F., M. Labrador, E. M. Baxter, T. I. Gerasimova, and V. G. Corces, 2002 Trans-
splicing as a novel mechanism to explain interallelic complementation in Drosophila.
Genetics 160: 1481–1487.

Mullins R. D., J. A. Heuser, and T. D. Pollard, 1998 The interaction of Arp2/3 complex with actin:
Nucleation, high affinity pointed end capping, and formation of branching networks of
filaments. Proc. Natl. Acad. Sci. U. S. A. 95: 6181–6186.
https://doi.org/10.1073/pnas.95.11.6181

Nakamura F., O. Heikkinen, O. T. Pentikäinen, T. M. Osborn, K. E. Kasza, *et al.*, 2009
Molecular basis of filamin A-FilGAP interaction and its impairment in congenital disorders
associated with Filamin A mutations. PLoS One 4.
https://doi.org/10.1371/journal.pone.0004928

Nakamura F., T. P. Stossel, and J. H. Hartwig, 2011 The filamins: Organizers of cell structure
and function. Cell Adh. Migr. 5: 160–169. https://doi.org/10.4161/cam.5.2.14401

Niederman R., P. C. Amrein, and J. Hartwig, 1983 Three-dimensional structure of actin
filaments and of an actin gel made with actin-binding protein. J. Cell Biol. 96: 1400–1413.
https://doi.org/10.1083/jcb.96.5.1400

Ohsako T., T. Horiuchi, T. Matsuo, S. Komaya, and T. Aigaki, 2003 Drosophila lola encodes a
family of BTB-transcription regulators with highly variable C-terminal domains containing
zinc finger motifs. Gene 311: 59–69. https://doi.org/10.1016/S0378-1119(03)00554-7

Ohta Y., and J. H. Hartwig, 1996 Phosphorylation of actin-binding protein 280 by growth factors
is mediated by p90 ribosomal protein S6 kinase. J. Biol. Chem. 271: 11858–11864.
https://doi.org/10.1074/jbc.271.20.11858

Ohta Y., N. Suzuki, S. Nakamura, J. H. Hartwig, and T. P. Stossel, 1999 The small GTPase
RalA targets filamin to induce filopodia. Proc. Natl. Acad. Sci. U. S. A. 96: 2122–2128.
https://doi.org/10.1073/pnas.96.5.2122

Ohta Y., J. H. Hartwig, and T. P. Stossel, 2006 FilGAP, a Rho- and ROCK-regulated GAP for
Rac binds filamin A to control actin remodelling. Nat. Cell Biol. 8: 803–814.
https://doi.org/10.1038/ncb1437

Olguín P., A. Glavic, and M. Mlodzik, 2011 Intertissue mechanical stress affects Frizzled-
mediated planar cell polarity in the Drosophila notum epidermis. Curr. Biol. 21: 236–42.
https://doi.org/10.1016/j.cub.2011.01.001

Oliva C., C. Molina-Fernandez, M. Maureira, N. Candia, E. L??pez, *et al.*, 2015 Hindsight
regulates photoreceptor axon targeting through transcriptional control of jitterbug/Filamin
and multiple genes involved in axon guidance in Drosophila. Dev. Neurobiol. 75: 1018–

1032. https://doi.org/10.1002/dneu.22271

Pi M., R. F. Spurney, Q. Tu, T. Hinson, and L. D. Quarles, 2002 Calcium-sensing receptor

activation of Rho involves filamin and rho-guanine nucleotide exchange factor.

Endocrinology 143: 3830–3838. https://doi.org/10.1210/en.2002-220240

Razinia Z., T. Mäkelä, J. Ylänne, and D. a Calderwood, 2012 Filamins in mechanosensing and

signaling. Annu. Rev. Biophys. 41: 227–46. https://doi.org/10.1146/annurev-biophys-

050511-102252

Rorth Pe., 1996 A modular misexpression screen in Drosophila detecting tissue-specific

phenotypes. Proc. Natl. Acad. Sci. 93: 12418–12422.

Sasaki E., A. T. Byrne, E. Phelan, D. W. Cox, and W. Reardon, 2019 A review of filamin A

mutations and associated interstitial lung disease. Eur. J. Pediatr. 178: 121–129.

https://doi.org/10.1007/s00431-018-3301-0

Savoy R. M., and P. M. Ghosh, 2013 The dual role of filamin A in cancer: Can't live with (too

much of) it, can't live without it. Endocr. Relat. Cancer 20: 341–356.

https://doi.org/10.1530/ERC-13-0364

Shao Q. Q., T. P. Zhang, W. J. Zhao, Z. W. Liu, L. You, *et al.*, 2016 Filamin A: Insights into its

Exact Role in Cancers. Pathol. Oncol. Res. 22: 245–252. https://doi.org/10.1007/s12253-

015-9980-1

Sokol N. S., and L. Cooley, 1999 Drosophila Filamin encoded by the cheerio locus is a

component of ovarian ring canals. Curr. Biol. 9: 1221–1230. https://doi.org/10.1016/S0960-

9822(99)80502-8

Sokol N. S., and L. Cooley, 2003 Drosophila filamin is required for follicle cell motility during

oogenesis. Dev. Biol. 260: 260–272. https://doi.org/10.1016/S0012-1606(03)00248-3

Song J., and M. A. Tanouye, 2006 Seizure suppression by shakB2, a gap junction mutation in

Drosophila. J. Neurophysiol. 95: 627–635. https://doi.org/10.1152/jn.01059.2004

Tepass U., 1996 Crumbs, a component of the apical membrane, is required for zonula adherens

formation in primary epithelia of Drosophila. Dev. Biol. 177: 217–225.

https://doi.org/10.1006/dbio.1996.0157

Thurmond J., J. L. Goodman, V. B. Strelets, H. Attrill, L. S. Gramates, *et al.*, 2019 FlyBase 2.0:

The next generation. Nucleic Acids Res. 47: D759–D765.

https://doi.org/10.1093/nar/gky1003

Tikhonov M., M. Utkina, O. Maksimenko, and P. Georgiev, 2018 Conserved sequences in the

Drosophila mod(mdg4) intron promote poly(A)-independent transcription termination and

trans-splicing. Nucleic Acids Res. 46: 10608–10618. https://doi.org/10.1093/nar/gky716

Tirupula K. C., S. S. Ithychanda, M. L. Mohan, S. V. Naga Prasad, J. Qin, *et al.*, 2015 G protein-
coupled receptors directly bind filamin A with high affinity and promote filamin
phosphorylation. Biochemistry 54: 6673–6683.
https://doi.org/10.1021/acs.biochem.5b00975

Ueda K., Y. Ohta, and H. Hosoya, 2003 The carboxy-terminal pleckstrin homology domain of
ROCK interacts with filamin-A. Biochem. Biophys. Res. Commun. 301: 886–890.
https://doi.org/10.1016/S0006-291X(03)00048-2

Valle-Pérez B. Del, V. G. Martínez, C. Lacasa-Salavert, A. Figueras, S. S. Shapiro, *et al.*, 2010
Filamin B plays a key role in vascular endothelial growth factor-induced endothelial cell
motility through its interaction with Rac-1 and Vav-2. J. Biol. Chem. 285: 10748–10760.
https://doi.org/10.1074/jbc.M109.062984

Wang Y., J. I. Kreisberg, R. G. Bedolla, M. Mikhailova, R. W. Devere White, *et al.*, 2007 A 90
kDa fragment of filamin A promotes Casodex-induced growth inhibition in Casodex-
resistant androgen receptor positive C4-2 prostate cancer cells. Oncogene 26: 6061–6070.
https://doi.org/10.1038/sj.onc.1210435

Wang Q., W. Zheng, Z. Wang, J. W. Yang, S. Hussein, *et al.*, 2015 Filamin-A increases the
stability and plasma membrane expression of polycystin-2. PLoS One 10: 1–19.
https://doi.org/10.1371/journal.pone.0123018

Wodarz A., U. Hinz, M. Engelbert, and E. Knust, 1995 Expression of crumbs confers apical
character on plasma membrane domains of ectodermal epithelia of drosophila. Cell 82:
67–76. https://doi.org/10.1016/0092-8674(95)90053-5

Wolosewick J. J., and J. Condeelis, 1986 Fine structure of gels prepared from an actin-binding
protein and actin: Comparison to cytoplasmic extracts and cortical cytoplasm in amoeboid
cells of Dictyostelium discoideum. J. Cell. Biochem. 30: 227–243.
https://doi.org/10.1002/jcb.240300305

Yue J., S. Huhn, and Z. Shen, 2013 Complex roles of filamin-A mediated cytoskeleton network
in cancer progression. Cell Biosci. 3: 1. https://doi.org/10.1186/2045-3701-3-7

1166