***Toxoplasma gondii* GRA28 is required for placenta-specific induction of the**

regulatory chemokine CCL22 in human and mouse.

Elizabeth N. Rudzki[1], Stephanie E. Ander[1,2], Rachel S. Coombs[1], Hisham S. Alrubaye[1], Leah F.

Cabo[1], Matthew L. Blank[1], Nicolas Gutierrez-Melo[1], JP Dubey[3], Carolyn B. Coyne[4] and Jon P.

Boyle[1]*

[1]Department of Biological Sciences, Dietrich School of Arts and Sciences, University of

Pittsburgh, Pittsburgh, PA. 15260.

[2]Department of Pediatrics, University of Pittsburgh School of Medicine, Pittsburgh, PA. 15260

[3]United States Department of Agriculture, Agricultural Research Service, Beltsville Agricultural

Research Center, Animal Parasitic Diseases Laboratory, Beltsville, Maryland, 20705.

[4]Department of Molecular Genetics and Microbiology, Duke University School of Medicine,

Durham, NC. 27708.

*Author to whom correspondence should be addressed:

Jon Patrick Boyle, PhD

Associate Professor

Department of Biological Sciences

Dietrich School of Arts and Sciences

University of Pittsburgh

4249 Fifth Avenue

Pittsburgh, PA. 15260

Email: boylej@pitt.edu

ABSTRACT

Toxoplasma gondii is an intracellular protozoan pathogen of humans that can cross the placenta

and result in adverse pregnancy outcomes and long-term birth defects. The mechanism used by

T. gondii to cross the placenta are unknown but complex interactions with the host immune

response are likely to play a role in dictating infection outcomes during pregnancy. Prior work

showed that *T. gondii* infection dramatically and specifically increases the secretion of the

immunomodulatory chemokine CCL22 in human placental cells during infection.. Given the

important role of this chemokine during pregnancy, we hypothesized that CCL22 induction was

driven by a specific *T. gondii*-secreted effector. Using a combination of bioinformatics and

molecular genetics, we have now identified *T. gondii* GRA28 as the gene product required for

CCL22 induction. GRA28 is secreted into the host cell where it localizes to the nucleus, and

deletion of this gene results in reduced CCL22 placental cells as well as a human monocyte cell

line. The impact of GRA28 on CCL22 production is also conserved in mouse immune and

placental cells both *in vitro* and *in vivo*. Moreover, parasites lacking GRA28 are impaired in their

ability to disseminate throughout the animal, suggesting a link between CCL22 induction and

the ability of the parasite to cause disease. Overall these data demonstrate a clear function for

GRA28 in altering the immunomodulatory landscape during infection of both placental and

peripheral immune cells, and show a clear impact of this immunomodulation on infection

outcome.

AUTHOR SUMMARY

Toxoplasma gondii is a globally ubiquitous pathogen that can cause severe disease in

HIV/AIDS patients and can also cross the placenta and infect the developing fetus. We have

found that placental and immune cells infected with *T. gondii* secrete significant amounts of a

chemokine (called "CCL22") that is critical for immune tolerance during pregnancy. In order to

better understand whether this is a response by the host or a process that is driven by the

parasite, we have identified a *T. gondii* gene that is absolutely required to induce CCL22

production in human cells, indicating that CCL22 production is a process driven almost entirely

by the parasite rather than the host. Consistent with its role in immune tolerance, we also found

that *T. gondii* parasites lacking this gene are less able to proliferate and disseminate throughout

the host. Taken together these data illustrate a direct relationship between CCL22 levels in the

infected host and a key parasite effector, and provide an interesting example of how *T. gondii*

can directly modulate host signaling pathways in order to facilitate its growth and dissemination.

INTRODUCTION

Toxoplasma gondii is an obligate intracellular parasite that is an important parasite of

humans and other animals. While this pathogen is particularly well-known to cause severe

disease in the immunocompromised, such as those with HIV/AIDS or undergoing

immunosuppression for organ transplants, *T. gondii* is also capable of crossing the placenta and

infecting the developing fetus, leading to a variety of infection outcomes ranging from

asymptomatic to severe (1). Importantly, even children born without symptoms are at high risk

for extensive health problems later in life, including ocular disease and neurological disorders

(2, 3). To date little is known about how *T. gondii* gains access to the fetal compartment and

how the host responds to the presence of parasites at the maternal-fetal interface.

Recently we (4) found that primary human trophoblast cells (derived from term

placentas) and 2nd trimester placental explants produced the chemokine CCL22 in response to

infection with *T. gondii* (4). Production of this chemokine was dependent on parasite invasion

and the dense granule effector trafficking gene product MYR1 (4). While the role of CCL22

during infection with *T. gondii* is poorly understood, this chemokine is a key molecular signal for

the recruitment of regulatory T cells which are well known for their role in suppressing immune

responses to tumors, leading to poor clinical outcomes (5, 6). Importantly disruption of T_{reg}

recruitment to tumors can lead to improved outcomes in animal models. For example, using

Ccl22 DNA vaccines in mice leads to misdirection of regulatory T cells and ultimately reduced

tumor growth (5). The role for CCL22 in healthy humans is less well understood, although it is

thought to subvert and/or modulate inflammatory responses and may be particularly important

for response resolution after pathogen clearance. CCL22 and regulatory T cells also play a

critical role during pregnancy, where they seem to govern immune tolerance (7) and regulation

of inflammation at the maternal-fetal interface. This regulatory role appears to be critical in

determining pregnancy outcome during pathogen-mediated immune activation (7, 8). Given the

important role played by CCL22 during pregnancy and our recent findings regarding the ability

of a congenitally acquired parasite to directly modulate production of this chemokine, we sought

to identify the parasite effector(s) responsible for this in order to determine the impact of CCL22

modulation on congenital transmission and pregnancy outcome during vertical transmission. To

do this we used a bioinformatic screen identify candidate genes and identified one

(TGGT1_231960) as being required for CCL22 induction in human and mouse cells. Overall

these data show that a specific effector is largely responsible for *T. gondii*-mediated CCL22

induction in a relatively small number of human and mouse cell types, and suggest that the

manipulation of CCL22 levels by GRA28 may influence the ability of *T. gondii* to disseminate to

throughout the host.

RESULTS

***Toxoplasma gondii* induces a monocyte-like cell line to produce the**

immunomodulatory chemokine CCL22.

Previous work established that placental explants and primary human trophoblasts infected with

T. gondii had increased *CCL22* transcript abundance and released more CCL22 protein into the

culture media compared to mock infected controls (4). Since we also found that not all cell types

produce CCL22 in response to infection (e.g., HFFs), we were interested in identifying a human

cell line that could be used as a more tractable model than placental cells to assay *T. gondii*-

driven CCL22 induction. THP-1 cells, a cell line derived from a patient with monocytic leukemia,

were a reasonable candidate given their origins in the myeloid lineage and known production of

CCL22 in response to a variety of stimuli (9, 10). We infected THP-1 cells with a type 1 *T. gondii*

strain (RH88 or RH:YFP;(4)) at a multiplicity of infection (MOI) of 3. Following 24 hours of

infection, supernatants were collected from each well. Human foreskin fibroblasts (HFFs) were

infected in parallel as negative controls. Mock treatments involved passing the parasite solution

through a 0.22 µM filter prior to exposure to the cells. Based on CCL22 ELISA, *T. gondii*

infection induced CCL22 in THP-1 cells, and as expected there was no CCL22 production from

mock-treated controls or *T. gondii*-infected HFFs (**Figure 1a**). We also infected primary

placental tissues in the same manner, and as expected villous tree explants and decidua taken

from 2nd trimester placentas produced significantly more CCL22 compared to mock-treated

controls (**Figure 1a**). In addition to the type 1 RH strain, other *T. gondii* strain types (Type

2:PRU and Type 3:CEP; Supplementary **Figure 1 Supplement 1a**) also induced secretion of

CCL22 from THP-1 cells, as did the nearest extant relative of *T. gondii*, *Hammondia hammondi*

(**Figure 1 Supplement 1b**). In contrast to *H. hammondi*, and just as we observed previously in

primary human placental cells (4), *Neospora caninum* has no effect on THP-1 production of

CCL22 (Supplementary **Figure 1 Supplement 1b**). These data provided strong support that the

mechanism of CCL22 induction was the same for THP-1 and placental cells.

We also determined if live parasites were required to induce CCL22 in THP-1 cells by

exposing host cells to parasites that were exposed to a variety of lethal treatments. As shown in

Figure 1b dead parasites failed to induce CCL22 production by THP-1 cells. We also pretreated

parasites and host cells with 10 µg/mL Cytochalasin-D (Cyt-D) to block invasion (11), and as

shown in **Figure 1c** Cyt-D treated parasites were significantly impaired in their ability to induce

CCL22, suggesting that active invasion was required for this phenomenon. We obtained similar

results with the inhibitor 4-BPB (**Figure 1d**), which also significantly blocked CCL22 production

by THP-1 cells at 0.5 and 1 µM. This drug blocks rhoptry and dense granule secretion from *T.*

gondii, but not microneme secretion(12), suggesting that the factor is not a microneme protein

(**Figure 1d**). We also infected THP-1 cells with *T. gondii* parasites that were deficient in the

dense granule trafficking protein MYR1 ((13); kind gift from John Boothroyd, Stanford

University) and compared them to TgRHΔ*MYR1*:MYR1$_c$ parasites. TgRHΔ*MYR1* parasites

failed to induce any detectable CCL22 from THP-1 cells while, as expected,

TgRHΔ*MYR1*:MYR1$_c$ parasites induced significantly more than mock-treated cells (**Figure 1e**).

We also observed a very tight correlation between parasite multiplicity of infection (MOI) and

CCL22 levels, suggesting that the signal was primarily driven by the parasite rather than the

host cell (**Figure 1f**). Based on these results we felt confident that the unknown secreted factor

driving CCL22 production in human primary placental cells was very likely the same as the one

driving it in the THP-1 cell line and chose THP-1 cells for screening candidate effectors given

their tractability in the laboratory.

Transcript abundance correlation analysis identifies a large group of putatively

MYR1-trafficked gene products.

As described previously (above and (4)), we have determined that primary human trophoblast

cells infected with *T. gondii* have a transcriptional signature that is characterized by the

production of immunomodulatory chemokines, with CCL22 being the most potently induced. To

identify candidate *T. gondii* genes responsible for this effect on placental cells, and since this

effect required the *T. gondii* effector translocation complex protein MYR1 (4), we hypothesized

that MYR1-dependent substrates would have highly correlated gene expression profiles across

diverse gene expression datasets. To test this hypothesis, we generated an "all vs. all"

correlation matrix of 396 *T. gondii* Affymetrix microarray datasets that we downloaded and

curated from the Gene Expression Omnibus (see Materials and Methods). Analysis of the entire

correlation matrix (shown in **Figure 2a** and downloadable at 10.6084/m9.figshare.16451832)

confirms this hypothesis for certain gene classes. For example, we identified one cluster

containing multiple SAG-related sequences (SRS) which are typically expressed at high levels

in bradyzoites (including SRS49 cluster members A, C and D; **Figure 2 Supplement 1A**) and

another containing 70 genes, 43 of which encode ribosomal subunits (**Figure 2 Supplement**

1a). Examination of the gene expression heatmaps across all 396 microarray analyses clearly

show distinct patterns of gene expression in these two clusters depending on life stage

treatment exposure (**Figure 2 Supplement 1a,b**).

We quantified the degree of transcript abundance correlation between 5 "bait" genes

(*MYR1*, *2* and *3*; (13, 14)) and the known *MYR1*-dependent substrates *TgGRA24* and *TgIST*

(15, 16)) and all other genes across all 396 expression datasets. We identified genes as

candidate MYR1 substrates if they had an average correlation with the 5 "bait" genes ≥0.7, a

dN/dS ratio ≥2, and the presence of a predicted signal peptide OR at least one transmembrane

domain. Using this set of filters we were left with 28 candidate genes (plus all 5 bait genes

which also met these cutoffs), including the known TgMYR-dependent substrate TgGRA25 (17).

Since all known MYR1 trafficked substrates are dense granule proteins, we eliminated any

surface antigens or soluble enzymes, leaving a number of confirmed dense granule proteins

(e.g., GRA4 and GRA8) and conserved hypothetical proteins. Importantly, when we examined

the correlation between the bait genes and all *T. gondii* genes annotated as "Dense granule"

either in the primary product notes or via user annotation, we found that not all dense granule-

encoding transcripts correlated highly with bait transcript levels (**Figure 3 Supplement 1a**),

indicating that our approach could discriminate between different classes of proteins secreted

from the same organelle. For example, while genes like *GRA32* (*TGME49_212300*) had

transcript levels with relatively high (>0.8) correlations with bait transcript levels, other genes

encoding GRA1, GRA2 and GRA11 paralogs had transcript levels that correlated much more

poorly with the bait genes. This is despite the fact that most of these dense granule encoding

genes have high transcript levels compared to other gene clusters as shown in the heat map in

Figure 3 Supplement 1a, demonstrating that our approach yielded an additional layer of

discrimination to categorize dense granule-trafficked gene products. Moreover, many of the co-

regulated genes are not yet annotated but based on our analysis one would predict that many

are likely to be dense granule protein derived secreted effectors or structural constituents of this

parasite organelle.

***T. gondii GRA28* is the gene responsible for CCL22 induction in human immune**

and placental cells.

When we specifically examined correlations between the bait genes listed above and *MYR1*, we

found that *MYR1* expression profiles were highly correlated at the transcriptional level with

MYR2/3 and *IST* (**Figure 2b,c** and **Figure 3a,** top), consistent with the idea that MYR1

substrates could be identified using this approach. After identifying a small list of candidate

genes (**Figure 3a,** bottom), we deleted each using CRISPR-CAS9 and screened for CCL22-

induction in THP-1 monocytes by ELISA. Among the five genes that we tested (including

GRA18 which was recently found to induce Ccl22 in mouse macrophages; (18)), we found that

GRA28 (*TGME49_231960*) was required for the induction of CCL22 secretion by infected THP-

1 monocytes (**Figure 3b**). We also found that Δ*Toxofilin* parasites had significantly reduced

levels of CCL22 induction (**Figure 3b**), albeit to a much lesser extent than *ΔGRA28* parasites.

We think it likely that this decrease is owed to the reduced invasion capacity of Δ*Toxofilin*

parasites rather than a direct impact of this gene product on host CCL22 production (19, 20).

To determine if GRA28 was responsible for CCL22 production by human placental cells,

we infected 2nd trimester human villous placental explants with WT and ΔGRA28 parasites and

observed a marked decrease in CCL22 production by explants exposed to *ΔGRA28* parasites

compared to WT (**Figure 3c**). To gain a broader understanding of the transcriptional networks

altered by *GRA28* we compared THP-1 cells infected with RHΔ*HPT:HPT* and RHΔ*GRA28* using

RNAseq. A relatively small number of transcripts had significantly altered abundance using

stringent statistical cutoffs (67 genes with Padj<0.0001 and abs(log$_2$FC≥2) are highlighted in

Figure 3d) and these included CCL22 as well as the chemokines XCL1 and XCL2. Interestingly

transcript abundance for CCL17, a chemokine that is often co-regulated with CCL22 (21, 22)

and which is induced in some cells along with CCL22 by the *T. gondii* effector GRA18 (18), was

not dependent on *GRA28* (**Figure 3d**). The majority of transcripts that were GRA28-dependent

were of higher abundance in WT compared to *ΔGRA28* parasites when using slightly relaxed

statistical cutoffs (263 higher, 33 lower; P$_{adj}$<0.05 and log$_2$FC≥1 or ≤1).

We performed pathway analysis on these sets of regulated genes using Ingenuity

pathway analysis (IPA) and identified host cell pathways that were either more or less induced

in WT *T. gondii*-infected cells compared to *ΔGRA28*-infected cells (**Figure 4**), including

Dendritic Cell Maturation, IL6 and IL8 signaling, and NFκB signaling (**Figure 4a**). When we

assessed the degree of gene overlap in these gene sets, we found that 10 of the pathways

contained the *JUN* and *FOS* genes (**Figure 4b**), indicating a potential role for AP-1 complex

targeted transcripts in GRA28-dependent transcriptional changes. We examined correlations

across these gene sets (after creating a matrix of presence/absence of each of the genes

shown in **Fig. 4b**) and identified two non-overlapping sets of genes. The larger cluster contains

multiple immunity-related genes while the smaller cluster contains genes involved in

proteoglycan synthesis (**Figure 4c**), including the *XYLT1* gene which encodes the enzyme that

adds UDP-Xylose to serine residues as a first step in glycosaminoglycan synthesis. When we

performed a similar analysis using the "upstream regulator" module in IPA, we identified a small

set of significant (Z-score ≥2; *P*<0.001) regulatory factors that were upstream of the GRA28-

dependent gene set, including multiple regulators associated with the NFκB pathway (**Figure 4**

Supplement 1a). Cluster (**Figure 4 Supplement 1b**) and downstream gene overlap (**Figure 4**

Supplement 1c) analyses further confirmed the *FOS* and *JUN* genes as contributing to the

signaling pathways that were GRA28-dependent, while also confirming a putative role for NFκB.

For example, the cluster with the most similar target gene overlap contains multiple genes in the

NFκB pathway (*NFKBIA*, *NFKB1*, *RELA*) (**Figure 4 Supplement 1c**). However, when we co-

transfected HEK293 cells with NFκB luciferase reporter constructs and a construct encoding the

first exon of *GRA28* (see below) we saw no increase in the levels of luciferase after GRA28

transfection in contrast to a known NFκB activating construct containing multiple Caspase

Activation and Recruitment Domains (CARDs; **Figure 4d**). This suggests that NFκB activation

may not play a role in CCL22 induction. Other candidate transcriptional mediators with GRA28-

dependent transcript levels are *FOS*, *JUN* and *IRF4* (**Figure 4 Supplement 1d**). Transcript

levels of *JUN* have been shown in numerous studies to increase in a variety of host cells after

infection with *T. gondii* (23, 24). To test whether GRA28 played a role in altering C-JUN

abundance during infection we infected THP-1 cells with WT or ΔGRA28 *T. gondii* parasites for

24 h and using semi-quantitative western blotting to quantify C-JUN protein levels. While

infection of THP-1 cells clearly increased C-JUN levels compared to mock-treated cells (**Figure**

4 Supplement 2), the presence or absence of GRA28 in the infecting strain had no significant

impact on C-JUN protein abundance. These data suggest that while JUN transcript levels

appear to be at least somewhat dependent on GRA28 in the infecting strain (**Figure 4**

Supplement 1d), this does not appear to be detectable at the protein level using western

blotting.

The first exon of *GRA28* is sufficient for induction of CCL22 in during parasite

infection of and ectopic expression in human cells.

The *GRA28* gene has been described previously as encoding a dense granule protein that was

capable of trafficking to the host cell nucleus during infection (25). However, the exact structure

of the GRA28-encoding gene was somewhat ambiguous based on its annotation in ToxoDB.

Specifically, while TGME49_231960 is predicted as a single exon gene spanning ~7.4 kb of

genomic sequence (**Figure 5a**), the annotated gene is shorter in TGGT1 (**Figure 5a**) and split

into two gene products in *T. gondii* strains VEG, FOU, ARI, VAND, MAS, CATPRC2 and P89.

The 5' end of the gene was consistently predicted across all annotated genes, including the

precise location of the first intron. When we performed *de novo* assembly of the *T. gondii* RH

transcriptome, we were unable to identify any assembled transcripts that spanned the entire

length of the TGME49_231960 prediction, most likely due to the fact that a 39 bp repeat in

between each of these transcripts disrupted the assembly process (repeat consensus

sequence: CAGCAGCAGCCACAAGGGWMTGTTGTGCATCAACCACTA; **Figure 5a**). However

it should be noted that when we examined recently released Oxford Nanopore long-read single

molecule sequencing of *T. gondii* transcripts that are available on ToxoDB.org there are multiple

reads that span this repeat region (select Nanopore reads shown in **Figure 5a**), suggesting that

the gene is at least similar to that predicted for ME49 in the *Toxoplasma* genome database.

Regardless, given the challenges associated with amplifying and cloning this repetitive region

we expressed an HA-tagged version of the first exon of GRA28 in *T. gondii* and observed

expression within both the parasites and HA signal in the nucleus of infected cells (**Figure 5b**).

Importantly, CCL22 induction could be restored in an RH*ΔGRA28* clone after bulk transfection

of the exon 1 GRA28-expression construct prior to infecting THP-1 cells (**Figure 5c**), confirming

the role of sequences present in the first exon of GRA28 in driving CCL22 production in human

cells. Similar results were obtained when we transiently expressed a construct containing the

entire genomic locus for the predicted *T. gondii* GT1 *GRA28* gene (light green bar, **Figure 5a**) in

RH*ΔGRA28* parasites (**Figure 5d**). In contrast, when the Exon 1 construct was expressed

transiently in *Neospora caninum* (strain NC-1; (26)), we did not observe any HA signal in the

infected host cell despite expression of the protein within the parasite (**Figure 5e**). When we

quantified host nuclear HA signal intensity (background subtracted and then normalized to

staining intensity within the parasite; see Materials and Methods) in infected host cells there was

a clear and significant (*P*=0.0012) difference in the amount of HA-derived signal in the host

nucleus when TgGRA28 was expressed in *T. gondii* compared to *N. caninum* (**Figure 5**

Supplement 1a). Close inspection of multiple images suggest that the trafficking of *T. gondii*

GRA28 within *N. caninum* itself may be distinct from how it traffics in *T. gondii*. For example HA

staining was observed mostly within the parasite for *N. caninum* but could be found both within

T. gondii and at the vacuole periphery (**Figure 5e** and **Figure 5 Supplement 1**). While clearly

N. caninum failed to traffic detectable amounts of GRA28 into the host cell, this could be due to

a) poor trafficking of the protein within the parasite such that it never gains proper access to

vacuolar export machinery components like MYR1 and/or b) poor trafficking from the parasite

into the host cell due to incompatibility with the *N. caninum* export machinery. Interestingly *N.*

caninum does not appear to have an intact *GRA28* gene in its genome (see the synteny map for

TGME49_231960 at ToxoDB.org), although it does have a *MYR1* ortholog which has been

shown to be sufficient to traffic secreted *T. gondii* proteins into the host nucleus. Finally,

GRA28-Exon1 (minus the residues encoding the predicted signal peptide) could be robustly

expressed in HeLa cells with a V5 tag where it trafficked to the host cell nucleus (**Figure 5f**) and

also was functional when expressed ectopically in THP-1 cells where it induced CCL22

secretion (**Figure 5f**).

GRA28 induction of Ccl22 is fully conserved in mice.

To determine whether parasite-driven induction of CCL22 is conserved in the murine model, we

compared WT and GRA28-deficient (*ΔGRA28*) parasites for their ability to induce this

chemokine *in vitro*, *ex vivo*, and *in vivo*. First, we infected mouse macrophages (RAW 264.7) *in*

vitro with type 1 strain (RH) *T. gondii* parasites (WT), or (RH) *ΔGRA28 T. gondii* parasites at

MOIs of 3. Based on Ccl22 ELISA, mouse macrophages not only release more Ccl22 protein

during *T. gondii* infection, but similar to human THP-1 cells this phenotype is also dependent on

the presence of *T. gondii* secreted protein GRA28 (**Figure 6a**). Next, we investigated whether

primary mouse tissues, specifically mouse placental tissue, also elicit this response to *T. gondii*

infection. Embryonic day 12.5 Swiss Webster mouse placentas were halved and distributed into

separate treatment groups. These placental explants were then infected *ex vivo* with 2.0 x 10^6

Type 1 strain (RH) *T. gondii* parasites (WT), (RH) *ΔGRA28 T. gondii* parasites, or mock

treatment. As shown in **Figure 6b,** primary mouse placental tissue also responds to *T. gondii*

infection by releasing Ccl22 protein in a GRA28-dependent manner. RNA was also extracted

from the infected placental samples and we performed RNAseq. As shown in **Figure 6c** the

number of transcripts that varied in a GRA28-dependent manner was markedly small,

suggesting that GRA28 is a highly specific inducer of *Ccl22* in mouse placental explants. Of the

three genes with significantly higher transcript levels (*Ccl22*, *Il12rb2*, *Ccr7*) in wild type

1nfections as compared to *ΔGRA28* infections, *Ccl22* was the most highly induced. These data

show conservation of the parasite-driven Ccl22 phenotype 1n primary mouse placental explants

at both a protein and transcript level. Finally, we investigated mouse *in vivo* Ccl22 responses to

T. gondii intraperitoneal infection. Female BALB/cJ mice (n = 3 for each treatment) were

infected with WT, *ΔGRA28*, or mock *T. gondii* treatments. We focused on early, acute infection

and performed Ccl22 ELISA on serum (**Figure 6d**) and peritoneal lavage fluid (**Figure 6e**).

These suggest *in vivo* Ccl22 protein levels are at least partially dependent on GRA28.

Moreover, while there was a significant amount of systemic Ccl22 protein detected in serum of

infected mice, even in the *ΔGRA28* parasite treatment, Ccl22 was almost undetectable in

peritoneal lavage fluid in *ΔGRA28*-infected mice. Overall, these data indicate the process

driving *T. gondii* GRA28 induced Ccl22 is similar, if not the same, in both mice and humans, and

that this parasite effector can mediate robust changes in Ccl22 production at the site of infection

and systemically.

GRA28-deficient parasites have distinct inflammatory and dissemination phenotypes in the acute and chronic phases of infection, respectively.

To determine the impact of *T. gondii* GRA28 *in vivo* we indexed differences in mouse behavior relevant to inflammatory responses and quantified differences in infection-induced weight loss and total morbidity after infection of BALB/cJ mice with either RH:WT or RH*ΔGRA28 T. gondii*. We observed no significant differences in morbidity or weight loss (**Figure 7a,b**). However, when we scored (**Figure 7 Supplement 1**) mice over the course of infection as to the extent of inflammation-induced behavioral changes, we observed significantly heightened fur ruffling in the *ΔGRA28*-infected mice on days 6 and 7 post-infection (**Figure 7c**), despite the fact that mortality was unchanged.

We also generated *ΔGRA28* parasites in a Type 2 *T. gondii* background that had been previously engineered to express luciferase and GFP (specifically ME49*ΔHPT:LUC*; (27, 28)) to permit non-invasive quantification of parasite burden and dissemination over the course of infection. For the ME49 strain infections we observed only minor and non-significant differences in mouse morbidity and weight loss (**Figure 8a,b**). However, during the acute phase of infection we observed slight differences in parasite burden between ME49*ΔHPT:LUC* (WT) and ME49*ΔGRA28*-infected mice, with burden being significantly *higher* in ME49*ΔGRA28* compared to WT on day 9 post-infection (**Figure 8c**). This difference was not due to experimental variation in parasite input between strains since parasite burden was indistinguishable during the first 6 days post-infection (**Figure 8c**). In contrast to these minor differences during the acute phase of infection, we observed more dramatic differences in parasite burden during the later stages of infection. Specifically, quantification of *in vivo* bioluminescence data taken dorsally on days 14 and 15 post-infection revealed that WT parasites were of much greater abundance in the brains compared to those infected with ME49*ΔGRA28* (**Figure 8d,e**).

DISCUSSION:

T. gondii-infected host cells have dramatically altered transcriptomes compared to uninfected cells, and effectors that are secreted from the parasite during invasion drive most, but not all, of these changes (29). To date, the vast majority of these parasite effectors are derived from the dense granule and rhoptry organelles. We previously identified that T. gondii induces the production of CCL22 in human placental trophoblasts, while human foreskin fibroblasts do not exhibit this chemokine induction during T. gondii infection (4). Additionally, our previous work has shown that this induction required parasite invasion and the effector chaperone-like T. gondii gene product MYR1 (4, 13). While T. gondii induces CCL22 during infection of a variety of cell types from both mice and humans (18, 30), including at the transcriptional level in mouse brain (31), placental cell CCL22 induction is driven by a highly specific parasite effector, GRA28. CCL22 production is considered to be an indication of M2 macrophage polarization, and macrophage polarization has been linked to strain-specific T. gondii effectors like ROP16 and GRA15 (32). The impact of GRA28 is distinct from these effectors because CCL22 induction occurs similarly in all three canonical strain types, and GRA28 does not alter the expression of other M2-associated genes (such as *IL4*, *IL10*, *IL13* or *ARG1*), in the cell types that we have assayed. The specificity of GRA28 for only a few target genes is novel compared to effectors like GRA15 and ROP16 (24, 33) that alter the abundance of hundreds of transcripts.

Transcriptional co-regulation has been used in other systems as a means to identify members of protein complexes (34), but to our knowledge this is the first time this approach has been successfully applied at this scale in T. gondii. We used 396 microarray datasets derived from multiple T. gondii life stages and experimental manipulations to provide enough variation to better distinguish subclusters within closely-related gene families. Genes encoding dense

granule proteins are among the most highly expressed in the *T. gondii* genome, making them

more difficult to separate from one another, but they still clustered into two distinct groups with

functional themes. The MYR1/GRA28 cluster harbored a handful of known secreted dense

granule effectors, while the other contained genes encoding dense granule structural proteins or

those that are secreted into the vacuole but do not traffic to the host cell. We anticipate that the

former cluster can be exploited further to identify additional MYR1-trafficked, and putatively

host-modulating, effectors while the latter has highlighted new candidates important in dense

granule structure or function within the parasite. The entire dataset is available for download as

a text file at Figshare (doi: 10.6084/m9.figshare.16451832) so that these data can be mined to

identify candidates for membership in other critical *T. gondii*-specific protein complexes.

GRA28 was previously shown to encode a dense granule protein secreted from the

parasite into the host cell where it trafficked to the host cell nucleus (25), but its impact on the

host cell was unknown. Its natural presence in the host nucleus during infection has also been

further confirmed using proteomics, where it was found to be one of the more abundant *T.*

gondii proteins in the nucleus of the infected host cell (35). The fact that it affects the abundance

of only a small number of chemokine-encoding genes at the transcriptional level suggests that it

modulates transcriptional activity via direct interactions with transcription factors and/or

upstream regulatory sequences. Other *T. gondii* effectors traffic to the host nucleus but this is

not always critical for function. For example, ROP16 localization to the host cell nucleus is

dispensable for its primary function of phosphorylating STAT6 which occurs in the cytoplasm of

the host cell (36). Other *T. gondii* effectors like IST (15, 37) and GRA24 (16) function within the

host cell nucleus, but many of these mediate changes in hundreds of transcripts via their

cooperation with existing transcriptional suppressors (IST; (15, 37)) or activators (GRA24; (16)).

It remains to be seen if the function of GRA28 can occur independent of nuclear trafficking or if

this ultimate localization is required for chemokine induction, but its specificity for downstream

genes raises the interesting hypothesis that it may function directly, possibly as a heterologous

transcription factor.

The signaling pathway governing GRA28 function is unknown but some clues can be

found in our pathway analyses which suggest a role for GRA28 in mediating changes in key

immunity-related host cell signaling pathways. The transcription factor genes *JUN*, *FOS* and

components of the NFκB complex were consistently linked to the GRA28-dependent host

transcripts. LPS is a well-known activator of both NFκB and C-Jun activity in THP-1 cells (38,

39), and this can occur via Toll-like receptor activation (40). However *T. gondii* induction of

CCL22 was not fully dependent upon host MYD88, since MYD88[(-/-)] THP-1 cells still produced

significant amounts of CCL22 in response to *T. gondii* infection (**Figure 1 Supplement 1c**). The

difference in CCL22 production by the MYD88[(-/-)] cells in comparison to the WT cells should also

be considered in light of the fact that the cell lines have different origins (and therefore distinct

passage histories which could have the more subtle effects shown on CCL22 production after

infection). A distinct cluster of GRA28-dependent host genes was identified that encoded gene

products involved in proteoglycan synthesis, including the rate-limiting enzyme XYLT1 . *T.*

gondii attachment to host cells is mediated by interactions between parasite adhesins and host

cell surface sulfated proteoglycans (PG) like heparan sulfate (41, 42)*,* and *T. gondii* adheres

poorly to cells with genetically or enzymatically depleted levels of surface sulfated proteoglycans

(41, 42). Therefore direct and/or indirect modulation of XYLT1 transcript levels by GRA28 may

serve to make infected cells susceptible to adhesion, and ultimately invasion, by *T. gondii* or any

other pathogens that depend on surface proteoglycans.

GRA28 had no impact on transcript levels of the gene encoding CCL17 which is

commonly co-regulated along with *CCL22*. Mouse macrophages infected with *T. gondii* produce

Ccl17 and Ccl22 and this is due, at least in part, to another *T. gondii* effector GRA18 (18). Using

the same *GRA18* knockout lines (kindly provided by the Bougdour lab) we found that GRA18

had no impact on CCL22 production at the transcriptional (not shown) or protein (**Figure 4**) level

in human THP-1 cells, suggesting that GRA18 and GRA28 have distinct targets. This is also

consistent with the observation that Ccl22 induction in RAW macrophages is only partially

dependent on GRA18 and β-Catenin signaling, in contrast to Ccl17 and Ccl24. Finally, in our

work we used lower MOIs (2-3 here compared to 5-6 in (18)). Regardless, GRA28 appears to

be the more potent modulator of Ccl22 production compared to GRA18, while Ccl17 appears to

be much more dependent on GRA18. It is exciting to speculate that *T. gondii GRA28* has

evolved to uniquely target CCL22 as a means to gain access to the fetal compartment since this

chemokine is potently induced in placental cells and this chemokine plays a role in immune

tolerance during pregnancy (7). However, as shown clearly in this study, GRA28 also alters

monocyte/macrophage CCL22 production, making it equally plausible that this intricate

molecular relationship developed first as a more generalized immune evasion (via suppression)

strategy.

The role of specific chemokines like CCL22 during *T. gondii* infection is poorly

understood but the discovery of *GRA28* allows this to be addressed more directly using *T.*

gondii ΔGRA28 parasites from different genetic backgrounds. Hypervirulent *T. gondii* RH strain

ΔGRA28 parasites caused inflammation-related behavioral changes earlier during infection in

mice, compared to mice infected with WT parasites, suggesting that GRA28 functions to

suppress inflammatory responses (likely due to induction of CCL22 although we did not test this

directly). This could arise via GRA28-mediated recruitment and/or activation of regulatory T cells

to the site of infection. These behavioral changes occurred without an effect on the acute

virulence phenotype as all mice succumbed to the infection with similar kinetics, which is

consistent with an impact of GRA28 on suppressing inflammatory responses without altering the

ability of the mouse to control parasite replication. However, after infections using the type 1

parasite genotype we observed a significant reduction in *ΔGRA28* parasite burden in the brain

compared to wild type parasites. This effect was unexpected given the fact that parasite burden

was statistically equivalent during the acute phase of infection, but points to a potential

important role for GRA28 in altering the host innate immune response in a manner that

increases host susceptibility to dissemination of *T. gondii* across critical barriers like that

guarding the CNS. *T. gondii* can infect blood-brain barrier epithelial cells as a means to cross

into the host CNS (43), so GRA28 may promote parasite survival at this critical interface by

recruitment of regulatory T cells or other cell types that might downregulate inflammatory

responses.

Summary: Taken together our data point to a specific role of *T. gondii* GRA28 in modulating

chemokine production in the infected cell. Importantly, this effect occurs only in certain cell

types, including cells from both human and mouse placenta. A relatively small number of host

chemokines are affected by parasites expressing this gene, and it plays a role in both

modulation of the inflammatory response (as evidenced by mouse behavior and appearance

during infection) and ultimately parasite dissemination to "privileged" sites like the CNS.

METHODS

Cell Culture

All cell and tissue cultures were incubated at 37°C and 5% CO_2. All media were supplemented

with 10% fetal bovine serum (FBS; Atlas Biologicals), 2 mM L-glutamine, and 50 mg/mL

penicillin-streptomycin. Human foreskin fibroblast (HFF) cells were grown in Dulbecco's

Modified Eagle Medium (DMEM; Gibco), Raw264.7 cells were grown in DMEM (Gibco) with 10

mM HEPES, and THP-1 cells were grown in RPMI 1640 medium (Corning). THP-1 cells were

assayed for viability using Trypan Blue staining (0.4%) (Gibco), counted, spun at 120 x g for 10

minutes at 24°C and medium replaced with supplemented DMEM prior to infection. All THP-1

cell numbers listed are based on trypan blue-negative cells.

Human Placental Explants

Human placental tissue from less than 24 weeks of gestation was obtained, cultured, and

infected with *T. gondii* as described previously (44).

Mouse Placental Explants

Mouse placental tissues were obtained by dissection of E12.5 or 18.5 Swiss Webster mice.

Upon removing the fetuses from the mother, the placentas were dissected away from other

tissues and placed into pre-warmed 37°C PBS. The placentas were washed 3x in fresh pre-

warmed PBS. Each placenta was then cut in half with sterilized surgical scissors and each half

was placed into a well on a plate with pre-warmed 37°C DMEM with 10 mM HEPES, 10% FBS ,

2 mM L-glutamine, and 50 mg/mL penicillin-streptomycin. Each placenta had one half-piece of

tissue represented in each treatment group. For *T. gondii* infections, isolated tissue was infected

immediately with $5.0 \times 10^5 - 2.0 \times 10^6$ parasites for ~24 hours.

Parasites

Type 1 (RH, GT1), Type 2 (Me49, Pru), Type 3 (Veg, CEP) *Toxoplasma gondii* tachyzoites and

sporozoites, *Neospora caninum* (NC-1) tachyzoites, and *Hammondia hammondi* (HhCatAmer

and HhCatEth1; (45, 46)) sporozoites were used in this study. Sporozoites were excysted from

sporulated oocysts as described (47) and either used immediately or grown for 24 h in human

foreskin fibroblasts prior to being used in controlled infections. Tachyzoites were maintained by

continual passage in human foreskin fibroblast (HFF) cultures incubated at 37°C and 5% CO_2 in

DMEM supplemented with 10% fetal bovine serum (FBS) (Atlas Biologicals), 2 mM L-glutamine,

and 50 mg/mL penicillin-streptomycin. The Rh*YFP* strain was a gift from David Roos (University

of Pennsylvania), the Rh*ΔMYR1* and Rh*ΔMYR1:MYR1$_c$* parasites (13) were a gift from John

Boothroyd (Stanford University), the PRUΔGRA18 and complemented knockout parasites were

shared by Alexandre Bougdour (18), and the PruΔ*Toxofilin* KO parasites (19) were a gift from

Melissa Lodoen (UC Irvine). For infections, infected monolayers were washed with fresh

cDMEM and then scraped and syringe lysed to release tachyzoites. These tachyzoites were

then passed through a 5 µM syringe filter and counted. Parasites were then centrifuged at 800 x

g for 10 minutes at 24°C and resuspended and diluted in cDMEM before being used in

infections. Mock treatments were produced by filtering the same parasites through a 0.22 µM

syringe filter and exposing host cells to the same volume of the filtrate as was used for parasite

infections. Freeze treatments were produced by subjecting the parasites to -80°C for 15

minutes, fixation treatments by exposure to 4% paraformaldehyde for 10 minutes followed by

washing in PBS, and sonication treatments by sonicating at 0°C using five 30-second bursts at

50 amps with 30-second cooling intervals in between bursts followed by microcentrifugation at

800 x g for 10 minutes to generate soluble (S) and pellet (P) fractions.

Invasion inhibitor assays

For Cytochalasin-D (Cyt-D) treatment, parasites were pre-treated with 10 µg/mL of Cyt-D in

cDMEM for 1 hour and then used to infect cells in the presence of 10 µg/mL of Cyt-D in cDMEM

for the duration of infection. Vehicle of Cyt-D is DMSO (40 µL per mL of cDMEM). For 4-

Bromophenacyl bromide (4-BPB) treatment, parasites were pre-treated with either 0.5 or 1 µM

of 4-BPB for 15 minutes. 4-BPB was dissolved directly in cDMEM. Parasites were then washed

twice with normal cDMEM with 10-minute 800 x g spin steps between each wash, and then

used to infect cells in the presence of normal cDMEM.

Plaque Assays

Parasites were serially diluted in media and used to infect monolayers so that each tissue

culture flask of HFFs received 100 parasites. These flasks were then incubated at 37°C in 5%

CO_2 undisturbed for 5-7 days. At the end of the incubation period, each flask was counted for

number of plaques present and parasite viability was calculated. Crystal violet staining was

used to count plaques as follows: the monolayer was washed with PBS and fixed for 5 minutes

with ethanol. Then crystal violet solution (12.5 g crystal violet in 125 mL ethanol mixed with 500

mL 1% ammonium oxalate in water) was introduced to the monolayer and allowed to stain for 5

minutes. The monolayer was then washed extensively with PBS and allowed to air dry prior to

counting plaques.

Candidate gene identification using transcript level correlation analysis

To identify candidate effectors for inducing CCL22 we exploited the fact that the CCL22

induction response in THP-1 and placental cells required the presence of the *T. gondii* effector

translocation protein MYR1 (13, 14). We hypothesized that MYR1-dependent effectors would

have similar transcript abundance profiles across diverse expression datasets. We downloaded

396 publicly available *T. gondii* microarray expression datasets from the Gene Expression

Omnibus platform hosted by the NCBI (48). We loaded and processed each CEL file using the

"affy" module implemented in R (49). Data were processed and normalized using the following

commands: bgcorrect.method = "rma", normalize.method = "quantiles", pmcorrect.method =

"pmonly", summary.method = "medianpolish". RMA-normalized data were exported, re-imported

into R and then transposed. An all-versus-all Pearson correlation matrix was generated using

the "cor" function from the R:Stats base module. Probe names on the Affymetrix array (in the

format of XX.mXXXXXX) were converted to current TGME49 gene models using data

downloaded from ToxoDB and the Vlookup function in Microsoft Excel. In some cases the

microarray annotations could not be matched to current TGME49 gene model names and are

shown as blanks in plots. This correlation matrix, the normalized array data used to generate it,

and a key to convert Affymetrix probe names current gene model names are all available on

Figshare (10.6084/m9.figshare.16451832). To analyze this correlation matrix we used

hierarchical clustering tools implemented in R including heatmap.2 (from the gplots package)

and the dendextend package. To identify candidate genes using this matrix we calculated the

mean correlations between 5 bait genes and all other queried genes from the microarray. The

bait genes were known to encode either components of the MYR complex themselves or known

TgMYR substrates (TgMYR1; (13), TgMYR2, TgMYR3 (14), TgGRA24 (16) and TgIST (15)).

Most of the candidate CCL22-inducing genes were identified based on having a) an average

correlation with the 5 "bait" genes listed above ≥0.7, b) a dN/dS ratio ≥2, c) and the presence of

a predicted signal peptide OR at least one transmembrane domain (which we reasoned could

be a cryptic signal peptide if the wrong start codon was chosen for the current gene annotation).

De novo transcript assembly

To identify and assemble transcripts coding for GRA28 we used *de novo* transcript assembler

Trinity ((50); version 2.6.6; default settings) using triplicate RNAseq datasets from WT *T. gondii*

RH parasites infecting THP-1 cells (see below). Assembled transcripts with similarity to the

predicted *GRA28* gene (TGME49_231960) were identified using BLASTN and BLASTX.

Primary plots were generated using GenePalette software (51) and then modified.

CRISPR-mediated gene disruption and validation of knockouts

The pSAG1::CAS9-U6::sgUPRT plasmid provided David Sibley (Addgene plasmid #54467;(52))

was modified using the Q5 Site-Directed Mutagenesis Kit (NEB) so that the gRNA sequence

was replaced with two restriction enzyme sites (sequence: GTTTAAACGGCCGGCC) for PseI

(NEB R0560S) and FseI (NEB R0588S). This modified plasmid was then used as the template

for all future Q5 reactions. Two unique gRNA sequences were created for each candidate gene

by utilizing the genomic sequences for *T. gondii* GT1 (toxodb.org) and E-CRISP (e-crisp.org)

using the ToxoDB-7.1.31 reference genome. A forward primer for each gRNA was created for

use with the modified pSAG plasmid, with the unique gRNA sequence followed by a section of

the plasmid scaffolding (GTTTTAGAGCTAGAAATAGCAAG). The reverse primer used with this

plasmid is AACTTGACATCCCCATTTAC. The gRNA sequences for the genes mentioned in this

study and the primers used to validate the knockouts are listed in **Supplementary Table 1**.

A plasmid was created for each gene of interest (GOI) using the modified pSAG plasmid

template and the Q5 Site-Directed Mutagenesis Kit (NEB) by following the manufacturer's

protocol with a few adaptations. The KLD enzyme step was extended to 60 minutes incubation

at room temperature, and following the KLD enzyme step the product was heated to 65°C for 20

minutes and then double digested with PseI and FseI in CutSmart buffer (NEB) for 60 minutes

at 37°C to remove any remaining plasmid that was not eliminated by the DpnI in the KLD step.

This digested product was then heated to 65°C for 20 minutes to deactivate the enzymes prior

to transformation, plasmid isolation and sequencing to validate insertion of the correct gRNA

sequence (pSAG:GOI:gRNA). Parasites were transfected with either a single gRNA plasmid or

equal amounts of plasmids encoding two gRNAs targeting the same gene. For validation of

knockouts after cloning, a clone was considered a knockout if PCR across a targeted cut site

failed or if a PCR reaction across the entire gene (just upstream and downstream of the start

and stop codons, respectively) failed. In some cases where all PCR reactions worked (indicating

that the plasmid failed to insert at the gRNA target site), the amplified band was sequenced and

a clone was considered a knockout if insertions/deletions were identified near the gRNA binding

that resulted in frame shifts and premature stop codons.

Parasite transfections

In general, transfections were performed using standard approaches. Briefly, parasite

suspensions were obtained by needle passage (25 and 27 gauge needles) and then pelleted for

10 minutes at 800 x g. Parasites (~2 x 10^7 per transfection) were re-suspended in Cytomix (120

mM KCl; 0.15 mM $CaCl_2$; 10 mM KPO_4; 25 mM Hepes, 2mM EDTA, 5mM $MgCl_2$; pH to 7.6)

containing GSH and ATP and electroporated at 1.6 Kv with a capacitance setting of 25 µF using

a BTX ECM600 Electroporator. Transfected parasites were then used to infect coverslips and/or

flasks of confluent HFFs and placed under appropriate selection. For candidate gene knockouts,

~2x10^7 *T. gondii* RHΔHPT parasites were transfected with ~30-50 µg of the relevant

pSAG:GOI:gRNA plasmid(s) (described above) along with 2-5 µg of an empty pGRA-HA-HPT

(53) plasmid. Parasites were placed under selection the next day and cloned by limiting dilution

after 2-3 passages. Individual clones were screened for gene deletion by PCR and sequencing

to permit identification of both target gene disruptions (via insertion of the pGRA-HA-HPT

plasmid at the CAS9 cut site) or mutation via DNA repair events at the CAS9 cut site. For HA-

tagging experiments, Type 1 (RH) *GRA28* exon 1 (residues 1-498) was C-terminally HA tagged

by cloning into the *T. gondii* expression plasmid pGRA-HA-HPT (53). This plasmid drives

protein expression using the highly active *GRA1* promoter. TgRHΔ*HPT* or *N. caninum* Liverpool

(NcLIVΔ*HPT*; (54)) parasites were transfected with ~40-60 µg of *GRA28* exon 1 plasmid and

cells were grown overnight in normal media. For analysis of transiently transfected parasites,

cells were only grown for 18 h post-transfection while for stable transfection parasites were

grown for 2-3 passages in media containing 50 µg/mL of mycophenolic acid and xanthine. Cells

were fixed with 4% PFA and permeabilized in 0.1%Triton/PBS. Samples were probed with anti-

HA rat monoclonal antibody (3F10 clone, Roche) diluted to 0.1 mg/mL in 0.1%Triton-PBS buffer

and washed four times in PBS. Samples were then incubated in 488 goat anti-rat (Life

Technologies Alexa Fluor H+L) followed by PBS washes. All samples were mounted in

Vectashield with DAPI (Vector laboratories). For genetic complementation, TgRHΔ*GRA28*

parasites were transfected the exon 1 construct used above or a construct amplified from

genomic DNA encompassing the start and stop codons of the GT1 version of GRA28

(TGGT1_231960). Expression plasmids (~30 µg) were co-transfected along ~5 µg of

pLIC_3xHA_DHFR* plasmid (kindly provided by Vern Carruthers; (55)), and populations were

placed under 1 µM pyrimethamine selection for 2-3 weeks, and then used to infect THP-1 cells

for 24 h as above followed by assays to quantify CCL22 in culture supernatants by ELISA.

Fluorescence image analysis

To compare signal intensity in the nucleus of host cells infected with either *T. gondii* or *N.*

caninum parasites transiently transfected with the HA-tagged GRA28 exon 1 construct, we

scanned stained coverslips for GRA28-HA-positive vacuoles, and then used Fiji (an

implementation of Imagej;) to calculate 1) the average HA signal intensity in the nucleus of the

infected host cell (AvgIntInf), 2) the average HA signal intensity in the nucleus of a neighboring,

uninfected host cell (AvgIntUninf) and 3) the average signal intensity of the parasite-containing

vacuole (AvgIntVacuole). We then used the following calculation to determine the normalized,

background-subtracted nuclear signaling intensity:

$$\frac{(AvgIntInf - AvgIntUninf)}{AvgIntVacuole}$$

Example images of this process are shown in **Figure 5** and **Figure 5 Supplement 1**. Data were

log10 transformed prior to performing a Student's T-test.

RNA-seq

RNA was isolated from cultures using the RNeasy Mini Kit (QIAGEN) and its associated RNase-

Free DNase digestion set (QIAGEN), following the manufacturer's protocol for mammalian cells.

An Agilent Bioanalyzer was used to check the quality of the RNA samples. Tru-Seq stranded

mRNA libraries were generated from 5-17 ng/µl of mRNA for THP-1 cells, and 50-120 ng/µL of

mRNA for murine PECs and placental explants, and sequenced with an Illumina NextSeq 500

sequencer. mRNA-Seq FASTQ reads were mapped to the human reference genome (Homo

sapiens v81; hg38) using default options on CLC Genomics Workbench 11 (Qiagen). Total gene

reads (with at least 1 read count) were exported from CLC Genomics Workbench and used for

DESeq2 (56) to perform differential expression analysis using methods outlined previously (e.g.,

(4)). Data were evaluated using principal component analysis (embedded in the DESeq2

package) and genes were deemed to be significantly expressed if the $\log_2$ fold-change was ≥ 1

or ≤ -1 and with a P_{adj} value <0.01. Gene set enrichment analysis (GSEA;(57)) and Ingenuity

Pathway Analysis (QIAGEN; (58)) software were used to compare gene sets that were

differentially regulated after infection with WT and Δ*GRA28* parasites.

CCL22 and Ccl22 ELISA

CCL22/Ccl22 ELISAs were performed on culture supernatants (undiluted or diluted when

necessary) using Immulon 4HBX flat bottom microtiter plates with the human CCL22/MDC

DuoSet ELISA (R&D Systems DY336) or the mouse Ccl22/Mdc DuoSet ELISA (R&D Systems

DY439) per the manufacturer's instructions.

Mouse Experiments with WT and *ΔGRA28* parasites

To determine the impact of *T. gondii* candidate effectors on mouse morbidity and cytokine

production, BALB/cJ mice from Jackson laboratories (4-6 week old, female) were injected

intraperitoneally with 200 μL of PBS containing 1.0 x 10^6 *T. gondii* tachyzoites, or 200 μL of 0.22

μM filtered parasite solution as a mock treatment. Mice were sacrificed at 48 hours post

infection (HPI), and peritoneal exudate cells (PECs) were collected by injecting 3 mL sterile PBS

into the abdominal cavity, rocking the mouse to mix the PBS, and siphoning the PBS solution

into a sterile conical tube. The solution was then centrifuged at 1,000 x g for 10 minutes at 24°C,

the supernatant was collected, and RNA was extracted from the pellet for RT-qPCR and RNA-

seq. Blood was collected in Sarstedt Microvette CB 300 Z tubes by cardiac puncture and

centrifuged at 10,000 x g for 5 minutes to separate the serum. Mice were infected with RH*ΔHPT*

(wild type), RH*ΔMYR1* or RH*ΔGRA28* parasites depending on the experiment.

To determine the impact of *GRA28* deletion on *T. gondii* proliferation and dissemination,

female BALB/cJ mice (4 weeks old) from Jackson laboratories were injected intraperitoneally

with 200 μL PBS containing 100 *T. gondii* tachyzoites. In one experiment five mice received an

injection of RH*ΔGRA28* parasites, while the other five received an injection of the transfection

control parasite that was transfected with empty vector (Rh*ΔHPT:HPT*). For behavioral indices

of inflammatory responses, photographs of the mice were taken dorsally and laterally every 4-6

hours for the entire duration of the infection. Mice were visually scored 0-3 based on the

presence of fur ruffling, the location of ruffling, and the presence of skin redness/irritation

(**Figure 7 Supplement 1**). **0)** No fur ruffling or red/irritated skin present. **1)** Mild ruffling present

predominantly located on the head and back of the neck. No red/irritated skin visible. **2)**

Moderate ruffling present - fur forms larger clumps and extends to the rest of the body. Skin

may be visible through the clumps but is not red or irritated. **3)** Severe ruffling characterized by

fur ruffling across the entire body with visibly red/irritated skin in between fur clumps.

In a second experiment mice were infected with 1000 tachyzoites of *T. gondii* strain

ME49:*LUCΔGRA28* or a passage-matched wild type strain (27, 28). Mice were imaged daily

after injection of D-Luciferin as described previously (28, 59) using an IVIS Lumina II *in vivo*

bioluminescence imaging system (with ventral imaging occurring on all days post-infection and

ventral and dorsal imaging occurring starting on day 10 post-infection). Animals were

anaesthetized using 2% isoflurane during the 4-8 minute imaging period (ventrally and dorsally

where applicable). When necessary, blood was collected via submandibular lancet puncture,

collected into Sarstedt Microvette CB 300 Z tubes and spun at 10,000 x g for 5 minutes to

separate the serum. Mice were monitored extensively over the course of infection for symptoms

of morbidity and humanely euthanized. All animal procedures were approved by the Division of

Laboratory Animal Resources and IACUC and our animal facilities are routinely inspected by

the USDA and local IACUC committee.

Statistics

All statistics were performed in Graphpad Prism for Windows (versions 7 or 9; GraphPad

Software, La Jolla, CA). For most two treatment assays, we used unpaired, two-tailed student's

T test, and for multi-treatment/condition experiments we used one- or two-way ANOVA followed

by multiple comparisons post-hoc tests. Individual comparisons are listed for each assay in the

text and figure legend, and only pre-planned comparisons were performed to minimize Type 1

error. *In vivo* bioluminescence data (total flux; photons/s) and nuclear staining intensity data

(comparing nuclear trafficking of *T. gondii* GRA28 when expressed in *T. gondii* or *N. caninum*)

were log$_{10}$-transformed prior to statistical analysis.

Acknowledgements

706   The authors would like to thank Dr. Alexandre Bougdour (Inserm; Grenoble, France) for

707   providing GRA18 knockout and complemented strains, Drs. John Boothroyd and Michael Panas

708   (Stanford University) for helpful discussions and providing MYR1 knockout and complemented

709   strains, and Dr. Peter Bradley (University of California-Los Angeles) for helpful discussions and

710   sharing reagents related to GRA28 that facilitated this work.

BIBLIOGRAPHY

1.   Fallahi S, Rostami A, Nourollahpour Shiadeh M, Behniafar H, Paktinat S. 2018. An

updated literature review on maternal-fetal and reproductive disorders of *Toxoplasma*

gondii infection. J Gynecol Obstet Hum Reprod 47:133–140.

2.   Stagno S, Reynolds DW, Amos CS, Dahle AJ, McCollister FP, Mohindra I, Ermocilla R,

Alford CA. 1977. Auditory and visual defects resulting from symptomatic and subclinical

congenital cytomegaloviral and *Toxoplasma* infections. Pediatrics 59:669–678.

3.   Fahnehjelm KT, Malm G, Ygge J, Engman ML, Maly E, Evengård B. 2000.

Ophthalmological findings in children with congenital toxoplasmosis. Report from a

Swedish prospective screening study of congenital toxoplasmosis with two years of follow-

up. Acta Ophthalmol Scand 78:569–575.

4.   Ander SE, Rudzki EN, Arora N, Sadovsky Y, Coyne CB, **Boyle JP**. 2018. Human

Placental Syncytiotrophoblasts Restrict *Toxoplasma gondii* Attachment and Replication

and Respond to Infection by Producing Immunomodulatory Chemokines. MBio 9.

5.   Klarquist J, Tobin K, Farhangi Oskuei P, Henning SW, Fernandez MF, Dellacecca ER,

Navarro FC, Eby JM, Chatterjee S, Mehrotra S, Clark JI, Le Poole IC. 2016. Ccl22 diverts

T regulatory cells and controls the growth of melanoma. Cancer Res 76:6230–6240.

6.   Layseca-Espinosa E, Korniotis S, Montandon R, Gras C, Bouillié M, Gonzalez-Amaro R,

Dy M, Zavala F. 2013. CCL22-producing CD8α- myeloid dendritic cells mediate regulatory

T cell recruitment in response to G-CSF treatment. J Immunol 191:2266–2272.

7.   Freier CP, Kuhn C, Rapp M, Endres S, Mayr D, Friese K, Anz D, Jeschke U. 2015.

Expression of CCL22 and infiltration by regulatory T cells are increased in the decidua of

human miscarriage placentas. Am J Reprod Immunol 74:216–227.

8.   Martinez de la Torre Y, Buracchi C, Borroni EM, Dupor J, Bonecchi R, Nebuloni M,

Pasqualini F, Doni A, Lauri E, Agostinis C, Bulla R, Cook DN, Haribabu B, Meroni P,

Rukavina D, Vago L, Tedesco F, Vecchi A, Lira SA, Locati M, Mantovani A. 2007.

Protection against inflammation- and autoantibody-caused fetal loss by the chemokine

decoy receptor D6. Proc Natl Acad Sci USA 104:2319–2324.

9.   Martinenaite E, Munir Ahmad S, Hansen M, Met Ö, Westergaard MW, Larsen SK,

Klausen TW, Donia M, Svane IM, Andersen MH. 2016. CCL22-specific T Cells:

Modulating the immunosuppressive tumor microenvironment. Oncoimmunology

5:e1238541.

10.   Kimura S, Tanimoto A, Wang K-Y, Shimajiri S, Guo X, Tasaki T, Yamada S, Sasaguri Y.

2012. Expression of macrophage-derived chemokine (CCL22) in atherosclerosis and

regulation by histamine via the H2 receptor. Pathol Int 62:675–683.

11.   Ryning FW, Remington JS. 1978. Effect of cytochalasin D on *Toxoplasma gondii* cell

entry. Infect Immun 20:739–743.

12.   Li L, Li X, Yan J. 2008. Alterations of concentrations of calcium and arachidonic acid and

agglutinations of microfilaments in host cells during *Toxoplasma gondii* invasion. Vet

Parasitol 157:21–33.

13.   Franco M, Panas MW, Marino ND, Lee M-CW, Buchholz KR, Kelly FD, Bednarski JJ,

Sleckman BP, Pourmand N, Boothroyd JC. 2016. A novel secreted protein, MYR1, is

central to *Toxoplasma*'s manipulation of host cells. MBio 7:e02231–15.

14.   Marino ND, Panas MW, Franco M, Theisen TC, Naor A, Rastogi S, Buchholz KR, Lorenzi
HA, Boothroyd JC. 2018. Identification of a novel protein complex essential for effector
translocation across the parasitophorous vacuole membrane of *Toxoplasma gondii*. PLoS
Pathog 14:e1006828.

15.   Olias P, Etheridge RD, Zhang Y, Holtzman MJ, Sibley LD. 2016. *Toxoplasma* Effector
Recruits the Mi-2/NuRD Complex to Repress STAT1 Transcription and Block IFN-γ-
Dependent Gene Expression. Cell Host Microbe 20:72–82.

16.   Braun L, Brenier-Pinchart M-P, Yogavel M, Curt-Varesano A, Curt-Bertini R-L, Hussain T,
Kieffer-Jaquinod S, Coute Y, Pelloux H, Tardieux I, Sharma A, Belrhali H, Bougdour A,
Hakimi M-A. 2013. A *Toxoplasma* dense granule protein, GRA24, modulates the early
immune response to infection by promoting a direct and sustained host p38 MAPK
activation. J Exp Med 210:2071–2086.

17.   Shastri AJ, Marino ND, Franco M, Lodoen MB, Boothroyd JC. 2014. GRA25 is a novel
virulence factor of *Toxoplasma gondii* and influences the host immune response. Infect
Immun 82:2595–2605.

18.   He H, Brenier-Pinchart M-P, Braun L, Kraut A, Touquet B, Couté Y, Tardieux I, Hakimi M-
A, Bougdour A. 2018. Characterization of a *Toxoplasma* effector uncovers an alternative
GSK3/β-catenin-regulatory pathway of inflammation. Elife 7.

19.   Lodoen MB, Gerke C, Boothroyd JC. 2010. A highly sensitive FRET-based approach
reveals secretion of the actin-binding protein toxofilin during *Toxoplasma gondii* infection.
Cell Microbiol 12:55–66.

20.   Delorme-Walker V, Abrivard M, Lagal V, Anderson K, Perazzi A, Gonzalez V, Page C,
Chauvet J, Ochoa W, Volkmann N, Hanein D, Tardieux I. 2012. Toxofilin upregulates the
host cortical actin cytoskeleton dynamics, facilitating *Toxoplasma* invasion. J Cell Sci
125:4333–4342.

21. Riezu-Boj J-I, Larrea E, Aldabe R, Guembe L, Casares N, Galeano E, Echeverria I,
Sarobe P, Herrero I, Sangro B, Prieto J, Lasarte J-J. 2011. Hepatitis C virus induces the
expression of CCL17 and CCL22 chemokines that attract regulatory T cells to the site of
infection. J Hepatol 54:422–431.

22. Jeong S-I, Choi B-M, Jang SI. 2010. Sulforaphane suppresses TARC/CCL17 and
MDC/CCL22 expression through heme oxygenase-1 and NF-κB in human keratinocytes.
Arch Pharm Res 33:1867–1876.

23. Wiley M, Teygong C, Phelps E, Radke J, Blader IJ. 2011. Serum response factor
regulates immediate early host gene expression in *Toxoplasma gondii*-infected host cells.
PLoS One 6:e18335.

24. Saeij JPJ, Coller S, **Boyle JP**, Jerome ME, White MW, Boothroyd JC. 2007. *Toxoplasma*
co-opts host gene expression by injection of a polymorphic kinase homologue. Nature
445:324–327.

25. Nadipuram SM, Kim EW, Vashisht AA, Lin AH, Bell HN, Coppens I, Wohlschlegel JA,
Bradley PJ. 2016. *In vivo* biotinylation of the *Toxoplasma* parasitophorous vacuole reveals
novel dense granule proteins important for parasite growth and pathogenesis. MBio 7.

26. English ED, Adomako-Ankomah Y, **Boyle JP**. 2015. Secreted effectors in *Toxoplasma*
gondii and related species: determinants of host range and pathogenesis? Parasite
Immunol 37:127–140.

27. Blank ML, Parker ML, Ramaswamy R, Powell CJ, English ED, Adomako-Ankomah Y,
Pernas LF, Workman SD, Boothroyd JC, Boulanger MJ, **Boyle JP**. 2018. A *Toxoplasma*
gondii locus required for the direct manipulation of host mitochondria has maintained
multiple ancestral functions. Mol Microbiol 108:519–535.

28. English ED, **Boyle JP**. 2018. Impact of Engineered Expression of Mitochondrial
Association Factor 1b on *Toxoplasma gondii* Infection and the Host Response in a Mouse
Model. mSphere 3.

29.   Hakimi M-A, Olias P, Sibley LD. 2017. *Toxoplasma* effectors targeting host signaling and

806   transcription. Clin Microbiol Rev 30:615–645.

807   30.   Lee CW, Sukhumavasi W, Denkers EY. 2007. Phosphoinositide-3-kinase-dependent,

808   MyD88-independent induction of CC-type chemokines characterizes the macrophage

809   response to *Toxoplasma gondii* strains with high virulence. Infect Immun 75:5788–5797.

810   31.   Hill RD, Gouffon JS, Saxton AM, Su C. 2012. Differential gene expression in mice infected

811   with distinct *Toxoplasma* strains. Infect Immun 80:968–974.

812   32.   Jensen KDC, Wang Y, Wojno EDT, Shastri AJ, Hu K, Cornel L, Boedec E, Ong Y-C,

813   Chien Y, Hunter CA, Boothroyd JC, Saeij JPJ. 2011. *Toxoplasma* polymorphic effectors

814   determine macrophage polarization and intestinal inflammation. Cell Host Microbe 9:472–

815   483.

816   33.   Rosowski EE, Lu D, Julien L, Rodda L, Gaiser RA, Jensen KDC, Saeij JPJ. 2011. Strain-

817   specific activation of the NF-kappaB pathway by GRA15, a novel *Toxoplasma gondii*

818   dense granule protein. J Exp Med 208:195–212.

819   34.   Lee JW, Zemojtel T, Shakhnovich E. 2009. Systems-level evidence of transcriptional co-

820   regulation of yeast protein complexes. J Comput Biol 16:331–339.

821   35.   Rosenberg A, Sibley LD. 2021. *Toxoplasma gondii* secreted effectors co-opt host

822   repressor complexes to inhibit necroptosis. Cell Host Microbe.

823   36.   Ong Y-C, Reese ML, Boothroyd JC. 2010. *Toxoplasma* rhoptry protein 16 (ROP16)

824   subverts host function by direct tyrosine phosphorylation of STAT6. J Biol Chem

825   285:28731–28740.

826   37.   Gay G, Braun L, Brenier-Pinchart M-P, Vollaire J, Josserand V, Bertini R-L, Varesano A,

827   Touquet B, De Bock P-J, Coute Y, Tardieux I, Bougdour A, Hakimi M-A. 2016.

828   *Toxoplasma gondii* TgIST co-opts host chromatin repressors dampening STAT1-

829   dependent gene regulation and IFN-γ-mediated host defenses. J Exp Med 213:1779–

830   1798.

831   38.   Gupta D, Wang Q, Vinson C, Dziarski R. 1999. Bacterial peptidoglycan induces CD14-
832   dependent activation of transcription factors CREB/ATF and AP-1. J Biol Chem
833   274:14012–14020.

834   39.   Hall AJ, Vos HL, Bertina RM. 1999. Lipopolysaccharide induction of tissue factor in THP-1
835   cells involves Jun protein phosphorylation and nuclear factor kappaB nuclear
836   translocation. J Biol Chem 274:376–383.

837   40.   Wan J, Shan Y, Fan Y, Fan C, Chen S, Sun J, Zhu L, Qin L, Yu M, Lin Z. 2016. NF-κB
838   inhibition attenuates LPS-induced TLR4 activation in monocyte cells. Mol Med Rep
839   14:4505–4510.

840   41.   Carruthers VB, Håkansson S, Giddings OK, Sibley LD. 2000. *Toxoplasma gondii* uses
841   sulfated proteoglycans for substrate and host cell attachment. Infect Immun 68:4005–
842   4011.

843   42.   Harper JM, Hoff EF, Carruthers VB. 2004. Multimerization of the *Toxoplasma gondii* MIC2
844   integrin-like A-domain is required for binding to heparin and human cells. Mol Biochem
845   Parasitol 134:201–212.

846   43.   Konradt C, Ueno N, Christian DA, Delong JH, Pritchard GH, Herz J, Bzik DJ, Koshy AA,
847   McGavern DB, Lodoen MB, Hunter CA. 2016. Endothelial cells are a replicative niche for
848   entry of *Toxoplasma gondii* to the central nervous system. Nat Microbiol 1:16001.

849   44.   Platt DJ, Smith AM, Arora N, Diamond MS, Coyne CB, Miner JJ. 2018. Zika virus-related
850   neurotropic flaviviruses infect human placental explants and cause fetal demise in mice.
851   Sci Transl Med 10.

852   45.   Dubey JP, Tilahun G, **Boyle JP**, Schares G, Verma SK, Ferreira LR, Oliveira S, Tiao N,
853   Darrington C, Gebreyes WA. 2013. Molecular and biological characterization of first
854   isolates of *Hammondia hammondi* from cats from Ethiopia. J Parasitol 99:614–618.

855   46.   Dubey JP, Sreekumar C. 2003. Redescription of *Hammondia hammondi* and its
856   differentiation from *Toxoplasma gondii*. Int J Parasitol 33:1437–1453.

857   47.   Sokol SL, Primack AS, Nair SC, Wong ZS, Tembo M, Verma SK, Cerqueira-Cezar CK,

858   Dubey JP, **Boyle JP**. 2018. Dissection of the *in vitro* developmental program of

859   *Hammondia hammondi* reveals a link between stress sensitivity and life cycle flexibility in

860   *Toxoplasma gondii*. Elife 7.

861   48.   Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, Marshall KA,

862   Phillippy KH, Sherman PM, Holko M, Yefanov A, Lee H, Zhang N, Robertson CL, Serova

863   N, Davis S, Soboleva A. 2013. NCBI GEO: archive for functional genomics data sets--

864   update. Nucleic Acids Res 41:D991–5.

865   49.   Gautier L, Cope L, Bolstad BM, Irizarry RA. 2004. affy--analysis of Affymetrix GeneChip

866   data at the probe level. Bioinformatics 20:307–315.

867   50.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,

868   Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, di Palma

869   F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A. 2011. Full-length

870   transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol

871   29:644–652.

872   51.   Smith AF, Posakony JW, Rebeiz M. 2017. Automated tools for comparative sequence

873   analysis of genic regions using the GenePalette application. Dev Biol 429:158–164.

874   52.   Shen B, Brown KM, Lee TD, Sibley LD. 2014. Efficient gene disruption in diverse strains

875   of *Toxoplasma gondii* using CRISPR/CAS9. MBio 5:e01114–14.

876   53.   Saeij JPJ, **Boyle JP**, Coller S, Taylor S, Sibley LD, Brooke-Powell ET, Ajioka JW,

877   Boothroyd JC. 2006. Polymorphic secreted kinases are key virulence factors in

878   toxoplasmosis. Science 314:1780–1783.

879   54.   Coombs RS, Blank ML, English ED, Adomako-Ankomah Y, Urama I-CS, Martin AT,

880   Yarovinsky F, **Boyle JP**. 2020. Immediate Interferon Gamma Induction Determines

881   Murine Host Compatibility Differences between *Toxoplasma gondii* and *Neospora*

882   *caninum*. Infect Immun 88.

883   55.   Huynh M-H, Carruthers VB. 2009. Tagging of endogenous genes in a *Toxoplasma gondii*

884   strain lacking Ku80. Eukaryotic Cell 8:530–539.

885   56.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and' ' dispersion

886   for RNA-seq data with DESeq2. Genome Biol 15:550.

887   57.   Croken MM, Qiu W, White MW, Kim K. 2014. Gene Set Enrichment Analysis (GSEA) of

888   *Toxoplasma gondii* expression datasets links cell cycle progression and the bradyzoite

889   developmental program. BMC Genomics 15:515.

890   58.   Krämer A, Green J, Pollard J, Tugendreich S. 2014. Causal analysis approaches in

891   Ingenuity Pathway Analysis. Bioinformatics 30:523–530.

892   59.   Saeij JPJ, **Boyle JP**, Grigg ME, Arrizabalaga G, Boothroyd JC. 2005. Bioluminescence

893   imaging of *Toxoplasma gondii* infection in living mice reveals dramatic differences

894   between strains. Infect Immun 73:695–702.

FIGURE LEGENDS:

Figure 1: A) THP-1 cells, Human Foreskin Fibroblasts (HFFs), and 2nd trimester placental samples of villous trees and decidual tissue were infected with a type I strain of *Toxoplasma gondii* (RH:YFP). Statistics on the THP-1 cell and HFF cell samples performed by ordinary Two-way ANOVA analysis with Tukey's multiple comparisons test. Statistics on the placental samples performed by two-tailed welch-corrected t-tests. B) Type I strain (RH) *T. gondii* parasites subjected to multiple treatments as described in methods. The soluble fraction of the sonicated treatment is denoted by S, the insoluble fraction is denoted by P. THP-1 cells were then exposed to either live parasites, or treated parasites. Statistics performed by multiple two-tailed welch-corrected t-test comparisons with the live parasite treatment $P \leq 0.0145$. C,D) Type I strain (RH) *T. gondii* parasites were treated with either Cytochalasin D (C), or 4-Bromophenacyl bromide (D) as described in methods. THP-1 cells were then infected with the respective parasite treatment. Statistics were performed by two-way ANOVA analysis and multiple comparisons post-hoc tests, where Cyt-D $P = 0.009$, and 4-BPB $P < 0.0001$. E) Type I strain (RH) *T. gondii* parasites deficient in Myr-1 (TgRHΔMyr-1) and their complement (TgRHΔMyr-1:Myr-1$_{comp}$) were used to infect THP-1 cells. (F) Type I strain (RH) *T. gondii* parasites were used to infect THP-1 cells at MOIs of: 20, 10, 8, 6, 4, 3, 1, 0.8, 0.4, 0.2, and 0.1. A, B, C, D) Respective cells/tissues were infected with a multiplicity of infection (MOI) of 3. Supernatants were collected at 24 hours post-infection unless indicated otherwise, and assayed by CCL22 ELISA.

Figure 2: Transcript correlation analyses to identify putative MYR1 substrates based on 396 publicly available microarray datasets. A) Gene-by-gene correlation data for *T. gondii* genes across 396 microarray datasets. A subset (3,217 genes with at least one sample having a normalized log2-transformed value ≥10) of the total number (8,058) of *T. gondii* genes are

shown for simplicity. Genes outlined in the green box indicate the cluster containing all of the

bait genes as well as candidate CCL22-inducing genes with the exception of *Toxofilin*. Dark

blue tick marks on each dendrogram indicate the location of all of the bait genes. Color scale

covers correlations ranging from -1 to +1. B) Subcluster containing *MYR2* and *GRA28*. C)

Subcluster containing *MYR1*, *TgIST*, *GRA18* and *GRA24*. Note: For B and C the color scale is

from 0.5 to 1.0 to highlight subcluster differences.

Figure 3: Identification of the *T. gondii* gene GRA28 as an inducer of CCL22 in human cells. A)

Gene expression correlations across 396 *T. gondii* expression microarrays between TgMYR1

and 4 additional "bait" genes (top) and 5 candidate CCL22-inducing effectors (bottom). dN/dS

ratios are also shown to illustrate the high level of positive selection acting on this class of

genes. B) Effect of deleting 5 candidate genes on CCL22 secretion in THP-1 cells, showing that

ΔGRA28 parasites induced significantly less CCL22 compared to wild type controls (>100 fold

reduction; ***P<0.0001). PRUΔ*Toxofilin* parasites also induced significantly lower levels of

CCL22 in THP-1 cells (1.4-fold reduction; **P<0.01). Each blue dot indicates a genetically

distinct knockout clone. C) *ΔGRA28* parasite clones also induce significantly less CCL22 from

primary human 2nd trimester placental villous explants. D) MA plot of RNAseq analysis

performed on THP-1 cells infected with WT or *ΔGRA28 T. gondii* (RH strain). CCL22 and the

chemokines CXL1 and CXL2 were the most highly GRA28-dependent transcripts, while a

handful (64) of other genes had significantly higher transcript abundance in WT parasites

compared to *ΔGRA28* parasites (Padj<0.001; Log2FC>2; blue symbols). CCL17 (arrow, green

symbol), a chemokine that is typically co-regulated with CCL22, did not show any evidence of

being induced by GRA28.

Figure 4. IPA analysis on THP-1 cells infected with WT or *ΔGRA28 T. gondii* parasites for 24h

showing canonical pathways that were differentially regulated (-log(*P*)≥2; Z score ≤ -2 or ≥ 2)

depending on the presence or absence of the GRA28 gene. A) Z scores for significant canonical

pathways. All were higher in WT compared to *ΔGRA28 T. gondii.* B) There was extensive

overlap of component genes within each canonical pathway, particularly for the genes *KL* and

those encoding components of the AP-1 transcription factor complex (*JUN* and *FOS*). A

heatmap of fold-difference in transcript abundance for cells infected with RH:WT and

RH*ΔGRA28* is shown. C) GRA28 is responsible for driving transcriptional changes in two major

gene clusters identified based on the degree of gene sharing between each canonical pathway

(clusters outlined in dotted green boxes). The larger cluster consists primarily of immunity-

related pathways while the smaller cluster consists of genes involved in proteoglycan synthesis.

D) Quantification of NFκB activation in 293T cells. Cells were transfected with NFκB firefly

luciferase plasmid, a consitutive renilla luciferase plasmid, as well as empty vector (EV), a

construct expressing a CARD domain (dsCARDS) or the first exon of *T. gondii* GRA28. While

the CARD domain construct induced firefly luciferase expression as expected, expression of *T.*

gondii GRA28 had no significant impact on firefly luciferase levels (letters indicate groups that

were not significantly different from one another following one way ANOVA and Tukey's multiple

comparisons post-hoc test.)

Figure 5: A) Schematic of the *GRA28* locus along with its gene prediction in the current

annotation of the *T. gondii* genome (www.toxodb.org). In addition to gene models from two *T.*

gondii strains (GT1 and ME49), 39 bp repeats and regions used in expression constructs are

shown in brown and light green. Map created using GenePalette software (see Materials and

Methods; (51)). B) Sequence encoding an N-terminal HA tag was inserted immediately after the

predicted signal peptide cleavage site in a GRA1-promoter driven version of *T. gondii* GRA28

Exon 1. When transiently transfected into *T. gondii* HA-tagged protein could be detected in the

parasites as well as the host cell nucleus. C) *ΔGRA28 T. gondii* parasites (RH strain) were

transiently transfected with empty pGRA-HA-HPT vector (EV) or the same construct described

in B encoding an HA-tagged version of *T. gondii* GRA28 Exon 1. After washing in cDMEM

parasites were used to infect freshly plated THP-1 cells for 24h and CCL22-levels were

quantified in culture supernatants using ELISA. Mock-treated cells were exposed to a sterile

filtered parasite preparation. D) E) The construct encoding HA-tagged *T. gondii* GRA28 Exon 1

(same as that used in B and C) was used to transfect *Neospora caninum*, a near relative of *T.*

gondii. HA staining revealed expression of this *T. gondii* GRA28 Exon 1 in *N. caninum* parasites

(visualized by HA staining) but in contrast to *T. gondii* we did not observe trafficking of GRA28

to the host cell nucleus when expressed in this strain. Quantification of nuclear HA-derived

signal is presented in **Figure 5** Supplementary **Figure 1**. F) Sequences encoding an N-terminal

V5 tag were inserted downstream of a Kozak consensus sequence and upstream of GRA28

Exon 1 (minus the signal peptide-encoding sequence). The construct was transfected into HeLa

cells and V5 staining was observed prominently in the nucleus of transfected cells. G)

Transfection of the construct in E directly into RAW 264.7 cells significantly induced CCL22

production as detected by ELISA. T-test was performed on $\log_{10}$-transformed data.

Figure 6. GRA28 induction of Ccl22 is conserved in mouse immune and placental tissues. A,B)

ΔGRA28 parasites induce significantly less Ccl22 secretion from RAW 264.7 macrophages (A)

and mouse placental explants (B) compared to wild type parasites. C) The number of host

genes besides Ccl22 that are GRA28-dependent in placental explants is relatively small,

suggesting that GRA28 is a highly specific inducer of Ccl22 in mouse placental tissue. D,E)

Mouse serum (D) and peritoneal lavage (E) levels of Ccl22 48 h-post-infection are dependent

on GRA28.

Figure 7. A) Mortality and B) weight loss does not significantly differ in mice infected with WT

and ΔGRA28 type 1 strain *T. gondii* parasites. C) Behavioral changes and fur ruffling

phenotypes associated with infection are exacerbated during the acute phase of infection for

mice infected with ΔGRA28 parasites compared to WT based on a phenotype scoring system. Specifically, mice infected with ΔGRA28 parasites (light grey lines and triangles) exhibited infection-related symptoms at earlier time points compared to those infected with WT parasites (black lines and circles). Curves beneath lines are the average across all mice at that time point (with lighter grey representing Δ*GRA28* knockout parasites and darker grey representing wild type). *: P<0.05 after Two Way ANOVA and followed by multiple comparisons at each time point.

Figure 8: Impact of the *GRA28* gene on proliferation and dissemination of Type II parasites expressing luciferase. Neither A) mortality nor B) mouse weight loss were significantly different in mice infected with WT or *ΔGRA28 T. gondii* Type II (ME49) strain parasites. C) Throughout the acute phase of infection parasite-derived bioluminescence of WT and *ΔGRA28* parasites in the peritoneal cavity was measured by imaging the animals ventrally. Burden was similar (P>0.05) between parasite strains at all time points except for D9 post-infection, when signal was significantly higher in *ΔGRA28*-infected mice compared to WT mice (*P*=0.014). D,E) Starting on D10 post-infection we imaged mice both ventrally and dorsally to visualize dissemination to and proliferation in the mouse brain. In contrast to the acute phase we observed a consistently lower level of parasite-derived bioluminescence in the brains of mice infected with *ΔGRA28* parasites compared to WT parasites on days 14 and 15 post-infection (*P*=0.016 and 0.0002, respectively). All statistical tests were performed on log2-transformed bioluminescent data.

SUPPLEMENTAL FIGURE LEGENDS:

Figure 1 Supplement 1: A) Type 1 (Rh88), Type 2 (Pru) and Type 3 (CTG) *T. gondii* strains all induce CCL22 in THP-1 cells. For mock cells were treated with 0.2 micron-filtered parasite suspensions. B) *H. hammondi* induces CCL22 production in THP-1 cells while *N. caninum* does

not when compared to mock-treated cells. C) The *T. gondii* gene MYR1 is required for CCL22

production by THP-1 cells after infection. *T. gondii* ΔMYR1 parasites were compared to

ΔMYR1:MYR1-complemented parasites and only the MYR1-complemented parasites induced

secretion of CCL22 from wild type (black data points; left) or MYD88-knockout (blue data points;

right) THP-1 cells. While MYD88 knockout cells produced significantly less CCL22 in response

to MYR1-complemented parasites, they still produced CCL22 at levels much greater than

ΔMYR1-infected cells or mock-treated cells. MYR1-knockout and complemented parasites were

kindly provided by John Boothroyd and Michael Panas, Stanford University.

1035

Figure 2 Supplement 1: Clusters of co-regulated genes share developmental expression

profiles and functional activities. A) Transcript abundance correlation analysis (left) and

clustered transcript abundance analysis (in RMA log2-normalized units) for 21 genes with

transcripts known to increase in abundance during the tachyzoite to bradyzoite transition,

including BAG1, LDH2 and enolase. Bar across the top of the expression heatmap indicates the

life cycle stage source for each of the samples. Pie chart indicates that 19 of the 21 transcripts

increase in abundance during pH-induced bradyzoite development according to the Bradyzoite

Differentiation (3-day Time Series) dataset on ToxoDB.org. B) Cluster containing multiple

ribosomal protein coding genes (for both the large and small subunits) showing high transcript

abundance in tachyzoites and bradyzoites but comparatively low transcript abundance in

samples taken from sporozoites and merozoites.

1047

Figure 3 Supplementary Figure 1: Transcript level correlation analysis with the 6 bait genes

used in this study (A) and clustered, normalized gene expression data (B) for all genes in the *T.*

gondii genome annotated with the term "dense granule" in the product name or user comments.

Dense granule protein coding genes fall into two major clusters in the correlation analysis, with

the top cluster containing the known secreted effectors GRA15, GRA24, TgIST, GRA7 and

GRA28.

1054

Figure 3 Supplementary Figure 2: Validation of knockouts generated for the present study (A-

D) of for GRA18 which was generated in another study (E; He *et al.*, eLife 2018;7:e39887).

Clones that were validated as knockouts and used in the CCL22-induction assays shown in

Figure 3b are labled with ***bold-italic font***. We validated using two paralell approaches for most

of the knockouts: (1) amplifying across the site targeted by the protospacer(s) encoded by the

transfected gRNA-expressing plasmid(s) where no amplification indicated a potential insertion of

plasmid sequence into that location and (2) amplifying across the entire coding sequence of the

gene, where no amplification also suggests insertion of the selectable marker and other plasmid

sequence at at least one of the protospacer sites. In some cases multiple protospacer encoding

plasmids were transfected into the same parasite population (and protospacer numbers are our

own internal nomenclature), and in these cases it was possible to have insertion/disruption at

both protospacer sites (and this ocurred in some instances). Primer sequences and gRNA

target sites are shown in **Table S1**. A) Validation of four TGGT1_201390 knockout clones

generated by batch transfection with gRNAs targeting two distinct sites in the TGGT1_201390

coding sequence (2 and 12). Left gel: Two primer sets (A and B) were used to amplify across

the site targeted by the protospacer, and MAF1b primers were used as a positive control.

Clones D11 and G11 likely had insertions at protospacer site 2 (as evidenced by the lack of

amplification with primer set A), while clone D2 likely had an insertion in protospacer site 12 as

evidenced by a lack of amplification with primer set B. Clone F2 had amplification of the correct

size at both protospacer sites (2 and 12), but we could not amplify the coding sequence from

the F2 clone (right gel, lane labeled F2), suggesting that the locus was disrupted in this strain as

well. All four of these knockout clones were assayed in biological triplicate for CCL22 induction

in THP-1 cells (**Figure 3b**). B) Validation of 3 GRA4 knockout clones out of 5 tested. Clones

A1:D6 and B2:C1 (where A1 and B1 indicate the parasites were from distinct transfections) had

a likely insertion in the GRA4 gene at protospacer site 0, while clone B2:B11 had a likely

insertion at both sites (0 and 26). C) Validation of 4 GRA8 knockout clones. Parasites were

transfected with a single gRNA expressing plasmid (targeting gRNA sequence 4) and PCR on

all 4 clones failed to amplify across the gRNA 4 target site. All amplifications across the gRNA

target site 27 worked as did the positive control amplification of MAF1b. D) Validation of 4

GRA28 knockout clones. Parasites were co-transfected with plasmids encoding gRNAs

targeting sites 5 and 26 and queried using primers targeting the entire locus (sets A and B in

this case), or flanking gRNA target sites 5 and 26 (sets C and D). PCR across the entire GRA28

locus for clones 1A4, 1D4 and 1D3 all failed to generate PCR products, suggesting that these 3

clones had disruptions in the *GRA28* gene. PCR on clone F3 with primer sets A and B gave a

product of the expected size. PCR across the gRNA target site 26 failed for clone 1A4, while

PCR across gRNA target site 5 failed for clones 1D4 and 1D3. Taken together these data

indicate that clones 1A4, 1D4 and 1D3 were all GRA28 knockouts via insertion of selectable

marker and accompanying plasmid sequences at the targeted gRNA sites. For clone 1F3 the

locus seemed to be intact, but when we sequenced PCR products similar to those amplified by

primer sets C and D (at gRNA target sits 5 and 26, respectively) we determined that clone 1F3

had a single base pair deletion at base 250 relative to the start codon of GRA28 which

introduced a stop codon 100 bp downstream of the indel (as well as multiple stop codons in

frame further downstream). Sequences across the gRNA 26 target site were identical to wild

type. The deletion near gRNA target site 5 was within the gRNA protospacer sequence itself,

just proximal to the PAM site (GTTCCGCTGGTGCCTT**C**ACC [TGG] was mutated to

GTTCCGCTGGTGCCTT_ACC [TGG]). Therefore we treated 1F3 as a GRA28-null parasite

strain. E) Validation of GRA18 knockouts received from the Bougdour lab: We received GRA18

knockout and wild type clones from the laboratory of Alexandre Bougdour and validated them

using PCR. In this case we generated primers to amplify across the entire GRA18 locus, which

was completely deleted using double homologous recombination with large sequences flanking

the entire coding region. As expected we could amplify across the entire GRA18 locus in

PRU:WT and GRA18-complemented strains, but failed to do so in the GRA18 knockout.

1107

Figure 4 Supplementary Figure 1: Ingenuity pathway analysis of THP-1 cells infected with

RH:WT or RH*ΔGRA28 T. gondii* identifies candidate upstream regulatory gene products that

may be driving GRA28-dependent differences in transcript abundance. A) Genes of higher

abundance in RH:WT-infected THP-1 cells were found to be significantly associated with

multiple immunity-related regulatory genes, including those related to the NFκB pathway (e.g.,

NFκBIA, NFκB1 and REL). B) Hierarchical cluster of correlations in the amount of target gene

overlap for each of the upstream regulators shown in A. A small cluster of the most highly

correlated genes contained multiple genes relevant to NFkB activation (outlined in dotted

green). C) As in B, most of the upstream regulators had the same downstream targets, and this

was most evident for the AP-1 transcription factor complex (encoded by *FOS* and *JUN* genes)

as well as *IL1B* and *ICAM1*. A heat map indicating fold-differences between RH:WT and

RH*ΔGRA28*-infected THP-1 cells for these downstream targets is also shown. D) Heatmap

showing transcript abundance for all transcriptional regulators that were found to be significantly

altered in infected THP-1 cells in a GRA28-dependent manner (P<0.001; $\log_2$(fold-

difference)≥1).

1123

Figure 4 Supplementary Figure 2: C-JUN protein levels are induced by *T. gondii* infection in

THP-1 cells but do not depend on the presence or GRA28 in the infecting strain. THP-1 cells

were infected with the indicated strains (or mock-treated by exposing them to the same WT

parasite suspensions as for infection but after sterile-filtering with a 0.2 µm filter) for 24 h and

then C-JUN protein level was quantiifed using western blotting. Histone H3 levels served as a

control and densitometry was used to calculate the C-JUN/Histone H3 ratio as a proxy for

normalized C-JUN abundance. Two replicates, with an N=3 wells of cells for infections and N=2

wells for mock, are shown, each having similar results.

1132

Figure 5 Supplementary Figure 1: A) Quantification of nuclear localization in *T. gondii* (N=5)

and *N. caninum* (N=3) vacuoles expressing Exon 1 of *T. gondii* HA-tagged GRA28. Data were

normalized for each image and HA-positive vacuole by subtracting mean HA-intensity in a

nucleus neighboring the infected cell from the mean GFP intensity in the nucleus of the infected

cell, and then dividing that by the mean HA-intensity of the parasite-containing vacuole.

Expression in *T. gondii* led to significantly ($P<0.01$; T-test on the log10-normalized data) higher

normalized intensity in the nucleus compared to when GRA28 was expressed in *N. caninum*.

B,C) Schematic illustrating how data in A were collected and calculated for *T. gondii* (B) and *N.*

caninum (C).

1142

Figure 7 Supplementary Figure 1: Visual representation of pathology index scores. All images

are the same individual from two viewpoints (lateral and dorsal) at four different timepoints of

infection. **0)** No fur ruffling or red/irritated skin present. **1)** Mild ruffling present predominantly

located on the head and back of the neck. No red/irritated skin visible. **2)** Moderate ruffling

present - fur forms larger clumps and extends to the rest of the body. Skin may be visible

through the clumps but is not red or irritated. **3)** Severe ruffling is characterized by ruffling

across the entire body with visibly red/irritated skin.

1150

A

Expression Correlation

0.0   0.5   1.0

99th %ile

Gene	dN/dS
TgIST	7.1
TgMYR3	3.3
TgMYR1	3.0
TgGRA24	2.7
TgMYR2	2.0
201390	6.6
TgGRA4	3.8
TgGRA8	2.4
TgGRA28	7.2
TgTfilin	3.6

Columns: TgMYR1, TgGRA24, TgIST, TgMYR3, TgMYR2

B Knockouts screened in THP-1 cells

CCL22-induction, Mean WT/KO ratio

Gene name: Tg201390, TgGRA4, TgGRA8, TgGRA28 (***), TgGRA18, TgTfilin (**)

C Human Placental Explants

CCL22 (pg/mL +/- SD)

Strain: Mock, RhΔHXGPRT, RhΔGRA28 1F3, RhΔGRA28 1A4

0.008, 0.013, 0.009

D

Log2(Fold Difference) WT versus KO

Log10 (normalized counts)

CCL22, XCL1, XCL2, CCL17

A
TGME49_231960
◻ 39 bp repeat: CAGCAGCA..AACCACTA
TGGT1_231960

Exon 1 Expression Construct
Genomic Expression Construct

Select Type I (RH)
tachyzoite Nanopore
RNAseq reads

B
GRA1 5' | SP | GRA28-Ex1 | HA

T. gondii (RH)
PHASE

GRA28-HA

GRA28-HA DAPI

C
Complementation:
Exon 1
***
CCL22 (pg/mL)
Mock ΔGRA28:EV ΔGRA28:GRA28

D
Complementation:
Exon1 and Genomic
CCL22 (pg/mL)
Mock WT ΔGRA28 ΔGRA28:Exon1 ΔGRA28:Genomic

E
GRA1 5' | SP | GRA28-Ex1 | HA

N. caninum (NC-1)
PHASE

GRA28-HA

GRA28-HA DAPI

F
CMV 5' | V5 | GRA28-Ex1

ACTIN

GRA28-V5 DAPI

G
P=0.0018
CCL22 (pg/mL)
EV GRA28-V5
Construct