Title

Novel epigenetic clock for fetal brain development predicts prenatal age for cellular stem cell models

and derived neurons.

Authors

Leonard C. Steg[1], Gemma L. Shireby[1], Jennifer Imm[1], Jonathan P. Davies[1], Alice Franklin[1], Robert

Flynn[1], Seema C. Namboori[1], Akshay Bhinge[1], Aaron R. Jeffries[1], Joe Burrage[1], Grant W. A. Neilson[1],

Emma M. Walker[1], Leo W. Perfect[2], Jack Price[2], Grainne McAlonan[3,4], Deepak P. Srivastava[2,4],

Nicholas J. Bray[5], Emma L. Cope[6], Kimberly M. Jones[6], Nicholas D. Allen[6], Ehsan Pishva[1,7], Emma L.

Dempster[1], Katie Lunnon[1], Jonathan Mill[1], & Eilis Hannon[1,*]

[1] University of Exeter Medical School, College of Medicine and Health, University of Exeter, Exeter,

UK.

[2] Department of Basic and Clinical Neuroscience, Institute of Psychiatry, Psychology and

Neuroscience, King's College London, UK.

[3] Department of Forensic and Neurodevelopmental Sciences, King's College London, London, UK.

[4] MRC Centre for Neurodevelopmental Disorders, King's College London, London, UK.

[5] Division of Psychological Medicine and Clinical Neurosciences, School of Medicine, Cardiff

University, Cardiff, UK.

[6] School of Biosciences, Cardiff University, Cardiff, UK.

[7] Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience

(MHeNS), Maastricht University, Maastricht, The Netherlands.

* Corresponding author: Eilis Hannon, University of Exeter Medical School, RILD Building Level 3,

Royal Devon and Exeter Hospital, Barrack Rd, Exeter. EX2 5DW. UK. E-mail: e.j.hannon@exeter.ac.uk

Keywords

Epigenetic clock, DNA methylation, fetal, neurodevelopment, induced pluripotent stem cells, iPSC-

derived neurons, neuronal precursor cells, DNAm clock

Abstract

Induced pluripotent stem cells (iPSCs) and their differentiated neurons (iPSC-neurons) are a widely

used cellular model in the research of the central nervous system. However, it is unknown how well

they capture age-associated processes, particularly given that pluripotent cells are only present

during the earliest stages of mammalian development. Epigenetic clocks utilize coordinated age-

associated changes in DNA methylation to make predictions that correlate strongly with

chronological age. It has been shown that the induction of pluripotency rejuvenates predicted

epigenetic age. As existing clocks are not optimized for the study of brain development, we

developed the fetal brain clock (FBC), a bespoke epigenetic clock trained in human prenatal brain

samples in order to investigate more precisely the epigenetic age of iPSCs and iPSC-neurons. The FBC

was tested in two independent validation cohorts across a total of 194 samples, confirming that the

FBC outperforms other established epigenetic clocks in fetal brain cohorts. We applied the FBC to

DNA methylation data from iPSCs and iPSC-derived neuronal precursor cells and neurons, finding

that these cell types are epigenetically characterized as having an early fetal age. Furthermore, while

differentiation from iPSCs to neurons significantly increases epigenetic age, iPSC-neurons are still

predicted as being fetal. Together our findings reiterate the need to better understand the

limitations of existing epigenetic clocks for answering biological research questions and highlight a

limitation of iPSC-neurons as a cellular model of age-related diseases.

MAIN TEXT

Introduction

Induced pluripotent stem cells (iPSCs) offer a unique cellular system to investigate disease in human-

derived cells. iPSCs are obtained by treating skin or blood cells with a set of core pluripotency

transcription factors that reprogram the cells to a pluripotent state [1]. Established iPSC lines have

the capacity to be further differentiated into specific cell types, including neurons, when treated

with the appropriate factors [2–4]. This is of particular interest for neuroscience, as the only

alternative cellular model for human neurons are immortalized cell lines. Because immortalized cell

lines retain some physiological properties of the cancerous cells they were derived from [5] they do

not fully recapitulate the neuronal phenotype. iPSC-derived neurons (iPSC-neurons), on the other

hand, express appropriate morphological and neurophysiological properties of neurons and subject

to different protocols can be differentiated into a wide range of specific neuronal subtypes [6]. iPSCs

and their neuronal derivatives have been widely used to research disorders of the central nervous

system, including developmental disorders such as autism and schizophrenia and age-related

diseases such as Alzheimer's disease (AD) and Parkinson's disease. However, the extent to which

iPSCs and especially iPSC-neurons capture age-associated processes is not known, which is

fundamental to the study of age-related diseases. Of specific relevance is the fact that pluripotent

cells only occur during the early stages of mammalian development and the effect of differentiation

from iPSCs towards neurons on the developmental or aging trajectory of the cellular model [7] has

yet to be adequately profiled.

Epigenetic mechanisms, such as DNA methylation (DNAm), are chemical processes that stably

regulate gene expression, and while they are sensitive to environmental stimuli they also underpin

key developmental processes [8–10]. There has been much interest and success in capitalizing on

these patterns of epigenetic variation to derive individual age predictions from a biological sample.

Age predictors based on DNAm, known as "epigenetic clocks" or "DNAm clocks", are widely used to

predict the "epigenetic age" of a sample. Epigenetic age, defined here as age predicted by an

epigenetic clock, correlates strongly with chronological age, albeit not perfectly, and it has been

hypothesized that the deviations from this prediction, referred to as age acceleration, are

meaningful in the context of disease [11, 12]. The most well-known epigenetic clock is the Horvath

multi-tissue clock (MTC) which was developed using a large number of samples (n > 8000) from 51

different tissues and cell types [13]. Overall, the MTC generates reliable predictions of chronological

age for most sample-types, although there are potential biases when using Horvath's clock in

samples derived from certain tissues, especially the brain [14, 15]. To this end, a number of new

DNAm clocks have been developed for specific tissue types, including whole blood [16] and cortex

[14], which demonstrate more accurate predictions within the specified tissue. A less established

refinement of epigenetics clocks is the application to specific developmental stages, with prenatal

samples excluded or underrepresented in most training datasets. Recently, clocks were developed to

predict gestational age (GA) of newborns, derived using pre- and perinatal DNAm data from blood

samples [17] or placental samples [18]. While existing epigenetic clocks have been shown to

accurately predict age in either postnatal brain samples (predominantly middle and older age) or

non-brain prenatal samples, these tools have not been thoroughly tested on prenatal brain samples,

and it is unknown whether they are able to delineate the earliest stages of brain development.

Previous analysis applying the MTC to DNAm data generated from iPSCs and their corresponding

primary cells from adult donors found that the induction of pluripotency reversed the aging process,

with iPSCs predicted as postnatal, with an epigenetic age close to zero (i.e. birth) [13]. As human

pluripotent cells only occur during prenatal development, we hypothesize that existing clocks are

not sensitive enough to accurately predict iPSCs at prenatal developmental stages. The inability to

accurately estimate age during this crucial stage of neurodevelopment limits our ability to profile

changes in epigenetic age induced by the differentiation of iPSCs into specific cell-types using

already established DNAm clocks. Here we present a novel DNAm clock developed using prenatal

brain samples that accurately predicts fetal age, outperforming other DNAm clocks in

neurodevelopmental samples. We then apply our clock to iPSCs, iPSC-derived neuronal progenitor

cells (NPCs) and iPSC-derived neurons, as well as in other cellular stem cell models and derived

neuronal cells, to characterize the epigenetic age of these cellular models before and during the

differentiation process.

Methods

6   All statistical analyses were performed using R version 3.5.2 (https://www.r-project.org/) [19].

Development of Fetal Brain Clock (FBC)

Description of fetal brain samples

To develop and profile the performance of the fetal brain clock (FBC), we collated a dataset of 258

fetal brain samples (see **Table S1**) of which 194 were processed by our group at the University of

Exeter as described previously [20] and 64 were a subset (age < 0 years) of a publicly available

dataset downloaded from the Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/;

GSE74193) [21]. Of the samples processed in Exeter, 154 overlap those included in [20] following

additional outlier filtering by principal component analysis, where DNAm was quantified using the

Illumina 450K DNA methylation array. The other 40 samples represent additional samples where

DNAm was quantified using the Illumina EPIC DNA methylation array using a standard protocol as

previously described [14].

Data pre-processing and quality control.

All datasets for which raw data was available were pre-processed following a standard quality

control (QC) and normalization pipeline as described before [14] using either the R package

wateRmelon [22] or *bigmelon* [23]. Briefly, samples with low signal intensities or incomplete bisulfite

conversion were excluded prior to applying the *pfilter*() function from the *wateRmelon* package,

excluding samples with >1 % of probes with a detection P value >0.05 and probes with >1 % of

samples with detection P value >0.05. This was followed by the exclusion of probes known to be

affected by SNPs or known to cross-hybridize [24]. QC was finished by quantile normalization using

the *dasen*() function of the packages *wateRmelon* or *bigmelon* [22, 23]. To harmonize the age

variable across datasets, age was converted into days post-conception (dpc), as it represents the

most precise unit of age available across the datasets. Where age was provided as weeks post-

conception it was transformed to days post-conception by dividing by 7, and where age was

reported in (negative) years it was transformed to days post-conception by multiplying by 365 and

adding 280. Of note, a few samples (15 out of 258) are actually defined as embryonic (GA < 63 dpc)

and not fetal.

Fetal brain clock development.

To create two separate datasets for the purpose of training and testing the FBC, 75% of the samples

from each dataset were randomly assigned into a training dataset (n = 193, age range = 37-184 dpc,

age median = 99 dpc), while the remaining 25% included in the testing dataset (n = 65, age range =

23-153 dpc, age median = 99 dpc) (**Figure S1, Table S1**). There was no overlap in samples between

the training and testing dataset. To simplify the FBC development, only probes available in all

samples after QC were taken forward (n = 385,069 probes). To develop the fetal brain clock we

applied an elastic net (EN) regression model, using the approach described by Horvath [13],

regressing chronological age against DNAm level of all available probes. The EN algorithm selects a

subset of DNA methylation probes that together produce the optimal prediction of the outcome, in

this case chronological age, by combining ridge and LASSO (Least Absolute Shrinkage and Selection

Operator) regression. Briefly, ridge regression penalizes the sum of squared coefficients while LASSO

penalizes the sum of the absolute values of coefficients. EN is a combination of both methods,

where the user specifies the extent of the mixing of the two methods as a number between 0 and 1,

in our application this was set to 0.5 [25]. EN was implemented with the R package *GLMnet* [26].

The shrinkage parameter lambda was calculated using 10-fold cross-validation on the data, which

resulted in a lambda of 3.27.

Statistical evaluation of FBC performance

To profile the performance of the FBC, we additionally tested three established DNAm clocks:

Horvath's multi-tissue clock (MTC) [13], Knight's Gestational Age clock (GAC) [17] and Lee's Control

Placental epigenetic clock (CPC) [18]. The clocks were applied using the *agep()* function of the

wateRmelon package [22], where the default estimates the MTC and other clocks (here the GAC and

CPC) can be estimated by providing the necessary coefficient and intercept values. The predictive

accuracy of each clock was profiled in each dataset by two measures: Pearson's correlation

coefficient with reported chronological age and root mean squared error (RMSE). To investigate

potential effects of sex on the predicted epigenetic age, linear models were fitted in the testing and

validation datasets with FBC predicted epigenetic age as dependent variable, chronological age and

sex as main effects and an interaction of chronological age and sex.

Validation of the Fetal Brain Clock (FBC) in additional fetal datasets and adult cortex

To further test the FBC, we used data from 96 additional fetal brain samples currently being

assessed by our group (unpublished data), none of which overlapped with either the training data or

test data described in the previous section, with DNAm quantified using the Illumina EPIC DNA

methylation array. QC and normalization were performed as described above. We also included

data from 33 fetal samples from two publicly available datasets on GEO (GSE116754 and GSE90871)

[27, 28], where DNAm was quantified using the Illumina 450K DNA methylation array. Pre-processing

and QC for the publicly available datasets was not performed in our lab as no raw data was available.

Age of all samples was converted to dpc as described above. The combined validation dataset has an

age range of 42 – 280 dpc with the median at 112 dpc (**Figure S1, Table S1**). To evaluate the

performance of the DNAm clocks in adult brain samples we utilized data from the Brains for

Dementia Research (BDR) cohort previously generated by our group [14]. Briefly, these data consist

of 1,221 samples from 632 donors (age range 41-104 years, median = 84 years), with DNA extracted

from the prefrontal cortex (n = 610) and occipital cortex (n = 611). DNAm was quantified using the

Illumina EPIC DNAm array, and were pre-processed using a standard QC pipeline as described in [14].

Statistical evaluation of FBC performance

The predictive accuracy of the FBC was profiled in each dataset by two measures: Pearson's

correlation coefficient with reported chronological age and root mean squared error (RMSE).

Testing of Fetal Brain Clock (FBC) in cellular samples

iPSC - neuron samples.

Five different DNAm datasets generated using iPSCs, iPSC-derived NPCs and iPSC-derived neurons

were used to characterize epigenetic age of the neuronal cell model, details of which can be found in

Table S1. For two of these datasets (*Imm, Price*) DNAm data was generated by our lab in Exeter,

where DNAm was quantified using the Illumina EPIC DNA methylation array. These were

supplemented by three publicly available datasets, downloaded from GEO (*Nazor*, GSE31848,

Sultanov, GSE105093, and *Fernández-Santiago*, GSE51921) consisting of Illumina 450K DNAm array

data [3, 4, 29]. References describing the origin of cell lines and the different methods used for cell

culture and differentiation are listed in **Table S1**. Pre-processing and QC for the *Nazor* and

Fernández-Santiago datasets was not performed in our lab as no raw data was available.

iPSC – motor neuron samples

A dataset comprised of 23 cellular samples with two iPSC samples and 21 derived motor neurons

was generated by our lab, with DNAm quantified using the Illumina EPIC DNAm array (**Table S1**).

These data were QC'd following the pipeline described above.

ESC – neuron samples

Two publicly available cellular datasets with NPCs and neurons derived from embryonic stem cells

(ESCs) were downloaded from GEO (*Nazor*, GSE31848, *Kim* GSE38214) [29, 30]. Both datasets consist

of data quantified using the Illumina 450K DNAm array. Pre-processing and QC for both ESC – neuron

datasets were not performed in our lab as no raw data was available.

Statistical comparison of cellular states.

The FBC was applied to DNAm data for all cellular samples available. To test for differences in

predicted epigenetic age between cell stages within each dataset, either two sample t-tests or

ANOVA followed by Tukey HSD multiple comparison (when three cell stages were available), were

used. To combine results across all iPSC - neuron datasets, a mixed effects linear model was fitted

with predicted epigenetic age as the dependent variable, a fixed effect for cell stage represented as

two dummy variables contrasting NPCs vs iPSCs and iPSC-neurons vs iPSCs as and a random effect

(i.e. random intercept) for dataset.

Results

Fetal brain clock outperforms existing DNAm clocks at predicting age of prenatal brain samples.

We applied EN regression to genome wide DNAm data from a subset of available prenatal brain

samples (n = 193; **Table S1** and **Figure S1**) to develop the fetal brain clock (FBC). 107 DNAm probes

were assigned non-zero coefficients and therefore were selected as the basis of the FBC (**Table S2**).

We found no overlap in the DNAm sites selected for the FBC and DNAm sites used in the other

established clocks tested in our analysis. Testing the FBC clock in an independent test dataset of fetal

brain samples (**Table S1** and **Figure S1**) to evaluate its performance we found a strong linear

relationship between chronological and predicted prenatal age (r = 0.80; **Figure 1A**) with the

majority of samples predicted within 15 days of their actual chronological age (RMSE = 14.84 dpc).

To benchmark the performance of our clock, we compared it to three existing DNAm clocks:

Horvath's MTC [13], Knight's GAC [17] and Lee's CPC [18]. These clocks were selected as they

represent either the most well-established algorithm with the broadest applicability (MTC) or were

specifically developed to predict pre- and perinatal gestational ages, albeit in non-brain tissue (GAC

and CPC). Of note, the MTC only predicted 27 fetal brain samples (41.2%) as prenatal (dpc < 280)

with a low correlation between chronological and predicted age (r_{MTC} = 0.06). This correlation is

much lower than those reported in the original manuscript when Horvath tested the clock in adult

samples [13], highlighting the challenges with extrapolating clocks to samples which were not well

represented in model development. By comparison, the GAC and CPC perform better than the MTC,

although they have smaller correlation coefficients (r_{GAC} = 0.52 and r_{CPC} = 0.76) and are associated

with a larger error ($RMSE_{GAC}$ = 21.32 and $RMSE_{CPC}$ = 60.08) than the FBC. Interestingly, while the

predictions from the GAC are more precise, it is not as effective at ranking the samples by age as the

CPC. Taken together, these results demonstrate that our novel FBC outperforms existing clocks at

predicting age in fetal brain samples, and therefore is the optimal tool available to profile the

epigenetic age in models of neuronal development. When applying clocks to the training data, the

three established clocks produce similar correlations and RMSEs as in the testing data. As expected,

the predictions of the FBC in the training data are more accurate than the predictions in the testing

data, reflecting overfitting of the model (**Figure S2**).

We further tested the FBC in an independent prenatal brain dataset (n = 129, **Table S1**), finding a

stronger linear relationship between chronological age and predicted epigenetic age (r = 0.87, **Figure**

2A), than in the test dataset (**Figure 1A**) but a larger error rate (RMSE = 26.36 dpc). On closer

inspection, we observed that this error is mainly driven by a subset of older samples (> 185dpc,

Figure 2A) in the validation dataset (**Table S1**), that are older than any of the samples in the training

data (**Figure 2**). If we limit our analysis to the samples whose chronological age overlaps the range of

ages used in the training data (37 - 185 dpc; n = 125) then the error is decreased to 18.94 dpc. The

performance of the established clocks in the validation dataset is also comparable to their

performance in the testing dataset, with smaller correlations and higher error compared to the FBC

and no sample with a predicted prenatal age by the MTC (**Figure 2B-D**).

Given our previous finding of divergent, sex-specific age trajectories at multiple DNAm sites during

prenatal development [20], we tested whether the FBC performed differently between males and

females in our testing dataset. Although this analysis initially indicated a significant difference in the

correlation with age between males and females ($P_{Sex*Age}$ = 0.0007), this relationship is likely driven

by outliers. Indeed, a sensitivity analysis excluding the two samples with youngest and oldest

predicted ages produced a non-significant result ($P_{Sex*Age}$ = 0.081). Repeating the analysis in the

validation dataset we find a small significant effect of sex on age in the full dataset ($P_{Sex*Age}$ =

0.00179), which was driven by the samples older than 185, all of them being female. As samples

older than 185 dpc produce inaccurate predictions, they could unfairly bias the analysis of potential

sex effects and after removing the them from the analysis, there is no longer a significant effect of

sex on the accuracy of the clock ($P_{Sex*Age}$ = 0.95).

Fetal and gestational age clocks are not able to predict adult ages in adult brain tissue.

All four clocks were additionally tested in an adult brain DNAm dataset (**Figure S3**). As expected, the

FBC performs poorly in this sample set, with all samples predicted as prenatal albeit at the older end

of the spectrum of ages in the training data (range of predicted ages 115-170 dpc). In contrast the

MTC performs the best (r_{MTC} = 0.65, $RMSE_{MTC}$ = 20.11 years) as it is the only clock we considered that

was developed using adult samples. As with the FBC, the GAC and CPC fail to produce predictions of

postnatal age, again reflecting the fact that they were also constructed using data from pre- or

perinatal samples.

Fetal brain clock captures differences in differentiation of cellular stem cell models towards neurons

Having demonstrated that our novel FBC is the optimal clock to profile age in prenatal brain samples,

we applied it to DNAm data from multiple cellular studies to characterize epigenetic age in iPSCs and

ESCs differentiating towards cortical neurons. All samples were estimated to have a fetal epigenetic

age, regardless of cell stage, cell line origin or differentiation protocol. Furthermore, they were

predicted to have a "young" fetal age (**Figure 3**) with the iPSCs having a mean predicted age of 75.6

dpc (SD = 6.9 dpc; n = 59), the NPCs having a mean age of 79.1 dpc (SD = 11.0 dpc; n = 8) and iPSC -

neurons having a mean predicted age of 83.2 dpc (SD = 8.87 dpc; n = 31). To test whether the

differentiation process had an effect on the epigenetic age predictions from the FBC we compared

the estimated ages between iPSCs and neurons, observing significant differences in all datasets

(**Figure 3A**), with neurons being older than iPSCs. For the *Imm* dataset, which included proliferative

NPCs as well as postmitotic neurons, we additionally found a significant difference between NPCs

and iPSC-neurons (Δ_{mean} = 20.0 dpc, P = 0.00039), but not between iPSCs and NPCs (Δ_{mean} = 10.0 dpc,

P = 0.24). In contrast, in the *Nazor* dataset, which only included iPSCs and NPCs, we did find a

significant difference in predicted epigenetic age between iPSCs and NPCs (Δ_{mean} = 10.5 dpc, P =

0.00734). Meta-analyzing the data across the five studies including iPSCs and iPSC-derived NPCs and

neurons, we found that iPSC-neurons were predicted to have a significantly advanced epigenetic age

compared to iPSCs of about 2 weeks (Δ_{mean} = 13.35 dpc, P = 2.13e-11) but no significant difference

was observed between iPSC and NPCs (Δ_{mean} = 4.33 dpc, P = 0.11).

As well as cortical neurons, we applied the FBC to DNAm data from a study on iPSCs and

differentiated motor neurons. The motor neurons had a mean predicted epigenetic age of 79.82 dpc

(SD = 6.82 dpc) which is slightly younger than the cortical neurons (mean$_{iPSC-neurons}$ = 83.2 dpc). When

comparing the predicted ages between the motor neurons and the iPSCs they were originally

derived from, we did not observe a significant difference (Δ_{mean} = 2.94 dpc, P = 0.825, **Figure 3C**).

Finally, we tested for effects on epigenetic age through the process of differentiation from

embryonic stem cells (ESCs) to NPCs and neurons, using two publicly available DNAm datasets (**Table**

S1). In both datasets we observed a significant increase of predicted epigenetic age from ESC to NPC

(**Figure 3D**). Additionally, in the *Kim* dataset, we were able to compare ESC derived neurons, and

found a significant difference between ESCs and neurons (Δ_{mean} = 55.42 dpc, P = 0.00015, **Figure 3D**)

but no change in epigenetic age from NPC to neuron (Δ_{mean} = -0.58 dpc, P = 0.95).

Discussion

In this study we established a novel epigenetic clock, the fetal brain clock (FBC), to characterize the

earliest stages of human neurodevelopment, and applied it to determine the epigenetic age of iPSCs

and ESCs and their derived NPCs and neurons. Epigenetic clocks have been widely applied, including

for the analysis of both in-vivo and in-vitro models of aging, where they have been shown to

correlate with hallmarks of the aging process [31–34]. However, their application has predominantly

been focused on studies involving adult samples. Given the lack of fetal brain samples in the

development of existing DNAm clocks, prior to this study, there was no optimal method for

estimating the age of fetal brain samples from DNAm data, limiting the ability to characterize iPSC-

derived neuronal models or other models of neuronal development. We show that, in two non-

overlapping independent validation datasets, the FBC generates predictions that correlate strongly

with chronological age in prenatal brain samples. Furthermore, it outperforms both a pan-tissue

epigenetic clock (Horvath's MTC), and epigenetic clocks focused on the same developmental stage,

but based on DNAm profiled in different tissues (Knight's GAC and Lee's CPC) [13, 17, 18]. The FBC

outperforms these clocks using both correlation and error statistics (RMSE), indicating that it is not

only better at ranking the samples, but it generates more precise estimates.

As the FBC was predominantly trained on second trimester brain samples, with some first trimester

samples, it made less accurate predictions when applied to third trimester samples and performed

extremely poorly in adult brain samples. Altogether, this reinforces the findings of previous studies

that have also demonstrated that the applicability of DNAm clocks is dependent on the

characteristics of the data are were trained on, with the tissue and age range of the training samples

of particular relevance [14, 15]. More specifically, we note that while the accuracy of a DNAm clock

is typically decreased in tissues not represented in its training data, clocks are completely limited to

predicting ages represented in the training data. If the true age of a tested sample lies outside of the

age range of the training data, the clock is unable to provide an appropriate prediction with the

magnitude of inaccuracy increasing as the true age of the sample becomes more extreme suggesting

that, in general, age range is more critical than tissue when training a clock.

Previous epigenetic clocks have shown that the predicted epigenetic age of iPSCs is significantly

lower than the cells from which they are reprogrammed and the chronological age of the donor at

sample donation [13]. The induction of pluripotency reprograms the epigenome, including at the loci

used in the clock algorithm, ultimately leading to a younger predicted epigenetic age. However, in

these analyses the predicted ages remain postnatal, which is unexpected as human pluripotent cells

only occur during the early stages of human development and hence we hypothesized that, with an

adequately calibrated clock, iPSCs would be expected to be estimated as being early fetal. Applying

the FBC to five datasets of iPSCs and iPSC-derived NPCs and neurons, we found this to be the case.

iPSCs were estimated as having a mean age of 75.6 dpc, fitting our hypothesis that they represent

first trimester developmental stages. These results align with studies that have reported

rejuvenation effects on the transcriptome, telomeres and mitochondria of iPSCs following

reprogramming [35–37]. In addition, we profiled the effect on predicted epigenetic age following the

differentiation of iPSCs towards neurons reporting a small but significant aging effect of 13 dpc. This

developmental stage coincides with fetal neurogenesis [38] and suggests that while differentiation

does induce an aging process, it does not accelerate iPSC-neurons to a postnatal state. Of note,

Mertens and colleagues found that while iPSCs lose age related transcriptomic signatures, induced

neurons (iNs; neurons directly reprogrammed from fibroblasts) keep their specific aging signatures

[36]. Therefore, it would be interesting to apply our FBC to iNs, iPSCs, iPSC-neurons and their

corresponding somatic tissues to verify whether age associated DNAm differences are also

preserved in iNs. In addition, we tested the FBC on ESC and ESC-derived neurons. Although our

sample size was small, the results also suggested that the differentiation process induced a small

aging effect as the ESCs were differentiated into neurons. Altogether, our results indicate that iPSC-

neurons may have limited utility for the study of age-related brain diseases, like Alzheimer's disease

or other dementias, as many molecular processes related to an aging phenotype may not be

recapitulated.

Epigenetic clocks have been utilized for a wide range of applications [34], with a predominant focus

on exploring the biological meaning of deviations between chronological age and epigenetic age. As

we have shown the cause of this deviation may result from the use of an inappropriate clock,

therefore the FBC is a critical tool for assessing whether epigenetic age acceleration during

neurodevelopment is associated with later life outcomes such as disease or in utero exposures (such

as maternal smoking). The FBC also has utility for determining the specific developmental stage a

model of neurodevelopment recapitulates (e.g. brain organoids or cellular neuronal models) and

how different exposures or genetic backgrounds may influence neurodevelopmental processes and

aging.

While a strength of our study is the development of a bespoke clock to optimally profile the

epigenetic age of human fetal brain samples, due to the training data predominantly containing

second trimester samples the FBC is most accurate for this period of neurodevelopment. We are

confident that the FBC has correctly predicted fetal epigenetic ages for the cell lines included in our

analysis as the vast majority of the stem cell models and their derived neurons were less than the

median age in the training data. This indicates that the predictions are not confounded by saturation

of the coefficients. Although we took advantage of previously published data to include all available

samples appropriate for addressing our research questions, some of the group sizes, in particular the

NPCs, were small and we were not powered to detect significant aging effects as a result of

differentiation from iPSCs to NPCs in all datasets. Furthermore, across the different studies there

was variation in the estimated prenatal age of each cell state; we hypothesize that these differences

results from subtle variation in differentiation protocols, timepoints of cell collection or the

definition of NPCs within the respective studies [39]. Despite this, we are confident in the

conclusions we report as these study specific effects were controlled for in our analysis.

In summary, we demonstrate that established DNAm clocks struggle to capture changes in

epigenetic age during neurodevelopment and for precise predictions a bespoke clock trained on

fetal brain data is required. Using the FBC to assess the epigenetic age of iPSCs and differentiated

neurons, we found that iPSCs and derived NPCs and neurons reflect early prenatal developmental

stages. Our findings question the suitability of the iPSC-neurons for the study of aging associated

processes.

List of abbreviations

Abbreviation	Definition
AD	Alzheimer's Disease
BDR	Brains for Dementia Research
CPC	Control Placental Clock
DNAm	DNA methylation
dpc	Days post-conception
EN	Elastic net
ESC	Embryonic Stem Cell
FBC	Fetal brain clock
GA	Gestational age
GAC	Gestation age clock
GEO	Gene Expression Omnibus
iN	Induced neuron
iPSC	Induced pluripotent stem cell
iPSC-neuron	iPSC-derived neuron
LASSO	Least Absolute Shrinkage and Selection Operator
MTC	Multi-tissue clock
NPC	Neuronal precursor cell
QC	Quality control
RMSE	Root mean squared error

7   **Declarations**

8   *Ethical approval*

Ethical approval for the collection of human fetal brain tissue acquired from the Human

Developmental Biology Resource was granted by the Royal Free Hospital research ethics committee

under reference 08/H0712/34 and Human Tissue Authority material storage license 12220. Ethical

approval for collection of human fetal brain tissue for the Medical Research Council Brain Bank was

granted under reference 08/MRE09/38. Ethical approval for the work with methylomic data of

human brain tissue from the BDR cohort was granted by the University of Exeter, College of

Medicine and Health Research Ethics Committee under reference Mar20/D/009Δ5. Ethical approval

for the Price cellular samples is as follows: participants were recruited and methods carried out in

accordance to the 'Patient iPSCs for Neurodevelopmental Disorders (PiNDs) study' (REC No

13/LO/1218). Informed consent was obtained from all subjects for participation in the PiNDs study.

Ethical approval for the PiNDs study was provided by the NHS Research Ethics Committee at the

South London and Maudsley (SLaM) NHS R&D Office.

Consent for publication

Not applicable

Availability of data and materials

Raw and normalized DNAm data for the human fetal brain samples used to test and train the fetal

brain clock have been submitted to the NCBI Gene Expression Omnibus (GEO;

http://www.ncbi.nlm.nih.gov/geo/) under accession numbers GSE58885, GSE74193, GSE157908,

GSE116754 and GSE90871. DNAm data for the unpublished fetal brain are available from the

corresponding author on request. DNAm data for the cellular datasets, are also available under

accession numbers GSE158089, GSE105093, GSE51921, GSE31848 and GSE38214. DNAm data for

the cellular dataset *Bhinge* is available from the corresponding author on request. The coefficients of

the FBC are included within the article in **Table S2**. The custom R scripts used for this study and to

apply the FBC are available at https://github.com/LSteg/EpigeneticFetalClock.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by a Simons Foundation (SFARI) grant (grant number: 573312) awarded to

JM and a Medical Research Council (MRC) project grant (grant number: MR/R005176/1) to JM. The

human embryonic and fetal material was provided by the Joint MRC Wellcome Trust Human

Developmental Biology Resource (http://www.hdbr.org), which is funded by Medical Research

Council grant G0700089 and Wellcome Trust grant GR082557. Adult brain DNAm data was

generated using tissue from the Brains for Dementia Research (BDR) cohort, which is jointly funded

by Alzheimer's Research UK and the Alzheimer's Society in association with the Medical Research

Council. The South West Dementia Brain Bank is part of the Brains for Dementia Research program,

jointly funded by Alzheimer's Research UK and Alzheimer's Society, and is also supported by BRACE

(Bristol Research into Alzheimer's and Care of the Elderly) and the Medical Research Council. The

Price cellular data was supported by grants from the European Autism Interventions (EU-AIMS): the

Innovative Medicines Initiative Joint Undertaking under grant agreement no. 115300, resources of

which are composed of financial contribution from the European Union's Seventh Framework

Programme (FP7/2007-2013) and EFPIA companies' in kind contribution (JP, DPS); StemBANCC:

support from the Innovative Medicines Initiative joint undertaking under grant 115439-2, whose

resources are composed of financial contribution from the European Union [FP7/2007-2013] and

EFPIA companies' in-kind contribution (JP, DPS). G.S. was supported by a PhD studentship from the

Alzheimer's Society. N.J.B. was supported by the Medical Research Council grant MR/L010674/2.

N.D.A. was supported by Medical Research Council grant N013255/1.

Authors' contributions

L.C.S, G.L.S. and E.H. designed the work; J.M., E.H., E.L.D., N.J.B, A.B., J.P., D.P.S., N.D.A. and K.L.

acquired funding; J.I., J.P.D., R.F., E.L.D., S.C.N., A.R.J., J.B., G.W.A.N., E.M.W., L.W.P., E.L.C., K.M.J.,

A.F., G.M., D.P.S., E.P., A.B., N.J.B. and N.D.A acquired and processed samples and data; L.C.S, G.L.S.

and E.H analyzed and interpreted data; L.C.S. and E.H. wrote the paper.

Acknowledgements

We would like to gratefully acknowledge all donors and their families for the tissue provided for this

study. We are grateful to Rosy Watkins, Paulina Nowosiad, Rupert Faraway and Roland Nagy for

generation and maintenance of *Price* iPSCs [41, 42].

Figure Legends

Figure 1 The Fetal Brain Clock (FBC) outperforms other DNAm clocks when applied to

neurodevelopmental samples. Shown are scatterplots comparing chronological age (x-axis; days

post-conception) against (y-axis; days post-conception) predicted epigenetic age calculated using **A**

Fetal Brain Clock (FBC), **B** Horvath's multi tissue clock (MTC), **C** Knight's Gestational Age Clock (GAC),

and **D** Lee's Control Placental Clock (CPC) in our fetal brain testing dataset (n = 65, age range = 23 –

153 dpc). The black line indicates the identity line of chronological and predicted epigenetic age and

represents a perfect prediction. Two statistics were calculated to evaluate the precision of each

DNAm clock: Pearson's correlation coefficient (r) and the root mean squared error (RMSE).

Figure 2 Validation of the Fetal Brain Clock in an independent fetal brain dataset. Shown are scatterplots comparing chronological age (x-axis; days post-conception (dpc)) against (y-axis; dpc) predicted epigenetic age calculated using **A** Fetal Brain Clock (FBC), **B** Horvath's Multi Tissue Clock (MTC), **C** Knight's Gestational Age Clock (GAC), and **D** Lee's Control Placental Clock (CPC) on data from an independent validation dataset (n = 129, age range = 42 - 280). Two statistics were calculated to evaluate the precision of each DNAm clock: Pearson's correlation coefficient (r) and the root mean squared error (RMSE). The dashed line in **A** indicates 185 dpc, which is the highest available age in the training dataset of the FBC.

Figure 3 Comparisons of predicted epigenetic age using the fetal brain clock between cellular models throughout differentiation states. A Boxplots comparing the distribution of predicted epigenetic age (days post-conception) in iPSCs and their derived NPCs and neurons, where each panel represents a different dataset. P values of Tukey HSD corrected ANOVA for the *Imm* dataset and two-sample t-tests for *Price*, *Fernández-Santiago* and *Sultanov* datasets are given. F. -S. = *Fernández-Santiago*. **B** Boxplots of predicted epigenetic age calculated using the FBC where samples are grouped by cell stage (n = 82, 30 iPSCs, 4 NPCs, 48 iPSC-neurons) and colored by dataset. P values from mixed effects model are given for differences between iPSCs and NPCs (non-significant) and iPSC and neurons. **C** Boxplots comparing the predicted epigenetic age in a cohort with iPSCs and derived motor neurons. P values of two-sample t-test are given. **D** Boxplots of predicted epigenetic age by FBC applied on two datasets including ESCs and their derived NPCs and neurons. P values of Tukey HSD corrected ANOVA for the *Kim* dataset and two-sample t-tests for the *Nazor* dataset are given

Additional Files

Additional File 1: Table S1 Summary of fetal, adult and cellular datasets.

Additional File 2: Table S2 Intercept and coefficients of the fetal brain clock.

Additional File 3: Figure S1 Histogram of age distribution. Chronological age of the fetal samples

measured in days post-conception in A) training data, B) testing data and C) validation data.

Additional File 4: Figure S2 Comparison of predictions from the four DNAm clocks in the training

data (n = 193). Shown are scatterplots comparing chronological age (x-axis) against age (y-axis)

predicted epigenetic age for **A** Fetal Brain Clock (FBC); **B** Horvath's Multi Tissue Clock (MTC); **C**

Knight's Gestational Age Clock (GAC); **D** Lee's Control Placental Clock (CPC) in the data used for

training of the FBC. Where necessary, predicted age was converted to days post-conception. The

black line indicates the identity line of chronological and predicted epigenetic age and represents a

perfect prediction. Two statistics were calculated to evaluate the precision of each DNAm clock:

Pearson's correlation coefficient (r) and the root mean squared error (RMSE).

Additional File 5: Figure S3 Comparison of predictions from the four DNAm clocks in adult brain

samples (n = 1221). Shown are scatterplots comparing chronological age (x-axis) against (y-axis)

predicted epigenetic age calculated using **A** Fetal Brain Clock (FBC); **B** Horvath's multi tissue clock

(MTC); **C** Knight's Gestational Age Clock (GAC); **D** Lee's Control Placental Clock (CPC) in an

independent adult brain dataset. Where necessary, predicted age was converted to years, where 0

indicates birth. The black line indicates the identity line of chronological and predicted epigenetic

age and represents a perfect prediction. Two statistics were calculated to evaluate the precision of

each DNAm clock: Pearson's correlation coefficient (r) and the root mean squared error (RMSE).

References

1. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S (2007) Induction of Pluripotent Stem Cells from Adult Human Fibroblasts by Defined Factors. Cell 131:861–872

2. Telezhkin V, Schnell C, Yarova P, et al (2016) Forced cell cycle exit and modulation of GABA A , CREB, and GSK3β signaling promote functional maturation of induced pluripotent stem cell-derived neurons. Am J Physiol Physiol 310:C520–C541

3. Fernández-Santiago R, Carballo-Carbajal I, Castellano G, et al (2015) Aberrant epigenome in iPSC -derived dopaminergic neurons from Parkinson's disease patients. EMBO Mol Med 7:1529–1546

4. Sultanov R, Lebedeva O, Arapidi G, Lagarkova M, Kiselev S (2018) Methylation profile of induced pluripotent stem cells generated by integration and integration-free approaches. Data Br 17:662–666

5. Gordon J, Amini S, White MK (2013) General overview of neuronal cell culture. Methods Mol Biol 1078:1–8

6. Arber C, Lovejoy C, Wray S (2017) Stem cell models of Alzheimer's disease: Progress and challenges. Alzheimer's Res Ther. https://doi.org/10.1186/s13195-017-0268-4

7. Morgani S, Nichols J, Hadjantonakis AK (2017) The many faces of Pluripotency: In vitro adaptations of a continuum of in vivo states. BMC Dev Biol 17:1–20

8. Monk D, Mackay DJG, Eggermann T, Maher ER, Riccio A (2019) Genomic imprinting disorders: lessons on how genome, epigenome and environment interact. Nat Rev Genet 20:235–248

9. Zeng Y, Chen T (2019) DNA methylation reprogramming during mammalian development. Genes (Basel). https://doi.org/10.3390/genes10040257

10. Smith ZD, Meissner A (2013) DNA methylation: Roles in mammalian development. Nat Rev Genet 14:204–220

11. Levine ME, Lu AT, Quach A, et al (2018) An epigenetic biomarker of aging for lifespan and healthspan. Aging (Albany NY) 10:573–591

12. Hillary RF, Stevenson AJ, McCartney DL, et al (2020) Epigenetic measures of ageing predict the prevalence and incidence of leading causes of death and disease burden. Clin Epigenetics 12:115

13. Horvath S (2013) DNA methylation age of human tissues and cell types. Genome Biol. https://doi.org/10.1186/gb-2013-14-10-r115

14. Shireby G, Davies J, Francis P, et al (2020) Recalibrating the Epigenetic Clock: Implications for Assessing Biological Age in the Human Cortex. bioRxiv 2020.04.27.063719

15. El Khoury LY, Gorrie-Stone T, Smart M, et al (2019) Systematic underestimation of the epigenetic clock and age acceleration in older subjects. Genome Biol 20:283

16. Zhang Q, Vallerga CL, Walker RM, et al (2019) Improved precision of epigenetic clock estimates across tissues and its implication for biological ageing. Genome Med 11:54

17. Knight AK, Craig JM, Theda C, et al (2016) An epigenetic clock for gestational age at birth based on blood methylation data. Genome Biol 17:1–11

18. Lee Y, Choufani S, Weksberg R, et al (2019) Placental epigenetic clocks: Estimating gestational age using placental DNA methylation levels. Aging (Albany NY) 11:4238–4253

19. R Development Core Team (2012) R: A Language and Environment for Statistical Computing.

20. Spiers H, Hannon E, Schalkwyk LC, Smith R, Wong CCY, O'Donovan MC, Bray NJ, Mill J (2015) Methylomic trajectories across human fetal brain development. Genome Res 25:338–352

21. Jaffe AE, Gao Y, Deep-Soboslay A, Tao R, Hyde TM, Weinberger DR, Kleinman JE (2015) Mapping DNA methylation across development, genotype and schizophrenia in the human frontal cortex. Nat Neurosci 19:40–47

22. Pidsley R, Y Wong CC, Volta M, Lunnon K, Mill J, Schalkwyk LC (2013) A data-driven approach to preprocessing Illumina 450K methylation array data. BMC Genomics. https://doi.org/10.1186/1471-2164-14-293

23. Gorrie-Stone TJ, Smart MC, Saffari A, Malki K, Hannon E, Burrage J, Mill J, Kumari M, Schalkwyk LC (2019) Bigmelon: Tools for analysing large DNA methylation datasets. Bioinformatics 35:981–986

24. Chen YA, Lemire M, Choufani S, Butcher DT, Grafodatskaya D, Zanke BW, Gallinger S, Hudson TJ, Weksberg R (2013) Discovery of cross-reactive probes and polymorphic CpGs in the Illumina Infinium HumanMethylation450 microarray. Epigenetics 8:203–209

25. Zou H, Hastie T (2005) Regularization and variable selection via the elastic net. J R Stat Soc Ser B Stat Methodol 67:301–320

26. Friedman J, Hastie T, Tibshirani R (2010) Regularization paths for generalized linear models via coordinate descent. J Stat Softw 33:1–22

27. Chatterton Z, Hartley BJ, Seok MH, et al (2017) In utero exposure to maternal smoking is associated with DNA methylation alterations and reduced neuronal content in the developing fetal brain. Epigenetics and Chromatin. https://doi.org/10.1186/s13072-017-0111-y

28. Colunga T, Hayworth M, Kreß S, et al (2019) Human Pluripotent Stem Cell-Derived Multipotent Vascular Progenitors of the Mesothelium Lineage Have Utility in Tissue Engineering and Repair. Cell Rep 26:2566-2579.e10

29. Nazor KL, Altun G, Lynch C, et al (2012) Recurrent variations in DNA methylation in human pluripotent stem cells and their differentiated derivatives. Cell Stem Cell 10:620–634

30. Kim M, Park YK, Kang TW, et al (2014) Dynamic changes in DNA methylation and hydroxymethylation when hES cells undergo differentiation toward a neuronal lineage. Hum Mol Genet 23:657–667

31. Liu Z, Leung D, Thrush K, Zhao W, Ratliff S, Tanaka T, Schmitz LL, Smith JA, Ferrucci L, Levine ME (2020) Underlying features of epigenetic aging clocks in vivo and in vitro. Aging Cell 19:e13229

32. Horvath S, Raj K (2018) DNA methylation-based biomarkers and the epigenetic clock theory of ageing. Nat Rev Genet 19:371–384

33. Wagner W (2019) The link between epigenetic clocks for aging and senescence. Front Genet. https://doi.org/10.3389/fgene.2019.00303

34. Ryan CP (2020) "Epigenetic clocks": Theory and applications in human biology. Am J Hum Biol e23488

35. Suhr ST, Chang EA, Tjong J, Alcasid N, Perkins GA, Goissis MD, Ellisman MH, Perez GI, Cibelli JB (2010) Mitochondrial Rejuvenation After Induced Pluripotency. PLoS One. https://doi.org/10.1371/journal.pone.0014095

36. Mertens J, Paquola ACM, Ku M, et al (2015) Directly Reprogrammed Human Neurons Retain Aging-Associated Transcriptomic Signatures and Reveal Age-Related Nucleocytoplasmic Defects. Cell Stem Cell 17:705–718

37. Lapasset L, Milhavet O, Prieur A, et al (2011) Rejuvenating senescent and centenarian human cells by reprogramming through the pluripotent state. Genes Dev 25:2248–2253

38. Lui JH, Hansen D V., Kriegstein AR (2011) Development and evolution of the human neocortex. Cell 146:18–36

39. Mattis VB, Tom C, Akimov S, et al (2014) HD iPSC-derived neural progenitors accumulate in culture and are susceptible to BDNF withdrawal due to glutamate toxicity. Hum Mol Genet 24:3257–3271

40. Bhinge A, Namboori SC, Zhang X, VanDongen AMJ, Stanton LW (2017) Genetic Correction of SOD1 Mutant iPSCs Reveals ERK and JNK Activated AP1 as a Driver of Neurodegeneration in Amyotrophic Lateral Sclerosis. Stem Cell Reports 8:856–869

41. Cocks G, Curran S, Gami P, et al (2014) The utility of patient specific induced pluripotent stem cells for the modelling of Autistic Spectrum Disorders. Psychopharmacology (Berl) 231:1079–1088

42. Warre-Cornish K, Perfect L, Nagy R, et al (2020) Interferon-γ signaling in human iPSC–derived neurons recapitulates neurodevelopmental disorder phenotypes. Sci Adv 6:eaay9506

A iPSC - Neuron Datasets

i. Imm

ii. Price

iii. Nazor

iv. F.–S.

v. Sultanov

B iPSC - Neuron Merged

C iPSC - Motor neuron

i. Bhinge

D ESC - Neuron

i. Nazor **ii. Kim**