Characterisation of growth plate dynamics in murine models of osteoarthritis

Hasmik J. Samvelyan[1,2], Kamel Madi[3], Anna E. Törnqvist[4,5], Behzad Javaheri[6], Katherine A.

Staines[1,2]

[1] School of Pharmacy and Biomolecular Sciences, University of Brighton, Brighton, UK

[2] Centre for Stress and Age-Related Disease, University of Brighton, Brighton, UK

[3] 3Dmagination Ltd, Harwell Campus, Didcot, UK

[4] Rheumatology and Bone Diseases Unit, Centre for Genomic and Experimental Medicine,

MRC Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK

[5] Centre for Bone and Arthritis Research, University of Gothenburg, Gothenburg, Sweden

[6] School of Mathematics, Computer Science and Engineering, City University of London,

London, UK

Corresponding authors

Katherine Ann Staines, University of Brighton, Lewes Road, Brighton BN2 4GJ

Email: k.staines@brighton.ac.uk Tel: 01273 642 094

Hasmik Jasmine Samvelyan, University of Brighton, Lewes Road, Brighton BN2 4GJ

Email: h.samvelyan@brighton.ac.uk Tel: 01273 641 671

Running title: Growth dynamics and osteoarthritis

Abstract

Objective: To investigate growth plate (GP) dynamics in surgical and loading murine models

of osteoarthritis (OA), to understand whether abnormalities in these dynamics predict those at

risk of OA.

Design: 8-week-old C57BL/6 male mice underwent destabilization of medial meniscus

(DMM) ($n=8$) surgery in right knee joints. Contralateral left knee joints had no intervention

(controls). In 16-week-old C57BL/6 male mice ($n=4$), OA was induced using non-invasive

mechanical loading of right knee joints with peak force of 11N. Non-loaded left knee joints

were internal controls. Chondrocyte transiency in tibial articular cartilage (AC) and GP was

examined by histology and immunohistochemistry. Tibial subchondral bone (SCB)

parameters were measured using microCT and correlated to GP bridging.

Results: Higher expression of chondrocyte hypertrophy markers; Col10a1 and MMP13 were

observed in tibial AC chondrocytes of DMM and loaded mice. In tibial GP, Col10a1 and

MMP13 expressions were widely dispersed in a significantly enlarged zone of proliferative

and hypertrophic chondrocytes. 3-dimensional quantification revealed enriched GP bridging

and higher bridge densities in medial compared to lateral tibiae of DMM and loaded knee

joints of the mice. GP dynamics were associated with increased SCB and epiphyseal

trabecular bone volume fraction (BV/TV; %) in medial tibiae of DMM and loaded knee joints

respectively.

Conclusions: Results confirm associations between aberrant chondrocyte hypertrophy marker

expression and OA pathology in a surgical and loaded murine model of OA. Spatial

variations in GP bridging formation revealed accelerated cartilage-bone transitions which

may contribute to anatomical variation in vulnerability to OA development in these models.

Keywords: osteoarthritis, cartilage, growth, chondrocyte, murine model

Introduction

Osteoarthritis (OA) is a chronic musculoskeletal disease and a leading cause of disability and major healthcare costs in the world. It is estimated that worldwide 10% of men and 18% of women aged over 60 years have symptomatic OA[1] and, therefore, the predicted increase in the ageing population and longevity will result in a greater occurrence of the disease. OA is a complex disease in which the pathogenesis, cellular and molecular mechanisms of initiation and progression are not completely understood. It is characterised by progressive loss of articular cartilage (AC), formation of osteophytes, subchondral bone (SCB) sclerosis, synovial proliferation and inflammation and lax tendons. These can ultimately lead to a loss of joint function, pain, reduced mobility and disability[2]. Despite the significant healthcare and economic burden there are few non-invasive therapies available to patients. Therefore, understanding the pathogenesis of OA and defining the molecular mechanisms underpinning AC degeneration can lead to the development of successful targeted and effective disease-modifying treatments.

Primary OA is described as naturally occurring OA affecting one joint (localised) or three or more joints (generalised), while secondary OA is associated with various causes and risk factors leading to the disease including trauma, obesity, diabetes, metabolic bone and congenital disorders[3]. AC degeneration is one of the main hallmarks of OA and previous research has largely sought to identify mechanisms underpinning its deterioration. Fully developed, uncalcified AC is populated by a single resident cell chondrocytes, which maintain a stable phenotype characterised by small cell size and expression of tenascin-C[4]. The inherent stability of AC chondrocytes ensures that dynamic events are restricted to assure lifelong articular integrity and healthy joint function. In contrast, epiphyseal growth plate (GP) chondrocytes have a transient phenotype to ensure long bone development

(endochondral ossification) and growth. GP chondrocytes undergo a differentiation sequence

of proliferation, maturation and hypertrophy. The final stage of chondrocyte hypertrophy

enables mineralisation of the cartilage extracellular matrix, vascular invasion and subsequent

replacement of the mineralised cartilage anlagen with bone[5]. These processes are coupled,

however, with sexual maturation the human GP undergoes progressive narrowing as bony

bridges form and span its width. This ultimately leads to complete GP closure and cessation

of human growth. Indeed, in humans the longitudinal bone growth stops with the onset of

puberty, the metaphysis then fuses with the epiphysis and the growth plate disappears [6]. In

mice, longitudinal bone growth does not cease at sexual maturity instead it slows

dramatically at puberty, but the growth plates do not completely fuse and disappear [7].

We have previously shown that in the STR/Ort mouse, a naturally occurring OA murine

model, AC chondrocytes transform from their inherently stable phenotype to a transient one,

characteristic of the chondrocytes in the GP. This was confirmed by immunolabelling for

chondrocyte hypertrophy markers; type X collagen (Col10a1) and matrix metalloproteinase

13 (MMP13)[8]. Further, we revealed accelerated longitudinal bone growth, aberrant

expression of growth plate markers (Col10a1 and MMP13) and increased growth plate

chondrocyte maturation in these mice. Consistent with this, using a novel synchrotron

computed tomography method we revealed enriched GP bone bridging in STR/Ort mouse

tibiae indicative of advanced GP closure which may underpin OA[8,9].

Despite this, interlinks between the differing chondrocyte phenotypes in the AC and the GP,

and the contribution that the GP may play in underpinning OA vulnerability is not yet

understood. Indeed, understanding this will inform strategies for maintaining musculoskeletal

health in ageing by potentially identifying whether GP dynamics may predict who is at risk of

OA in later life and ultimately developing targeted OA treatments. Therefore, we

hypothesised that surgically (destabilisation the medial meniscus [DMM]) or non-surgically

(mechanical loading) induced OA onset in C57BL/6 mice is linked to altered growth plate

dynamics.

Method

Animals

Male C57BL/6 wild type mice at 7 weeks of age (young adult) were obtained from Charles

River Laboratories Inc. (Margate, UK). The mice were acclimatised to their surroundings for

seven days. All mice were allowed free access to water and maintenance diet ad libitum

(Special Diet Services, Witham, UK) in a 12-hour light/dark cycle at a room temperature of

21 ± 2^{0}C and relative humidity of $55 \pm 10\%$. All procedures complied with the United

Kingdom Animals (Scientific Procedures) Act 1986 and were approved by The University of

Edinburgh Roslin Institute's Animal Users and Research Ethics Committees. All analyses

were conducted blindly to minimise the effects of subjective bias.

The destabilisation of the medial meniscus (DMM)

A group of eight 8-week old wild type C57/BL6 male mice underwent DMM surgeries to

induce OA-like changes in the right knee joints under isoflurane-induced anaesthesia. We

chose not to performed SHAM (placebo) surgery on the left contralateral knee of the animals

based upon animal welfare grounds since it was previously shown that there is no difference

in OA scores between SHAM-operated and non-operated knee joints[10,11]. Animals were

monitored for unexpected adverse effects to reduce suffering and distress. Following

transection of the medial meniscotibial ligament (MMTL) to destabilise the medial meniscus,

the skin was closed and anaesthesia reversed[10]. Eight weeks later, mice were sacrificed by

exsanguination and confirmation of death by cervical dislocation. The knee joints of all the

mice were dissected, fixed in 4% paraformaldehyde for 24 hours at 4°C, and then stored in

70% ethanol.

In vivo loading of the knee joint

The right knee joints of 16-week-old wild type C57BL/6 male mice ($n = 4$) were subjected to

non-invasive, dynamic axial mechanical loading under the isofluorane-induced anaesthesia

(liquid isofluorane was vaporised to a concentration of 4% and maintained at a concentration

of 2% with oxygen) for 7 min/day, 3 alternate days a week for 2 weeks according to the

protocols described in the previous studies [12,13]. The left knee joints were non-loaded internal

controls in these animals. Briefly, using a servo-electric materials testing machine

(Electroforce 3100, Bose, UK), axial compressive loads were applied through the right knee

joint via customised concave cups which held the knee and ankle joints flexed and the tibiae

vertically.

The tibia was held in place by continuous static preload of 0.5N onto which dynamic loads

were superimposed in a series of 40 trapezoidal shaped waveform cycles with steep up and

down ramps and a peak force of 11N for 0.05 seconds (0.025 seconds rise and fall time; 9.9

seconds baseline hold time between periods of peak loading). The right and left knees were

dissected 3 days after the final loading episode. Mice were sacrificed by exsanguination and

confirmation of death by cervical dislocation. Knee joints were fixed in 4%

paraformaldehyde for 24 hours at $4^{o}C$ before being stored in 70% ethanol.

Micro-computed (microCT) tomography and 3-dimensional (3D) bridging analysis

Scans were performed with an 1172 X-Ray microtomograph (Bruker MicroCT, Kontich,

Belgium) to evaluate the SCB and GP bridging. High-resolution scans with an isotropic voxel

size of 5 µm were acquired (50 kV, 200 µA, 0.5 mm aluminium filter, 0.6° rotation angle).

The projection images were reconstructed and binarised with a threshold of 0 to 0.16, ring

artefact reduction was set at 10 and beam hardening correction at 0% using the SkyScan

NRecon software package (v1.6.9.4, Bruker MicroCT). The images then were realigned

vertically using DataViewer software (v1.5.1.2 64-bit, Bruker MicroCT) to ensure similar

orientation for analysis. Hand-drawn regions of interests (ROI) of the SCB and epiphyseal

trabecular bone for each tibial lateral and medial compartments were selected[14]. The

structural parameters of tibial SCB plate and epiphyseal trabecular bone were calculated

using 3D algorithms of SkyScan CTAn software (Bruker MicroCT) including SCB (SCB

BV/TV; %) and trabecular bone volume fraction (Tb. BV/TV; %) and correlated to GP

bridging analysis using a 3D synchrotron-computed tomography quantification method as

previously described[15]. Briefly, microCT scans of the tibiae were segmented using a region-

growing algorithm within the Avizo® (V8.0, VSG, Burlington, VT, USA) software. The

central points of each bony bridges were identified and projected on the tibial joint surface.

The distribution of the areal number density of bridges (N, the number of bridges per 256 µm

$\times$ 256 µm window; $d = m/V$) is then calculated and superimposed on the tibial joint surface

(each bridge has a colour that represents the areal number density at the bridge location). The

SCB plate and epiphyseal trabecular bone thickness (Th; mm) was determined and colour-

coded thickness images were generated using the Avizo® software[15].

Histological analysis

The left and right knee joints of all the mice were decalcified in 10%

ethylenediaminetetraacetic acid (EDTA) solution, wax-embedded at Leica EG1160 Tissue

Embedding Station and 6 µm coronal sections cut using Leica RM2135 manual microtome.

Immunohistochemistry

Immunohistochemical analysis of chondrocyte transiency markers in tibial AC and GP was

performed on 6 µm coronal sections using anti-matrix metalloproteinase 13 (anti–MMP13)

(1:200 dilution; Abcam) or anti-collagen type X (anti-Col10a1) (1:100 dilution; Abcam)

antibodies. As a control, an equal concentration of rabbit IgG was used. For

immunohistochemical localisation of MMP13 and Col10a1, sections were dewaxed in xylene

and rehydrated. Sections were incubated at 37 °C for 30 min in 1mg/ml trypsin for antigen

demasking. Endogenous peroxidases were blocked by treatment with 0.3% H_2O_2 in methanol

(Sigma) for 30 min at room temperature. The Vectastain ABC universal detection kit (Vector

Laboratories, Peterborough, UK) was used to detect the biotinylated secondary antibody

(Anti-Mouse IgG Reagent) after incubation for 30 min at room temperature according to the

manufacturer's instructions. Diaminobenzidine (DAB) solution used to detect the location of

antigens. The sections were finally dehydrated, counterstained with haematoxylin and

mounted in DePeX. All sections to be compared were immunolabelled at the same time to

standardise conditions and minimise any differences in antibody incubation times.

GP zone analysis

The stained sections from the joints of 4 individual mice from each experimental group were

used to measure the width of the GP proliferating and hypertrophic zones, as well as the total

GP width, measured at 10 different points along the length of the GP in tibiae, using a light

microscope and ImageJ software.

Statistical analysis

All analyses were performed with GraphPad Prism software 6.0f version (GraphPad Inc, La

Jolla, CA, USA) using a two-sided 0.05 level of significance. The results were presented as

the mean ± standard error of the mean (SEM). The Normal distribution of data was assessed

using the Shapiro-Wilk normality test. For comparing two groups (experimental with control,

or medial with the lateral compartment of tibiae), two-tail Student's *t*-test (paired or unpaired)

was used. For comparing more than two groups, two-way ANOVA (analysis of variance) was

used with Tukey post-hoc test.

Results

Transient chondrocyte behaviour in the tibial AC of DMM and loaded C57BL/6 young adult

male mice

We first sought to confirm whether loss of AC in C57BL/6 mice with surgically and loading

induced OA was associated with the expression of markers of transient chondrocyte

phenotype. Immunohistochemistry analysis showed higher expression levels of well-

established chondrocyte hypertrophy markers; Col10a1 and MMP13, observed in tibial AC of

C57BL/6 mice that have undergone DMM surgery or mechanical loading compared with

non-operated and non-loaded control left tibiae, respectively (Figure 1A, 1B). The expression

pattern of Col10a1 was largely restricted to hypertrophic chondrocytes in the uncalcified zone

of the AC of unaffected condyles of non-operated (Figure 1A) and non-loaded (Figure 1B)

mouse left joints as expected[16]. Whereas, the immunolabeling of Col10a1 was more

widespread throughout the extracellular matrix (ECM) of the AC in affected right joints of

DMM (Figure 1A) and loaded (Figure 1B) mice. Similarly, immunohistochemistry analysis

showed positive MMP13 labelling in both superficial and deep articular chondrocytes in the

right joints of DMM and loaded C57BL/6 male mice compared to the control knee joints

(Figure 1A and B). These findings confirm an aberrant deployment of transient chondrocytes

in uncalcified AC.

Dysfunctional GP morphology in DMM and loaded tibiae of C57BL/6 young adult male mice

In the tibial GP of mice with surgically and loading induced OA, Col10a1 expression was

more greatly and widely dispersed throughout the zones of proliferative and hypertrophic

chondrocytes compared with their controls (Figure 2A and 2B). Indeed, immunolabeling for

Col10a1 revealed the expected localisation in the GP of non-operated and non-loaded mouse

tibiae, limited primarily to the hypertrophic zone and underlying adjacent metaphyseal bone

(Figure 2A and 2B). This disrupted the distribution of a GP zone marker was also evident for

MMP13 in the GP of DMM and loaded C57BL/6 mouse tibiae compared to their controls

(Figure 2A and 2B).

GP zone analysis revealed significantly enlarged proliferative and hypertrophic zones of

chondrocytes in both DMM (p=0.002 and p<0.0001, respectively) and loaded (both

p<0.0001) tibiae of C57BL/6 mice compared to their controls, and significantly increased

total GP width (both p<0.0001) (Figure 3). Together, the results may indicate associations

between dysfunctional GP morphology and marker expression, and OA development.

Associations between GP bridging in DMM and loaded tibiae of C57BL/6 young adult male

mice and OA development

To further correlate aberrant longitudinal GP dynamics, GP bridging and OA development in

these C57BL/6 young adult male mice, we used our newly developed 3D method to quantify

bony bridges across the tibial epiphysis of DMM and loaded mice (Figure 4). 3D

quantification revealed a significantly higher number of GP bridges in medial compared to

lateral tibiae that underwent DMM surgeries (306 ± 32 versus 196 ± 33; p=0.03) (Figure 4B

and 4E). This significant difference was not observed in non-operated tibiae (326 ± 32 versus

254 ± 32; p=0.14) (Figure 4A and 4E). Similarly, significantly enriched GP bridging was

evident in the medial compartment of loaded tibiae in comparison to the lateral compartment

(731 ± 19 versus 532 ± 56; p=0.01) (Figure 4D and 4E), and in those of non-loaded tibiae,

although less pronounced than in the loaded right knee joints (782 ± 38 versus 614 ± 27;

p=0.02) (Figure 4C and 4E). However, no significant differences in GP bridge numbers and

densities were observed between interventions (DMM versus non-operated, and loaded

versus non-loaded) at this time point in either the medial or lateral compartment. Anatomical

variations were observed however with clusters forming in the medial anterior of loaded

tibiae versus those more in the central and posterior of non-loaded tibiae (Figure 4C and 4D).

Similarly, in the DMM tibiae, GP bridges were more widespread across the tibiae than in

non-operated which were predominantly observed around the periphery (Figure 4A and 4B).

These results were consistent with the areal bridge density analysis. The mean areal bridge

densities were significantly greater in medial compared to the lateral compartment of DMM

(8.2 ± 0.9 versus 10.3 ± 0.9; p=0.01) and loaded tibiae (16 ± 0.9 versus 20.3 ± 0.6; p=0.003)

(Figure 4F). However, no significant differences in the mean areal bridge densities were

observed between interventions (DMM versus non-operated, and loaded versus non-loaded).

MicroCT analysis and 3D visualisation of SCB plate and epiphyseal trabecular bone

To establish whether aberrant GP dynamics are associated with the SCB plate and epiphyseal

trabecular bone abnormalities after the DMM surgery and mechanical loading in our mice,

we performed microCT analysis and determined local thickness of SCB plate and epiphyseal

trabecular bone. The SCB plate volume fraction (SCB BV/TV) was significantly higher in

medial compared to lateral compartment of DMM and non-operated tibiae (DMM: SCB

BV/TV DMM: 29.42 ± 3.1% versus 36.72 ± 4.5%, p=0.03, non-operated: 35.67 ± 0.7%

versus 42.57 ± 1.54%, p=0.004) (Figure 5A). No significant differences were observed

between DMM and non-operated tibia (Figure 5A), or in the epiphyseal trabecular bone

volume fraction (Tb. BV/TV; Figure 5B). Conversely, in the loaded and non-loaded tibiae,

epiphyseal trabecular bone volume fraction (Tb. BV/TV) was significantly increased in the

medial compared to lateral compartment (loaded: Tb. BV/TV 63.13 ± 1.2% versus 76.61 ±

2.7%, p=0.004, non-loaded: Tb. BV/TV 64.58 ± 1.4% versus 79.1 ± 3.2%, p=0.005) (Figure

5D). No significant differences were observed between interventions, or in the SCB BV/TV

(Figure 5C). Colour-coded SCB plate thickness analysis revealed anatomical variation in

SCB plate thickness between DMM and non-operated, and loaded and non-loaded plates

(Figure 6A and 6B). Differences in the non-loaded and loaded SCB plates were particularly

apparent (Figure 6B), and correlated with the clusters of higher density GP bridges previously

observed (Figure 4C and 4D). Epiphyseal trabecular bone thickness was not significantly

altered in response to either invasive or non-invasive intervention (data not shown; Figure 6C

and 6D).

Discussion

This study reveals altered GP dynamics in both surgical (DMM) and non-invasive loading

murine *in vivo* models of OA, which with findings described above may indicate an

endochondral defect in AC and GP cartilage in these mouse models of OA. Our data show

changes in AC of the knee joints of these mice consistent with the aberrant deployment of

hypertrophic chondrocytes. This is associated with disrupted GP morphology, increased GP

chondrocyte differentiation indicated by widespread expression of chondrocyte hypertrophy

markers and increased GP zone widths. Moreover, we have discovered enriched GP bony

bridging indicative of premature GP fusion and accelerated growth cessation in the medial

compartment of tibiae of these mice. These bridging events are spatially correlated with

increases in SCB thickness. These data reveal that altered GP dynamics and spatial

differences in GP bridging may contribute to an anatomical variation in vulnerability to OA

development in surgical and loaded murine models of OA.

The STR/ort OA murine model is predisposed to developing spontaneous idiopathic OA

whilst its nearest available parental strain, the CBA mouse, has a very low susceptibility

which makes them effective controls for the studies[17]. We have previously shown that

aberrant deployment of transient chondrocyte behaviour, consistent with re-initiation of

endochondral processes, occurs in uncalcified AC of STR/ort mouse knee joints compared to

CBA controls[8]. Here we extend these studies to look at the expression of transient markers in

other stratifications of OA. Surgically induced DMM model is widely used for target

validation studies or evaluation of the pathophysiological roles of many molecules in OA.

Following DMM, medial displacement of the medial meniscus in a mouse knee joint provides

a smaller area to transmit the weight-bearing forces and leads to an increased local

mechanical stress[18]. Whereas, in a cyclic AC tibial compression model, the non-invasive

dynamic mechanical loading applied to the mouse tibia through the knee and ankle joints,

modifies AC structure locally through a mechanoadaptive homeostatic response contributing

to OA development[19]. Our findings in both surgical and non-invasive loading C57BL/6

mouse models are consistent with the evidence of the role of chondrocyte phenotype

alterations in OA pathology. Hypertrophic chondrocytes in the calcified cartilage and GP of

the healthy joints express Col10a1[17–19]. The calcified cartilage acts to protect the uncalcified

AC through maintaining its ECM in an unmineralised state and the stability of the AC.

However, hypertrophic differentiation of these chondrocytes contributes to AC matrix

degradation, calcification and vascular invasion resulting in the demise of the AC[23].

Consistent with this, the expression of Col10a1, as examined using immunohistochemistry,

has been observed throughout AC in the joints of our both DMM and loaded mice. Further,

the higher expression level of another marker of chondrocyte hypertrophy; MMP13 has been

detected in superficial and uncalcified chondrocytes in the AC of DMM and loaded mice

compared to their non-operated and non-loaded left knee joints. Indeed, cartilage degradation

observed in OA has been attributed to an elevated production of proteolytic enzymes among

which MMP13 has a major role[24,25]. Studies using transgenic mice deficient in catabolic

transcription factors that induce hypertrophic differentiation revealed that animals were

protected against surgically and chemically induced OA further highlighting the role of

transient chondrocytes in AC degradation[26,27].

The results of the present study indicate aberrant widespread expression of Col10a1 and

MMP13 in the GP of DMM and loaded tibiae of C56BL/6 mice, compared to the GP of non-

operated and non-loaded control knees. Indeed, the histomorphometric analysis also revealed

that these mice display a significantly enlarged zone of proliferative and hypertrophic

chondrocytes, compared with the chondrocytes in the GP of control tibiae of both models.

Similarly, a significantly increased cumulative GP width was observed in these models.

Longitudinal bone growth is determined by the modifying number of chondrocytes in the

proliferative zone of GP, rate of their proliferation, the extent of chondrocyte hypertrophy

and controlled synthesis and degradation of ECM throughout the GP[28]. Altered growth rate

and mechanical modulation of GP function appear to result from complex interactions of

changes in the states of these chondrocytes, as does the rate of growth plate closure due to the

formation of bone bridges forming and spanning the width of the GP.

Our report of increased GP chondrocyte dynamics in these DMM and loaded C57BL/6 mice

is further strengthened by our data acquired using a 3D quantification method of bony

bridging across the tibial epiphysis, which showed premature GP closure in the medial

compartment of the tibiae in all mice examined, regardless of intervention. This was

indicated by significantly enriched spatial localisation of GP bone bridging clustering and

number in the medial compared to the lateral compartments of DMM mice compared to non-

operated mice. This is consistent with our previous work in the STR/Ort mouse model of

spontaneous OA in which we observed similar spatial variations. We postulate that the

formation of these bridges may be accelerated by local factors like instigating altered

mechanoadaptive response and that these spatial variations in GP bridging may disclose the

anatomical vulnerability to OA. These findings are supported by the previous studies

suggesting associations between local mechanical stress caused by medial displacement of

the medial meniscus[29] or cyclic tibial AC compression[19] and GP function in C57BL/6 mice.

They are further supported by our previous work in which we revealed, by finite element

modelling, that GP bridges act to dissipate stresses upon loading to the overlying SCB and

thus suggest that this contributes to OA seen in these models. These studies have, for the first

time, shown that accelerated GP closure is indicative of modified growth dynamics in the

right tibiae of both surgical and loading murine models of OA. This is consistent with our

microCT analysis of spatial variations in SCB plate thickening in these animals, and its

correlation to GP bridges. We observed statistically significant changes in subchondral bone

parameters in medial compared to lateral compartment including subchondral bone volume

fraction in the tibiae of DMM mice and epiphyseal trabecular bone volume fraction in the

tibiae of loaded mice. It was previously shown that SCB thickening may be intensified by

adjacent AC lesions following mechanical trauma thus indicating a special link between

changes in SCB architecture and AC lesions in OA pathology[14].

However, the limitation of this study is that we only examined the GP at one specific time

point for each model and therefore more time points are required to understand how these GP

bridges temporally affect SCB changes and OA pathology. With its controllability, the

intermittent non-invasive mechanical loading model will allow in future to distinguish

between short- and long-term effects of various cyclic loading regimens on subchondral and

trabecular bone parameters as well as AC integrity, GP dynamics and correlate these to

initiation and progression of human OA. Indeed, it is known that in the loading model, the

short-term intervention is not sufficient to induce changes in subchondral thickness and thus

longer intervention times for both the DMM and loading models may also prove useful in

pursuit of understanding these relationships[14].

Taken together, our studies indicate that accelerated GP chondrocyte dynamics and bridging

events may contribute to OA pathology in both surgical and loading C57BL/6 mouse OA

models and that similar osteoarthritic pathological changes happen in different *in vivo* models

of secondary (post-traumatic) OA[18] through inducing direct (DMM) or indirect (mechanical

loading) injuries to the joints. The GP bony bridging analysis may signify accelerated

cartilage-bone transition and growth cessation in these affected bones advancing our

understanding of GP closure mechanisms and how these contribute to the health of the joint.

Further, our work yields more insights into the changes in the micro-mechanical environment

of the GP and cells, specifically chondrocytes, within the GP.

Nonetheless, whether modified GP trajectories are prevalent in human OA is yet to be

defined and studies determining associations between GP dynamics during adolescence and

OA development in human patients would allow elucidation of the pathogenesis of OA and

will ultimately enable the development of novel and specific therapeutic interventions.

Acknowledgements

The authors would like to thank The Roslin Institute of The University of Edinburgh for the

assistance with the care of animal models in this study.

Author contributions

All authors contributed to the study design, analysis and interpretation of the data, drafted,

critically reviewed, edited and approved the version of the manuscript for publication.

Role of the funding source

We are grateful to Medical Research Council (to KAS; MR/R022240/1), Tenovus Scotland,

the Swedish Research Council (to AET; 2013–455) for funding. The funding sources did not

influence on the study design, data collection, analyses, interpretation, writing or submission

of the manuscript.

Conflict of interest

The research leading to these results has received technical support from 3Dmagination Ltd,

Didcot, UK. KM is co-founder and director of 3Dmagination Ltd in Oxfordshire, UK, a

company which provides training and consultancy in 3D and 4D imaging.

References

1.   Murray CJ., Lopez AD. Global Burden of Disease and Injur Y Series the Global

Burden of Disease. *Oms*. 1996:1-46. doi:10.1088/1742-6596/707/1/012025

2.   Goldring MB, Goldring SR. Osteoarthritis. *J Cell Physiol*. 2007;213(3):626-634.

doi:10.1002/jcp.21258

3.   Altman R, Asch E, Bloch D, et al. Development of criteria for the classification and

reporting of osteoarthritis: Classification of osteoarthritis of the knee. *Arthritis Rheum*.

1986;29(8):1039-1049. doi:10.1002/art.1780290816

4.   Pacifici M, Koyama E, Shibukawa Y, et al. Cellular and molecular mechanisms of

synovial joint and articular cartilage formation. *Ann N Y Acad Sci*. 2006;1068(1):74-

86. doi:10.1196/annals.1346.010

5.   Staines KA, MacRae VE, Farquharson C. The importance of the SIBLING family of

proteins on skeletal mineralisation and bone remodelling. *J Endocrinol*.

2012;214(3):241-255. doi:10.1530/JOE-12-0143

6.   Parfitt AM. Misconceptions (1): Epiphyseal fusion causes cessation of growth. *Bone*.

2002;30(2):337-339. doi:10.1016/S8756-3282(01)00668-8

7.   Glatt V, Canalis E, Stadmeyer L, Bouxsein ML. Age-related changes in trabecular

architecture differ in female and male C57BL/6J mice. *J Bone Miner Res*.

2007;22(8):1197-1207. doi:10.1359/jbmr.070507

8.   Staines KA, Madi K, Mirczuk SM, et al. Endochondral Growth Defect and

Deployment of Transient Chondrocyte Behaviors Underlie Osteoarthritis Onset in a

Natural Murine Model. *Arthritis Rheumatol*. 2016;68(4):880-891.

doi:10.1002/art.39508

9.   Madi K, Staines KA, Bay BK, et al. In situ characterization of nanoscale strains in

loaded whole joints via synchrotron X-ray tomography. *Nat Biomed Eng*. 2019;4:343–

354. doi:10.1038/s41551-019-0477-1

10. Glasson BVSc SS, Blanchet BS TJ, Morris DVM EA. The surgical destabilization of

the medial meniscus (DMM) model of osteoarthritis in the 129/SvEv mouse.

Osteoarthr Cart. 2007;15:1061-1069. doi:10.1016/j.joca.2007.03.006

11. Staines KA, Ikpegbu E, Törnqvist AE, et al. Conditional deletion of E11/podoplanin in

bone protects against load-induced osteoarthritis. *BMC Musculoskelet Disord*.

2019;20(1):1-11. doi:10.1186/s12891-019-2731-9

12. Sugiyama T, Price JS, Lanyon LE. Functional adaptation to mechanical loading in

both cortical and cancellous bone is controlled locally and is confined to the loaded

bones. *Bone*. 2010;46(2):314-321. doi:10.1016/j.bone.2009.08.054

13. Poulet B, Hamilton RW, Shefelbine S, Pitsillides AA. Characterizing a novel and

adjustable noninvasive murine joint loading model. *Arthritis Rheum*. 2011;63(1):137-

147. doi:10.1002/art.27765

14. Poulet B, de Souza R, Kent A V, et al. Intermittent applied mechanical loading induces

subchondral bone thickening that may be intensified locally by contiguous articular

cartilage lesions. *Osteoarthr Cartil*. 2015;23(6):940-948.

doi:10.1016/j.joca.2015.01.012

15. Staines KA, Madi K, Javaheri B, Lee PD, Pitsillides AA. A computed

microtomography method for understanding epiphyseal growth plate fusion. *Front

Mater*. 2018;4. doi:10.3389/fmats.2017.00048

16.   Chambers MG, Kuffner T, Cowan SK, Cheah KSE, Mason RM. Expression of

434   collagen and aggrecan genes in normal and osteoarthritic murine knee joints.

435   *Osteoarthr Cartil*. 2002;10(1):51-61. doi:10.1053/joca.2001.0481

436   17.   Poulet B, Westerhof TAT, Hamilton RW, Shefelbine SJ, Pitsillides AA. Spontaneous

437   osteoarthritis in Str/ort mice is unlikely due to greater vulnerability to mechanical

438   trauma. *Osteoarthr Cartil*. 2013;21(5):756-763. doi:10.1016/j.joca.2013.02.652

439   18.   Samvelyan HJ, Hughes D, Stevens C, Staines KA. Models of Osteoarthritis: Relevance

440   and New Insights. *Calcif Tissue Int*. 2020. doi:10.1007/s00223-020-00670-x

441   19.   Poulet B, Hamilton RW, Shefelbine S, Pitsillides AA. Characterizing a novel and

442   adjustable noninvasive murine joint loading model. *Arthritis Rheum*. 2011;63(1):137-

443   147. doi:10.1002/art.27765

444   20.   Atkins GJ, Rowe PS, Lim HP, et al. Sclerostin is a locally acting regulator of late-

445   osteoblast/preosteocyte differentiation and regulates mineralization through a MEPE-

446   ASARM-dependent mechanism. *J Bone Miner Res*. 2011;26(7):1425-1436.

447   doi:10.1002/jbmr.345

448   21.   Lai Y, Bai X, Zhao Y, et al. ADAMTS-7 forms a positive feedback loop with TNF-α

449   in the pathogenesis of osteoarthritis. *Ann Rheum Dis*. 2014;73(8):1575-1584.

450   doi:10.1136/annrheumdis-2013-203561

451   22.   Lu Y, Ding M, Li N, et al. Col10a1-runx2 transgenic mice with delayed chondrocyte

452   maturation are less susceptible to developing osteoarthritis. *Am J Transl Res*.

453   2014;6(6):736-745.

23.   Ea HK, Nguyen C, Bazin D, et al. Articular cartilage calcification in osteoarthritis:

Insights into crystal-induced stress. *Arthritis Rheum*. 2011;63(1):10-18.

doi:10.1002/art.27761

24. Ortega N, Behonick DJ, Werb Z. Matrix remodeling during endochondral ossification.

Trends Cell Biol. 2004;14(2):86-93. doi:10.1016/j.tcb.2003.12.003

25. Stickens D, Behonick DJ, Ortega N, et al. Altered endochondral bone development in

matrix metalloproteinase 13-deficient mice. *Development*. 2004;131(23):5883-5895.

doi:10.1242/dev.01461

26. Saito T, Fukai A, Mabuchi A, et al. Transcriptional regulation of endochondral

ossification by HIF-2α during skeletal growth and osteoarthritis development. *Nat*

Med. 2010;16(6):678-686. doi:10.1038/nm.2146

27. Yang S, Kim J, Ryu JH, et al. Hypoxia-inducible factor-2α is a catabolic regulator of

osteoarthritic cartilage destruction. *Nat Med*. 2010;16(6):687-693.

doi:10.1038/nm.2153

28. Breur GJ, Lapierre MD, Kazmierczak K, Stechuchak KM, McCabe GP. The domain of

hypertrophic chondrocytes in growth plates growing at different rates. *Calcif Tissue*

Int. 1997;61(5):418-425. doi:10.1007/s002239900358

29. Huang H, Skelly JD, Ayers DC, Song J. Age-dependent Changes in the Articular

Cartilage and Subchondral Bone of C57BL/6 Mice after Surgical Destabilization of

Medial Meniscus. *Sci Rep*. 2017;7:1-9. doi:10.1038/srep42294

 Figure legends

 Figure 1A, 1B. Immunohistochemical labelling for type X collagen (Col10a1) and matrix

 metalloproteinase (MMP13) in the articular cartilage of non-operated and DMM (A), or non-

 loaded and mechanically loaded (B) knee joints of C57BL/6 male mice. Arrows indicate

 examples of positive staining in the tibiae. Images are representative of results in 3 individual

 mice. Scale bar = 200μm

 Figure 2A, 2B. Immunohistochemical labelling for type X collagen (Col10a1) and matrix

 metalloproteinase (MMP13) in the growth plate of DMM and non-operated (A), or

 mechanically loaded and non-loaded (B) knee joints of C57BL/6 male mice. Arrows indicate

 examples of positive staining in the tibiae. Images are representative of results in 3 individual

 mice. Scale bar = 200μm. PZ = proliferative zone; HZ = hypertrophic zone

 Figure 3. Growth plate zone width of non-operated and DMM, or non-loaded and loaded

 knee joints of C57BL/6 male mice. Ten measurements per section were obtained along the

 length of the growth plate of the tibiae (n=4 mice for each experimental group). PZ =

 proliferative zone; HZ = hypertrophic zone **** $p<0.0001$

 Figure 4A, 4B, 4C, 4D, 4E, 4F. Location and areal densities of bridges across the growth

 plate projected on the medial (M) and lateral (L) tibial joint surface in non-operated (A),

 DMM (B), non-loaded (C) and loaded (D) tibiae of mice at 16 and 18 weeks of age, number

 of bridges in lateral and medial tibiae of non-operated compared to DMM and non-loaded

 compared to loaded tibiae of mice (E), Areal density (d) of bridges in medial compared to

 lateral tibiae, defined as the number of bridges per 256 mm x 256 mm window of non-

 operated and DMM or non-loaded and loaded knee joints (F). Bars represent mean ± SEM.

 Group sizes were $n = 8$ for non-operated and DMM-operated mice and $n = 4$ for non-loaded

and loaded mice. * $p<0.05$ ** $p<0.01$ *** $p<0.001$

Figure 5A, 5B, 5C, 5D. MicroCT analysis of the epiphyseal region of the lateral and medial

tibiae in non-operated controls and DMM-operated knee joints subchondral bone volume

fraction (SCB BV/TV) (A) and epiphyseal trabecular bone volume fraction (Tb. BV/TV) (B).

MicroCT analysis of the epiphyseal region of the medial and lateral tibiae in non-loaded

controls and loaded knee joints subchondral bone volume fraction (SCB BV/TV) (C) and

epiphyseal trabecular bone volume fraction (Tb. BV/TV) (D). Bars represent mean ± SEM.

Group sizes were $n = 8$ for non-operated and DMM-operated mice and $n = 4$ for non-loaded

and loaded mice. * $p<0.05$ ** $p<0.01$

Figure 6A, 6B, 6C, 6D. Representative colour coded images of lateral and medial

subchondral bone plate thickness of non-operated and DMM-operated tibiae of mice (A).

Representative colour coded images of lateral and medial subchondral bone plate thickness of

non-loaded and loaded tibiae of mice (B). Representative colour coded images of epiphyseal

trabecular bone thickness of non-operated and DMM-operated (C), or non-loaded and loaded

tibiae of mice (D)