**Characterisation of growth plate dynamics in murine models of osteoarthritis**

Hasmik J. Samvelyan[1,2], Kamel Madi[3], Anna E. Törnqvist[4,5], Behzad Javaheri[6], Katherine A.

Staines[1,2]

[1] School of Pharmacy and Biomolecular Sciences, University of Brighton, Brighton, UK

[2] Centre for Stress and Age-Related Disease, University of Brighton, Brighton, UK

[3] 3Dmagination Ltd, Harwell Campus, Didcot, UK

[4] Rheumatology and Bone Diseases Unit, Centre for Genomic and Experimental Medicine,

MRC Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK

[5] Centre for Bone and Arthritis Research, University of Gothenburg, Gothenburg, Sweden

[6] School of Mathematics, Computer Science and Engineering, City University of London,

London, UK

Corresponding author

Hasmik Jasmine Samvelyan, University of Brighton, Lewes Road, Brighton BN2 4GJ

Email: h.samvelyan@brighton.ac.uk Tel: 01273 641 671

**Abstract**

*Background:* The purpose of this study was to investigate growth plate dynamics in surgical and loading murine models of osteoarthritis, to understand whether abnormalities in these dynamics predict osteoarthritis vulnerability.

*Methods:* 8-week-old C57BL/6 male mice underwent destabilisation of medial meniscus (DMM) ($n = 8$) surgery in right knee joints. Contralateral left knee joints had no intervention (controls). In 16-week-old C57BL/6 male mice ($n = 6$), osteoarthritis was induced using non-invasive mechanical loading of right knee joints with peak force of 11N. Non-loaded left knee joints were internal controls. Chondrocyte transiency in tibial articular cartilage and growth plate was confirmed by histology and immunohistochemistry. Tibial subchondral bone parameters were measured using microCT and correlated to 3D GP bridging analysis.

*Results:* Higher expression of chondrocyte hypertrophy markers; Col10a1 and MMP13 were observed in tibial articular cartilage chondrocytes of DMM and loaded mice. In tibial growth plate, Col10a1 and MMP13 expressions were widely dispersed in a significantly enlarged zones of proliferative and hypertrophic chondrocytes in DMM ($p=0.002$ and $p<0.0001$, respectively) and loaded (both $p<0.0001$) tibiae of mice compared to their controls. 3D quantification revealed enriched growth plate bridging and higher bridge densities in medial compared to lateral tibiae of DMM and loaded knee joints of the mice. Growth plate dynamics were associated with higher subchondral bone volume fraction (BV/TV; %) in medial tibiae of DMM and loaded knee joints and epiphyseal trabecular bone volume fraction in medial tibiae of loaded knee joints.

*Conclusions:* The results confirm articular cartilage chondrocyte transiency in a surgical and loaded murine model of osteoarthritis. Herein, we reveal for the first time spatial variation of

45　growth plate bridging in surgical and loaded osteoarthritis models and how these may

46　contribute to anatomical variation in vulnerability of osteoarthritis development.

47　**Keywords:** articular cartilage, chondrocyte, growth, murine model, osteoarthritis,

48　subchondral bone

## Background

Osteoarthritis (OA) is a chronic musculoskeletal disease and a leading cause of disability and

major healthcare costs in the world. It is estimated that worldwide 10% of men and 18% of

women aged over 60 years have symptomatic OA (1) and, therefore, the predicted increase in

the ageing population and longevity will result in a greater occurrence of the disease. OA is a

complex disease in which the pathogenesis, cellular and molecular mechanisms of initiation

and progression are not completely understood. It is characterised by progressive loss of

articular cartilage (AC), formation of osteophytes, subchondral bone (SCB) sclerosis,

synovial proliferation, inflammation and lax tendons. These can ultimately lead to a loss of

joint function, pain, reduced mobility and disability (2). Despite the significant healthcare and

economic burden there are few non-invasive therapies available to patients. Therefore,

understanding the pathogenesis of OA and defining the molecular mechanisms underpinning

AC degeneration can lead to the development of successful targeted and effective disease-

modifying treatments.

Primary OA is described as naturally occurring OA affecting one joint (localised) or three or

more joints (generalised), while secondary OA is associated with various causes and risk

factors leading to the disease including trauma, obesity, diabetes, metabolic bone and

congenital disorders (3). AC degeneration is one of the main hallmarks of OA and previous

research has largely sought to identify mechanisms underpinning its deterioration. Fully

developed, uncalcified AC is populated by a single resident cell chondrocytes, which

maintain a stable phenotype characterised by small cell size and expression of tenascin-C (4).

The inherent stability of AC chondrocytes ensures that dynamic events are restricted to assure

lifelong articular integrity and healthy joint function. In contrast, epiphyseal growth plate

(GP) chondrocytes have a transient phenotype to ensure long bone development

(endochondral ossification) and growth. GP chondrocytes undergo a differentiation sequence

of proliferation, maturation and hypertrophy. The final stage of chondrocyte hypertrophy

enables mineralisation of the cartilage extracellular matrix, vascular invasion and subsequent

replacement of the mineralised cartilage anlagen with bone (5). These processes are coupled,

however, with sexual maturation the human GP undergoes progressive narrowing as bony

bridges form and span its width. This ultimately leads to complete GP closure and cessation

of human growth. Indeed, in humans the longitudinal bone growth stops with the onset of

puberty, the metaphysis then fuses with the epiphysis and the GP disappears (6). In mice,

longitudinal bone growth does not cease at sexual maturity instead it slows dramatically at

puberty, but the GPs do not completely fuse and disappear (7).

We have previously shown that in the STR/Ort mouse, a naturally occurring OA murine

model, AC chondrocytes transform from their inherently stable phenotype to a transient one,

characteristic of the chondrocytes in the GP. This was confirmed by immunolabelling for

chondrocyte hypertrophy markers; type X collagen (Col10a1) and matrix metalloproteinase

13 (MMP13)(8). Further, we revealed accelerated longitudinal bone growth, aberrant

expression of GP markers (Col10a1 and MMP13) and increased GP chondrocyte maturation

in these mice. Consistent with this, using a novel synchrotron computed tomography method

we revealed enriched GP bone bridging in STR/Ort mouse tibiae indicative of advanced GP

closure which may underpin OA (8,9).

Despite this, interlinks between the differing chondrocyte phenotypes in the AC and the GP,

and the precise contribution that the GP and its fusion mechanisms may play in underpinning

OA vulnerability is not yet understood. The aim of this study was to investigate whether

surgically (destabilisation the medial meniscus [DMM]) or non-surgically (mechanical

loading) induced OA onset in C57BL/6 mice is linked to altered GP dynamics. Indeed,

understanding this will inform strategies for maintaining musculoskeletal health in ageing by

potentially identifying whether GP dynamics may predict who is at risk of OA in later life

and ultimately developing targeted OA treatments.

**Methods**

*Animals*

Male C57BL/6 wild type mice at 7 weeks of age (young adult) were obtained from Charles

River Laboratories Inc. (Margate, UK). The mice were acclimatised to their surroundings for

seven days. All mice were allowed free access to water and maintenance diet ad libitum

(Special Diet Services, Witham, UK) in a 12-hour light/dark cycle at a room temperature of

$21 \pm 2^0$C and relative humidity of $55 \pm 10\%$.

*The destabilisation of the medial meniscus (DMM)*

8-week old wild type C57/BL6 male mice underwent DMM surgeries to induce OA-like

changes in the right knee joints under isoflurane-induced anaesthesia ($n = 8$/group). We chose

not to performed SHAM (placebo) surgery on the left contralateral knee of the animals based

upon animal welfare grounds in keeping with the 3Rs, since it was previously shown that

there is no difference in OA scores between SHAM-operated and non-operated knee joints

(10,11).  Animals were monitored for unexpected adverse effects to reduce suffering and

distress. Following transection of the medial meniscotibial ligament (MMTL) to destabilise

the medial meniscus, the skin was closed and anaesthesia reversed (10). Eight weeks later,

mice were culled by exsanguination and confirmation of death by cervical dislocation. The

knee joints of all the mice were dissected, fixed in 4% paraformaldehyde for 24 hours at $4^o$C,

and then stored in 70% ethanol.

*In vivo loading of the knee joint*

The right knee joints of 16-week-old wild type C57BL/6 male mice ($n = 6$) were subjected to

non-invasive, dynamic axial mechanical loading under the isofluorane-induced anaesthesia

(liquid isofluorane was vaporised to a concentration of 4% and maintained at a concentration

of 2% with oxygen) for 7 min/day, 3 alternate days a week for 2 weeks according to the

protocols described in the previous studies (12,13). The left knee joints were non-loaded

internal controls in these animals. Briefly, using a servo-electric materials testing machine

(Electroforce 3100, Bose, UK), axial compressive loads were applied through the right knee

joint via customised concave cups which held the knee and ankle joints flexed and the tibiae

vertically.

The tibia was held in place by continuous static preload of 0.5N onto which dynamic loads

were superimposed in a series of 40 trapezoidal shaped waveform cycles with steep up and

down ramps and a peak force of 11N for 0.05 seconds (0.025 seconds rise and fall time; 9.9

seconds baseline hold time between periods of peak loading). The right and left knees were

dissected 3 days after the final loading episode. Mice were culled by exsanguination and

confirmation of death by cervical dislocation. Knee joints were fixed in 4%

paraformaldehyde for 24 hours at 4$^{o}$C before being stored in 70% ethanol.

*Micro-computed (microCT) tomography and 3-dimensional (3D) bridging analysis*

Scans were performed with an 1172 X-Ray microtomograph (Bruker MicroCT, Kontich,

Belgium) to evaluate the SCB and GP bridging. High-resolution scans with an isotropic voxel

size of 5 µm were acquired (50 kV, 200 µA, 0.5 mm aluminium filter, 0.6° rotation angle).

The projection images were reconstructed and binarised with a threshold of 0 to 0.16, ring

artefact reduction was set at 10 and beam hardening correction at 0% using the SkyScan

NRecon software package (v1.6.9.4, Bruker MicroCT). The images then were realigned

vertically using DataViewer software (v1.5.1.2 64-bit, Bruker MicroCT) to ensure similar

orientation for analysis. Hand-drawn regions of interests (ROI) of the SCB and epiphyseal

trabecular bone for each tibial lateral and medial compartments were selected (14). The

structural parameters of tibial SCB plate and epiphyseal trabecular bone were calculated

using 3D algorithms of SkyScan CTAn software (Bruker MicroCT) including SCB (SCB

BV/TV; %) and trabecular bone volume fraction (Tb. BV/TV; %) and correlated to GP

bridging analysis using a 3D synchrotron-computed tomography quantification method as

previously described (15). Briefly, microCT scans of the tibiae were segmented using a

region-growing algorithm within the Avizo® (V8.0, VSG, Burlington, VT, USA) software.

The central points of each bony bridges were identified and projected on the tibial joint

surface. The distribution of the areal number density of bridges (N, the number of bridges per

256 µm $\times$ 256 µm window; $d = m/V$) is then calculated and superimposed on the tibial joint

surface (each bridge has a colour that represents the areal number density at the bridge

location). The SCB plate and epiphyseal trabecular bone thickness (Th; mm) was determined

and colour-coded thickness images were generated using the Avizo® software(15).

*Histological analysis*

The left and right knee joints of all the mice were decalcified in 10%

ethylenediaminetetraacetic acid (EDTA) solution, wax-embedded at Leica EG1160 Tissue

Embedding Station and 6 µm coronal sections cut using Leica RM2135 manual microtome.

*Immunohistochemistry*

Immunohistochemical analysis of chondrocyte transiency markers in tibial AC and GP was

performed on 6 µm coronal sections from the middle region of the knee joint, using anti-

matrix metalloproteinase 13 (anti–MMP13) (1:200 dilution; Abcam) or anti-collagen type X

(anti-Col10a1) (1:100 dilution; Abcam) antibodies. As a control, an equal concentration of

rabbit IgG was used. For immunohistochemical localisation of MMP13 and Col10a1,

sections were dewaxed in xylene and rehydrated. Sections were incubated at 37 °C for

30 min in 1mg/ml trypsin for antigen demasking. Endogenous peroxidases were blocked by

treatment with 0.3% $H_2O_2$ in methanol (Sigma) for 30 min at room temperature. The

Vectastain ABC universal detection kit (Vector Laboratories, Peterborough, UK) was used to

detect the biotinylated secondary antibody (Anti-Mouse IgG Reagent) after incubation for 30

min at room temperature according to the manufacturer's instructions. Diaminobenzidine

(DAB) solution used to detect the location of antigens. The sections were finally dehydrated,

counterstained with haematoxylin and mounted in DePeX. All sections to be compared were

immunolabelled at the same time to standardise conditions and minimise any differences in

antibody incubation times. For each group, immunolabelling was performed on four

individual animals per experimental group and representative images taken using a light

microscope.

*Growth plate (GP) zone analysis*

The width of the GP proliferating and hypertrophic zones, as well as the total GP width, were

measured at 10 different points along the length of the GP in 6μm coronal sections from the

middle region of the knee joint of four individual animals per experimental group, using a

light microscope and ImageJ software.

*Statistical analysis*

All analyses were performed with GraphPad Prism software 6.0f version (GraphPad Inc, La

Jolla, CA, USA) using a two-sided 0.05 level of significance. All analyses were conducted

blindly to minimise the effects of subjective bias. The results were presented as the mean ±

standard error of the mean (SEM). The Normal distribution of data was assessed using the

Shapiro-Wilk normality test. For comparing two groups (experimental with control, or medial

with the lateral compartment of tibiae), two-tail Student's *t*-test (paired or unpaired) was

used. For comparing more than two groups, two-way ANOVA (analysis of variance) was

used with Tukey post-hoc test.

**Results**

*Transient chondrocyte behaviour in the tibial AC of DMM and loaded C57BL/6 young adult*

*male mice*

We first sought to confirm whether loss of AC in C57BL/6 mice with surgically and loading

induced OA was associated with the expression of markers of transient chondrocyte

phenotype. In accordance with previous studies, immunohistochemistry analysis showed

higher expression levels of well-established chondrocyte hypertrophy markers; Col10a1 and

MMP13, observed in tibial AC of C57BL/6 mice that have undergone DMM surgery or

mechanical loading compared with non-operated and non-loaded control left tibiae,

respectively (Fig. 1a and b). The expression pattern of Col10a1 was largely restricted to

hypertrophic chondrocytes in the uncalcified zone of the AC of unaffected condyles of non-

operated (Fig. 1a) and non-loaded (Fig. 1b) mouse left joints as expected (16). Whereas, the

immunolabeling of Col10a1 was more widespread throughout the extracellular matrix (ECM)

of the AC in affected right joints of DMM (Fig. 1a) and loaded (Fig. 1b) mice. Similarly,

immunohistochemistry analysis showed positive MMP13 labelling in both superficial and

deep articular chondrocytes in the right joints of DMM and loaded C57BL/6 male mice

compared to the control knee joints (Fig. 1a and b). These findings confirm an aberrant

deployment of transient chondrocytes in uncalcified AC.

*Dysfunctional GP dynamics in DMM and loaded tibiae of C57BL/6 young adult male mice*

To determine the GP phenotype in these OA models, we first completed histological

assessment of GP zones. Our analysis revealed significantly enlarged proliferative and

hypertrophic zones of chondrocytes in both DMM ($p=0.002$ and $p<0.0001$, respectively) and

loaded (both $p<0.0001$) tibiae of C57BL/6 mice compared to their controls, and significantly

increased total GP width (both $p<0.0001$) (Fig. 2a). This is suggestive of aberrant GP

dynamics in these OA models.

We next determined the expression of known chondrocyte transiency markers in the GPs. In

the tibial GP of mice with surgically and loading induced OA, Col10a1 expression was more

greatly and widely dispersed throughout the zones of proliferative and hypertrophic

chondrocytes compared with their controls (Fig. 2b and c). Indeed, immunolabeling for

Col10a1 revealed the expected localisation in the GP of non-operated and non-loaded mouse

tibiae, limited primarily to the hypertrophic zone and underlying adjacent metaphyseal bone

(Fig. 2b and c). This disrupted the distribution of a GP zone marker was also evident for

MMP13 in the GP of DMM and loaded C57BL/6 mouse tibiae compared to their controls

(Fig. 2b and c). Together, the results may indicate associations between dysfunctional GP

morphology and marker expression, and disease development in these murine models of OA.

*Associations between GP bridging in DMM and loaded tibiae of C57BL/6 young adult male*

*mice and OA development*

To further correlate aberrant longitudinal GP dynamics, GP bridging and OA development in

these models, we used our newly developed 3D method to quantify GP bone bridges across

the tibial epiphysis of DMM and loaded mice for the first time (Fig. 3). 3D quantification

revealed a significantly higher number of GP bridges in medial compared to lateral tibiae that

underwent DMM surgeries (196 $\pm$ 33 versus 306 $\pm$ 32; $p$=0.03) (Fig. 3b and e). This

significant difference was not observed in non-operated tibiae (254 $\pm$ 32 versus 326 $\pm$ 32;

$p$=0.14) (Fig. 3a and e). Similarly, significantly enriched GP bridging was evident in the

medial compartment of loaded tibiae in comparison to the lateral compartment (539 $\pm$ 36

versus 745 $\pm$ 18; $p$=0.0009) (Fig. 3d and e), and in those of non-loaded tibiae, although less

pronounced than in the loaded right knee joints (596 $\pm$ 22 versus 726 $\pm$ 44; $p$=0.04) (Fig. 3c

and e). However, no significant differences in GP bridge numbers and densities were

observed between interventions (DMM versus non-operated, and loaded versus non-loaded)

at this time point in either the medial or lateral compartment. These results were consistent

with the areal bridge density analysis. The mean areal bridge densities were significantly

greater in medial compared to the lateral compartment of DMM ($8.2 \pm 0.9$ versus $10.3 \pm 0.9$;

$p$=0.01) and loaded tibiae ($16.3 \pm 0.6$ versus $20.6 \pm 0.5$; $p$<0.0001) (Fig. 3f). However, no

significant differences in the mean areal bridge densities were observed between

interventions (DMM versus non-operated, and loaded versus non-loaded).

Anatomical variations were observed however in all samples, with focal high density clusters

forming in the anterior region of the medial loaded tibiae versus those more in the central and

posterior of non-loaded tibiae (Fig. 3c and d). Similarly, in the DMM tibiae, GP bridges were

more widespread across the tibiae than in non-operated which were predominantly observed

around the periphery (Fig. 3a and b). Together, these results suggest that murine OA models

exhibit spatial variations in GP bridging in comparison to controls.

*MicroCT analysis and 3D visualisation of SCB plate and epiphyseal trabecular bone*

To establish whether spatial variations in GP bridging are associated with the SCB plate and

epiphyseal trabecular bone abnormalities after the DMM surgery and mechanical loading in

our mice, we performed microCT analysis and determined local thickness of SCB plate and

epiphyseal trabecular bone. The SCB plate volume fraction (SCB BV/TV) was significantly

higher in medial compared to lateral compartment of DMM and non-operated tibiae (DMM:

SCB BV/TV $29.4 \pm 3.1\%$ versus $36.7 \pm 4.5\%$, $p$=0.03, non-operated: SCB BV/TV $35.7 \pm$

$0.7\%$ versus $42.6 \pm 1.5\%$, $p$=0.004) (Fig. 4a). No significant differences were observed

between DMM and non-operated tibia (Fig. 4a), or in the epiphyseal trabecular bone volume

fraction (Tb. BV/TV; Fig. 4b). Conversely, in the loaded and non-loaded tibiae, epiphyseal

trabecular bone volume fraction (Tb. BV/TV) was significantly increased in the medial

compared to lateral compartment (loaded: Tb. BV/TV 60.7 ± 1.8% versus 73.4 ± 2.8%,

$p$=0.005, non-loaded: Tb. BV/TV 61.4 ± 2.2% versus 77.4 ± 2.5%, $p$=0.0005). No significant

differences were observed between interventions (Fig. 4d). The SCB plate volume fraction

(SCB BV/TV) was significantly higher in medial compared to lateral compartment of loaded

tibiae (40.1 ± 1.7% versus 47.1 ± 1.6%, $p$=0.02), but not that of non-loaded tibiae (42.1 ±

1.1% versus 45.8 ± 1.6%, $p$=0.34) or between interventions (Fig. 4c).

Colour-coded SCB plate thickness analysis revealed anatomical variation in SCB plate

thickness between DMM and non-operated, and loaded and non-loaded plates (Fig. 4e and

4f). Differences in the non-loaded and loaded SCB plates were particularly apparent (Fig. 4f),

and correlated with the clusters of higher density of GP bridges previously observed (Fig. 3c

and d). Epiphyseal trabecular bone thickness was not significantly altered in response to

either invasive or non-invasive intervention (data not shown; Fig. 4g and h).

**Discussion**

Together, the data presented herein reveals altered GP dynamics in both surgical (DMM) and

non-invasive loading murine *in vivo* models of OA. Our data confirm previous studies

showing changes in AC of the knee joints of these mice consistent with the aberrant

deployment of chondrocyte transiency, and reveal disrupted GP morphology, increased GP

chondrocyte differentiation indicated by widespread expression of chondrocyte hypertrophy

markers and increased GP zone widths in these models. Moreover, we have utilised a recently

developed 3D model to discover for the first time anatomical variations in GP bone bridging,

indicative of GP fusion, which are spatially correlated with increase in SCB thickness. These

data reveal that altered GP dynamics and spatial differences in GP bridging may contribute to

an anatomical variation in vulnerability to OA development in surgical and loaded murine

models of OA.

which with findings described above may indicate an endochondral defect in AC and GP

cartilage in these mouse models of OA. These findings build upon our previous work in the

STR/ort OA murine model which is predisposed to developing spontaneous idiopathic OA

whilst its nearest available parental strain, the CBA mouse, has a very low susceptibility

which makes them effective controls for the studies (17). We have previously shown that

aberrant deployment of transient chondrocyte behaviour, consistent with re-initiation of

endochondral processes, occurs in uncalcified AC of STR/ort mouse knee joints compared to

CBA controls (8). Here we extend these studies to look at the expression of transient markers

in other stratifications of OA using murine models. Surgically induced DMM model is widely

used for target validation studies or evaluation of the pathophysiological roles of many

molecules in OA. Following DMM, medial displacement of the medial meniscus in a mouse

knee joint provides a smaller area to transmit the weight-bearing forces and leads to an

increased local mechanical stress (18). Whereas, in a cyclic AC tibial compression model, the

non-invasive dynamic mechanical loading applied to the mouse tibia through the knee and

ankle joints, modifies AC structure locally through a mechanoadaptive homeostatic response

contributing to OA development (19). Hypertrophic chondrocytes in the calcified cartilage

and GP of the healthy joints express Col10a1 (17,19). The calcified cartilage acts to protect

the uncalcified AC through maintaining its ECM in an unmineralised state and the stability of

the AC. However, hypertrophic differentiation of these chondrocytes contributes to AC

matrix degradation, calcification and vascular invasion resulting in the demise of the AC

(23).  Consistent  with  this,  the  expression  of  Col10a1,  as  examined  using

immunohistochemistry, has been observed throughout the AC in the joints of our both DMM

and loaded mice. Further, the higher expression level of another marker of chondrocyte

hypertrophy; MMP13 has been detected in superficial and uncalcified chondrocytes in the

AC of DMM and loaded mice compared to their non-operated and non-loaded left knee

joints. Indeed, cartilage degradation observed in OA has been attributed to an elevated

production of proteolytic enzymes among which MMP13 has a major role (24,25). Studies

using transgenic mice deficient in catabolic transcription factors that induce hypertrophic

differentiation revealed that animals were protected against surgically and chemically

induced OA further highlighting the role of transient chondrocytes in AC degradation

(26,27).

The results of the present study indicate that in these murine OA models, there is a

significantly enlarged zone of GP proliferative and hypertrophic chondrocytes, compared

with the chondrocyte zones in the GP of control tibiae of both models. Similarly, a

significantly increased cumulative GP width was observed in these models. This is associated

with aberrant widespread expression of Col10a1 and MMP13 in the GP of DMM and loaded

tibiae of C56BL/6 mice, compared to the GP of non-operated and non-loaded control knees.

Longitudinal bone growth is determined by the modifying number of chondrocytes in the

proliferative zone of GP, rate of their proliferation, the extent of chondrocyte hypertrophy

and controlled synthesis and degradation of ECM throughout the GP (28). Altered growth

rate and mechanical modulation of GP function appear to result from complex interactions of

changes in the states of these chondrocytes, as does the rate of GP closure due to the

formation of bone bridges forming and spanning the width of the GP.

Our report of aberrant GP chondrocyte dynamics in these DMM and loaded C57BL/6 mice is

further strengthened by our data acquired using a 3D quantification method of bone bridging

across the tibial epiphysis (15). Here we show significantly enriched spatial localisation of

GP bone bridging clustering and number in the tibial medial compared to the lateral

compartments of DMM mice in comparison non-operated mice. We reveal focal clusters of

higher density bridges in different anatomical regions of the tibiae, which was correlated with

increased SCB thickness. This is consistent with our previous work in the STR/Ort mouse

model of spontaneous OA in which we observed similar spatial variations. We postulate that

the formation of these bridges may be accelerated by local factors like instigating altered

mechanoadaptive response and that these spatial variations in GP bridging may disclose the

anatomical vulnerability to OA. These findings are supported by the previous studies

suggesting associations between local mechanical stress caused by medial displacement of

the medial meniscus (29) or cyclic tibial AC compression (19) and GP function in C57BL/6

mice. They are further supported by our previous work in which we revealed, by finite

element modelling, that GP bridges act to dissipate stresses upon loading to the overlying

SCB and thus suggest that this contributes to OA seen in these models (15). Indeed the GP is

subject to a variety of mechanical forces placed upon it, and strain distributions are

inhomogeneous through the GP with the hypertrophic zone exhibiting low mechanical

stiffness. The precise interplay between GP bridging and mechanics has yet to be determines,

specifically in the context of OA and how this may increase OA vulnerability.

Identification of modified GP dynamics underpinning human OA would allow elucidation

into OA predisposition and ultimately enable the development of novel and specific

therapeutic interventions. Our recent work in the MRC National Survey of Health and

Development (NSHD) found that increased height in childhood was associated, albeit

modestly, with lower odds of knee OA at age 53 years, as was adult achieved height (30).

Concurrent with this, a recent study from offspring in the Avon Longitudinal Study of

Parents and Children (ALSPAC) found height tempo (corresponding to pubertal timing) to be

strongly associated with the hip shape models which may be related to hip future risk

OA(31). Further, several known OA susceptibility single nucleotide polymorphisms (SNPs)

have been associated with hip shape in perimenopausal women in the ALSPAC(32). Most

intriguingly, when combined with data from other cohorts, the eight SNPs independently

associated with hip shape were intimately associated with the process of endochondral

ossification (33). Together these data provide further tantalising evidence that there are

associations between GP dynamics during adolescence and OA development.

The limitation of this study is that we only examined the GP at one specific time point for

each model and therefore more time points are required to fully understand how these GP

bridges temporally affect SCB changes and OA pathology. With its controllability, the

intermittent non-invasive mechanical loading model will allow in future to distinguish

between short- and long-term effects of various cyclic loading regimens on SCB and

trabecular bone parameters as well as AC integrity, GP dynamics and allow correlation of

these to initiation and progression of human OA. Indeed, it is known that in the loading

model, the short-term intervention is not sufficient to induce significant changes in

subchondral thickness and thus longer intervention times for both the DMM and loading

models may also prove useful in pursuit of understanding these relationships (14).

**Conclusions**

Our studies indicate that aberrant GP dynamics may contribute to OA pathology in both

surgical and loading mouse OA models, and that similar GP-related osteoarthritic

pathological changes happen in different *in vivo* models of secondary (post-traumatic) OA

(18) through inducing direct (DMM) or indirect (mechanical loading) injuries to the joints.

We reveal for the first time spatial variation of GP bridging in these OA models which may

contribute to anatomical variation in vulnerability of OA development. Our GP bony bridging

analysis may signify accelerated cartilage-bone transition in these affected joints, advancing

our understanding of GP closure mechanisms and how these contribute to the health of the

joint. Further, our work yields more insights into the changes in the micro-mechanical

environment of the GP and chondrocytes within the GP. This work extends our knowledge on

the contribution of the GP in OA development, which is vital if we are to reduce the burden

of this global disease.

**List of abbreviations**

AC, articular cartilage; Col10a1, type X collagen; DAB, Diaminobenzidine; DMM,

destabilisation of medial meniscus; ECM, extracellular matrix; GP, growth plate; MMP13,

matrix metalloproteinase 13; MMTL, meniscotibial ligament; OA, osteoarthritis; SCB,

subchondral bone.

**Declarations**

*Ethics approval and consent to participate*

All procedures complied with the United Kingdom Animals (Scientific Procedures) Act 1986

and were approved by The University of Edinburgh Roslin Institute's Animal Users and

Research Ethics Committees.

*Consent for publication*

Not applicable

*Availability of data and materials*

The datasets used and/or analysed during the current study are available from the

corresponding author on reasonable request.

*Competing interests*

The research leading to these results has received technical support from 3Dmagination Ltd,

Didcot, UK. KM is co-founder and director of 3Dmagination Ltd in Oxfordshire, UK, a

company which provides training and consultancy in 3D and 4D imaging.

*Funding*

We are grateful to Medical Research Council (to KAS; MR/R022240/1), Tenovus Scotland*,*

the Swedish Research Council (to AET; 2013–455) for funding. The funding sources did not

influence on the design of the study, collection, analyses, interpretation of data, writing or

submission of the manuscript.

*Authors' contributions*

All authors contributed to the study design, analysis and interpretation of the data, drafted,

critically reviewed, edited and approved the version of the manuscript for publication.

*Acknowledgements*

The authors would like to thank The Roslin Institute of The University of Edinburgh for the

assistance with the care of animal models in this study.

**References**

1.     Murray CJ., Lopez AD. Global Burden of Disease and Injur Y Series the Global

442           Burden of Disease. Oms. 1996;1–46.

443     2.     Goldring MB, Goldring SR. Osteoarthritis. J Cell Physiol. 2007;213(3):626–34.

444     3.     Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K, et al. Development of

445           criteria for the classification and reporting of osteoarthritis: Classification of

446           osteoarthritis of the knee. Arthritis Rheum. 1986;29(8):1039–49.

447     4.     Pacifici M, Koyama E, Shibukawa Y, Wu C, Tamamura Y, Enomoto-Iwamoto M, et

448           al. Cellular and molecular mechanisms of synovial joint and articular cartilage

449           formation. Ann N Y Acad Sci. 2006;1068(1):74–86.

450     5.     Staines KA, MacRae VE, Farquharson C. The importance of the SIBLING family of

451  proteins on skeletal mineralisation and bone remodelling. J Endocrinol.

452  2012;214(3):241–55.

453 6. Parfitt AM. Misconceptions (1): Epiphyseal fusion causes cessation of growth. Bone.

454  2002;30(2):337–9.

455 7. Glatt V, Canalis E, Stadmeyer L, Bouxsein ML. Age-related changes in trabecular

456  architecture differ in female and male C57BL/6J mice. J Bone Miner Res.

457  2007;22(8):1197–207.

458 8. Staines KA, Madi K, Mirczuk SM, Parker S, Burleigh A, Poulet B, et al. Endochondral

459  Growth Defect and Deployment of Transient Chondrocyte Behaviors Underlie

460  Osteoarthritis Onset in a Natural Murine Model. Arthritis Rheumatol. 2016;68(4):880–

461  91.

462 9. Madi K, Staines KA, Bay BK, Javaheri B, Geng H, Bodey AJ, et al. In situ

463  characterization of nanoscale strains in loaded whole joints via synchrotron X-ray

464  tomography. Nat Biomed Eng. 2019;4:343–354.

465 10. Glasson BVSc SS, Blanchet BS TJ, Morris DVM EA. The surgical destabilization of

466  the medial meniscus (DMM) model of osteoarthritis in the 129/SvEv mouse.

467  Osteoarthr Cart. 2007;15:1061–9.

468 11. Staines KA, Ikpegbu E, Törnqvist AE, Dillon S, Javaheri B, Amin AK, et al.

469  Conditional deletion of E11/podoplanin in bone protects against load-induced

470  osteoarthritis. BMC Musculoskelet Disord. 2019;20(1):1–11.

471 12. Sugiyama T, Price JS, Lanyon LE. Functional adaptation to mechanical loading in

472  both cortical and cancellous bone is controlled locally and is confined to the loaded

bones. Bone. 2010;46(2):314–21.

13.    Poulet B, Hamilton RW, Shefelbine S, Pitsillides AA. Characterizing a novel and

adjustable noninvasive murine joint loading model. Arthritis Rheum. 2010/10/01.

2011;63(1):137–47.

14.    Poulet B, de Souza R, Kent A V, Saxon L, Barker O, Wilson A, et al. Intermittent

applied mechanical loading induces subchondral bone thickening that may be

intensified locally by contiguous articular cartilage lesions. Osteoarthr Cartil. 2015

Jun;23(6):940–8.

15.    Staines KA, Madi K, Javaheri B, Lee PD, Pitsillides AA. A computed

microtomography method for understanding epiphyseal growth plate fusion. Front

Mater. 2018;4(January).

16.    Chambers MG, Kuffner T, Cowan SK, Cheah KSE, Mason RM. Expression of

collagen and aggrecan genes in normal and osteoarthritic murine knee joints.

Osteoarthr Cartil. 2002;10(1):51–61.

17.    Poulet B, Westerhof TAT, Hamilton RW, Shefelbine SJ, Pitsillides AA. Spontaneous

osteoarthritis in Str/ort mice is unlikely due to greater vulnerability to mechanical

trauma. Osteoarthr Cartil. 2013;21(5):756–63.

18.    Samvelyan HJ, Hughes D, Stevens C, Staines KA. Models of Osteoarthritis: Relevance

and New Insights. Calcif Tissue Int [Internet]. 2020;(0123456789). Available from:

https://doi.org/10.1007/s00223-020-00670-x

19.    Poulet B, Hamilton RW, Shefelbine S, Pitsillides AA. Characterizing a novel and

adjustable noninvasive murine joint loading model. Arthritis Rheum. 2011;63(1):137–

47.

20.    Atkins GJ, Rowe PS, Lim HP, Welldon KJ, Ormsby R, Wijenayaka AR, et al.

497             Sclerostin is a locally acting regulator of late-osteoblast/preosteocyte differentiation

498             and regulates mineralization through a MEPE-ASARM-dependent mechanism. J Bone

499             Miner Res. 2011;26(7):1425–36.

500     21.    Lai Y, Bai X, Zhao Y, Tian Q, Liu B, Lin EA, et al. ADAMTS-7 forms a positive

501             feedback loop with TNF-α in the pathogenesis of osteoarthritis. Ann Rheum Dis.

502             2014;73(8):1575–84.

503     22.    Lu Y, Ding M, Li N, Wang Q, Li J, Li X, et al. Col10a1-runx2 transgenic mice with

504             delayed chondrocyte maturation are less susceptible to developing osteoarthritis. Am J

505             Transl Res. 2014;6(6):736–45.

506     23.    Ea HK, Nguyen C, Bazin D, Bianchi A, Guicheux J, Reboul P, et al. Articular cartilage

507             calcification in osteoarthritis: Insights into crystal-induced stress. Arthritis Rheum.

508             2011;63(1):10–8.

509     24.    Ortega N, Behonick DJ, Werb Z. Matrix remodeling during endochondral ossification.

510             Trends Cell Biol. 2004;14(2):86–93.

511     25.    Stickens D, Behonick DJ, Ortega N, Heyer B, Hartenstein B, Yu Y, et al. Altered

512             endochondral bone development in matrix metalloproteinase 13-deficient mice.

513             Development. 2004;131(23):5883–95.

514     26.    Saito T, Fukai A, Mabuchi A, Ikeda T, Yano F, Ohba S, et al. Transcriptional

515             regulation of endochondral ossification by HIF-2α during skeletal growth and

516             osteoarthritis development. Nat Med. 2010;16(6):678–86.

27.   Yang S, Kim J, Ryu JH, Oh H, Chun CH, Kim BJ, et al. Hypoxia-inducible factor-2α

is a catabolic regulator of osteoarthritic cartilage destruction. Nat Med.

2010;16(6):687–93.

28.   Breur GJ, Lapierre MD, Kazmierczak K, Stechuchak KM, McCabe GP. The domain of

hypertrophic chondrocytes in growth plates growing at different rates. Calcif Tissue

Int. 1997;61(5):418–25.

29.   Huang H, Skelly JD, Ayers DC, Song J. Age-dependent Changes in the Articular

Cartilage and Subchondral Bone of C57BL/6 Mice after Surgical Destabilization of

Medial Meniscus. Sci Rep [Internet]. 2017;7(February):1–9. Available from:

http://dx.doi.org/10.1038/srep42294

30.   Staines KA, Hardy RJ, Samvelyan HJ, Ward KA, Cooper R. Life course longitudinal

growth and risk of knee osteoarthritis at age 53 years: evidence from the 1946 British

birth cohort study. 2020 Aug; Available from:

https://doi.org/10.1101%2F2020.08.18.20177485

31.   Frysz M, Gregory JS, Aspden RM, Paternoster L, Tobias JH. The effect of pubertal

timing, as reflected by height tempo, on proximal femur shape: Findings from a

population-based study in adolescents. Bone. 2020;131.

32.   Baird DA, Paternoster L, Gregory JS, Faber BG, Saunders FR, Giuraniuc C V., et al.

Investigation of the Relationship Between Susceptibility Loci for Hip Osteoarthritis

and Dual X-Ray Absorptiometry–Derived Hip Shape in a Population-Based Cohort of

Perimenopausal Women. Arthritis Rheumatol. 2018;70(12):1984–93.

33.   Baird DA, Evans DS, Kamanu FK, Gregory JS, Saunders FR, Giuraniuc C V., et al.

Identification of Novel Loci Associated With Hip Shape: A Meta-Analysis of

Genomewide Association Studies. J Bone Miner Res. 2019;34(2):241–51.

**Figure legends**

**Fig. 1. Immunohistochemical labelling in the tibial articular cartilage of mouse knee**

**joints**

Immunohistochemical labelling for type X collagen (Col10a1) and matrix metalloproteinase

(MMP13) in the articular cartilage of non-operated and DMM (**a**), or non-loaded and

mechanically loaded (**b**) knee joints of C57BL/6 male mice. Arrows indicate examples of

positive staining in the tibiae. Images are representative of results in 3 individual mice. Scale

bar = 200μm

**Fig. 2. Growth plate dynamics in the tibial growth plate of mouse knee joints**

Growth plate zone width of non-operated and DMM, or non-loaded and loaded knee joints of

C57BL/6 male mice (**a**). Ten measurements per section were obtained along the length of the

tibial growth plate in the middle region of the knee joint (n=4 mice for each experimental

group). Immunohistochemical labelling for type X collagen (Col10a1) and matrix

metalloproteinase (MMP13) in the growth plate of DMM and non-operated (**b**), or

mechanically loaded and non-loaded (**c**) knee joints of C57BL/6 male mice. Arrows indicate

examples of positive staining in the tibiae. Images are representative of results in 4 individual

mice. Scale bar = 200μm. PZ = proliferative zone; HZ = hypertrophic zone **** *p*<0.0001

**Fig. 3. Bridge location and bridge areal densities across the tibial growth plate of mice**

Location and areal densities of bridges across the growth plate projected on the medial (M)

and lateral (L) tibial joint surface in non-operated (**a**), DMM (**b**), non-loaded (**c**) and loaded

(**d**) tibiae of mice at 16 and 18 weeks of age, number of bridges in lateral and medial tibiae of

non-operated compared to DMM and non-loaded compared to loaded tibiae of mice (**e**),

Areal density (d) of bridges in medial compared to lateral tibiae, defined as the number of

bridges per 256 mm x 256 mm window of non-operated and DMM or non-loaded and loaded

knee joints (**f**). Bars represent mean ± SEM. Group sizes were $n = 8$ for non-operated and

DMM-operated mice and $n = 6$ for non-loaded and loaded mice. * $p < 0.05$ ** $p < 0.01$ ***

$p < 0.001$  **** $p < 0.0001$

**Fig. 4. MicroCT analysis of epiphyseal region of medial and lateral tibiae of mice**

MicroCT analysis of the epiphyseal region of the lateral and medial tibiae in non-operated

controls and DMM-operated knee joints subchondral bone volume fraction (SCB BV/TV) (**a**)

and epiphyseal trabecular bone volume fraction (Tb. BV/TV) (**b**). MicroCT analysis of the

epiphyseal region of the medial and lateral tibiae in non-loaded controls and loaded knee

joints subchondral bone volume fraction (SCB BV/TV) (**c**) and epiphyseal trabecular bone

volume fraction (Tb. BV/TV) (**d**). Representative colour coded images of lateral and medial

subchondral bone plate thickness of non-operated and DMM-operated tibiae of mice (**e**).

Representative colour coded images of lateral and medial subchondral bone plate thickness of

non-loaded and loaded tibiae of mice (**f**). Representative colour coded images of epiphyseal

trabecular bone thickness of non-operated and DMM-operated (**g**), or non-loaded and loaded

tibiae of mice (**h**). Bars represent mean ± SEM. Group sizes were $n = 8$ for non-operated and

DMM-operated mice and $n = 6$ for non-loaded and loaded mice.  * $p < 0.05$ ** $p < 0.01$ ***

$p < 0.001$

**Non-operated**

a

L

M

0    8.5    17
Areal density, d

**DMM**

b

M

L

**Non-loaded**

c

L

M

2    21.5    41
Areal density, d

**Loaded**

d

M

L

e

☐ Lateral
■ Medial

Number of bridges

Non-operated          DMM
                        *

Number of bridges

Non-loaded          Loaded
    *              ***

f

☐ Lateral
■ Medial

Areal density (d)

15
10
5
0

Non-operated          DMM
    ***              **

Areal density (d)

Non-loaded          Loaded
  ****              ****