"Sifarchaeota" a novel Asgard phylum capable of polysaccharide degradation and

anaerobic methylotrophy

Ibrahim F. Farag, Rui Zhao, Jennifer F. Biddle

School of Marine Science and Policy, University of Delaware, 1044 College Drive, Lewes DE

19958, USA.

Abstract

The Asgard superphylum is a deeply branching monophyletic group of Archaea, recently

described as some of the closest relatives of the eukaryotic ancestor. The wide application of

genomic analyses from metagenome sequencing has established six distinct phyla, whose

genomes encode for diverse metabolic capacities and play important biogeochemical and

ecological roles in marine sediments. Here, we describe two metagenome-assembled genomes

(MAGs) recovered from deep marine sediments off Costa Rica margin, defining a novel lineage

phylogenetically married to Thorarchaeota, as such we propose the name "Sifarchaeota" for this

phylum. The two "Sifarchaeota" MAGs encode for an anaerobic methylotrophy pathway

enabling the utilization of C1-C3 compounds (methanol and methylamines) to synthesize acetyl

CoA. Also, the MAGs showed a remarkable saccharolytic capabilities compared to other Asgard

lineages and encoded for diverse classes of carbohydrate active enzymes (CAZymes) targeting

different mono-, di- and oligosaccharides. Comparative genomic analysis based on the full

metabolic profiles of Asgard lineages revealed the close relation between "Sifarchaeota" and

Odinarchaeota MAGs, which suggested a similar metabolic potentials and ecological roles.

Furthermore, we identified multiple potential horizontal gene transfer (HGT) events from

different bacterial donors within "Sifarchaetoa" MAGs, which hypothetically expanded

"Sifarchaeota" capacities for substrate utilization, energy production and niche adaptation.

Introduction

Deep marine sediments are the home of multiple poorly described archaeal lineages, most

of which are yet uncultured (1)(2). Recently, the discovery of Asgard archaea in benthic

environments has generated great interest in novel lineages from marine sediments. Additionally,

greater attention directed towards studying Asgard archaea is also in part to understanding

eukaryogenesis, as this superphylum harbors the most closely related archaeal group to

eukaryotes and their genomes encode for multiple homologs of eukaryotic proteins (3)(4). So far,

there are 6 established Asgard phyla: Lokiarchaeota, Thorarchaeota, Odinarchaeota,

Heimdallarchaeota, Helarchaeota and Gerdarchaeota (4)(5)(6). The number of novel lineages yet

to be found is at this point unknown. This raises the need for more genome resolved metagenome

surveys to recover genomes of these archaeal lineages, decipher their metabolic capacities and

place them in the context of microbial ecology. Previous studies targeting the deep sediment

from the Costa Rica (CR) margin subseafloor have shown the presence of diverse archaeal

communities (7)(8). Among these archaeal lineages, members of Asgard superphyla were highly

abundant at multiple depths, making up 17% of the archaeal communities present (7)(8). In this

study on the Costa Rica margin subseafloor, we employ genome resolved metagenomics to

describe the metabolic potential of the genomes belonging to a new Asgard phylum.

Phylogenomic analysis placed the sequences of the new Asgard genomes as a sister clade to the

sequences of Thorarchaeota and we propose the name "Sifarchaeota" to describe this new

Asgard phylum. Comparative analysis shows distinct differences between Sifarchaeota and

previously reported Asgard archaea in terms of substrate utilization, energy production and niche

adaptation strategies. Finally, we detect multiple potential horizontal gene transfer (HGT) events

from different bacterial donors expanding the substrate utilization, energy production and

secondary metabolite production capacities of the Sifarchaeota.

Methods

Sample collection

Samples were collected during International Ocean Drilling Program (IODP) Expedition

334, Site U1379B on the Costa Rican Margin. Details on sample location and sampling

methodology have been previously described (7)(9). Microbiology samples (whole-round cores)

were collected on board and frozen immediately at -80°C. They were shipped to the Gulf Core

Repository (College Station, Texas) on dry ice and stored at -80°C until shipping to the Biddle lab

(Lewes, Delaware) on dry ice and further storage at -80°C. Metagenome sequencing data were

generated from four silty clay sediment horizons (2H-1, 2H-2, 2H-5A, and 2H-5B) in the depth

interval of 2-9 mbsf, within the sulfate reduction zone (7).

DNA extraction, library construction and sequencing

DNA for metagenomic sequencing was extracted from ~7 g sediment (~0.7 g sediment in

10 individual lysis tubes) using PowerSoil DNA Isolation Kit (Qiagen) following the

manufacturer's instructions, except for the following minor modification: the lysing tubes were

incubated in water bath of 60°C for 15 min prior to beading beating on a MP machine at speed 6

for 45 seconds. The DNA extracts were iteratively eluted from the 10 spin columns into a final of

100 μL of double distilled H_2O for further analysis. Metagenomic libraries were prepared and

sequenced (150 bp paired-end) on an Illumina NextSeq 500 sequencer at the Genome Sequencing

& Genotype Center at the University of Delaware.

Assembly and genome binning

The raw sequencing data were processed with Trimmomatic v.0.36 (10) to remove Illumina

adapters and low quality reads ("SLIDINGWINDOW:10:25"). The quality-controlled reads from

the eight samples were de novo co-assembled into contigs using Megahit v.1.1.2 (11) with the k-

mer length varying from 27 to 117. Contigs longer than 1000 bp were automatically binned using

MaxBin2 (12) and Metabat2 (13), and the best quality ones were selected using DAS_Tool (14)

with the default parameters. The resulting MAGs were quality assessed using CheckM (15) and

taxonomically classified using GTDBTk v1.3.0 (16) using the default parameters. Genome bins of

>50% completeness were manually refined using the gbtools (17) based on the GC content,

taxonomic assignments, and differential coverages in different samples. Coverages of contigs in

each sample were determined by mapping trimmed reads onto the contigs using BBMap v.37.61

(18). Taxonomy of contigs were assigned according to the taxonomy of the single-copy marker

genes in contigs identified using a script modified from blobology (19) and classified by BLASTn.

SSU rRNA sequences in contigs were identified using Barrnap (Seeman 2015, Github), and

classified using VSEARCH with the SILVA 132 release (20) as the reference.

To improve the quality of the two novel Asgard archaea MAGs, we recruited quality-

controlled reads using BBMap from 2H-2, because the highest genome coverages of these two

MAGs were detected in this particular sample. The recruited reads were then re-assembled using

SPAdes v.3.12.0 (21) using default parameters. After removal of contigs shorter than 1 kb, the

resulting scaffolds were visualized and re-binned manually using gbtools (17) as described above.

The quality of the resulting Asgard archaea genomes were checked using CheckM v.1.0.7 (15)

with the "lineage_wf" option.

Concatenated ribosomal protein phylogeny

To determine the phylogenetic affiliations of the two Sifarchaeota MAGs in the Archaea domain, we performed a thorough phylogenomic analysis based on the concatenation of 16 ribosomal proteins (L2, L3, L4, L5, L6, L14, L15, L16, L18, L22, L24, S3, S8, S10, S17, and S19). Reference genomes were selected from all the major archaeal phyla (3-5 for each) included in the GTDB database (22), except for the Asgard superphylum for which all available genomes were included. Ribosomal protein sequences were detected in Anvi'o (23) using the respective HMM profiles, aligned using MUSCLE v3.8.31(24), and concatenated. The maximum likelihood phylogenetic tree was reconstructed using IQ-Tree (v1.6.6) (25) (located on the CIPRES web server) (26) with VT + F + R10 as the best-fit substitution model selected by ModelFinder (27), and single branch location was tested using 1000 ultrafast bootstraps and approximate Bayesian computation (28). Branches with bootstrap support >80% were marked by black circles. In addition to phylogenomic analysis, we also calculated the average nucleotide identity (ANI) using FastANI (29) and average amino acid identity (AAI) using CompareM (https://github.com/dparks1134/CompareM) with default settings between these novel Asgard MAGs and other public available ones (i.e., those included in the GTDB database), to further explore the novelty of these MAGs.

Metabolic reconstruction

Amino acid sequences encoded by the Sifarchaeota MAGs were predicted using Prodigal v2.6.3 (30) applying the default parameters and using translation table 11. The resulting amino acid sequences were screened using HMMsearch tool (31) against custom HMM databases (32) representing the key genes for specific metabolic pathways to understand the potential metabolic capacities and their ecological roles of Sifarchaeota. The presence/absence profiles of the

metabolic pathways and their completion levels were further assessed through querying the

predicted amino acids against KEGG database using BlastKoala tool (33). Carbohydrate-active

enzymes encoded by the Sifarchaeota MAGs were analyzed using dbCAN-fam-HMMs (v6)

database (34). Proteases, peptidases, and peptidase inhibitors encoded by the MAGS were detected

via USEARCH-ublast tool (35) against the MEROPS database v12.1(36). Finally, the predicted

amino acid sequences were queried against the TCDB database (37) using USEARCH-ublast tool

(35) to identify the potential transporters.

Genome centric comparative analysis for the Asgard superphylum

We compared the total metabolic profiles of Sifarchaetoa MAGs with representatives from

other Asgard lineages including (Odinarchaetoa, Thorarchaeota, Lokiarchaeota and

Heimdallarcaheota) to identify the key metabolic differences between each of the phyla within the

Asgard superphylum. We queried the representative MAGs of each of the Asgard lineages against

KOfam database (KEGG release 94.1) via KofamKOALA webtool (38) and using evalue 10^{-3}.

Then, the Asgard MAGs were clustered based on the presence/absence profiles of the identified

KOfams, shared between at least 3 genomes, using clustergrammer web-based tool (39) and

applying the Euclidean distance and average linkage type.

Horizontal gene transfer (HGT) analysis

The HGT events were detected through querying Sifarchaeota predicted amino acids

against KEGG database (KEGG release 94.1) via GhostKoala web interface (33). Proteins with

KO annotation, non-Asgard hits, and a bit score >100 were considered as potential HGT candidate

proteins. Then, the candidate set of proteins were queried against (nr) and UniProtKB/swissprot

databases and all proteins showing Asgard hits as one of the top hits were removed from any

further analysis. Finally, each candidate protein is aligned to reference set of proteins collected via

Annotree (40) using the corresponding KO entry and the HGT events were confirmed through

creating approximately-maximum-likelihood phylogenetic tree for each candidate protein using

FastTree v2.1 (41). An outline for the HGT detection pipeline is illustrated in Supplementary

Figure 1.

Results

MAG construction and phylogenomic analysis

We reconstructed 3 MAGs belonging to a potentially novel Asgard phylum with moderate

completion levels (67-80%) and very low contamination levels (0.93-1.9%) (Table 1). The

taxonomic affiliations of these potentially novel Asgard MAGs were tested using 16 ribosomal

proteins phylogenomic tree and it showed that both MAGs clustered together and formed a sister

lineage to MAGs belonging to the phylum Thorarchaeota (Figure 1). The phylogenetic analysis

using a set of 122 archaea specific marker proteins implemented in GTDB-tk confirmed that these

MAGs are belonging to a potentially novel Asgard archaeal lineage. To confirm the unique

positions of these candidate novel Asgard lineage, we calculated the average nucleotide identities

(AAI) and average amino acid identities (ANI) between the 3 novel Asgard MAGs and other

MAGs representing the other Asgard lineages including (Lokiarchaeota, Thorarchaeota,

Heimdallarchaeota and Odinarchaeota). On average, the 3 novel Asgard MAGs showed low AAI

values when compared to the other Asgard MAGs (<50%) (Supplementary Tables 1 and 2).

Two of the new Asgard MAGs, bins 190 and 142, showed a very high similarity with AAI

98.81% and ANI 99.52%, as such, we focused all our subsequent analysis on two MAGs, bins 042

and 142, since they were the most complete and unique MAGs in this study.

Description of the taxa

Although the two Sifarchaeota MAGs, bin042 and bin142, shared many metabolic

similarities, we could not describe them using one type strain due to the significant phylogenetic

distances between the two MAGs (Figure 1) and the genomic differences as described using ANI

and AAI values (Supplementary Tables 1 and 2). Therefore, we used two type strains to describe

the two putative Sifarchaeota lineages.

Candidatus Sifarchaeotum costaricensis **(*costaricensis* of or from Costa Rica).** Type material is

the genome designated as bin042 representing '*Candidatus* Sifarchaeotum costaricensis'.

Candidatus Sifarchaeotum subterraneus **(*subterraneus* latin name of subsurface).** Type material

is the genome designated as bin142 representing '*Candidatus* Sifarchaeotum subterraneus'.

Based on these genera, we further propose the name of a new Asgard phylum, the phylum

'*Candidatus* Sifarchaeota phylum nov.'

Metabolic reconstruction of Sifarchaeota MAGs showed remarkable saccharolytic capacities

and potential anaerobic methylotrophy

Previous reports showed that Asgard genomes encode for wide range of protein and fatty

acids degradation capacities (42). So far, most of the known Asgard genomes showed limited

saccharolytic capacities emphasized by their low genomic densities of CAZymes and sugar

transporters. However, Sifarchaeota showed high abundance and diversity of Carbohydrate Active

Enzymes (CAZymes) encoded by their MAGs specifically targeting sugars varying in

complexities from low (C1-C3) to moderate (C4-C6), including mono-, di- and oligo- saccharides

(Supplementary Figure 1, Supplementary Tables 3 and 5). Interestingly, CAZyme analysis

revealed the presence of different glycoside hydrolases (GH) families including cellulases and

endoglucanases (GH5 and GH9), cyclomaltodextrinase (GH13), α-glucosidase (GH63), β-1,4-

mannooligosaccharide phosphorylase (GH130) and β-L-arabinofuranosidase (GH142) targeting

cellobiose, maltose and maltooligosaccharides, alpha-glucosaccharides, glucose/mannose and

arabinosaccharides, respectively. We also identified dedicated sugar transporters mediating the

transfer of these sugars inside the Sifarchaeota cells, where the degradation and fermentation

processes take place (Supplementary Table 6). Metabolic reconstruction of Sifarchaeota MAGs

predict a general fermentative anaerobic life style with multiple anaerobic respiration capabilities

emphasized by the presence of incomplete reverse TCA cycle, presence of various fermentation

pathways capable of producing different fermentative products including acetate, acetoin and

butanediol under strictly anaerobic conditions as well as the potential capabilities to anaerobically

respire sulfate to sulfite and nitrite to ammonia. Notably, the degradation capacities for proteins,

peptides and amino acids as well as amino acid metabolism are limited in Sifarchaeota compared

to the other benthic archaea. Hence, Sifarchaeota may make up the short supply of fixed nitrogen

via reducing nitrite to ammonia and encoding for amidases to extract fixed nitrogen from nitrogen

containing amides like formamide.

Sifarchaeota MAGs showed the capacity to utilize various C1 compounds including formate

and methanol. For example, methanol is metabolized using an anaerobic methylotrophy pathway

previously described (8), suggesting the potential widespread distribution of this pathway among

benthic marine archaea to metabolize methylated compounds. This potential anaerobic

methylotrophic capacity was inferred via detecting incomplete methylotrophic methanogenesis

pathway, where the key genes encoding for the methyl coenzyme reductase complex were

completely absent. Besides, genes for incomplete Wood Ljundahl (WL) pathway were identified,

where only the genes of the carbonyl branch were present and the genes of the methyl branch were

completely absent. Detecting these set of genes in Sifarchaeota MAGs suggests the presence of

anaerobic methylotrophic capability, enabling Sifarchaeota to recycle methyl groups within

methanol and other methylated compounds. Then, these methyl groups are transferred to

tetrahydrofolate complex replacing the function of the methyl branch of WL pathway and

ultimately producing acetyl CoA.

Comparative genomic analyses between different Asgard lineages shows diverse metabolic

features and life style patterns

To understand the key metabolic differences between the Asgard lineages, we conducted a

comprehensive genome-centric analyses. As described in the methods section, we parsed 13

different MAGs (2 MAGs obtained from this study and 11 publicly available MAGs) belonging

to 5 different Asgard phyla (Lokiarchaeota, Thorarchaeota, Heimdallarchaeota, Odinarchaeota and

Sifarchaeota) against KOfam database using an HMM search tool (Supplementary Table 7. The

analysis output focused on two aspects (1) exploring the range of life style diversity within the

Asgard superphylum and (2) comparing the metabolic capabilities of different Asgard lineages.

Overall, the comparative analyses grouped the Asgard superphylum into 4 distinct clusters

based on their whole metabolic profiles. Notably, all the MAGs within the Asgard superphylum

suggested a similar fermentative anaerobic life styles, where the core genes for reverse TCA cycle,

various fermentation pathways were present as well as the absence of oxidative phosphorylation

and oxygen tolerance related genes. Here we describe the clusters and the features that drive their

clustering.

The **first cluster** grouped the 2 Sifarchaeota MAGs (bin42 and 142), obtained from this

study, with Candidatus Odinarchaetoa archaeon LCB4. Sifarchaeota MAGs shared multiple

metabolic similarities with the Odinarchaeota MAGs including limited amino acid and fatty acids

metabolic potentials, evident saccharolytic activities, emphasized by high density of CAZyme

encoding genes targeting different mono-, di- and oligosaccharide. Also, we could only identify

genes encoding for the oxidative branch of the pentose phosphate pathway, producing

Phosphoribosyl pyrophosphate (PRPP), which eventually channeled to the purine and pyrimidine

metabolic pathways to be used for nucleotide and nucleic acid biosynthesis. Also, MAGs within

this cluster encoded for large number of genes involved in C1 metabolism including formate,

methylamines and methanol. Genes encoding for formate dehydrogenase, methanol and

methylamine specific corrinoid protein:coenzyme M methyltransferases were present, which

suggest the potential capability of both lineages to utilize formate, methylamines and methanol as

carbon sources, respectively.

The **second cluster** grouped Lokiarchaeota and Thorarchaeota MAGs together. Unlike the

previous cluster, MAGs belonging to that cluster are characterized by their proteins and peptide

degrading capabilities emphasized by the presence of high number of proteases and peptidases

encoding genes ranging in density from (126-193 proteins/Mbs) and belonging to different

families of serine and metalloproteases. Unlike other Asgard groups, Thorarchaeota and

Lokiarchaeota showed the capacities to synthesize different amino acids including nonpolar amino

acids (e.g. isoleucine, leucine, valine and alanine), aromatic amino acids (e.g. tryptophan and

phenylalanine) via the shikimate pathway and charged amino acids (e.g. glutamate and lysine).

Similar to other Asgard archaea, MAGs belonging to Thor- and Loki- archaeota encoded for genes

mediating the metabolism of C1 compounds (e.g. formate), however, the absence of

methyltransferases encoding genes excluded the use methylated compounds as one of the potential

substrates.

Finally, **the third and fourth clusters** included different Heimdallarchaeota MAGs. The

separation between Heimdallarchaeota based on their metabolic profiles suggests the presence of

fundamental metabolic differences between the members of Heimdallarchaeota phylum and

supports the previous findings that described Heimdallarcaheota as a polyphyletic group (3)(6).

This raises the need for wider sampling efforts targeting Heimdallarchaeota genomes to fully

resolve their phylogenetic position and evolutionary history. However, in this study we grouped

all the Heimdallarchaeota MAGs and treated them as one phylum and designed a model that

highlights the difference between them and the other Asgard lineages. Notably, all

Heimdallarcahetoa MAGs showed the capacity to utilize proteins and short chain fatty acids as

carbon sources, while polysaccharide degradation was less supported. Similar to Loki- and

Throarchaeota MAGs, Heimdallarcahetoa MAGs encoded for high number of peptidases with

coding densities (110-210 proteins/Mbs) and belonging to diverse families of proteases and

peptidases including serine, metallo, cysteine and threonine peptidases (Supplementary Table 3).

Also, Heimdallarcahetoa MAGs showed the capacity to metabolize and synthesize nonpolar amino

acids (e.g. alanine, glycine, and threonine). Among all the Asgard MAGs included in this analysis,

only Heimdallarchaeota MAGs encoded for enzymes mediating beta-oxidation pathway including

acyl CoA dehydrogenase, enoyl CoA hydratase and hydroxyacyl CoA dehydrogenase, suggesting

their potential to use short-chain fatty acids (SCFA) as carbon and energy sources. Similar to all

other Asgards, Heimdallarchaeota encoded for incomplete WL pathway, which could have a role

in metabolizing C1 compounds like formate and formaldehyde.

Due to the limited access to the surrounding microbial community composition and

environmental conditions as well as the underrepresentation of the some of the lineages (only 1

MAG from Odinarchaeota), we could not assess the exact reasons for these diverse substrate

preferences between the Asgard phyla (e.g. polysaccharides in Sifarchaeota, proteins in Thor- and

Loki-archaeota and short-chain fatty acids in Heimdallarchaeota). Also, we are unable to conclude

at this point if these findings could be generalized for all the members within each phylum or it is

just limited to the lineages/MAGs included in the study.

Role of horizontal gene transfers (HGT) in enhancing Sifarchaeota metabolic capacities and

niche adaptations

We investigated the role of HGT in expanding the metabolic capacities, substrate

utilization and niche adaptation of the Sifarchaeota phylum. We traced the origin of each HGT

event and identified the extent of the spread of each gene within the Asgard superphylum

(Supplementary Figure 2). We successfully identified a total of 65 HGT events in Sifarchaeota, 12

(0.65% of the total proteins) and 53 (1.34% of the total proteins) events in bin042 and bin142,

respectively. Most likely, the majority of the identified HGT events were lineage specific (58,

89.2%) and only few similar events were detected in other Asgard lineages (7, 11.8%). Also, we

identified the potential donors for most of the horizontally transferred genes, ~90% of which were

of bacterial origins. The major bacterial contributors for the horizontally transferred genes were

Firmicutes (~30%), Chloroflexi (~15%), Proteobacteria (~13%), Cyanobacteria (5%) and other

bacterial lineages (~30%). Only a small fraction (~10%) of the horizontally transferred genes were

of archaeal origin outside of the Asgard superphylum (Figure 5 and Supplementary Table 8).

We classified HGT events based on the how widespread the transferred genes were among

the different Asgard phylum, as well as other archaeal lineages. Accordingly, the HGT events were

classified into lineage-specific, phylum-specific and domain-wide events. In the lineage specific

event, the closest relatives of the transferred genes were bacteria and the genes were exclusively

found in Sifarchaetoa MAGs and no orthologs were found in other Asgard or any other archaeal

phyla (e.g. butanediol dehydrogenase). In the phylum-specific event, the closest relatives of the

transferred genes were bacteria and the genes were found in multiple Asgard phyla (e.g. enoyl

CoA hydratase). In the domain wide event, the closest relatives of the transferred genes were

bacteria and the genes were found in different archaeal phyla (e.g. arsenate reductase *arsC*

thioredoxin) (Figure 5). In general, the functional annotation of the transferred genes showed that

these genes are involved in augmenting the Sifarchaeota metabolic repertoire. The majority of the

transferred genes fall within two metabolic modules: butanoate metabolism and biosynthesis of

secondary metabolites. In the butanoate metabolism, the majority of transferred genes were

encoding for butane diol dehydrogenases and glutaconate CoA-transferase subunits mediating the

key steps in pyruvate and acetate formation from butane diol and hydroxy glutaryl CoA,

respectively.

While the majority of the genes involved in the biosynthesis of secondary metabolites

were part of the porphyrin and chlorophyll metabolism and terpenoid biosynthesis (e.g.

anaerobic magnesium-protoporphyrin IX monomethyl ester cyclase and 1,4-dihydroxy-2-

naphthoate polyprenyltransferase). Interestingly, multiple genes involved in niche adaptation

may be acquired from the surrounding bacterial communities. These genes including

formamidase and MtaA/CmuA family methyltransferase which potentially enable Sifarchaeota

to utilize amide containing compounds and methylated compounds as nitrogen and carbon

sources, respectively. Also, a gene encodes for arsenate reductase was potentially acquired from

candidate division KSB1 bacterium, potentially allowing Sifarchaeota to use arsenate containing

compounds as final electron acceptors.

Discussion

In this study, we recovered three MAGs from deep Costa Rica sediments belonging to a new

Asgard phylum, forming a sister clade to MAGs belonging to Thorarchaeota. We propose the name

Sifarchaeota for this phylum, named after the Norse goddess, Sif, wife of Thor. Putative collective

metabolic profiles of the Sifarchaeota MAGs showed remarkable differences in the life style and

niche adaptations compared to the other Asgard members. We predict a saccharolytic,

fermentation-based life style with limited amino acid and fatty acid metabolism, whereas most of

the Asgard archaea identified before this study were known for their peptide degradation and short

chain fatty acid oxidation capacities (3)(5)(42). We detected genes encoding for incomplete

methanogenesis pathways coupled with the carbonyl branch of WL pathway suggesting the

capability of Sifarchaeota to perform anaerobic methylotrophy enabling the utilization of various

methylated compounds (e.g. methanol and methylamines). The widespread of anaerobic

methylotrophy in multiple benthic archaea highlights the importance of this pathway as an

effective strategy to utilize various methylated compounds commonly encountered in the marine

sediment niches (8)(43). On the other hand, Sifarchaeota MAGs shared similar potential

biogeochemical functions with other Asgard archaea including the presence of nitrite reductase

(*nirBD*) genes, putatively enabling Sifarchaeota members to reduce nitrite to ammonia as well as

genes encoding for sulfate adenylyltransferase (*sat*) and phosphoadenosine phosphosulfate

reductase (*cysH*), signifying their putative capability to perform assimilatory sulfate reduction and

sulfate activation. These shared characteristics between Asgard genomes confirm the significant

roles of different Asgard lineages in nitrogen and sulfur biogeochemical cycles in marine sediment

environments.

We also gauged the role of HGT in shaping the evolution of Sifarchaeaota. Our analysis

suggests that HGT events either added novel genes to the Sifarchaeota pangenome that impart new

functions e.g. butanoate metabolism and biosynthesis of secondary metabolites or enabled the

utilization of alternative non-organic compounds as electron source e.g. arsenate reductase.

Moreover, we explored the range of horizontally transferred gene donors and we concluded that

HGT is not limited to a specific phylogenetic group and probably acquired from the surrounding

bacterial communities, normally present in deep marine sediments. Finally, 91% of HGT events

are Sifarchaeota lineage specific and probably took place relatively recently during the course of

evolution, after the diversification of Sifarchaeota from the other Asgard lineages. We identified

only 9% of the events happened earlier to the full diversification of Sifarchaeota from other

Asgards, as well as other archaeal lineages. This could be a plausible explanation for the presence

of shared functions of non-archaeal origin between most of Asgard lineages.

Acknowledgements
354
355 　We would like to thank the shipboard scientists and crew of IODP Expedition 334 for

356 collecting these sediments, and the shorebased curators of the Gulf Coast Repository for their

357 faithful stewardship of precious frozen subsurface samples. We also want to thank our high-

358 performance computation cluster administrator, Karol Miaskiewicz, for his tireless work. This

359 work was supported by the WM Keck Foundation award to JFB.

Figure 1. Maximum-likelihood phylogenetic tree of archaea genomes based on concatenated 16 ribosomal proteins. This tree was inferred using IQ-TREE v1.6.10 with the LG+R7 model and 1000 ultrafast bootstraps. The Sifarchaeota MAGs recovered in this study is highlighted in red. Lineages of the Asgard superphylum are expanded, while the other linages were collapsed, if possible. The scale bar shows estimated sequence substitutions per residue.

Figure 2. Metabolic reconstruction of the key metabolic pathways encoded by the Sifarchaeota MAGs. Central metabolic pathways are shown in gray boxes, carbon fixation pathways (WL and rTCA cycles) are shown in pink, electron transport chain (ETC) proteins are shown in yellow, fermentation products are shown in blue boxes, enzymes and enzyme complexes are shown in orange circles, energy carriers are shown in red, and metabolite and amino acid transporters are shown in light green.

Figure 3. Comparative analysis between MAGs representing different Asgard members (A) Heatmap clustering the different Asgard MAGs (X-axis) based on their total metabolic profiles as predicted by Kofam database (Y-axis). The clustering was performed using Euclidean distance and complete linkage methods. (B) Metabolic model illustrates the key metabolic features identified in Sifarchaeota cluster. (C) Metabolic model illustrates the key metabolic features identified in Lokiarchaeota and Thorarchaeota cluster. (D) Metabolic model illustrates the key metabolic features identified in Heimdallarchaeota clusters.

Figure 4. Summary of HGT events detected in the Sifarchaeota MAGs. (A) Different functional modules of horizontally transferred genes. (B) Major donors of the horizontally transferred genes.

(A) Lineage specific

(B) Phylum specific

(C) Domain wide event

Figure 5. Examples for horizontally transferred genes in Sifarchaeota MAGs. (A) Lineage specific HGT event. Maximum likelihood phylogenetic tree of butanediol dehydrogenase gene sequences. Shaded areas correspond to the potential source organims. The tree was constructed on the basis of butanediol dehydrogenase gene sequences using FastTree. Reference sequences were obtained using AnnoTree (K00004). (B) Phylum specific HGT event. Maximum likelihood phylogenetic tree of enoyl CoA hydratase gene sequences. The tree was constructed on the basis of enoyl CoA hydratase gene sequences using FastTree. Reference sequences were obtained using AnnoTree (K01692). (C) Domain wide HGT event. Maximum likelihood phylogenetic tree of arsenate reductase ArsC thioredoxin gene sequences. The tree was constructed on the basis of arsenate reductase ArsC thioredoxin gene sequences using FastTree. Reference sequences were obtained using AnnoTree (K03741).

Table 1. Details of the Sifarchaeota MAGs analyzed in this study.

Bin ID	Completion	Contamination	Genome Size (Mbps)	Estimated Genome Size (Mbps)	Strain heterogeneity	# contigs
Bin.042_1	67.76	0.93	1.8	2.39	0	301
Bin_190	74.4	1.9	1.9	2.38	0	786
Bin_142	81.4	1.9	2.01	2.39	0	300

References

1. Teske A, Sørensen KB. 2008. Uncultured archaea in deep marine subsurface sediments: have we caught them all? ISME J 2:3–18.

2. Solden L, Lloyd K, Wrighton K. 2016. The bright side of microbial dark matter: lessons learned from the uncultivated majority. Curr Opin Microbiol 31:217–226.

3. Spang A, Saw JH, Jørgensen SL, Zaremba-Niedzwiedzka K, Martijn J, Lind AE, van Eijk R, Schleper C, Guy L, Ettema TJG. 2015. Complex archaea that bridge the gap between prokaryotes and eukaryotes. Nature 521:173–179.

4. Zaremba-Niedzwiedzka K, Caceres EF, Saw JH, Bäckström D, Juzokaite L, Vancaester E, Seitz KW, Anantharaman K, Starnawski P, Kjeldsen KU, Stott MB, Nunoura T, Banfield JF, Schramm A, Baker BJ, Spang A, Ettema TJG. 2017. Asgard archaea illuminate the origin of eukaryotic cellular complexity. Nature 541:353–358.

5. Seitz KW, Dombrowski N, Eme L, Spang A, Lombard J, Sieber JR, Teske AP, Ettema TJG, Baker BJ. 2019. Asgard archaea capable of anaerobic hydrocarbon cycling. Nat Commun 10:1822.

6. Cai M, Liu Y, Yin X, Zhou Z, Friedrich MW, Richter-Heitmann T, Nimzyk R, Kulkarni A, Wang X, Li W, Pan J, Yang Y, Gu J-D, Li M. 2020. Diverse Asgard archaea including the novel phylum Gerdarchaeota participate in organic matter degradation. Sci China Life Sci 63:886–897.

7. Amanda Martino, Matthew E. Rhodes, Rosa León-Zayas, Isabella E. Valente, Jennifer F. Biddle and Christopher H. House. 2019. Microbial Diversity in Sub-Seafloor Sediments from the Costa Rica Margin. Geosciences 9:218.

8. Farag IF, Biddle JF, Zhao R, Martino AJ, House CH, León-Zayas RI. 2020. Metabolic potentials of archaeal lineages resolved from metagenomes of deep Costa Rica sediments. ISME J 14:1345–1358.

9. Vannucchi, P.; Ujiie, K.; Stroncik, N.; IODP Exp. 334 Scientific Party; Yatheesh, V. 2013. IODP expedition 334: An investigation of the sedimentary record, fluid flow and state of stress on top of the seismogenic zone of an erosive subduction margin. Scientific Drilling vol.15:23–30.

10. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114–2120.

11. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. 2015. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics 31:1674–1676.

12. Wu Y-W, Tang Y-H, Tringe SG, Simmons BA, Singer SW. 2014. MaxBin: an automated binning method to recover individual genomes from metagenomes using an expectation-maximization algorithm. Microbiome 2:26.

13. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. PeerJ 7:e7359.

14. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, Banfield JF. 2018. Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. Nat Microbiol 3:836–843.

15. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. Genome Res 25:1043–1055.

16. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2019. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. Bioinformatics https://doi.org/10.1093/bioinformatics/btz848.

17. Seah BKB, Gruber-Vodicka HR. 2015. gbtools: Interactive Visualization of Metagenome Bins in R. Frontiers in Microbiology 6:1451.

18. Bushnell B. 2014. BBMap: A Fast, Accurate, Splice-Aware Aligner.

19. Kumar S, Jones M, Koutsovoulos G, Clarke M, Blaxter M. 2013. Blobology: exploring raw genome data for contaminants, symbionts and parasites using taxon-annotated GC-coverage plots. Frontiers in Genetics 4:237.

20. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res 41:D590-596.

21. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,

Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477.

22. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz P. 2018. A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. Nature Biotechnology 36:996–1004.

23. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, Delmont TO. 2015. Anvi'o: an advanced analysis and visualization platform for 'omics data. PeerJ 3:e1319.

24. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32:1792–1797.

25. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol 32:268–274.

26. Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. 2010 Gateway Computing Environments Workshop (GCE) 1–8.

27. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods 14:587–589.

28. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the Ultrafast Bootstrap Approximation. Mol Biol Evol 35:518–522.

29. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun 9:5114.

30. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics 11:119.

31. Johnson LS, Eddy SR, Portugaly E. 2010. Hidden Markov model speed heuristic and iterative HMM search procedure. BMC Bioinformatics 11:431.

32. Beckmann S, Farag IF, Zhao R, Christman GD, Prouty NG, Biddle JF. 2020. Evidence for an expanded repertoire of electron acceptors for the anaerobic oxidation of methane in authigenic carbonates in the Atlantic and Pacific Ocean. preprint, Microbiology.

33. Kanehisa M, Sato Y, Morishima K. 2016. BlastKOALA and GhostKOALA: KEGG Tools for Functional Characterization of Genome and Metagenome Sequences. J Mol Biol 428:726–731.

34. Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. 2012. dbCAN: a web resource for automated carbohydrate-active enzyme annotation. Nucleic Acids Res 40:W445-451.

35. Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics 26:2460–2461.

36. Rawlings ND, Barrett AJ, Thomas PD, Huang X, Bateman A, Finn RD. 2018. The MEROPS database of proteolytic enzymes, their substrates and inhibitors in 2017 and a comparison with peptidases in the PANTHER database. Nucleic Acids Res 46:D624–D632.

37. Saier MH, Reddy VS, Tsu BV, Ahmed MS, Li C, Moreno-Hagelsieb G. 2016. The Transporter Classification Database (TCDB): recent advances. Nucleic Acids Res 44:D372-379.

38. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata H. 2020. KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Bioinformatics 36:2251–2252.

39. Fernandez NF, Gundersen GW, Rahman A, Grimes ML, Rikova K, Hornbeck P, Ma'ayan A. 2017. Clustergrammer, a web-based heatmap visualization and analysis tool for high-dimensional biological data. Sci Data 4:170151.

40. Mendler K, Chen H, Parks DH, Lobb B, Hug LA, Doxey AC. 2019. AnnoTree: visualization and exploration of a functionally annotated microbial tree of life. Nucleic Acids Res 47:4442–4448.

41. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood trees for large alignments. PLoS ONE 5:e9490.

42. MacLeod F, Kindler GS, Wong HL, Chen R, Burns BP. 2019. Asgard archaea: Diversity, function, and evolutionary implications in a range of microbiomes. AIMS microbiology 5:48–61.

43. Anda VD, Chen L-X, Dombrowski N, Hua Z, Jiang H-C, Banfield J, Li W-J, Baker B. 2020.

Brockarchaeota, a novel archaeal lineage capable of methylotrophy. preprint, In Review.

Supplementary Figure 1. Relative densities (numbers per 1 Mb) of peptidases and CAZymes encoded by the different Asgard MAGs.

Supplementary Figure 2. Workflow diagram describes the steps followed to identify HGT

events in Sifarchaeota MAGs.