1   **The Oncogenic Function of PLAGL2 is Mediated via Specific Target Genes Through a**

2   **Wnt-independent Mechanism in Colorectal Cancer**

4   Anthony D. Fischer[2], Daniel A. Veronese Paniagua[3], Shriya Swaminathan[3], Hajime Kashima[1,]

5   Deborah L. Rubin[1], Blair B. Madison[1]

6   1. Department of Medicine, Division of Gastroenterology, Washington University School of

7   Medicine, Saint Louis, MO 63110.

8   2. Department of Genetics, Washington University School of Medicine, Saint Louis, MO

9   63110.

3. Washington University School of Medicine, Saint Louis, MO 63110.

Corresponding Author:

Blair B. Madison, Ph.D.

Assistant Professor

Washington University School of Medicine

Department of Medicine

Division of Gastroenterology

660 S. Euclid Ave.

Campus Box 8124

CSRB NT 923

St Louis MO 63110

Ph: 267-496-6670

Email: bbmadison@gmail.com

Running Title: PLAGL2 Drives Colorectal Epithelial Transformation Via Wnt-Independent

Pathways

Keywords: Colorectal cancer, PLAGL2, IGF2, ASCL2, Wnt, intestinal epithelium

ABSTRACT

Colorectal cancer (CRC) tumorigenesis and progression are linked to common

oncogenic mutations, especially in the tumor suppressor *APC*, whose loss triggers the

deregulation of TCF4/β-Catenin activity. CRC tumorigenesis is also driven by multiple epi-

mutational modifiers, such as transcriptional regulators. We describe the common (and near-

universal) activation of the zinc finger transcription factor and Let-7 target PLAGL2 in CRC and

find that it is a key driver of intestinal epithelial transformation. PLAGL2 drives proliferation, cell

cycle progression, and anchorage-independent growth in CRC cell lines and non-transformed

intestinal cells. Investigating effects of PLAGL2 on downstream pathways revealed very modest

effects on canonical Wnt signaling. Alternatively, we find pronounced effects on the direct

PLAGL2 target genes *IGF2,* a fetal growth factor, and *ASCL2*, an intestinal stem cell-specific

bHLH transcription factor. Inactivation of PLAGL2 in CRC cell lines has pronounced effects on

ASCL2 reporter activity. Furthermore, ASCL2 expression can partially rescue deficits of

proliferation and cell cycle progression caused by depletion of PLAGL2 in CRC cell lines. Thus,

the oncogenic effects of PLAGL2 appear to be mediated via core stem cell and onco-fetal

pathways, with minimal effects on downstream Wnt signaling.

INTRODUCTION

Colorectal cancer (CRC) is the third most common of all human malignancies and is the

second leading cause of cancer-related deaths, after lung/bronchus cancer

(seer.cancer.gov/statfacts). For over 20 years, the role of canonical Wnt signaling has been

front and center for this malignancy, given the key role of the gatekeeper tumor suppressor,

APC, which encodes a key scaffold protein in the β-Catenin destruction complex [1, 2].

Mutations in other Wnt pathway components, such as *AXIN2*, *CTNNB1*, and *RSPO* genes [3]

can also lead to hyperactive Wnt signaling in CRC. Mutations in *TP53*, *SMAD4*, and *KRAS* are

also very common in CRC, underscoring the roles of pathways regulating genome integrity,

TGFβ/SMAD, and MAPK signaling. However, additional pathways are relevant in CRC

pathogenesis, especially those pathways that drive an immature, fetal, or stem cell expression

signature; such signatures predict an aggressive phenotype and poor prognosis [4, 5].

The Let-7 family of microRNAs (miRNAs) are well known for their key role in repressing

naïve cellular states, controlling proliferation, and for maintaining cellular differentiation [6-9].

Consistent with this role Let-7 depletion promotes stem cell fate in intestinal epithelial cells [10]

while down-regulation of Let-7 miRNA levels (or compromised Let-7 activity) fuels CRC

carcinogenesis [10-12], with similar pro-oncogenic effects in many other malignancies [13-17].

Targets repressed by the Let-7 miRNA family are often part of proto-typical onco-fetal pathways

that are frequently re-activated in a multitude of malignancies [8], including CRC [18, 19]. Such

targets include *IGF2BP1*, *IGF2BP2*, *HMGA2*, *MYCN*, and a target we have recently

characterized, *PLAGL2* [20]. We have previously documented the integral role of the Let-7

target HMGA2 in driving tumorigenesis in mouse models of intestine-specific Let-7 depletion

[10], although HMGA2 alone does not appear to drive stem cell fate in intestinal epithelial cells

[10]. In contrast, we discovered that PLAGL2 clearly drives stem cell fate in intestinal organoids,

and directly activates the key stem cell-specific transcription factor, ASCL2 [20]. ASCL2 is

critical for establishing and maintaining intestinal stem cell fate [21, 22].

Previous studies have demonstrated some oncogenic roles for PLAGL2 in CRC cell

lines, with some documented effects on features of cellular transformation, *in vitro* [23-27].

Although these effects are hypothesized to be mediated via PLAGL2 enhancement of canonical

Wnt signaling [23-27], the effects on Wnt signaling are often modest and the specific roles for

such effects have not been determined. Studies in human CRC and glioma cells have revealed

that PLAGL2 directly activates expression of *WNT6* [27, 28]. Despite this, effects on

downstream Wnt signaling were not documented following manipulation of either PLAGL2 or

WNT6 in CRC cell lines [27], whereas a clear effect was seen in neural stem cells [28]. This

may reflect the commonplace ligand-independent hyperactivation of canonical Wnt signaling

that occurs in CRC, e.g. from mutations in *APC*, which likely obscures (or renders irrelevant)

any effects of individual upstream Wnt ligands. Further insight into the relationship between

PLAGL2 and Wnt signaling was gained from experiments where we manipulated PLAGL2 levels

in intestinal organoids [20]. PLAGL2 over-expression conferred Wnt-independent growth, but

did not consistently augment Wnt signaling [20]. These results contrast the PLAGL2 studies

described above, which hypothesize an obligatory downstream role for Wnt signaling in the

context of cellular transformation. In non-transformed cells (organoids) we did find that factors

(perhaps Wnt ligands) secreted from PLAGL2-overexpressing organoids enhanced a Wnt

reporter in co-cultured organoids [20]. Consistent with this, PLAGL2 over-expression robustly

activated expression of several Wnt ligands in organoids [20].

To investigate oncogenic mechanisms downstream of PLAGL2 we examined the

potential of PLAGL2 to drive and maintain features of cellular transformation, and how such

effects were mediated by direct PLAGL2 target genes. We find that PLAGL2 is a commonly up-

regulated factor in CRC and has significant transforming properties. PLAGL2 effects on ASCL2-

mediated signaling appear much more robust than effects on Wnt signaling. Through a close

examination of TCF4/β-Catenin target genes and a TOP-tdT Wnt reporter, we find that PLAGL2

has minimal effects on canonical Wnt signaling. Consistent with these findings, over-expression

of the PLAGL2 target genes *ASCL2* and *IGF2* (but not constitutive activation of Wnt signaling)

can rescue proliferation defects caused by PLAGL2 inactivation.

RESULTS

We first examined expression of *PLAGL2* in matched colonic adenocarcinomas to

compare expression between tumors and non-malignant tissue from the same individual.

PLAGL2 expression was usually undetectable in non-tumor tissue (Fig. 1A), whereas in tumor

tissue *PLAGL2* was universally up-regulated (Fig. 1A). Consistent with this, analysis of TCGA

RNA-seq data for colorectal cancer revealed the common and robust up-regulation of *PLAGL2*

in tumor tissue (Fig. 1B). Expression analysis in CRC cell lines revealed *PLAGL2* expression in

all CRC cell lines tested, with robust expression in HEK293T cells (Fig. 1C). Further

examination of TCGA data revealed that *PLAGL2* over-expression is a frequent feature of

chromosomal instability (CIN) microsatellite-stable (MSS) tumors (P = 3.7e-17), a class that

makes up the majority (~80%) of CRCs. This is consistent with data from TCGA [29-31], where

we find that the only mutations significantly associated with *PLAGL2* over-expression are *TP53*

(P < 0.0001) and *APC* (P <0.01) (Fig. 1D), which are hallmarks of CIN tumors [29]. This close

association with *APC* loss could underlie observations that PLAGL2 expression correlates with

β-Catenin levels in tumors [25]. As expected, *PLAGL2* up-regulation is rarely seen in tumors

with mutations in *BRAF, ACVR2A, TGFBR2, MSH6,* or *MSH3* (P < 0.0001), which are typical in

MSI tumors [29].

To determine effects on disease outcomes, tumors were stratified for expression of

PLAGL2 using data from two CRC TCGA cohorts, and analyzed for overall survival. In both

cohorts, high expression of PLAGL2 in tumors predicted significantly reduced survival of CRC

patients (Fig 1E, F). Thus, PLAGL2 is commonly turned on in CRC tumors, with high-level

expression likely driving more aggressive disease progression.

We previously generated PLAGL2-mutant DLD1 cell lines [20] using SRIRACCHA [32]

and here we examined their proliferation via a co-expressed H2BGFP reporter, which easily

enabled their enumeration. Each mutant clone exhibited significant deficits in proliferation

compared to non-mutant DLD1 parent cells (Fig. 2A). While defined mutant clones generated

with site-specific nucleases (such as CRISPR) can produce robust loss-of-function models, we

also examined *PLAGL2* loss using CRISPR/Cas9 mutagenesis in a polyclonal population of

mutant Caco2 and HT29 cells. Targeted cells were monitored using a transposon expressing a

nuclear GFP reporter along with guide RNAs (gRNAs) against *PLAGL2* (Fig. 2B). This

transposon was delivered to cells already stably expressing EspCas9. In both Caco2 (Fig. 2C)

and HT29 (Fig. 2D) cells, proliferation was significantly reduced in cells expressing the *PLAGL2-*

specific gRNA. Lastly, shRNA knockdown of *PLAGL2* was pursued by a similar strategy with an

shRNA-expressing transposon, along with a nuclear GFP reporter (Fig. 2E). DLD1 cell lines

established with this transposon revealed several shRNAs that robustly depleted *PLAGL2*

mRNA (Fig. 2F). In Caco2 cells expressing shRNA #3 and #4 (Fig. 2G), proliferation was

significantly reduced, especially for shRNA #4 (Fig. 2H). Effects of *PLAGL2* knockdown were

evaluated in the context of the FUCCI cell cycle reporter [33], delivered to cells using a

transposon vector. In both Caco2 (Fig. 2I) and HT29 (Fig. 2J) cells, the FUCCI reporter revealed

that *PLAGL2* knockdown decreased the number of cells in the $S/G_2/M$ phases of the cell cycle,

but increased the number of cells in G_0/G_1. In sum, PLAGL2 depletion in CRC cell lines

compromises proliferation and cell cycle progression.

Migration, invasion, and the ability to survive anchorage-independent growth are salient

hallmarks of transformed cells. In transwell assays we evaluated migration in the DLD1 mutant

clones and found that migration is severely compromised in both mutants (Fig. 3A, D). Invasion

through a layer of Matrigel was also remarkably reduced in DLD1 mutants, especially mutant #1

(Fig. 3B, E, F). Knock-down of PLAGL2 in Caco2 cells also reduced migration in transwell

assays, especially for shRNA #4 (Fig. 3C). Lastly, *PLAGL2* was over-expressed in IEC6 cells to

evaluate anchorage independent growth in soft agar. *PLAGL2* O/E consistently enabled the

growth and formation of colonies (Fig. 3G, H), demonstrating that PLAGL2 can confer

resistance to anoikis-mediated cell death, a key property of transformed cells.

We next investigated the role of specific PLAGL2 target genes. *IGF2* encodes a critical

fetal growth factor that has been demonstrated to be a direct PLAG1 and PLAGL2 target gene

[34-36], but its role downstream of PLAGL2 in the context of cellular transformation has not

been investigated. Here we find that *IGF2* expression is significantly reduced in *PLAGL2* mutant

DLD1 clones (Fig. 4A) and following shRNA-mediated knockdown (Fig. 4B). In SW480 cells we

performed SRIRACCHA-enriched mutagenesis of *PLAGL2* with CRISPR/Cas9, delivered via

transposon vector. RNA was extracted from a mixed polyclonal population of *PLAGL2* mutants

for expression analysis, and mutagenesis of *PLAGL2* was confirmed by Illumina sequencing

(Fig. S1). In this polyclonal population of *PLAGL2* mutants, *IGF2* expression was significantly

reduced (Fig. 4C). In defined clonal mutants, generated using CRISPR/Cas9 (also delivered via

transposon vector) in mouse organoids [20] *Igf2* expression was also significantly reduced.

Thus, *PLAGL2* is required for expression of *IGF2* in both transformed and non-transformed

intestinal epithelial cells. Over-expression of PLAGL2 in mouse organoids [20] resulted in robust

dose-dependent up-regulation of *Igf2* mRNA (Fig. 4E) and IGF2 protein (Fig. 4F). In sum,

PLAGL2 appears necessary and sufficient to drive *IGF2* expression. To gauge effects of IGF2

alone, the human IGF2 cDNA was over-expressed in mouse intestinal enteroids via transposon

transgenesis (Fig. 4G). IGF2 expression triggered organoid hyperplasia and the formation of

large cysts (Fig. 4H, I), which phenocopies the cyst-like appearance of PLAGL2-overexpressing

enteroids [20].

170   To investigate functional roles of PLAGL2 target genes, the downstream effectors

ASCL2 and *IGF2* were further examined in tumors and transformed CRC cell lines. We

examined CRC tumors from the TCGA for expression correlation between *PLAGL2* and ASCL2

target genes (*ASCL2, KLHDC4, OLFM4, RNF43, LGR5, MYB, NR2E3, SMOC2, OSBPL5,* and

SOX9). Nine out of ten ASCL2 targets showed positive correlation with *PLAGL2* in all three

TCGA CRC cohorts, while most targets were also positively correlated with *PLAGL2* in stomach

adenocarcinoma and hepatocellular carcinoma tumors (Fig. 5A). Consistent with this, CRC cell

lines stratified for the lowest vs. highest quintile of PLAGL2 expression, as determined by

previous RNA-seq studies [37], revealed that expression of *ASCL2* is significantly higher in

PLAGL2-high CRC cell lines (Fig. 5B). SRIRACCHA-enriched mutagenesis of *PLAGL2* in

SW480 cells (Fig. S1) also revealed depletion of ASCL2 mRNA levels. This is all consistent with

a role for PLAGL2 in the direct transcriptional activation of *ASCL2*, as previously demonstrated

in intestinal organoids and CRC cell lines [20]. However, to quantitatively gauge the impact on

ASCL2 activity in CRC cell lines we used the ASCL2 reporter [38], adapted for transposon-

mediated expression of tdTomato (STAR-tdT, Fig. 5D). Co-transfection with shRNA vectors or

an ASCL2 vector revealed robust activation by ASCL2 and modest reduction by shRNAs

against *PLAGL2* (Fig. 5E). After establishing stable transgenic lines, shRNA knockdown of

PLAGL2 caused a pronounced reduction in STAR-tdT reporter activity in SW480, HT29, and

Caco2 cells (Fig. 5F-H).

If ASCL2 and/or IGF2 are critical drivers downstream of PLAGL2, then we expect that

their expression would rescue growth following PLAGL2 loss-of-function. We first examined

proliferation using the FUCCI reporter following knock-down of PLAGL2 and rescue with ASCL2

and/or IGF2. Only ASCL2 was able to restore cell cycle progression in Caco2 cells while neither

IGF2 nor constitutively active β-Catenin (*CTNNB1^{S33Y}*) expression was sufficient (Fig. 5I).

Because of their expression of H2BGFP, DLD1 *PLAGL2* mutants could not be assessed using

the FUCCI reporter. However, in *PLAGL2* DLD1 mutant line #1, ASCL2 and IGF2 individually

augmented clone size following stable transfection, and when co-transfected together we

observed synergistic effects between ASCL2 and IGF2 on growth in this *PLAGL2* mutant (Fig.

5J), suggesting a role for both target genes. To examine a more diverse array of *PLAGL2*

mutants, we performed SRIRACCHA-mediated mutagenesis [32] in Caco2 CRC cell lines while

simultaneously providing PB-mediated transgenic expression of either IGF2 or ASCL2. If either

IGF2 or ASCL2 is able to compensate for loss of PLAGL2, then we would expect increased

survival and growth of clones with *PLAGL2* loss-of-function mutations if such clones also

express exogenous IGF2 or ASCL2. Thus, a rescue would be evident in a higher proportion of

PLAGL2 mutants. Exogenous IGF2 expression does not result in a higher frequency of *PLAGL2*

mutations, but surprisingly, ASCL2 expression results in a significantly lower frequency of

PLAGL2 mutations and higher proportion of non-coding mutations (Fig. 5K). In a 2-step

experiment where lines were first established that already over-expressed ASCL2 or IGF2,

PLAGL2 mutagenesis was not tolerated in Caco2 cells already over-expressing ASCL2; i.e. no

clones were recovered, unlike IGF2 over-expressing or vector controls. This suggests that any

oncogenic effects of ASCL2 depend on PLAGL2, and that these two factors may function as

obligate partners.

Previous studies have reported that PLAGL2 augments Wnt signaling [23, 24, 27],

perhaps via the transcriptional activation of Wnt ligand genes [28]. To explore effects of

PLAGL2 on canonical Wnt signaling, we compared expression levels of *PLAGL2* mRNA with 36

Wnt target genes among 7 cancer datasets from the TCGA (Fig. 6A). While some targets

showed a positive correlation, many targets also showed anti-correlation with PLAGL2, and we

observed no overall trend towards a positive association, except perhaps a slight positive

correlation among liver hepatocellular carcinoma samples (Fig. 6B). In addition, an examination

of fold changes of these Wnt targets in RNA-seq data from PLAGL2-expressing intestinal

organoids showed no clear trend (Fig. 6C). In contrast, data sets directly manipulating Wnt

signaling showed a clear decrease in Wnt target expression following β-Catenin knock-down or

dnTCF4 expression, while stimulation of Wnt signaling in intestinal organoids (via GSK3β

inhibition) showed an induction of Wnt target gene expression (Fig. 6C). Canonical Wnt

signaling is also frequently measured via heterologous reporters. We used our TOP-tdT reporter

[20] to quantify Wnt signaling in the context of PLAGL2 knock-down. Co-transfection of dnTCF4

dramatically reduced signal from this reporter, as expected (Fig. 6D). After establishing stable

shRNA expression in the context of the TOP-tdT reporter, we observe a small decrease in

normalized reporter signal in SW480 and Caco2 cells (Fig. 6E, F), but not in HT29 cells (Fig.

6G). Representative images from Caco2 experiments confirm modest effects on TOP-tdT signal

(Fig. 6H). Despite subtle effects in SW480 and Caco2 cells, no significant effects were observed

on non-phosphorylated ("active") β-Catenin or total β-Catenin protein levels in these cells (Fig.

6I, J). In DLD1 *PLAGL2* mutant cell lines, similar effects were observed. Values were quantified

from these immunoblot experiments (Fig. 6 K-M). Expression analysis of Wnt target genes also

did not reveal any clear trend following *PLAGL2* knock-down (Fig. 6N-P). SRIRACCHA-

mediated mutagenesis of *PLAGL2* in SW480 cells (Fig. S1) and RT-PCR revealed a slight trend

of reduced Wnt target gene expression, but only *ETS2* mRNA levels were significantly reduced

(Fig. 6Q, R). In sum, the data only support a small effect of *PLAGL2* on Wnt signaling in CRC

cells.

DISCUSSION

We find that PLAGL2 is a transcriptional regulator of cellular transformation in colorectal

cancer, being able to drive hallmark features, including unchecked proliferation, migration,

invasion, and anchorage independent growth. The normal physiological role of PLAGL2 is likely

integral to a developmental pathway considering its high expression in the fetal intestinal

anlagen and early postnatal intestine, but low expression in adult tissue [39]. *Plagl2*$^{-/-}$ mice

demonstrate defects in intestinal epithelial differentiation and function, suggesting that PLAGL2

plays a critical role during fetal intestinal development [39]. Similar to many other Let-7 targets,

such as *HMGA2*, *PLAGL2* appears to exhibit typical features of an onco-fetal gene, considering

its frequent re-activation in CRC and other malignancies [20, 28, 40-42]. However, there has

been little insight into the role of direct transcriptional targets of PLAGL2, either during fetal

development or carcinogenesis.

In possible oncogenic roles, PLAGL2 has been shown to directly activate transcription of

the thrombopoietin receptor (*MPL*) [43], which is a key receptor necessary for megakaryocyte

and platelet formation [44]. MPL is also a proto-oncogene that can drive hematopoietic cell

proliferation when it (or the truncated *v-mpl* oncogene) is over-expressed [45, 46]. Relevant to

PLAGL2, MPL is implicated in driving transformation downstream of PLAGL2 in acute myeloid

leukemia (AML) [43], a malignancy in which increased expression of MPL marks a particularly

aggressive subset [47]. However, even though MPL has been shown to be expressed on a

subset of CRC cells that have metastatic tropism for the liver and lung [48, 49], *MPL* expression

is not induced by PLAGL2 in intestinal organoids [20] and *MPL* expression does not correlate

with *PLAGL2* expression in CRC [29]. Thus, *MPL* may not be a relevant oncogenic PLAGL2

target gene in CRC. Other potential PLAGL2 targets, such as *NIP3* and *P73* [50, 51], appear to

have tumor-suppressive properties, while another documented target, surfactant protein C

(*SPC*) [52], has unknown relevance to tumorigenesis and/or cellular transformation.

We have previously identified *ASCL2* as a direct target gene of PLAGL2. ASCL2 is a

bHLH transcription factor that directly supports *LGR5* expression [22], drives IESC fate [21, 22,

53], and promotes an aggressive phenotype in CRC [53-55]. Regarding this effect in CRC,

ASCL2 represses expression of *CDX2*, a transcription factor with a well-described role in

positively driving intestinal epithelial differentiation [54]. ASCL2 also represses expression of

miR-200 miRNAs, which may underlie effects on a mesenchymal phenotype and/or EMT in

cancer, which is repressed by the miR-200 family of miRNAs [55]. Studies have also suggested

that ASCL2 may play a role in augmenting the tumor-initiating capacity of CRC cells [56, 57],

although this has not yet been carefully examined. Ultimately, the role of *ASCL2* may prove to

be pleiotropic, as these studies suggest, with variable effects on differentiation, EMT, and tumor

initiating potential.

Surprisingly, in our studies, ASCL2 over-expression confers a survival disadvantage to

PLAGL2 mutants in the Caco2 cell line, even though we see that cell cycle progression is

promoted by ASCL2 rescue in the context of PLAGL2 knock-down. Thus, ASCL2 alone may

have different effects than when co-expressed with PLAGL2 at high levels; such effects of

ASCL2 may repress cellular transformation when PLAGL2 levels are low. Previous studies

seeking to identify ASCL2 co-immunoprecipitating proteins included the identification of

PLAGL2, as determined by mass spectrometry [38], although such direct interaction was not

verified by other methods. If direct interaction can occur between these factors, then the effects

we see on the STAR-tdT reporter in CRC cell lines could possibly reflect direct roles for

PLAGL2 on ASCL2-responsive regulatory elements. While follow-up studies are needed,

PLAGL2 and ASCL2 may biochemically cooperate to drive specific oncogenic targets – targets

that may differ from those that are transcriptionally activated individually by each individual

factor.

For effects on tumorigenesis, a gain-of-function role for ASCL2 (e.g. in the context of

ASCL2 up-regulation) is not clear. Modest entopic over-expression of *Ascl2*, at levels 2-3-fold

higher than non-transgenic mice, via a BAC transgene, does not accelerate CRC tumorigenesis

in the context of the *Apc^{Min}* allele [58], a tumor model in which mice develop adenomatous

polyps in the small intestine and colon. However, DNA replication (as measured by

bromodeoxyuridine incorporation) is significantly elevated in the intestinal epithelium of these

Ascl2 transgenic mice, suggesting that elevated levels of ASCL2 can drive cell cycle

progression [58]. Consistent with these *in vivo* findings, we observe accelerated cell cycle

progression following ASCL2 over-expression in the context of PLAGL2 knock-down. In contrast

to SW480 cells, which express very low levels of ASCL2 mRNA and protein, Caco2 cells

express very high levels of ASCL2 [54]. Thus, ASCL2 may play a differential role downstream of

PLAGL2, depending on the individual tumor or cancer cell line. In light of *in vivo* studies

described above, *ASCL2* may only have cancer-promoting effects when *PLAGL2* is also

coordinately up-regulated and/or activated. The co-expression of *PLAGL2* with *ASCL2* is

striking in CRC [20], so the context for cooperation certainly exists in tumors.

The PLAGL2 paralog, PLAG1, is reported to transcriptionally activate the IGF2 promoter

[34, 36], with over-expression experiments in NIH-3T3 and HEK293 cells suggesting that

PLAGL2 may also positively regulate *IGF2* expression. IGF2 (a known driver of tumor

progression) is over-expressed in ~15% of CRCs and also activates the PI3K-AKT pathway [59-

65]**.** IGF2 up-regulation also occurs in the context of wild-type *PTEN*, *PIK3CA*, *BRAF*, and

KRAS [29], suggesting that IGF2 over-expression can functionally substitute for common PI3K-

AKT-activating mutations. Here we provide data in primary intestinal organoids and human CRC

cell lines that PLAGL2 is necessary and sufficient to drive *IGF2* expression. Following knock-out

or knock-down of *PLAGL2* in CRC cell lines, we observe that *IGF2* expression is depleted up to

95%, and in a CRC cell line such as Caco2, which is known to express very high levels of IGF2

[66-68]), severe reduction of IGF2 has consequential effects on cellular growth. Evidence for

this exists through studies of Caco2 cells, which are sensitive to both a neutralizing anti-IGF2

antibody [66] and an IGF1R/INSR inhibitor [68], which blocks IGF2 or the receptors through

which IGF2 signals, respectively. However, we cannot rescue cellular proliferation and/or

survival following *PLAGL2* mutagenesis in Caco2 cells via over-expression of IGF2. Thus, the

growth defects caused by PLAGL2 loss are due to other effectors, besides IGF2. In *PLAGL2*

mutant DLD1 cells ASCL2 and IGF2 each alone have modest effects on clone proliferation, but

cooperative/synergistic effects when co-expressed. Therefore, in some contexts these PLAGL2

targets cooperate, whereas in other contexts, such as Caco2 cells, we do not see evidence of

cooperation. Thus, the oncogenic dependency of PLAGL2 on IGF2 and ASCL2 is context-

dependent, varying between tumors and/or tumor types.

Finally, our data suggest a minor role for Wnt signaling downstream of PLAGL2 in CRC.

While previous studies indicate that WNT6 is a PLAGL2 target gene [27, 28], and our previous

studies show that PLAGL2 activates the expression of Wnt genes (i.e. *Wnt9b*, *Wnt4*, *Wnt10a*,

and *Wnt5a*) in primary intestinal epithelial cells [20], our studies here in CRC cell lines indicate

that canonical Wnt signaling (via TCF4/β-Catenin) are only modestly affected by PLAGL2.

Additionally, in available data from TCGA we find that PLAGL2 levels do not correlate with a

Wnt signature, as measured by the induction of Wnt target genes. If the effects of PLAGL2 on

Wnt signaling are routed via Wnt ligands (such as WNT6), then a minor effect on Wnt activation

is not unexpected, since the vast majority of colorectal cancers possess mutations in either

APC, *AXIN2*, or *CTNNB1* [29] — mutations that lead to ligand-independent Wnt pathway

activation. Other PLAGL2 target genes, besides Wnt ligands themselves, could be intracellular

modifiers of the canonical Wnt signaling pathway. However, given the modest effects of

PLAGL2 on canonical Wnt signaling in CRC cells, the net effect of any such hypothetical targets

is likely to be relatively small. This underscores the need to identify additional PLAGL2 target

genes, especially those involved in fetal growth pathways and carcinogenesis; identifying such

targets will help illuminate the onco-fetal pathways downstream of PLAGL2.

Figure Legends

Figure 1. *PLAGL2* is turned on in the majority of colorectal cancers and is predictive of

overall survival. A) *PLAGL2* mRNA expression in 10 pairs of colon adenocarcinomas, with

matched non-tumor adjacent tissue, as previously described [10]. **B)** *PLAGL2* expression in

non-tumor and colorectal adenocarcinoma tumors from the TCGA PanCancer Atlas cohort [69]

C) RT-PCR assay quantifying relative levels of *PLAGL2* mRNA in colorectal cancer cell lines

and HEK293T cells. **D)** Oncoprint showing mutation and expression profiles of commonly

mutated cancer genes, including *PLAGL2*, from the TCGA PanCancer Atlas of colorectal

adenocarcinomas. **E)** Overall survival of CRC patients stratified for high expression of PLAGL2

compared to all other tumors, based on microarray expression data [29]. **F)** Overall survival of

CRC patients stratified for high expression of PLAGL2 compared to low expression based on

microarray expression data (TCGA Provisional). Statistical significance was evaluated using

Student's paired t-test **(A)**, Student's one-tailed t-test **(B)**, or a Mantel-Cox log-rank test **(E, F)**.

Figure 2. Knock-down or knock-out of *PLAGL2* compromises proliferation and cell cycle

progression in CRC cell lines. A) Proliferation of stable CRISPR-mutated *PLAGL2* mutant

DLD1 clones [20] compared to parental DLD1 cells. **B)** PB vector for stable expression of

gRNAs and NLS-GFP, with Puro resistance. Caco2 **(C)** or HT29 **(D)** cell lines stably expressing

EspCas9 were transfected with a PB vector expressing nuclear GFP and a non-specific (NS)

gRNA or a gRNA against PLAGL2, and then growth was monitored. **E)** PB vector for stable

expression of shRNAs and NLS-GFP, with Puro resistance. **F)** Knock-down (KD) of PLAGL2

was assessed by RT-PCR in DLD1 cells **(F)** and also confirmed in Caco2 **(G). H)** Proliferation of

Caco2 cells stably transfected with PB vectors expressing nuclear GFP and non-specific (NS) or

PLAGL2-specific shRNAs. Cell cycle status using the FUCCI [33] reporter co-transfected with a

non-fluorescent shRNA PB vector was evaluated in Caco2 **(I)** and HT29 **(J)** CRC cells. For **A-D,**

and **H** proliferation was quantified through enumeration of fluorescently labeled nuclei

expressing NLS-GFP. Statistical significance (*$p < 0.05$ or **$p < 0.01$) was evaluated through an

ordinary one-way ANOVA and Dunnett's multiple comparisons post-hoc test.

Figure 3. PLAGL2 drives migration, invasion, and colony formation in soft agar. Assays

measuring migration **(A)** or invasion through Matrigel **(B)** of stable CRISPR-mutated *PLAGL2*

mutant DLD1 clones [20] compared to parental DLD1. **C)** Migration assay of Caco2 cells stably

transfected with PB vectors expressing nuclear GFP and a non-specific (NS) or a PLAGL2-

specific shRNA. Images of GFP-expressing cells that have migrated through porous trans-well

membranes 24 hours after plating for parental DLD1 cells **(D)**, DLD1 mutant clone #1 **(E)** or

mutant clone #2 **(F)**. Soft agar colony forming assay for IEC6 cells transfected with a PB empty

vector **(G)** or a PB vector expressing PLAGL2 **(H)**. Statistical significance (*$p < 0.05$ or **$p < 0.01$)

was evaluated through an ordinary one-way ANOVA and Dunnett's multiple comparisons post-

hoc test.

Figure 4. PLAGL2 drives IGF2 expression in CRC cells and intestinal organoids and

partially rescues growth phenotype in *PLAGL2* mutant CRC cells. A) RT-PCR for *IGF2* in

DLD1 *PLAGL2* mutant clones. **B)** RT-PCR for *IGF2* in Caco2 cells following shRNA KD of

PLAGL2. **C)** RT-PCR for *IGF2* following CRISPR/Cas mutagenesis using SRIRACCHA [32] and

Hygromycin selection. **D)** RT-PCR for *Igf2* in *Plagl2* knockout mouse intestinal enteroids [20]

compared to WT parental enteroids. **E)** RT-PCR for *Igf2* in 3 lines of mouse intestinal enteroids

expressing low, medium, or high levels of human HA-tagged PLAGL2 [20] compared to empty

vector control enteroids. **F)** Immunoblot for HA-tag and IGF2 in low and high-expressing

PLAGL2 O/E enteroid lines. **G)** RT-PCR for IGF2 mRNA following O/E in enteroids. **H+I)** Cyst-

like morphology in IGF2 O/E enteroids. For comparison of more than 2 conditions the statistical

significance (*p<0.05 or **p<0.01) was evaluated through an ordinary one-way ANOVA and

Dunnett's multiple comparisons post-hoc test. For pair-wise comparison, Welch's t-test was

performed to determine statistical significance (*p<0.05 or **p<0.01).

Figure 5. PLAGL2 activates an ASCL2-dependent program in CRC cells. A) Plots of

correlation coefficients of linear regressions between *PLAGL2* and ten ASCL2 direct target

genes (*ASCL2, KLHDC4, OLFM4, RNF43, LGR5, MYB, NR2E3, SMOC2, OSBPL5,* and *SOX9*)

for seven TCGA datasets, including 3 from CRC adenocarcinoma (AdCA), 2 from stomach

AdCA, and 2 from liver hepatocellular carcinoma (HCC). **B)** Colorectal cancer cell lines (N=154

from [37]) were parsed into bottom and top quintiles for *PLAGL2* expression and then evaluated

for *ASCL2* mRNA expression. **C)** EspCas9-mediated mutagenesis using SRIRACCHA [32] and

gRNAs directed against LacZ or *PLAGL2*, followed by selection with hygromycin and RT-PCR

for *ASCL2* mRNA. **D)** Depiction of the STAR ASCL2 reporter construct. **E)** Activity of the stem

cell ASCL2 reporter (STAR-tdT) reporter [38] 48 hours after transfection in SW480 CRC cells

that were also co-transfected along with vectors providing *PLAGL2* shRNA-mediated KD or

ASCL2 O/E. As expected ASCL2 O/E augments STAR-tdT reporter activity. After selection with

Puromycin for 6-8 days for stable *PLAGL2* shRNA expression STAR-tdT reporter activity

(integral RFP fluorescence) was measured in SW480 **(F)**, HT29 **(G)**, and Caco2 **(H)** CRC cell

lines. RFP levels were normalized to integral GFP fluorescence constitutively expressed by the

PB shRNA vector. **I) FUCCI** analysis of cell cycling in Caco2 cells after shRNA mediated KD of

PLAGL2 and exogenous expression of PLAGL2, ASCL2, IGF2, ASCL2 and IGF2, or a

constitutively active β-catenin. **J)** Rescue of DLD1 *PLAGL2* mutant clone #1 with IGF2 and/or

ASCL2 O/E and assessment of colony size following selection with G418. **K)** Mutation profile of

PLAGL2 targeted with SRIRACCHA using eSpCas9 with simultaneous over-expression of

ASCL2 or IGF2. For comparison of more than 2 conditions the statistical significance (*p<0.05

or **p<0.01) was evaluated through an ordinary one-way ANOVA and Dunnett's multiple

comparisons post-hoc test. For pair-wise comparison, Welch's t-test was performed to

determine statistical significance (*p<0.05 or **p<0.01).

Figure 6. Minimal effects on canonical Wnt signaling following inactivation of PLAGL2. A)

Heatmap of correlation coefficients for linear regressions between *PLAGL2* and 36 TCF4/β-

Catenin target genes for seven TCGA datasets, including 3 from CRC adenocarcinoma (AdCA),

2 from stomach AdCA, and 2 from liver hepatocellular carcinoma (HCC). **B)** Dot/scatter plots of

data from **(A). C)** Violin plots of fold changes among 28 TCF4/β-Catenin target genes for

conditions as follows: 1) between PLAGL2-low vs. PLAGL2-high CRC AdCA tumors (TCGA), 2)

between control and KD of *CTNNB1* (β-Catenin) or inducible O/E of a dnTCF4 in LS174T CRC

cell lines [70], and 3) GSK3β inhibition and Wnt/TCF4/β-Catenin activation in intestinal

organoids following treatment with 5 μM CHIR99021 [71]. **D)** Activity of the Wnt Super 8x

TopFlash reporter [72] driving expression of tdT (TOP-tdT) 48 hours after transfection in SW480

CRC cells that were also co-transfected along with vectors providing *PLAGL2* shRNA-mediated

KD or dnTCF4 O/E. As expected, dnTCF4 O/E reduces signal from the TOP-tdT reporter and

reduces the number of RFP-positive cells. After selection with Puromycin for 6-8 days for stable

PLAGL2 shRNA expression TOP-tdT reporter activity (integral RFP fluorescence) was

measured in SW480 **(E)**, Caco2 **(F)**, and HT29 **(G)** CRC cell lines. RFP levels were normalized

to integral GFP fluorescence constitutively expressed by the PB shRNA vector. **H)**

Representative images of fluorescent Caco2 cells after 6 days of selection with 10 μg/mL

puromycin. Active (non-phosphorylated) and total β-Catenin levels were measured following

PLAGL2 KD in SW480 **(I)**, Caco2 **(J)**, and *PLAGL2* mutant DLD1 clones **(K)**. Protein levels were

quantified and normalized to tubulin **(L-N)**. Direct TCF4/β-Catenin target gene expression levels

were measured by RT-PCR in SW480 and Caco2 cells following PLAGL2 KD **(O, P)**, in *PLAGL2*

mutant DLD1 clones **(Q)**, and following SRIRACCHA mediated mutagenesis of *PLAGL2* in

SW480 cells **(R)**. Statistical significance (*p<0.05 or **p<0.01) was evaluated through an

ordinary one-way ANOVA and Dunnett's multiple comparisons post-hoc test.

Figure S1. Indel and substitution mutations in SW480 cells around three target

SRIRACCHA based CRISPR/Cas9 cleavage sites in *PLAGL2*. A) Mutation distribution

around cleavage sites in *PLAGL2* gene with NS gRNA, showing no events. **B)** Mutation

distribution around the three target cleavage sites in *PLAGL2* **C)** Example allele profile of

SRIRACCHA reads from these *PLAGL2* mutations. Note, the top hit represents unmodified

PLAGL2.

MATERIALS AND METHODS

RT-PCR

CRC tumor samples, along with paired non-cancerous controls, were obtained from Siteman

Cancer Center Tissue Procurement Core and were previously described [10]. RNA was

extracted with TRIzol (ThermoFisher Scientific) and further purified using the RNeasy RNA

Clean-up Kit (Qiagen) for these tumor samples. For cell lines or enteroids, RNA was prepared

using TRIzol. RT reactions were performed with 3-4 μg total RNA using oligo-dT and

SuperScript III RT (ThermoFisher Scientific). QPCR was performed as previously described

[20], and expression levels normalized to *PPIA* and *B2M* for human specimens or human cell

lines. For RT-PCR using RNA from mouse organoids or cell lines expression levels were

normalized to *Hprt* and *Tbp.*

TCGA Analysis

TCGA data was examined using the cBioPortal for Cancer Genomics [30, 31], hosted by the

Center for Molecular Oncology at MSKCC. Oncoprint analysis (Fig. 1D) was performed for CRC

AdCA tumors from the PanCan study [73]. Overall survival rates among CRC patients from the

TCGA were determined from analysis of microarray data from two cohorts parsed for

expression of *PLAGL2*. For one CRC dataset [29] PLAGL2-high tumors were those with greater

than 3 SD above the mean for PLAGL2 expression, while another CRC dataset (TCGA-

provisional) *PLAGL2*-high tumors were defined as those with greater than 2 SD above the mean

for *PLAGL2* expression and *PLAGL2*-low tumors were defined as those with more than 0.25 SD

below the mean for *PLAGL2* expression.

PLAGL2 Mutagenesis with CRISPR/eSpCas9 and SRIRACCHA

For expression of eSpCas9 [74], the ORF from eSpCas9 was amplified from eSpCas9(1.1),

which was a gift from Feng Zhang (Addgene plasmid # 71814) and cloned into BII-ChBtW,

which is identical to BII-ChPtW (which confers Puromycin resistance through PAC), but confers

Blasticidin resistance. Using 8 µL Lipofectamine 2000 (ThermoFisher Scientific) this vector, BII-

ChBtW-eSpCas9, was introduced into Caco2 and HT29 cells by transfecting 1.6 µg of

transposon along with 400 ng of pCMV-hyPBase [75], a generous gift from Dr. Allan Bradley

(Wellcome Sanger Institute). Selection with 10 µg/ml Blasticidin was initiated 48 hours after

transfection and continued for 7 days, whereupon cells were expanded for transfection with BII-

gR-PnGW. The BII-gR-PnGW vector contains a U6-driven gRNA cassette cloned from pX335,

which was a gift from Feng Zhang (Addgene plasmid # 42335, [76]). This U6-driven gRNA

module was cloned 5' of the CMV/hEf1a promoter at a unique *Sfi*I site, and was modified to

contain two *BsmB*I sites for cloning gRNA oligos, in place of the existing *Bbs*I sites. The BII-gR-

PnGW also constitutively expresses GFP-NLS for visualization of transduced cells.

Transfections of the BII-gR-PnGW vector were performed in quadruplicate in 24-well plates 24

hours after plating 5×10^4 cells per well. Cells were transfected as above using 400 ng BII-gR-

PnGW and 100 ng pCMV-hyPBase, selected with 5 µg/mL Puromycin for 4 days (HT29) or 10

µg/mL Puromycin for 10 days (Caco2). Fluorescence was measured starting 48 hours after

transfection. A non-specific gRNA (GGAGACGCTGACCCGTCTCT) was used as a control for

comparison with a PLAGL2-targeting gRNA (GTTCACCGCAAGGACCATCTG) [20].

SRIRACCHA-mediated mutagenesis was performed using the BII-gR-PtW-eSpCas9 vector,

which contains a gRNA-expression cassette (as in BII-gR-PnGW) and confers resistance to

Puromycin. SW480 (8x10^5 cells) or Caco2 (3x10^5 cell) were plated in triplicate or

quadruplicate in 6-well plates 24 hours prior to transfection. For one-step transfection with

SRIRACCHA components, 600 ng of the BII-gR-PtW-eSpCas9 vector with gRNAs specific for

either PLAGL2 (GTTCACCGCAAGGACCATCTG) or LacZ (GAAGGCGGCGGGCCATTACC)

were transfected along with 600 ng of the BII-C3H target vector [32] containing target

sequences either for PLAGL2 or LacZ cloned at the 3' end of the PAC ORF, 500 ng of pBS-PtH,

and 300 ng of pCMV-hyPBase. SW480 cells were transfected using JetPrime (Polyplus Inc.) (2

µL/µg DNA) per manufacturer instructions while Caco2 cells were transfected using

Lipofectamine 2000 (ThermoFisher Inc.) (4 µL/µg DNA) per manufacturer instructions. Cells

were selected with Puromycin for 5 days, then switched to Hygromycin selection for 10 days.

RNA was then isolated using TRIzol or, alternatively, both RNA and DNA were isolated using

the Allprep DNA/RNA Mini Kit (Qiagen Inc.). Mutation analysis was performed either on cDNA

(SW480) or gRNA (Caco2). Mutation of *PLAGL2* was determined by amplicon sequencing on

the Illumina platform by the Center for Genomic Sciences (Washington University) and analysis

was performed using CRISPResso [77]. For transfection with SRIRACCHA components in

rescue experiments, Caco2 cells were transfected in a one- or two-step manner. For the two-

step method, 3×10^5 cells in a 6-well plate that had been seeded the previous day were

transfected with 800 ng BII-C3H-PLAGL2-T1 vector containing the PLAGL2 targeting gRNA

sequence, 800 ng of either BII-ChBtW (empty vector), BII-ChBtW-ASCL2 (overexpressing

ASCL2), or BII-ChBtW-IGF2 (overexpressing *IGF2*), and 400 of pCMV-hyPBase. Transfection

medium was removed in 24 hours and replaced with growth medium. Growth medium was

changed in a further 24 hours for 10 μg/mL Puromycin and 10 μg/mL Blasticidin containing

growth medium, and cells were selected for resistance for 8 days with regular media changes.

Once cells reached confluency again, they were trypsinized and removed from the plate, and 3

x 10^5 cells were again seeded, this time into three identical wells of a 6-well plate. The following

day, cells were transfected with 1,600 ng of the BII-gR-PtW-eSpCas9-PLAGL2-T1R vector, the

Cas9 activity surrogate vector, 750 ng of the pBS-PtH vector, which contains the hygromycin

resistance cassette, and 250 ng of pCMV-hyPBase. Transfection medium was removed and

changed as before, and 48 hours post transfection medium containing 400 μg/mL Hygromycin

was added. Cells were selected for Hygromycin resistance for 13 days.

For the one-step method, 3 x 10^5 cells seeded the previous day into a 6-well plate were

transfected with 400 ng BII-C3H-PLAGL2-T1 vector containing the PLAGL2 targeting gRNA

sequence, 400 ng of either BII-ChBtW (empty vector), BII-ChBtW-ASCL2 (overexpressing

ASCL2), or BII-ChBtW-IGF2 (overexpressing *IGF2*), 500 ng of the BII-gR-PtW-eSpCas9-

PLAGL2-T1R vector, 300 ng of the pBS-PtH vector, and 400 ng of the pCMV-hyPBase vector.

Transfection medium was removed and changed again in 24 hours, and 48 hours post

transfection cells were selected with 10 μg/mL Puromycin and 10 μg/mL Blasticidin for 5 days,

before changing to medium containing 400 μg/mL Hygromycin for 11 days.

For both methods, after Hygromycin selection was finished (cells reached confluency), gDNA

was extracted and mutation of *PLAGL2* was again determined by amplicon sequencing on the

Illumina platform by the Center for Genomic Sciences (Washington University) and subsequent

analysis by CRISPResso.

Cell Proliferation Assays

Cellular proliferation was quantified by automated microscopy of GFP-positive cells on a Biotek

Cytation3 using a 4x objective, where cell numbers were gauged by enumerating individual cells

or by quantifying total integrated fluorescence per well. DLD1 *PLAGL2* mutants [20] express

H2BGFP while other cell lines express nuclear GFP following stable transfection with BII-

ShPnGW or BII-gR-PnGW. Cellular fluorescence was imaged every 1-2 days and media

changed every 2-3 days.

Cell Cycle Analysis with FUCCI Reporter

A *Piggybac* version of the FUCCI reporter (BII-ChPtW-iresFUCCI) was constructed by PCR

amplification of the Clover-Geminin-ires-mKO2-Cdt fragment from pLL3.7 (Addgene Plasmid

#83841, [33]) and insertion into unique *BsmB*I sites within the *Piggybac* vector BII-ChPtW. The

BII-ChPtW vector was constructed by insertion of the Woodchuck Hepatitis Virus (WHP)

Posttranscriptional Regulatory Element (WPRE) downstream of the BsmBI site in the previously

described BII-ChPt vector [20]. Transfections of the BII-ChPtW-iresFUCCI vector were

performed in quadruplicate in 24-well plates 24 hours after plating 5×10^4 cells per well. Cells

were transfected as above using 400 ng of PB transposon and 100 ng pCMV-hyPBase,

selected with 5 µg/mL Puromycin for 4 days (HT29) or 10 µg/mL Puromycin for 10 days

(Caco2).

PLAGL2 shRNA-mediated Knockdown

For protein knockdown, an shRNA expression *Piggybac* vector, BII-ShPnGW was constructed

by insertion of a GFP-NLS cDNA into the BsmBI cloning sites of BII-ChPtW. A U6-driven shRNA

module was then inserted at a unique *Sfi*I site upstream of the CMV/hEf1a promoter, with

*BsmB*I cloning sites for insertion of unique shRNA double-stranded oligos. The BII-ShPiRW

vector was constructed for FUCCI experiments. This vector is identical to BII-ShPnGW, but

expresses the iRFP702 near-IR protein [78], which was first codon optimized, synthesized and

cloned from a gBlock fragment (IDT Inc.). Alternatively, for rescue experiments (Fig. 6), the

shRNA module from BII-ShPnGW was cloned as an *Nsi*I-*Age*I fragment into those unique

restriction sites in BII-ChPtW-iresFUCCI to generate BII-Sh-FUCCI. The Broad Gene

Perturbation Portal was queried for identification of candidate shRNAs against *PLAGL2*, and

oligos prepared identical to the strategy employed for shRNA expression from the pLKO.1

vector [79], except overhangs were modified for directional cloning into our BII-ShPnGW vector,

with 5' overhangs ACCG and AAAA at the termini of each dsDNA oligo. The following four

PLAGL2-specific target sequences were cloned in this manner: shRNA #1:

TTCAGGCTCTAGGATCGATTC, shRNA #2: CCGTAGGACTTCAGGTATTAT, shRNA #3:

TTGGATGACCTCTAGAGAAAT, shRNA #4: GCAGGAGAGAAGGCCTTTATT. A luciferase-

specific shRNA was used as a control, with target sequence TCACAGAATCGTCGTATGCAG.

The BII-ShPnGW vector with each specific shRNA was transfected into DLD1 cells in triplicate

in 6-well plates, with 1.6 µg of transposon transfected along with 400 ng of pCMV-hyPBase.

Selection was initiated 48 hours after transfection using 10 µg/mL Puromycin for Caco2 cells, or

5 µg/mL for other cell lines, and continued for 4-7 days. Cells were then cultured for 1-2 days in

medium without Puromycin prior to harvesting RNA or protein for RT-PCR or immunoblot.

Migration and Invasion Assays

For migration assays 1×10^5 cells were plated in 0.5 mL DMEM in Falcon Fluoroblok trans-well

inserts containing 8 µm pores, which were placed into Falcon 24-well plates, with each well

containing 0.75 mL DMEM and 10% FBS. Plates were cultured 24 to 48 hours and then imaged

for GFP fluorescence using the Biotek Cytation3. For invasion assays Falcon Fluoroblok trans-

well inserts were coated with cold 25% Matrigel (75% DMEM), which was then solidified for one

hour at 37°C. Cells were then added to transwell inserts and cultured and imaged as above for

migration assays. DLD1 mutant clones (#1 and #2) and parental cells express H2BGFP for

visualization (Strubberg), while the shRNA knock-down vector (BII-ShPnGW) provided nuclear-

localized GFP for enumeration of migratory cells by microscopy using the Biotek Cytation3.

Soft Agar Colony Formation Assays

IEC6 cells were cultured in Advanced-DMEM/F12 (ThermoFisher Inc.) containing 5% FBS,

Glutamax (ThermoFisher Inc.), HEPES, N2 Supplement (ThermoFisher Inc.), Plasmocin

(Invivogen Inc.), and Penicillin/Streptomycin. Over-expression vectors were prepared by cloning

the PLAGL2 coding sequence into BII-ChBtW and IEC6 cells transfected in 6 well plates with

1600 ng of this plasmid and 400 ng of pCMV-hyPBase using Lipofectamine 2000

(ThermoFisher Inc.). Cells were select 48 hours after transfection with medium containing 4

µg/ml Blasticidin and selected for 5 days. To assay anoikis resistance and anchorage-

independent proliferation IEC6 cells were grown in 0.35% low-melt agarose (Lonza SeaPlaque).

A base layer of agarose was prepared by melting pre-sterilized 1.2% agarose in dH2O at 75°C,

and cooling for 60 minutes at 42°C. This was then mixed with pre-heated 0.22 µm-filtered 2x

DMEM containing 20% FBS, Sodium Bicarbonate, Sodium Pyruvate, Glutamax (ThermoFisher

Inc.), HEPES, 2x N2 Supplement (ThermoFisher Inc.), Plasmocin (Invivogen Inc.), and

Penicillin/Streptomycin. This mixture (1 mL) was applied to the bottom of an ultra-low

attachment 6-well plate (Greiner Bio-One Inc.) and allowed to cool for 5 minutes at room

temperature. Cells were detached with 0.05% Trypsin-EDTA and counted and placed on ice.

Pre-sterilized 0.7% agarose was prepared as above, equilibrated for 60 minutes at 42°C, and

mixed with pre-warmed growth medium, and then incubated 30 minutes at 37°C. Cells (8×10^4)

were added to 3.2 mL of this 0.35% agarose, mixed well, and then 1 mL overlaid onto each well

containing a base layer of 0.6% agarose in 6-well plates. Plates were then incubated 20 minutes

at room temperature and then placed into a 37°C incubator with 6% CO2. The next day 1 mL of

complete culture medium was added. Colonies were counted 3 weeks following plating.

Organoid Culture and Transfection

Mouse small intestine organoids (enteroids) were generated from jejunum crypts from

C57BL/6 mice (8-12 weeks) and cultured and transfected as previously described [20]. *Plagl2*

mutant/KO and PLAGL2 over-expressing (O/E) mouse enteroids were previously described

[20]. IGF2 O/E enteroids were generated by cloning the human IGF2 ORF into the BII-ChBtW

vector via *Bsm*BI sites, and transfection in mouse enteroids using 800 ng of this BII-ChBtW-

IGF2 vector along with 200 ng of pCMV-hyPBase. Enteroid transfections were performed as

previously described [20]. Multiple (n=20-50) Blasticidin-resistance clones were obtained and

pooled for stable propagation of a polyclonal line. IGF2 expression was assayed by RT-PCR

and morphology documented using the Cytation3 microscopy platform and a 4x objective.

ASCL2 and Wnt Reporter Assays

The ASCL2 reporter was previously described [38]. Briefly, this reporter was adapted for

Piggybac transposon-mediated gene delivery by cloning ASCL2 regulatory elements

downstream of 2 polyadenylation signals (one synthetic and one from the SV40 TK gene) and

upstream of a tandem tomato (tdT) fluorescent protein ORF and NLS signal, followed by the

bovine growth hormone (bGH) polyadenylation signal, all flanked by HSIV core insulators from

the chicken *HBBA* gene. For transfection, SW480 (2x10^5 cells), HT29 (2x10^5 cells), or Caco2

(1x10^5 cells) were plated in triplicate or quadruplicate in 24-well plates 24 hours prior to

transfection. Cells were transfected with 200 ng of BII-STAR-tdT, 200 ng of BII-ShPnGW, and

100 ng of pCMV-hyPBase. The BII-ShPnGW vector contained shRNA sequences directed

against a non-specific target or the *PLAGL2* mRNA, as described above. For a positive control

of reporter activity BII-STAR-tdT was co-transfected with 200 ng of BII-ChBtW-ASCL2 for

ASCL2 O/E. Red fluorescence was quantified 48 hours after transfection using the Cytation3

platform. Cells were then selected with Puromycin for 5 days and then both GFP and RFP

fluorescence were quantified using the Cytation3. The BII-TOP-tdT reporter for the canonical

Wnt pathway was described previously [20], but was modified for nuclear expression with a C-

terminal NLS. This vector was transfected and visualized in SW480, HT29, and Caco2 cells as

described above for BII-STAR-tdT. For a positive control of this fluorescent Wnt reporter, BII-

ChBtW-dnTCF4 was co-transfected with BII-TOP-tdT. This dominant negative TCF4 variant was

described previously [80] and was modified here for expression via transposon. Red

fluorescence was quantified 48 hours after transfection using the Cytation3 (Agilent-Biotek Inc.)

platform and subsequently after antibiotic selection, as described above for the BII-STAR-tdT

reporter.

FUCCI Rescue Experiments

For rescuing cell cycle effects caused by PLAGL2 knockdown the FUCCI vector was modified

for simultaneous shRNA expression by cloning the *Nsi*I – *Age*I fragment from BII-ShPnGW,

which includes shRNA components, into BII-ChPtW-iresFUCCI to create the BII-shFUCCI

vector. Caco2 were plated in quadruplicate for transfection with BII-shFUCCI along with BII-

ChBtW vectors for O/E of PLAGL2, ASCL2, IGF2, or CTNNB1^{S33Y}. The constitutively active

beta-catenin mutant (CTNNB1^{S33Y}) was described previously [81]. Beginning 48 hours after

transfection cells were selected with Puromycin and Blasticidin for 5 days and then RFP and

GFP fluorescence were quantified using the Cytation3.

665
Immunoblots and Protein Quantification

For analysis of active and total β-Catenin levels upon *PLAGL2* mRNA KD, Caco2 and SW480

cells in a 6 well dish were transfected with the shRNA construct and selected as previously

described, in triplicate. After the initial selection and subsequent expansion of cells to near

confluency, cells were harvested and lysate was extracted using Cell Lysis Buffer (Cell

Signaling Technology) according to manufacturer's specifications. Briefly, a 1X concentration of

this buffer was made with the addition of 2X protease inhibitor cocktail (Millipore Sigma), 1 mM

sodium fluoride, 1 mM sodium pyrophosphate, and 5 mM activated sodium orthovanadate.

Cells were rinsed in 1x PBS in the plate, and 500 μL of the lysis buffer was added. The plate

was incubated on ice for 5 minutes. The cells and buffer were then removed to a 1.7 mL

microcentrifuge tube and sonicated for 10 seconds. Tubes were centrifuged at 14,000 x g for

10 minutes, and supernatant was removed to a separate tube. Protein concentration was

determined using a microplate and BCA assay kit (Thermofisher) as per manufacturer's

specifications. Equal amounts of protein (nominally 50 μg) from each sample were separated

on 4-20% gradient Bis-Tris SDS-PAGE gels (GenScript), and protein was then electroblotted

onto low autofluorescence PVDF membrane (Bio-Rad). The membrane was blocked in 1X PBS

+ 5% BSA (Millipore Sigma), and probed overnight with primary rabbit antibodies raised against

total or active β-Catenin (Cell Signaling Technology), and primary mouse antibodies raised

against α-tubulin (Santa Cruz Biotechnology) as a loading control. The next day, the primary

antibodies were washed off and the blot was incubated with anti-mouse DyLight 680 conjugated

and anti-rabbit DyLight 800 conjugated secondary antibodies for an hour. The secondary

antibodies were washed off and signal was captured using the LiCor Odyssey CLx Near-

Infrared Imaging System, using the 800 channel to capture β-Catenin signal and the 700

channel to capture α-tubulin signal. Quantitation of signal was performed using ImageStudio

Lite (LiCor), using the software's "average" method of background subtraction. β-Catenin signal

was normalized using α-tubulin signal, and average normalized β-Catenin signal, standard

deviation, and statistical significance were calculated and plotted using GraphPad Prism 8.

693
694   **REFERENCES**
695

1.   MacDonald BT, Tamai K, He X. Wnt/beta-catenin signaling: components, mechanisms,

and diseases. Dev Cell. 2009;17(1):9-26. Epub 2009/07/22. doi: S1534-5807(09)00257-3 [pii]

10.1016/j.devcel.2009.06.016. PubMed PMID: 19619488.

2.   Kinzler KW, Nilbert MC, Su LK, Vogelstein B, Bryan TM, Levy DB, et al. Identification of

FAP locus genes from chromosome 5q21. Science. 1991;253(5020):661-5. Epub 1991/08/09.

doi: 10.1126/science.1651562. PubMed PMID: 1651562.

3.   Seshagiri S, Stawiski EW, Durinck S, Modrusan Z, Storm EE, Conboy CB, et al.

Recurrent R-spondin fusions in colon cancer. Nature. 2012;488(7413):660-4. doi:

10.1038/nature11282. PubMed PMID: 22895193; PubMed Central PMCID: PMC3690621.

4.   Ben-Porath I, Thomson MW, Carey VJ, Ge R, Bell GW, Regev A, et al. An embryonic

stem cell-like gene expression signature in poorly differentiated aggressive human tumors. Nat

Genet. 2008;40(5):499-507. doi: 10.1038/ng.127. PubMed PMID: 18443585; PubMed Central

PMCID: PMC2912221.

5.   Merlos-Suarez A, Barriga FM, Jung P, Iglesias M, Cespedes MV, Rossell D, et al. The

intestinal stem cell signature identifies colorectal cancer stem cells and predicts disease

relapse. Cell Stem Cell. 2011;8(5):511-24. doi: 10.1016/j.stem.2011.02.020. PubMed PMID:

21419747.

6.   Johnson CD, Esquela-Kerscher A, Stefani G, Byrom M, Kelnar K, Ovcharenko D, et al.

The let-7 microRNA represses cell proliferation pathways in human cells. Cancer Res.

2007;67(16):7713-22. Epub 2007/08/19. doi: 67/16/7713 [pii]

10.1158/0008-5472.CAN-07-1083. PubMed PMID: 17699775.

7.   Gurtan AM, Ravi A, Rahl PB, Bosson AD, JnBaptiste CK, Bhutkar A, et al. Let-7

represses Nr6a1 and a mid-gestation developmental program in adult fibroblasts. Genes Dev.

2013;27(8):941-54. doi: 10.1101/gad.215376.113. PubMed PMID: 23630078; PubMed Central

PMCID: PMC3650230.

8.   Boyerinas B, Park SM, Shomron N, Hedegaard MM, Vinther J, Andersen JS, et al.

Identification of let-7-regulated oncofetal genes. Cancer Res. 2008;68(8):2587-91. Epub

2008/04/17. doi: 68/8/2587 [pii]

10.1158/0008-5472.CAN-08-0264. PubMed PMID: 18413726.

9.   Boyerinas B, Park SM, Hau A, Murmann AE, Peter ME. The role of let-7 in cell

differentiation and cancer. Endocr Relat Cancer. 2010;17(1):F19-36. doi: 10.1677/ERC-09-

0184. PubMed PMID: 19779035.

10.   Madison BB, Jeganathan AN, Mizuno R, Winslow MM, Castells A, Cuatrecasas M, et al.

Let-7 Represses Carcinogenesis and a Stem Cell Phenotype in the Intestine via Regulation of

Hmga2. PLoS Genet. 2015;11(8):e1005408. doi: 10.1371/journal.pgen.1005408. PubMed

PMID: 26244988; PubMed Central PMCID: PMC4526516.

11.   King CE, Wang L, Winograd R, Madison BB, Mongroo PS, Johnstone CN, et al. LIN28B

fosters colon cancer migration, invasion and transformation through let-7-dependent and -

independent mechanisms. Oncogene. 2011;30(40):4185-93. Epub 2011/06/01. doi:

10.1038/onc.2011.131. PubMed PMID: 21625210; PubMed Central PMCID: PMC3165068.

12.   Madison BB, Liu Q, Zhong X, Hahn CM, Lin N, Emmett MJ, et al. LIN28B promotes

growth and tumorigenesis of the intestinal epithelium via Let-7. Genes Dev. 2013;27(20):2233-

45. doi: 10.1101/gad.224659.113. PubMed PMID: 24142874; PubMed Central PMCID:

PMC3814644.

13.   Kumar MS, Erkeland SJ, Pester RE, Chen CY, Ebert MS, Sharp PA, et al. Suppression

of non-small cell lung tumor development by the let-7 microRNA family. Proc Natl Acad Sci U S

A. 2008;105(10):3903-8. Epub 2008/03/01. doi: 0712321105 [pii]

10.1073/pnas.0712321105. PubMed PMID: 18308936.

14.   Yu F, Yao H, Zhu P, Zhang X, Pan Q, Gong C, et al. let-7 regulates self renewal and

tumorigenicity of breast cancer cells. Cell. 2007;131(6):1109-23. doi: 10.1016/j.cell.2007.10.054.

PubMed PMID: 18083101.

15.   Ricarte-Filho JC, Fuziwara CS, Yamashita AS, Rezende E, da-Silva MJ, Kimura ET.

Effects of let-7 microRNA on Cell Growth and Differentiation of Papillary Thyroid Cancer. Transl

Oncol. 2009;2(4):236-41. Epub 2009/12/04. PubMed PMID: 19956384.

16.   Nadiminty N, Tummala R, Lou W, Zhu Y, Shi XB, Zou JX, et al. MicroRNA let-7c is

downregulated in prostate cancer and suppresses prostate cancer growth. PLoS One.

2012;7(3):e32832. Epub 2012/04/06. doi: 10.1371/journal.pone.0032832. PubMed PMID:

22479342; PubMed Central PMCID: PMC3316551.

17.   Qian P, Zuo Z, Wu Z, Meng X, Li G, Wu Z, et al. Pivotal role of reduced let-7g

expression in breast cancer invasion and metastasis. Cancer Res. 2011;71(20):6463-74. doi:

10.1158/0008-5472.CAN-11-1322. PubMed PMID: 21868760.

18.   Ye S, Song W, Xu X, Zhao X, Yang L. IGF2BP2 promotes colorectal cancer cell

proliferation and survival through interfering with RAF-1 degradation by miR-195. FEBS Lett.

2016;590(11):1641-50. doi: 10.1002/1873-3468.12205. PubMed PMID: 27153315.

19.   Wang X, Liu X, Li AY, Chen L, Lai L, Lin HH, et al. Overexpression of HMGA2 promotes

metastasis and impacts survival of colorectal cancers. Clin Cancer Res. 2011;17(8):2570-80.

doi: 10.1158/1078-0432.CCR-10-2542. PubMed PMID: 21252160; PubMed Central PMCID:

PMC3079060.

20.   Strubberg AS, Veronese-Paniagua DA, Zhao TT, L. D, Pritchard T, Bayguinov P, et al.

The Zinc Finger Transcription Factor PLAGL2 Enhances Stem Cell Fate and Activates

Expression of ASCL2 in Intestinal Epithelial Cells. Stem cell reports. 2018;*In Press*.

21.   van der Flier LG, van Gijn ME, Hatzis P, Kujala P, Haegebarth A, Stange DE, et al.

Transcription factor achaete scute-like 2 controls intestinal stem cell fate. Cell. 2009;136(5):903-

12. Epub 2009/03/10. doi: S0092-8674(09)00079-8 [pii]

10.1016/j.cell.2009.01.031. PubMed PMID: 19269367.

22.   Schuijers J, Junker JP, Mokry M, Hatzis P, Koo BK, Sasselli V, et al. Ascl2 acts as an R-

spondin/Wnt-responsive switch to control stemness in intestinal crypts. Cell Stem Cell.

2015;16(2):158-70. doi: 10.1016/j.stem.2014.12.006. PubMed PMID: 25620640.

23.   Wang YP, Guo PT, Zhu Z, Zhang H, Xu Y, Chen YZ, et al. Pleomorphic adenoma gene

like-2 induces epithelial-mesenchymal transition via Wnt/beta-catenin signaling pathway in

human colorectal adenocarcinoma. Oncol Rep. 2017. doi: 10.3892/or.2017.5485. PubMed

PMID: 28259923.

24.   Zhou J, Liu H, Zhang L, Liu X, Zhang C, Wang Y, et al. DJ-1 promotes colorectal cancer

progression through activating PLAGL2/Wnt/BMP4 axis. Cell Death Dis. 2018;9(9):865. Epub

2018/08/31. doi: 10.1038/s41419-018-0883-4. PubMed PMID: 30158634; PubMed Central

PMCID: PMCPMC6115399.

25.   Liu X, Chen X, Zeng K, Xu M, He B, Pan Y, et al. DNA-methylation-mediated silencing of

miR-486-5p promotes colorectal cancer proliferation and migration through activation of

PLAGL2/IGF2/beta-catenin signal pathways. Cell Death Dis. 2018;9(10):1037. Epub

2018/10/12. doi: 10.1038/s41419-018-1105-9. PubMed PMID: 30305607; PubMed Central

PMCID: PMCPMC6180105.

26.   Li D, Lin C, Li N, Du Y, Yang C, Bai Y, et al. PLAGL2 and POFUT1 are regulated by an

evolutionarily conserved bidirectional promoter and are collaboratively involved in colorectal

cancer by maintaining stemness. EBioMedicine. 2019;45:124-38. Epub 2019/07/08. doi:

10.1016/j.ebiom.2019.06.051. PubMed PMID: 31279780; PubMed Central PMCID:

PMCPMC6642334.

27.   Li N, Li D, Du Y, Su C, Yang C, Lin C, et al. Overexpressed PLAGL2 transcriptionally

activates Wnt6 and promotes cancer development in colorectal cancer. Oncol Rep.

2019;41(2):875-84. Epub 2018/12/12. doi: 10.3892/or.2018.6914. PubMed PMID: 30535429;

PubMed Central PMCID: PMCPMC6313070.

28.   Zheng H, Ying H, Wiedemeyer R, Yan H, Quayle SN, Ivanova EV, et al. PLAGL2

regulates Wnt signaling to impede differentiation in neural stem cells and gliomas. Cancer Cell.

2010;17(5):497-509. doi: 10.1016/j.ccr.2010.03.020. PubMed PMID: 20478531; PubMed

Central PMCID: PMC2900858.

29.   Cancer Genome Atlas N. Comprehensive molecular characterization of human colon

and rectal cancer. Nature. 2012;487(7407):330-7. doi: 10.1038/nature11252. PubMed PMID:

22810696; PubMed Central PMCID: PMC3401966.

30.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer

genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer

discovery. 2012;2(5):401-4. doi: 10.1158/2159-8290.CD-12-0095. PubMed PMID: 22588877;

PubMed Central PMCID: PMC3956037.

31.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative

analysis of complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal.

2013;6(269):pl1. doi: 10.1126/scisignal.2004088. PubMed PMID: 23550210; PubMed Central

PMCID: PMC4160307.

32.   Wen Y, Liao G, Pritchard T, Zhao TT, Connelly JP, Pruett-Miller SM, et al. A Stable but

Reversible Integrated Surrogate Reporter for Assaying CRISPR/Cas9-Stimulated Homology-

directed Repair. J Biol Chem. 2017. doi: 10.1074/jbc.M117.777722. PubMed PMID: 28228480.

33.   Bajar BT, Lam AJ, Badiee RK, Oh YH, Chu J, Zhou XX, et al. Fluorescent indicators for

simultaneous reporting of all four cell cycle phases. Nat Methods. 2016;13(12):993-6. Epub

2016/11/01. doi: 10.1038/nmeth.4045. PubMed PMID: 27798610; PubMed Central PMCID:

PMCPMC5548384.

34.   Akhtar M, Holmgren C, Gondor A, Vesterlund M, Kanduri C, Larsson C, et al. Cell type

and context-specific function of PLAG1 for IGF2 P3 promoter activity. Int J Oncol.

2012;41(6):1959-66. doi: 10.3892/ijo.2012.1641. PubMed PMID: 23023303; PubMed Central

PMCID: PMC3583874.

35.   Hensen K, Van Valckenborgh IC, Kas K, Van de Ven WJ, Voz ML. The tumorigenic

diversity of the three PLAG family members is associated with different DNA binding capacities.

Cancer Res. 2002;62(5):1510-7. PubMed PMID: 11888928.

36.   Voz ML, Agten NS, Van de Ven WJ, Kas K. PLAG1, the main translocation target in

pleomorphic adenoma of the salivary glands, is a positive regulator of IGF-II. Cancer Res.

2000;60(1):106-13. PubMed PMID: 10646861.

37.   Medico E, Russo M, Picco G, Cancelliere C, Valtorta E, Corti G, et al. The molecular

landscape of colorectal cancer cell lines unveils clinically actionable kinase targets. Nat

Commun. 2015;6:7002. doi: 10.1038/ncomms8002. PubMed PMID: 25926053.

38.   Oost KC, van Voorthuijsen L, Fumagalli A, Lindeboom RGH, Sprangers J, Omerzu M, et

al. Specific Labeling of Stem Cell Activity in Human Colorectal Organoids Using an ASCL2-

Responsive Minigene. Cell Rep. 2018;22(6):1600-14. doi: 10.1016/j.celrep.2018.01.033.

PubMed PMID: 29425513; PubMed Central PMCID: PMCPMC5847189.

39.   Van Dyck F, Braem CV, Chen Z, Declercq J, Deckers R, Kim BM, et al. Loss of the

PlagL2 transcription factor affects lacteal uptake of chylomicrons. Cell metabolism.

2007;6(5):406-13. doi: 10.1016/j.cmet.2007.09.010. PubMed PMID: 17983586.

40.   Landrette SF, Kuo YH, Hensen K, Barjesteh van Waalwijk van Doorn-Khosrovani S,

Perrat PN, Van de Ven WJ, et al. Plag1 and Plagl2 are oncogenes that induce acute myeloid

leukemia in cooperation with Cbfb-MYH11. Blood. 2005;105(7):2900-7. doi: 10.1182/blood-

2004-09-3630. PubMed PMID: 15585652.

41.   Liu B, Lu C, Song YX, Gao P, Sun JX, Chen XW, et al. The role of pleomorphic

adenoma gene-like 2 in gastrointestinal cancer development, progression, and prognosis.

International journal of clinical and experimental pathology. 2014;7(6):3089-100. PubMed PMID:

25031728; PubMed Central PMCID: PMC4097215.

42.   Guo J, Wang M, Wang Z, Liu X. Overexpression of Pleomorphic Adenoma Gene-Like 2

Is a Novel Poor Prognostic Marker of Prostate Cancer. PLoS One. 2016;11(8):e0158667. doi:

10.1371/journal.pone.0158667. PubMed PMID: 27537362; PubMed Central PMCID:

PMCPMC4990332.

43.   Landrette SF, Madera D, He F, Castilla LH. The transcription factor PlagL2 activates Mpl

transcription and signaling in hematopoietic progenitor and leukemia cells. Leukemia.

2011;25(4):655-62. doi: 10.1038/leu.2010.301. PubMed PMID: 21263445; PubMed Central

PMCID: PMC3076538.

44.   de Sauvage FJ, Hass PE, Spencer SD, Malloy BE, Gurney AL, Spencer SA, et al.

Stimulation of megakaryocytopoiesis and thrombopoiesis by the c-Mpl ligand. Nature.

1994;369(6481):533-8. doi: 10.1038/369533a0. PubMed PMID: 8202154.

45.   Souyri M, Vigon I, Penciolelli JF, Heard JM, Tambourin P, Wendling F. A putative

truncated cytokine receptor gene transduced by the myeloproliferative leukemia virus

immortalizes hematopoietic progenitors. Cell. 1990;63(6):1137-47. doi: 10.1016/0092-

8674(90)90410-g. PubMed PMID: 2175677.

46.   Skoda RC, Seldin DC, Chiang MK, Peichel CL, Vogt TF, Leder P. Murine c-mpl: a

member of the hematopoietic growth factor receptor superfamily that transduces a proliferative

signal. EMBO J. 1993;12(7):2645-53. PubMed PMID: 8334987; PubMed Central PMCID:

PMCPMC413511.

47.   Vigon I, Dreyfus F, Melle J, Viguie F, Ribrag V, Cocault L, et al. Expression of the c-mpl

proto-oncogene in human hematologic malignancies. Blood. 1993;82(3):877-83. PubMed PMID:

8393355.

48.   Gao W, Chen L, Ma Z, Du Z, Zhao Z, Hu Z, et al. Isolation and phenotypic

characterization of colorectal cancer stem cells with organ-specific metastatic potential.

Gastroenterology. 2013;145(3):636-46 e5. doi: 10.1053/j.gastro.2013.05.049. PubMed PMID:

23747337.

49.   Wu Z, Wei D, Gao W, Xu Y, Hu Z, Ma Z, et al. TPO-Induced Metabolic Reprogramming

Drives Liver Metastasis of Colorectal Cancer CD110+ Tumor-Initiating Cells. Cell Stem Cell.

2015;17(1):47-59. doi: 10.1016/j.stem.2015.05.016. PubMed PMID: 26140605.

50.   Mizutani A, Furukawa T, Adachi Y, Ikehara S, Taketani S. A zinc-finger protein,

PLAGL2, induces the expression of a proapoptotic protein Nip3, leading to cellular apoptosis. J

Biol Chem. 2002;277(18):15851-8. doi: 10.1074/jbc.M111431200. PubMed PMID: 11832486.

51.   Hanks TS, Gauss KA. Pleomorphic adenoma gene-like 2 regulates expression of the

p53 family member, p73, and induces cell cycle block and apoptosis in human promonocytic

U937 cells. Apoptosis. 2012;17(3):236-47. doi: 10.1007/s10495-011-0672-3. PubMed PMID:

22076304.

52.   Guo Y, Yang MC, Weissler JC, Yang YS. PLAGL2 translocation and SP-C promoter

activity--a cellular response of lung cells to hypoxia. Biochem Biophys Res Commun.

2007;360(3):659-65. doi: 10.1016/j.bbrc.2007.06.106. PubMed PMID: 17618602; PubMed

Central PMCID: PMCPMC2084061.

53.   Giakountis A, Moulos P, Zarkou V, Oikonomou C, Harokopos V, Hatzigeorgiou AG, et al.

A Positive Regulatory Loop between a Wnt-Regulated Non-coding RNA and ASCL2 Controls

Intestinal Stem Cell Fate. Cell Rep. 2016;15(12):2588-96. doi: 10.1016/j.celrep.2016.05.038.

PubMed PMID: 27292638.

54.   Shang Y, Pan Q, Chen L, Ye J, Zhong X, Li X, et al. Achaete scute-like 2 suppresses

CDX2 expression and inhibits intestinal neoplastic epithelial cell differentiation. Oncotarget.

2015;6(31):30993-1006. doi: 10.18632/oncotarget.5206. PubMed PMID: 26307678; PubMed

Central PMCID: PMC4741583.

55.   Tian Y, Pan Q, Shang Y, Zhu R, Ye J, Liu Y, et al. MicroRNA-200 (miR-200) cluster

regulation by achaete scute-like 2 (Ascl2): impact on the epithelial-mesenchymal transition in

colon cancer cells. J Biol Chem. 2014;289(52):36101-15. doi: 10.1074/jbc.M114.598383.

PubMed PMID: 25371200; PubMed Central PMCID: PMC4276874.

56.   Ye J, Liu S, Shang Y, Chen H, Wang R. R-spondin1/Wnt-enhanced Ascl2 autoregulation

controls the self-renewal of colorectal cancer progenitor cells. Cell Cycle. 2018;17(8):1014-25.

doi: 10.1080/15384101.2018.1469874. PubMed PMID: 29886802; PubMed Central PMCID:

PMCPMC6103697.

57.   Wei X, Ye J, Shang Y, Chen H, Liu S, Liu L, et al. Ascl2 activation by YAP1/KLF5

ensures the self-renewability of colon cancer progenitor cells. Oncotarget. 2017;8(65):109301-

18. doi: 10.18632/oncotarget.22673. PubMed PMID: 29312609; PubMed Central PMCID:

PMCPMC5752522.

58.   Reed KR, Tunster SJ, Young M, Carrico A, John RM, Clarke AR. Entopic

overexpression of Ascl2 does not accelerate tumourigenesis in ApcMin mice. Gut.

2012;61(10):1435-8. doi: 10.1136/gutjnl-2011-300842. PubMed PMID: 22138533.

59.   Cui H, Cruz-Correa M, Giardiello FM, Hutcheon DF, Kafonek DR, Brandenburg S, et al.

Loss of IGF2 imprinting: a potential marker of colorectal cancer risk. Science.

2003;299(5613):1753-5. doi: 10.1126/science.1080902. PubMed PMID: 12637750.

60.   Kaneda A, Wang CJ, Cheong R, Timp W, Onyango P, Wen B, et al. Enhanced

sensitivity to IGF-II signaling links loss of imprinting of IGF2 to increased cell proliferation and

tumor risk. Proc Natl Acad Sci U S A. 2007;104(52):20926-31. doi: 10.1073/pnas.0710359105.

PubMed PMID: 18087038; PubMed Central PMCID: PMC2409243.

61.   Sakatani T, Kaneda A, Iacobuzio-Donahue CA, Carter MG, de Boom Witzel S, Okano H,

et al. Loss of imprinting of Igf2 alters intestinal maturation and tumorigenesis in mice. Science.

2005;307(5717):1976-8. Epub 2005/02/26. doi: 1108080 [pii]

10.1126/science.1108080. PubMed PMID: 15731405.

62.   Zanella ER, Galimi F, Sassi F, G. M, Cottino F, Leto SM, et al. IGF2 is an actionable

target that identifies a distinct subpopulation of colorectal cancer patients with marginal

response to anti-EGFR therapies. Sci Transl Med. 2015;7(272):272ra12. Epub 1/28/2015. doi:

10.1126.

63.   Kinouchi Y, Hiwatashi N, Higashioka S, Nagashima F, Chida M, Toyota T. Relaxation of

imprinting of the insulin-like growth factor II gene in colorectal cancer. Cancer Lett.

1996;107(1):105-8. PubMed PMID: 8913273.

64.   Nishihara S, Hayashida T, Mitsuya K, Schulz TC, Ikeguchi M, Kaibara N, et al. Multipoint

imprinting analysis in sporadic colorectal cancers with and without microsatellite instability. Int J

Oncol. 2000;17(2):317-22. PubMed PMID: 10891541.

65.   Nakagawa H, Chadwick RB, Peltomaki P, Plass C, Nakamura Y, de La Chapelle A. Loss

of imprinting of the insulin-like growth factor II gene occurs by biallelic methylation in a core

region of H19-associated CTCF-binding sites in colorectal cancer. Proc Natl Acad Sci U S A.

2001;98(2):591-6. doi: 10.1073/pnas.011528698. PubMed PMID: 11120891; PubMed Central

PMCID: PMC14632.

66.   Zhong H, Fazenbaker C, Chen C, Breen S, Huang J, Yao X, et al. Overproduction of

IGF-2 drives a subset of colorectal cancer cells, which specifically respond to an anti-IGF

therapeutic antibody and combination therapies. Oncogene. 2016. doi: 10.1038/onc.2016.248.

PubMed PMID: 27399333.

67.   Berg KCG, Eide PW, Eilertsen IA, Johannessen B, Bruun J, Danielsen SA, et al. Multi-

omics of 34 colorectal cancer cell lines - a resource for biomedical studies. Mol Cancer.

2017;16(1):116. doi: 10.1186/s12943-017-0691-y. PubMed PMID: 28683746; PubMed Central

PMCID: PMC5498998.

68.   Sanderson MP, Hofmann MH, Garin-Chesa P, Schweifer N, Wernitznig A, Fischer S, et

al. The IGF1R/INSR Inhibitor BI 885578 Selectively Inhibits Growth of IGF2-Overexpressing

Colorectal Cancer Tumors and Potentiates the Efficacy of Anti-VEGF Therapy. Mol Cancer

Ther. 2017;16(10):2223-33. doi: 10.1158/1535-7163.MCT-17-0336. PubMed PMID: 28729397.

69.   Cancer Genome Atlas Research N, Weinstein JN, Collisson EA, Mills GB, Shaw KR,

Ozenberger BA, et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nat Genet.

2013;45(10):1113-20. doi: 10.1038/ng.2764. PubMed PMID: 24071849; PubMed Central

PMCID: PMCPMC3919969.

70.   Mokry M, Hatzis P, Schuijers J, Lansu N, Ruzius FP, Clevers H, et al. Integrated

genome-wide analysis of transcription factor occupancy, RNA polymerase II binding and steady-

state RNA levels identify differentially regulated functional gene classes. Nucleic Acids Res.

2012;40(1):148-58. doi: 10.1093/nar/gkr720. PubMed PMID: 21914722; PubMed Central

PMCID: PMC3245935.

71.   Muta Y, Fujita Y, Sumiyama K, Sakurai A, Taketo MM, Chiba T, et al. Composite

regulation of ERK activity dynamics underlying tumour-specific traits in the intestine. Nat

Commun. 2018;9(1):2174. doi: 10.1038/s41467-018-04527-8. PubMed PMID: 29872037;

PubMed Central PMCID: PMCPMC5988836.

72.   Veeman MT, Slusarski DC, Kaykas A, Louie SH, Moon RT. Zebrafish prickle, a

modulator of noncanonical Wnt/Fz signaling, regulates gastrulation movements. Curr Biol.

2003;13(8):680-5. PubMed PMID: 12699626.

73.   Hoadley KA, Yau C, Hinoue T, Wolf DM, Lazar AJ, Drill E, et al. Cell-of-Origin Patterns

Dominate the Molecular Classification of 10,000 Tumors from 33 Types of Cancer. Cell.

2018;173(2):291-304 e6. doi: 10.1016/j.cell.2018.03.022. PubMed PMID: 29625048; PubMed

Central PMCID: PMCPMC5957518.

74.   Slaymaker IM, Gao L, Zetsche B, Scott DA, Yan WX, Zhang F. Rationally engineered

Cas9 nucleases with improved specificity. Science. 2016;351(6268):84-8. doi:

10.1126/science.aad5227. PubMed PMID: 26628643; PubMed Central PMCID: PMC4714946.

75.   Yusa K, Zhou L, Li MA, Bradley A, Craig NL. A hyperactive piggyBac transposase for

mammalian applications. Proceedings of the National Academy of Sciences of the United States

of America. 2011;108(4):1531-6. Epub 2011/01/06. doi: 10.1073/pnas.1008322108. PubMed

PMID: 21205896; PubMed Central PMCID: PMC3029773.

76.   Cong L, Ran FA, Cox D, Lin S, Barretto R, Habib N, et al. Multiplex genome engineering

using CRISPR/Cas systems. Science. 2013;339(6121):819-23. doi: 10.1126/science.1231143.

PubMed PMID: 23287718; PubMed Central PMCID: PMCPMC3795411.

77.   Pinello L, Canver MC, Hoban MD, Orkin SH, Kohn DB, Bauer DE, et al. Analyzing

CRISPR genome-editing experiments with CRISPResso. Nat Biotechnol. 2016;34(7):695-7.

Epub 2016/07/13. doi: 10.1038/nbt.3583. PubMed PMID: 27404874; PubMed Central PMCID:

PMCPMC5242601.

78.   Shcherbakova DM, Verkhusha VV. Near-infrared fluorescent proteins for multicolor in

vivo imaging. Nat Methods. 2013;10(8):751-4. doi: 10.1038/nmeth.2521. PubMed PMID:

23770755; PubMed Central PMCID: PMCPMC3737237.

79.   Moffat J, Grueneberg DA, Yang X, Kim SY, Kloepfer AM, Hinkle G, et al. A lentiviral

RNAi library for human and mouse genes applied to an arrayed viral high-content screen. Cell.

2006;124(6):1283-98. doi: 10.1016/j.cell.2006.01.040. PubMed PMID: 16564017.

80.   Korinek V, Barker N, Morin PJ, van Wichen D, de Weger R, Kinzler KW, et al.

Constitutive transcriptional activation by a beta-catenin-Tcf complex in APC-/- colon carcinoma.

Science. 1997;275(5307):1784-7. Epub 1997/03/21. PubMed PMID: 9065401.

81.   Kolligs FT, Hu G, Dang CV, Fearon ER. Neoplastic transformation of RK3E by mutant

beta-catenin requires deregulation of Tcf/Lef transcription but not activation of c-myc

expression. Mol Cell Biol. 1999;19(8):5696-706. PubMed PMID: 10409758; PubMed Central

PMCID: PMC84421.

A. DLD1 - Migration

B. DLD1 - Invasion

C. Caco2 - Migration

D. Parental **E.** *PLAGL2 Mut #1* **F.** *PLAGL2 Mut #2*

IEC6
Soft
Agar
Colony
Form.

G. Vector **H.** *PLAGL2 O/E*

A. *IGF2* mRNA DLD1

B. *IGF2* mRNA - Caco2

C. *PLAGL2* SRIRACCHA SW480 *IGF2* mRNA

D. *Igf2* mRNA Enteroids

E. *Igf2* mRNA Enteroids

F. *PLAGL2* O/E

Vec. Low High

PLAGL2 (anti-HA)

IGF2

1.00 2.43 2.39

TUB

G. *IGF2* mRNA Enteroids

H. Empty Vector

I. IGF2 O/E