GIANI: open-source software for automated analysis of 3D microscopy images

David J. Barry[1*], Claudia Gerri[2], Donald M. Bell[1], Rocco D'Antuono[1], Kathy K. Niakan[2,3]

1 Crick Advanced Light Microscopy, Francis Crick Institute, London, NW1 1ST, UK

2 Human Embryo and Stem Cell Laboratory, Francis Crick Institute, London, NW1 1ST, UK

3 The Centre for Trophoblast Research, Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge, CB2 3EG, UK

* david.barry@crick.ac.uk

Abstract

The study of cellular and developmental processes in physiologically relevant three-dimensional (3D) systems facilitates an understanding of mechanisms underlying cell fate, disease and injury. While cutting-edge microscopy technologies permit the routine acquisition of 3D datasets, there is currently a limited number of open-source software packages to analyse such images. Here we describe GIANI (djpbarry.github.io/Giani), new software for the analysis of 3D images, implemented as a plugin for the popular FIJI platform. The design primarily facilitates segmentation of nuclei and cells, followed by quantification of morphology and protein expression. GIANI enables routine and reproducible batch-processing of large numbers of images and also comes with scripting and command line tools, allowing users to incorporate its functionality into their own scripts and also run GIANI on a high-performance computing cluster. We demonstrate the utility of GIANI by quantifying cell morphology and protein expression in confocal images of mouse early embryos and by segmenting nuclei from light sheet microscopy images of the flour beetle embryo. We also validate the performance of the software using simulated data. More generally, we anticipate that GIANI will be a useful tool for researchers in a variety of biomedical fields.

Introduction

The ability to routinely acquire multi-dimensional datasets with modern microscopy techniques is transforming, among other fields, cell biology, developmental biology and cancer research. There has long been an acceptance that two-dimensional (2D) cell cultures may not accurately recreate behaviours

found in complex, three-dimensional (3D) *in vivo* environments [1]. Commonly-used 3D culture formats include, but are not limited to, populations of single cells in organotypic matrices, spheroid models, tissue sections or whole embryos and organisms.

However, the development of software for the quantitative analysis of such data has not kept pace with imaging advances and there is now a pressing need for automated solutions [2]. Manual annotation of such data is not feasible, due to the time required to do so. Commercial packages, such as Imaris (Bitplane) and Vison4D (Arivis) provide excellent visualisation functionality and are also equipped with analysis tools. However, licenses for such software are expensive and they also rely on proprietary file formats. Furthermore, the closed-source nature of such software prevents detailed interrogation of specific calculations and processes.

There are numerous excellent, freely-available bioimage analysis tools in the open source domain. However, their support for 3D analysis is either limited (for example, several CellProfiler [3] modules are not yet compatible with 3D images) or challenging to execute and/or automate for the uninitiated (FIJI [4], Icy [5]). This can lead to undesirable compromises being made, such as 2D slices from a 3D volume being analysed individually, blinding the analysis to information in adjacent slices. Alternatively, 3D data may be compressed into 2D via projection, which, consequently, artificially reduces distances between objects and can lead to spurious results. A number of very useful open-source MATLAB (MathWorks, Cambridge, UK) -based tools have also been implemented, most notably MINS, which has been effectively used to analyse embryo datasets [6] and LOBSTER [7]. However, these require the purchase of a MATLAB license.

We have therefore developed GIANI (**G**eneral **I**mage **A**nalysis of $\qquad$ 32
Nuclei-based **I**mages; https://djpbarry.github.io/Giani), a generally-applicable, $\qquad$ 33
open source tool, implemented as a plugin for the widely-used image analysis $\qquad$ 34
platform, FIJI. With an emphasis on detection and segmentation of cells in 3D $\qquad$ 35
microscopy images, GIANI has been implemented specifically with $\qquad$ 36
batch-processing in mind. While an understanding of fundamental concepts of $\qquad$ 37
bioimage analysis is beneficial, GIANI's user interface has been implemented in $\qquad$ 38
a wizard format to facilitate use by non-specialists and is fully documented $\qquad$ 39
(https://github.com/djpbarry/Giani/wiki). Analysis protocols may be reproduced $\qquad$ 40
by loading a single, small parameter file. $\qquad$ 41

The utility of GIANI is illustrated here using three examples. In the first, we $\qquad$ 42
generated a series of simulated datasets to evaluate the accuracy of $\qquad$ 43
segmentations produced by GIANI using known "ground truths". In the second $\qquad$ 44
proof-of-concept we use a series of mouse preimplantation embryo datasets and $\qquad$ 45
demonstrate the ability of GIANI to detect variations in morphology and protein $\qquad$ 46
expression in different experimental conditions. Finally, to show that GIANI can $\qquad$ 47
also be used on much larger datasets, we present segmentations of nuclei from $\qquad$ 48
light sheet microscopy images of the flour beetle embryo. It should be noted that $\qquad$ 49
GIANI can be used to quantify 3D images from a range of cellular and $\qquad$ 50
developmental contexts and we anticipate that it will be a useful tool to $\qquad$ 51
automate quantification of a wide variety of complex imaging data. $\qquad$ 52

Materials and methods

Simulated data generation

Simulated data sets were generated using an extension of a previously described method [8]. Simulated nuclei positions were generated as previously described [8], then cell membranes were approximated using a Euclidean Distance Map constructed around nuclei. The simulated images were then convolved with a Gaussian point-spread function, sub-sampled and noise added from a gamma distribution. The complete code for simulated image generation is available on GitHub (https://github.com/djpbarry/Embryo-Generator).

Metrics for segmentation quality assessment

We calculated the cell count error E_c, as

$$E_c = |N_g - N_{gt}| \tag{1}$$

where N_{gt} is the actual number of cells present in a simulated embryo and N_g is the number counted by GIANI. Nuclear (E_{nl}) and cell (E_{cl}) centroid localisation errors were calculated based on the euclidean distance between the known ground truth centroids and the centroids of the segmentations produced by GIANI.

Mouse zygote collection

Four- to eight-week-old (C57BL6 $\times$ CBA) F1 female mice were super-ovulated using injection of 5 IU of pregnant mare serum gonadotrophin (PMSG;

Sigma-Aldrich). Forty-eight hours after PMSG injection, 5 IU of human   72

chorionic gonadotrophin (HCG; Sigma-Aldrich) was administered.   73

Superovulated females were set up for mating with eight-week-old or older   74

(C57BL6 × CBA) F1 males. Mice were maintained on a 12 h light–dark cycle.   75

Mouse zygotes were isolated in EmbryoMax FHM mouse embryo media   76

(Sigma-Aldrich; MR-122-D) under mineral oil (Origio; ART-4008-5P) and   77

cumulus cells were removed with hyaluronidase (Sigma-Aldrich; H4272). All   78

animal research was performed in compliance with the UK Home Office   79

Licence Number 70/8560.   80

Mouse embryo culture   81

Mouse embryos were cultured in drops of pre-equilibrated Global medium   82

(LifeGlobal; LGGG-20) supplemented with 5 mg/ml protein supplement   83

(LifeGlobal; LGPS-605) and overlaid with mineral oil (Origio; ART-4008-5P).   84

Preimplantation embryos were incubated at 37°C and 5.5% CO_2 and cultured up   85

to the day of analysis.   86

Inhibitor treatment   87

Inhibitor experiment was performed as previously described [9]. Briefly, the   88

aPKC inhibitor CRT0276121 (Cancer Research Technology LTD) was dissolved   89

in DMSO to 10 mM stock concentration and diluted to the optimal   90

concentration of 8 μM in pre-equilibrated embryo culture media. Mouse   91

embryos were incubated in pre-equilibrated media with 8 μM of CRT0276121   92

from 4-cell to morula stage. Control mouse embryos were developed in   93

pre-equilibrated media where volume-matched DMSO was added.   94

Immunofluorescence

Embryos were fixed with freshly prepared 4% paraformaldehyde in PBS that was pre-chilled at 4°C. Embryo fixation was performed for 20 min at RT and then the embryos were transferred through 3 washes of 1X PBS with 0.1% Tween-20 to remove residual paraformaldehyde. Embryos were permeabilized with 1X PBS with 0.5% Triton X-100 and then blocked in blocking solution (3% BSA in 1X PBS with 0.2% Triton X-100) for 2 h at RT on a rotating shaker. Then, embryos were incubated with primary antibodies (listed in Table 1) diluted in blocking solution overnight at 4°C on rotating shaker.

Table 1. Primary antibodies used in this study and the dilution at which they were used.

Antibody	Supplier	Catalogue Number	Dilution
YAP1	Abnova	H00010413-M01	1:50
GATA3	R&D	AF2605	1:200
E-CADHERIN	Life Technologies	131900	1:400

The following day, embryos were washed in 1X PBS with 0.2% Triton X-100 for 20 min at RT on rotating shaker and then incubated with secondary antibodies diluted in blocking solution for 1 h at RT on rotating shaker in the dark. Next, embryos were washed in 1X PBS with 0.2% Triton X-100 for 20 min at RT on rotating shaker. Finally, embryos were placed in 1X PBS with 0.1% Tween-20 with Vectashield with DAPI mounting medium (Vector Lab; H-1200) (1:30 dilution). Embryos were placed on μ-Slide 8 well dishes (Ibidi; 80826) for confocal imaging.

Image Acquisition

All images were acquired on a Leica SP5 laser scanning confocal microscope using a Leica 1.3 NA 63x HCX PL APO CS glycerol objective and a voxel size

of approximately 0.1 x 0.1 x 1.0 μm in x, y and z, respectively. ₁₁₅

Software ₁₁₆

GIANI was written using Java 8 as a plug-in for FIJI [4], making extensive use ₁₁₇
of the underlying FIJI and ImageJ [10] libraries. A number of other open-source ₁₁₈
projects are leveraged. Reading of image data is facilitated by interfacing to ₁₁₉
Bio-Formats [11], making GIANI compatible with a wide range of file formats ₁₂₀
(https://docs.openmicroscopy.org/bio-formats/6.7.0/supported-formats.html). ₁₂₁
Detection of nuclear blobs makes use of either TrackMate's spot detector [12] or ₁₂₂
FeatureJ (https://imagescience.org/meijering/software/featurej). Segmentation ₁₂₃
of cells and nuclei takes advantage of the marker-controlled watershed ₁₂₄
functionality in MorphoLibJ [13] and 3D Image Suite [14]. The browsing of ₁₂₅
results is based upon the 3D Region of Interest (ROI) Manager from 3D Image ₁₂₆
Suite. Complete source code, documentation and test data are available online ₁₂₇
(https://djpbarry.github.io/Giani). ₁₂₈

Statistical Analysis ₁₂₉

All statistical analyses in this study were performed using RStudio ₁₃₀
(www.rstudio.com). The tests performed in this paper are two-sample Wilcoxon ₁₃₁
tests (also know as Mann-Whitney). All box plots show the median and ₁₃₂
inter-quartile range, with the whiskers extending 1.5 times the inter-quartile ₁₃₃
range from the 25th and 75th percentiles. Unless otherwise stated, each dot in ₁₃₄
dot plots represents a single cell. ₁₃₅

Results & Discussion

Software implementation

The design philosophy behind GIANI is inspired by CellProfiler. It is assumed that the user wishes to detect "primary objects" of some sort (typically cell nuclei), followed by the subsequent segmentation of "secondary objects" (typically cells) and then wishes to measure either the morphology of said objects, or the expression of a fluorescent signal within these objects. The principal difference in the case of GIANI is that, in order to facilitate ease of use, the order of steps in the pipeline is fixed, although some flexibility is present where necessary (Fig 1).

Fig 1. Overview of the algorithms underpinning GIANI. All images represent a single slice of a 3D stack and are for illustrative purposes only. A: An example dataset, consisting of a mouse embryo showing DAPI (blue) and E-Cadherin (red). GIANI can accept as input any image data that is readable by Bio-Formats [11]. B: Nuclear are first approximated using one of two blob detectors - Laplacian of Gaussian (left) or Hessian (right). C: Gaussian filtering is used to suppress noise in the channels used for nuclear and cell segmentation. D: The nuclear channel is then subjected to top-hat filtering to remove background - contrast has been increased to illustrate the effect of the filter. E: Nuclear segmentation is achieved using a marker-controlled watershed approach, with the background-subtracted image from D serving as the input and centroids from B serving as the seeds. F: Cell segmentation is achieved using the same marker-controlled watershed approach, with the filtered image from C serving as the input and nuclei segmentations from E serving as the seeds.

File Formats

GIANI can accept as input any images that are readable by Bio-Formats [11], which, at the time of writing, can read almost 160 different formats. It is expected that this data will be three-dimensional in nature, containing any number of different channels. As a minimum requirement, in order to facilitate

accurate segmentation, these channels must contain some form of marker for both nuclei and cells. The nuclear marker must be a volume marker, while the cell marker can be either membrane-localised or volumetric in nature, although the former is preferred for more accurate segmentation. In the example pipeline illustrated in Fig 1, a nuclear volume marker (DAPI) and cell membrane marker (E-Cadherin) are used.

Detection of Nuclear Centres

The nuclear channel is first subjected to one of two forms of blob detection to estimate the centroid location of each nucleus (Fig 1B). The first option is Laplacian of Gaussian (LoG) blob detection, which involves the application of an LoG filter to the image and then identifying local extrema (Fig 1B, left panels). This is implemented in GIANI by using the spot detector of TrackMate [12], which is based on LoG detection. The advantage of this approach is that it is relatively fast and generic. The disadvantage is it assumes the objects to be detected are approximately Gaussian in nature, so will likely perform poorly for irregularly shaped nuclei.

For this reason, a second option is given for nuclear centroid detection, based on the eigenvalues of the image's Hessian matrix (Fig 1B, right panels) - this is similar to the approach used for nuclear detection in MINS [6]. The Hessian is a square matrix of second-order partial derivatives describing the local curvature in the image. For a detailed explanation on how Hessian eigenvalues can be used to detect blobs in images, see [15]. This is implemented in GIANI by using FeatureJ, a sub-component of the ImageScience plugin (https://imagescience.org/meijering/software/featurej). This approach has the advantage of being capable of detecting "blobs" of any shape, such as irregularly

shaped nuclei. The main disadvantage is that it is much more computationally 176
demanding than the simpler LoG detector, so will take longer to run. 177

Suppression of Noise and Background 178

Prior to full segmentation of nuclei, Gaussian filtering is employed to smooth 179
any noise that may have been present in the input image (Fig 1C). To both 180
enhance the nuclei and homogenise the image background, the output of the 181
Gaussian filtering operation is input into a top-hat filter (Fig 1D), implemented 182
using MorphoLibJ [13]. Top-hat filtering can be computationally expensive for 183
large datasets, so GIANI includes an option to downsize datasets by a specified 184
factor prior to this step - the dataset is then restored to its original size 185
post-filtering. 186

Segmentation of Nuclei 187

Nuclei are fully segmented using a marker-controlled watershed approach 188
(Fig 1E). GIANI uses MorphoLibJ to achieve this, with the previously-detected 189
nuclear centroids serving as seeds and the Euclidean Distance Transform of the 190
centroid mask, calculated using the 3D ImageJ Suite [14], serving as the image 191
to be flooded. In addition, the top-hat filtered nuclei channel is thresholded 192
(using one of FIJI's in-built thresholding algorithms, specified by the user) to act 193
as a mask, restricting the overall extent of the segmentation. 194

Segmentation of Cells 195

Cells are segmented using the same marker-controlled watershed approach 196
(Fig 1F), with the previously-segmented nuclei serving as seeds and the distance 197
transform of the nuclei serving as the image to be flooded. The 198

Gaussian-filtered cell channel (Fig 1C) is thresholded to act as a mask in this case. The nature of the distance transform used is dependent on whether a membrane marker or volume marker is used to identify the cells. In the case of a volume marker, a standard Euclidean Distance Transform is used. In the case of a membrane marker, a modified distance transform is used, similar to that described by [16], where the distance between adjacent voxels, d, is calculated based on a Riemannian metric defined in terms of the image I and a regularisation parameter, λ:

$$d = \frac{\nabla I + \lambda}{1 + \lambda} \tag{2}$$

It can be seen that, as $lim_{\lambda \to \infty}$, d tends towards the Euclidean distance.

Analysis of simulated embryos

To first validate the performance of GIANI, we generated a series of 189 "simulated embryos" exhibiting different levels of signal-to-noise ratio and cell density (Fig 2A). Simulated data has the significant advantage of having a known "ground truth" (Fig 2B). That is, because the images are generated artificially, we know what the "correct" segmentation should look like. This permits us to compare the segmentation results generated with any piece of analysis software with the known "true" values.

The number of false positives and false negatives produced by GIANI is consistently extremely low, although cell count errors do generally increase with cell density and decreasing signal-to-noise ratio (Fig 2C, F). At high signal-to-noise ratios and low cell densities, GIANI localises nuclei to within

Fig 2. Validation of GIANI using simulated embryo data. In each of the heat maps, a single tile represents the average of three simulated embryos. The GIANI settings used to produce this data are provided as File S3. A: A 2D slice of an exemplar 3D simulated embryo. B: The ground truth segmentation of 'A'. C - H: Absolute errors in cell counts (E_c, calculated according to Eq 1), nuclear centroid localisation error (E_{nl}) and cell centroid localisation error (E_{cl}) produced by GIANI for simulated embryos with the indicated number of cells and signal-to-noise ratios (SNR). Results were obtained using either GIANI's basic (C - E) or advanced (F - H) nuclear detector.

approximately 600 nm of their true positions (Fig 2D, G). Even at high cell $\qquad$ 220

densities and low signal-to-noise ratios, this error never exceeded 1 μm (the $\qquad$ 221

simulated nuclei are ellipsoidal with axis dimensions of approximately 10.0 x $\qquad$ 222

7.5 x 7.5 μm). In addition to nuclear localisation error, we also quantified the $\qquad$ 223

cell centroid localisation error on a per cell basis (Fig 2E, H). $\qquad$ 224

To benchmark GIANI against other available software, we analysed the same $\qquad$ 225

simulated embryos using Imaris (Fig S1). This demonstrated that the $\qquad$ 226

performance of GIANI is comparable to Imaris when it comes to nuclear $\qquad$ 227

localisation (Fig S1 B, D) and that GIANI is superior in detecting the correct $\qquad$ 228

number of cells (Fig S1 A, C). We also performed a limited evaluation of $\qquad$ 229

CellProfiler on a subset of the same simulated data, which resulted in errors far $\qquad$ 230

larger than those produced by GIANI (Fig S2). $\qquad$ 231

The accurate quantitative analysis of a multi-cellular image dataset, such as $\qquad$ 232

an embryo, is dependent on the correct identification and segmentation of nuclei. $\qquad$ 233

While GIANI presently employs generic blob detectors for this purpose, which $\qquad$ 234

is similar to other previously-described methods [6], other approaches could be $\qquad$ 235

incorporated [17–21]. In particular, the ability to load pre-trained neural $\qquad$ 236

networks, by leveraging DeepImageJ [22], for example, or machine-learning $\qquad$ 237

classifiers trained with, for example, Weka [23] or Ilastik [24], may be added in $\qquad$ 238

a future release. $\qquad$ 239

In addition to the convolution of the ground truth data, the simulation process $_{240}$ also takes into consideration depth of slices within a sample when modelling $_{241}$ fluorescent intensity - deeper slices will be more prone to scattering effects and $_{242}$ therefor exhibit lower intensity. This means that finding a single global $_{243}$ thresholding strategy that does not under-segment the dimmer cells while also $_{244}$ not over-segmenting the brighter cells is challenging. Taking all this into $_{245}$ consideration, any metric used to assess the accuracy of cell segmentations will $_{246}$ have its limitations. $_{247}$

There are strategies that could be employed to mitigate against these factors. $_{248}$ For example, some form of adaptive thresholding, whereby the intensity $_{249}$ threshold changes according to z-location, could be used. However, while an $_{250}$ optimal adaptive thresholding strategy could be found for a given dataset (such $_{251}$ as the simulated data used in this study), implementing a universally-applicable $_{252}$ strategy would be difficult. One of the principle design aims of GIANI was $_{253}$ simplicity of use, which does not allow for the incorporation of a variety of $_{254}$ case-specific segmentation approaches. $_{255}$

Analysis of mouse embryos $_{256}$

We subsequently applied GIANI to the analysis of two populations of mouse $_{257}$ preimplantation embryos at the morula stage, one control ($n = 18$) and one $_{258}$ treated with a small molecule inhibitor [9] ($n = 20$). At this stage, two distinct $_{259}$ cell populations are discernible: inner and outer cells. In subsequent cell $_{260}$ divisions, a blastocyst is formed, whereby the inner cells give rise to an inner $_{261}$ cell mass (ICM), and the outer cells become the trophectoderm (TE), a polarized $_{262}$ epithelium that will form fetal components of the placenta [25]. $_{263}$

At the morula stage, inner and outer cells display different polarisation states, $_{264}$

which influence their cell fate acquisition. The outer cells acquire an apical ₂₆₅ domain, enriched with the atypical protein kinase C (aPKC) [26]. In the polar ₂₆₆ outer cells, aPKC prevents activation of downstream Hippo pathway kinases, ₂₆₇ Large tumor suppressor kinases 1/2 (LATS1/2) [27]. Consequently, in outer ₂₆₈ cells, YAP1 accumulates in the nucleus, where it promotes the expression of ₂₆₉ GATA3 [28]. In contrast, in the apolar inner cells, the activation of the Hippo ₂₇₀ pathway results in YAP1 cytoplasmic retention, thus maintaining the inner cells ₂₇₁ in an unspecified state [27, 29, 30]. ₂₇₂

We therefore divided cells within each embryo into an 'inner' and 'outer' ₂₇₃ population (Fig 3A, B). This division is based on the distance of the detected ₂₇₄ nuclear centroid from the embryo centroid. A cell was classified as 'inner' if the ₂₇₅ following condition held true: ₂₇₆

$$\frac{D_i}{D_m} < D_T \tag{3}$$

where D_i is the Euclidean distance of cell i from the embryo centroid, D_m is the ₂₇₇ maximum Euclidean distance from any cell in the same embryo to the embryo ₂₇₈ centroid and D_T is an arbitrarily set distance threshold. For the purposes of this ₂₇₉ study, we chose a value of 0.5 for D_T - it was found that modifying this value ₂₈₀ slightly ($\pm 10\%$) did not significantly impact the results (data not shown). The ₂₈₁ embryo centroid was calculated as the average of all nuclear centroids. ₂₈₂

GIANI shows differences in cell morphology, YAP1 and GATA3 ₂₈₃
expression within inner and outer cells in control embryos ₂₈₄

We then applied GIANI to detect variations in morphology and protein ₂₈₅ expression in mouse preimplantation embryos. We began with a morphological ₂₈₆

Fig 3. GIANI reveals differences in morphology and protein expression in mouse embryos. Unless otherwise stated, each dot represents a single cell. The GIANI settings used to produce this data are provided as File S4. A. Illustration of the division of embryo cells into 'outer' (red) and 'inner' (green) sub-populations. The embryo centroid is indicated by the white square. The blue circle has radius D_m (see Eq 3) and indicates the distance from the embryo centroid to the most distant nucleus centroid. The radius of the yellow circle is $D_T \times D_m$ (Eq 3). B. The number of cells in each embryo divided into outer and inner sub-populations using a value of 0.5 for D_T in control embryos. Each dot represents a single embryo ($n_{control} = 18$). C: Volume of nuclei in inner/outer populations in control embryos. D: Volume of cells in inner/outer populations in control embryos. E: Ratio of cell-to-nuclear volume in inner and outer cells in control embryos. F: Nuclear/cytoplasmic ratio of YAP1 expression in control embryos. G: Difference in expression profiles of nuclear GATA3 expression, normalised to DAPI, in control and treated embryos ($n_{treated} = 20$). H: Difference in nuclear/cytoplasmic ratiometric expression profiles of YAP1 in control and treated embryos.

analysis of cells in control embryos. No significant difference was found   287

between median nuclear volumes in inner and outer cell populations   288

(Fig 3C; $p = 0.861$, Wilcoxon rank sum test). A difference in median cell   289

volume between the two populations was also identified, although it fell slightly   290

short of being statistically significant (Fig 3D; $p = 0.069$). However, a   291

comparison of cell-to-nucleus volume ratios revealed a significant difference   292

between median values, with outer cells having a proportionately greater   293

cytoplasmic volume than inner cells (Fig 3E; $p < 0.001$). We also confirmed   294

previous analysis of YAP1 translocation [31], illustrating that nuclear   295

localisation is significantly higher in outer versus inner cells (Fig 3F;   296

$p < 0.001$). Similarly, GATA3 expression was shown to be higher in outer   297

compared to inner cells (Fig 3G; $p < 0.0001$).   298

GIANI reveals differences in YAP1 and GATA3 expression in mouse embryos after pharmacological treatment

To further demonstrate the utility of GIANI, we sought to analyse GATA3 and YAP1 expression after treating mouse embryos with a small molecular inhibitor against aPKC, the upstream regulator of YAP1 and GATA3. The aPKC inhibitor, CRT0276121, has previously been confirmed to specifically inhibit aPKC in various biological and cellular contexts [32–34]. Specifically, in mouse pre-implantation embryos, aPKC inhibition has been recently shown to efficiently abrogate YAP1 and GATA3 expression in outer cells [9].

Expression of GATA3 in the nucleus (normalised to DAPI to correct for diminished signal intensity with increasing sample depth) was found to be similarly low in the inner cells of both control and treated embryos (Fig 3G; $p = 0.841$). However, GATA3 expression in outer cells was found to be significantly lower in the treated embryos relative to the control group ($p < 0.002$). Normalised expression of GATA3 was also found to be significantly higher in outer cells of control embryos relative to their inner cells ($p < 0.001$), while the same was not true of treated embryos ($p = 0.171$). In addition, differences in distribution are evident between control and treated embryos, with two distinct populations evident in both inner and outer cells in control populations (Fig 3G).

While no statistically significant difference in nuclear YAP1 expression (normalised to cytoplasmic expression) between control and treated cells were observed when comparing inner cells (Fig 3H; $p = 0.609$), a large difference was observed in outer cells ($p < 0.0001$). However, nuclear/cytoplasmic YAP1 expression was still higher in the outer cells of treated embryos versus inner ($p < 0.0001$). Altogether, this demonstrates that GIANI allows for the

automated quantification of expression differences between cells following   325

perturbation.   326

Analysis of blastocysts   327

Finally, to illustrate that GIANI can be run successfully on later stage mouse   328

embryos, we analysed two examples of blastocysts and compared with Imaris.   329

While the resultant segmentations are less than perfect, a qualitative assessment   330

(in the absence of ground truth segmentations) shows that the results produced   331

by GIANI are at least comparable to those obtained using Imaris (Fig S3).   332

Analysis of larger datasets   333

Analysis of large simulated dataset   334

To test GIANI's ability to handle more complex datasets, we generated a large   335

simulated volume consisting of approximately 2,000 nuclei (File S1). We   336

compared the segmentations generated by GIANI with those generated by   337

Imaris (Fig 4). A qualitative assessment shows that while both software   338

successfully segment the vast majority of cells, errors are apparent, particularly   339

in cases were nuclei are highly clustered (Fig 4A). We attempted to quantify   340

these success rates using a number of different metrics.   341

Fig 4. Demonstration of GIANI with a large simulated dataset. The GIANI
and Imaris settings used to produce this data are provided as File S5.
A: Illustration of the segmentations produced by GIANI and Imaris on a large
simulated dataset (File S1). The top row shows a single slice of each 3D volume,
while the bottom row shows the magnified views of the boxes in the top row
images. Scale bars are all equivalent to 20 μm. B: Distribution of localisation
errors produced by both Imaris and GIANI in detecting the simulated nuclei in
File S1. C: Relationship between localisation error for detected nuclei and the
distance of each nucleus to its nearest neighbour. D: Influence of distance of
nuclei to their nearest neighbour on successful detection.

We first compared the distribution of localisation errors produced by GIANI ₃₄₂
and Imaris (Fig 4B). While in both cases nuclei were localised to within ₃₄₃
approximately 1 μm or less of their true locations most of the time, the ₃₄₄
localisation errors produced by Imaris were, overall, marginally lower. But, ₃₄₅
Imaris had a greater tendency to produce outliers (localisation errors above ₃₄₆
approximately 2μm). ₃₄₇

We next looked for any link between localisation error and nuclear density. ₃₄₈
To do so, we examined the relationship between the localisation error for each ₃₄₉
successfully detected nucleus and the distance between that nucleus and its ₃₅₀
nearest neighbour (calculated as the distance between nuclear centroids; Fig 4C). ₃₅₁
While the relationship between these two variables is highly non-linear, it is ₃₅₂
apparent that, for both GIANI and Imaris, a trend exists - localisation error ₃₅₃
increases as the distance between nuclei decreases. ₃₅₄

Finally, we compared the success rates of GIANI and Imaris in detecting the ₃₅₅
simulated nuclei and investigated whether nuclear density influenced the ₃₅₆
likelihood of successful nuclear detection (Fig 4D). We found that while both ₃₅₇
GIANI and Imaris successfully detected the vast majority of nuclei, the success ₃₅₈
rate was slightly higher for GIANI (93.3% versus 89.0%). There was also a ₃₅₉
clear relationship between inter-nuclear distance and likelihood of successful ₃₆₀
detection. Of the 85 nuclei that both Imaris and GIANI failed to detect, 95% ₃₆₁
were 10 μm or less from their nearest neighbour. Given that the longest ₃₆₂
dimension of the simulated nuclei is approximately 10 μm, this is perhaps not an ₃₆₃
unreasonable result - two such nuclei that are less than 10 μm apart at their ₃₆₄
centres are likely to significantly overlap. ₃₆₅

Analysis of a dataset derived from light sheet microscopy

To further emphasise the capacity of GIANI to analyse large volumes of data, 367
we tested it on a *Tribolium castaneum* embryo dataset derived from light sheet 368
microscopy (File S2), again benchmarking against Imaris (Fig 5). We obtained 369
this data from the Cell Tracking Challenge 370
(http://celltrackingchallenge.net/3d-datasets) and, unfortunately, the ground truth 371
data available is very limited (annotations for only three cells is provided). 372
However, the intention here is to demonstrate that GIANI is capable of 373
segmenting large numbers of nuclei in such data (approximately 5,600 were 374
detected) and the results are comparable to those obtained using market-leading 375
commercial software (Imaris). 376

Fig 5. Demonstration of GIANI on a large light sheet microscopy dataset.
Illustrations of the segmentations produced by GIANI and Imaris on a *Tribolium castaneum* embryo dataset derived from light sheet microscopy (File S2). Two different slices of the 3D volumes are shown, at approximately 46 μm (A) and 247 μm (B), to illustrate the variation in nuclei morphologies at different depths. In each case, the top row shows the relevant slice of each 3D volume, while the bottom row shows the magnified views of the boxes in the top row images. Approximately 5,600 nuclei were detected in the full volume. Scale bars are all equivalent to 20 μm. The GIANI and Imaris settings used to produce this data are provided as File S6.

Future developments 377

Given the ongoing interest in studying cells in "native" 3D extracellular 378
environments [35], future extension to the capabilities of GIANI will include the 379
ability to analyse time-lapse data, as there is currently a lack of open source 380
tools for the quantification of 3D cell migration [36]. The incorporation of 381
additional functionality from TrackMate (and MaMuT [37] or Mastodon) will 382
be explored to facilitate 3D cell tracking. 383

Future development will also include the replacement of the 3D ROI ⁠384
Manager interface for visualizing results with a new, custom-built interface. At ⁠385
present, segmented objects are saved solely as FIJI ROI files, but it is intended ⁠386
that support for other formats (such as .stl, .ply, .obj and .x3d) will be added to ⁠387
allow the import of objects into a variety of different software. ⁠388

More generally, with a view to improving and optimising performance, ⁠389
further use of ImgLib2 [38] will be incorporated in future releases. Taking ⁠390
advantage of GPU acceleration is also an aim, most likely by exploiting ⁠391
CLIJ [39]. At present, the analysis of a single simulated embryo used in this ⁠392
study (approx. 2.1 GB) takes between approximately 30 and 60 minutes, ⁠393
depending on the number of CPUs available (Fig S4). ⁠394

Conclusion ⁠395

We have used GIANI to quantitatively analyse mouse embryos in 3D. This ⁠396
analysis has revealed differences in morphology and protein expression between ⁠397
different experimental conditions. Analysis of simulated ground truth data was ⁠398
used to confirm the validity of these results. Further development of GIANI is ⁠399
planned, with the specific aim of improving segmentations in noisy and dense ⁠400
fields of view, common in 3D images of cells. Extension to timelapse analysis is ⁠401
also planned. GIANI is freely available on GitHub (github.com/djpbarry/Giani) ⁠402
and we anticipate that it will be a useful resource for the community to perform ⁠403
routine, automated quantification of complex imaging data. ⁠404

Supporting information

Fig S1   Accuracy of simulated nuclei detection and localisation by GIANI and Imaris. In each of the heat maps, a single tile represents the average of three simulated embryos. The GIANI and Imaris settings used to produce this data are provided as File S7. A: Absolute errors in cell counts (E_c, calculated according to Eq 1) produced by GIANI for simulated embryos with the indicated number of cells and signal-to-noise ratios (SNR). B: Absolute errors in nuclear centroid localisation (E_{nl}) produced by GIANI. C: Absolute errors in cell counts produced by Imaris. D: Absolute errors in nuclear centroid localisation produced by Imaris.

Fig S2   Accuracy of simulated nuclei and cell detection and localisation by GIANI and CellProfiler compared. The data shown in A - C is a subset of that shown in Fig 2F - H. In D - F, each tile represents a single simulated embryo, which was reduced in size to 512 x 512 x 112 voxels prior to running the pipeline. A: Absolute errors in cell counts (E_c, calculated according to Eq 1) produced by GIANI for simulated embryos with the indicated number of cells and signal-to-noise ratios (SNR). B: Absolute errors in nuclear centroid localisation (E_{nl}) produced by GIANI. C: Absolute errors in cell centroid localisation error (E_{cl}) produced by GIANI. D: Absolute errors in cell counts produced by CellProfiler. E: Absolute errors in nuclear centroid localisation produced by CellProfiler. F: Absolute errors in cell centroid localisation error produced by CellProfiler. The CellProfiler pipeline used and raw data are provided in File S8.

Fig S3   Comparison of mouse blastocyst nuclear segmentations between 428
GIANI and Imaris.. A single slice of (A) expanded and (B) hatching 429
blastocysts are shown, together with the segmentations produced by Imaris and 430
GIANI. Scale bars are all equivalent to 20 μm. The GIANI and Imaris settings 431
used to produce this data, together with the raw image data, are provided as File 432
S9. 433

Fig S4   Execution time of GIANI is independent of cell number, but 434
decreases with increasing CPU availability. A: The length of time taken by 435
GIANI to analyse simulated embryos versus the number of cells in the embryo. 436
Each point represents the execution time for a single embryo - approximately 20 437
embryos were analysed for each cell number. The blue line represents a moving 438
average calculated with LOESS smoothing. The grey bands represent the 95% 439
confidence interval. B: The length of time taken by GIANI to analyse simulated 440
embryos consisting of 30 cells versus the number of available CPUs. Each point 441
represents the execution time for a single embryo - approximately 20 embryos 442
were analysed for each CPU number. The blue line represents a moving average 443
calculated with LOESS smoothing. The grey bands represent the 95% 444
confidence interval. 445

File S1   Large simulated image of nuclei. There are approximately 2000 446
nuclei in total, each being approximately 10 μm long and 7.5 μm wide. The 447
stack dimensions are 1500 x 1500 x 300 voxels, with each voxel being 0.4 x 0.4 448
x 2.0 μm, and a total data size of 2.5 GB. Available to download from: 449
https://dx.doi.org/10.5281/zenodo.5270244 450

File S2 *Tribolium castaneum* **embryo dataset derived from light sheet** 451
microscopy. The dataset is freely available on the Cell Tracking Challenge 452
website (http://celltrackingchallenge.net/3d-datasets) - only the first timepoint 453
was used in the present study. The stack dimensions are 965 x 1871 x 991 voxels, 454
with each voxel being 0.381 x 0.381 x 0.381 μm, and a total data size of 3.3 GB. 455
Available to download from: https://dx.doi.org/10.5281/zenodo.5270323 456

File S3 GIANI settings used to produce the data in Fig 2 457

File S4 GIANI settings used to produce the data in Fig 3 458

File S5 GIANI and Imaris settings used to produce the data in Fig 4 459

File S6 GIANI and Imaris settings used to produce the data in Fig 5 460

File S7 GIANI and Imaris settings used to produce the data in Fig S1 461

File S8 CellProfiler pipeline and data used to produce Fig S2. Available 462
to download from: https://dx.doi.org/10.5281/zenodo.5286507 463

File S9 GIANI and Imaris settings and data used to produce Fig S3. 464
Available to download from: https://dx.doi.org/10.5281/zenodo.5286670 465

Acknowledgments 466

Thanks to Kurt Anderson for his constructive comments on the manuscript. This 467
work was supported by the Francis Crick Institute, which receives its core 468
funding from Cancer Research UK (FC001120, FC001999), the UK Medical 469

References

1. Duval K, Grover H, Han LH, Mou Y, Pegoraro AF, Fredberg J, et al. Modeling Physiological Events in 2D vs. 3D Cell Culture. Physiology. 2017;32(4):266–277.

2. Meijering E, Carpenter AE, Peng H, Hamprecht FA, Olivo-Marin JC. Imagining the future of bioimage analysis. Nature Biotechnology. 2016;34(12):1250–1255.

3. Lamprecht MR, Sabatini DM, Carpenter AE. CellProfiler™: free, versatile software for automated biological image analysis. BioTechniques. 2007;42(1):71–75.

4. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source platform for biological-image analysis. Nature Methods. 2012;9:676.

5. de Chaumont F, Dallongeville S, Chenouard N, Hervé N, Pop S, Provoost T, et al. Icy: an open bioimage informatics platform for extended reproducible research. Nature Methods. 2012;9:690.

6. Lou X, Kang M, Xenopoulos P, Muñoz-Descalzo S, Hadjantonakis AK. A Rapid and Efficient 2D/3D Nuclear Segmentation Method for Analysis of Early Mouse Embryo and Stem Cell Image Data. Stem Cell Reports. 2014;2(3):382 – 397.

7. Tosi S, Bardia L, Filgueira MJ, Calon A, Colombelli J. LOBSTER: An environment to design bioimage analysis workflows for large and complex fluorescence microscopy data. Bioinformatics. 2019;.

8. Rajasekaran B, Uriu K, Valentin G, Tinevez JY, Oates AC. Object Segmentation and Ground Truth in 3D Embryonic Imaging. PLOS ONE. 2016;11(6):1–17.

9. Gerri C, McCarthy A, Alanis-Lobato G, Demtschenko A, Bruneau A, Loubersac S, et al. Initiation of a conserved trophectoderm program in human, cow and mouse embryos. Nature. 2020;587(7834):443–447.

10. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nature methods. 2012;9:671–675.

11. Linkert M, Rueden CT, Allan C, Burel JM, Moore W, Patterson A, et al. Metadata matters: access to image data in the real world. The Journal of Cell Biology. 2010;189(5):777–782.

12. Tinevez JY, Perry N, Schindelin J, Hoopes GM, Reynolds GD, Laplantine E, et al. TrackMate: An open and extensible platform for single-particle tracking. Methods. 2017;115:80 – 90.

13. Legland D, Arganda-Carreras I, Andrey P. MorphoLibJ: integrated library and plugins for mathematical morphology with ImageJ. Bioinformatics. 2016;32(22):3532–3534.

14. Ollion J, Cochennec J, Loll F, Escudé C, Boudier T. TANGO: a generic tool for high-throughput 3D image analysis for studying nuclear organization. Bioinformatics. 2013;29:1840–1841.

15. Cautun M, van de Weygaert R, Jones BJT. NEXUS: tracing the cosmic web connection. Monthly Notices of the Royal Astronomical Society. 2012;429(2):1286–1308. doi:10.1093/mnras/sts416.

16. Jones TR, Carpenter A, Golland P. Voronoi-Based Segmentation of Cells on Image Manifolds. In: Liu Y, Jiang T, Zhang C, editors. Computer Vision for Biomedical Image Applications. Berlin, Heidelberg: Springer Berlin Heidelberg; 2005. p. 535–543.

17. Kostrykin L, Schnörr C, Rohr K. Globally optimal segmentation of cell nuclei in fluorescence microscopy images using shape and intensity information. Medical Image Analysis. 2019;58:101536.

18. Weigert M, Schmidt U, Haase R, Sugawara K, Myers G. Star-convex Polyhedra for 3D Object Detection and Segmentation in Microscopy; 2019.

19. Blin G, Sadurska D, Portero Migueles R, Chen N, Watson JA, Lowell S. Nessys: A new set of tools for the automated detection of nuclei within intact tissues and dense 3D cultures. PLOS Biology. 2019;17(8):1–29.

20. Caicedo JC, Goodman A, Karhohs KW, Cimini BA, Ackerman J, Haghighi M, et al. Nucleus segmentation across imaging experiments: the 2018 Data Science Bowl. Nature Methods. 2019;.

21. Yang L, Ghosh RP, Franklin JM, Chen S, You C, Narayan RR, et al. NuSeT: A deep learning tool for reliably separating and analyzing crowded cells. PLOS Computational Biology. 2020;16(9):1–20.

22. Gómez-de Mariscal E, García-López-de Haro C, Ouyang W, Donati L, Lundberg E, Unser M, et al. DeepImageJ: A user-friendly environment to run deep learning models in ImageJ. bioRxiv. 2021;doi:10.1101/799270.

23. Arganda-Carreras I, Kaynig V, Rueden C, Eliceiri KW, Schindelin J, Cardona A, et al. Trainable Weka Segmentation: a machine learning tool for microscopy pixel classification. Bioinformatics. 2017;33(15):2424–2426.

24. Berg S, Kutra D, Kroeger T, Straehle CN, Kausler BX, Haubold C, et al. ilastik: interactive machine learning for (bio)image analysis. Nature Methods. 2019;.

25. Cockburn K, Rossant J. Making the blastocyst: lessons from the mouse. The Journal of Clinical Investigation. 2010;120(4):995–1003.

26. Plusa B, Frankenberg S, Chalmers A, Hadjantonakis AK, Moore CA, Papalopulu N, et al. Downregulation of Par3 and aPKC function directs cells towards the ICM in the preimplantation mouse embryo. Journal of Cell Science. 2005;118(3):505–515.

27. Hirate Y, Hirahara S, Inoue KI, Suzuki A, Alarcon VB, Akimoto K, et al. Polarity-dependent distribution of angiomotin localizes Hippo signaling in preimplantation embryos. Current biology. 2013;23(23791731):1181–1194.

28. Ralston A, Cox BJ, Nishioka N, Sasaki H, Chea E, Rugg-Gunn P, et al. Gata3 regulates trophoblast development downstream of Tead4 and in parallel to Cdx2. Development. 2010;137(3):395–403.

29. Cockburn K, Biechele S, Garner J, Rossant J. The Hippo Pathway Member Nf2 Is Required for Inner Cell Mass Specification. Current Biology. 2013;23(13):1195 – 1201.

30. Frum T, Murphy TM, Ralston A. HIPPO signaling resolves embryonic cell fate conflicts during establishment of pluripotency in vivo. eLife. 2018;7(30526858):e42298.

31. Nishioka N, ichi Inoue K, Adachi K, Kiyonari H, Ota M, Ralston A, et al. The Hippo Signaling Pathway Components Lats and Yap Pattern Tead4 Activity to Distinguish Mouse Trophectoderm from Inner Cell Mass. Developmental Cell. 2009;16(3):398 – 410.

32. Kjær S, Linch M, Purkiss A, Kostelecky B, Knowles P, Rosse C, et al. Adenosine-binding motif mimicry and cellular effects of a thieno[2,3-d]pyrimidine-based chemical inhibitor of atypical protein kinase C isoenzymes. Biochemical Journal. 2013;451(2):329–342.

33. Rodriguez J, Peglion F, Martin J, Hubatsch L, Reich J, Hirani N, et al. aPKC Cycles between Functionally Distinct PAR Protein Assemblies to Drive Cell Polarity. Developmental Cell. 2020;42(4):400–415.e9.

34. Aguilar-Aragon M, Elbediwy A, Foglizzo V, Fletcher GC, Li VSW, Thompson BJ. Pak1 Kinase Maintains Apical Membrane Identity in Epithelia. Cell Reports. 2020;22(7):1639–1646.

35. Yamada KM, Sixt M. Mechanisms of 3D cell migration. Nature Reviews Molecular Cell Biology. 2019;.

36. Masuzzo P, Troys MV, Ampe C, Martens L. Taking Aim at Moving Targets in Computational Cell Migration. Trends in Cell Biology. 2016;26(2):88 – 110.

37. Wolff C, Tinevez JY, Pietzsch T, Stamataki E, Harich B, Guignard L, et al. Multi-view light-sheet imaging and tracking with the MaMuT software reveals the cell lineage of a direct developing arthropod limb. eLife. 2018;7:e34410.

38. Pietzsch T, Preibisch S, Tomančák P, Saalfeld S. ImgLib2—generic image processing in Java. Bioinformatics. 2012;28(22):3009–3011.

39. Haase R, Royer LA, Steinbach P, Schmidt D, Dibrov A, Schmidt U, et al. CLIJ: GPU-accelerated image processing for everyone. Nature Methods. 2020;17(1):5–6.

Figure 1

A

B

C
GIANI Cell Count Error

D
GIANI Nuclear Localisation Error (μm)

E
GIANI Cell Localisation Error (μm)

F
GIANI Cell Count Error

G
GIANI Nuclear Localisation Error (μm)

H
GIANI Cell Localisation Error (μm)

Figure 2

Figure 3

Figure 4

A

Raw   Imaris   GIANI

B

Raw   Imaris   GIANI

Figure 5

Supplementary Figure 1

Supplementary Figure 2

A

B

Raw   Imaris   GIANI

Supplementary Figure 3

Supplementary Figure 4