Cortical synaptic diversity naturally arises with heterogeneous neural populations

Jacob L. Yates[1], Benjamin Scholl[2]

[1]University of Maryland, College Park, Maryland, USA
[2]University of Pennsylvania, Philadelphia, PA, USA

Running title: Synaptic diversity results from heterogeneity

Corresponding author: Benjamin Scholl, scholl.ben@gmail.com

Number of pages: 22
Number of figures: 7
Number of tables: 0
Number of words Abstract: 148
Number of words Introduction: 723
Number of words Discussion: 988

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgements: Jacob Yates is an Open Philanthropy fellow of the Life Sciences Research Foundation. Benjamin Scholl is supported by K99EY031137.

We thank Alex Huk, Richard Lange, Sabya Shivkumar, and Krishnan Padmanabhan for helpful comments and discussion.

Author contributions statement: J.Y and B.S. conceived of the experiment. J.Y. derived analytic solutions, J.Y and B.S. preformed analysis, J.Y and B.S. wrote the manuscript.

Abstract

Synaptic inputs onto single cortical neurons *in vivo* exhibit substantial functional diversity with respect to sensory-driven activity. However, it is unclear what this diversity reflects, appearing counter-productive in generating tuned responses to specific stimuli. We propose that functional diversity naturally arises if neurons extract information encoded from noisy input populations. Focusing on a single sensory variable, orientation, we construct a probabilistic decoder that estimates orientation from the responses of a realistic hypothetical input population of neurons. Analytically derived weights exhibit diversity when input populations consist of noisy, correlated, and heterogeneous neurons. Weight diversity was necessary to accurately decode orientation. Further, *in silico* weight diversity matched the functional heterogeneity of dendritic spines imaged *in vivo*. This suggests that synaptic diversity is expected when information is extracted from realistic input populations, highlighting the importance of studying weighting structures in population coding theory and consideration in pursuits of the cortical connectome.

Introduction

Cortical neurons are driven by large populations of excitatory synaptic inputs. Synaptic populations ultimately shape how sensory signals are encoded, decoded, or transformed. The sensory representation or functional properties of an excitatory input population will define and constrain the operations a neuron can perform. Electrophysiological and anatomical studies have suggested that excitatory connectivity exhibits functional specificity (Cossell et al., 2015; Ko et al., 2011; Lee et al., 2016; Reid and Alonso, 1995). In contrast, with the ability to capture the activity of presynaptic populations onto single spiny cortical neurons, a different picture has emerged: presynaptic populations exhibit functional diversity, deviating from canonical connectivity rules (Scholl and Fitzpatrick, 2020). Functional diversity within input populations has been observed in a variety of mammalian species, from rodents to primates, and for a variety of sensory stimuli (Chen et al., 2013, 2011; Iacaruso et al., 2017; Jia et al., 2011, 2010; Kerlin et al., 2019; Scholl et al., 2017; Wertz et al., 2015; Wilson et al., 2018, 2016). This apparent discrepancy challenges our understanding about how synaptic inputs drive the selective outputs of cortical neuron and leads to a fundamental question: What *should* the functional diversity of inputs be if neurons accurately represent sensory stimuli?

To answer this question, we borrow from population coding theory; starting with the idea that to accurately represent sensory signals, cortical neurons must decode the activity of

upstream populations. This decoding is likely accomplished by combing signals across neural populations (Graf et al., 2011; Jazayeri and Movshon, 2006). Many studies have examined how sensory variables might be decoded from cortical populations (Butts and Goldman, 2006; Graf et al., 2011; Shamir and Sompolinsky, 2006), an endeavor increasingly applied to larger population sizes with innovative recording techniques (Rumyantsev et al., 2020; Stringer et al., 2019). These decoding approaches are often used as a tool to quantify the information about a stimulus available in a neural population, carrying the assumption that downstream areas could be performing such a process (Berens et al., 2011; DiCarlo et al., 2012). Real neural decoders must be composed of individual neurons, driven by a set of presynaptic inputs, akin to a decoder's weights over a given input population. Few studies have explicitly examined the weight structure of population decoders (Jazayeri and Movshon, 2006; Rust et al., 2006; Zavitz and Price, 2019). This is partly due to the fact that a decoder's weights will depend on the statistics of the input population and it has not been clear what can be interpreted from those weights. For example, the functional homogeneity of a neuron population and strength of neuronal correlations will directly impact a downstream decoder and determine the weight structure (Ecker et al., 2011; Kohn et al., 2016; Shamir, 2014).

In this paper, we investigate what the weights look like for a simple population decoder and how they compare to real synaptic inputs measured *in vivo*. Focusing on a single sensory variable, orientation, we derive the maximum-likelihood readout for a simulated input population that encodes stimuli with tuning curves plus noise (Ecker et al., 2011 and others). Under reasonable assumptions, the decoder weights can be interpreted the synaptic connectivity between the input population and the downstream decoder neurons. This allows us to examine how synaptic connectivity depends on properties of the input population and to directly compare population decoders to synaptic input measured *in vivo*. Here we test a hypothesis that an optimal decoder will show substantial heterogeneity in its synaptic weights given biologically realistic input population. We find that when input populations are homogeneous, the expected synaptic excitatory inputs closely resemble the somatic output. However, with a heterogeneous input population, the expected inputs onto readout neurons exhibit functional diversity. We then compare the orientation tuning of simulated inputs with large populations of dendritic spines (excitatory synaptic inputs) onto individual neurons of ferret primary visual cortex (V1), recorded with two-photon calcium imaging *in vivo* (Wilson et al., 2018, 2016). This comparison revealed strikingly similar diversity in the orientation tuning of dendritic spines and simulated inputs. The similarity between the synaptic populations of actual V1 neurons and the optimal neural decoder suggest that diversity and heterogeneity observed in dendritic spines across sensory cortices are,

in fact, expected when considering realistic noisy populations of diverse inputs and reflect the nature of how information is propagated through neural circuits in the presence of noise.

Results

Following several decades of work on population codes, we derive a Bayesian decoder to report the identity of a visual stimulus represented by a neural population (Fig. 1). With this framework, and given the specifics of the *encoding* population, we can analytically derive the optimal *decoding* weights of a population of readout neurons. Here, we use "optimal" to refer to the maximum-likelihood solution. Previous work has shown that a population of neurons could, in principle, perform such probabilistic decoding with weighted summation and divisive normalization, as long as their inputs exhibit Poisson-like noise (Jazayeri and Movshon, 2006; Ma et al., 2006). Starting from that basic framework, we derived a decoder that represents the probability that a particular stimulus orientation was present given the responses of a large population of upstream, input neurons (P_{IN}). This decoder has weight vectors for each possible stimulus which integrate across P_{IN} and are passed through simple nonlinearities (normalized summation and an exponential nonlinearity). As we will show below, given specific assumptions about the variability in P_{IN}, the weights over P_{IN} depend systematically on the tuning functions and covariance of P_{IN}. Following a characterization of this decoding framework, we will make direct comparisons with real data: defining an effective "synaptic input population" (P_{SYN}) as nonzero, positive weights over P_{IN}. Although our strategy applies to any one-dimensional stimulus variable, we describe this model in the context of the orientation of drifting gratings presented to V1 neurons, for a direct comparison with *in vivo* measurements.

A neural population as a probabilistic decoder

A probabilistic decoder reports the probability that a particular stimulus orientation, θ_k, was present given the spiking responses of an input population, $\boldsymbol{R}$. This can be expressed as a normalized exponential function of the log-likelihood plus the log prior for each θ_k,

$$p(\theta_k | R) = \frac{p(R|\theta_k)p(\theta_k)}{\sum_{i=1}^{K} p(R|\theta_i)\,p(\theta_i)} = \frac{e^{L(\theta_k)}}{\sum_{i=1}^{j} e^{L(\theta_i)}}$$

where

$$L(\theta_k) = ln\big(p(R|\theta_k)\big) + ln\big(p(\theta_k)\big)$$

The likelihood, $p(\boldsymbol{R} | \theta_k)$, is the probability of the observed responses in an input population given the stimulus k and $p(\theta_k)$ is the prior probability of that stimulus class. If $p(\boldsymbol{R} | \theta_k)$ is in the exponential

family, then $L(\theta_k)$ becomes as a weighted sum of the population response vector plus an offset, which can be estimated numerically via multinomial logistic regression (Ma et al., 2006). For simplicity we assume the input population has a response that is a function of the stimulus plus Gaussian noise, and equal covariance across all stimulus conditions. Although this assumption deviates from real neural activity, in this simple case, the weights and offset can be solved analytically (see Methods) and, as will be shown below, such a simple model makes substantial headway in explaining the biological phenomena. Under the Gaussian assumption, the decoder amounts to:

$$p(\theta_k|R) = \frac{e^{\left(R^T w_k + \beta_k\right)}}{\sum_i^K e^{\left(R^T w_i + \beta_i\right)}} \qquad (1)$$

where

$$w_k = Q^{-1} f(\theta_k)$$

$$\beta_k = -\left(\frac{1}{2}\right) f(\theta_k)^T Q^{-1} f(\theta_k) + \ln p(\theta_k)$$

Here, $f(\theta_k)$ is the mean input-population response to stimulus orientation, k, K is the total number of orientations, and Q is the covariance matrix. The covariance term captures the influence of each neuron's response variance (diagonal elements) and the variability shared with other neurons (off-diagonal elements). Intuitively, in the absence of covariability (i.e., off-diagonal elements are zero), the weights are proportional to the signal-to-noise ratio of the neuron (the mean divided by the variance). The term $R^T w_k$ is the dot product between the population response and weights. The second term, β_k, is an offset for each stimulus. It is also worth noting that if the covariance depends on the stimulus, the optimal readout is no longer a linear function of R and is quadratic, which can be interpreted as a complex-cell (Pagan et al., 2016) and is a potentially fruitful future direction.

In this study, we focus on the weights of the Gaussian, equal covariance decoder in order to examine how synaptic tuning from such a simple decoder would arise. Because the optimal weights have an analytic solution (eq 1), we can see how they depend on the parameters of P_{IN}. The simplifying assumptions we use to derive the maximum-likelihood weights help build intuitions about what can be expected in biological circuits, and linear weights such as these could be learned by real neural systems (Dayan and Abbott, 2001). A key difference here from prior work is that rather than focus on discrimination, we treat orientation estimation as a multiclass identification problem, discretizing θ such that for each possible θ_k, there is a separate weight

vector. Thus, in this derivation, the optimal weights depend on the tuning curves themselves, not the derivative.

Characteristics of a neural population decoder

To understand how synaptic weights depend on input statistics, we derived maximum-likelihood weights for input populations, P_{IN}, with different tuning and covariance. To generate P_{IN} we simulated N neurons responding to K oriented stimuli ($\theta = [-\pi/2:\text{pi}/2K:\pi/2]$). We briefly describe the construction of P_{IN} (full details are described in the Methods). Each neuron is defined by a tuning function and noise term, describing trial-by-trial variability, which are summed to generate stimulus-driven responses. We compared two types of input populations: homogeneous and heterogenous. A homogeneous P_{IN} consists of shifted copies of a single tuning curve (Fig. 2a). Heterogenous P_{IN} were generated to match measurements from macaque V1 (Ringach et al., 2002). This P_{IN} consisted of tuning curves closely resembling V1 physiology in terms of the variation in peak firing rate, bandwidth, and baseline firing rate (Fig. 2d). Limited-range correlations were included such that the noise correlation between two neurons depends on the difference in their tuning preferences (Ecker et al., 2011; Kohn et al., 2016).

Statistics of P_{IN} responses, **R,** will dictate the weight structure for neurons in a decoding population. For a homogeneous P_{IN}, the weights are smooth across orientation space and exhibit three primary features: a prominent peak about the preferred orientation of the output tuning, slight negative weights for orientations just outside the preferred, and near-zero weights at orthogonal orientations (Fig. 2c). With more realistic tuning diversity (heterogenous P_{IN}), optimal weights are no longer smooth (Fig. 2d). While the optimal weights appear to roughly have the same overall shape as for homogeneous P_{IN}, there is considerable positive and negative weighting across orientation space. Despite substantial changes in optimal weight vectors, the decoder output (i.e. somatic response) tuning was narrow (Fig. 2f), similar to the output for the homogeneous case (Fig. 2e).

To explore the importance of decoding weight diversity, we imposed a smoothing penalty on the weight vectors (Park and Pillow, 2011) and calculated cross-validated decoder accuracy using the mean-squared error between the maximum a posteriori estimation and true stimulus (see Methods). Different degrees of smoothing are shown for an example set of weights in Figure 3a. We simulated a range of population sizes ($N = 2 - 2048$) and correlations ($c_o = 0, 0.25, 0.50$). Without noise-correlations, the accuracy of all decoders increases with population size, with a homogenous P_{IN} preforming best (Fig. 3b). However, in the presence of noise-correlations,

accuracy saturates for large homogenous P_{IN} (Fig. 3c-d). As previously shown (Ecker et al., 2011), accuracy for heterogeneous populations with limited-range correlations does not saturate (Fig. 3c-d). However, this is dependent on the weights exhibiting diversity. Smoothing the weights for heterogeneous P_{IN} caused saturation and decreased accuracy (Fig. 3c-d), demonstrating that weight amplitude diversity in analytically derived weights distributions are critical for the decoder performance.

For the remainder of simulations in this study, we used set $N = 1000$ for P_{IN} and set $c_o = 0.20$.

Simulating excitatory weight tuning

In order to compare analytically derived weights with the synaptic inputs onto V1 neurons measured *in vivo*, we generated excitatory synaptic input populations (P_{SYN}) from the weights. Under the assumption that synaptic integration is linear, two synapses of equal weight are the same as one synapse with double that weight. This creates a degeneracy where synapse count and size trade off. Because current spine imaging techniques typically capture large synapses (Scholl et al., 2019a), we can assume size is fixed and convert the derived weights into a frequency distribution of 'synaptic' inputs (Fig. 4a). The tuning curve for such a synapse is the tuning curve of the input and thus, the synaptic input population, $P_{SYN,}$ is the input population resampled with probabilities given by the derived weights. An example P_{SYN} for a single decoder neuron is shown in Figure 4b (drawn from the heterogeneous P_{IN} in Figure 2). P_{SYN} in this example displays some specificity in orientation tuning relative to the somatic output, indicated by a larger proportion of simulated synapses with similar orientation preference as the somatic output ($\phi_k = 0°$) of the decoder neuron.

Empirical distribution of dendritic spine tuning consistent with decoding of heterogeneous input population

We analyzed two-photon calcium imaging of soma and corresponding dendritic spines on individual neurons in ferret V1 during the presentation of oriented drifting gratings (Wilson et al., 2018, 2016). While our model draws from a P_{IN} matched to measurements from macaque V1, the orientation tuning of ferret neurons in layer 2/3 V1, as measured by two-photon cellular imaging, exhibit a similar range in selectivity (Wilson et al., 2017). Visually responsive and isolated dendritic spines (see Methods) typically exhibited diverse orientation tuning relative to the somatic output,

although some individual cells show greater overall diversity (Fig. 5b) than others (Fig. 5a). To characterize P_{SYN} diversity, both for real dendritic spines and simulated inputs, we computed the Pearson correlation coefficient between the tuning curves of individual inputs and the corresponding somatic output. For these comparisons, we sampled orientation space in our model to match our empirical measurements (22.5 deg increments) and the number of total excitatory inputs recovered for each simulated downstream neuron was set to 100 to match the average number of visually-driven spines recorded for each ferret V1 neuron (n = 28, n = 103.3 ± 37.8 spines/cell). Simulations were run 1000 times.

　　　Across all simulated inputs, spine-soma tuning correlation was higher for homogeneous P_{SYN} compared to heterogeneous P_{SYN} (median r_{hom} = 0.72, median r_{hom} = 0.24, n = 90000; Fig. 5c). Tuning correlation between all imaged dendritic spines and soma was also low (median r_{cell} = 0.35, n = 2,892 spines from 28 cells), more closely resembling our model with a heterogenous P_{IN}. As somatic orientation selectivity (i.e., tuning bandwidth) varies for single cells in ferret V1 (Goris et al., 2015; Wilson et al., 2016), we next examined the average input-output tuning correlation across individual cells (Fig. 5d). A homogeneous model exhibited more specificity in inputs that a heterogeneous model (median r_{hom} = 0.64; median r_{het} = 0.20, n = 9000). Across our population of ferret V1 cells we observed similar spine-soma correlation as the heterogeneous simulation (median r_{cell} = 0.21, n = 28). These data were not statistically different from a heterogeneous P_{SYN} (p = 0.49, Mann-Whitney test), while homogeneous P_{SYN} showed significantly larger correlations (p < 0.001, Mann-Whitney test). It is worth noting that a small percentage of imaged cells (17.9%, n = 5/28) had synaptic populations whose mean tuning correlation were within the 95% confidence interval of the homogeneous model distribution. Additionally, some cells had negative average correlations with their spines, which never occurred in the models. This implies nonlinearities between the spines and soma.

　　　We also examined the average orientation preference difference between input and output within single cells (Fig. 5e). As for tuning correlation, there was far more synaptic specificity for homogeneous, compared to heterogeneous, P_{SYN} (median $\Delta\theta_{hom}$ = 17.2˚, median $\Delta\theta_{het}$ = 44.1˚). Dendritic spines in ferret V1 cells displayed significantly less orientation matching than the homogeneous model (median $\Delta\theta_{cell}$ = 32.1˚; p = 0.029 Mann-Whitney test), but also exhibited significantly greater matching than the heterogeneous model (p < 0.001 Mann-Whitney test). Taken together, these results support the idea that a decoding framework naturally produces functional diversity of synaptic inputs that is observed in individual V1 neurons.

Functional properties of measured and simulated synaptic inputs

As a heterogenous P_{IN}, with biologically realistic parameters, closely matched our *in vivo* data from ferret V1, we explored the functional properties of dendritic spines and decoder weights (i.e. P_{SYN}) in greater detail. We focus on two important characteristics of inputs onto V1 neurons: orientation tuning preference and selectivity. Inputs can exhibit a broad range of preferences relative to the output; some contribute directly to the soma, while the purpose of others (e.g. orthogonally-tuned) is less understood (Scholl and Fitzpatrick, 2020). V1 neurons also range in their tuning selectivity to orientation, which can be characterized by a vector strength index (see Methods). Selective inputs are proposed to provide more information about encoded stimulus variables (Seriès et al., 2004; Shamir and Sompolinsky, 2006; Zavitz and Price, 2019), and unselective (or poorly selective) inputs could convey information through their covariance with selective neurons (Zylberberg, 2017).

For each neuron in our dataset, we measured the orientation preference difference and selectivity of dendritic spines, and calculated the joint probability distribution, pooling across each to generate an average (Fig. 6a). There was an overall bias for spines co-tuned to the soma, but tuning selectivity varied broadly for all orientation preference differences. However, orthogonally-tuned spines exhibited greater selectivity than those co-tuned to the soma ($\Delta\theta_{ori}$ > 67.5°: median = 0.42, $\Delta\theta_{ori}$ < 22.5°: median = 0.31; p < 0.001, Mann-Whitney test). To explore decoder excitatory weights, we simulated a heterogeneous P_{SYN} (same parameters as in Figure 2, 100 iterations) for a single decoder neuron. Here we found that co-tuned excitatory weights occurred most frequently, but orthogonally-tuned excitatory weights exhibited slightly less selectivity ($\Delta\theta_{ori}$ > 67.5°: median = 0.28, $\Delta\theta_{ori}$ < 22.5°: median = 0.35; p < 0.001, Mann-Whitney test). Clearly, some aspects of our observations *in vivo* are not captured by our simple model and remain to be elucidated. Taken together, our simulation of heterogenous P_{IN} suggest the collection of orthogonally-tuned excitatory inputs in cortical neurons *in vivo* are not aberrant. Instead, the synaptic architecture of layer 2/3 neurons in ferret visual cortex are likely optimized for the readout of downstream populations tuned to orientation.

Discussion

We used a population decoding framework (Jazayeri and Movshon, 2006; Kohn et al., 2016; Pouget et al., 2000; Shamir, 2014) to understand possible functional sources of synaptic diversity. We find that even simple decoders exhibit substantial heterogeneity in their weights as long as the inputs are noisy, correlated neural populations with heterogeneous orientation tuning.

We argue that this could naturally explain the heterogeneity in synaptic inputs measured in vivo if neural populations are decoding information from upstream populations. We compared two neural decoders: one with homogenous input (Jazayeri and Movshon, 2006) and one with heterogenous input (Ecker et al., 2011). We show that empirical measurements from dendritic spines recorded within individual cortical neurons in ferret V1 exhibit a similar amount of diversity in orientation tuning as simulated inputs (i.e. excitatory weights) from heterogeneous input populations. It may appear trivial that heterogeneous input populations would produce heterogeneous weights, but it was neither immediately obvious that the weights would not be smooth nor that neurons with orthogonal tuning would receive weights. Orthogonally-tuned or non-preferred inputs are often considered to be aberrant; to be pruned away during experience-dependent plasticity or development (Holtmaat and Svoboda, 2009). Our decoding approach suggests these inputs are purposeful and emerge through development as cortical circuits learn the statistics of their inputs (Avitan and Goodhill, 2018). Taken together, our results shed light on synaptic diversity that has been puzzling from recent measurements (Scholl and Fitzpatrick, 2020), suggesting that it is, in fact, expected given known properties of the input population.

We believe our study is a significant step forward in combing population coding theory (Averbeck et al., 2006; Pouget et al., 2000) with novel biological measurements in highly organized visual systems (Wilson et al., 2016). The ability to measure receptive field properties and statistics of sensory-driven responses of synapses *in vivo* provides a new testing bed for population codes. The individual neurons which synapses converge on are the real components of what has long been a hypothetical downstream population decoder. While we did not set out to build a computational or biophysical model of a neuron, we believe simplistic approaches such as ours are fruitful for understanding basic principles. We hope future studies build upon this modeling framework, exploring quadratic decoders for example (Pagan et al., 2016), and work towards richer visual stimuli and neural models (Chalk et al., 2017).

Inhibition (i.e. inhibitory weights) were not explicitly studied in this work. We focused on excitatory weights as there are now a number of studies reporting the functional diversity of dendritic spines on cortical neurons (Chen et al., 2013, 2011; Iacaruso et al., 2017; Jia et al., 2010, 2011; Kerlin et al., 2019; Scholl et al., 2017, 2019a; Wertz et al., 2015; Wilson et al., 2016, 2018), which are the recipients of excitatory afferents. Few techniques currently exist to access inhibitory populations *in vivo* (Scholl and Fitzpatrick, 2020). Nevertheless, we were curious about the functional properties of simulated inhibitory weights, which exhibit similar diversity as excitatory weights (Fig. 2e). Applying the same procedure as in Figure 6b on putative inhibitory weights, we found that the joint probability distribution for selectivity and orientation preference

difference was slightly different than excitatory weights (Fig. 7). First, inhibitory weights were more broadly distributed across orientation space, consistent with a recent study measuring layer 2/3 inhibitory inputs in ferret V1 with optogenetic mapping and electrophysiology (Scholl et al., 2019b). Second, inhibitory weights were largely selective for orientation, which is what is to be expected in the V1 of carnivores, where inhibitory interneurons are selective for visual features (Keller and Martin, 2015; Wilson et al., 2017), but difficult to relate to mouse V1, where most are unselective (Hofer et al., 2011; Kerlin et al., 2010; Runyan et al., 2010; Scholl et al., 2015). Third, there is an overall trend for inhibitory weights to be slightly co-tuned with the somatic output or excitatory weight structure, and this has been suggested to be a fundamental principle underlying cortical circuits. Fourth, inhibitory weights exhibited an inverse relationship compared to the excitatory weight structure: more selective inputs were weighted more heavily for orientations differing form the somatic (decoder neuron) output. Going forward, it will be crucial to develop techniques and measure functional properties of presynaptic inhibitory populations, alongside coalesced excitatory populations for validation of different modeling frameworks.

To limit complexity, our decoder did not account for many aspects of cortical networks such as stimulus-dependent correlations or recurrent connections. In the case of stimulus dependent covariance, the optimal decoder is no longer linear, however, that decoder closely resembles a complex cell (Pagan et al., 2016). Extending this decoder to include realistic noise has been used to capture many nonlinear features of neural responses including divisive normalization, gain control and contrast-dependent temporal dynamics, all of which fall naturally from a normative framework (Chalk et al., 2017). These more sophisticated approaches may be able to make predictions about the synaptic organization itself, whereby local clusters of synapses act as nonlinear subunits (Ujfalussy et al., 2018).

Finally, it is important to emphasize that our model does not describe a cortical transformation. Instead, to limit complexity, we focused on the propagation of orientation selectivity from one neural population to another, akin to the propagation of basic receptive field properties from V1 to higher-visual areas. Our approach was an to provide a starting point for predicting the tuning diversity of synaptic input populations as compared to the tuning output or downstream cells. However, this model could be modified to study the convergence and transformation of cortical inputs. An obvious case study would be complex cells in layer 2/3 V1 (Hubel and Wiesel, 1962; Movshon et al., 1978; Spitzer and Hochstein, 1988), which are thought to integrate across presynaptic cells with similar oriented receptive field with offset spatial subunits to produce polarity invariance. This extension would be better suited for a nonlinear quadratic decoder (Pagan et al., 2016), rather than the linear one used here.

Figure Legends

Figure 1: A population decoding framework to study synaptic diversity.
An upstream population of neurons are tuned for a single stimulus variable (orientation) (*top*). This input population is readout by downstream decoder neurons (*bottom*). Downstream neurons decode stimulus identify by reading out spikes from the upstream input population. Each decoder neuron is defined by a set weights (*middle*) over the upstream population, which are summed and rectified to produce an output.

Figure 2: Model simulations with homogenous and heterogeneous input populations.
(a) Orientation tuning of a homogenous input population. Shown is a subset of the total population (n = 20/1000). Ordinate is orientation preference, restricted between -90^o and 90^o. (b) Derived weights for a single decoder neuron (preferring 0^o) reading out the homogenous (*blue*) input population in (a). Weights for homogenous populations smoothly vary over orientation space. (c) Response output of the decoder neuron whose weights are shown in (b). (d-f) Same as in (a-c) for a heterogeneous input population with moderate correlation (c_o = 0.25). Note that decoder weights for heterogeneous input populations are not smooth.

Figure 3: Decoder performance of heterogeneous input populations depends on population size, correlations, and weight diversity.
(a) Example weight distribution for a decoder neuron reading out a heterogeneous input population (*top*). Shown are the effects of progressively smoothing weights. Ordinate is orientation preference, restricted between -90^o and 90^o. (b) Decoder performance (inverse mean-squared-error) plotted for homogenous and heterogeneous input populations of increasing size. Simulations here include no correlations (c_o = 0). Shading indicates standard error. (c) Same as in (b) for input populations with moderate correlation (c_o = 0.25). (d) Same as in (b) for input populations with stronger correlation (c_o = 0.50).

Figure 4: Simulation of synaptic populations from decoder neuron weight distributions. (a) Example weight distribution for a single decoder neuron tuned to 0^o (*left*). Ordinates are orientation preference, restricted between -90^o and 90^o. Dashed line separates excitatory (positive) and inhibitory (negative) weights. Excitatory weight distribution over the input population is transformed into a frequency distribution, whereby greater amplitude equates to greater frequency of occurrence (*right*). (b) Example simulated synaptic population (n = 100 spines) from

the weight distribution in (a). Shown are the orientation tuning curves of each simulated synapse (normalized).

Figure 5: Orientation tuning diversity of dendritic spine populations in ferret V1 match simulations with correlated, heterogeneous input populations.

(a) Two-photon standard-deviation projection of example dendrite and spines recorded from a single cell (*left*). Inset: Two-photon standard-deviation projection of corresponding soma. Scale bar is 10 microns. Orientation tuning of soma (*top*) and all visually-responsive dendritic spines from this single cell (n = 159) are shown (*right*). Spine responses are normalized. Orientation preferences are shown relative to the somatic preference (aligned to $0°$). (b) Same as in (a) for another example cell (n = 162 visually-responsive spines). (c) Cumulative distributions of tuning correlation between individual dendritic spines or simulated synaptic inputs with corresponding somatic tuning or decoder output. Shown are correlations of simulations of homogenous (*blue*) or heterogeneous (*red*) input populations, compared to empirical data (*gray*). (d) Distributions of average tuning correlation between synaptic input and somatic output across measured cells (28). Also shown are distributions of average tuning correlation for simulated cells. Triangles denote median values for each distribution. (e) Distributions of average synaptic orientation preference, relative to somatic output, for empirical data and simulations. Triangles denote median values for each distribution.

Figure 6: Excitatory input joint distribution of tuning preference and selectivity. (a) Joint probability distribution of dendritic spine orientation preference difference with the soma and tuning selectivity. Shown is the average frequency across all cells (n = 28). 2D pixel values are average frequency value (darker = larger). Also shown are the marginal distributions for tuning preference (top) and selectivity (right). (b) Same as in (a) for simulated excitatory inputs. Simulation is the same as in Figure 2 for heterogenous input population. Excitatory inputs (100) extracted for a single decoder neuron tuned to $0°$. Average calculated over 100 simulations.

Figure 7: Inhibitory input joint distribution of tuning preference and selectivity. (a) Joint probability distribution of simulated inhibitory inputs (i.e. negative weights) spine orientation preference difference with the soma and tuning selectivity. Simulation is the same as in Figure 2 for heterogenous input population. Excitatory inputs (100) extracted for a single decoder neuron tuned to $0°$. Average calculated over 100 simulations. Also shown are marginal distributions for tuning preference (top) and selectivity (right).

Methods

All procedures were performed according to NIH guidelines and approved by the Institutional Animal Care and Use Committee at Max Planck Florida Institute for Neuroscience.

Derivation for a Bayesian probabilistic decoder

We construct a probabilistic decoder, represented by a population of neurons, that reports or estimates the identity of a stimulus from the spiking response of an input population of neurons. We assume Poisson-like noise, an input population with responses that are a function of the stimulus, $f(\theta_k)$, plus Gaussian noise, $f(\theta_k)$, and the covariance (Q) is equal for all stimulus conditions (Q) such that $Q = Q_k = Q_i$. Then, the posterior distribution can be written as

$$p(\theta_k | \boldsymbol{R}) = \frac{p(\theta_k)\, N(f(\theta_k), \boldsymbol{Q})}{\sum_i^K p(\theta_i)\, N(f(\theta_i), \boldsymbol{Q})}$$

where a bivariate Gaussian is

$$N(f(\theta_k), \boldsymbol{Q}) = ((2\pi)^n \boldsymbol{Q})^{-\frac{1}{2}}\, e^{\left(\left(-\frac{1}{2}\right)(\boldsymbol{R}-f(\theta_k))^T \boldsymbol{Q}^{-1}(\boldsymbol{R}-f(\theta_k))\right)}$$

This can be expanded and simplified such that

$$p(\theta_k | R) = \frac{p(\theta_k)((2\pi)^n \boldsymbol{Q})^{-\frac{1}{2}}\, e^{\left(\left(-\frac{1}{2}\right)(\boldsymbol{R}-f(\theta_k))^T \boldsymbol{Q}^{-1}(\boldsymbol{R}-f(\theta_k))\right)}}{\sum_i^K p(\theta_i)((2\pi)^n \boldsymbol{Q})^{-\frac{1}{2}}\, e^{\left(\left(-\frac{1}{2}\right)(\boldsymbol{R}-f(\theta_i))^T \boldsymbol{Q}^{-1}(\boldsymbol{R}-f(\theta_i))\right)}}$$

$$p(\theta_k | R) = \frac{p(\theta_k)\, e^{\left(\left(-\frac{1}{2}\right)\left(\boldsymbol{R}^T \boldsymbol{Q}^{-1}\boldsymbol{R} - f(\theta_k)^T \boldsymbol{Q}^{-1}\boldsymbol{R} - \boldsymbol{R}^T \boldsymbol{Q}^{-1}f(\theta_k) + f(\theta_k)^T \boldsymbol{Q}^{-1}f(\theta_k)\right)\right)}}{\sum_i^K p(\theta_i)\, e^{\left(\left(-\frac{1}{2}\right)\left(\boldsymbol{R}^T \boldsymbol{Q}^{-1}\boldsymbol{R} - f(\theta_i)^T \boldsymbol{Q}^{-1}\boldsymbol{R} - \boldsymbol{R}^T \boldsymbol{Q}^{-1}f(\theta_i) + f(\theta_i)^T \boldsymbol{Q}^{-1}f(\theta_i)\right)\right)}}$$

$$p(\theta_k | R) = \frac{p(\theta_k)\, e^{\left(-\frac{1}{2}\right)\boldsymbol{R}^T \boldsymbol{Q}^{-1}\boldsymbol{R}}\, e^{\left(f(\theta_k)^T \boldsymbol{Q}^{-1}\boldsymbol{R} + \left(-\frac{1}{2}\right)f(\theta_k)^T \boldsymbol{Q}^{-1}f(\theta_k)\right)}}{\sum_i^K p(\theta_i)\, e^{\left(-\frac{1}{2}\right)\boldsymbol{R}^T \boldsymbol{Q}^{-1}\boldsymbol{R}}\, e^{\left(f(\theta_i)^T \boldsymbol{Q}^{-1}\boldsymbol{R} + \left(-\frac{1}{2}\right)f(\theta_i)^T \boldsymbol{Q}^{-1}f(\theta_i)\right)}}$$

$$p(\theta_k | R) = \frac{e^{\left(\boldsymbol{R}^T \boldsymbol{Q}^{-1}f(\theta_k) + \left(-\frac{1}{2}\right)f(\theta_k)^T \boldsymbol{Q}^{-1}f(\theta_k) + \ln p(\theta_k)\right)}}{\sum_i^K e^{\left(\boldsymbol{R}^T \boldsymbol{Q}^{-1}f(\theta_i) + \left(-\frac{1}{2}\right)f(\theta_i)^T \boldsymbol{Q}^{-1}f(\theta_i) + \ln p(\theta_i)\right)}}$$

$$p(\theta_k | R) = \frac{e^{\left(\boldsymbol{R}^T w_k + \beta_k\right)}}{\sum_i^K e^{\left(\boldsymbol{R}^T w_i + \beta_i\right)}}$$

where

$$w_k = \boldsymbol{Q}^{-1} f(\theta_k)$$

$$\beta_k = -\left(\frac{1}{2}\right) f(\theta_k)^T Q^{-1} f(\theta_k) + \ln p(\theta_k)$$

Here, w are the weights over k for each neuron in the decoder population and β is an offset term over k. Importantly, because we assume Gaussian input, with this formulation, w and β are derived closed form.

Input population model

To generate input populations (P_{IN}), we simulated N neurons responding to a stimulus characterized by orientation ($\theta_k = [-\pi/2 : \pi/2K : \pi/2]$). The response of each neuron, r, depends on a tuning function, $f(\theta)$, and a noise term, ε, describing trial-to-trial variability:

$$r_n = f_n(\theta) + \varepsilon_n(\theta)$$

Noise is correlated across the population, generated from a multivariate Gaussian distribution with zero mean. Orientation tuning functions were defined as:

$$f_n(\theta) = \alpha_n + \beta_n e^{\kappa_n \left[\cos(\theta - \phi_n)^2 - 1 \right]}$$

Here, α is the baseline firing rate, β scales the tuned response, κ scales the tuning bandwidth, and ϕ is the orientation preference of each neuron. For homogeneous P_{IN} all parameters except ϕ were fixed: $(\alpha, \beta, \kappa) = (0, 5, 4)$. For heterogenous P_{IN}, we sampled parameters to match measurements from macaque V1 (Ringach et al., 2002). Tuning bandwidth was generated by converting half-width at $1/\sqrt{2}$ height (γ) values from a lognormal distribution ($\mu = -1, \sigma = 0.3$):

$$\kappa = -\log\left(\sqrt{2}\right) / \left(\cos\left(\gamma^2\right) - 1\right)$$

Limited-range correlations were included so neural noise correlation depends on tuning preference difference (Ecker et al., 2011). A correlation matrix, C, was specified by the difference between preferred orientations of neurons and the maximum pairwise correlation, c_o:

$$A_{ij} = c_o e^{-\left| \delta(\phi_i - \phi_j) \right|}$$

where δ is the circular difference and

$$C_{ij} = A_{ij} + (1 - c_o)I$$

where I is the identity matrix of size N. We scaled the correlation matrix by the mean firing rate of each neuron to produce Poisson-like noise (Ecker et al., 2011).

Derived weights for a given P_{IN} were artificially smoothed using the following equation (Park and Pillow, 2011):

$$S_{ij} \mathrel{+}= e^{-\left(\rho_1 + \left(\frac{\delta(\phi_i - \phi_j)}{\rho_2} \right) \right)}$$

Here, S^+ is the pseudoinverse of S, $\delta(\phi_i - \phi_j)$ is the difference between preferred orientations of neurons, ρ_1 scales the amplitude of smoothing, and ρ_2 scales functional range of smoothing.

Population decoder estimation accuracy

Decoding accuracy was calculated with the mean-squared-error of the maximum a posterior probability estimate across t simulated trials of each stimulus (k):

$$error(k) = \left(\frac{1}{t}\right) \sum_{1}^{t} arg\left(e^{i(MAP(w_k) - \theta_k)}\right)^2$$

Here, w_k are the weights for a given decoder neuron and θ_k is the true stimulus.

Viral Injections

Briefly, female ferrets aged P18-23 (Marshall Farms) were anesthetized with isoflurane (delivered in O2). Atropine was administered and a 1:1 mixture of lidocaine and bupivacaine was administered SQ. Animals were maintained at an internal temperature of 37 degrees Celsius. Under sterile surgical conditions, a small craniotomy (0.8 mm diameter) was made over the visual cortex (7-8mm lateral and 2-3mm anterior to lambda). A mixture of diluted AAV1.hSyn.Cre (1:25000 to 1:50000) and AAV1.Syn.FLEX.GCaMP6s (UPenn) was injected (125 - 202.5 nL) through beveled glass micropipettes (10-15 micron outer diameter) at 600, 400, and 200 microns below the pia. Finally, the craniotomy was filled with sterile agarose (Type IIIa, Sigma-Aldrich) and the incision site was sutured.

Cranial Window

After 3-5 weeks of expression, ferrets were anesthetized with 50mg/kg ketamine and isoflurane. Atropine and bupivacaine were administered, animals were placed on a feedback-controlled heating pad to maintain an internal temperature of 37 degrees Celsius, and intubated to be artificially respirated. Isoflurane was delivered throughout the surgical procedure to maintain a surgical plane of anesthesia. An intravenous cannula was placed to deliver fluids. Tidal CO2, external temperature, and internal temperature were continuously monitored. The scalp was retracted and a custom titanium headplate adhered to the skull (Metabond, Parkell). A craniotomy was performed and the dura retracted to reveal the cortex. One piece of custom cover-glass (3mm diameter, 0.7mm thickness, Warner Instruments) adhered using optical adhesive (71, Norland

Products) to custom insert was placed onto the brain to dampen biological motion during imaging. A 1:1 mixture of tropicamide ophthalmic solution (Akorn) and phenylephrine hydrochloride ophthalmic solution (Akorn) was applied to both eyes to dilate the pupils and retract the nictating membranes. Contact lenses were inserted to protect the eyes. Upon completion of the surgical procedure, isoflurane was gradually reduced and pancuronium (2 mg/kg/hr) was delivered IV.

Visual Stimuli

Visual stimuli were generated using Psychopy (Peirce, 2007). The monitor was placed 25 cm from the animal. Receptive field locations for each cell were hand mapped and the spatial frequency optimized (range: 0.04 to 0.25 cpd). For each soma and dendritic segment, square-wave drifting gratings were presented at 22.5 degree increments (2 second duration, 1 second ISI, 8-10 trials for each field of view).

Two photon imaging

Two photon imaging was performed on a Bergamo II microscope (Thorlabs) running Scanimage (Pologruto et al., 2003) (Vidrio Technologies) with 940nm dispersion-compensated excitation provided by an Insight DS+ (Spectraphysics). For spine and axon imaging, power after the objective was limited to < 50 mW. Cells were selected for imaging on the basis of their position relative to large blood vessels, responsiveness to visual stimulation, and lack of prolonged calcium transients resulting from over-expression of GCaMP6s. Images were collected at 30 Hz using bidirectional scanning with 512x512 pixel resolution or with custom ROIs (frame rate range: 22 - 50 Hz). Somatic imaging was performed with a resolution of 2.05 - 10.24 pixels/micron. Dendritic spine imaging was performed with a resolution of 6.10 -15.36 pixels/micron.

Two Photon Imaging Analysis

Imaging data were excluded from analysis if motion along the z-axis was detected. Dendrite images were corrected for in-plane motion via a 2D cross-correlation based approach in MATLAB or using a piecewise non-rigid motion correction algorithm (Pnevmatikakis and Giovannucci, 2017). ROIs (region of interest) were drawn in ImageJ; dendritic ROIs spanned contiguous dendritic segments and spine ROIs were fit with custom software. Mean pixel values for ROIs were computed over the imaging time series and imported into MATLAB (Hiner et al.,

2017; Sage et al., 2012). $\Delta F/F_o$ was computed by computing F_o with time-averaged median or percentile filter (10th percentile). For spine signals, we subtracted a scaled version of the dendritic signal to remove back-propagating action potentials as performed previously (Wilson et al., 2016). $\Delta F/F_o$ traces were synchronized to stimulus triggers sent from Psychopy and collected by Spike2. Spines were included for analysis if the SNR of the preferred response exceeded 2 median absolute deviations above the baseline noise (measured during the blank) and were weakly correlated with the dendritic signal (Spearman's correlation, $r < 0.4$). Some spine traces contained negative events after subtraction, so correlations were computed ignoring negative values. We then normalized each spine's responses so that each spine had equal weight. Preferred orientation for each spine was calculated by fitting responses with a Gaussian tuning curve using lsqcurvefit (Matlab). Tuning selectivity was measured as the vector strength index (v) for each neuron's response:

$$v_i = \frac{\sqrt{\sum (r_i cos\theta_k)^2 + \sum (r_i sin\theta_k)^2}}{\sum r_i}$$

Here r is the mean responses over the orientations (θ_k) presented for each spine (i). Note, this same index is used to characterize simulated input selectivity.

Analysis

To compare input tuning (derived synaptic population or measured dendritic spine population) with output tuning (downstream readout or measured somatic tuning) we computed the Pearson Correlation coefficient (Matlab). This correlation was computed on trial-averaged responses across different orientations. For dendritic spines and soma, measured responses across stimulus presentation trials were averaged. For simulated synaptic populations and corresponding downstream readout neuron, we simulated trials by adding noise to each synaptic tuning curve.

Code Availability

Matlab code to generate input and readout populations used are provided: https://github.com/schollben/SpineProbablisticModel2020

Bibliography

Averbeck, B.B., Latham, P.E., Pouget, A., 2006. Neural correlations, population coding and computation. Nat. Rev. Neurosci. 7, 358–366. doi:10.1038/nrn1888

Avitan, L., Goodhill, G.J., 2018. Code under construction: neural coding over development. Trends Neurosci. 41, 599–609. doi:10.1016/j.tins.2018.05.011

Berens, P., Ecker, A.S., Gerwinn, S., Tolias, A.S., Bethge, M., 2011. Reassessing optimal neural population codes with neurometric functions. Proc. Natl. Acad. Sci. USA 108, 4423–4428. doi:10.1073/pnas.1015904108

Butts, D.A., Goldman, M.S., 2006. Tuning curves, neuronal variability, and sensory coding. PLoS Biol. 4, e92. doi:10.1371/journal.pbio.0040092

Chalk, M., Masset, P., Deneve, S., Gutkin, B., 2017. Sensory noise predicts divisive reshaping of receptive fields. PLoS Comput. Biol. 13, e1005582. doi:10.1371/journal.pcbi.1005582

Chen, T.-W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., Looger, L.L., Svoboda, K., Kim, D.S., 2013. Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature 499, 295–300. doi:10.1038/nature12354

Chen, X., Leischner, U., Rochefort, N.L., Nelken, I., Konnerth, A., 2011. Functional mapping of single spines in cortical neurons in vivo. Nature 475, 501–505. doi:10.1038/nature10193

Cossell, L., Iacaruso, M.F., Muir, D.R., Houlton, R., Sader, E.N., Ko, H., Hofer, S.B., Mrsic-Flogel, T.D., 2015. Functional organization of excitatory synaptic strength in primary visual cortex. Nature 518, 399–403. doi:10.1038/nature14182

Dayan, P., Abbott, L.F., 2001. Theoretical neuroscience: computational and mathematical modeling of neural systems.

Deneve, S., Latham, P.E., Pouget, A., 1999. Reading population codes: a neural implementation of ideal observers. Nat. Neurosci. 2, 740–745. doi:10.1038/11205

DiCarlo, J.J., Zoccolan, D., Rust, N.C., 2012. How does the brain solve visual object recognition? Neuron 73, 415–434. doi:10.1016/j.neuron.2012.01.010

Ecker, A.S., Berens, P., Tolias, A.S., Bethge, M., 2011. The effect of noise correlations in populations of diversely tuned neurons. J. Neurosci. 31, 14272–14283. doi:10.1523/JNEUROSCI.2539-11.2011

Goris, R.L.T., Simoncelli, E.P., Movshon, J.A., 2015. Origin and function of tuning diversity in macaque visual cortex. Neuron 88, 819–831. doi:10.1016/j.neuron.2015.10.009

Graf, A.B.A., Kohn, A., Jazayeri, M., Movshon, J.A., 2011. Decoding the activity of neuronal populations in macaque primary visual cortex. Nat. Neurosci. 14, 239–245. doi:10.1038/nn.2733

Hiner, M.C., Rueden, C.T., Eliceiri, K.W., 2017. ImageJ-MATLAB: a bidirectional framework for scientific image analysis interoperability. Bioinformatics 33, 629–630. doi:10.1093/bioinformatics/btw681

Hofer, S.B., Ko, H., Pichler, B., Vogelstein, J., Ros, H., Zeng, H., Lein, E., Lesica, N.A., Mrsic-Flogel, T.D., 2011. Differential connectivity and response dynamics of excitatory and inhibitory neurons in visual cortex. Nat. Neurosci. 14, 1045–1052. doi:10.1038/nn.2876

Holtmaat, A., Svoboda, K., 2009. Experience-dependent structural synaptic plasticity in the mammalian brain. Nat. Rev. Neurosci. 10, 647–658. doi:10.1038/nrn2699

Hubel, D.H., Wiesel, T.N., 1962. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. J. Physiol. (Lond.) 160, 106–154. doi:10.1113/jphysiol.1962.sp006837

Iacaruso, M.F., Gasler, I.T., Hofer, S.B., 2017. Synaptic organization of visual space in primary visual cortex. Nature 547, 449–452. doi:10.1038/nature23019

Jazayeri, M., Movshon, J.A., 2006. Optimal representation of sensory information by neural populations. Nat. Neurosci. 9, 690–696. doi:10.1038/nn1691

Jia, H., Rochefort, N.L., Chen, X., Konnerth, A., 2010. Dendritic organization of sensory input to cortical neurons in vivo. Nature 464, 1307–1312. doi:10.1038/nature08947

Jia, H., Rochefort, N.L., Chen, X., Konnerth, A., 2011. In vivo two-photon imaging of sensory-evoked dendritic calcium signals in cortical neurons. Nat. Protoc. 6, 28–35. doi:10.1038/nprot.2010.169

Keller, A.J., Martin, K.A.C., 2015. Local Circuits for Contrast Normalization and Adaptation Investigated with Two-Photon Imaging in Cat Primary Visual Cortex. J. Neurosci. 35, 10078–10087. doi:10.1523/JNEUROSCI.0906-15.2015

Kerlin, A., Boaz, M., Flickinger, D., MacLennan, B.J., Dean, M.B., Davis, C., Spruston, N., Svoboda, K., 2019. Functional clustering of dendritic activity during decision-making. Elife 8. doi:10.7554/eLife.46966

Kerlin, A.M., Andermann, M.L., Berezovskii, V.K., Reid, R.C., 2010. Broadly tuned response properties of diverse inhibitory neuron subtypes in mouse visual cortex. Neuron 67, 858–871. doi:10.1016/j.neuron.2010.08.002

Ko, H., Hofer, S.B., Pichler, B., Buchanan, K.A., Sjöström, P.J., Mrsic-Flogel, T.D., 2011. Functional specificity of local synaptic connections in neocortical networks. Nature 473, 87–91. doi:10.1038/nature09880

Kohn, A., Coen-Cagli, R., Kanitscheider, I., Pouget, A., 2016. Correlations and neuronal population information. Annu. Rev. Neurosci. 39, 237–256. doi:10.1146/annurev-neuro-070815-013851

Lee, W.-C.A., Bonin, V., Reed, M., Graham, B.J., Hood, G., Glattfelder, K., Reid, R.C., 2016. Anatomy and function of an excitatory network in the visual cortex. Nature 532, 370–374. doi:10.1038/nature17192

Ma, W.J., Beck, J.M., Latham, P.E., Pouget, A., 2006. Bayesian inference with probabilistic population codes. Nat. Neurosci. 9, 1432–1438. doi:10.1038/nn1790

Movshon, J.A., Thompson, I.D., Tolhurst, D.J., 1978. Receptive field organization of complex cells in the cat's striate cortex. J. Physiol. (Lond.) 283, 79–99. doi:10.1113/jphysiol.1978.sp012489

Pagan, M., Simoncelli, E.P., Rust, N.C., 2016. Neural Quadratic Discriminant Analysis: Nonlinear Decoding with V1-Like Computation. Neural Comput. 28, 2291–2319.

doi:10.1162/NECO_a_00890

Park, M., Pillow, J.W., 2011. Receptive field inference with localized priors. PLoS Comput. Biol. 7, e1002219. doi:10.1371/journal.pcbi.1002219

Peirce, J.W., 2007. PsychoPy--Psychophysics software in Python. J. Neurosci. Methods 162, 8–13. doi:10.1016/j.jneumeth.2006.11.017

Pnevmatikakis, E.A., Giovannucci, A., 2017. NoRMCorre: An online algorithm for piecewise rigid motion correction of calcium imaging data. J. Neurosci. Methods 291, 83–94. doi:10.1016/j.jneumeth.2017.07.031

Pologruto, T.A., Sabatini, B.L., Svoboda, K., 2003. ScanImage: flexible software for operating laser scanning microscopes. Biomed Eng Online 2, 13. doi:10.1186/1475-925X-2-13

Pouget, A., Dayan, P., Zemel, R., 2000. Information processing with population codes. Nat. Rev. Neurosci. 1, 125–132. doi:10.1038/35039062

Reid, R.C., Alonso, J.M., 1995. Specificity of monosynaptic connections from thalamus to visual cortex. Nature 378, 281–284. doi:10.1038/378281a0

Ringach, D.L., Shapley, R.M., Hawken, M.J., 2002. Orientation selectivity in macaque V1: diversity and laminar dependence. J. Neurosci. 22, 5639–5651. doi:20026567

Rumyantsev, O.I., Lecoq, J.A., Hernandez, O., Zhang, Y., Savall, J., Chrapkiewicz, R., Li, J., Zeng, H., Ganguli, S., Schnitzer, M.J., 2020. Fundamental bounds on the fidelity of sensory cortical coding. Nature 580, 100–105. doi:10.1038/s41586-020-2130-2

Runyan, C.A., Schummers, J., Van Wart, A., Kuhlman, S.J., Wilson, N.R., Huang, Z.J., Sur, M., 2010. Response features of parvalbumin-expressing interneurons suggest precise roles for subtypes of inhibition in visual cortex. Neuron 67, 847–857. doi:10.1016/j.neuron.2010.08.006

Rust, N.C., Mante, V., Simoncelli, E.P., Movshon, J.A., 2006. How MT cells analyze the motion of visual patterns. Nat. Neurosci. 9, 1421–1431. doi:10.1038/nn1786

Sage, D., Prodanov, D., Tinevez, J.Y., 2012. MIJ: making interoperability between ImageJ and Matlab possible. bigwww.epfl.ch.

Scholl, B., Fitzpatrick, D., 2020. Cortical synaptic architecture supports flexible sensory computations. Curr. Opin. Neurobiol. 64, 41–45. doi:10.1016/j.conb.2020.01.013

Scholl, B., Pattadkal, J.J., Dilly, G.A., Priebe, N.J., Zemelman, B.V., 2015. Local integration accounts for weak selectivity of mouse neocortical parvalbumin interneurons. Neuron 87, 424–436. doi:10.1016/j.neuron.2015.06.030

Scholl, B., Thomas, C.I., Ryan, M.A., Kamasawa, N., Fitzpatrick, D., 2019a. Cortical neuron response selectivity derives from strength in numbers of synapses. BioRxiv. doi:10.1101/2019.12.24.887422

Scholl, B., Wilson, D.E., Fitzpatrick, D., 2017. Local Order within Global Disorder: Synaptic Architecture of Visual Space. Neuron 96, 1127–1138.e4. doi:10.1016/j.neuron.2017.10.017

Scholl, B., Wilson, D.E., Jaepel, J., Fitzpatrick, D., 2019b. Functional logic of layer 2/3 inhibitory connectivity in the ferret visual cortex. Neuron 104, 451–457.e3.

doi:10.1016/j.neuron.2019.08.004

Seriès, P., Latham, P.E., Pouget, A., 2004. Tuning curve sharpening for orientation selectivity: coding efficiency and the impact of correlations. Nat. Neurosci. 7, 1129–1135. doi:10.1038/nn1321

Seung, H.S., Sompolinsky, H., 1993. Simple models for reading neuronal population codes. Proc. Natl. Acad. Sci. USA 90, 10749–10753. doi:10.1073/pnas.90.22.10749

Shamir, M., 2014. Emerging principles of population coding: in search for the neural code. Curr. Opin. Neurobiol. 25, 140–148. doi:10.1016/j.conb.2014.01.002

Shamir, M., Sompolinsky, H., 2006. Implications of neuronal diversity on population coding. Neural Comput. 18, 1951–1986. doi:10.1162/neco.2006.18.8.1951

Spitzer, H., Hochstein, S., 1988. Complex-cell receptive field models. Prog. Neurobiol.

Stringer, C., Michaelos, M., Pachitariu, M., 2019. High precision coding in visual cortex. BioRxiv.

Ujfalussy, B.B., Makara, J.K., Lengyel, M., Branco, T., 2018. Global and Multiplexed Dendritic Computations under In Vivo-like Conditions. Neuron 100, 579–592.e5. doi:10.1016/j.neuron.2018.08.032

Wertz, A., Trenholm, S., Yonehara, K., Hillier, D., Raics, Z., Leinweber, M., Szalay, G., Ghanem, A., Keller, G., Rózsa, B., Conzelmann, K.-K., Roska, B., 2015. PRESYNAPTIC NETWORKS. Single-cell-initiated monosynaptic tracing reveals layer-specific cortical network modules. Science 349, 70–74. doi:10.1126/science.aab1687

Wilson, D.E., Scholl, B., Fitzpatrick, D., 2018. Differential tuning of excitation and inhibition shapes direction selectivity in ferret visual cortex. Nature 560, 97–101. doi:10.1038/s41586-018-0354-1

Wilson, D.E., Smith, G.B., Jacob, A.L., Walker, T., Dimidschstein, J., Fishell, G., Fitzpatrick, D., 2017. Gabaergic neurons in ferret visual cortex participate in functionally specific networks. Neuron 93, 1058–1065.e4. doi:10.1016/j.neuron.2017.02.035

Wilson, D.E., Whitney, D.E., Scholl, B., Fitzpatrick, D., 2016. Orientation selectivity and the functional clustering of synaptic inputs in primary visual cortex. Nat. Neurosci. 19, 1003–1009. doi:10.1038/nn.4323

Zavitz, E., Price, N.S.C., 2019. Weighting neurons by selectivity produces near-optimal population codes. J. Neurophysiol. 121, 1924–1937. doi:10.1152/jn.00504.2018

Zylberberg, J., 2017. Untuned but not irrelevant: A role for untuned neurons in sensory information coding. BioRxiv. doi:10.1101/134379

firing rate

input population (P_{IN})

orientation space (Θ)

weight amplitudes

excitatory synaptic
population (P_{SYN})

weights (*w*)

orientation space (Θ)

Σ

output population (*r*)
(decoder neurons)

a Example Weights

$\rho_1 = 0$
$\rho_2 = 0$

$\rho_1 = 0.1$
$\rho_2 = 1$

$\rho_1 = 0.2$
$\rho_2 = 2$

$\rho_1 = 1$
$\rho_2 = 10$

b Homogenous — Hetereogenous — $c_o = 0$

c $c_o = 0.25$

d $c_o = 0.50$

a

b

Simulated Synaptic Population

a Two-Photon Spine Data

b Heterogenous P_{IN}

a

Hetereogenous P_{IN}