High-throughput transcriptomic analysis of human primary hepatocyte spheroids

exposed to per- and polyfluoroalkyl substances (PFAS) as a platform for relative

potency characterization

A. Rowan-Carroll[1], A. Reardon[1], K. Leingartner[1], R. Gagné[1], A. Williams[1], M.J. Meier[1],

B. Kuo [1], J. Bourdon-Lacombe[2], I. Moffat[2], R. Carrier[2], A. Nong[1], L. Lorusso[3], S.S.

Ferguson[4], E. Atlas[1 *], C. Yauk[5,1*]

[1]Environmental Health Science and Research Bureau, Healthy Environments and

Consumer Safety Branch (HECSB), Health Canada, [2]Water and Air Quality Bureau,

HECSB, Health Canada, [3]Chemicals and Environmental Health Management Bureau,

HECSB, Health Canada, [4]US National Institute of Environmental Health Sciences,

[5]Dept. of Biology, University of Ottawa.

* To Whom correspondence should be addressed: carole.yauk@uottawa.ca and

ella.atlas@canada.ca

Keywords: TempO-Seq, PFAS, liver spheroids, benchmark concentration, new

approach methodology, bioactivity exposure ratio

This entire data set is publicly available at: at
https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE144775

Abstract

Per- and poly-fluoroalkyl substances (PFAS) are widely found in the environment because of

their extensive use and persistence. Although several PFAS are well studied, most lack

toxicity data to inform human health hazard and risk assessment. This study focussed on four

model PFAS: perfluorooctanoic acid (PFOA; 8 carbon), perfluorobutane sulfonate (PFBS; 4

carbon), perfluorooctane sulfonate (PFOS; 8 carbon), and perfluorodecane sulfonate (PFDS;

10 carbon). Human primary liver cell spheroids (pooled from 10 donors) were exposed to 10

concentrations of each PFAS and analyzed at four time-points. The approach aimed to: (1)

identify gene expression changes mediated by the PFAS; (2) identify similarities in biological

responses; (3) compare PFAS potency through benchmark concentration analysis; and (4)

derive bioactivity exposure ratios (ratio of the concentration at which biological responses

occur, relative to daily human exposure). All PFAS induced transcriptional changes in

cholesterol biosynthesis and lipid metabolism pathways, and predicted PPARα activation.

PFOS exhibited the most transcriptional activity and had a highly similar gene expression

profile to PFDS. PFBS induced the least transcriptional changes and the highest benchmark

concentration (i.e., was the least potent). The data indicate that these PFAS may have

common molecular targets and toxicities, but that PFOS and PFDS are the most similar. The

transcriptomic bioactivity exposure ratios derived here for PFOA and PFOS were comparable

to those derived using rodent apical endpoints in risk assessments. These data provide a

baseline level of toxicity for comparison with other known PFAS using this testing strategy.

Introduction

Per- and poly-fluoroalkyl substances (PFAS) are a class of chemicals ubiquitously found in the environment. Their unique hydrophobic and lipophobic properties have led to their use in many consumer products and their persistence, and bioaccumulation potential, led to their detection in wildlife and humans worldwide (Giesy and Kanna 2001; Armitage *et al.*, 2006; 2009a, Wania, 2007). Although perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) have been discontinued from North American production, they continue to be found in drinking water and at contaminated sites within Canada (Health Canada., 2018a and b). Moreover, there are thousands of PFAS that remain in production, with little toxicological data available for the vast majority (OECD/UNEP Global PFC Group, 2018).

The adverse health effects of PFOS and PFOA in humans are well documented, including increased cholesterol levels (Erikson *et al.*, 2013; Geiger *et al.*, 2014), decreased human fertility (Bach *et al.*, 2016), and reduced immune response (NTP Monograph, 2016; Kielsen *et al.*, 2016; Health Canada, 2018a and b). Recently, the US Environmental Protection Agency developed an action plan to address PFAS in drinking water, general exposure reduction, safety in commerce, and support for communications (EPA, 2020).

The effects of several PFAS have been extensively studied in animal models. For example, wild-type and PPARα-null mice treated with several PFAS exhibit steatosis, which is characterised by an accumulation of lipids in hepatocytes, leading to liver disease (Das *et al.*, 2017). Furthermore, rats dosed with PFOS (up to 1.14 mg/kg/day)

and offspring of mice dosed with PFOA (up to 5 mg/kg/day) developed liver cancer

(Bjork *et al.*, 2011; Filgo *et al.*, 2015).

Increased exposure to PFAS has been associated with adverse liver outcomes

and the development of steatosis. Using NHANES data, increased PFAS exposure was

associated with alterations in select liver biomarkers, suggesting they play a role in the

progression of non-alcoholic fatty liver disease (Jain and Ducatman, 2019). Both *in vitro*

and *in vivo* studies suggest that the mechanism by which these compounds act involves

increased intracellular triglycerides levels and activation of nuclear receptors (Louisse *et*

al. 2020, Schlezinger *et al.,* 2020). Given these noted adverse liver effects, the liver was

identified as the critical epidemiological endpoint in Health Canada's existing regulatory

guidelines on PFAS (Health Canada, 2018a and b). In humans, median serum levels

can reach 80 ng/ml for PFOA and 22 ng/ml for PFOS and these exposure levels are

correlated with increased blood lipid levels (Steenland *et al.*, 2009). Conversely, a

review from Chang et al. (2014) found no apparent causal association with cancer in

human occupational exposures to PFOA that are an order of magnitude higher (serum

levels ranging from 160 to 2880 ng/ml) than environmental exposures.

Acquiring information on data-poor substances for risk assessment has been a

challenge for regulatory agencies worldwide given the cost and time required to conduct

traditional toxicological research. To accelerate the pace of chemical risk assessments,

global efforts have focused on increasing the use of New Approach Methodologies

(NAMs) (Kavlock *et al.*, 2018; Krewski *et al.*, 2020). *In silico* predictions, high-throughput

screening, novel *in vitro* models, *in vitro* to *in vivo* extrapolation (IVIVE), and read-

across are some of the NAMs with increasing use in risk assessment for chemicals with

limited data. High-throughput transcriptomics is a particularly powerful method for

obtaining information about the broad scope of biological perturbations that result from

chemical exposures. New technologies now enable gene expression analyses directly

from cell lysates, which significantly increases the ease and speed with which

transcriptomic data can be produced (Trejo *et al.*, 2019).

To address the need for efficient assessment of data-poor PFAS, we sought to

explore transcriptional responses across exposure time with a broad concentration

range in order to refine a streamlined set of exposure conditions for PFAS evaluations.

We developed an approach to broadly screen for PFAS-induced biological perturbations

through global gene expression profiling of a primary human liver cell spheroid model. A

main objective was to establish baseline transcriptomic profiles and an analytical

pipeline for future evaluations of data-poor PFAS. Spheroids were exposed to nine

concentrations of PFAS and sampled at four time points. To establish our approach, we

focused on four prototype PFAS: short-chain perfluorobutane sulfonate (PFBS), long-

chain perfluorodecane sulfonate (PFDS), and PFOA and PFOS. Microscopy was used

to evaluate phenotypic changes in the spheroids, and the lactose dehydrogenase

activity assay was used to assess cytotoxicity over time. The TempO-Seq platform from

BioSpyder using the NIEHS S1500 gene set was applied for transcriptomic analysis

(House *et al.*, 2017; Mav *et al.*, 2018). PFAS were compared based on the induction of

differentially expressed genes, transcriptional patterns measured by hierarchical

clustering and principal component analysis, pathway enrichment, upstream regulator

enrichment, and transcriptomic benchmark concentrations. Bioactivity-exposure ratios

for PFOS and PFOA were calculated by comparing human administered equivalent

dose (the daily dose required to achieve the *in vitro* concentration of interest) to human

exposure levels. A small bioactivity exposure ratio suggests that biological perturbations

occur at concentrations that approach external human exposure levels. This initial

research project was used to inform experimental design and analytical approach for

analysis of a second larger group of data-poor PFAS (Reardon *et al.*, BioRxiv

doi: https://doi.org/10.1101/2020.10.20.347328).

Methods

Chemicals and preparation of working solutions

PFOS (95% purity CAS 1763-23-1), PFOA (95% purity CAS 335-67-1), PFBS

(95% purity CAS 375-73-5) and Cyclosporine A (CsA) were purchased from Sigma-

Aldrich (Oakville, ON) and PFDS was purchased from Toronto Research Chemicals

(95% purity CAS 335-77-3 Toronto, ON). Chemicals were dissolved in dimethylsulfoxide

(DMSO, Sigma-Aldrich, Oakville, ON) to prepare working stock solutions ranging in

concentration from 0.02 to 30 mM. Each PFAS was tested at 10 different concentrations

(0 to 100 µM), in quadruplicate experiments. For microscopic characterization, a 30 mM

working stock of CsA in DMSO was also prepared. The final concentration of DMSO in

the media was 0.1% for all PFAS at all concentrations up to 50 µM; the DMSO was

0.17% for 50 µM and 0.3% for 100 µM. The %DMSO varied at the two top

concentrations because chemical stocks were diluted to 30 mM for all PFAS due to

solubility. All experimental concentrations of PFAS were matched to vehicle only

(DMSO) time-matched and concentration-matched controls; e.g., PFAS at 100 µM

(0.3% DMSO) was matched to 0.3% DMSO controls from the same time point.

Cell Culture

3D InSight™ Human Liver Microtissues were purchased from InSphero (Brunswick, ME) in a 96 well format, with a single spheroid per well. These spheroids are a co-culture model pooled from 10 different human liver donors and are a metabolically active system comprised of isolated primary hepatocytes and Kupffer cells. Upon arrival, the media were replaced with serum free cell medium (InSphero Human Liver Maintenance Medium–Tox) and spheroids were allowed to acclimate for 24 hours prior to PFAS exposures. The Human Liver Maintenance Medium–Tox contains no additives or serum and is designed for testing of drug induced liver injury (DILI) compounds. The media contains no proteins or albumin; however, the spheroids themselves do secrete albumin into the media.

Liver spheroids were cultured under sterile conditions with incubation at 37 $^\circ$C, 5% CO_2 as per the materials and instructions provided by the manufacturer. Liver spheroids were exposed for 1, 4, 10 or 14 days to PFBS, PFOA, PFOS, PFDS, or DMSO control diluted in culture medium. Final exposure concentrations were 0.02, 0.1, 0.2, 1, 2, 10, 20, 50, and 100 µM of each of the four PFAS chemicals. The entire experiment included eight 96-well plates of spheroids. Each plate contained every concentration of each PFAS alongside a matched DMSO control for every sample. On each plate there were also 18 DMSO controls: four 0.3% DMSO (for 100 µM concentrations), four 0.17% DMSO (for 50 µM concentrations), and ten 0.1% DMSO (for all of the remaining concentrations). There were two 96-well plates for every time point. PFAS or DMSO, at the desired concentrations, were replenished each time the media were changed (every 3 days).

The highest concentration tested was 100 µM, as per the US EPA's ToxCast

program. Although the top concentration is high, it is useful to have a top concentration

yielding a robust transcriptomic response for benchmark concentration modeling and

this provides important context analogous to LD_{50} for guideline toxicology studies. From

the top concentration we scaled down in total by 3 logs to the lowest, most

environmentally relevant concentration (Health Canada., 2017; 2018a and b). Media

were replaced every 3 days with freshly diluted chemical, and spent media were

reserved for cytotoxicity analysis. Spheroids designated for microscopy were treated

with 0.1% DMSO; 48 hours prior to the day-14 time point half of these spheroids were

exposed to 30 µM CsA for immunostaining. Spheroids designated for TempO-Seq

library building and sequencing were washed with Dulbecco's phosphate-buffered saline

(DPBS; Thermo Fisher Scientific Waltham, MA) and lysed using 5-7 µl of 2X TempO-

Seq lysis buffer (BioSpyder Technologies Inc Carlsbad, CA). Liver spheroids were

triturated and allowed to complete lysis for approximately 10 minutes before freezing at

-80 $^\circ$C. Spheroids designated for TempO-Seq library building and sequencing analysis

were collected after exposure on day-1, day-4, day-10, and day-14.

Cytotoxicity Assessment

Cytotoxicity was measured from spent culture media collected every 3 days

during media replenishment (i.e., on days 3, 6, 9, 12, and then on day-14 at study

termination, versus days 1, 4, 10, and 14 for transcriptomic analyses below) using the

lactate dehydrogenase assay (LDH-Glo Cytotoxicity Assay Promega Corp Madison,

USA), as per manufacturer's instructions. Briefly, media were diluted 1:10 in storage

buffer (200 mM Tris-HCl, pH 7.3, containing 10% glycerol and 1% bovine serum

albumin) and kept frozen at -80 $^\circ$C until analysis. For the assay, thawed samples were

diluted with an equal volume of LDH Detection Reagent and equilibrated for 60 minutes

at room temperature before reading the luminescence (GloMax 96 Microplate

Luminometer, Promega Corp Madison, WI USA) in relative luminescence units (RLU).

Cytotoxicity was determined after performing statistical analysis of the RLU data after

blank subtraction in Sigma Plot 13™, using the Kruskal-Wallis one-way analysis of

variance on ranks, followed by the Dunn's multiple comparison versus the control group

for each time point, with n = 4 for all time points, with exceptions on Day-14 (n = 2 in

some cases as indicated). The sample RLU was converted to a ratio by division with the

average RLU of the DMSO controls and the mean and standard deviations of these

ratios were calculated for each treatment and time point. The release of LDH was

expressed as a fold-change over DMSO controls. Overall cytotoxicity was determined

by combining three metrics of cell viability: LDH release greater than 10-fold over the

control, low read counts from Next Generation Sequencing (described further below),

and visual examination of cell morphology.

Microscopy

Spheroid whole mounts were prepared on Day-4 and Day-14 according to a

protocol provided by InSphero Inc. (Technical Protocol 008 c. 2015), with modifications.

Briefly, spheroids were fixed with 4% paraformaldehyde in phosphate buffered saline

(PBS) for 1 hour, permeabilized with 0.2% Triton-X100 in PBS for 30 minutes, and

blocked with 5% bovine serum albumin in PBS for 1 hour at room temperature. On day-

4, the spheroids were incubated with CD-68 antibody (1:100; Santa Cruz CD68 (E-11):

sc-17832 conjugated to AlexaFluor 594) overnight at 4 °C, followed by incubation with

1:40 phalloidin (Life Technologies 12379 conjugated to AlexaFluor 488 Carlsbad, CA)

and DAPI (2 µg/ml; Sigma D8417 Oakville, ON) in PBS for 45 minutes at room

temperature. This was performed to ensure spheroid integrity and cell composition.

On Day-14, spheroids were stained with CYP3A4 primary antibody (1:00; Santa

Cruz CYP3A4 (HL3): sc-53850 conjugated to AlexaFluor 647 Dallas, TX) phalloidin and

DAPI as described above, or with Nile Red (1 µg/ml; Sigma 19123 Oakville, ON) and

DAPI (2 µg/ml; Sigma D8417 Oakville, ON) in PBS for 30 minutes at room temperature.

After antibody incubation and three washes with PBS, spheroids were transferred to

glass microscope slides and mounted with Prolong Gold Mounting Medium (Invitrogen

P36934, Burlington, ON). Confocal laser-scanning immunofluorescence microscopy

was performed using a Leica TCS SP8 microscope.

TempO-Seq Library Building and Next Generation Sequencing

Gene expression was measured using the human TempO-Seq S1500 panel

(BioSpyder Technologies Inc. Carlsbad, CA) (Mav *et al.*, 2018). This targeted RNA-seq

panel is comprised of 3000 genes selected by the NIEHS to represent a biologically diverse

set of pathways that are toxicologically responsive (https://federalregister.gov/a/2015-

08529). For TempO-Seq analysis, liver spheroids were exposed to nine concentrations of

the four PFAS and solvent control (DMSO), over the four time points (days 1, 4, 10, and 14).

Spheroids were lysed *in situ* using 2 x TempO-Seq lysis buffer diluted with an equal amount

of PBS. Lysates and positive technical controls (Human Universal Reference RNA - uhrRNA

Agilent Cat# 740000, Santa Clara, California and Human Brain Total RNA brRNA -

ThermoFisher AM7962, Waltham, Massachusetts), as well as no-cell negative controls (1X

TempO-Seq lysis buffer alone) were hybridized to the detector oligo mix following the

manufacturer's instructions (TEMPO-SEQ Human Tox +Surrogate with Standard

Attenuation Transcriptome Kit (96 Samples) BioSpyder Technologies, Inc. Hybridization

was followed by nuclease digestion of excess oligos, detector oligo ligation, and

amplification of the product with the tagged primers according to manufacturer instructions.

At the time of amplification, each sample was ligated to a sample-specific bar code that

allows for combining samples for sequencing purposes. Labelled amplicons were then

pooled and purified using NucleoSpin Gel and PCR Clean-up kits, (Takara Bio USA, Inc.

Mountain View, CA). Libraries were sequenced in-house using a NextSeq 500 High-

Throughput Sequencing System (Illumina San Diego, California) using 50 cycles from a

75-cycle high throughput flow cell. A median read depth of 2 million reads/sample was

achieved.

Data quality overview and removal of transcriptomic outliers

The gene expression data are available in the NCBI's Gene Expression Omnibus

(Bioproject number PRJNA604830). Initial quality assessment was done on water-only

controls and reference standards included on every TempO-Seq plate. We first

confirmed very little signal (< 0.3%) from samples with no cells/RNA. In addition, we

confirmed that commercial human universal reference RNA (two per plate – 16 in total)

showed high Spearman correlation coefficients (> 0.94) across all plate comparisons

indicating a high degree of technical reproducibility across the plates.

The median read depth for the study was two million reads with a minimum read

depth of 400,000 for retained samples. The median range of % mapped reads across

experimental space using BioSpyder's script was: day-1: 71%, day-4: 62%, day-10:

49%, and day-14: 53%. The cytotoxicity information was paired with the number of

reads recovered during the TempO-Seq experiments to identify overtly cytotoxic

concentrations for removal from the transcriptomic data analysis. The conditions

removed from subsequent analyses were: 100 µM PFOA on days 10 and 14, and

50 µM on day-14; and 50 and 100 µM PFOS from all time points.

Data processing and identification of differentially expressed genes

Reads were extracted from the bcl files and demultiplexed (i.e., assigned to

respective sample files) with bcl2fastq v. 2.20.0.42. The fastq files were processed with

the "pete. star. script_v3.0" supplied by BioSpyder. Briefly, the script uses star v.2.5 to

align the reads and the count function from QuasR to extract the feature counts

specified in a gtf file from the aligned reads.

Samples with low read counts (library sizes < 100,000 reads) were removed from

the study. As described above, read counts were also used to identify overtly cytotoxic

concentrations alongside LDH assay results as samples with low read counts were

often associated with high levels of cytotoxicity as measured in the LDH assay. Probes

with low counts were flagged as absent genes and removed (probes that did not exceed

a median of 5 counts in at least one group). Dendrogram plots (Supplemental Files1-1

through 1-9) were generated on all the remaining genes in R using the hclust () function

with complete-linkage for each time point using a distance metric defined as 1-

Spearman correlation. Samples that clustered as singletons when cutting the

dendrograms at the 0.1 dissimilarity were removed from the study. To account for

differences in the percentage of DMSO used, the data were normalized to matched

controls and rescaled to the average of the 0.1% DMSO dose group. Boxplots of the

log2 CPM (counts per million) were plotted to ensure that the samples within each

treatment group had similar medians and interquartile ranges (Supplemental Files 2

through 5). This analysis led to the removal of a total of 105 samples, such that the

sample size of the experiment was at least n = 2 (four concentrations in total had n = 2;

41 had n = 3; and 70 had n = 4) for every concentration tested (Supplemental File 6A)

with 18 DMSO controls per plate. Supplemental File 6B summarizes the number of

controls pre- and post-filtering.

Analyses to identify differentially expressed genes were conducted on probes

using raw counts with the default parameters of DESeq2 v1.30. Here, default

parameters include using Cook's distance to suppress p-values for probes that had one

or more samples deemed to be an outlier or an influential observation at the 0.99 cut-

off. Time points were analyzed independently, as data across time points were

generated from different sequencing runs. The full model used in each analysis included

a treatment effect and a blocking effect indicating the index plate. Using the full model,

pairwise comparisons were conducted for each chemical concentration combination to

the appropriate DMSO control (i.e., matched to the relevant %DMSO). Probes reaching

the threshold of a false discovery rate (FDR) adjusted p-value < 0.05 and an absolute

fold change > 1.5 (on a linear scale) were flagged (probes passing filters) and retained

for further analyses. Post-hoc collapsing of probes mapping to the same gene was

conducted to generate the table of differentially expressed genes to represent unique

genes only.

Hierarchal clustering and principal component analysis of differentially

expressed genes

Heatmaps were generated with the pheatmapfunction from R with both the row and

column distances defined as 1-Spearman correlation. Values were averaged between

replicates; CPM mean values of probes passing filters were averaged between replicates

within each group. Resulting values were subsequently log2 transformed. The principal

components were calculated with the prcompR function. The percentages of variation on the

axes were gathered from the "importance" field produced by the summary function applied

to the resulting object of the prcomp function. The work above was done with R v.3.6.1.

Further details on pipeline analysis are available in Supplemental File 7. Principal

component analysis plots were generated using the probes passing filters (Supplemental

File 8A). This led to the identify of an additional DMSO outlier. The DeSeq2 analysis was

repeated and the PCA was replotted (Supplemental File 8B; Supplemental File 6B has the

total number DMSO controls used in the study). Hierarchical clustering plots for each time

point are also included as Supplemental Files 8 C and 8D. Heatmaps were generated with

the pheatmapfunction from R with both the row and column distances defined as 1-

Spearman correlation with complete linkage. Note that the summary of the number of

replicates (n) included for the benchmark concentration analysis are the same as those

used to identify differentially expressed genes (Supplemental File 6A and B).

Functional analysis using Ingenuity Pathway Analysis

Functional analysis of the differentially expressed genes was conducted using

Ingenuity Pathway Analysis (IPA, Qiagen, Redwood City, CA, USA) using probes with

FDR-corrected p-value < 0.05, absolute fold change > 1.5 and user supplied probes for

background. For probes that mapped to the same gene, the probe with the maximum

absolute fold change was used to represent the gene. Right-tailed Fisher's exact tests

were used to calculate the probability of enrichment of biological functions, canonical

pathways, and upstream regulators. The background tab was set to "user-defined",

which allows for enrichment comparisons against the TempO-Seq panel only. The z-

score, a statistical predictor of relationship between a direct observation and a

prediction of activation/suppression of a particular pathway was set to +/- |2.0|, which is

considered a significant cut-off (Ek *et al.*, 2015).

Benchmark concentration – Gene-level analysis

Best practices as recommended by the US EPA's expert panel on genomic

benchmark dose analysis were followed (National Toxicology Program Report, 2018).

Benchmark concentration modeling was conducted using BMDExpress v.2.3, a free

software package that is available for download

(https://github.com/auerbachs/BMDExpress-2/releases). Additional information on

BMDExpress is available (https://github.com/auerbachs/BMDExpress-

2/wiki/Benchmark-Dose-Analysis). Normalized counts per million of the mapped reads

for genes were used for this analysis. Samples identified to be cytotoxic or outliers were

removed prior to benchmark concentration analysis. Data were imported into

BMDExpress and subject to pre-filtering using the Williams' Trend Test with 100

permutations and a 1.5-fold change (FC) cutoff to select genes that exhibit significant

monotonically increasing or monotonically decreasing concentration-response ($p <$

0.05). Data exhibiting an increasing or decreasing change in response were modeled

using Hill, Power, Linear, Polynomial (2° and 3°), and Exponential (2, 3, 4, 5) models to

identify potential concentration-response relationships and best fit. For linear and

polynomial models, the Nested Chi-square test was used to determine best-fit models

followed by a comparison of Akaike information criterion (AIC) for nested models (Hill

and Power) with a goodness of fit p-value of 0.1. Hill models were flagged if the 'k'

parameter was less than 1/3 the lowest (non-solvent control) concentration, and in the

case of a flagged Hill model, the next best-fit model with a goodness of fit p-value > 0.05

was selected (Hill *et al.*, 2018). Additional parameters included a benchmark response

factor of 1 standard deviation (SD) relative to the response of controls, a confidence

level of 0.95, power restricted to ≥ 1, and maximum of 250 iterations (maximum

iterations represent a convergence criterion for the model). Applied filters to best-fit

models excluded data with highly uncertain benchmark concentrations ($p < 0.1$, BMC

Upper threshold/ BMC Lower Threshold ≥ 40) as well as data producing benchmark

concentration values > the highest non-cytotoxic concentration (e.g., 100 µM). Gene

accumulation plots showing the distribution of gene-level benchmark concentrations

across the range of concentrations for best-fit models were exported from BMDExpress.

The BMD "project file (.bm2)" files are available through a link within the supplementary

information (Supplementary File 9).

Potency comparison of the PFAS was conducted using the median benchmark

concentration of all filtered genes adhering to best-fit models and a bootstrap R-script

producing 95% confidence intervals. Potencies were also compared through visual

inspection of gene accumulation plots. In addition, both the 5th percentile gene

benchmark concentration and the lowest pathway median gene benchmark

concentration were used to derive a bioactivity exposure ratio for PFOS and PFOA

(described below).

Derivation of the Bioactivity Exposure Ratio

Administered equivalent doses (AEDs) for PFOS and PFOA were estimated from

toxicokinetics information described in Wambaugh et al. (2013). Wambaugh et al.

performed simulations using a three compartmental model based on points of departure

collected from various studies to obtain the steady-state concentrations (Css). Their

population Monte Carlo simulation included a distributed oral uptake rate to consider for

variability. We chose the Css and lowest observe effect level (LOEL) values for each of

the PFOS and PFOA from non-cancer liver rodent points of departure, which were most

consistent with the toxicokinetics analyses in this manuscript.

Transcriptional benchmark concentration values were converted to administered

equivalent doses (AED) using the following equation:

$$AED = Benchmark\ Concentration\ (\mu M) \times \frac{LOEL}{Css}$$

375   For ease of calculation, the formula was reduced to:

$$AED = \frac{Benchmark\ Concentration\ (\mu M)}{Conversion\ Factor}$$

Where the $Conversion\ Factor = \frac{Css}{LOEL}$

The resulting conversion factors to compute administered equivalent dose for

each of the PFOS and PFOA are 19.47 and 46.4 mg/L per mg/kg/day, respectively.

To derive a bioactivity exposure ratio, the administered equivalent dose (mg/kg-

BW/day) were compared to the upper limit of the daily US population exposure (95th

percentile) for PFOA (4.50×10^{-04} mg/kg-BW/day) and PFOS (4.76×10^{-04} mg/kg-BW/day)

obtained from the US EPA comptox dashboard (https://comptox.epa.gov/dashboard,

accessed: January 2020). The other PFAS within the current investigation (PFBS and

PFDS) were excluded due to limited data availability.

Two approaches to bioactivity exposure ratio derivation were used. In **Approach**

1, the ratio was calculated using the lower bound 5[th] percentile administered equivalent

dose from the gene-level benchmark concentration analysis that was rounded to the

nearest gene with an integer of greater than or equal value (i.e., the ceiling) and

compared to the population exposure. This methodology has been used in previous

work to represent a lower bound estimate of *in vivo* adverse effects within the ToxCast

database in order to derive a bioactivity exposure ratio (Paul Friedman *et al.*, 2020). In

Approach 2, bioactivity exposure ratio was derived based on data demonstrating a

strong correlation between the dose at which apical effects *in vivo* occur and the lowest

(most sensitive) *in vivo* pathway benchmark concentration (Thomas *et al.*, 2013). The

pathway level analysis was conducted using the built-in Defined Category Analysis

function within BMDExpress. Pathway-level estimates were derived from genes meeting

the criteria of best-fit models mapped to probes from the IPA database (downloaded

October 2019). Estimates were removed if they contained genes with a benchmark

concentration > highest concentration from the category descriptive statistics,

benchmark concentration with a goodness of fit p-value < 0.1, or a BMCU/BMCL value

> 40. Exported pathway level estimates included additional filters limiting probe-sets to

pathways with ≥ 3 genes passing all previous filters and with ≥ 5% of pathway genes

affected. Each approach was then compared to reference administered equivalent dose

values derived from *in vivo* animal data that were used to determine drinking water

guidelines for PFOA and PFOS (Health Canada, 2018 a and b) for the purpose of risk

assessment.

Results

Human primary liver cell spheroids microscopic evaluation

410   In order to assess the composition of the liver spheroid model, approximately 25

spheroids were stained with CD68, a marker of Kupffer cells, and phalloidin to stain for

actin (Figure 1A). As anticipated, the spheroids stained positive for CD68, confirming

the presence of this native cell type that is part of the normal liver composition. Actin

staining showed the intact morphology of the spheroids and the hepatocytes within. The

spheroids were likewise treated with a known inducer of steatosis, Cyclosporine A

(CsA) for 48 hrs, and then processed for microscopic evaluation (Figure 1A). Treatment

with 30 µM CsA induced expression of CYP3A4 (Figure 1B), a member of

the cytochrome P450 oxidizing family of enzymes responsible for the oxidation of

toxicants (Zanger and Schwab, 2013), and increased lipid accumulation as assessed by

Nile Red staining (Figure 1B panel D). These results confirm that the spheroid

hepatocytes were responding as expected.

Cytotoxicity assessment

423   The LDH-Glo assay was used to measure cytotoxicity over the course of the

experiment. Cytotoxicity was evaluated at every media change (Days 3, 6, 9, 12, and

14). At each time point, concentrations that surpassed a threshold of 10-fold LDH

release compared to the solvent controls were considered overtly cytotoxic. In addition

to the LDH-Glo assay, cytotoxicity was evaluated based on read counts (TempO-Seq)

and spheroid morphology (microscopy). Once LDH thresholds were surpassed, this

cytotoxic concentration was eliminated from all time points thereafter.

Exposure of the liver cell spheroids to PFOA, the sole carboxylic acid of the

group, did not induce significant cytotoxicity over time at 0.02 – 20 µM concentrations

(Figure 2A). The 100 µM PFOA treatment resulted in an increase in LDH release in the

supernatants beginning on day-6 (13.9-fold over control). By day-9, LDH levels

decreased due to the decline in viable cells remaining. Data from the day-10 and day-14

time points of the 100 µM PFOA dose were excluded from further analyses based on

these cytotoxicity results. The 50 µM concentration at day-14 was also eliminated

because visual inspection revealed disintegration of the spheroids at this concentration

and time point.

PFOS was the most cytotoxic of the chemicals tested based on the magnitude of

LDH leakage and the speed at which cell death occurred at the high concentrations

(Figure 2C). The onset of cytotoxic effects occurred at 50 µM and 100 µM PFOS,

causing dramatic cell death as early as 24 hrs post-exposure. The RLU ratios declined

at later time points because of fewer viable cells as cell death progressed. After day-6,

the LDH release for this concentration was no longer different from the control, due to

the low number of surviving cells (visual qualitative inspection, data not shown). For the

50 µM PFOS treatment, an initial 50-fold increase in cytotoxicity was observed after

day-1. Thus, the 50 µM and 100 µM concentrations were both excluded from the gene

expression and benchmark concentration analysis. No overt cytotoxicity was observed

for the shorter (PFBS) or longer (PFDS) chain PFAS (Figures 2B and 2D respectively).

Identification of differentially expressed genes and hierarchical clustering

Table 1 summarizes the numbers of differentially expressed genes (FDR adj p-

value ≤ 0.05 and |FC| ≥ 1.5) for each PFAS. As anticipated, the number of differentially

expressed genes generally showed a trend toward increasing response with

concentration. PFOS exposure induced the greatest number of differentially expressed

genes; whereas, PFBS induced very few differentially expressed genes across time,

even at the highest concentration (largest number of differentially expressed genes was

76 on day-10 at 100 µM). Comparing all PFAS at 20 µM, the highest non-cytotoxic

concentration in common across all of the chemicals, revealed that the rank order for

which PFAS induced the most transcriptional changes was consistent across time in the

following order: PFOS > PFDS > PFOA > PFBS. There were minimal transcriptional

alterations at the lowest concentrations; however, there was some indication for the

possibility of non-monotonic responses (e.g., 85 differentially expressed genes for

PFOS at 0.1 µM on day-1; and 30 genes for PFDS at 0.1 µM on day-10, Table 1). Note

that a detailed pathway analysis of PFOS (presented below) over each time point is

depicted within the supplementary information and suggests some differences in

responses at lower (i.e., 0.1 µM) compared to higher (i.e., 10 and 20 µM) exposures

over specific time points. Interpretation of these observations is beyond the scope of the

current study but will be the subject of future investigation. Nevertheless, analysis of the

differentially expressed genes for each PFAS at their top non-cytotoxic concentrations

(Supplemental File 10 Panel A) and at a uniform concentration of 20 µM (Supplemental

File 10B) showed few genes in common. Interestingly, Perilipin-2 (PLIN2), a lipid droplet

associated protein involved in the storage of fats, appeared in several of the commonly

affected genes.

Principal component analysis was conducted on the differentially expressed

genes at each time point to explore general relationships in the transcriptional profiles

induced by the PFAS (Figure 3; Supplemental File 8B). Generally, principal component

1 (the main driver of variability) appears to be based on PFAS concentration, with

higher concentrations to the right on the plot and lower concentrations together on the

left. The analysis demonstrates the more marginal impact of PFBS on the liver spheroid

transcriptome. The high concentrations of PFBS tend to group more closely with the low

concentrations and controls (in particular after day-1). In addition, higher concentrations

of PFOS and PFDS were more frequently in close proximity to each other than they

were to PFOA, suggesting more similar profiles. However, by day-14, principal

component 2 separates PFOS and PFDS. Note that the top concentration here for both

PFOS and PFOA is 20 µM.

Functional analysis and upstream regulators

Pathway (i.e., gene set) analyses were conducted in IPA using the differentially

expressed genes from the 20 µM concentration (selected because it was the top non-

cytotoxic concentration in common across all the PFAS) (Figure 4A), or the top non-

cytotoxic concentration (Figure 4B), for each chemical and time points. This analysis

revealed that most pathways affected by PFAS were downregulated based on IPA

predictions that factor in the directionality of the expression changes. Overall, there was

a remarkably similar pattern of pathway enrichment for PFOS and PFDS that was

different from PFOA and PFBS.

PFOS and PFDS pathway analyses suggest that the exposures cause inhibition

of nicotine and melatonin degradation pathways as well as downregulation of

cholesterol biosynthesis pathways and LXL/RXR activity, which all appear to persist

across time. In contrast, PFOA treatment appears to strongly activate stearate

biosynthesis and fatty acid beta oxidation pathways. Unlike the other PFAS, PFBS

displayed such small proportions of gene expression changes that it did not perturb

many canonical pathways up to 20 µM. However, the 100 µM concentration of PFBS

elicited a strong upregulation of transcript in several pathways at the 10 and 14-day time

points (Figure 4B) including fatty acid oxidation, stearate biosynthesis, nicotine

degradation, melatonin degradation, acetone degradation, estrogen biosynthesis, and

bupropion degradation. Supplemental File 11 A & B illustrates a less stringent pathway

analysis for all PFAS over all time points (no z-score applied).

A detailed overview of perturbed pathways over all concentrations and time

points for each PFAS is beyond the scope of this study, which is geared toward analysis

of general trends/patterns across the data. However, concentration-response for the

pathway analysis is shown for PFOS as an illustration (Supplemental File 12). This

analysis supports the general trend of increasing pathway perturbations at higher

concentrations (with the exception of 0.1 µM on day-1), and consistency in response of

specific pathways across concentration and time including inhibition of cholesterol

biosynthesis, LXR/RXR activation, melatonin degradation, and nicotine degradation.

The upstream analysis was restricted to more stringent criteria (z-score filter of

absolute +/- |3.0| and a Benjamini Hochberg adjusted p-value of 0.05) to focus on the

most impacted regulators representing potential key mediators of the differentially

expressed genes and to provide additional insight into mode of action. Peroxisome

proliferator-activated receptor alpha (PPARα) was predicted to be an activated

upstream regulator for all four PFAS at all time points (Figure 5). Furthermore, PPARα

was predicted to have increased activity in all PFAS exposures to varying degrees. Of

the predicted inhibited upstream regulators, Copper-transporting ATPase 2 (ATP7B),

sterol regulatory element-binding transcription factor 1 and 2 (SREBF 1 and SREBF 2),

sterol regulatory element-binding protein cleavage-activating protein (SCAP), and

Insulin Receptor (INSR) were all predicted to be inhibited at all-time points with PFOS

and PFDS. Interestingly, these upstream regulators were also predicted to be inhibited

(albeit more weakly) by PFOA at the day-4 and day-10 time points. Overall, there was a

strikingly consistent pattern of upstream regulator activation/inhibition between PFOS

and PFDS that differed from the other two PFAS. Although this stringent approach

limited the numbers of regulators analyzed, it was a useful to begin examining the PFAS

targets.

Benchmark concentration analysis

To compare potencies across the PFAS, we conducted a transcriptional

benchmark concentration analysis with a benchmark response of one standard

deviation. To broadly compare potencies, median benchmark concentrations for all

genes that could be obtained from best-fit models within BMDExpress were derived

along with a bootstrapped 95% confidence interval (Figure 6). An overview of the

proportion of genes that were best fit to each model analyzed is shown in the

supplementary information (Supplemental File 13). PFOS consistently had a median

benchmark concentration between 8.2 to 9.1 µM with narrow confidence intervals,

across all time points. There was some variability across time in the benchmark

concentrations of the other PFAS. The median benchmark concentrations of both PFOA

and PFDS on day-1 were significantly higher than PFOS (non-overlapping confidence

intervals); however, by day-4 (for PFDS), and day-14 (for PFOA), both of these PFAS

were within the same range as PFOS, suggesting similar potencies. In contrast, PFBS

was significantly less potent with benchmark concentrations ranging from 46.3 to 69.7

µM (Supplemental File 14).

Gene accumulation plots show the benchmark concentrations for each of the

genes that could be modeled (Figure 7). Note that overtly cytotoxic concentrations were

not modeled; thus, some curves terminate at lower concentrations than others (i.e., 20

µM for all PFOS time points, and 20 µM for PFOA for day- 14). The plot demonstrates

that of those genes that could be modeled, the transcriptional benchmark

concentrations for PFOS were generally lower overall than the other PFAS (i.e., shifted

furthest to the left on the x-axis). Similar to the overall BMC median gene comparisons

above, PFDS and PFOS accumulation curves in the low concentration ranges are more

overlapping by day-4 and later, suggesting similar potencies. By comparison, PFBS

transcripts had higher benchmark concentrations relative to the other three PFAS,

indicating that it was the least potent in inducing transcriptional changes (most right

shifted). Notably, transcriptional activity for PFOS, PFOA, and PFDS converge at the

origin, indicating that the 'most sensitive' gene transcripts are induced in a similar

concentration range.

Bioactivity Exposure Ratio

The bioactivity exposure ratio (BER) is the ratio of the administered equivalent

dose for a new approach methodology (NAM) point of departure to the daily population

exposure for a given chemical. A large bioactivity exposure ratio value is an indication of

lower risk, whereas a lower value indicates potentially increased risk to human health

(Wetmore *et al.*, 2015b; Paul Friedman *et al.*, 2020). The use of transcriptomics in risk

assessment is a relatively new concept and there is lack of consistency in how to derive

the transcriptomic point of departure. Recognizing this lack of standardization, Farmahin

et al., (2017) demonstrated general consistency (within 10x) across a variety of

methods used for transcriptomic point of departure determination. Herein we compare

two of the most conservative approaches (applying precautionary practices) for deriving

a transcriptomic point of departure. For **Approach 1**, the applied point of departure is

comparable to those approaches used previously for ToxCast data: the 5[th] percentile

gene benchmark concentration (Paul Friedman *et al.*, 2020). **Approach 2** was based on

data demonstrating a strong correlation between the dose at which apical effects *in vivo*

occur and the lowest (most sensitive) *in vivo* pathway benchmark concentration

(Thomas *et al.*, 2013). Bioactivity exposure ratios were derived by dividing the

administered equivalent dose (for each transcriptomic point of departure approach) by

exposure estimates taken from the US population (Figure 8). The bioactivity exposure

ratios for each approach (approach 1, blue diamond; approach 2, red diamond) were

then compared to a range of endpoints derived from animal exposure data used in risk

assessment guidelines (green crossbar) at each time point for PFOA and PFOS (Table

2). As previously noted, PFBS and PFDS were excluded from bioactivity exposure ratio

calculation due to lack of available data.

For PFOA, administered equivalent dose values slightly increased from day-1 to

day-4 and were lowest by day-14 for both point of departure approaches (Figure 8). The

calculated bioactivity exposure ratios for approach 1 were within 2-orders of magnitude

of the population exposure, with a range 178 to 848, except for day-14 with a bioactivity

exposure ratio of 26 (Table 2). The second approach generally decreased the bioactivity

exposure ratios for longer treatment times (day-10 and day-14 bioactivity exposure

ratios of 95 and 27, respectively) compared to shorter treatment times (day-1 and day-4

bioactivity exposure ratios of 111 and 160, respectively) (Table 2). The reference

bioactivity exposure ratio for PFOA (105) was derived using the lowest endpoint in

Health Canada's drinking water guidelines: the BMD_{10} for induction of hepatocellular

hypertrophy in rats (Health Canada, 2018a). This animal *in vivo* bioactivity exposure

ratio was within the same order of magnitude for days 1, 4, and 10, but was about 4x

higher than the bioactivity exposure ratios for both approaches 1 and 2 on day-14

(Table 2).

For PFOS, administered equivalent dose values remained relatively constant

over treatment time for both approaches (Figure 8). In addition, the bioactivity exposure

ratios (Table 2) for approach 1 (range of 112 to 184) were highly similar to approach 2

(range 124 to 211), with approach 1 being marginally more conservative. The reference

bioactivity exposure ratio for PFOS (44) was derived using the lowest endpoint from

Health Canada's drinking water guidelines: the no-observed-adverse-effect-level

(NOAEL) for induction of hepatocellular hypertrophy in rats (Health Canada, 2018b).

This animal *in vivo* data-derived bioactivity exposure ratio was within an order of

magnitude of both transcriptomic points of departure, but was consistently lower than

the NAM-based ratio.

Discussion

This study investigates the utility of a primary human liver cell spheroid model

paired with transcriptional profiling as an alternative approach to toxicological

assessment of PFAS. Four prototype PFAS were examined using a detailed

concentration-response analysis across four time points. The extent of transcriptional

changes induced by PFOS was larger than any of the other PFAS, with transcriptional

perturbations at matched concentrations in the order PFOS > PFDS > PFOA > PFBS.

Although PFOS had the lowest benchmark concentrations at the early time points, the

transcriptomic benchmark concentrations of PFOS, PFDS, and PFOA were overlapping

at the later time points, indicating the importance of transcriptional time-series analyses

in potency comparisons. In addition, the lowest transcriptomic benchmark

concentrations were highly similar across these three PFAS, suggesting fairly equal

potencies. Pathway and upstream regulator analysis revealed that although PFOS and

PFDS transcriptional profiles were most similar to each other, all of the PFAS appeared

to activate PPARα. Indeed, PFOS and PFDS induced remarkably similar transcriptional

alterations at the pathway level. In contrast, the short-chain PFAS (PFBS) induced

marginal changes in transcriptional activity and was much less potent. These data serve

as a foundational data set for comparison of additional data-poor PFAS and for

developing approaches in this human liver cell model in future studies.

Functional analysis of the differentially expressed genes suggests that PPARα

plays a role in mediating the toxic effects of PFAS in human liver spheroids. PPARα

was predicted to be an 'activated' upstream across all of the PFAS and time points.

PPARs are nuclear receptors involved in the regulation of genes involved in energy

homeostasis and lipid metabolism (Janssen *et al.*, 2015; Liss and Finck, 2017). Dose-

dependent increases in PPARα-regulated genes with PFOS exposure have been

observed in rodents (Ye *et al.*, 2012). Likewise, increased PPARα activity in rodents is

associated with exposure to both PFOA and PFOS (Rosen *et al.*, 2008, 2010).

However, these studies also reveal that PPARα knock-out mice have altered lipid

metabolism pathways following PFOS and PFOA exposure, similar to the wild type

animals, indicating PPARα-independent mechanisms as well (Rosen *et al.*, 2008, 2010;

Abbott *et al.*, 2009). Our data support that PPARα activity is increased in response to

PFAS exposure in human liver spheroids, but that there are other upstream regulators

that are likewise affected.

PFOA, PFOS, and PFDS induced changes in gene expression in our study that

are suggestive of inhibition of several upstream regulators: ATP7B, SREBF1 and 2,

SCAP, and INSR. Inhibition of ATP7B, an ATPase copper protein transporter required

for the transport of copper to the bile, is associated with copper accumulation in

hepatocytes (Bartee and Lutsenko, 2007). Inhibition of sterol regulatory binding factors

1 and 2 (SREBF 1 and 2), and sterol regulatory element-binding protein cleavage –

activating protein (SCAP), which are both a part of the cholesterol, fatty acids, and

triglycerides synthesis pathways (Moslehi and Hamidi-zad, 2018), suggests

downregulation of lipogenesis. Perturbation of these pathways is potentially adaptive;

i.e., as PPARα is activated, SREBP1, 2 and SCAP may be downregulated in an attempt

to reach homeostasis of lipid accumulation. Finally, suppression of the insulin receptor

(INSR), an activator of the SREBF 1 transcription factor, suggests that PFAS exposure

affects the insulin signalling pathway and thus impairs normal lipogenesis further. Over

time, continued disruption of cholesterol efflux into the intestine may ultimately lead to

liver disease (Vrins *et al.*, 2012; Utzschneider and Kahn, 2006).

PFOA exhibited a broadly different pattern of transcriptional pathway alterations

at the concentrations examined than the other PFAS. PFOS and PFDS exerted major

effects on the transcription of genes involved in cholesterol biosynthesis. In contrast,

PFOA upregulated transcripts involved in the stearate biosynthesis I pathway, in parallel

with the fatty acid-ß oxidation pathway. Das *et al*. (2017) found that exposure of wild

type and PPARα-null mice to PFOA resulted in increased fatty acid accumulation in the

liver, which is aligned with our findings. Upregulation of the fatty acid-ß oxidation

pathway after PFOA exposure has also been reported in Balb/c mice (Yu *et al.*, 2016)

and in Wistar rats (Haughom and Spydevold, 1992). Although upregulation of the

stearate biosynthesis pathway, at the same time as upregulation of the oxidation or

breakdown of fatty acids, seems counterproductive, we speculate that PFOA is causing

dysregulation of pathways involved in lipid homeostasis in the liver. Overall, although

fairly equal in transcriptional potency, PFOA appears to operate through different

mechanisms to PFOS and PFDS, suggesting that it may be advisable to consider this

carboxylate separately from PFAS with sulfonic acid functional groups.

PFBS is clearly less potent at inducing transcriptional changes than the other

PFAS. Indeed, only the highest concentrations of PFBS and later time points had

significant transcriptional activity. Our results are consistent with an analysis by Naile *et*

al. (2011) of 10 PFAS in rat hepatocytes that revealed that shorter chain PFAS cause

less transcriptional perturbation.

Our transcriptomic findings are also consistent with demonstrated effects of

PFAS on cholesterol, lipid synthesis, and metabolism in other model organisms.

Although data for most PFAS are limited, rodent exposure to PFOS and PFOA is known

to induce changes in fatty acid production, cholesterol synthesis, and lipid homeostasis

leading to cholestasis/steatosis (Wang *et al.*, 2014; Bijland *et al.*, 2011; Du *et al.*, 2009;

Peden-Adams *et al.*, 2009; Wan *et al.*, 2012). In keeping with these effects, obese

participants from the National Health and Nutrition Examination Survey (2011-2014)

exhibit a positive association of PFOA, PFHxS, and PFNA exposure with major liver

biomarkers of steatosis (Jain and Ducatman, 2019). These findings support the

relevance of our results for exposed humans. Overall, our short-term *in vitro*

transcriptomic study points to the same genes/pathways/disease outcomes as predicted

by *in vivo* studies and associated human data (Guruge *et al.,* 2006*;* Jin *et al.,* 2020). Our

results support an effect of PFAS on lipogenesis in human liver cells that warrants

further research.

Benchmark concentration analysis was conducted to compare potencies of the

four PFAS. Based on median gene benchmark concentration as a central estimate of

the transcriptional activity, PFOS initially appeared to be the most potent; however, at

the later timepoints, PFDS and PFOA had similar potencies to PFOS. The same pattern

is observed in the gene accumulation plots, which show the initial PFOS curve to be left

shifted relative to the other PFAS, but with PFDS and PFOA curves more super-

imposed over PFOS at the later time points. It is important to bear in mind that our

potency analysis is based on nominal concentrations of the PFAS exposures with these

in vitro models. Cell permeability and uptake of PFOS has been found to be much faster

than PFBS in human lung epithelial and murine preadipocyte embryonic fibroblasts

(Sanchez Garcia *et al.*, 2018), and PFOS uptake efficiency was 20 times higher than

PFBS in an *in vitro* HEK293 cell model (Zhao *et al.*, 2017). Furthermore, both active and

passive transport mechanisms for PFOA have been identified in isolated rat

hepatocytes (Han *et al.*, 2008) and in the human colorectal adenocarcinoma cell line

Caco-2 (Kimura *et al.*, 2017). In humans, PFOS accumulates within the liver to a greater

extent than PFBS, PFOA, and PFDS (Pérez *et al.*, 2013). Based on the premise of

selective uptake and concentrations of different PFAS, it is reasonable to speculate that

differences in PFAS potencies over time may be influenced in part by the kinetics of

cellular uptake and cell permeability. Zhao et al. (2015; 2017) found that several PFAS

are transported into hepatocytes via Na+ / taurocholate co-transporting polypeptide

(NTCP) and that human organic anion transporting polypeptides (OATPs), which are

expressed in the spheroids in our study (data not shown), contributed to the

enterohepatic circulation of certain PFAS and extend their half-lives. Thus, one

important limitation of our study was that we did not measure the concentration of PFAS

within the cells in the spheroids or the free fraction of the chemicals. Future studies will

examine internal concentrations of PFAS within the spheroid cells over time.

Pharmacokinetic and exposure information were only available for PFOS and

PFOA to derive bioactivity exposure ratios. The two approaches used (Approach 1: 5[th]

percentile gene benchmark concentration and Approach 2: lowest pathway benchmark

concentration) yielded results that were within 10-fold of the apical endpoint bioactivity

exposure ratio, lending confidence to these approaches. Several of the lowest pathway

benchmark concentrations across the PFAS are pathophysiologically relevant, including

mevalonate pathway I, fatty acid activation, and stearate biosynthesis. These pathways

are also consistent with the literature indicating that both PFOS (Reif *et al.*, 2010) and

PFOA (Das *et al.*, 2017) cause hepatic steatosis (i.e., fatty liver disease), providing

biological plausibility that these transcriptomic points of departure are meaningful.

Steatosis was highlighted as an important endpoint within the US EPA's Integrated Risk

Information System (IRIS) on PFOS/PFOA and is recognized as a key step in liver

disease progression (Kaiser *et al.*, 2012). Wetmore *et al.* (2015b) proposed that a

bioactivity exposure ratio cut-off > 100 suggests that the administered equivalent dose

of biological activity is significantly different from daily population exposure estimates.

Our study yielded bioactivity exposure ratios that were close to or less than <100,

indicating that there is a relatively narrow margin between highly exposed members of

the US population and activity *in vitro* for PFOS and PFOA. We note that the

assumptions used within our approach to derive a value for the bioactivity exposure

ratio from the administered equivalent dose do not account for the bioaccumulation from

repeat low-dose exposures, which may be a limitation; this was a consideration when

interpreting the values of the bioactivity exposure ratio within the investigation.

Liver concentration values were estimated from median values in Perez *et al.*, 2013,

and were equivalent to 0.002, 0.08, and 0.01 µM for PFBS, PFOS, and PFOA,

respectively; PFDS could not be compared as observations were below detection limits

within this study (Perez *et a*l., 2013). Since serum concentration measured from the

same area were 10 to 5 times lower for the population (Ericson *et al.*, 2007), the liver

concentration can be used as an adequate marker of exposure against the benchmark

concentrations values. The lowest median gene the benchmark concentrations within

our observations were 46.3, 8.2, and 6.4 µM for PFBS, PFOS, and PFOA respectively.

The PFBS estimates from the literature are several orders of magnitude different from

the lowest median gene benchmark concentration in our findings but there was a <

1000-fold difference for PFOA, and approximately 100-fold difference in ratio for PFOS.

Considering a more conservative estimate of the margin of potential exposure, the

median benchmark concentration of the gene-set from the lowest, most sensitive

pathway, yields ratios of 0.74 and 1.2 for PFOS and PFOA respectively, suggesting

that changes in gene expression at current exposure levels within this population are

likely.

High-throughput transcriptomics is of considerable value because it broadly

informs global biological perturbations induced by chemical exposures and can be

readily used to compare similarities/differences in biological activities and potencies

across chemicals. This study produced a baseline transcriptomic data set and pipeline

to be applied in future studies to study data-poor PFAS. We demonstrate several

approaches that can be used to inform hazard, mode of action, and potency.

Acknowledgements

We thank Dr. Francesco Marchetti, Dr. Marc Beal, and Dr. Scott Auerbach for

their helpful commentary and review of this research, the US EPA for their valuable

consultations, as well as BioSpyder for their contributions on this project. We would like

to thank Dr. Iain Lambert, Carleton University, for logistical support for this project.

Funding

We acknowledge the Water and Air Quality Bureau, HECSB, Health Canada, the

Chemicals and Environmental Health Management Bureau, HECSB, Health Canada,

and the Environmental Health Sciences & Research Bureau, HECSB, Health Canada

who jointly funded this research.

References

Abbott, B.D., Wolf, C.J., Das, K.P., Zehr, R.D., Schmid, J.E., Lindstrom, A.B., Strynar,

M.J., and Lau, C. (2009). Developmental toxicity of perfluorooctane sulfonate

(PFOS) is not dependent on expression of peroxisome proliferator activated

receptor-alpha (PPARα) in the mouse. *Reproductive Toxicology*, **27**, 258–265.

Armitage, J., Cousins, I.T., Buck, R.C., Prevedouros, K., Russell, M.H., Macleod, M.,

and Korzeniowski, S.H. (2006). Modeling global-scale fate and transport of

perfluorooctanoate emitted from direct sources. *Environmental Science and*

Technology, **40**, 6969–6975.

Armitage, J.M., Macleod, M., and Cousins, I.T. (2009a). Comparative assessment of the

global fate and transport pathways of long-chain perfluorocarboxylic acids (PFCAs)

and perfluorocarboxylates (PFCs) emitted from direct sources. *Environmental*

Science and Technology, **43**, 5830–5836.

Bach, C.C., Vested, A., Jørgensen, K.T., Bonde, J.P.E., Henriksen, T.B., Toft, G.

(2016). Perfluoroalkyl and polyfluoroalkyl substances and measures of human

fertility: a systematic review. *Critical Reviews in Toxicology*, **46**, 735–755

Bartee, M.Y. and Lutsenko, S. (2007). Hepatic copper-transporting ATPase ATP7B:

Function and inactivation at the molecular and cellular level. *BioMetals*, **20**, 627–

637.

Bijland, S., Rensen, P.C.N., Pieterman, E.J., Maas, A.C.E., van der Hoorn, J.W., van

Erk, M.J., Havekes, L.M., van Dijk, K.W., Chang, S.C., Ehresman, D.J., *et al.*

(2011). Perfluoroalkyl sulfonates cause alkyl chain length-dependent hepatic

steatosis and hypolipidemia mainly by impairing lipoprotein production in APOE*3-

leiden CETP mice. *Toxicological Sciences*, **123**, 290–303.

Bjork, J.A., Butenhoff, J.L., and Wallace, K.B. (2011). Multiplicity of nuclear receptor

activation by PFOA and PFOS in primary human and rodent hepatocytes.

Toxicology, **288**, 8–17.

Chang, E.T., Adami, H.O., Boffetta, P., Cole, P., Starr, T.B., and Mandel, J.S. (2014). A

critical review of perfluorooctanoate and perfluorooctanesulfonate exposure and

cancer risk in humans. *Critical Reviews in Toxicology*, **44**, 1–81.

Das, K.P., Wood, C.R., Lin, M.J., Starkov, A.A., Lau, C., Wallace, K.B., Corton, J.C.,

and Abbott, B.D. (2017). Perfluoroalkyl acids-induced liver steatosis: Effects on

genes controlling lipid homeostasis. *Toxicology*, **378**, 37–52.

Du, Y., Shi, X., Liu, C., Yu, K., and Zhou, B. (2009). Chronic effects of water-borne

PFOS exposure on growth, survival and hepatotoxicity in zebrafish: A partial life-

cycle test. *Chemosphere*, **74**, 723–729.

Ek, C.J., Nathanielsz, P., Li, C., and Mallard, C. (2015). Transcriptomal changes and

functional annotation of the developing nonhuman primate choroid plexus. *Frontiers*

in Neuroscience, **9**.

Environmental Protection Agency. (2020). EPA'S PFAS Action Plan: Program Update

February 2020.

Ericson, I., Gómez, M., Nadal, M., van Bavel, B., Lindström, G., Domingo, J.L. (2007).

Perfluorinated chemicals in blood of residents in Catalonia (Spain) in relation to age

and gender: A pilot study. *Environment International*, **33**, 616–623.

Eriksen, K.T., Raaschou-Nielsen, O., McLaughlin, J.K., Lipworth, L., Tjønneland, A.,

Overvad, K., and Sørensen, M. (2013). Association between Plasma PFOA and

PFOS Levels and Total Cholesterol in a Middle-Aged Danish Population. *PLoS*

ONE, **8**, e56969.

Farmahin, R., Williams, A., Kuo, B., Chepelev, N.L., Thomas, R.S., Barton-Maclaren,

T.S., Curran, I.H., Nong, A., Wade, M.G., and Yauk, C.L. (2017). Recommended

approaches in the application of toxicogenomics to derive points of departure for

chemical risk assessment. *Archives of Toxicology*, **91**, 2045–2065.

Filgo, A.J., Quist, E.M., Hoenerhoff, M.J., Brix, A.E., Kissling, G.E., and Fenton, S.E.

(2015). Perfluorooctanoic Acid (PFOA)–induced Liver Lesions in Two Strains of

Mice Following Developmental Exposures: PPARα Is Not Required. *Toxicologic*

Pathology, **43**, 558–568.

Geiger, S.D., Xiao, J., Ducatman, A., Frisbee, S., Innes, K., and Shankar, A. (2014).

The association between PFOA, PFOS and serum lipid levels in adolescents.

Chemosphere, **98**, 78–83.

Giesy, J.P., Kanna, K. (2001). Global Distribution of Perfluorooctane Sulfonate in

Wildlife. *Environmental Science & Technology*, **35**, 1339-1342.

Guruge, K.S., L. W. Y., Yeung, N., Yamanaka, S., Miyazaki, P.K. S., Lam. J.P., Giesy,

P. D., Jones, N., Yamashita. (2006). Gene Expression Profiles in Rat Liver Treated

With Perfluorooctanoic Acid (PFOA). *Toxicological sciences* **89**, 93–107.

Han, X., Yang, C.H., Snajdr, S.I., Nabb, D.L., and Mingoia, R.T. (2008). Uptake of

perfluorooctanoate in freshly isolated hepatocytes from male and female rats.

Toxicology Letters, **181**, 81–86.

Haughom, B. and Spydevold, O. (1992). The mechanism underlying the hypolipemic

effect of perfluorooctanoic acid (PFOA), perfluorooctane sulphonic acid (PFOSA)

and clofibric acid. *Biochimica et Biophysica Acta,* **1128**, 65-72.

Health Canada. (2017). Water talk: drinking water screening values: perfluoroalkylated

substances. Health Canada.

Health Canada (2018a). Guidelines for Canadian Drinking Water Quality Guideline

Technical Document Perfluorooctanoic Acid (PFOA).

Health Canada (2018b). Guidelines for Canadian Drinking Water Quality: Guideline

Technical Document – Perfluorooctane Sulfonate (PFOS).

Hill, C.E., Myers, J.P., and Vandenberg, L.N. (2018). Nonmonotonic dose–Response

curves occur in dose ranges that are relevant to regulatory decision-making. *Dose-*

Response, **16**.

House, J.S., Grimm, F.A., Jima, D.D., Zhou, Y.H., Rusyn, I., and Wright, F.A. (2017). A

pipeline for high-throughput concentration response modeling of gene expression

for toxicogenomics. *Frontiers in Genetics*, **8**.

Jain, R.B. and Ducatman, A. (2019). Selective Associations of Recent Low

Concentrations of Perfluoroalkyl Substances with Liver Function Biomarkers:

NHANES 2011 to 2014 Data on US Adults Aged ≥20 Years. *Journal of*

Occupational and Environmental Medicine, **61**, 293–302.

Janssen, A.W.F., Betzel, B., Stoopen, G., Berends, F.J., Janssen, I.M., Peijnenburg,

A.A., and Kersten, S. (2015). The impact of PPARaα activation on whole genome

gene expression in human precision cut liver slices. *BMC Genomics*, **16**,760-773.

Jin, R., McConnell R., Cioffi C., Xu S., Walker D.W., Stratakis N., Jones D.P., Miller

G.W., Peng C., Conti D.V., Vos M.B., Chatzi L. (2020). *Environmental International.*

134.

Kaiser, J.P., Lipscomb, J.C., and Wesselkamper, S.C. (2012). Putative mechanisms of

environmental chemical-induced steatosis. *International Journal of Toxicology*, **31**,

551–563.

Kavlock, R.J., Bahadori, T., Barton-Maclaren, T.S., Gwinn, M.R., Rasenberg, M., and

Thomas, R.S. (2018). Accelerating the Pace of Chemical Risk Assessment.

Chemical Research in Toxicology, **31**, 287–290.

Kielsen, K., Shamim, Z., Ryder, L.P., Nielsen, F., Grandjean, P., Budtz-Jørgensen, E.,

and Heilmann, C. (2016). Antibody response to booster vaccination with tetanus

and diphtheria in adults exposed to perfluorinated alkylates. *Journal of*

Immunotoxicology, **13**, 270–273.

Kimura, O., Fujii, Y., Haraguchi, K., Kato, Y., Ohta, C., Koga, N., and Endo, T. (2017).

Uptake of perfluorooctanoic acid by Caco-2 cells: Involvement of organic anion

transporting polypeptides. *Toxicology Letters*, **277**, 18–23.

Krewski, D., Andersen, M.E., Tyshenko, M.G., Krishnan, K., Hartung, T., Boekelheide,

K., Wambaugh, J.F., Jones, D., Whelan, M., Thomas, R., *et al.* (2020). Toxicity

testing in the 21st century: progress in the past decade and future perspectives.

Archives of Toxicology, **94**.

Liss, K.H.H. and Finck, B.N. (2017). PPARs and nonalcoholic fatty liver disease.

Biochimie, **136**, 65–74.

Louisse, J., Rijkers, D., Stoopen, G., Janssen, A., Staats, M., Hoogenboom, R.,

Kersten, S., Peijnenburg, A. (2020). Perfluorooctanoic acid (PFOA),

perfluorooctanesulfonic acid (PFOS), and perfluorononanoic acid (PFNA) increase

triglyceride levels and decrease cholesterogenic gene expression in human

HepaRG liver cells. *Archives of Toxicology*, **94**, 3137–3155.

Mav, D., Shah, R.R., Howard, B.E., Auerbach, S.S., Bushel, P.R., Collins, J.B., Gerhold,

D.L., Judson, R.S., Karmaus, A.L., Maull, E.A., *et al.* (2018). A hybrid gene

selection approach to create the S1500+ targeted gene sets for use in high-

throughput transcriptomics. *PLoS ONE*, **13**.

Moslehi, A. and Hamidi-zad, Z. (2018). Role of SREBPs in Liver Diseases: A Mini-

review. *Journal of Clinical and Translational Hepatology*, **6**, 1–7.

Naile, J., Wiseman, S., Bachtold, K., Jone, P.D., Giesy, J.P. (2012). Transcriptional

effects of perfluorinated compounds in rat hepatoma cells. *Chemosphere*, **86**, 270–

277.

National Toxicology Program Monograph Immunotoxicity Associated with Exposure to

Perfluorooctanoic Acid or Perfluorooctane Sulfonate. (2016).

National Toxicology Program Research Report on Program Approach To Genomic

Dose-Response Modeling. (2018).

Organisation for Economic Co-operation and Development (2018). Environment

Directorate Joint Meeting of The Chemicals Committee And The Working Party On

Chemicals, Pesticides and Biotechnology Towards a New Comprehensive Global

(Database of Per-and-Polyfluroalkyl Substances (PFASs): Summary Report on

Updating The OECD 2007 List of Per-and-Polyfluroalkyl Substances (PFASs)

Series On Risk Management. No. 39

Paul Friedman, P.K., Gagne, M., Loo, L.H., Karamertzanis, P., Netzeva, T., Sobanski,

T., Franzosa, J.A., Richard, A.M., Lougee, R.R., Gissi, A., *et al.* (2020). Utility of in

Vitro Bioactivity as a Lower Bound Estimate of in Vivo Adverse Effect Levels and in

Risk-Based Prioritization. *Toxicological Sciences*, **173**, 202–225.

Peden-Adams, M.M., Stuckey, J.E., Gaworecki, K.M., Berger-Ritchie, J., Bryant, K.,

Jodice, P.G., Scott, T.R., Ferrario, J.B., Guan, B., Vigo, C., *et al.* (2009).

Developmental toxicity in white leghorn chickens following in ovo exposure to

perfluorooctane sulfonate (PFOS). *Reproductive Toxicology*, **27**, 307–318.

Pérez, F., Nadal, M., Navarro-Ortega, A., Fàbrega, F., Domingo, J.L., Barceló, D., and

Farré, M. (2013). Accumulation of perfluoroalkyl substances in human tissues.

Environment International, **59**, 354–362.

Reif, D.M., Martin, M.T., Tan, S.W., Houck, K.A., Judson, R.S., Richard, A.M., Knudsen,

T.B., Dix, D.J., and Kavlock, R.J. (2010). Endocrine profiling and prioritization of

environmental chemicals using ToxCast data. *Environmental Health Perspectives*,

118, 1714–1720.

Rosen, M.B., Abbott, B.D., Wolf, D.C., Corton, J.C., Wood, C.R., Schmid, J.E., Das,

K.P., Zehr, R.D., Blair, E.T., and Lau, C. (2008). Gene Profiling in the Livers of

Wild-type and PPARα-Null Mice Exposed to Perfluorooctanoic Acid. *Toxicologic*

Pathology, **36**, 592–607.

Rosen, M.B., Schmid, J.R., Corton, J.C., Zehr, R.D., Das, K.P., Abbott, B.D., and Lau,

C. (2010). Gene expression profiling in wild-type and PPARα-null mice exposed to

perfluorooctane sulfonate reveals PPARα-independent effects. *PPAR Research,*

2010.

Sanchez Garcia, D., Sjödin, M., Hellstrandh, M., Norinder, U., Nikiforova, V., Lindberg,

J., Wincent, E., Bergman, Å., Cotgreave, I., and Munic Kos, V. (2018). Cellular

accumulation and lipid binding of perfluorinated alkylated substances (PFASs) – A

comparison with lysosomotropic drugs. *Chemico-Biological Interactions*, **281**, 1–10.

Schlezinger, J.J., Puckett, H., Oliver, J., Nielsen, G., Heiger-Bernays, W., Webster, T.F.

(2020). Perfluorooctanoic acid activates multiple nuclear receptor pathways and

skews expression of genes regulating cholesterol homeostasis in liver of

humanized PPARα mice fed an American diet. *Toxicology and Applied*

Pharmacology, **405***, 115204.*

Steenland, K., Tinker, S., Frisbee, S., Ducatman, A., Vaccarino, V. (2009). Association

of Perfluorooctanoic Acid and Perfluorooctane Sulfonate With Serum Lipids Among

Adults Living Near a Chemical Plant. American Journal of Epidemiology, **170**,

1268-1278.

Thomas, R.R., Wesselkamper, S.C., Wang, N.C.Y., Zhao, J.J., Petersen, D.D.,

Lambert, J.C., Cote, I., Yang, L., Healy, E., Black, M.B., *et al.* (2013). Temporal

concordance between apical and transcriptional points of departure for chemical

risk assessment. *Toxicological Sciences*, **134**, 180–194.

Trejo, C.L., Babić, M., Imler, E., Gonzalez, M., Bibikov, S.I., Shepard, P.J.,

VanSteenhouse, H.C., Yeakley, J.M., and Seligmann, B.E. (2019). Extraction-free

whole transcriptome gene expression analysis of FFPE sections and histology-

directed subareas of tissue. *PLoS ONE*, **14**.

Utzschneider, K.M. and Kahn, S.E. (2006). Review: The role of insulin resistance in

nonalcoholic fatty liver disease. *Journal of Clinical Endocrinology and Metabolism*,

91, 4753–4761.

Vrins, C.L.J., Ottenhoff, R., van den Oever, K., de Waart, D.R., Kruyt, J.K., Zhao, Y.,

van Berkel, T.J.C., Havekes, L.M., Aerts, J.M., van Eck, M., *et al.* (2012). Trans-

intestinal cholesterol efflux is not mediated through high density lipoprotein. *Journal*

of Lipid Research, **53**, 2017–2023.

Wambaugh, J.F., Setzer, R.W., Pitruzzello, A.M., Liu, J., Reif, D.M., Kleinstreuer, N.C.,

Wang, N.C.Y., Sipes, N., Martin, M., Das, K., *et al.* (2013). Dosimetric anchoring of

In vivo and In vitro studies for perfluorooctanoate and perfluorooctanesulfonate.

Toxicological Sciences, **136**, 308–327.

Wan, H.T., Zhao, Y.G., Wei, X., Hui, K.Y., Giesy, J.P., and Wong, C.K.C. (2012). PFOS-

induced hepatic steatosis, the mechanistic actions on β-oxidation and lipid

transport. *Biochimica et Biophysica Acta - General Subjects*, **1820**, 1092–1101.

Wang, L., Wang, Y., Liang, Y., Li, J., Liu, Y., Zhang, J., Zhang, A., Fu, J., and Jiang, G.

(2014). PFOS induced lipid metabolism disturbances in BALB/c mice through

inhibition of low density lipoproteins excretion. *Scientific Reports*, **4**, 4582.

Wania, F. (2007). A global mass balance analysis of the source of perfluorocarboxylic

acids in the Arctic Ocean. *Environmental Science and Technology*, **41**, 4529–4535.

Wetmore, B.A., Wambaugh, J.F., Allen, B., Ferguson, S.S., Sochaski, M.A., Setzer,

R.W., Houck, K.A., Strope, C.L., Cantwell, K., Judson, R.S., *et al.* (2015b).

Incorporating high-throughput exposure predictions with dosimetry-adjusted in vitro

bioactivity to inform chemical toxicity testing. *Toxicological Sciences*, **148**, 121–136.

Ye, L., Zhao, B., Yuan, K., Chu, Y., Li, C., Zhao, C., Lian, Q.Q., and Ge, R.S. (2012).

Gene expression profiling in fetal rat lung during gestational perfluorooctane

sulfonate exposure. *Toxicology Letters*, **209**, 270–276.

Yu, N., Wei, S., Li, M., Yang, J., Li, K., Jin, L., Xie, Y., Giesy, J.P., Zhang, X., and Yu, H.

(2016). Effects of Perfluorooctanoic Acid on Metabolic Profiles in Brain and Liver of

Mouse Revealed by a High-throughput Targeted Metabolomics Approach. *Scientific*

Reports, **6**, 23963.

Zanger, U.M. and Schwab, M. (2013). Cytochrome P450 enzymes in drug metabolism:

Regulation of gene expression, enzyme activities, and impact of genetic variation.

Pharmacology and Therapeutics, **138**, 103–141.

Zhao, W., Zitzow, J.D., Ehresman, D.J., Chang, S.C., Butenhoff, J.L., Forster, J., and

Hagenbuch, B. (2015). Na+/taurocholate cotransporting polypeptide and apical

sodium-dependent bile acid transporter are involved in the disposition of

perfluoroalkyl sulfonates in humans and rats. *Toxicological Sciences*, **146**, 363–

373.

Zhao, W., Zitzow, J.D., Weaver, Y., Ehresman, D.J., Chang, S.C., Butenhoff, J.L., and

Hagenbuch, B. (2017). Organic anion transporting polypeptides contribute to the

disposition of perfluoroalkyl acids in humans and rats. *Toxicological Sciences*, **156**,

84–95.

Tables

Table 1. Number of differentially expressed genes in the PFAS-treated spheroids

relative to controls for each time point and concentration.

Note. Sample size was 3 to 4 for each concentration, except 10 µM PFBS, day-4; 0.02

and 100 µM PFDS day-10; and 10 µM PFOA day-4 (where n = 2). Supplemental Files

10 A and B illustrate genes in common at top non-cytotoxic concentrations and at 20

µM. These differentially expressed genes are based on unique genes; any gene with

more than one probe for the same gene was counted only once (probes collapsed).

1006

Table 2. Administered Equivalent Dose (AEDs) and bioactivity exposure ratio (BER)

derived from Approaches 1 and 2 as well as reference values from animal models using

Health Canada drinking water guidelines for PFOA and PFOS.

Animal bioactivity exposure ratio = (Animal AED / Population exposure level)

Approach 1 bioactivity exposure ratio = (Lowest 5th percentile gene AED / Population

exposure level).

Approach 2 bioactivity exposure ratio = (Median gene-level BMD of lowest pathway

AED / Population exposure level).

1015

Figures

Figure 1. (A) Liver spheroid morphology after 4 days of treatment with 0.1% DMSO.

Immunofluorescence staining with CD-68 for Kupffer cells (yellow), phalloidin for F-actin

filaments (green), and DAPI for nuclei (blue), scale = 100 µm. (B) Effects of 48-hour

exposure to Cyclosporine A on spheroid morphology compared to DMSO control.

Microscopy images are reflective of expected detoxification and increased activity of the

detoxification/biotransformation of the spheroids. (A and C) Immunofluorescence

staining for: CYP3A4–(red), F-actin filaments stained with phalloidin (green), and nuclei

with DAPI (blue). (B and D) Immunofluorescence staining for: neutral lipids (green), and

nuclei with DAPI (blue). 400x magnification.

Figure 2. Lactate dehydrogenase (LDH) release as an indicator of cytotoxicity following

treatment with a) PFOA, b) PFBS, c) PFOS, or d) PFDS over a 14-day time course (n =

4 with the exception of day-14, n =2). Plots demonstrate the means +/- standard

deviation of the ratio of sample relative luminescence units (RLU) to mean DMSO RLU.

* indicates the RLU was significantly higher than the DMSO control group ($p < 0.05$).

Note that after reaching a peak, LDH levels decline over time consistent with reduced

numbers of viable cells or evolving tolerance to chemical exposures in the spheroids.

Figure 3. Principal Component Analysis conducted on the probes passing filters (FDR

adjusted p-value < 0.05 and an absolute fold change > 1.5 on a linear scale) at each of

the four time points. Each point represents the individual replicates. The horizontal and

vertical green lines at zero were added to identify the four quadrants in each of the four

plots. Sample size was n = 3-4 for all exposure groups, except for n = 2 on day-4 PFBS

10 µM and PFOA 10 µM, and n = 2 on day-10 PFDS 0.02 and 100 µM. Additional

principal component analyses explored each chemical individually within a time point

(Supplemental File 8A, B). Hierarchical cluster analysis was also performed for each

time point and can be found in Supplemental Files 8C, and 8D.

Figure 4. Pathway analysis of the differentially expressed genes for all time points at the

20 µM concentration (A) or for the highest non-cytotoxic concentrations (B). For (B)

concentrations were 20 µM for PFOA and PFOS, and 100 µM for PFBS and PFDS.

Filters for this analysis were set to z-score ≥ absolute 2.0 and Benjamin–Hochberg

adjusted p-value ≤ 0.05. Blue represents inhibited pathways and orange represents

activated pathways. n = 3-4 for all exposure groups, except for n = 2 on day-10 PFDS

100 µM. Supplemental File 11 A & B is a less stringent pathway analysis for all PFAS

over all time points (no z-score applied). Supplemental File 12 illustrates a more

detailed pathway analysis for PFOS over all time points and concentrations.

Figure 5. Upstream regulator predictions from the differentially expressed genes for the

top non-cytotoxic concentration of each PFAS. The z-score filters were set to 3.0 with a

Benjamini Hochberg adj p-value of 0.05 for this analysis. Blue represents inhibited

upstream regulators and orange represents activated regulators. n = 3-4 for all

exposure groups, except for n = 2 on day-10 PFDS 100 µM.

Figure 6. Median overall gene benchmark concentration (BMC) (colored dot) for all the genes

that could be modeled and 95% confidence intervals for each of the PFAS. n = 3-4 for all

treatment groups, except for n = 2 on day- 4 for PFBS 10 µM and PFOA 10 µM, and day-10

PFDS 0.02 and 100 µM. Supplemental File 14 shows overall benchmark concentrations (95

% CI) by day. A summary of the distribution for all of the modeled genes exposed to PFAS

over 1, 4, 10, and 14 days is in Supplemental File 13, and a summary of sample replicates is

in Supplemental File 6.

Figure 7. Gene accumulation plots showing the benchmark concentrations (BMC) for each

gene that could be modeled for each of the PFAS across time points. n = 3-4 for all exposure

groups, except for n = 2 for day-4 PFBS 10 µM and PFOA 10 µM and day-10 PFDS, 0.02 and

100 µM. Supplemental File 14 summarizes the benchmark concentrations (and 95 % CI) by

day. A summary of the distribution for all of the modeled genes exposed to PFAS over 1, 4, 10,

and 14 days is in Supplemental File 13, and a summary of sample replicates is in

Supplemental File 6.

Figure 8. Boxplots of administered equivalent doses (mg/kg-BW/day) of all transcripts modeled

for PFOA and PFOS are shown. These were used to determine the Bioactivity Exposure Ratio

(BER) based on either the lowest 5th percentile administered equivalent dose (blue diamond)

or the median administered equivalent dose of the gene-set from the most sensitive pathway

(red diamond). These were compared with the estimated daily exposure of the upper limit of

the daily US population exposure (95th percentile) obtained from the US EPA CompTox

dashboard (red dash). A range of administered equivalent doses was included as a

representation of apical points of departure that were derived from non-cancer animal in vivo

data (green crossbar) from Health Canada drinking water guidelines (Health Canada 2018a,

and 2018b).

Tables

Table 1. Number of differentially expressed genes in the PFAS-treated spheroids relative to controls for each time point and concentration.

PFAS	DAY	0.02µM	0.1µM	0.2µM	1µM	2µM	10µM	20µM	50µM	100µM
PFOA	1	0	8	36	8	19	79	69	227	465
	4	22	30	35	41	14	42	100	184	782
	10	14	7	2	4	10	82	101	593	cytotoxic
	14	4	0	8	9	6	76	96	cytotoxic	cytotoxic
PFBS	1	0	1	0	5	49	5	0	44	73
	4	0	5	20	20	0	0	8	0	59
	10	2	1	0	1	2	0	7	54	76
	14	0	0	6	0	1	2	0	2	52
PFOS	1	1	85	3	6	51	167	277	cytotoxic	cytotoxic
	4	6	12	30	48	62	251	313	cytotoxic	cytotoxic
	10	2	7	4	6	60	163	466	cytotoxic	cytotoxic
	14	1	0	3	1	6	249	398	cytotoxic	cytotoxic
PFDS	1	0	1	0	6	22	59	81	177	186
	4	0	17	15	7	20	340	247	171	273
	10	0	30	0	43	40	134	175	232	231
	14	1	3	1	18	100	182	147	211	340

Note. Sample size was 3 to 4 for each concentration, except: 10 µM PFBS, day-4; 0.02 and 100 µM PFDS day-10; and 10 µM PFOA day-4 (where n = 2). Supplemental Files 10 A and B illustrate genes in common at top non-cytotoxic concentrations and at 20 µM. These differentially expressed genes are based on unique genes; any gene with more than one probe for the same gene was counted only once (probes collapsed).

Table 2. Administered Equivalent Dose (AEDs) and bioactivity exposure ratio (BER) derived from Approaches 1 and 2 as well as reference values from animal models using Health Canada drinking water guidelines for PFOA and PFOS.

| | Day | Population Exposure (mg/kgBW/day) | Reference Values (Animal) Health Canada guidelines | | Approach 1 Lowest 5th percentile | | Approach 2 Median AED of the gene-set from the most sensitive pathway | | Pathway |
			AED (mg/kgBW/day)	BER	AED (mg/kgBW/day)	BER	AED (mg/kgBW/day)	BER	
PFOS	01	4.76E-04	0.021	44	0.053	112	0.059	124	Mevalonate Pathway I
	04	4.76E-04	0.021	44	0.056	118	0.092	192	Mevalonate Pathway I
	10	4.76E-04	0.021	44	0.054	113	0.066	138	Antigen Presentation Pathway
	14	4.76E-04	0.021	44	0.088	184	0.101	211	Cholesterol Biosynthesis I
PFOA	01	4.50E-04	0.050	105	0.124	275	0.049	111	Fatty Acid Activation
	04	4.50E-04	0.050	105	0.382	848	0.072	160	Ketogenesis
	10	4.50E-04	0.050	105	0.080	178	0.043	95	Ethanol Degradation IV
	14	4.50E-04	0.050	105	0.012	26	0.012	27	Stearate Biosynthesis I (Animals)

Animal bioactivity exposure ratio = (Animal AED / Population exposure level)
Approach 1 bioactivity exposure ratio = (Lowest 5th percentile gene AED / Population exposure level).
Approach 2 bioactivity exposure ratio = (Median gene-level BMD of lowest pathway AED / Population exposure level).

A

B

Figure 1. (A) Liver spheroid morphology after 4 days of treatment with 0.1% DMSO. Immunofluorescence staining with CD-68 for Kupffer cells (yellow), phalloidin for F-actin filaments (green), and DAPI for nuclei (blue), scale = 100 μm. (B) Effects of 48-hour exposure to Cyclosporine A on spheroid morphology compared to DMSO control. Microscopy images are reflective of expected detoxification and increased activity of the detoxification/biotransformation of the spheroids. (A and C) Immunofluorescence staining for: CYP3A4–(red), F-actin filaments stained with phalloidin (green), and nuclei with DAPI (blue). (B and D) Immunofluorescence staining for: neutral lipids (green), and nuclei with DAPI (blue). 400x magnification.

Figure 2. Lactate dehydrogenase (LDH) release as an indicator of cytotoxicity following treatment with a) PFOA, b) PFBS, c) PFOS, or d) PFDS over a 14-day time course (n = 4 with the exception of day-14, n =2). Plots demonstrate the means +/- standard deviation of the ratio of sample relative luminescence units (RLU) to mean DMSO RLU. * indicates the RLU was significantly higher than the DMSO control group (p < 0.05). Note that after reaching a peak, LDH levels decline over time consistent with reduced numbers of viable cells or evolving tolerance to chemical exposures in the spheroids. An RLU to RLU DMSO ratio greater than 10 was considered to be overtly cytotoxic.

Figure 3. Principal Component Analysis conducted on the probes passing filters (FDR adjusted p-value < 0.05 and an absolute fold change > 1.5 on a linear scale) at each of the four time points. Each point represents the individual replicates. The horizontal and vertical green lines at zero were added to identify the four quadrants in each of the four plots. Sample size was n = 3-4 for all exposure groups, except for n = 2 on day-4 PFBS 10 µM and PFOA 10 µM, and n = 2 on day-10 PFDS 0.02 and 100 µM. Additional principal component analyses explored each chemical individually within a time point (Supplemental File 8A, B). Hierarchical cluster analysis was also performed for each time point and can be found in Supplemental Files 8C, and 8D.

A

Activation z-score
-3.606   3.000

PFOA 20μM DAY 1
PFOA 20μM DAY 4
PFOA 20μM DAY 10
PFOA 20μM DAY 14
PFBS 20μM DAY 1
PFBS 20μM DAY 4
PFBS 20μM DAY 10
PFBS 20μM DAY 14
PFOS 20μM DAY 1

LPS/IL-1 Mediated Inhibition of RXR Function
EIF2 Signaling
Stearate Biosynthesis I (Animals)
Fatty Acid β-oxidation I
Tryptophan Degradation III (Eukaryotic)
Glutaryl-CoA Degradation
Systemic Lupus Erythematosus In T Cell Signaling Pathway
Complement System
Zymosterol Biosynthesis
CD28 Signaling in T Helper Cells
Calcium-induced T Lymphocyte Apoptosis
Neuroinflammation Signaling Pathway
Dendritic Cell Maturation
iCOS-iCOSL Signaling in T Helper Cells
Xenobiotic Metabolism CAR Signaling Pathway
Nicotine Degradation II
Nicotine Degradation III
Superpathway of Melatonin Degradation
Melatonin Degradation I
Superpathway of Cholesterol Biosynthesis
Cholesterol Biosynthesis II (via 24,25-dihydrolanosterol)
Cholesterol Biosynthesis III (via Desmosterol)
Cholesterol Biosynthesis I
LXR/RXR Activation
Superpathway of Geranylgeranyldiphosphate Biosynthesis I
Mevalonate Pathway I
PKCθ Signaling in T Lymphocytes
Role of NFAT in Regulation of the Immune Response
Serotonin Degradation
Acetone Degradation I (to Methylglyoxal)
Estrogen Biosynthesis
Bupropion Degradation

Figure 4. Pathway analysis of the differentially expressed genes for all time points at the 20 µM concentration (A) or for the highest non-cytotoxic concentrations (B). For (B) concentrations were 20 µM for PFOA and PFOS, and 100 µM for PFBS and PFDS. Filters for this analysis were set to z-score ≥ absolute 2.0 and Benjamin–Hochberg adjusted p-value ≤ 0.05. Blue represents inhibited pathways and orange represents activated pathways. n = 3-4 for all exposure groups, except for n = 2 on day-10 PFDS 100 µM. Supplemental File 11 A & B illustrates a less stringent pathway analysis for all PFAS over all time points (no z-score applied). Supplemental File 12 illustrates a more detailed pathway analysis for PFOS over all time points and concentrations.

Figure 5. Upstream regulator predictions from the differentially expressed genes for the top non-cytotoxic concentration of each PFAS. The z-score filters were set to 3.0 with a Benjamini Hochberg adj p-value of 0.05 for this analysis. Blue represents inhibited upstream regulators and orange represents activated regulators. n = 3-4 for all exposure groups, except for n = 2 on day-10 PFDS 100 µM.

Figure 6. Median overall gene benchmark concentration (BMC) (colored dot) for all the genes that could be modeled and 95% confidence intervals for each of the PFAS. n = 3-4 for all treatment groups, except for n = 2 on day- 4 for PFBS 10 µM and PFOA 10 µM, and day-10 PFDS 0.02 and 100 µM. Supplemental File 14 shows overall benchmark concentrations (95 % CI) by day. A summary of the distribution for all of the modeled genes exposed to PFAS over 1, 4, 10, and 14 days is in Supplemental File 13, and a summary of sample replicates is in Supplemental File 6.

Figure 7. Gene accumulation plots showing the benchmark concentrations (BMC) for each gene that could be modeled for each of the PFAS across time points. n = 3-4 for all exposure groups, except for n = 2 for day-4 PFBS 10 µM and PFOA 10 µM and day-10 PFDS, 0.02 and 100 µM. Supplemental File 14 summarizes the benchmark concentrations (and 95 % CI) by day. A summary of the distribution for all of the modeled genes exposed to PFAS over 1, 4, 10, and 14 days is in Supplemental File 13, and a summary of sample replicates is in Supplemental File 6.

Figure 8. Boxplots of administered equivalent doses (mg/kg-BW/day) of all transcripts modeled for PFOA and PFOS are shown. These were used to determine the Bioactivity Exposure Ratio (BER) based on either the lowest 5th percentile administered equivalent dose (blue diamond) or the median administered equivalent dose of the gene-set from the most sensitive pathway (red diamond). These were compared with the estimated daily exposure of the upper limit of the daily US population exposure (95th percentile) obtained from the US EPA CompTox dashboard (red dash). A range of administered equivalent doses was included as a representation of apical points of departure that were derived from non-cancer animal in vivo data (green crossbar) from Health Canada drinking water guidelines (Health Canada 2018a, and 2018b).