# Predicting COVID-19 Severity with a Specific Nucleocapsid Antibody plus Disease Risk Factor Score

**Authors:**

S. Sen[1†], E.C. Sanders[2†], K.N. Gabriel[1†], B.M. Miller[2], H.M. Isoda[2], G.S. Salcedo[2], J.E. Garrido[1], R.P. Dyer[1], R. Nakajima[3], A. Jain[3], A.-M. Caldaruse[7], A.M. Santos[2], K. Bhuvan[2], D.F. Tifrea[4], J.L. Ricks-Oddie[5,6], P.L. Felgner[3], R.A. Edwards[4], S. Majumdar[2], and G.A. Weiss[1,2,7*]

**Author Affiliations:**

[1] Department of Molecular Biology & Biochemistry, University of California, Irvine, Irvine CA 92697-3900 USA
[2] Department of Chemistry, University of California, Irvine, Irvine CA 92697-2025 USA
[3] Department of Physiology and Biophysics, University of California, Irvine, Irvine CA 92697-4560 USA
[4] Department of Pathology and Laboratory Medicine, University of California, Irvine, Irvine CA 92697-4800 USA
[5] Center for Statistical Consulting, Department of Statistics, University of California, Irvine, Irvine CA 92697-1250 USA
[6] Biostatics, Epidemiology and Research Design Unit, Institute for Clinical and Translational Sciences, University of California, Irvine, Irvine CA 92697-4094 USA
[7] Department of Pharmaceutical Sciences, University of California, Irvine, Irvine CA 92697-3958 USA

[†] S. Sen, E.C. Sanders, and K.N. Gabriel contributed equally. Author order was determined in order of increasing seniority.
* Address correspondence to gweiss@uci.edu

**Corresponding Author:**

*GAW e-mail: gweiss@uci.edu, Tel: +1-949-824-5566, address: 1102 Natural Sciences 2, Irvine, CA 92697-2025, USA

**Preprint Information:**

**Author Contributions:**

S.S., E.C.S., K.N.G., S.M., P.L.F., and G.A.W. designed research; S.S., E.C.S., K.N.G., B.M.M., H.M.I., G.S.S., J.E.G., R.P.D., A.M.S., K.B., R.N., and A.J. performed research; S.S., E.C.S, K.N.G., and G.A.W. analyzed data; J.L.R. advised on statistical analysis;

40 D.F.T. and R.A.E. collected patient samples and advised on patient clinical data
41 analysis; and S.S., E.C.S., K.N.G., S.M., and G.A.W. wrote the manuscript.
42
43

44 **Competing Interest Statement:**
45
The authors declare the following competing financial interest(s): P.L.F., R.N., and A.J.
have a financial interest in a company, Nanommune Inc., that is commercializing the
COVAM technology. Nanommune partners with Sino Biological Inc. (Beijing, China) for
expression and purification of COVAM antigens used in this study. The terms of this
arrangement have been reviewed and approved by the University of California, Irvine in
accordance with its conflict of interest policies.
52
53
54

55 **Keywords:**
56 coronaviruses | SARS-CoV-2 | prognostic | epitope mapping | phage display

58 **This PDF file includes:**

59       Main Text
60       Figures 1 to 4
61
62

**Abstract:**

Effective methods for predicting COVID-19 disease trajectories are urgently

needed. Here, ELISA and coronavirus antigen microarray (COVAM) analysis mapped

antibody epitopes in the plasma of COVID-19 patients (n = 86) experiencing a wide-

range of disease states. The experiments identified antibodies to a 21-residue epitope

from nucleocapsid (termed Ep9) associated with severe disease, including admission to

the ICU, requirement for ventilators, or death. Importantly, anti-Ep9 antibodies can be

detected within six days post-symptom onset and sometimes within one day.

Furthermore, anti-Ep9 antibodies correlate with various comorbidities and hallmarks of

immune hyperactivity. We introduce a simple-to-calculate, disease risk factor score to

quantitate each patient's comorbidities and age. For patients with anti-Ep9 antibodies,

scores above 3.0 predict more severe disease outcomes with a 13.42 Likelihood Ratio

(96.7% specificity). The results lay the groundwork for a new type of COVID-19

prognostic to allow early identification and triage of high-risk patients. Such information

could guide more effective therapeutic intervention.

**Significance statement:**

The COVID-19 pandemic has resulted in over two million deaths worldwide.

Despite efforts to fight the virus, the disease continues to overwhelm hospitals with

severely ill patients. Diagnosis of COVID-19 is readily accomplished through a multitude

of reliable testing platforms; however, prognostic prediction remains elusive. To this

end, we identified a short epitope from the SARS-CoV-2 nucleocapsid protein and also

a disease risk factor score based upon comorbidities and age. The presence of

antibodies specifically binding to this epitope plus a score cutoff can predict severe

COVID-19 outcomes with 96.7% specificity.

**Main Text:**

**Introduction**

The COVID-19 pandemic has triggered an ongoing global health crisis. More

than 108.2 million confirmed cases and 2.3 million deaths have been reported

worldwide as of February 16, 2021 (1). The virus that causes COVID-19, severe acute

respiratory syndrome coronavirus (SARS-CoV-2), belongs to the same family of viruses

responsible for respiratory illness linked to recent epidemics – severe acute respiratory

syndrome (SARS-CoV-1 termed SARS here) in 2002-2003 and Middle East respiratory

syndrome (MERS) in 2012 (2). The current and previous outbreaks suggest

coronaviruses will remain viruses of concern for global health.

Many risk factors and comorbidities, including age, sex, hypertension, diabetes,

and obesity, can influence COVID-19 patient outcomes (3). Analysis of patient immune

parameters has linked disease severity to elevated levels of biomarkers for

inflammation (c-reactive protein and cardiac troponin I), organ damage (aspartate

aminotransferase, abbreviated AST, and hypoalbuminemia), immune hyperactivity (IL-6

and IL-10), and clotting (D-dimer) (4). Mortality in COVID-19 is often caused by multi-

organ injury and severe pneumonia attributed to an excessive immune response,

termed a cytokine storm (5). Given the rapid and wide spectrum of COVID-19 disease

progression, a more precise prognostic linking disease risk factors and specific immune

responses can potentially predict disease trajectories and guide interventions.

One hypothesis to explain differences in severity of COVID-19 implicates weakly

binding, non-neutralizing antibodies (Abs) to SARS-CoV-2 proteins (6). However, the

potential harm of these suboptimal Abs in COVID-19 patient outcomes remains ill-

defined. Furthermore, a recent review on antibody-dependent enhancement of SARS-

CoV-2 stated, "At present, there are no known clinical findings, immunological assays or

biomarkers that can differentiate any severe infection from immune-enhanced disease,

whether by measuring antibodies, T cells or intrinsic host responses (7)." This

conclusion inspired our study.

SARS-CoV-2 encodes four major structural proteins – spike (S), nucleocapsid

(N), membrane (M), and envelope (E). The S, N, and M proteins from SARS elicit an

Ab-based immune response (8, 9). The Ab response and its effects on disease

progression in SARS-CoV-2 remain under investigation (10, 11). Bioinformatics has

predicted >55 Ab binding epitope regions from SARS-CoV-2 (12–17). The epitopes for

N, M or E proteins are less well-characterized than for S protein. Several studies have

reported comprehensive epitope mapping of the antibody response to SARS-CoV-2

(18–21). Here, we sought to characterize epitopes from SARS-CoV-2 and their

correlations with disease severity. ELISAs with phage-displayed epitopes (phage

ELISAs) and coronavirus antigen microarray (COVAM) analysis (22) examined plasma

samples from COVID-19 patients (n = 86). The results demonstrate that Abs to a

specific epitope from N protein plus disease risk factors strongly correlate with COVID-

19 disease severity.

**Results**

*Design and production of candidate epitopes*

Twenty-one putative SARS-CoV-2 epitopes were predicted through

bioinformatics (12–14) and structure-based analysis. The candidate epitopes span the

S, N, M, or E proteins and are on average 34 amino acids in length (**Fig. 1** and **Table**

**S1**). These epitopes were phage-displayed as fragments of the full-length protein and

were likely unstructured. Here, epitope refers to the predicted region of the antigenic

protein recognized by the antibody's paratope. The structure of S protein bound to a

neutralizing antibody (23, 24) provided the starting point for 12 of these antibody

epitopes. Epitopes were designed to potentially isolate even suboptimal Abs binding to

small portions of these structural proteins; such suboptimal Abs were hypothesized to

provide insight into disease severity. After display of each potential epitope on the

surface of phage, the quality of the epitopes was evaluated by PCR, DNA sequencing,

and QC ELISA (**Fig. S1**). A total of 18 phage-displayed, putative epitopes passed

quality control PCR, and were selected for further study.

146
**Fig. 1. Predicted SARS-CoV-2 epitopes examined by phage ELISA.** Structural
models (gray) of the SARS-CoV-2 **A**) S, **B**) N, **C**) M, or **D**) E proteins illustrate our
epitope design (colored). Sequence Ep13* has the mutation D614G, which increases
the fitness of SARS-CoV-2 (25,26). The depicted structural models were derived from
an S protein X-ray structure (PDB: 6VXX) (23) or computation modeling of N, M, and E
proteins (Protein Gene Bank: QHD43423, QHD43419, and QHD43418, respectively)
(27). **Table S1** provides sequences, sources, and rationale for selections.
154

*Mapping epitope binding to anti-SARS-CoV-2 Abs*

Plasma from COVID-19 patients was subjected to ELISAs with the phage-

displayed SARS-CoV-2 epitopes **(Fig. 2A)**. Unless otherwise indicated (e.g., healthy

controls), plasma refers to samples from PCR-verified, COVID-19 patients. In this initial

assay, plasma was pooled, diluted 100-fold, and coated on a microtiter plate (3 pools of

n = 5 patients per pool). Nonspecific interactions were blocked (ChonBlock), and phage-

displayed epitopes were added for ELISA. The resultant data were normalized by signal

from the corresponding negative control (phage without a displayed epitope). Seven

candidate epitopes from the pooled patients were further investigated with a larger

number of individual patient samples (n = 28) (**Fig. 2B**). The strongest, reproducible

binding was observed for three epitopes from M (Ep6), N (Ep9), and S (Ep20) proteins.

Additional COVID-19 plasma samples were profiled for binding to these three epitopes

(n = 86 total) (**Fig. 2B**).

Only the Ep9 epitope from N protein demonstrated robust, statistically significant

antibody binding in 27% of patients (n = 186) (**Figs. 2B** and **S2**). Of these patients, 100

did not have corresponding health information and were not analyzed further in this

report. To test non-phage displayed epitopes, dose-dependent binding to Ep9 fused to

eGFP (eGFP-Ep9) or to full-length N protein demonstrates that αEp9 IgGs bind its

antigen with $EC_{50}$ = 3.22 nM (95% CI = 2.49 to 4.14 nM). This experiment examines

plasma samples with the highest IgG response against the N protein in the COVAM

assay. Patients without αEp9 Abs have roughly the same level of binding to N protein as

observed for αEp9 Abs binding to Ep9. However, such αEp9 Abs appear to add to N

protein binding by antibodies; we observe approximately two-fold increase in apparent

antibody binding levels for N protein if the patient also has αEp9 Abs (**Fig. S3**).

Therefore, the αEp9 response we report cannot solely be due to N protein antigenicity.

In patients for whom longitudinal samples were available, the highest levels of αEp9

Abs were observed at days 1 to 14 post-symptom onset (n = 11) and were detectable

within 6 days (**Fig. 2C**). In four of these patients, αEp9 Abs persisted after day 14.

183
**Fig. 2. Mapping COVID-19 patient antibody responses with phage-displayed**
**SARS-CoV-2 epitopes. A)** This phage ELISA with the indicated epitopes (x-axis)
examined plasma pooled from patients (n = 3 pools of 5 patients each, 2 technical
replicates). STOP4 is the phage negative control. **B)** The epitopes with the highest
signals were then further examined by ELISA with plasma from individual patients (n as
indicated). **C)** With samples from individual patients (designated as P# and by color)
collected at the indicated times, αEp9 Abs were measured. The subset of patients
shown here comprise all samples for which longitudinal data was available. **D)** Phage
ELISA with samples from patients with strong αEp9 Ab response (two from the
longitudinal study and one from the patient population) examines cross-reactive binding
to Ep9 or Ep9 orthologs from the indicated coronaviruses (x-axis, 3 technical replicates).
The arrow on the y-axis and gray line (panels B and C) represents the negative control
used for normalizing the data. Error bars represent SEM (panels A, B, and D) or range
of two measurements (panel C).
198

199 *Cross-reactivity of αEp9 Abs against orthologous epitopes from other coronaviruses*

200 Next, the cross-reactivity of αEp9 Abs was examined with Ep9-orthologs from

201 four phylogenetically related coronaviruses known to infect humans (**Fig. S4A**).

202 Specifically, plasma with αEp9 Abs (n = 3) and pooled plasma from healthy individuals

203 (n = 5) were assayed. The Ep9 epitopes from SARS-CoV-2 and SARS have 90% amino

acid sequence homology. Unsurprisingly, this high degree of similarity resulted in a cross-reactive Ep9 epitope, and a strong antibody response was observed to Ep9 epitopes from both viruses (**Fig. 2D**). The coronaviruses, MERS, HKU-1, and NL63 have 52%, 43%, and 8% sequence homology to SARS-CoV-2 Ep9, respectively (**Fig. S4B**). These more distantly related orthologs exhibited no cross-reactivity with the αEp9 Abs. Furthermore, no response was observed to Ep9 in pooled plasma from healthy individuals.

The protein microarray COVAM analysis is a high-throughput serological test for SARS-CoV-2 Ab cross-reactivity with a panel of 61 antigens from 23 strains of 10 respiratory tract infection-causing viruses (22). In this assay, each antigen was printed onto microarrays, probed with human plasma, and analyzed with an ArrayCam imager. COVAM distinguishes between IgG and IgM Abs binding to the full-length N protein (**Fig. S5** and **S6**, respectively). Thus, the COVAM analysis complements the phage ELISA by expanding the scope of antigens surveyed and adding Ab serotype information. The ELISA and COVAM data both demonstrate that αEp9 Abs are highly specific for lineage B betacoronaviruses, and unlikely to be found in patients before their infection with SARS-CoV-2.

*More severe disease and poorer outcomes for αEp9 patients*

Direct comparison of data with full-length N protein from COVAM and Ep9 phage ELISA (n = 40 patients assayed with both techniques) reveals five unique categories of patients (**Fig. 3A**). To enable this comparison, raw data from each assay was normalized as a percentage of the negative control. Category 1 consists of patients

without Abs to the N protein. The next categories include patients with IgMs (Category

2) or IgGs (Category 3) binding to N protein, but not Ep9, termed non-Ep9 αN Abs.

Category 4 includes patients with αEp9 Abs (both IgMs and IgGs). Category 5 patients

have exclusively IgG αEp9 Abs. The αEp9 Abs are only found in patients with IgMs or

IgGs against full-length N protein from the COVAM assay; the COVAM analysis thus

independently corroborate the phage ELISAs (**Fig. 3A**).

Interestingly, the patients with αEp9 Abs suffer more prolonged illness and worse

clinical outcomes compared to patients with non-Ep9 αN Abs or no αN Abs. In this

study, severe COVID-19 cases are defined as resulting in death or requiring admission

to the ICU or intubation. The fraction of severe COVID-19 cases was 2.5 times higher in

αEp9 Abs patients than non-Ep9 αN Abs patients (**Fig. 3B, yellow panel**); the

differences in proportions of severe and non-severe αN-positive patients with or without

αEp9 Abs are statistically significant ($p < 0.030$, Fisher's exact test). Patients without αN

Abs (Category 1) had less severe symptoms. The αEp9 Abs patients also had longer

durations of symptoms and hospital stays relative to non-Ep9 αN Abs and no αN Abs

patients (**Figs. 3C** and **D**). A larger data set of patient plasma analyzed by phage ELISA

confirmed this conclusion ($p < 0.0013$, Fisher's exact test) (**Fig. 3B, blue panel**). Our

data further demonstrates that asymptomatic COVID-19 patients (n = 3) also tested

negative for αEp9 Abs (**Table S2**). The data also reveals early seroconversion of αEp9

IgGs (**Fig. 3E**), but not αEp9 IgMs (**Fig. 3F**).

**Fig. 3. Patients with αEp9 Abs have more severe disease. A)** Normalized and categorized data from measurements by COVAM (IgMs in yellow, IgGs in green) and Ep9 phage ELISA (blue). ANOVA comparing COVAM to ELISA with Dunnett's multiple comparisons yields p-values of **<0.01, ****<0.0001, or ns: not significant. **B)** Disease severity (color) binned by antibody response (COVAM in yellow, or ELISA in blue). Statistical analysis reveals significant differences between distributions of severe and non-severe disease comparing patient categories, $p<0.01$ ($\chi^2$) and $p<0.001$ (Fisher's exact test) for COVAM and ELISA, respectively. Patients with αEp9 Abs are **C)** symptomatic for longer durations and **D)** spend more days in the hospital than those with other αN Abs or no αN Abs. ANOVA with Tukey's multiple comparisons yields p-values of *<0.05 and **<0.01. One outlier (black) (ROUT = 0.1%) was omitted from statistical calculations for panels C and D. **E)** The αN IgG appear at high levels early in the course of disease only for αEp9-positive patients, but are lower in non-Ep9, αN-positive patients. After >15 days post symptom onset, αN IgG levels increase for both groups of patients. **F)** However, IgM levels do not change significantly. Error bars depict SEM with the indicated number of patients (n, numbers above columns).

*Strong correlation between disease severity and comorbidities in patients with αEp9 Abs*

We compared risk factors, clinical parameters, and disease outcomes among patients with αEp9 Abs (n = 23) (**Figs. 4A** and **S7**). A *disease risk factor score* (DRFS) was developed to evaluate the relationship between clinical preconditions and disease severity in patients with αEp9 Abs. The DRFS quantifies a patient's age, sex, and pre-

existing health conditions associated with COVID-19 disease severity and mortality.

Risk factors include hypertension, diabetes, obesity, cancer, and chronic conditions of

the following: cardiac, cerebrovascular, kidney, and pulmonary (28–31). Using the *age*

*score* from the Charlson Comorbidity Index (32) yields a patient's DRFS as:

$$DRFS = Σ \text{ (\# of risk factors) + (age score)}$$

where each risk factor is valued as either 0 or 1 if absent or present, respectively. The

DRFS of patients with αEp9 Abs strongly correlates with COVID-19 disease severity

(Pearson's r = 0.72, p-value <0.0001, and $R^2$= 0.52) (**Fig. 4A**). The correlation in

patients without αEp9 Abs is weak (r = 0.30, p-value = 0.089, $R^2$= 0.018) (**Fig. 4A**).

Amongst patients with αEp9 Abs (n = 23), a DRFS ≥3 can determine disease severity

with 92.3% sensitivity (1/13 false negatives) and 80% specificity (2/10 false positives)

(**Fig. 4B**). In the entire study cohort (n = 86), patients with αEp9 Abs and DRFS ≥3 (n =

11) have severe disease with a high degree of specificity (96.7%) and a sensitivity of

44%. Notably, DRFS predicts disease severity only for patients with αEp9 Abs (n = 23),

and patients without such Abs (n = 63) had no correlation with disease outcomes.

Examining key contributors to high DRFS, the presence of αEp9 Abs correlates

with more severe disease in patients who have hypertension, diabetes, or age >50

years. Such correlation is not observed for patients lacking αEp9 Abs (**Fig. 4C**). Such

risk factors are prevalent at roughly the same percentages in both populations of

patients (**Table S2**). Thus, these risk factors are particularly acute for patients with αEp9

Abs.

293
**Fig. 4. Correlation between disease severity and risk factors in patients with αEp9 Abs. A)** The relationship between DRFS and disease severity of COVID-19 patients with αEp9 Abs (blue) or no αEp9 Abs (gray). Each data point represents one patient. The solid lines indicate linear regression fits with 95% confidence intervals (dotted lines), and Pearson's r-value as noted. **B)** Correlation of disease severity with DRFS in patients with αEp9 Abs. The data depicts a significant correlation between DRFS and disease severity in patients with αEp9 Abs (blue), but not in patients lacking αEp Abs (gray). In αEp9 patients, a DRFS threshold of 3.0 can predict severe disease (red). Two-tailed, parametric t-tests were conducted to compare non-severe and severe disease outcomes of patients with and without αEp9 Abs, where ****p<0.0001. The error bars represent SD with the indicated n. **C)** The color-indicated risk factors (diabetes, hypertension, and age score) are depicted on the x-axis as the fractions of patients in each disease severity category (y-axis). Numbers indicate total patients (n) without αEp9 Abs (left) or with αEp9 Abs (right). The prevalence of risk factors (colors) increases with disease severity in patients with αEp9 Abs, but not in patients without these Abs. **D)** Patients with αEp9 Abs and DRFS ≥3 are predisposed to increased COVID-19 severity and poorer outcomes.

*High levels of inflammatory cytokine and tissue damage markers in patients with αEp9*

*Abs*

COVID-19 patients can have elevated serum concentrations of >20 inflammatory cytokines and chemokines (33). However, information on the cytokine levels and the

association with tissue damage and worse COVID-19 outcomes have been inconsistent

(33–35). For patients with IL-6 concentrations measured in plasma, patients with (n = 8)

or without (n = 11) αEp9 Abs were compared. Interestingly, the comparison uncovered

a strong positive sigmoidal association between IL-6 and AST unique to patients with

αEp9 Abs ($R^2$ = 0.968, Spearman's r = 1.0, p-value <0.0001, n = 8) (Blue line, **Fig.**

**S8A**); correlation of IL-6 and AST in patients with αEp9 Abs remains strong even after

removal of the data point at the highest IL-6 concentration. Conversely, a slight negative

trend is observed in patients lacking αEp9 Abs (Spearman's r = -0.575, p-value=

0.0612, n = 13). Thus, the presence of αEp9 Abs can disambiguate the sometimes

contradictory association of IL-6 with disease severity.

**Discussion**

This study introduces a two-step test as a prognostic for predicting COVID-19

disease severity and its worst outcomes. Specifically, αEp9 Abs can effectively predict

severe disease (specificity 83.6%). However, combining presence of αEp9 Abs with

DRFS ≥3 provides much higher specificity (96.7%) for predicting severe disease.

Previously, αN IgGs have been recognized as a focal site for an antibody response (18,

19, 21, 36) and associated with disease severity and poor outcomes (11, 36, 37).

The present investigation expands on previous reports that recognize various

regions of the RNA binding domain of N protein as focal sites for anti-SARS-CoV-2

antibody response. For example, the phage display-based VirScan identified an epitope

region spanning residues 141-196 and microarrays further isolated peptides including

residues  134-171,  155-171,  153-190,  and  153-171  (18,  19,  21).  The  above

investigations, however, do not find correlations between any these epitopes and

disease severity. Our results are confirmed by observations from a patient cohort in

Singapore, which identify an epitope (residues 153-170) very similar to Ep9 (residues

152-172) and shows a correlation between antibody response against the epitope and

pneumonia and the tissue damage markers (CRP and LDH) (20). In our investigation,

we examine in-depth patient clinical histories, test results, disease outcomes ranging

from asymptomatic to fatal, and longer longitudinal profiling post-symptom onset, to

determine the association of a larger subset of markers and risk factors. Such data

allows calculation of the DRFS. Together with the presence of αEp9 Abs, patient DRFS

allows early discrimination of severe from non-severe disease outcomes. Additionally,

fine epitope mapping demonstrates that αEp9 Abs strongly and uniquely correlate with

COVID-19 disease severity relative to other αN Abs.

We hypothesize that the underlying mechanism relating αEp9 Abs to increased

disease severity involves an overzealous immune response. Specifically, we observe

early seroconversion and strong early upregulation of αEp9 IgGs (**Fig. 3E**). Similar IgG

observations have been correlated with poor viral neutralization and clearance, resulting

in increased COVID-19 severity (10, 37, 38). Also, high levels of IL-6 are observed for

αEp9-positive patients with increased levels of the tissue damage marker AST; this

correlation does not exist for patients lacking αEp9 Abs (**Fig. S8A**). The sensitivity to IL-

6 concentration before AST-monitored organ damage suggests anti-IL-6 therapeutics

could be an effective management in the early and rapidly progressive stages of

respiratory distress for αEp9-positive patients (33, 39–43). Since binding to N protein by

αEp9 antibodies is unlikely to enhance uptake of SARS-CoV-2, an antibody-dependent

enhancement mechanism could invoke antigen uptake by macrophages. This

mechanism could stimulate complement activation and the cytokine storm observed

here as elevated IL-6 response. Further investigation is required to determine the basis

for increased disease severity in αEp9 patients.

The data demonstrate that αEp9 positive patients with DRFS ≥3 are 13.42 times

(Likelihood Ratio) more likely to have severe COVID-19 disease symptoms within the

study cohort (n = 86). The presence of αEp9 without DRFS is less effective as a

prognostic (Likelihood Ratio of 3.17). Despite its high specificity (96.7%), the sensitivity

of this two-step test is 44% (n = 86). However, this test could predict a subset of

patients with a specific immune response (i.e., early IgG response and IL-6 dependent

immune hyperactivity), and could suggest targeted treatment options (e.g., targeting IL-

6 and its pathways).

Importantly, αEp9 Abs appear early in the course of disease. Thus, such a

prognostic could outperform traditional markers for the cytokine storm such as IL-6,

which appears 6-8 days after symptom onset (33, 41); all plasma collected from αEp9

positive patients (n = 7, **Fig. 2C**) between 1 to 6 days post-symptoms onset

demonstrate detectable levels of αEp9 IgG (≥ 2 fold over negative control). Early

detection of αEp9 Abs in patients could be used to triage and treat COVID-19 prior to

the onset of its most severe symptoms; delayed treatments of IL-6 targeting drugs can

decrease their efficacy or be counterproductive (33, 39–44) (**Fig. S8B**). The αEp9 Ab

biomarker could identify patients most likely to benefit from anti-IL-6 therapeutics and

avoid ineffective treatments.

This study demonstrates the usefulness of fine epitope mapping, but the

following limitations should be noted. Short linear epitopes, unlike conformational

epitopes in larger domains, might not resemble the tertiary structure of an antigen. Post-

translational modifications, such as glycosylation were omitted for the phage-displayed

S protein epitopes; the COVAM antigens, however, are produced in baculovirus or

HEK-293 cells, which could glycosylate the antigens. Our analysis is largely based upon

a population of 86 COVID-19 patients and 5 healthy individuals, with the majority of

Hispanic descent. The conclusions could be further strengthened with follow-up

investigations in a larger population. Additionally, the population examined here only

included three asymptomatic individuals, and additional testing is required to verify

absence of αEp9 Abs in such patients. The sample size of patients with multiple

antibody targets was too limited to allow correlation analysis; future investigations could

examine associations between αEp9 and other Abs. Abs recognizing other SARS-CoV-

2 structural proteins could also exhibit similar characteristics to αEp9 Abs.

Existing diagnostic platforms could readily be adapted to test for αEp9 Abs, and

the DRFS calculation is quite simple to implement (e.g., assay with eGFP-Ep9 fusion

demonstrated here). As shown here, αEp9 Abs do not recognize orthologous

sequences from closely related coronaviruses, providing good specificity for αEp9 as a

prognostic. Previous studies have shown that the high homology of N protein among

related coronaviruses can lead to high false positive rates in serodiagnostics with full-

length N antigen (45). Thus, the two-step prognostic reported here could mitigate the

worst outcomes of COVID-19, particularly for patients at high risk.

**Materials and Methods:**

Detailed materials and methods for cloning, phage purification, patient sample collection, plasma phage-antibody ELISA, serum COVAM, and statistical analysis are described in the *SI Appendix.*

**Acknowledgements:**

We thank Dr. Hung Fan and Dr. Donald Forthal for helpful conversations and the patients who donated samples. We gratefully acknowledge the support of the UCI COVID-19 Basic, Translational and Clinical Research Fund (CRAFT), the Allergan Foundation, and UCOP Emergency COVID-19 Research Seed Funding. S.S. was supported by a Public Impact Fellowship from the UCI Graduate Division. K.N.G. was supported by a National Science Foundation Graduate Research Fellowship Program (DGE-1839285). G.S.S and A.M.S thank the Minority Access to Research Careers (MARC) Program, funded by the NIH (GM-69337). J.L.R. was supported by the National Center for Research Resources and the National Center for Advancing Translational Sciences from the NIH (TR001414). D.F.T and R.A.E. were supported by the Experimental Tissue Resource, funded by the Chao Family NCI-Comprehensive Cancer Center Support Grant from the NCI (P30CA062203).

**Abbreviations:**

COVID-19, Coronavirus disease 2019
ELISA, enzyme-linked immunosorbent assay
COVAM, coronavirus antigen microarray
Ep9, epitope 9
ICU, intensive care unit
SARS-CoV-2, severe acute respiratory syndrome coronavirus
SARS-CoV-1, severe acute respiratory syndrome
MERS, Middle East respiratory syndrome
AST, aspartate aminotransferase
IL-6, interleukin 6
IL-10, interleukin 10
Abs, antibodies
S protein, SARS-CoV-2 spike glycoprotein
N protein, SARS-CoV-2 nucleocapsid protein
M protein, SARS-CoV-2 membrane protein
E protein, SARS-CoV-2 envelope protein
Phage ELISA, ELISA with phage-displayed epitopes
PCR, polymerase chain reaction
QC ELISA, quality control ELISA
PDB, protein data bank
HKU-1, human coronavirus HKU1
NL63, human coronavirus NL63
IgG, immunoglobulin G
IgM, immunoglobulin M
αEp9 Abs, anti-Ep9 antibodies
ANOVA, analysis of variance
DRFS, disease risk factor score
457

**References:**

1.    World Health Organization, "Coronavirus Disease (COVID-19)" (2021)
https:/doi.org/10.1097/jcn.0000000000000710.

2.    A. Llanes, *et al.*, Betacoronavirus genomes: How genomic information has been
used to deal with past outbreaks and the covid-19 pandemic. *Int. J. Mol. Sci.* **21**,
1–28 (2020).

3.    S. Richardson, *et al.*, Presenting Characteristics, Comorbidities, and Outcomes
among 5700 Patients Hospitalized with COVID-19 in the New York City Area. *J.*
*Am. Med. Assoc.* **323**, 2052–2059 (2020).

4.    B. Gallo Marin, *et al.*, Predictors of COVID-19 severity: A literature review. *Rev.*
*Med. Virol.* (2020).

5.    P. Song, W. Li, J. Xie, Y. Hou, C. You, Cytokine storm induced by SARS-CoV-2.
*Clin. Chim. Acta* **509**, 280–287 (2020).

6.    A. Iwasaki, Y. Yang, The potential danger of suboptimal antibody responses in

COVID-19. *Nat. Rev. Immunol.* **20**, 339–341 (2020).

7.    A. M. Arvin, *et al.*, A perspective on potential antibody-dependent enhancement of
SARS-CoV-2. *Nature* **584**, 353–363 (2020).

8.    L. R. Huang, *et al.*, Evaluation of antibody responses against SARS coronaviral
nucleocapsid or spike proteins by immunoblotting or ELISA. *J. Med. Virol.* **73**,
338–346 (2004).

9.    Y. He, Y. Zhou, P. Siddiqui, J. Niu, S. Jiang, Identification of immunodominant
epitopes on the membrane protein of the severe acute respiratory syndrome-
associated coronavirus. *J. Clin. Microbiol.* **43**, 3718–3726 (2005).

10.    Q. X. Long, *et al.*, Antibody responses to SARS-CoV-2 in patients with COVID-19.
*Nat. Med.* **26**, 845–848 (2020).

11.    M. Batra, *et al.*, Role of IgG against N-protein of SARS-CoV2 in COVID19 clinical
outcomes. *medRxiv* (2020).

12.    E. Fast, R. B. Altman, B. Chen, Potential T-cell and B-cell Epitopes of 2019-nCoV.
*bioRxiv* (2020).

13.    A. Grifoni, *et al.*, A Sequence Homology and Bioinformatic Approach Can Predict
Candidate Targets for Immune Responses to SARS-CoV-2. *Cell Host Microbe* **27**,
671–680 (2020).

14.    V. Baruah, S. Bose, Immunoinformatics-aided identification of T cell and B cell
epitopes in the surface glycoprotein of 2019-nCoV. *J. Med. Virol.* **92**, 495–500
(2020).

15.    M. Zheng, L. Song, Novel antibody epitopes dominate the antigenicity of spike
glycoprotein in SARS-CoV-2 compared to SARS-CoV. *Cell. Mol. Immunol.* **17**,
536–538 (2020).

16.    B. Tilocca, *et al.*, Comparative computational analysis of SARS-CoV-2
nucleocapsid protein epitopes in taxonomically related coronaviruses. *Microbes*
*Infect.* **22**, 188–194 (2020).

17.    A. Rakib, *et al.*, Immunoinformatics-guided design of an epitope-based vaccine
against severe acute respiratory syndrome coronavirus 2 spike glycoprotein.
*Comput. Biol. Med.* **124**, 103967–103983 (2020).

18.    L. Wang, *et al.*, Serological Responses to Human Virome Define Clinical
Outcomes of Italian Patients Infected with SARS-CoV-2. *medRxiv* (2020).

19.    C. R. Zamecnik, *et al.*, ReScan, a Multiplex Diagnostic Pipeline, Pans Human
Sera for SARS-CoV-2 Antigens. *Cell Reports Med.* **1**, 100123–100133 (2020).

20.    S. Naqiah Amrun, *et al.*, Linear B-cell epitopes in the spike and nucleocapsid
proteins as markers of SARS-CoV-2 exposure and disease severity.
*EBioMedicine* **58**, 102911–102919 (2020).

21.    A. Musicò, *et al.*, SARS-CoV-2 epitope mapping on microarrays highlights strong
immune-response to n protein region. *Vaccines* **9**, 1–11 (2021).

22.    P. N. Hedde, *et al.*, A modular microarray imaging system for highly specific

COVID-19 antibody testing. *Lab Chip* **20**, 3302–3309 (2020).

23. A. C. Walls, *et al.*, Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* **181**, 281–292 (2020).

24. J. Lan, *et al.*, Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature* **581**, 215–220 (2020).

25. R. S. Baric, Emergence of a Highly Fit SARS-CoV-2 Variant. *N. Engl. J. Med.* **383**, 2684–2686 (2020).

26. J. A. Plante, *et al.*, Spike mutation D614G alters SARS-CoV-2 fitness. *Nature*, 1–6 (2020).

27. W. Zheng, *et al.*, Deep-learning contact-map guided protein structure prediction in CASP13. *Proteins Struct. Funct. Bioinforma.* **87**, 1149–1164 (2019).

28. W. Tian, *et al.*, Predictors of mortality in hospitalized COVID-19 patients: A systematic review and meta-analysis. *J. Med. Virol.* **92**, 1875–1883 (2020).

29. A. Simonnet, *et al.*, High Prevalence of Obesity in Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) Requiring Invasive Mechanical Ventilation. *Obesity* **28**, 1195–1199 (2020).

30. L. Y. W. Lee, *et al.*, COVID-19 prevalence and mortality in patients with cancer and the effect of primary tumour subtype and patient demographics: a prospective cohort study. *Lancet Oncol.* **21**, 1309–1316 (2020).

31. L. Zhang, *et al.*, Clinical Course and Mortality of Stroke Patients With Coronavirus Disease 2019 in Wuhan, China. *Stroke* **51**, 2674–2682 (2020).

32. M. E. Charlson, P. Pompei, K. L. Ales, C. R. MacKenzie, A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *J. Chronic Dis.* **40**, 373–383 (1987).

33. L. Lu, *et al.*, Preventing Mortality in COVID-19 Patients: Which Cytokine to Target in a Raging Storm? *Front. Cell Dev. Biol.* **8**, 1–8 (2020).

34. M. J. Cummings, *et al.*, Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *Lancet* **395**, 1763–1770 (2020).

35. Y. Yang, *et al.*, Plasma IP-10 and MCP-3 levels are highly associated with disease severity and predict the progression of COVID-19. *J. Allergy Clin. Immunol.* **146**, 119–127 (2020).

36. E. Shrock, *et al.*, Viral epitope profiling of COVID-19 patients reveals cross-reactivity and correlates of severity. *Science (80-. ).* **370**, 1–22 (2020).

37. W. Tan, *et al.*, Viral Kinetics and Antibody Responses in Patients with COVID-19. *medRxiv* (2020).

38. H. wei Jiang, *et al.*, SARS-CoV-2 proteome microarray for global profiling of COVID-19 specific IgG and IgM responses. *Nat. Commun.* **11**, 3581 (2020).

39. M. Benucci, *et al.*, COVID-19 pneumonia treated with Sarilumab: A clinical series

of eight patients. *J. Med. Virol.* **92**, 2368–2370 (2020).

40.  E. J. Kooistra, *et al.*, Anakinra treatment in critically ill COVID-19 patients: a
prospective cohort study. *Crit. Care* **24**, 1–12 (2020).

41.  A. Langer-Gould, *et al.*, Early Identification of COVID-19 Cytokine Storm and
Treatment with Anakinra or Tocilizumab. *Int. J. Infect. Dis.* **99**, 291–297 (2020).

42.  L. Castelnovo, *et al.*, Anti-IL6 treatment of serious COVID-19 disease. *Medicine
(Baltimore).* **100**, 1–6 (2021).

43.  C. Salama, *et al.*, Tocilizumab in Patients Hospitalized with Covid-19 Pneumonia.
*N. Engl. J. Med.* **384**, 20–30 (2021).

44.  V. C. Veiga, *et al.*, Effect of tocilizumab on clinical outcomes at 15 days in patients
with severe or critical coronavirus disease 2019: randomised controlled trial. *BMJ*
**372**, n84 (2021).

45.  Y. Yamaoka, *et al.*, Whole Nucleocapsid Protein of Severe Acute Respiratory
Syndrome Coronavirus 2 May Cause False-Positive Results in Serological
Assays. *Clin. Infect. Dis.* (2020).

46.  K. Murase, *et al.*, EF-Tu binding peptides identified, dissected, and affinity
optimized by phage display. *Chem. Biol.* **10**, 161–168 (2003).

47.  A. Bhasin, *et al.*, Virus Bioresistor (VBR) for Detection of Bladder Cancer Marker
DJ-1 in Urine at 10 pM in One Minute. *Anal. Chem.* **92**, 6654–6666 (2020).

48.  B. Pastorino, F. Touret, M. Gilles, X. De Lamballerie, R. N. Charrel, Heat
Inactivation of Different Types of SARS-CoV-2 Samples: What Protocols for
Biosafety, Molecular Detection and Serological Diagnostics? *Viruses* **12**, 1–8
(2020).

49.  S. Khan, *et al.*, Use of an influenza antigen microarray to measure the breadth of
serum antibodies across virus subtypes. *J. Vis. Exp.* **2019**, 1–8 (2019).

50.  A. Jain, *et al.*, Evaluation of Quantum dot immunofluorescence and a digital
CMOS imaging system as an alternative to conventional organic fluorescence
dyes and laser scanning for quantifying protein microarrays. *Protemoics* **16**,
1271–1279 (2016).

51.  R. Nakajima, *et al.*, Protein Microarray Analysis of the Specificity and Cross-
Reactivity of Influenza Virus Hemagglutinin-Specific Antibodies. *mSphere* **3**, 1–15
(2018).

52.  M. M. Mukaka, Statistics Corner: A guide to appropriate use of Correlation
coefficient in medical research. *Malawi Med. J.* **24**, 69–71 (2012).

53.  B. Korber, *et al.*, Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G
Increases Infectivity of the COVID-19 Virus. *Cell* **182**, 812–827 (2020).

**SI Appendix:** General methods and additional experimental data can be found in the *SI*

*Appendix.*

590 Figs. S1 to S8

591 Tables S1 to S4

592 References (46–53)