The role of behavioural flexibility in primate diversification

Maria J.A. Creighton[a], Dan A. Greenberg[b], Simon M. Reader[a] & Arne Ø. Mooers[b]

[a]Department of Biology, McGill University, Montreal, Quebec, Canada

[b]Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia, Canada

*Author for correspondence: maria.creighton@mail.mcgill.ca; ph: +1-514 398 7020; McGill University, Stewart Biology Building 1205 ave Docteur Penfield, Montreal, QC Canada, H3A 1B1 ORCID: https://orcid.org/0000-0003-1337-1946

DECLARATIONS OF INTEREST: None

Identifying the factors that influence species diversification is fundamental to our understanding of the evolutionary processes underlying extant biodiversity. Behavioural innovation, coupled with the social transmission of new behaviours, has been proposed to increase rates of evolutionary diversification, as novel behaviours expose populations to new selective regimes. Thus, it is believed that behavioural flexibility may play an important role in driving evolutionary diversification across animals. We test this hypothesis within the primates, a taxonomic group with considerable among-lineage variation in both species diversity and behavioural flexibility. We employ a time cut-off in our phylogeny to help account for biases associated with recent taxonomic reclassifications and compare three alternative measures of diversification rate that consider different phylogenetic depths. We find that the presence of behavioural innovation and social learning are positively correlated with diversification rates among primate genera, but not at shallower phylogenetic depths. Given that we find stronger associations when examining older rather than more recent diversification events, we suggest that extinction resistance, as opposed to speciation, may be an important mechanism linking behavioural flexibility and primate diversification. Our results contrast with work linking behavioural flexibility with diversification of birds at various phylogenetic depths. We offer a possible dispersal-mediated explanation for these conflicting patterns, such that the influence behavioural flexibility plays in dictating evolutionary trajectories differs across clades. Our results suggest that behavioural flexibility may act through several different pathways to shape the evolutionary trajectories of lineages.

KEYWORDS: behavioural drive; behavioural flexibility; evolutionary diversification;

extinction; primates; speciation

Extant species diversity is remarkably variable across the Tree of Life (Willis, 1922; Williams,

1964). For clades of the same age, differences in net diversification rate (i.e. speciation rate

minus extinction rate) ultimately drive differences in clade size. Both the external environment

(e.g. Badgley, 2010; Kozak & Wiens, 2010; Frey, 2010) and intrinsic features of a lineage (e.g.

Raikow, 1986; Heard & Hauser, 1995) can influence net diversification. Despite ongoing study,

there remains considerable debate over the factors that lead to differences in diversification rate

among lineages, and their relative importance (e.g. Lewontin, 1983; West-Eberhard, 1989; Isaac,

et al., 2005; Rabosky, 2009; see review by Wiens, 2017).

Plasticity has been repeatedly proposed to play a major role in shaping evolutionary trajectories

in general, and speciation in particular (Baldwin, 1902; Lewontin, 1983; Bateson, 1988; Wcislo,

1989; Odling-Smee, et al., 2003; West-Eberhard, 2003; Pelletier, et al., 2009; also see review in

Pfennig, et al., 2010), and theoretical modelling supports its potential influence (e.g. Hinton &

Nowlan, 1987; Anderson, 1995; Behera & Nanjundiah, 1995; Ancel, 1999; 2000). Behavioural

development and expression often allow for more rapid responses than other forms of plasticity

such as induced morphological changes (Duckworth, 2009; Snell-Rood, 2013; West-Eberhard,

2003). Thus, plasticity of behaviour has been hypothesized to play a particularly important role

in influencing evolutionary trajectories of animal lineages (Wyles, et al., 1983; Wilson, 1985;

West-Eberhard, 2003). The propensity to adopt new behaviours can greatly and quickly alter

ecological niches, exposing populations to new selective regimes. This can lead to increased trait

disparity among populations (the "behavioural drive hypothesis"; Wyles, et al., 1983; Wilson,

1985); importantly, divergent selection can also lead to eventual speciation. As a result, it has

been suggested that behaviourally flexible taxa (i.e. those taxa exhibiting high propensities for

behavioural change due to, for example, learning or readiness to transition to new conditions; Sol

& Lefebvre, 2000; Audet & Lefebvre, 2017) may undergo faster rates of speciation, and thus

overall net rates of evolutionary diversification, compared to less flexible taxa (Sol, et al., 2005;

Grant & Grant, 2008; Price, 2008; Sol & Price, 2008; Tebbich, et al., 2010). However, despite

support from theoretical modelling (e.g. Price, et al., 2003; Lachlan & Servedio, 2004; Lapiedra,

et al., 2013), the idea that behavioural flexibility enhances diversification rates remains

contested. Some dispute the extent to which behaviour plays an active role in dictating animal

diversity (e.g. Scott-Phillips, et al., 2014). Meanwhile, an alternative hypothesis posits that

behavioural flexibility may inhibit, rather than enhance, species diversification: populations that

can utilize new resources or transition to new environments are shielding their genomes from

bouts of strong directional selection (Bogert, 1949; Huey, et al., 2003).

Duckworth (2009) suggests that behavioural flexibility could both dampen and promote

evolutionary rates depending on the time scale. Under this proposed framework, behavioural

flexibility inhibits evolution on short time scales by buffering against abrupt environmental

changes that may otherwise result in a population bottleneck or strong bouts of directional

selection. Over longer time scales, the same behavioural shift can lead to speciation, either by

setting the stage for allopatric speciation or by exposing the newly situated population to novel

selective regimes (Huey, et al., 2003; Losos, et al., 2004; Duckworth, 2009; see example in

Muñoz & Losos, 2018).

Previous studies have provided support for behavioural flexibility both driving (e.g. Yeh, 2004;

Yeh & Price, 2004; Tebbich, et al., 2010, Riesch, et al., 2012; Foote, et al., 2016) and inhibiting

(e.g. Losos, et al., 2004; Weber, et al., 2004; Shultz, et al., 2005; Gonzalez-Voyer, et al., 2016)

evolution. However, many of these studies have primarily considered the effects of behavioural

flexibility on microevolutionary change at short time scales (e.g. recent speciation events or

population decline). One notable exception is in birds, where behavioural flexibility has been

associated with multiple estimates of macroevolutionary lineage diversification (Nicolakakis, et

al., 2003; Sol, et al., 2005 – also see Sol, 2003; Sayol, et al., 2019). These studies have employed

two proposed correlates of behavioural flexibility: brain size relative to body size (a structural

measure presumed and shown elsewhere to be associated with behavioural flexibility; e.g.

Lefebvre, et al., 2004) and innovation rate (a behavioural measure) taken from literature surveys.

Both large relative brain size and high innovation rates were associated with heightened

diversification in birds (Nicolakakis, et al., 2003; Sol, et al., 2005; Sayol, et al., 2019), consistent

with the idea that behavioural flexibility positively impacts diversification. Importantly, such

tests have yet to be applied across other taxa, making it difficult to generalize the role of

behaviour in shaping the Tree of Life.

Here, we explore the relationship between four proxies of behavioural flexibility and several

measures of diversification rate in primates, a taxonomic group with considerable among-lineage

variation in both behavioural flexibility (Reader & Laland, 2002; Reader, et al., 2011) and extant

species diversity (Upham, et al., 2019a; 2019b). Variables associated with diversity of other taxa

(e.g. geographic range size and latitude) have been shown to be poor predictors of primate

diversification (Arbour & Santana, 2017; Upham, et al., 2019a), leaving a great deal of what

shapes extant primate diversity unexplained. We focus on two behavioural measures of

behavioural flexibility, the presence or absence of published reports of innovation and of social

learning, and two brain size measures widely thought to be associated with ability to exhibit

flexible behaviours. Consistent with what has been reported for birds (Nicolakakis, et al., 2003;

Sol, et al., 2005; Sayol, et al., 2019) and Duckworth (2009)'s proposal that behavioural

flexibility promotes diversification events over longer evolutionary time scales, we predict that

our separate measures of behavioural flexibility will covary positively with diversification rates

across primate lineages. To examine how this association changes at different phylogenetic

depths, we examine how behavioural flexibility correlates with diversification over both

shallower and deeper time depths in our phylogeny. Understanding how behaviour and ecology

may interact to shape evolutionary patterns provides a glimpse into some of the processes that

have shaped primate biological diversity and could, in turn, dictate future diversity.

METHODS:

Data

Diversification Rate

Estimating diversification rates is challenging because it depends on an accurate assessment of

both the taxonomic richness and divergence time of a given lineage. A previous study testing

whether behavioural flexibility drives shallow divergences used subspecies per species as a

measure of subspecific diversification in birds (Sol, et al., 2005). Using this subspecies metric

could introduce considerable bias when applied in primates, however, as primate taxonomic

richness has changed drastically over recent decades (Tattersall, 2007; Groves, 2014; Rylands &

Mittermeier, 2014), with much of this change attributed to application of the 'phylogenetic

species concept' (PSC) and its tendency to raise former subspecies and variants to the full

species rank (Tattersall, 2007). Importantly, it has been suggested that these elevations in

subspecies status have been biased toward certain taxa (Isaac, et al., 2004), which would lead to

inconsistent estimates of species versus subspecies richness across lineages. Studies using other

estimates of primate diversification (i.e. diversification analyses using TreePar; Stadler, 2011)

have been hindered by the applications of the PSC, particularly when it comes to accurately

estimating shallow divergences (Springer, et al., 2012). Modern primate phylogenies are also not

reflective of modern primate taxonomies, as they include some phylogenetic species and omit

others, preventing us from using recently described evolutionary rate measures (see, e.g., Jetz, et

al., 2012; Mitchell & Rabosky, 2017) that rely on a comprehensive phylogeny with consistent

species designations among clades. Instead, we used well-resolved "lineages" that putatively

reflect stable species complexes. We started with the most widely used, dated primate tree

publicly available at the time of this study, the GenBank taxonomy consensus tree provided on

the 10kTrees website (version 3) (Arnold, et al., 2010), containing 301 tips. We then created a

time cut-off in the tree at the time when we determined a majority of robust biological species

described in Honacki et al. (1982) had evolved (1.1mya). We chose Honacki et al. (1982) as it

was the last major primate taxonomy published before the introduction of the PSC (Cracraft,

1983). After creating this time cut-off, we subsequently eliminated shallow divergences

occurring after 1.1mya from the consensus tree (see Figure S1). Each branch in the tree that was

extant at 1.1mya was retained in the tree and designated as a stable "lineage". We then

additionally pruned species from this phylogeny that were no longer recognized by modern

taxonomies (IUCN/SSC Primate Specialist Group, 2018). This resulted in 241 identifiable

lineages to compare in terms of taxonomic richness and divergence times. Using the most

recently published primate species and subspecies list from the IUCN/SSC Primate Specialist

Group (2018), we referenced taxonomic and phylogenetic works to assign each of the 705

species and subspecies to one of these 241 lineages. This allowed us to assign each lineage an

agnostic "taxon richness" score (i.e. the sum of all monotypic species and subspecies in a

lineage; see Figure S1) that accounts for the discrepancies in subspecies elevations across

lineages. Each species or subspecies listed by the IUCN/SSC Primate Specialist Group (2018)

was also assigned to one of the 66 genera named in our 10kTrees phylogeny (Arnold, et al.,

2010). After eliminating two individual species and one genus in our tree that could not be

resolved using these methods (see supplementary material), our study considered 703 taxa

(species or subspecies) assigned to 239 lineages and 65 genera.

To estimate lineage diversification rate we used the method-of-moments estimator (Magallon &

Sanderson, 2001) that divides the natural log of "taxon richness" (species and subspecies) by

lineage stem age to produce a 'Taxa per Lineage Diversification Rate' that should be less biased

by recent subspecies elevations. This method was repeated at the genus-level where the natural

log of "taxon richness" for each genus was divided by the stem age of that genus (hereafter

'Taxa per Genus Diversification Rate'). We note that all log transformations referenced hereafter

refer to natural log transformations ($\log_e$). Lastly, we created a second, and perhaps more

conservative estimate of genus diversification rate where richness scores were generated using

the number of lineages (n=239) per genus, rather than the number of taxa described by the

IUCN/SSC Primate Specialist Group (2018), hereafter 'Lineage per Genus Diversification Rate'.

A few genera were not monophyletic on our tree, and we considered these on a case-by-case

basis ultimately removing five from our genus-level analysis (see the supplementary material).

We illustrate these methods and present associated calculations using genus *Aotus* as an example

in the supplementary material to provide some considerations on the potential uses and

limitations of this approach.

Behavioural Proxies of Behavioural Flexibility

We focused on two key behaviours to infer behavioural flexibility: innovation and social

learning. Innovation and social learning are both important in determining the macroevolutionary

effects of behavioural flexibility because multiple individuals must acquire an innovation

through social learning or independent innovation to have population-level effects (Wyles, et al.,

1983; Wilson, 1985; Duckworth, 2009). In addition to facilitating the transmission of

innovations throughout a population, social learning can also be a valuable measure of

population-level behavioural flexibility on its own as it reflects the ability of individuals within a

population to pick up behaviours that are novel to them but not necessarily novel to the

population. We note that social learning and innovation are taxonomically widespread (Reader &

Biro, 2010; Snell-Rood, et al., 2015). However, we assume that species with no published

accounts of social learning or innovation are likely relying on these behaviours infrequently.

Moreover, the innovation and social learning data used here have been positively associated with

other measures of behavioural flexibility (e.g. brain size measures; Reader, et al., 2011;

Navarrete, et al., 2016). For innovation, we focus on technical innovations (classified as those

involving tool use following Navarrete, et al., 2016) because these more easily defined

behaviours have been linked to complex cognition (Overington, et al., 2009), and reports of other

classes of innovation (e.g. food type innovation) can be highly influenced by opportunistic

events (Ducatez, et al., 2015). Combined with the fact that taxa with reports of technical

innovation also tended to be those with evidence of other innovation types (see data in Navarrete,

et al., 2016), this likely makes technical innovation a robust estimate of innovativeness across

primates.

Counts of innovation and social learning per lineage came from Reader et al. (2011) and

Navarrete et al. (2016). Reader et al. (2011) established this dataset through a survey of over

4000 published articles for examples of social learning and behavioural innovation. Reader et al.

(2011, p. 1018) define an innovation as the tendency to "discover novel solutions to

environmental or social problems". These data were later subdivided into different innovation

categories by Navarrete et al. (2016), including 'technical' innovations, defined as innovations

involving tool use. Reader et al. (2011, p. 1018) define social learning as the tendency to "learn

skills and acquire information from others", including instances of social learning from both kin

and unrelated individuals. Social learning was often inferred from observational data in the

original reports.

As an alternative to treating behavioural data as a binary metric (e.g. presence or absence of

innovativeness in Ducatez, et al., 2020) some studies have used "rate" measures of behaviours:

residuals from a log-log plot of the total number of recorded instances of a behaviour (e.g. social

learning) and an estimate of research effort (e.g. the number of papers published on that taxa; e.g.

Sol, et al., 2005; Reader, et al., 2011; Navarrete, et al., 2016; Ducatez, et al., 2020). However, the

choice of how to model the relationship between total number of recorded innovation or social

learning instances and research effort matters considerably when creating these residual rate

measures. For our data, the relationship between the total number of innovation or social learning

instances and research effort was non-linear. This made the choice of model structure non-trivial,

with different models proving difficult to justify over one another. Residuals from these models

additionally showed further structural issues, including failure to meet assumptions about

homoscedasticity. Thus, we opted to use binary measures, which allowed us to use data

imputation methods to minimize biases caused by under-studied taxa, statistically account for

potential biases associated with summarizing behavioural data at higher taxonomic levels, and

run simulations to address the assumptions underlying our analyses. We therefore assigned each

lineage or genus binary scores of 1 (presence of innovation or social learning) or 0 (absence of

innovation or social learning). Further considerations regarding the use of literature-based

estimates of behavioural flexibility can be found in the supplementary material.

Structural Proxies of Behavioural Flexibility

Literature-based evidence for behavioural flexibility across taxa has its limitations and so we

chose to also consider structural correlates of behavioural flexibility. It is widely thought that

particular brain regions are associated with flexible behaviour – particularly the neocortex (see,

e.g., Keverne, et al., 1996; Mikhalevich, et al., 2017) and cerebellum (Vandervert, 2003;

Vandervert, et al., 2007; Barton, 2012). Therefore, in addition to total brain size (relative to body

mass), we considered the sum of neocortical and cerebellar volumes relative to rest of total brain

volume as another proxy for behavioural flexibility. Using both behavioural measures and

structural correlates of behavioural flexibility, we were able to compare the consistency of results

across different proxies for behavioural flexibility.

Lineage-level estimates for all brain measures were calculated by taking the geometric mean of

taxon volumes for each lineage. Brain volume relative to body size (hereafter 'relative brain

volume') was estimated by regressing lineage-level estimates of logarithmic endocranial volume,

in cm^3, (ECV) (Powell, et al., 2017) (hereafter 'brain volume') as a function of logarithmic body

mass, in grams (Jones, et al., 2009), and retaining the residuals (Dunbar & Schultz, 2007).

Notably, previous studies (e.g. Sayol, et al., 2019) have used phylogenetically corrected residuals

to account for an effect of body size when using similar brain size measures. However, for our

data, residuals obtained from phylogenetic models were skewed such that lineages with larger

body mass consistently had larger residual brain volumes (see Figure S2), and thus we opted to

use ordinary least square residuals that were more orthogonal to lineage body size. Consistent

with relative brain volume, neocortex and cerebellum volume relative to rest-of-brain volume

(hereafter 'relative neocortex and cerebellum volume') was estimated by taking the residuals

from a log-log regression of the combined neocortex and cerebellum volumes on the rest-of-

brain volumes (i.e. total brain volume minus neocortex and cerebellum volumes) taken from

Navarrete et al. (2018) and the compilation in DeCasien & Higham (2019). Genus-level

estimates for structural proxies of flexibility were calculated by taking the mean of the lineage

estimates within each genus. Further details, and considerations regarding the use of structural

proxies of behavioural flexibility, and residual brain measures can be found in the supplementary

materials along with correlation coefficients for all predictor variables (Table S1).

Phylogenetic signal of all predictor and response variables are reported in the supplementary

material (Table S2).

Analysis

All analyses used R version 3.6.3 (R Core Team, 2020).

Trait Imputation

While we collected data for our predictor variables from the most comprehensive datasets and

compilations available, research biases and the persistent reassessment of primate taxonomy has

resulted in inconsistent data coverage across lineages, and there were still many lineages that

were missing data (see Figure S3). To maximize our evolutionary inferences on diversification

and allow for the inclusion of data-poor lineages, we chose to impute missing predictor variables

using phylogenetic imputation methods (see supplementary material for details and reports of

predictive accuracy from model cross-validation; Table S3 and Figure S4). Data on relative

neocortex and cerebellum volume were sparse and unevenly distributed across the phylogeny

(82.4% of lineages missing data; Table S3), making it infeasible to reliably impute missing

values. We thus only ran models of relative neocortex and cerebellum volume on the original,

non-imputed dataset. All of the regressions we report below were repeated for the original, non-

imputed datasets (see Results and the supplementary material Tables S4 to S6) and except as

noted gave similar results.

Predictors of Diversification

To assess the relationship between our measures of behavioural flexibility and diversification

rate at the lineage-level (Taxa per Lineage Diversification Rate) we used phylogenetic

generalized least squares (PGLS) regressions implemented using the "gls" function in the nlme

package (Pinheiro, et al., 2020), including the 10kTrees consensus tree for our 239 defined

lineages as the phylogenetic backbone. PGLS is a common regression method used to investigate

evolutionary associations while accounting for the fact that closely-related lineages tend to be

similar (e.g. in body size, life history and ecology; see Freckleton, et al., 2002). Continuous data

were log-transformed, centred with respect to the mean, and scaled by 2 standard deviations in

all models for both the lineage and genus-level analyses to make effect sizes comparable to those

reported for binary variables (Gelman, 2008). After imputing missing values, our dataset

contained 54 lineages scored as having evidence of social learning (scored as 1) and 28 lineages

scored as having evidence of technical innovation (scored as 1). Wyles et al. (1983) predicted

accelerated evolution in species with a dual capacity for innovation and social propagation of

new behaviours, therefore, we also tested a combined measure of technical innovation and social

learning. In this combined measure 26 lineages with the presence of both behaviours were scored

as 1, and those exhibiting only one or neither behaviour were scored as 0.

To assess the relationship between our measures of behavioural flexibility and diversification

rate deeper in the tree we repeated the same analysis at the genus-level using two different

estimates of diversification rate: Taxa per Genus Diversification Rate and the more conservative

Lineage per Genus Diversification Rate. After imputing missing values, our dataset of 60 genera

contained 21 genera scored as having evidence of social learning, 9 genera scored as having

evidence of technical innovation and 8 genera with evidence of both behaviours.

Body mass and attendant life history traits have been predicted to impact diversification rates in

some taxa, albeit with conflicting results (see, e.g., Cardillo, et al., 2003; Paradis, 2005;

Fontanillas, et al., 2007; Thomas, et al., 2010), and body mass is closely correlated with many

primate life history traits (e.g. age at first reproduction, maximum lifespan; Charnov & Berrigan,

1993; Purvis, et al., 2003; Street, et al., 2017). To examine whether our results could stem from

confounding effects of body mass and its correlates, we ran PGLS analyses to test body mass as

an independent predictor of our diversification rate measures. Results from these tests were non-

significant across all measures of diversification (see supplementary material Tables S4 to S6).

We re-ran all PGLS analyses reported above after eliminating the great apes, which are

characterized by slow life histories, which may counteract the evolutionary effects of

behavioural flexibility by slowing their evolutionary response time (Price, et al., 2003). For tests

that proved significant, we then reran the same test 100 times over a sample of 1000 trees from

the tree block available in 10kTrees (Arnold, et al., 2010) to confirm results were consistent after

accounting for phylogenetic uncertainty.

Genus-Level Simulations

Genera were considered behaviourally flexible if any of their daughter lineages had evidence of

innovation or social learning, which potentially introduces a statistical bias if more lineage-rich

genera (which will generally have higher diversification rates) are more likely by chance to

include at least one lineage that expresses technical innovation or social learning. To account for

this possible bias, we simulated the neutral evolution of technical innovation and social learning

across the primate phylogeny 1000 times using the symmetrical rate Mk model of discrete trait

evolution (Lewis, 2001). We opted to model the evolution of these behaviours under a

symmetrical model of trait evolution since with smaller datasets such as ours, there is little power

to prefer asymmetric models (Mooers & Schluter, 1999). From these stochastic distributions of

the two traits, we repeated our analyses of diversification rate and created a distribution of

expected effect sizes under a null evolutionary scenario. More diverse genera may also be more

likely, by chance, to have well-studied lineages, which could create a bias toward observing the

presence of innovation or social learning in diverse genera (Ducatez & Lefebvre, 2014).

Although data did not suggest that lineage-rich clades were more likely to have intensely

investigated lineages in our dataset (see Figures S5 and S6), we nonetheless opted to take

research effort into account in our analysis and expanded upon our first simulation to consider a

scenario where the presence of technical innovation or social learning may go unobserved if

insufficient research effort was directed at a lineage. We took the 1000 simulations of neutral

evolution of technical innovation and social learning, and then randomly 'evolved' research

effort for each simulation, independently of the evolution of technical innovation and social

learning. We then converted the presence of these behaviours to absences in our simulated

datasets if a lineage's 'evolved' research effort was below the minimum threshold of studies for

a lineage with observed technical innovation (as it contained the higher research effort threshold

than social learning; Figure S7). By repeating our analyses of diversification rate with these new

simulated datasets where lineages with low research effort were assigned 'hidden states', we

were able to see if a bias towards having better studied lineages in diverse genera could drive a

positive association between behavioural proxies of flexibility and diversification rate by chance

and independent of biological mechanisms. Importantly, this simulation makes the assumption

that research effort and behavioural flexibility are independent; if this is not true such that

behaviourally flexible lineages attract research effort (see discussion in the 'Research Effort

Bias' section of the supplementary material) then the results of this simulation would actually be

conservative. We detail both simulations, along with considerations about using research effort

as a covariate in binary models (Ducatez, et al., 2020), in the supplementary material.

Ethical Note

This research was comparative and was based on data available in previously published

literature.

RESULTS:

Lineage-Level Predictors of Diversification

We found no support for an association between behavioural flexibility and diversification rates

when testing our measures of behavioural flexibility at the lineage-level (results summarized in

Figures 1 and S8; Table S4). Social learning (p= 0.171), technical innovation (p= 0.792), the

combined presence of technical innovation and social learning (p= 0.979), relative brain volume

(p= 0.215), relative neocortex and cerebellum volume (p= 0.664), and body mass (p= 0.764;

Table S4) did not exhibit noteworthy associations with Taxa per Lineage Diversification Rate in

either direction.

Genus-Level Predictors of Diversification

Technical innovation (p= 0.577), social learning (p= 0.442), the combined presence of technical

innovation and social learning (p= 0.411), relative neocortex and cerebellum volume (p= 0.930),

and body mass (p= 0.204) did not exhibit noteworthy associations with Taxa per Genus

Diversification Rate (Figures 1 and S8; Table S5). Relative brain volume was insignificantly (p=

0.108), but positively, related to diversification rate, and this relationship became significant

when we considered the non-imputed dataset (p= 0.004).

In contrast to the other diversification measures, Lineage per Genus Diversification Rate was

positively associated with all three behavioural measures of behavioural flexibility (results

summarized in Figures 1 and S8; Table S6). Genera with records of technical innovation were

shown to have a faster mean Lineage per Genus Diversification Rate (0.136 lineages my^{-1}) than

those with no reports (0.070 lineages my^{-1}; ß [95% CI] =0.066 [0.002-0.130]; p= 0.048). Genera

with records of social learning similarly exhibited a faster mean Lineage per Genus

Diversification Rate (0.137 lineages my^{-1}) than those without (0.049 lineages my^{-1}; ß [95% CI]

=0.088 [0.043-0.132]; p< 0.001). Genera with reports of both technical innovation and social

learning also exhibited a faster mean Lineage per Genus Diversification Rate (0.153 lineages my^{-}

^{1}) compared to those with evidence for only one or neither behaviour (0.068 lineages my^{-1}; ß

[95% CI] =0.085 [0.018-0.151]; p= 0.015). These results were consistent with results run over a

sample of 1000 trees from the posterior distribution, i.e. "tree block", available from 10kTrees

(Arnold, et al., 2010), indicating results were not biased by our choice of phylogeny (Figure S9).

Based on our simulation testing for a lineage-richness sampling bias, we found that these effect

sizes were unlikely to be due to chance alone for technical innovation (p= 0.028), social learning

(p= 0.002) and the combined presence of both behaviours (p= 0.008) (see Figure S10 and

supplementary results). After expanding our simulation to consider a research effort bias,

significance of these effects remained for social learning (p= 0.019) and its combined presence

with technical innovation (p= 0.027), but the effect of technical innovation alone was no longer

nominally significant (p= 0.109) (see Figure S11 and supplementary results). No structural

proxies of behavioural flexibility shared notable associations with Lineage per Genus

Diversification Rate (Figures 1 and S8; Table S6).

Results of all PGLS analyses reported above were qualitatively consistent after removing great

apes (see supplementary material Tables S7 to S9).

Figure 1: Results from PGLS analyses showing the association of proxies of behavioural

flexibility with Taxa per Lineage Diversification Rate (DR), Taxa per Genus Diversification

Rate and Lineage per Genus Diversification Rate across primates. 95% confidence intervals are

represented by horizontal lines around the associated value. Diamonds indicate diversification

rates estimated with species/subspecies richness, and squares indicate diversification rates

estimated with lineage richness. Results presented include imputed data when available and brain

measures were standardized (ln x/(2sd)). Significance indicated as: +P≤0.1; *P≤0.05; **P≤0.01.

DISCUSSION:

We find little to no compelling support for an association between our proxy measures of

behavioural flexibility and recent primate diversification rates of young species and subspecies,

however, we do find some evidence supporting a positive association when looking at the

diversification of older primate lineages. This pattern could be explained in several ways, and

interpretation of these results depend on the assumptions one makes about our different measures

of diversification rate.

One benefit of our study design is that it allowed us to consider taxa that are commonly

overlooked (i.e. subspecies and species omitted from higher order phylogenies), many of which

likely represent very recent splitting events. The weak associations with behavioural flexibility

that we observed at shallower time scales could be explained in several ways. On one hand, this

pattern could reflect biases in describing taxa (species and subspecies) among groups. While our

time cut-off, and use of both species and subspecies, mitigates against biases associated with the

elevation of subspecies under the PSC, if less flexible species are more likely to have a larger

number of taxa described overall (e.g. based on regional biases in designating species or

subspecies that covary with locations of inflexible clades) then this could obscure underlying

biological patterns. If this is the case, then behavioural flexibility may indeed enhance

diversification by acting on speciation, and we observe stronger associations when ignoring

shallow splits because our diversification rate metric calculated using lineage-richness is less

biased by this phenomenon. On the other hand, if we take our results at face value, a pattern of

stronger associations deeper in the phylogeny could indicate that time plays an even larger role

in the relationship between behavioural flexibility and diversification than previously suggested

by Duckworth (2009).

Under Mayr's 'ephemeral speciation model' (Mayr, 1963; Rosenblum, et al., 2012) and the

related 'ephemeral divergence hypothesis' (Futuyma, 1979; 2010), divergence can occur rapidly

and often, but many newly diversifying lineages do not persist, instead being eradicated via

extinction or 'reabsorption' by hybridization (see, e.g., Rosenblum, et al., 2012). It is possible

that a number of the described species and subspecies used here (from the IUCN/SSC Primate

Specialist Group, 2018) represent such ephemeral diversification events, especially considering

that the PSC has facilitated the splitting of very closely related populations. If so, our results

could be explained if behavioural flexibility buffers against extinction through behavioural shifts.

Behavioural flexibility would then promote diversification through bolstering lineage persistence

rather than the rate of splitting; and would be revealed when comparing the accumulation of

lineages that have escaped extinction (i.e. looking deeper in the tree). This would be

complementary to findings from Arbour & Santana (2017), who show that decreased extinction

preceded a shift to increased evolutionary rates in the most speciose primate family

(Cercopithecidae), and with evidence suggesting behavioural flexibility is beneficial for

population persistence in birds (e.g. Shultz, et al., 2005; Rossmanith, et al., 2006; Sol, et al.,

2007, Ducatez, et al., 2020). Under this scenario, the positive associations we find are not the

result of behavioural flexibility enhancing evolutionary rates by facilitating divergence events,

and instead indicate that extinction resistance may be an important mechanism linking

behavioural flexibility and primate diversification. Our results stand in contrast with consistent

reports of behavioural flexibility enhancing diversification of birds even at shallow phylogenetic

depths (Nicolakakis, et al., 2003; Sol, et al., 2005; Sayol, et al., 2019).

A potential explanation for differing effects of behavioural flexibility in birds versus primates

could be that the heightened dispersal capabilities (i.e. flight) of birds makes behavioural

flexibility particularly beneficial for these taxa when it comes to establishing in new

environments. This could cause behavioural flexibility to enhance bird speciation via increased

success in dispersal events while also buffering against extinction – leading to the strong effects

that have been observed on their net rates of evolution (Nicolakakis, et al., 2003; Sol, 2003; Sol,

et al., 2005; Sayol, et al., 2019). Comparatively, primates have much more limited dispersal

capabilities and thus behavioural flexibility may buffer against extinction while having little

effect on promoting allopatric establishment and subsequent speciation. If true, it is possible that

behavioural flexibility affects evolutionary trajectories via different mechanisms in these clades.

Directly testing whether behavioural flexibility buffers against extinction may be difficult

because measuring extinction rates at macroevolutionary scales is notoriously imprecise

(Rabosky, 2010; Louca & Pennell, 2020). Future studies could test proxies of primate

behavioural flexibility against estimates of contemporary extinction vulnerability (see, e.g.,

Nicolakakis, et al., 2003; Ducatez, et al., 2020). Another prediction concerns the rate of genetic

divergence within flexible versus less flexible, recent clades (e.g. subspecies within species): our

conjecture would be that these would not be different, because flexibility does not lead to

increased divergence over the short term.

Notably, not all of our results are consistent with a significantly positive association between

behavioural flexibility and diversification of older primate lineages. While the non-significant

association between Lineage per Genus Diversification Rate and neocortex and cerebellum

volume might be explained by data limitations (discussed below), the non-significant association

with relative brain volume illustrates that not all proxies of behavioural flexibility capture the

same thing. While relative brain volume is a known structural correlate of measures of primate

and avian behavioural flexibility (Lefebvre, et al., 2004), the brain has many functions and thus

is a less direct measure of behavioural flexibility than behavioural measures. Measures of brain

size have been linked to diversification rate in birds (Nicolakakis, et al., 2003; Sol, et al., 2005;

Sayol, et al., 2019), but notably the sample for birds is much larger, and is likely to contain a

broader range of variation, which would make it easier to detect a brain size effect. Additionally,

results of our simulation accounting for potential research effort biases indicate that a positive

association between technical innovation and diversification rate in primate genera could be

driven by a research effort bias. However, this simulation assumes that research effort and

behavioural flexibility are independent and, if instead research effort tends to be directed toward

groups where innovation is expected, then this research bias may actually have a low

contribution to the observed relationship between technical innovation and diversification rate.

Thus, our results paint a complicated picture, and emphasize the need to compare methodologies

and measures, and avoid a focus on single explanations for complex phenomena.

Our results provide some support for primate behavioural flexibility being associated with older

as opposed to more recent diversification events, however, there are several caveats on the limits

of our inferences. First, one measure thought to be associated with behavioural flexibility –

neocortex and cerebellum volume – had very little data available for lineages reported as

inflexible (e.g. among lemurs, tarsiers, titis and sakis). This meant our ability to test this measure

as a driver of diversification was limited. Additionally, to examine the relationship between

different measures of behavioural flexibility and diversification rate, we had to run multiple tests,

increasing the likelihood of observing Type I error. Our study is also limited by the lineages

present in the phylogeny. Available primate phylogenies, (e.g. Arnold, et al., 2010; Upham, et

al., 2019b) do not reflect the most recent taxon lists, and such limited taxonomic scope prevented

us from using diversification rate estimates that require a fully-resolved tree (e.g. the DR

measures from Jetz, et al., 2012). Lastly, many factors will contribute to true diversification, and

although we test two (e.g. body mass and a "great ape effect"; see supplemental results), there are

many biological and ecological factors (e.g. geographic discontinuities, habitat affiliations, or

presence on islands) yet to be explored that could mediate any relationship between behavioural

flexibility and primate diversification rate.

While our results provide some evidence for the hypothesis that behavioural flexibility drives

diversification of primate lineages, they raise important questions about its underlying

mechanisms. Importantly, our results are not consistent in supporting a significant association

between behavioural flexibility and diversification rate, and depend on the proxy measures

employed. The positive associations we do find may point to behavioural flexibility dampening

extinction of young primate lineages, rather than accelerating diversification via splitting of

behaviourally shifted individuals/populations. Thus, expanding tests of whether and how

behavioural flexibility is associated with diversification to other taxa may help us interpret the

results we report here.

ACKNOWLEDGEMENTS: This work was supported by the Natural Sciences and Engineering

Research Council of Canada (NSERC) Discovery Grants Program (SMR and AOM, grants

#2017-04720 and #2019-04950) and the Canada Foundation for Innovation (SMR, grant

#29433). MJAC was supported by awards from the Biodiversity, Ecosystem Services, and

Sustainability (BESS) program and McGill University. We thank the Reader and Guigueno labs,

Brian Leung, and Hans Larsson at McGill, William Wcislo, and the Crawford Lab for

Evolutionary Studies at SFU for feedback on study design and interpretation, as well as Isabella

Capellini and three anonymous reviewers for comments on previous versions.

Data and code will be made available through Dryad or a similar repository.

REFERENCES:

Anderson, R. W. (1995). Learning and evolution: A quantitative genetics approach. Journal of

Theoretical Biology, 175: 89–101.

Ancel, L. W. (1999). A quantitative model of the Simpson–Baldwin effect. Journal of

Theoretical Biology, 196: 197–209.

Ancel, L. W. (2000). Undermining the Baldwin expediting effect: does phenotypic plasticity

accelerate evolution? Theoretical Population Biology, 58: 307–319.

Arbour, J. H., & Santana, S. E. (2017). A major shift in diversification rate helps explain

macroevolutionary patterns in primate species diversity. Evolution, 71: 1600–1613.

Arnold, C., Matthews, L. J., & Nunn, C. L. (2010). The 10kTrees website: a new online resource

for primate phylogeny. Evolutionary Anthropology, 19: 114–118.

Audet, J. N., & Lefebvre, L. (2017). What's flexible in behavioural flexibility? Behavioural

Ecology, 28: 943–947.

Badgley, C. (2010). Tectonics, topography, and mammalian diversity. Ecography, 33: 220–231.

Baldwin, J. M. (1902). Development and evolution. MacMillan, New York.

Barton, R. A. (2012). Embodied cognitive evolution and the cerebellum. Philosophical

Transactions of the Royal Society B: Biological Sciences, 367: 2097–2107.

Bateson, P. (1988). The active role of behaviour in evolution. In M. −W. Ho & S. W. Fox (Eds.),

Evolutionary processes and metaphors (p. 191–207). John Wiley and Sons, Ltd., New York.

Behera, N., & Nanjundiah, V. (1995). An investigation into the role of phenotypic plasticity in

evolution. Journal of Theoretical Biology, 172: 225–234.

Bogert, C. M. (1949). Thermoregulation in reptiles, a factor in evolution. Evolution, 3: 195–211.

Cardillo, M., Huxtable, J. S., & Bromham, L. (2003). Geographic range size, life history and

rates of diversification in Australian mammals. Journal of Evolutionary Biology, 16: 282–

288.

Charnov, E. L., & Berrigan, D. (1993). Why do female primates have such long lifespans and so

few babies? Or life in the slow lane. Evolutionary Anthropology, 1: 191–194.

DeCasien, A. R., & Higham, J. P. (2019). Primate mosaic brain evolution reflects selection on

sensory and cognitive specialization. Nature Ecology & Evolution, 3: 1483–1493.

Ducatez, S., Clavel, J., & Lefebvre, L. (2015). Ecological generalism and behavioural

innovation in birds: technical intelligence or the simple incorporation of new foods?

Journal of Animal Ecology, 84: 79–89.

Ducatez, S., & Lefebvre, L. (2014). Patterns of research effort in birds. PLoS One, 9: e89955.

Ducatez, S., Sol, D., Sayol, F., & Lefebvre, L. (2020). Behavioural plasticity is associated

with reduced extinction risk in birds. Nature Ecology & Evolution, 4: 788–793.

Duckworth, R. A. (2009). The role of behaviour in evolution: a search for mechanism.

Evolutionary Ecology, 23: 513–531.

Dunbar, R. I., & Shultz, S. (2007). Understanding primate brain evolution. Philosophical

Transactions of the Royal Society B: Biological Sciences, 362: 649–658.

Fontanillas, E., Welch, J. J., Thomas, J. A., & Bromham, L. (2007). The influence of body size

and net diversification rate on molecular evolution during the radiation of animal phyla.

BMC Evolutionary Biology, 7: 95.

Foote, A. D., Vijay, N., Ávila-Arcos, M. C., Baird, R. W., Durban, J. W., Fumagalli, M., … &

Wolf, J. B. W. (2016). Genome-culture coevolution promotes rapid divergence of killer

whale ecotypes. Nature Communications, 7: 11693.

Freckleton, R. P., Harvey, P. H., & Pagel, M. (2002). Phylogenetic analysis and comparative

data: a test and review of evidence. The American Naturalist, 160: 712–726.

Frey, M. A. (2010). The relative importance of geography and ecology in species diversification:

evidence from a tropical marine intertidal snail (*Nerita*). Journal of Biogeography, 37: 1515–

1528.

Futuyma, D. J. (1979). Evolutionary biology. Sinauer Associates, Sunderland.

Futuyma, D. J. (2010). Evolutionary constraint and ecological consequences. Evolution, 64:

1865–1884.

Gelman, A. (2008). Scaling regression inputs by dividing by two standard deviations. Statistics

in Medicine, 27: 2865–2873.

Gonzalez-Voyer, A., González-Suárez, M., Vilà, C., & Revilla, E. (2016). Larger brain size

indirectly increases vulnerability to extinction in mammals. Evolution, 70: 1364–1375.

Grant, P. R., & Grant, B. R. (2008). How and why species multiply: the radiation of Darwin's

finches. Princeton University Press, Princeton.

Groves, C. P. (2014). Primate taxonomy: inflation or real? Annual Review of Anthropology, 43:

27–36.

Heard, S. B., & Hauser, D. L. (1995). Key evolutionary innovations and their ecological

mechanisms. Historical Biology, 10: 151–173.

Hinton, G. E., & Nowlan, S. J. (1987). How learning can guide evolution. Complex Systems, 1:

495–502.

Honacki, J. H., Kinman, K. E., & Koeppl, J. W. (1982). Mammal species of the world: a

taxonomic and geographic reference. Allen Press, Inc. and Association of Systematic

Collections, Lawrence.

Huey, R. B., Hertz, P. E., & Sinervo, B. (2003). Behavioural drive versus behavioural inertia in

evolution: a null model approach. The American Naturalist, 161: 357–366.

Isaac, N. J., Jones, K. E., Gittleman, J. L., & Purvis, A. (2005). Correlates of species richness in

mammals: body size, life history, and ecology. The American Naturalist, 165: 600–607.

Isaac, N. J., Mallet, J., & Mace, G. M. (2004). Taxonomic inflation: its influence on

macroecology and conservation. Trends in Ecology & Evolution, 19: 464–469.

IUCN/SSC Primate Specialist Group. (2018). Retrieved from http://www.primate-

sg.org/red_list_threat_status/ [accessed 10 November 2018].

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global diversity

of birds in space and time. Nature, 491: 444–448.

Jones, K. E., Bielby, J., Cardillo, M., Fritz, S. A., O'Dell, J., Orme, C. D. L., ... & Connolly, C.

(2009). PanTHERIA: a species-level database of life history, ecology, and geography of

extant and recently extinct mammals: Ecology, 90: 2648–2648.

Louca, S., & Pennell, M. W. (2020). Extant timetrees are consistent with a myriad of

diversification histories. Nature, 580: 502–505.

Keverne, E. B., Martel, F. L., & Nevison, C. M. (1996). Primate brain evolution: genetic and

functional considerations. Proceedings of the Royal Society B: Biological Sciences, 263:

689–696.

Kozak, K. H., & Wiens, J. J. (2010). Accelerated rates of climatic-niche evolution underlie rapid

species diversification. Ecology Letters, 13: 1378–1389.

Lachlan, R. F., & Servedio, M. R. (2004). Song learning accelerates allopatric speciation.

Evolution, 58: 2049–2063.

Lapiedra, O., Sol, D., Carranza, S., & Beaulieu, J. M. (2013). Behavioural changes and the

adaptive diversification of pigeons and doves. Proceedings of the Royal Society B:

Biological Sciences, 280: 20122893.

Lefebvre, L., Reader, S. M., & Sol, D. (2004). Brains, innovations and evolution in birds and

primates. Brain, Behaviour and Evolution, 63: 233–246.

Lewis, P. O. (2001). A likelihood approach to estimating phylogeny from discrete morphological

character data. Systematic Biology, 50: 913–925.

Lewontin, R. C. (1983). Gene, organism and environment. In D. S. Bendall (Eds.), Evolution

from molecules to men (p. 273–285). Cambridge University Press, New York.

Losos, J. B., Schoener, T. W., & Spiller, D. A. (2004). Predator-induced behaviour shifts and

natural selection in field-experimental lizard populations. Nature, 432: 505–508.

Magallon, S., & Sanderson, M. J. (2001). Absolute diversification rates in angiosperm clades.

Evolution, 55: 1762–1780.

Mayr, E. (1963). Animal species and evolution. Belknap Press, Cambridge.

Mikhalevich, I., Powell, R., & Logan, C. (2017). Is behavioural flexibility evidence of

cognitive complexity? How evolution can inform comparative cognition. Interface

Focus, 7: 20160121.

Mitchell, J. S., & Rabosky, D. L. (2017). Bayesian model selection with BAMM: effects of

the model prior on the inferred number of diversification shifts. Methods in Ecology

and Evolution, 8: 37–46.

Mooers, A. Ø., & Schluter, D. (1999). Reconstructing ancestor states with maximum likelihood:

support for one-and two-rate models. Systematic Biology, 48: 623–633.

Muñoz, M. M., & Losos, J. B. (2018). Thermoregulatory behaviour simultaneously promotes

and forestalls evolution in a tropical lizard. The American Naturalist, 191: E15–E26.

Navarrete, A. F., Blezer, E. L., Pagnotta, M., de Viet, E. S., Todorov, O. S., Lindenfors, P., ...

& Reader, S. M. (2018). Primate brain anatomy: new volumetric MRI measurements

for neuroanatomical studies (with Erratum). Brain, Behaviour and Evolution, 91: 109–

117.

Navarrete, A. F., Reader, S. M., Street, S. E., Whalen, A., & Laland, K. N. (2016). The

coevolution of innovation and technical intelligence in primates. Philosophical

Transactions of the Royal Society B: Biological Sciences, 371: 20150186.

Nicolakakis, N., Lefebvre, L. & Sol, D. (2003). Behavioural flexibility predicts species

richness in birds, but not extinction risk. Animal Behaviour, 65: 445–452.

Odling-Smee, F. J., Laland, K. N., & Feldman, M. W. (2003). Niche construction: the

neglected process in evolution. Princeton University Press, Princeton.

Overington, S. E., Morand-Ferron, J., Boogert, N. J., & Lefebvre, L. (2009). Technical

innovations drive the relationship between innovativeness and residual brain size in

birds. Animal Behaviour, 78: 1001–1010.

Paradis, E. (2005). Statistical analysis of diversification with species traits. Evolution, 59: 1–

12.

Pelletier, F., Garant, D., & Hendry, A. P. (2009). Eco-evolutionary dynamics. Philosophical

Transactions of the Royal Society B: Biological Sciences, 364: 1483–1489.

Pfennig, D. W., Wund, M. A., Snell-Rood, E. C., Cruickshank, T., Schlichting, C. D., &

Moczek, A. P. (2010). Phenotypic plasticity's impacts on diversification and

speciation. Trends in Ecology & Evolution, 25: 459–467.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Core Team. (2020). Linear and nonlinear

mixed effects models. R package version 3.1-144. https://CRAN.R-

project.org/package=nlme.

Powell, L. E., Isler, K., & Barton, R. A. (2017). Re-evaluating the link between brain size and

behavioural ecology in primates. Proceedings of the Royal Society B: Biological

Sciences, 284: 20171765.

Price, T. (2008). Speciation in birds. Roberts and Company, Greenwood Village.

Price, T. D., Qvarnström, A., & Irwin, D. E. (2003). The role of phenotypic plasticity in

driving genetic evolution. Proceedings of the Royal Society B: Biological Sciences,

270: 1433–1440.

Purvis, A., Webster, A. J., Agapow, P. M., Jones, K. E., & Isaac, N. J. (2003). Primate life

histories and phylogeny. In P. M. Kappeler & M. E. Pereira (Eds.), Primate life

histories and socioecology (p. 25–40). University of Chicago Press, Chicago.

R Core Team. (2020). R: A Language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/

Rabosky, D. L. (2009). Ecological limits and diversification rate: alternative paradigms to

explain the variation in species richness among clades and regions. Ecology Letters,

12: 735–743.

Rabosky, D. L. (2010). Extinction rates should not be estimated from molecular phylogenies.

Evolution, 64: 1816–1824.

Raikow, R. J. (1986). Why are there so many kinds of passerine birds? Systematic Zoology,

35: 255–259.

Reader, S. M., & Biro, D. (2010). Experimental identification of social learning in wild

animals. Learning & Behaviour, 38: 265–283.

Reader, S. M., Hager, Y., & Laland, K. N. (2011). The evolution of primate general and

cultural intelligence. Philosophical Transactions of the Royal Society B, 366: 1017–

1027.

Reader, S. M., & Laland, K. N. (2002). Social intelligence, innovation, and enhanced brain

size in primates. Proceedings of the National Academy of Sciences, 99: 4436–4441.

Riesch, R., Barrett-Lennard, L. G., Ellis, G. M., Ford, J. K. B., & Deecke, V. B. (2012). Cultural

traditions and the evolution of reproductive isolation: ecological speciation in killer whales?

Biological Journal of the Linnean Society, 106: 1–17.

Rosenblum, E. B., Sarver, B. A., Brown, J. W., Des Roches, S., Hardwick, K. M., Hether, T. D.,

... & Harmon, L. J. (2012). Goldilocks meets Santa Rosalia: an ephemeral speciation model

explains patterns of diversification across time scales. Evolutionary Biology, 39: 255–261.

Rossmanith, E., Grimm, V., Blaum, N., & Jeltsch, F. (2006). Behavioural flexibility in the

mating system buffers population extinction: lessons from the lesser spotted woodpecker

Picoides minor. Journal of Animal Ecology, 75: 540–548.

Rylands, A. B., & Mittermeier, R. A. (2014). Primate taxonomy: species and conservation.

Evolutionary Anthropology, 23: 8–10.

Sayol, F., Lapiedra, O., Ducatez, S., & Sol, D. (2019). Larger brains spur species diversification

in birds. Evolution, 73: 2085–2093.

Scott-Phillips, T. C., Laland, K. N., Shuker, D. M., Dickins, T. E., & West, S. A. (2014). The

niche construction perspective. A critical appraisal. Evolution, 68: 1231–1243.

Shultz, S. B., Bradbury, R. L., Evans, K. D., Gregory, R., & Blackburn, T. M. (2005). Brain size

and resource specialization predict long-term population trends in British birds. Proceedings

of the Royal Society B: Biological Sciences, 272: 2305–2311.

Snell-Rood, E. (2013). An overview of the evolutionary causes and consequences of behavioural

plasticity. Animal Behaviour, 85: 1004–1011.

Snell-Rood, E., Swanson, E., & Jaumann, S. (2015). Insects as a model system to understand the

evolutionary implications of innovation. In A. B. Kaufman & J. C. Kaufman (Eds.), Animal

creativity and innovation (p. 459–478). Academic Press, London.

Sol, D. (2003). Behavioural flexibility: a neglected issue in the ecological and evolutionary

literature? In S. M. Reader & K. N. Laland (Eds.), Animal innovation (p. 63–82). Oxford

University Press, Oxford.

Sol, D., & Lefebvre, L. (2000). Behavioural flexibility predicts invasion success in birds

introduced to New Zealand. Oikos, 90: 599–605.

Sol, D., & Price, T. D. (2008). Brain size and the diversification of body size in birds. The

American Naturalist, 172: 170–177.

Sol, D., Stirling, D. G., & Lefebvre, L. (2005). Behavioural drive or behavioural inhibition in

evolution: subspecific diversification in holarctic passerines. Evolution, 59: 2669–

2677.

Sol, D., Székely, T., Liker, A., & Lefebvre, L. (2007). Big-brained birds survive better in

nature. Proceedings of the Royal Society B: Biological Sciences, 274: 763–769.

Springer, M. S., Meredith, R. W., Gatesy, J., Emerling, C. A., Park, J., Rabosky, D. L., ... &

Fisher, C. A. (2012). Macroevolutionary dynamics and historical biogeography of

primate diversification inferred from a species supermatrix. PLoS One, 7: e49521.

Stadler, T. (2011). Mammalian phylogeny reveals recent diversification rate shifts.

Proceedings of the National Academy of Sciences, 108: 6187–6192.

Street, S. E., Navarrete, A. F., Reader, S. M., & Laland, K. N. (2017). Coevolution of cultural

intelligence, extended life history, sociality, and brain size in primates. Proceedings of

the National Academy of Sciences, 114: 7908-7914.

Tattersall, I. (2007). Madagascar's lemurs: cryptic diversity or taxonomic inflation?

Evolutionary Anthropology, 16: 12–23.

Tebbich, S., Sterelny, K., & Teschke, I. (2010). The tale of the finch: adaptive radiation and

behavioural flexibility. Philosophical Transactions of the Royal Society B: Biological

Sciences, 365: 1099–1109.

Thomas, J. A., Welch, J. J., Lanfear, R., & Bromham, L. (2010). A generation time effect on

the rate of molecular evolution in invertebrates. Molecular Biology and Evolution, 27:

1173–1180.

Upham, N., Esselstyn, J. A., & Jetz, W. (2019a). Ecological causes of uneven diversification

and richness in the mammal tree of life (Preprint). bioRxiv, 504803.

Upham, N. S., Esselstyn, J. A., & Jetz, W. (2019b). Inferring the mammal tree: Species-level

sets of phylogenies for questions in ecology, evolution, and conservation. PLoS

Biology, 17: 1–44.

Vandervert, L. (2003). The neurophysiological basis of innovation. In L. Shavinina (Eds.),

The international handbook on innovation (p. 17–30). Elsevier Science, Oxford.

Vandervert, L. R., Schimpf, P. H., & Liu, H. (2007). How working memory and the

cerebellum collaborate to produce creativity and innovation. Creativity Research

Journal, 19: 1–18.

Wcislo, W. T. (1989). Behavioural environments and evolutionary change. Annual Review of

Ecology and Systematics, 20: 137–169.

Weber, E. U., Shafir, S., & Blais, A.-R. (2004). Predicting risk sensitivity in humans and lower

animals: risk as variance or coefficient of variation. Psychological Review, 111: 430–445.

West-Eberhard, M. J. (1989). Phenotypic plasticity and the origins of diversity. Annual review of

Ecology and Systematics, 20: 249–278.

West-Eberhard, M. J. (2003). Developmental plasticity and evolution. Oxford University

Press, Oxford.

Wiens, J. J. (2017). What explains patterns of biodiversity across the Tree of Life? New

research is revealing the causes of the dramatic variation in species numbers across

branches of the Tree of Life. BioEssays, 39: 1600128.

Williams, C. B. (1964). Patterns in the balance of nature and related problems of quantitative

ecology. Academic Press, London & New York.

Willis, J. C. (1922). Area and age: A study of geographical distribution and origin in species.

Cambridge University Press, Cambridge.

Wilson, A. C. (1985). The molecular basis of evolution. Scientific American, 253: 164–173.

Wyles, J. S., Kunkel, J. G., & Wilson, A. C. (1983). Birds, behaviour, and anatomical

evolution. Proceedings of the National Academy of Sciences, 80: 4394–4397.

Yeh, P. J. (2004). Rapid evolution of a sexually selected trait following population

establishment in a novel habitat. Evolution, 58: 166–174.

Yeh, P. J., & Price, T. D. (2004). Adaptive phenotypic plasticity and the successful colonization

of a novel environment. The American Naturalist, 164: 531–542.