Respiratory tract explant infection dynamics of influenza A virus in California sea lions,

northern elephant seals, and rhesus macaques

Hongwei Liu[1] , Magdalena Plancarte[1] , Erin. E. Ball[1] , Christopher M. Weiss[1], Omar Gonzales-

Viera[1], Karen Holcomb[1], Zhong-Min Ma[2], A. Mark Allen[2], J. Rachel Reader[2], Pádraig J.

Duignan[3], Barbie Halaska[3], Zenab Khan[4], Divya Kriti[4], Jayeeta Dutta[4], Harm van Bakel[4],

Kenneth Jackson[1], Patricia A. Pesavento[1], Walter M. Boyce[1], Lark L. Coffey[1#]

[1]University of California, Davis, School of Veterinary Medicine, Department of Pathology, Microbiology

and Immunology

[2]University of California, Davis, California National Primate Research Center

[3]The Marine Mammal Center, 2000 Bunker Road, Sausalito, California

[4]Icahn School of Medicine at Mount Sinai, New York, New York

[#]Corresponding author

lcoffey@ucdavis.edu

ORCID numbers:

Lark L. Coffey : 0000-0002-0718-5146

Erin E. Ball: 0000-0002-8464-6612

Karen M. Holcomb: 0000-0002-9403-3756

Hongwei Liu: 0000-0002-8342-2545

Abstract

To understand susceptibility of wild California sea lions and Northern elephant seals to influenza A virus (IAV), we developed an *ex vivo* respiratory explant model and used it to compare infection kinetics for multiple IAV subtypes. We first established the approach using explants from colonized rhesus macaques, a model for human IAV. Trachea, bronchi, and lungs from 11 California sea lions, 2 Northern elephant seals and 10 rhesus macaques were inoculated within 24 hours post-mortem with 6 strains representing 4 IAV subtypes. Explants from the 3 species showed similar IAV infection kinetics with peak viral titers 48-72 hours post-inoculation that increased by 2-4 $\log_{10}$ plaque forming units (PFU)/explant relative to the inoculum. Immunohistochemistry localized IAV infection to apical epithelial cells. These results demonstrate that respiratory tissue explants from wild marine mammals support IAV infection. In the absence of the ability to perform experimental infections of marine mammals, this *ex vivo* culture of respiratory tissues mirrors the *in vivo* environment and serves as a tool to study IAV susceptibility, host-range, and tissue tropism.

Importance

Although influenza A virus can infect marine mammals, a dearth of marine mammal cell lines and ethical and logistical challenges prohibiting experimental infections of living marine mammals means that little is known about IAV infection kinetics in these species. We circumvented these limitations by adapting a respiratory tract explant model first to establish the approach with rhesus macaques and then for use with explants from wild marine mammals euthanized for non-respiratory medical conditions. We observed that multiple strains representing 4 IAV subtypes infected trachea, bronchi, and lungs of macaques and marine mammals with variable peak titers and kinetics. This *ex vivo* model can define infection dynamics for IAV in marine mammals. Further, use of explants from animals euthanized for other reasons reduces use of animals in research.

Introduction

Influenza A viruses (IAV) are important etiologies of respiratory disease in humans and especially affect the elderly, infants, and people with immunodeficiencies and chronic respiratory disease. Dwarfed in 2020 by SARS-CoV-2, IAV are a significant cause of morbidity annually, producing about 500,000 deaths worldwide each year (1). IAV possess a wide host range that includes birds, horses, pigs, and humans (2, 3). Marine mammals can also be

infected, sometimes with strains from human pandemics (4, 5). The viral genetic and host

factors that affect cross-species transmission by IAV, especially from birds to mammals

including humans, have been extensively studied (5–9). However, mechanisms of zoonotic IAV

transmission from avian or human to other mammalian species, including marine mammals,

are less well understood (6).

IAV was first identified in North American marine mammals in 1979, when an H7N7 epizootic

killed 500 harbor seals and caused hemorrhagic pneumonia in others (10–12). Since then, IAV

infection and sometimes disease has been identified in several marine mammal species,

including mass mortalities in harbor seals near Cape Cod, Massachusetts, USA, attributed to

H7N7, H4N5, or H4N6, and detection of antibody to multiple H and N subtypes in several seal

species (13–20). On the West Coast, surveillance from 2009 to 2015 (4, 21, 22), and from

2016 to present (unpublished) in multiple marine mammal species from California shows

variable IAV exposure. Seroprevalence in sea otters and Northern elephant seals is higher than

in sympatric harbor seals and California sea lions, and sea otters are exposed to avian- and

human-origin IAV, including pandemic H1N1 (4, 21). Despite frequent antibody detection,

isolation of IAV from marine mammals has been limited. In California, only pandemic H1N1 has

been isolated from 2 Northern elephant seals in 2010 (4). Together, these surveillance data

suggest that marine mammals can be infected with some of the same IAV subtypes that cause

human epidemics. However, in the absence of *in vivo* study capabilities in marine mammals,

infection dynamics and subtype-specific susceptibility remain unknown. Therefore, there is a

need to develop approaches to study the infection biology of IAV in these species.

Given that the respiratory tract is the initial site of IAV infection, defining infection dynamics and

cellular tropism in respiratory tissues is important for assessing species susceptibility. However,

in vivo systems for studying susceptibility of marine mammals to IAV are not available. As an

alternative, *ex vivo* explants from respiratory tract tissues can mimic the physiological

microenvironment of a respiratory tract. *Ex vivo* systems in human and animals (excluding

marine mammals) have been successfully developed and used to study host innate responses,

infection dynamics, viral genetic determinants of infection, antiviral drug treatments, and

pathogenesis of human and animal IAV and other respiratory viruses (23–28). The goal of this

study, therefore, was to expand these existing models for use in marine mammals to study

susceptibility and infection dynamics of IAV strains of mammalian and avian origin to assess

the potential for interspecies transmission, and to providing a new approach for studying the

biology and pathogenesis of IAV in marine mammals. Given that marine mammal tissues are

only opportunistically available from wild animals treated at The California Marine Mammal

Center in Sausalito, CA, USA, we first established the *ex vivo* explant system using rhesus

macaques that are more regularly available from the California National Primate Research

Center, Davis, CA, USA. Rhesus macaques represent a valuable model for understanding IAV

infection dynamics in the human respiratory tract due to similar structure, physiology, and

mucosal immunity. Additionally, IAV infection dynamics have not been studied in rhesus

macaques, although they are a model for human IAV infection (29) and are used for testing

vaccine candidates (30, 31).

We used trachea, bronchi, and lung explants from rhesus macaques to compare the

susceptibility, infection kinetics, and tissue tropism of 6 strains of IAV from 4 subtypes to

validate the utility of this system. After establishing the explant approach with IAV in rhesus

macaques, we used it to study IAV infection dynamics and tropism in California sea lions and

Northern elephant seals. We observed that both rhesus macaques and marine mammals are

susceptible to all 6 IAV strains, and that rhesus macaque and California sea lion respiratory

tract explants exhibit temporal, tissue, and IAV subtype-dependent IAV infection kinetics.

Material and Methods

Ethics Statement

This project was conducted with approval from the United States National Marine Fisheries Service

Marine Mammal permit # 18786-04. The University of California, Davis is accredited by the

Association for Assessment and Accreditation Laboratory Animal Care International (AAALAC).

Animal care was performed in compliance with the 2011 Guide for the Care and Use of Laboratory

Animals provided by the Institute for Laboratory Animal Research. Rhesus macaque studies were

approved by the University of California, Davis IACUC under protocol #19760.

Cell culture

Madin-Darby Canine Kidney (MDCK) cells (ATCC #CCL-34) were maintained at 37°C and 5%

CO_2 in MDCK-growth-medium (Iscove's Modified Dulbecco's Medium (IMDM), supplemented

with 5% FBS, 0.1% sodium bicarbonate, penicillin and streptomycin).

Influenza A virus propagation and titration

We used IAV strains of avian and marine mammal origin for explant studies (**Table 1**). The harbor

seal H3N8 (HS/H3N8) IAV strain was obtained from Dr. Hon Ip, (National Wildlife Health Center,

Madison Wisconsin). The other strains were isolated at UC Davis and used in this study. Avian

allantoic fluid virus stocks of 6 strains were amplified in Madin-Darby canine kidney (MDCK) cells to

obtain sufficient titers and volumes for use as inocula. Viral titers of the 6 inocula were determined by

plaque assay using MDCK cells. Briefly, MDCK cells were grown in 6-well plates to 80-90%

confluence in MDCK growth medium. Viral stocks were serially ten-fold diluted in viral-growth-medium

(Minimum Essential Medium (MEM), supplemented with 0.5% BSA, 0.1% sodium bicarbonate, 10

mM Hepes, penicillin and streptomycin). The MDCK cells were washed 3 times with Dulbecco's

phosphate buffered saline (DPBS) prior to inoculation with 200 ul of diluted virus samples. After 1

hour incubation at 37°C with 5% CO_2, 3 ml viral-growth-medium containing 1 ug/ml N-p-tosyl-L-

phenylalaninechloromethyl ketone-treated (TPCK) and 0.5% agarose was added to each well. After a

48 hour incubation at 37°C with 5% CO_2, MDCK cells were fixed with 4% formaldehyde and stained

with 0.05% crystal violet.

In vitro infection kinetics of IAV

Infection kinetics for the 6 IAV strains were assessed in MDCK cells. Cells were grown at 37°C

with 5% CO_2 to 6 x 10^5 cells/well in 24-well plates and inoculated in triplicate with 6 x 10^3 PFU

virus of IAV, representing a multiplicity of infection (MOI) of 0.01. After 1 hour incubation, each

well was washed with 1 ml of DPBS 3 times to remove unbound virus. Next, viral growth

medium supplemented with 1 ug/ml TPCK trypsin for IAV was added to each well. The

supernatant was sampled at 1, 24, 48, and 72 hours post-inoculation and stored at -80°C in

virus growth medium. Viral titers were assessed by plaque assay.

Titration to quantify infectious IAV

Plaques were counted against a white background and the titer in plaque forming units (PFU) was

determined by counting wells with individual plaques. Two to 3 dilutions of each explant were tested

once. Viral titers were recorded as the reciprocal of the highest dilution where plaques are noted and

represented as PFU per ml for liquid samples or PFU per explant for tissues. The limit of detection

(LOD) of the plaque assay was 50 PFU/ml or explant.

Explant sources, collection and processing

Respiratory tract tissues including trachea, bronchi, and lungs were used for explant studies **(Table**
2). Tissues were obtained from Indian origin rhesus macaques (*Macaca mulatta*), euthanized due to
medical conditions, who were born and raised at the California National Primate Research Center,
University of California, Davis, CA. The respiratory tract tissues of marine mammals were from wild
California sea lions (*Zalophus californianus*) and Northern elephant seals (*Mirounga angustirostris*)
who stranded on beaches and were euthanized due to medical conditions at The Marine Mammal
Center in Sausalito, CA. The Marine Mammal Center rehabilitates stranded marine mammals. Some
animals that present with severe conditions are not able to be returned to the wild and are therefore
euthanized. To reduce confounding effects of respiratory disease on IAV infection, we intentionally
excluded animals whose necropsy reports indicated respiratory disease as the primary or sole reason
for euthanasia. Serum harvested from blood collected at necropsy from macaques and marine
mammals were tested for IAV antibody directed against a conserved epitope of the nucleoprotein
using an enzyme-linked immunosorbent assay (ELISA) ID-Screen® Influenza A Antibody Competition
Multi-species kit, (IDvet, Grabels, France) following the manufacturer's instructions. Positive and
negative controls included in the kit were also tested on each plate. An ELX808 BioTek
Spectrophotometer (BioTek Instruments, Winooski, VT) was used to measure absorbance. Sera was
considered positive for IAV antibody when the ratio of the absorbance of the test sample to the
negative control was less than 0.45, as previously established (21, 22). Marine mammals were also
tested for IAV RNA by reverse transcription polymerase chain reaction (RT-PCR). Nasal and rectal
swabs were collected from each California sea lion and Northern elephant seal by veterinary staff at
The Marine Mammal Center upon entry to the facility. Swabs were placed in vials containing 1.5 ml of
viral transport media (VTM). Samples were refrigerated for up to one week prior to shipping to the
laboratory. Once received at the laboratory, swab samples were processed the same day or stored at
−80°C. RNA was extracted from swab samples using the MagMAX-96 AI/ND Viral RNA Isolation Kit
(Applied Biosystems, Foster City, CA) and a KingFisher Magnetic Particle Processor (Thermo
Scientific, Waltham, MA). Extracted RNAs from swab samples were subjected to an established IAV
RT-PCR (32) targeting a conserved region of the matrix gene using the AgPath-IDᴛᴍ One Step RT-
PCR mix (Applied Biosystems, Foster City, CA) and an ABI 7500 real-time PCR System (Applied
Biosystems, Foster City, CA). The positive control, a cell culture isolate of ES/H1N1, and a negative
control, VTM, were also tested on each plate. Samples with a cycle threshold (Ct) value <45 were
considered positive (33). Tissues from all three species were stored for up to 24 hours in Roswell
Park Memorial Institute-1640 (RPMI) medium (Gibco) at 4°C. During tissue preparations, lungworms
(*Parafilaroides decorus*) were observed, either in airways (i.e., bronchioles/bronchi) or sometimes free
in the lung parenchyma (in which case they were occasionally associated with mild inflammation), in
all California sea lions. Areas of tissues without worms were selected as explants. No worms were
observed grossly in explants during the IAV infection studies. Prior to preparation for infection studies,

tissues were separated and washed 4-6 times in 50 ml conical tubes containing 30 ml of RPMI

medium for 2-5 min each at 27°C. The final wash was performed in RPMI-growth-medium (RPMI

medium supplemented with 100 U/ml each of penicillin and 100 ug/ml streptomycin (Gibco) and 1x

antibiotic-antimycotic (Gibco)).

Preparation of tracheal and bronchi explants

A simplified *ex vivo* culture procedure was derived from the methods described by Nunes *et al.*

(Nunes et al, 2010). After washing in RPMI medium, the surrounding connective tissue exterior

to the tracheal and bronchi cartilage was removed. Trachea were cut into O-rings horizontally

where each slice contained about 0.5 cm cartilage. Each O-ring consisted of the respiratory

mucosa, epithelial cell layer, and underlying cartilage. O-rings were further cut into small pieces

of approximately 0.5 cm x 0.5 cm. Bronchi were cut similarly. Explants were implanted singly

onto agarose plugs (RPMI-growth-medium containing 0.5 % agarose) in 24-well plates with the

mucosal surface facing up. Explants were maintained in a humidified 37°C incubator with 5%

CO_2 for up to 7 days.

Preparation of lung, liver, and kidney explants

Lung tissues were used to investigate IAV tissue specificity. The liver and kidney of two rhesus

macaques were also used as non-target tissue controls where IAV infection was not expected.

Processing of these tissues was the same as for trachea and bronchi, where tissues were cut

into pieces of approximately 0.5 cm^3 after washing. Each piece was implanted onto agarose

plugs and incubated using the same conditions as for trachea and bronchi.

Ex vivo IAV infection kinetics

To evaluate IAV infection kinetics *ex vivo*, trachea, bronchi, and lung explants were inoculated

in triplicate (when sufficient tissue was available) with 1 x 10^4 PFU of one of each of the 6 IAV

strains in 10 ul viral growth medium onto the epithelial surface of each section for rhesus

macaques. The inoculum volume for marine mammals was adjusted to 20 ul at a dose of 2 x

10^4 PFU for each IAV strain. Viral growth medium was used in the mock-infected control

explants. Inoculated explants were sampled at 1 hour post inoculation (hpi) and every 24 hours

up to 72 hpi (or as indicated in graphs) for viral quantification and histology. For viral

quantification, each infected explant was directly immersed in 0.5 ml RPMI-growth-medium in a

2 ml centrifuge tube with a 5 mm glass bead and homogenized in a Mixer Mill MM300 (Retsch,

Leeds, UK) at 30 hertz for 4 minutes at 22°C followed by centrifugation at 16,000 g for 4

minutes and storage at -80°C. Titers of the released progeny virus from explants were

determined by plaque assay. Titers are reported as the geometric mean of triplicate explants at

each time point.

Stability assays

To determine whether IAV detected after inoculation was due to productive virus infection in

respiratory tissues, we performed viral stability assays in the absence of tissue sections.

Medium containing 1×10^4 PFU/ml of each of the 6 IAV strains was added to 24-well tissue

culture plates and supernatant was collected at 0, 24 and 48 hpi for the determination of viral

titers by plaque assay. As a second step, the stability assays were also performed using non-

target tissues (liver and kidney) to confirm that the viral titers detected in respiratory tissues

reflect productive infection in target cells. Liver and kidney explants were inoculated with 1 x

10^4 PFU of each of the 6 IAV strains in 10 ul RPMI growth medium and assayed by the same

methods used for respiratory tract tissues.

IAV immunohistochemistry

Tracheal, bronchial, and lung explants from a subset of IAV-inoculated animals (M11, M13, SL08, and

SL12) were fixed in 10% buffered formalin and embedded in paraffin. Explants not treated with IAV

and incubated only with rabbit IgG (Invitrogen) as primary antibody were included as negative isotype

controls for background staining. These animals were selected since they showed higher IAV titers

than others. Antigen retrieval was performed on 5 um sections via incubation in AR10 (Biogenex) in a

digital decloaking chamber (Biocare Medical) for 2 minutes at 125°C, followed by cooling to 90°C for

10 minutes, rinsing with water, and a final Tris-buffered saline with 0.05% Tween 20 rinse. Tissue

sections were exposed to the primary rabbit anti-influenza A virus nucleoprotein antibody (LSBio), at a

ratio of 1:250 with antibody diluent (Dako). Tris-buffered saline with 0.05% Tween 20 was used for all

washes. Nonspecific binding sites were blocked with 5% bovine serum albumin (Jackson

ImmunoResearch). Binding of the primary antibody was detected using Envision rabbit polymer with

AEC as the chromogen (Dako). Each tissue section was evaluated independently by two pathologists.

Sections from slides were visualized with a Zeiss Imager Z1 (Carl Zeiss). Digital images were

captured and analyzed using Openlab software (Improvisation). Cells with nuclear immunoreactivity

were considered positive. The positive cells in the epithelium of the trachea were manually counted.

The area of the tracheal epithelium was measured. The number of positive cells was presented as

cells per square millimeter of the epithelium.

Influenza A virus genome sequencing

IAV RNA from inocula and homogenized explant samples harvested 48 to 96 hpi from selected

animals for all 3 species were sequenced. Viral RNAs were extracted using the Magmax-96

AI/ND Viral RNA isolation kit (Applied Biosystems). Viral RNA extracts were used as a template

for multi-segment RT-PCR reactions to generate all 8 genomic segments of IAV using

procedures described previously (22, 34). Consensus sequences were generated at the Icahn

School of Medicine at Mount Sinai, as described previously (35). Genome sequences from

explant samples were compared to inocula.

Statistical analyses

Statistical analyses were conducted using GraphPad Prism software version 8. Two-way

ANOVA with Dunnett's multiple comparisons tests were performed to compare the mean viral

titers in explants at 24, 48 and 72 hpi to the mean titer 1 hpi as well as mean viral titers in

MDCK cells for all 6 IAV strains at 24, 48 and 72 hpi. IAV titers in tissue explants were plotted

at 0, 24, 48, and 72 hpi and mean area under the curve (AUC) was calculated for all titers

above the assay limit of detection. Within individual tissues, mean AUC was compared by one-

way ANOVA using Tukey's method for multiple comparisons between all IAV strains. AUC for

California sea lions grouped by reason for euthanasia (respiratory versus non-respiratory) were

compared by Mann-Whitney rank test. Mean titers for IAV strains and tissues after inoculation

of rhesus macaque and California sea lion explants were computed using R (36). Welch

ANOVA with the Games-Howell post-hoc tests were used to compare changes in mean $\log_{10}$

viral titers from 1 to 24, 48, and 72 hpi across IAV strains, tissues within species, and by IAV

strain and tissue pairs between species.

Data availability

Sequencing data are available at the CEIRS Data Processing and Coordinating Center (DPCC)

Project Identifier: SP4-Boyce_5004 and Submission_ID: 1136547734004 and at GenBank accession

numbers MW132167-MW132399.

Results

Influenza A virus strain selection

We compared relative infection kinetics for 6 IAV strains (**Table 1**) used in this study. Since

marine mammals and water birds share the same near shore environments where cross-

species transmission can occur, we used strains isolated from both marine mammals and birds.

The strains used also represent common IAV subtypes. Further, the elephant seal (ES)/H1N1

strain shows high genetic similarity with human pandemic H1N1 (4). Strains were isolated from

an elephant seal (ES, N=1), or a harbor seal (HS, N=1), or mallard ducks (m, N=4), and they

share similar passage histories in embryonated chicken eggs and MDCK cells.

323 *Influenza A virus strains exhibit differential infection kinetics in immortalized cells*

Given that cell infection kinetics for 4 of the strains had not previously been established, we first

performed growth assays for each of the 6 IAV strains after inoculation into MDCK cells in

triplicate at a MOI of 0.01 (**Figure 1**). The H3N8 strain isolated from a harbor seal (HS/H3N8),

exhibited the slowest and lowest growth kinetics, peaking at 10^5 PFU/ml 48 hpi. The H1N1

strains showed the fastest and highest growth kinetics, peaking at $>10^7$ PFU/ml by 24 hpi. The

viral infection kinetics of the other 4 strains tested, 3 of which were isolated from mallard ducks

and 1 from a harbor seal, were intermediate between HS/H3N8 and the 2 H1N1 strains. Titers

for all strains were significantly higher than those for HS/H3N8 at one or more time points

between 24-72 hpi (two way ANOVA). Given these data show that *in vitro* infection kinetics in

an immortalized cell line are different for the 6 IAV strains used in this study, we sought to

determine whether the growth kinetics of these strains also differed in explant tissues from the

3 animal species.

337 *Influenza A viruses infect rhesus macaque respiratory tract explants*

Given that marine mammal tissues are only opportunistically available from wild animals

treated at The California Marine Mammal Center in Sausalito, CA, USA, we first established the

ex vivo explant system using rhesus macaques that are more regularly available from the

California National Primate Research Center, Davis, CA, USA (**Figure 2A**) . The rhesus

macaques were bred and then euthanized at the California National Primate Research Center

due to non-respiratory medical conditions. Although tissues were used within 6 h post-mortem,

we microscopically visualized movement of glass beads by cilia placed atop tracheas from

some macaques and verified viability at 24, 48, and 72 hours, similar to Nunes *et al.* (23). To

determine whether rhesus macaque explants support IAV infection and to define infection

kinetics, we measured viral titers for 7 days (from 1 to 168 hpi) in explants from the first animal,

an 11-year old female macaque inoculated with HS/H3N8 (**Figure 2B**) who had no detectable

IAV antibody in sera at the time of necropsy. We defined infection as detection of a kinetic

increase in infectious IAV titer above the inoculum, 10^4 PFU. Titers increased over time in all 3

tissues and exceeded inocula from 48-168 hpi in trachea, from 24-120 hpi in bronchi, and from

72-96 hpi in the lung. Maximal titers $>10^5$ PFU/tissue were detected in all 3 tissue types.

Decreases in titers over time (i.e. 120-168 hpi) were likely concomitant with death of target cells

in explants. By contrast, when 10^4 PFU of IAV strains were inoculated into medium in the

absence of explants (**Figure 2C**) or into non-IAV target kidney (**Figure 2D**) or liver (**Figure 2E**)

from a 3.5 year old macaque, no IAV was detected above the limit of detection (50

PFU/explant) at 24 or 48 hpi, which indicates IAV are not viable over time absent IAV-

susceptible tissues. Together these results demonstrate that rhesus macaque respiratory tract

explants support productive infection by IAV.

Influenza A virus exhibits strain-specific infection patterns in ex vivo respiratory tract

tissues

Rhesus macaques: To determine whether rhesus macaques share similar susceptibilities to

different IAV subtypes, we inoculated respiratory tract explants from 10 rhesus macaques

(**Table 2**) with 6 virus strains. Five of the macaques were females and five were males and they

ranged from 1 to 16 years of age. Most of the macaques were euthanized due to injuries

sustained due to non-respiratory conditions including trauma or chronic diarrhea. All macaques

tested IAV seronegative by ELISA in serum from blood collected at necropsy (**Table 2**). We

focused on 0 to 72 hpi since increases in IAV titers were observed over that period in the

preliminary experiment (**Figure 2B**). We first determined kinetics of mean IAV titers in explants

from trachea, bronchi, and lung from all rhesus macaques considered together (**Figure 3A-C**).

Mean IAV infection kinetics increased from 0 to 72 hpi in all 3 explant tissues and reached

highest levels at 72 hpi for most strains. For most IAV strains, slower kinetics and peak titers

that were 10^{2-3} PFU/explant lower were observed in lungs compared to trachea and bronchi.

The mean area under the curve (AUC) was significantly higher for H3N8 and H7N5 strains

compared to H1N1 and H5N2 strains in trachea and bronchi and trended higher in the lung,

although not significantly (**Figure 3D**). Infection kinetics in explants from a representative

rhesus macaque (M15) are shown in detail (**Figure 4**) and parallel strain differences were

observed in explants from the mean of all 10 animals. Infection kinetics in individual rhesus

macaques over time (**Figure 5A**) also reveal the pattern of lower titers in the lung compared to

trachea or bronchi. To verify the histologic integrity of explants, hematoxylin and eosin (H&E)

staining of tracheal (**Figure 6A-D**) and bronchial explants (not shown) from 2 randomly

selected macaques was performed. Microscopically, rhesus macaque tracheal and bronchial

explants appeared viable for 48-72 hpi. Normal architecture of ciliated columnar respiratory

epithelium, underlying lamina propria, submucosa, and cartilage were generally maintained.

Conversely, bronchioles and alveoli were less well preserved. After 48-72 hpi, explants

exhibited variable but progressive loss of cilia at the apical epithelial surface, scattered vacuolar

degeneration or single-cell necrosis of respiratory epithelial cells, and occasional small foci of

epithelial cell loss, though these lesions were also occasionally observed at 0 or 24 hpi. No

explants showed evidence of cytopathic effects after IAV inoculation. Together, these results

show that multiple IAV subtypes infect viable respiratory tract explants from rhesus macaques

with variable kinetics and peak titers.

Marine mammals: We next examined whether respiratory tract explants from marine mammals are

susceptible to infection with the same IAV strains used to infect rhesus macaques, including the 2

strains of marine mammal origin. Explant tissues were obtained from wild California sea lions (N=11)

and Northern elephant seals (N=2) euthanized due to non-respiratory medical conditions at The

Marine Mammal Center in Sausalito, CA. Marine mammals were mostly male and ranged in age from

weanling to adult (**Table 2**). All animals were recovered stranded on beaches. The causes of death

varied and were sometimes ascribed to an infectious or toxic etiology but were not typically due to

respiratory disease. To assess whether Tissues from animals with gross respiratory tract pathologies

at necropsy were also excluded from this study. All California sea lions had a mild burden of a lung

worm, *Parafilaroides decorus*, which is found as a normal occurrence in healthy pinnipeds (37).

Explants were sectioned to avoid worms and no worms were observed grossly during the IAV

infection studies. All marine mammals tested IAV seronegative by ELISA and IAV RNA negative in

nasal and rectal swabs at intake to The Marine Mammal Center by RT-PCR (**Table 2**). Although

tissues were used within 24 h post-mortem, we confirmed ciliary viability by microscopically visualizing

movement of polystyrene beads floating atop cultured California sea lion tracheal tissue from 0 to 72

h. Viability was evidenced by clearance of beads to the side of explants within 60 minutes. As in

rhesus macaques, IAV infection kinetics for most strains increased in California sea lions from 0 to 72

hpi in all 3 tissues and reached highest levels at 72 hpi for most strains (**Figure 3E-G**). Similar to

rhesus macaques, the mean area under the curve (AUC) was significantly higher for H3N8 and H7N5

strains compared to H1N1 and H5N2 strains in trachea and bronchi while the AUC in lung explants

from sea lions was only higher for H7N5 (**Figure 3H**). Detailed data (**Figure 7**) from 1 representative

California sea lion (SL12) paralleled patterns from the mean of all 11 animals. Infection kinetics in

individual marine mammals over time (**Figure 5B**) showed higher titers in the bronchi relative to

trachea or lung. Contrary to our expectation that IAV of marine mammal origin would produce high

titers in marine mammal explants, the strain from an elephant seal (ES/H1N1) produced lower and

slower kinetics (**Figure 3E-G, 5B**) than many of the strains isolated from mallards. Like rhesus

macaque explants, California sea lion tracheobronchial explants maintained a relatively normal

microscopic appearance for 48-72 hours (**Figure 6E-H**), with poor preservation of bronchioles and

alveoli and progressive loss of normal architecture over time. All California sea lion tissues also

exhibited variable degrees of pre-existing tracheobronchitis, likely associated with their nematode

burden. Although California sea lions with respiratory conditions or gross respiratory pathologies were

excluded from this study, we also analyzed whether systemic conditions including domoic acid toxicity

or septicemia may have exacerbated IAV susceptibility. We assessed AUC for IAV infection kinetics in

sea lions euthanized for non-systemic and non-respiratory conditions including malnourishment or

leptospirosis (N=6, SL01,SL03-07) versus systemic conditions including domoic acid toxicity or

septicemia (N=5, SL02, SL08-13). The AUC infection kinetics between the 2 groups of animals were

not significantly different in any of the three respiratory tissues (data not shown). These analyses

suggest that IAV infection kinetics in wild California sea lion explants are not impacted by systemic

conditions. Together, these results show that, similar to rhesus macaques, multiple IAV subtypes

infect viable respiratory tract explants from marine mammals with variable kinetics and peak titers. To

further determine whether the differences in mean IAV kinetics were supported statistically over time,

we examined IAV infection dynamics in 24 hour windows.

IAV Kinetics Over Time: We tested whether explants inoculated with different IAV strains

experience different mean changes in titer from 1 to 72 hpi. Considering all 3 tissue types

together, the $\log_{10}$ change in viral titer was calculated for three time frames: 1 versus 24, 1

versus 48, and 1 versus 72 hpi for rhesus macaques and California sea lions (**Table 3**).

Northern elephant seals were not included in analyses since explants from only two animals

were available, although they produced infectious IAV after inoculation with some strains

(**Figure 5B**). Infection kinetics of the six IAV strains differed for each time frame considered. In

rhesus macaques, mean titer changes for HS/H3N8, m/H3N8, and m/H7N5 did not differ

significantly from each other over any of the time frames. m/H1N1 and mH5/N2 did not differ

significantly from each other over any of the time frames but were different from HS/H3N8,

m/H3N8 and m/H7N5. By contrast, the change in ES/H1N1 viral titers significantly differed from

all other strains except m/H1N1 from 1 to 48 hpi. In California sea lions, HS/H3N8 and m/H7N5

shared similar mean changes in $\log_{10}$ titers in all time frames. For m/H1N1 and HS/H3N8, the

mean change in $\log_{10}$ titers did not vary significantly from 1 to 24 hpi but did vary from 1 to 48

and 1 to 72 hpi. The mean change in $\log_{10}$ viral titers between m/H1N1 and m/H7N5 were only

different from 1 to 48 hpi and m/H5N2 and m/H7N5 differed from 1 to 48 and 1 to 72 hpi. These

results show that mean changes in IAV titers over 24 hour windows in respiratory tract explants

from both rhesus macaques and California sea lions vary with IAV strain.

IAV Kinetics by Tissue: To examine whether tissue type associated with IAV titer, we analyzed

all 6 IAV strains together (**Table 4**). Infection kinetics in bronchi and trachea from rhesus

macaques were significantly higher than in the lung over all 24 hour windows from 1 to 72 hpi.

In California sea lions, infection kinetics in bronchi and trachea were significantly higher than in

the lung at 24 hpi, while infection kinetics in bronchi were significantly higher than in the trachea

and lung at 48 and 72 hpi. Together, these data support tissue-specific IAV infectivity, where

bronchi and trachea from both species support production of higher IAV titers than lung

explants.

IAV Kinetics by Species: To evaluate whether species associates with infection kinetics for the

6 IAV strains, we compared changes in mean $\log_{10}$ titer between rhesus macaque and

California sea lion explants (**Table 5**). The mean change in titer of both H3N8 strains and

m/H7N5 was significantly higher in trachea at all time points and in bronchi to 48 hpi in rhesus

macaques compared to California sea lions. Changes in mean $\log_{10}$ titers for m/H3N8 in the

lung from 1 to 48 and 1 to 72 hpi were higher in rhesus macaques compared to California sea

lions. Together these data show that rhesus macaque explants produce higher titers compared

to California sea lions for most IAV strains used here.

Cellular tropism of influenza A virus in rhesus macaque and California sea lion explants

We used immunohistochemistry to assess the cellular tropism of IAV in rhesus macaque and

California sea lion respiratory tissue explants (**Figure 8**, **Table 6)**. At least 22 sections from each

tissue type for 2 rhesus macaques (M11 and M13) and at least 56 sections from each tissue type for 2

California sea lions (SL08 and SL12) were evaluated. All examined sections exhibited variable

degrees of nonspecific cytoplasmic, membranous and/or background staining; as such, only cells with

strong nuclear immunoreactivity were considered positive. For all explants from both species and all

IAV subtypes, positive staining for viral nucleoprotein was limited to epithelial cells of the trachea and

bronchi; representative slides are shown (**Figure 8A-D**). Staining for IAV nucleoprotein was not

observed in non-inoculated explants or IHC antibody-treated negative isotype controls from rhesus

macaque **(Figure 8E-F)** or California sea lion **(Figure 8G-H)** trachea. Pneumocytes within the lung

were IAV negative. Infection was primarily localized to apical epithelial cells, particularly tracheal and

bronchial ciliated columnar cells and goblet cells, while basal cells were largely spared. Rhesus

macaque explants from M11 and M13 infected with H3N8 strains exhibited higher relative proportions

of positive tracheal and bronchial epithelial cells 24 and 48 hpi, even though IAV titers at those times

for most strains exceeded 10^4 PFU/explant 24 hpi (**Figure 5A**). The explants from M11 infected with

m/H7N5 exhibited the largest variation in relative proportion of positive cells between 24 and 48 hpi.

The number of infected respiratory epithelial cells generally increased between 24 and 48 hpi, which

is consistent with infection kinetics where IAV titers increased during that period. While positive nuclei

were less frequent within California sea lion explants, those infected with H3N8 strains also exhibited

higher relative proportions of positive tracheal and bronchial epithelial cells at 24 and 48 hpi, which is

consistent with data from rhesus macaques. These results demonstrate that IAV infects the apical

epithelial cells of upper respiratory tract explants in rhesus macaques and California sea lions.

498 *Genetic changes in influenza A viruses sequenced from explants*

499 IAV genomes in inoculated explants from 2 rhesus macaques, 3 California sea lions, and 1 Northern

500 elephant seal were compared to sequences of the strains used as inocula (**Table 7**). Sequence

501 comparisons showed that all genomes from rhesus macaque explants were 100% identical at the

502 consensus level (mutations occurring on 50% or more of viral RNAs) to their corresponding inoculum

503 (data not shown). Six nonsynonymous nucleotide substitutions in the HA, NA, and PA genes were

504 detected in California sea lion and Northern elephant seal explants inoculated with different IAV

505 subtypes (**Table 8**). Inocula contained a mixture of bases including the mutant nucleotide at each of

506 these loci, indicating that none of the mutations developed *de novo* in explants (data not shown). One

507 synonymous nucleotide substitution, PB2 1368 in m/H7N5, was detected in bronchi from all 3

508 California sea lions at 48 and 72 hpi. The nucleotide affecting this substitution was present at a

509 minority frequency (41%) in the inoculum, indicating that it did not develop *de novo* but increased to

510 consensus frequency in explants by 48 hpi (data not shown). Together these data show that no *de*

511 *novo* IAV mutations were detected in any strain during *ex vivo* respiratory tract explant infection from

512 selected rhesus macaques and marine mammals.

513

514 **Discussion**

515 We adapted an *ex vivo* respiratory tract infection model to study susceptibility and to define

516 comparative infection kinetics of IAV in colony-raised rhesus macaques and wild marine

517 mammals. We observed that IAV exhibits temporal, subtype- and species-dependent infection

518 kinetics in explants from both rhesus macaques and California sea lions. Although the relative

519 infection kinetics for the six strains was similar for explants from rhesus macaques and

520 California sea lions, similar patterns were not observed in MDCK cells. This suggests that

521 immortalized cell lines may not accurately represent infection phenotypes *in vivo*, further

522 underscoring the value of using *ex vivo* systems.

523 E*x vivo* respiratory tract models have been used for studying respiratory pathogens of humans,

524 swine, bovines, canines, and equines (23, 25–28, 38–40). *Ex vivo* cultures of respiratory

525 tissues provide a close resemblance to the respiratory tract *in vivo* by: 1) shared polarity, where

526 the basolateral surface is exposed to culture medium and the apical cell surface is exposed to

527 air, 2) shared cell positioning *in situ* which maintains virus receptor distribution, 3) possession

528 of multiple cell types and states of differentiation, 4) maintenance of the three dimensional

529 integrity and architecture of a tissue, unlike cell monocultures, and 5) ciliary activity to preserve

530 mucociliary clearance. Given these similarities to living IAV hosts, explants are valuable tools

for assessing host susceptibility, infection kinetics, and pathogenesis of respiratory pathogens.

Ex vivo systems also better assess infectivity compared to cell monoculture binding assays that

only measure virus-receptor affinity (25, 27). In addition, tissues collected from a single animal

can be divided into pieces and used to compare relative infectivity of different viral strains while

holding the host constant. Last, use of explants from animals euthanized for other reasons

reduced the number of animals used in research, following the principles of reduction,

replacement, and refinement.

Limitations of the *ex vivo* approach include restricted availability of tissues, short periods of viability,

and inter-animal variability in IAV susceptibility. The absence of a blood supply also constitutes a

weakness since recruitment of immune cells to the infection site cannot occur. However, such a pitfall

might be co-opted in future studies to understand the effects of innate and intrinsic immunity without

immune cell influx. Use of explants from wild marine mammals euthanized for non-respiratory

conditions, some of which present systemically, is another limitation of this study. Some of the marine

mammals used in this study showed evidence of exposure to toxins, malnourishment, or congenital

defects, factors that may modify IAV susceptibility. However, systemic conditions including toxin

exposure did not correspond to observed differences in IAV infection capacity in California sea lion

explants in the current study. Given that the only marine mammals available for this study were those

euthanized for other conditions, we unfortunately do not have the opportunity to study tissues from

healthy animals, especially since both California sea lions and elephant seals are protected by the

Marine Mammal Protection Act.

Lower IAV titers in *ex vivo rhesus* macaque and marine mammal lungs relative to trachea and bronchi

support the upper respiratory tract as the primary site of virus infection with the virus strains used in

this study. The ability of IAV to produce higher titers in the upper respiratory tract may reflect

adaptation of the virus to infect cell targets in closest anatomic proximity to its entry point and

shedding site, the respiratory mucosa. Lower susceptibility of the lung to IAV infection could result

from decreased receptor expression in that tissue. We did not identify or enumerate IAV receptor

expression in explants in this study to confirm this hypothesis. An alternate possibility is that lungs of

both species decayed more quickly than the trachea and bronchi, rendering them less susceptible to

IAV infection. However, we feel accelerated lung decay is unlikely given our gross and histologic

observations showed similar architectural integrity in all 3 tissues to 72 hpi (data not shown).

Alternately, lower IAV lung infectivity could result from increased resident immune cell activation and

response in that tissue.

The infection kinetics of harbor seal-origin H3N8 and mallard-origin H7N5 in California sea lion
respiratory tract explant tissues were superior to other subtype strains isolated from mallards
and elephant seal-origin H1N1. Since all strains share similar passage histories, infection
kinetic differences are not likely due to varying passage or changes in consensus sequence
since all were nearly identical to those of their unpassaged progenitors. Our observation that
the elephant seal H1N1 infection kinetics were lower than other strains in California sea lions
suggests that having a marine mammal as the isolate source does not necessarily translate to
higher marine mammal explant infectivity. The lack of augmented infectivity in explants of the
species from which the isolate was made has also been observed in swine, where strains of the
same subtype isolated from both swine and humans showed similar infection kinetics (41).
Further, some human-isolated IAV show great inter-strain variability in infection kinetics in
human explants (27). In spite of the different infection kinetics across strains, the 6 strains
used here, representing 4 IAV subtypes, productively infected at least one explant type for both
species. These data suggest that any of these subtypes could also infect marine mammals in
the wild, consistent with serologic data showing exposure to many subtypes (13–16, 18, 19).
Together, data presented here show that the marine mammal *ex vivo* culture of respiratory
tissues is a tool to study IAV susceptibility, host-range, and tissue tropism.
582

583 **ACKNOWLEDGEMENTS**
584 We are grateful to the veterinary and research staff at The Marine Mammal Center for coordinating
585 the collection and transfer of fresh tissue samples to our laboratory. Sample possession was granted
586 by a Letter of Authorization from the National Oceanic and Atmospheric Administration National
587 Marine Fisheries Service.
588

FUNDING ACKNOWLEDGEMENTS
Funding support was provided by NIH NIAID #HHSN272201400008C to WB and LLC and from the
University of California, Davis Office of Research to PAP and HL. EEB was supported by a
Comparative Medical Science Training Program T32 fellowship # OD011147. We thank Kathy West at
the California National Primate Research Center for generating the rhesus macaque image used in
figures and Ana L. Ramirez for generating the other images used in figures.
595

596 **COMPETING INTERESTS**
597 The authors declare that there are no competing interests.
598

599 **AUTHOR CONTRIBUTIONS**

Conceptualization: HL, MP, WB, LLC, Methodology: HL, CMW, LLC, BH, Investigation: HL, MP, CMW, OG-V, AMA, EEB, ZMM, KMH, KJ, PAP, ZK, DK. Writing-original draft: HL, LLC. Writing-review and editing: HL, CMW, MP, EEB, ZMM, HB, BH, KMH, WB, PAP, LLC. Visualization: HL, CMW, EEB, PAP, KJ, LLC. Supervision and project administration: WB, HB, LLC. Funding acquisition: WB, HL, PAP, LLC.

REFERENCES

1. Krammer F, Smith GJD, Fouchier RAM, Peiris M, Kedzierska K, Doherty PC, Palese P, Shaw ML, Treanor J, Webster RG, García-Sastre A. 2018. Influenza. Nat Rev Dis Prim. Nature Publishing Group. 4(1):3.

2. Landolt GA, Olsen CW. 2007. Up to new tricks - a review of cross-species transmission of influenza A viruses. Anim Health Res Rev 8(1):1-21

3. Taubenberger JK, Kash JC. 2010. Influenza virus evolution, host adaptation, and pandemic formation. Cell Host Microbe. 7(6):440-51.

4. Goldstein T, Mena I, Anthony SJ, Medina R, Robinson PW, Greig DJ, Costa DP, Lipkin WI, Garcia-Sastre A, Boyce WM. 2013. Pandemic H1N1 Influenza Isolated from Free-Ranging Northern Elephant Seals in 2010 off the Central California Coast. PLoS One. 8(5):e62259.

5. Karlsson EA, Ip HS, Hall JS, Yoon SW, Johnson J, Beck MA, Webby RJ, Schultz-Cherry S. 2014. Respiratory transmission of an avian H3N8 influenza virus isolated from a harbour seal. Nat Commun. 5:4791.

6. Reperant LA, Kuiken T, Osterhaus ADME. 2012. Adaptive pathways of zoonotic influenza viruses: From exposure to establishment in humans. Vaccine. 30(30):4419-34.

7. Kaplan BS, Webby RJ. 2013. The avian and mammalian host range of highly pathogenic avian H5N1 influenza. Virus Res. 178(1): 3–11.

8. Sikkema RS, Freidl GS, de Bruin E, Koopmans M. 2016. Weighing serological evidence of human exposure to animal influenza viruses − A literature review. Eurosurveillance. 21(44):30388.

9. Suttie A, Deng YM, Greenhill AR, Dussart P, Horwood PF, Karlsson EA. 2019. Inventory of molecular markers affecting biological characteristics of avian influenza A viruses. Virus Genes. 55(6):739-768.

10. Lang G, Gagnon A, Geraci JR. 1981. Isolation of an influenza A virus from seals. Arch Virol.

11. Webster RG, Hinshaw VS, Bean WJ, Van Wyke KL, Geraci JR, St. Aubin DJ, Petursson G. 1981. Characterization of an influenza A virus from seals. Virology. 113(2):712-24.

12. Geraci JR, Aubin DJS, Barker IK, Webster RG, Hinshaw VS, Bean WJ, Ruhnke HL, Prescott JH, Early G, Baker AS, Madoff S, Schooley RT. 1982. Mass mortality of harbor seals: Pneumonia associated with influenza A virus. Science 215(4536):1129-31

13. Ohishi K, Kishida N, Ninomiya A, Kida H, Takada Y, Miyazaki N, Boltonov AN, Maruyama T.
2004. Antibodies to human-related H3 influenza A virus in Baikal seals (*Phoca sibirica*) and
ringed seals (*Phoca hispida*) in Russia. Microbiol Immunol. 48(11):905-9.

14. Fereidouni S, Munoz O, Von Dobschuetz S, De Nardi M. 2014. Influenza virus infection of
marine mammals. Ecohealth. 13(1):161-70.

15. Puryear WB, Keogh M, Hill N, Moxley J, Josephson E, Davis KR, Bandoro C, Lidgard D,
Bogomolni A, Levin M, Lang S, Hammill M, Bowen D, Johnston DW, Romano T, Waring G,
Runstadler J. 2016. Prevalence of influenza A virus in live-captured North Atlantic gray seals: A
possible wild reservoir. Emerg Microbes Infect. 5(8):e81.

16. Stuen S, Have P, Osterhaus AD, Arnemo JM, Moustgaard A. 1994. Serological investigation of
virus infections in harp seals (*Phoca groenlandica*) and hooded seals (*Cystophora cristata*). Vet
Rec. 134(19):502-3.

17. Nielsen O, Clavijo A, Boughen JA. 2001. Serologic evidence of influenza A infection in marine
mammals of Arctic Canada. J Wildl Dis. 37(4):820-5.

18. Greig DJ, Gulland FMD, Smith WA, Conrad PA, Field CL, Fleetwood M, Harvey JT, Ip HS,
Jang S, Packham A, Wheeler E, Hall AJ. 2014. Surveillance for zoonotic and selected
pathogens in harbor seals phoca vitulina from central California. Dis Aquat Organ. 111(2):93-
106.

19. Danner GR, Mcgregor MW, Zarnke RL, Olsen CW. 1998. Serologic evidence of influenza virus
infection in a ringed seal (*Phoca hispida*) from Alaska. Mar Mammal Sci.14(2):380-384.

20. Venkatesh D, Bianco C, Núñez A, Collins R, Thorpe D, Reid SM, Brookes SM, Essen S,
McGinn N, Seekings J, Cooper J, Brown IH, Lewis NS. 2020. Detection of H3N8 influenza A
virus with multiple mammalian-adaptive mutations in a rescued Grey seal (*Halichoerus grypus*)
pup. Virus Evol. 18;6(1):veaa016.

21. Capuano AM, Miller M, Stallknecht DE, Moriarty M, Plancarte M, Dodd E, Batac F, Boyce WM.
2017. Serologic detection of subtype-specific antibodies to influenza A viruses in southern sea
otters (*Enhydra lutris nereis*). J Wildl Dis. 53(4):906-910.

22. Ramey AM, Hill NJ, Cline T, Plancarte M, De La Cruz S, Casazza ML, Ackerman JT, Fleskes
JP, Vickers TW, Reeves AB, Gulland F, Fontaine C, Prosser DJ, Runstadler JA, Boyce WM.
2017. Surveillance for highly pathogenic influenza A viruses in California during 2014-2015
provides insights into viral evolutionary pathways and the spatiotemporal extent of viruses in
the Pacific Americas Flyway. Emerg Microbes Infect. 6(9):e80.

23. Nunes SF, Murcia PR, Tiley LS, Brown IH, Tucker AW, Maskell DJ, Wood JLN. 2010. An *ex
vivo* swine tracheal organ culture for the study of influenza infection. Influenza Other Respi
Viruses. 4(1):7-15

24. Löndt BZ, Brookes SM, Nash BJ, Núñez A, Stagg DA, Brown IH. 2013. The infectivity of

pandemic 2009 H1N1 and avian influenza viruses for pigs: An assessment by *ex vivo*

respiratory tract organ culture. Influenza Other Respi Viruses.7(3):393-402.

25.   Chan RWY, Chan MCW, Nicholls JM, Malik Peiris JS. 2013. Use of *ex vivo* and *in vitro* cultures

of the human respiratory tract to study the tropism and host responses of highly pathogenic

avian influenza A (H5N1) and other influenza viruses. Virus Res. 178(1):133-45.

26.   Nicholas B, Staples KJ, Moese S, Meldrum E, Ward J, Dennison P, Havelock T, Hinks TSC,

Amer K, Woo E, Chamberlain M, Singh N, North M, Pink S, Wilkinson TMA, Djukanović R.

2015. A novel lung explant model for the *ex vivo* study of efficacy and mechanisms of anti-

influenza drugs. J Immunol. 194(12):6144-54.

27.   Hui KPY, Ching RHH, Chan SKH, Nicholls JM, Sachs N, Clevers H, Peiris JSM, Chan MCW.

2018. Tropism, replication competence, and innate immune responses of influenza virus: an

analysis of human airway organoids and ex-vivo bronchus cultures. Lancet Respir Med.

6(11):846-854.

28.   Hui KPY, Cheung MC, Perera RAPM, Ng KC, Bui CHT, Ho JCW, Ng MMT, Kuok DIT, Shih KC,

Tsao SW, Poon LLM, Peiris M, Nicholls JM, Chan MCW. 2020. Tropism, replication

competence, and innate immune responses of the coronavirus SARS-CoV-2 in human

respiratory tract and conjunctiva: an analysis in *ex-vivo* and *in-vitro* cultures. Lancet Respir Med

8(7):687–695.

29.   Shinya K, Gao Y, Cilloniz C, Suzuki Y, Fujie M, Deng G, Zhu Q, Fan S, Makino A, Muramoto Y,

Fukuyama S, Tamura D, Noda T, Eisfeld AJ, Katze MG, Chen H, Kawaoka Y. 2012. Integrated

clinical, pathologic, virologic, and transcriptomic analysis of H5N1 influenza virus-induced viral

pneumonia in the rhesus macaque. J Virol. 86(11):6055-66.

30.   Carroll TD, Jegaskanda S, Matzinger SR, Fritts L, McChesney MB, Kent SJ, Fairman J, Miller

CJ. 2018. A lipid/DNA adjuvant-inactivated influenza virus vaccine protects rhesus macaques

from uncontrolled virus replication after heterosubtypic influenza a virus challenge. J Infect Dis.

218(6):856-867.

31.   Skinner JA, Zurawski SM, Sugimoto C, Vinet-Oliphant H, Vinod P, Xue Y, Russell-Lodrigue K,

Albrecht RA, García-Sastre A, Salazar AM, Roy CJ, Kuroda MJ, Oh SK, Zurawski G. 2014.

Immunologic characterization of a rhesus macaque H1N1 challenge model for candidate

influenza virus vaccine assessment. Clin Vaccine Immunol. 21(12):1668-80.

32.   Spackman E, Senne DA, Myers TJ, Bulaga LL, Garber LP, Perdue ML, Lohman K, Daum LT,

Suarez DL. 2002. Development of a real-time reverse transcriptase PCR assay for type A

influenza virus and the avian H5 and H7 hemagglutinin subtypes. J Clin Microbiol. 40(9):3256-

60.

33.   Spackman E. 2014. Avian Influenza Virus Detection and Quantitation by Real-Time RT-PCR, p.

vol 1161. *In* Animal Influenza Virus. Methods in Molecular Biology (Methods and Protocols).

Hu, New York, NY.

34. Mena I, Nelson MI, Quezada-Monroy F, Dutta J, Cortes-Fernández R, Lara-Puente JH, Castro-
Peralta F, Cunha LF, Trovão NS, Lozano-Dubernard B, Rambaut A, Van Bakel H, García-
Sastre A. 2016. Origins of the 2009 H1N1 influenza pandemic in swine in Mexico. Elife.
5:e16777.

35. Venkatesh D, Poen MJ, Bestebroer TM, Scheuer RD, Vuong O, Chkhaidze M, Machablishvili
A, Mamuchadze J, Ninua L, Fedorova NB, Halpin RA, Lin X, Ransier A, Stockwell TB,
Wentworth DE, Kriti D, Dutta J, van Bakel H, Puranik A, Slomka MJ, Essen S, Brown IH,
Fouchier RAM, Lewis NS. 2018. Avian influenza viruses in wild birds: virus evolution in a
multihost ecosystem. J Virol. 92(15):e00433-18.

36. R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria.gbif.org

37. Greig DJ, Gulland FMD, Kreuder C. 2005. A decade of live California sea Lion (*Zalophus*
californianus) strandings along the Central California coast: causes and trends, 1991-2000.
Aquat Mamm. 31(1):11-22.

38. Vandekerckhove A, Glorieux S, Broeck W Van den, Gryspeerdt A, van der Meulen KM,
Nauwynck HJ. 2009. *In vitro* culture of equine respiratory mucosa explants. Vet J. 181(3):280-
7.

39. Priestnall SL, Mitchell JA, Brooks HW, Brownlie J, Erles K. 2009. Quantification of mRNA
encoding cytokines and chemokines and assessment of ciliary function in canine tracheal
epithelium during infection with canine respiratory coronavirus (CRCoV). Vet Immunol
Immunopathol. 127(1-2):38-46.

40. Niesalla HS, Dale A, Slater JD, Scholes SFE, Archer J, Maskell DJ, Tucker AW. 2009. Critical
assessment of an *in vitro* bovine respiratory organ culture system: A model of bovine
herpesvirus-1 infection. J Virol Methods. 158(1-2):123-9.

41. Van Poucke SGM, Nicholls JM, Nauwynck HJ, Van Reeth K. 2010. Replication of avian, human
and swine influenza viruses in porcine respiratory explants and association with sialic acid
distribution. Virol J. 7:38.

 FIGURE LEGENDS
745

Figure 1. *In vitro* infection kinetics of 6 influenza A virus strains in Madin-Darby canine kidney
cells. A MOI of 0.01 was used. The geometric mean virus titer from triplicate wells at each time point
was plotted as PFU per milliliter ± geometric standard deviation. The dotted line represents the limit of
detection (LOD) of 500 PFU/ml. Asterisks show comparisons of mean titers between HS/H3N8 and all
other strains analyzed by two-way ANOVA of log transformed values where * is $p < 0.001$ and ** is p
< 0.0001. Strain names refer to the original source of the virus isolate followed by the hemagglutinin
and neuraminidase subtype. HS is harbor seal. m is mallard. ES is elephant seal.
753

Figure 2. (A) Experimental design for *ex vivo* rhesus macaque respiratory tract explant
inoculations with influenza A virus (IAV). **(B)** Seven day infection kinetics of IAV HS/H3N8 in
explants from an 11 year old animal that was serologically IAV non-reactive at necropsy. **(C)** IAV titers
in the absence of explants to evaluate stability of infectious virus from 0 to 48 hpi. IAV titers in kidney
(D) and liver **(E)**, tissues that are not IAV targets from a 3.5 year old animal from 0 to 48 hpi. Each bar
shows the measurement from a single explant. The dotted black line shows the limit of detection, 50
PFU/explant. The red line shows the inoculum. hpi is hours post inoculation. Strain names refer to the
original source of the virus isolate followed by the hemagglutinin and neuraminidase subtype. HS is
harbor seal. m is mallard. ES is elephant seal.
763

Figure 3: **Ex vivo influenza A virus infection kinetics.** Mean kinetics in 10 rhesus macaque (**A-C**)
and 11 California sea lions (**E-G**) and areas under the infection curve (AUC) (**D, H**) in *ex vivo*
respiratory tract trachea, bronchi, and lung explants. Error bars show standard deviations. The dotted
black line shows the limit of detection, 50 PFU/explant. Colors in squares in D and H show differences
in mean AUC by strain analyzed using one-way ANOVA tests where the darker the color, the smaller
the p-value. Strain names refer to the original source of the virus isolate followed by the hemagglutinin
and neuraminidase subtype. HS is harbor seal. m is mallard. ES is elephant seal.
771

Figure 4. Influenza A virus infection kinetics from 1 to 72 hpi in respiratory tract explants from a 4 year
old male rhesus macaque (M15 in Table 2) inoculated with $1X10^4$ PFU of 6 viral strains. Viral titers are
represented as the geometric mean and geometric standard deviation. Three explants at each time
point were titrated independently and the mean of the triplicates is represented by the middle
horizontal line. Asterisks show p values (two-way ANOVA with Dunnett's multiple comparisons)
comparing titers at 24, 48 or 72 hpi to 1 hpi, respectively. **$p \leq 0.005$, ***$p < 0.001$, and ****$p < 0.0001$.
The dashed line indicates limit of detection, 50 PFU/explant. Strain names refer to the original source
of the virus isolate followed by the hemagglutinin and neuraminidase subtype. HS is harbor seal. m is
mallard. ES is elephant seal.
781

Figure 5. Comparison of *ex vivo* influenza A virus infection kinetics in respiratory tract explants
from individual animals. (A) IAV titers in explants from 10 rhesus macaques and **(B)** 11 California
sea lions and 2 Northern elephant seals inoculated with 1×10^4 PFU and 2×10^4 PFU of each of the 6
IAV strains over a 72 hour period. The kinetics of viral infection were determined by plaque assays of
homogenized explants, where each square represents the titer from 10^1 (blue) to 10^6 (yellow) in
PFU/explant. The limit of detection was 50 PFU/explant. Each explant was titrated once at 2-3
dilutions. Strain names refer to the original source of the virus isolate followed by the hemagglutinin

and neuraminidase subtype (Table 1). HS is harbor seal. m is mallard. ES is elephant seal. Boxes with 'X' indicate the sample was not available for titration.

Figure 6. Representative light photomicrograph of sections of (A-D) rhesus macaque M11 and (E-H) California sea lion SL08 *ex vivo* tracheal explants at 0 (A, E), 24 (B, F), 48 (C, G), and 72 (D, H) hours. Cilia (black arrows) are present on the apical surface of epithelial cells. Sections were stained with hematoxylin and eosin (400x magnification); bar=50um. RE= respiratory epithelium; LP= lamina propria.

Figure 7. Influenza A virus infection kinetics from 1 to 96 hpi in respiratory tract explants from an adult female California sea lion (SL12 in Table 2) inoculated with $2X10^4$ PFU of 6 IAV strains. Viral titers are represented as the geometric mean and geometric standard deviation. Three explants at each time point were titrated independently and the mean of the triplicates is represented by the middle horizontal line. Asterisks show p values (two-way ANOVA with Dunnett's multiple comparisons) comparing titers at 24, 48, 72 or 96 hpi to 1 hpi, respectively. **p < 0.005, ***p < 0.001, and ****p < 0.0001. The dashed line indicates limit of detection, 50 PFU/explant. Strain names refer to the original source of the virus isolate followed by the hemagglutinin and neuraminidase subtype. HS is harbor seal. m is mallard. ES is elephant seal.

Figure 8. Immunohistochemical (IHC) staining for influenza A virus in rhesus macaque and California sea lion tracheal and bronchial explants. (A) Bronchi from rhesus macaques 48 hours post inoculation (hpi) with **(A)** m/H7N5 (animal M11) or **(B)** HS/H3N8 (animal M13); **(C)** Trachea from California sea lion SL12 48 hpi with HS/H3N8; **(D)** Trachea from California sea lion SL08 24 hpi with m/H3N8; Bronchi from rhesus macaque M13, 48 hpi treated with **(E)** growth medium only or **(F)** with HS/H3N8 and stained with isotype IgG; Trachea from SL12 24 hpi treated with **(G)** growth medium only or **(H)** with HS/H3N8 and stained with isotype IgG. IHC stain used an antibody that labels IAV nucleoprotein. Positive respiratory epithelial cells exhibit strong nuclear immunoreactivity (red-brown nuclear staining; arrows). Positive staining is primarily localized to apical epithelial cells, with relative sparing of basal cells.

Table 1. IAV strains used for explant infection experiments. ES is elephant seal, HS is harbor seal, m is mallard. N/A indicates strain was isolated at UC Davis and first described in this study. MDCK is Madin-Darby canine kidney cell. PFU is plaque forming unit.

Virus ID	Strain name	Subtype	Passage number	MDCK or Vero E6 titer (PFU/ml)	GenBank Accession number	Reference
ES/H1N1	A/Elephant seal/California/1/2010	H1N1	p4	1.3×10^8	JX865419-JX865426	Goldstein et al. 2013 (4)
HS/H3N8	A/harbor seal/New Hampshire/179629/2011	H3N8	p3	1.8×10^6	KJ467564-KJ467571	Karlsson et al. 2014 (5)
m/H3N8	A/mallard/California/1475/2010	H3N8	p2	6×10^6	CY120501-CY120508	N/A
m/H1N1	A/mallard/California/3134/2010	H1N1	p2	4×10^7	CY120611-CY120618	N/A
m/H5N2	A/mallard/California/2396/2010	H5N2	p2	4×10^7	CY120587-CY120594	N/A
m/H7N5	A/mallard/California/1390/2010	H7N5	p2	1×10^7	CY120555-CY120562	N/A

Table 2. Explant sources for IAV *ex vivo* infection studies. Tissues were derived from 10 rhesus macaques (*Macaca mulatta*), 11 California sea lions (*Zalophus californianus*) and 2 northern elephant seals (*Mirounga angustirostris)*. M is macaque, SL is sea lion, ES is elephant seal, y is year, m is month.

Animal	Species	Sex	Age	Reason for euthanasia	IAV ELISA	IAV RT-PCR
M12	*Macaca mulatta*	male	1y	chronic diarrhea	negative	not performed
M13	*Macaca mulatta*	male	3y4m	lameness due to trauma	negative	not performed
M15	*Macaca mulatta*	male	4y5m	leg trauma	negative	not performed
M16	*Macaca mulatta*	male	3y6m	chronic diarrhea	negative	not performed
M17	*Macaca mulatta*	male	4y5m	chronic diarrhea	negative	not performed
M11	*Macaca mulatta*	female	5y11m	chronic diarrhea	negative	not performed
M14	*Macaca mulatta*	female	16y6m	endometriosis	negative	not performed
M19	*Macaca mulatta*	female	1y6m	chronic diarrhea	negative	not performed
M21	*Macaca mulatta*	female	4y	leg trauma	negative	not performed
M20	*Macaca mulatta*	female	3y11m	liver amyloid, diarrhea, bloating	negative	not performed
SL01	*Zalophus californianus*	male	yearling	malnourished, peritonitis	negative	negative
SL02	*Zalophus californianus*	male	yearling	leptospirosis, mild pneumonia	negative	negative
SL03	*Zalophus californianus*	male	yearling	leptospirosis	negative	negative
SL04	*Zalophus californianus*	male	juvenile	leptospirosis	negative	negative
SL05	*Zalophus californianus*	male	subadult	leptospirosis, seizures	negative	negative
SL06	*Zalophus californianus*	male	subadult	leptospirosis	negative	negative
SL07	*Zalophus californianus*	male	subadult	leptospirosis	negative	negative
SL08	*Zalophus californianus*	male	yearling	asphyxia	negative	negative
SL09	*Zalophus californianus*	female	adult	domoic acid toxicity	negative	negative
SL12	*Zalophus californianus*	female	adult	septicemia	negative	negative
SL13	*Zalophus californianus*	male	yearling	domoic acid toxicity	negative	negative
ES10	*Mirounga angustirostris*	male	yearling	tachypnea, tachycardia	negative	negative
ES11	*Mirounga angustirostris*	male	weanling	blindness; congenital defect	negative	negative

Table 3. Mean change in $\log_{10}$ titer of influenza virus strains after inoculation in rhesus macaque or California sea lion respiratory tract explant tissues. Means for the same time frame followed by a common letter are not significantly different by the Games-Howell test at the 5% level of significance. Values in parentheses show standard deviations. The symbols indicate that the mean change in titer differs significantly for at least one strain according to Welch's ANOVA at 5% level of significance for: *1 vs 24 hpi F=15.417, df=(5,91.649), p-value=5.55e-11 for rhesus macaques and F=4.666, df=(5,189.21), p-value=4.85e-04 for California sea lions, the † denotes 1 vs 48 hpi (F=25.823, df=(5,91.569), p-value=3.59e-16) for rhesus macaques and (F=5.784, df=(5,182.59), p-value=5.55e-05) for California sea lions, and the ‡ denotes 72 hpi (F=23.351, df=(5,91.826), p-value=4.58e-15) for rhesus macaques and (F=6.636, df=(5,189.35), p-value=1.02e-05) for California sea lions. Strain names refer to the original source of the virus isolate followed by the hemagglutinin and neuraminidase subtype. HS is harbor seal. m is mallard. ES is elephant seal.

	RHESUS MACAQUES			CALIFORNIA SEA LIONS		
	1 vs 24 hpi*	1 vs 48 hpi†	1 vs 72 hpi‡	1 vs 24 hpi *	1 vs 48 hpi †	1 vs 72 hpi ‡
HS/H3N8	1.2 (1.1) [a]	1.8 (1.5) [ab]	2.2 (1.5) [ad]	0.4 (0.8) [e]	0.6 (1.2) [e]	0.8 (1.4) [e]
m/H3N8	0.9 (1.1) [ac]	1.8 (1.1) [ac]	0.9 (1.4) [ab]	-0.04 (0.7) [abc]	-0.07 (1.1) [abc]	0.2 (1.3) [abc]
ES/H1N1	-0.1 (0.6) [b]	-0.1 (0.8) [e]	0.1 (0.9) [e]	-0.1 (0.5) [ab]	-0.1 (0.8) [ab]	-0.1 (0.9) [ab]
m/H1N1	0.2 (0.7) [b]	0.3 (1.0) [de]	0.9 (1.2) [c]	0.03 (0.6) [bcde]	-0.05 (0.7) [bd]	0.2 (1.0) [bcd]
m/H5N2	0.3 (0.7) [bc]	1.0 (1.4) [bcd]	1.5 (1.4) [bcd]	-0.02 (0.5) [ad]	-0.07 (0.6) [ad]	0.2 (1.0) [ad]
m/H7N5	1.1 (0.8) [a]	1.9 (1.0) [a]	2.4 (1.2) [a]	0.3 (0.9) [cde]	0.5 (1.2) [ce]	0.8 (1.5) [ce]

Table 4. Mean change in log$_{10}$ titer of influenza virus in respiratory tract tissues from rhesus macaques and California sea lions. Means for the same time frame followed by a common letter are not significantly different by the Games-Howell test at the 5% level of significance. Values in parentheses show standard deviations. The symbols indicate that the mean change in titer differs significantly for at least one tissue according to Welch's ANOVA at 5% level of significance for: * 1 vs 24 hpi (F=28.071, df=(2, 127.39), p-value=7.93e-11) in rhesus macaques, (F=15.369, df=(2, 269.45), p-value=5.96e-05) for California sea lions, † 1 vs 48 hpi (F=13.641, df=(2, 129.47), p-value=4.02e-6) in rhesus macaques, (F=10.105, df=(2, 257.71), p-value=5.96e-05) in California sea lions, ‡ 1 vs 72 hpi (F=7.8203, df=(2, 133.05), p-value=6.16e-4) for rhesus macaques, (F=11.083, df=(2, 268.89), p-value=2.37e-0.5) for California sea lions.

Tissue	RHESUS MACAQUES			CALIFORNIA SEA LIONS		
	1 vs 24 hpi*	1 vs 48 hpi†	1 vs 72 hpi‡	1 vs 24 hpi*	1 vs 48 hpi†	1 vs 72 hpi‡
Bronchi	1.0 (0.9) ^{a}	1.5 (1.4) ^{a}	1.7 (1.5) ^{a}	0.3 (0.8) ^{a}	0.5 (1.3) ^{a}	0.8 (1.5) ^{a}
Trachea	0.8 (1.0) ^{a}	1.4 (1.5) ^{a}	1.9 (1.5) ^{a}	0.1 (0.7) ^{a}	0.1 (0.8) ^{b}	0.2 (1.0) ^{b}
Lung	0.1 (0.7) ^{b}	0.5 (1.1) ^{b}	1.0 (1.4) ^{b}	-0.2 (0.6) ^{b}	-0.1 (0.8) ^{b}	0.1 (1.1) ^{b}

Table 5. **Difference in mean log$_{10}$ change for respiratory tract explant tissues from California sea lions versus rhesus macaques**. Differences are show as p-values after the Games-Howell post-hoc test. Significant p-values ($p < 0.05$) are in bold. Strain names refer to the original source of the virus isolate followed by the hemagglutinin and neuraminidase subtype. HS is harbor seal. m is mallard. ES is elephant seal. hpi is hours post inoculation.

Timeframe	IAV Strain	Trachea	Bronchi	Lung
1-24 hpi	HS/H3N8	**-1.26 (0.011)**	**-1.13 (0.023)**	-0.11 (0.989)
	m/H3N8	**-1.26 (0.003)**	**-1.40 (<0.001)**	-0.29 (0.695)
	ES/H1N1	-0.03 (1.00)	-0.13 (0.967)	-0.10 (0.996)
	m/H1N1	-0.14 (0.959)	-0.32 (0.475)	-0.03 (1.00)
	m/H5N2	-0.41 (0.362)	-0.45 (0.388)	-0.07 (0.997)
	m/H7N5	**-1.02 (0.047)**	-0.62 (0.266)	-0.46 (0.551)
1-48 hpi	HS/H3N8	**-1.95 (0.002)**	-1.27 (0.120)	-0.18 (0.982)
	m/H3N8	**-2.21 (<0.001)**	**-1.50 (0.009)**	**-1.50 (0.006)**
	ES/H1N1	-0.02 (1.00)	0.25 (0.963)	-0.45 (0.572)
	m/H1N1	-0.74 (0.160)	-0.48 (0.714)	0.01 (1.00)
	m/H5N2	-0.99 (0.100)	**-1.50 (0.024)**	-0.25 (0.918)
	m/H7N5	**-1.75 (<0.001)**	**-1.50 (0.001)**	-0.71 (0.540)
1-72 hpi	HS/H3N8	**-2.41 (<0.001)**	**-1.74 (0.015)**	-0.49 (0.711)
	m/H3N8	**-2.02 (<0.001)**	-1.44 (0.085)	**-1.87 (0.006)**
	ES/H1N1	-0.05 (1.00)	0.01 (1.00)	-0.48 (0.464)
	m/H1N1	-0.83 (0.265)	-1.06 (0.131)	-0.03 (1.00)
	m/H5N2	**-2.11 (<0.001)**	-1.23 (0.083)	-0.32 (0.893)
	m/H7N5	**-2.16 (<0.001)**	-1.08 (0.229)	-1.11 (0.335)

Table 6. **Immunohistochemical detection of influenza A virus in macaque and California sea lion respiratory tissue explants.** Trachea, bronchi and lung were examined for each animal. At least 22 sections from each tissue type were examined for the rhesus macaques and at least 56 sections from each tissue type were examined for the California sea lions. Strain names refer to the original source of the virus isolate followed by the hemagglutinin and neuraminidase subtype. HS is harbor seal. m is mallard. ES is elephant seal. hpi is hours post inoculation.

Animal	IAV positive epithelial cells/mm^2							
	M11		M13		SL08		SL12	
hpi	24	48	24	48	24	48	24	48
HS/H3N8	458	519	71	224	0	0	51	21
m/H3N8	349	544	0	1	2	0	0	26
ES/H1N1	0	0	0	0	0	0	0	0
m/H1N1	0	8	0	0	0	0	0	0
m/H5N2	7	19	0	0	0	0	0	0
m/H7N5	50	626	9	0	0	0	8	<5

Table 7. IAV inoculated rhesus macaque and marine mammal respiratory tract explants from which whole genome IAV sequences were obtained at indicated times post-inoculation and compared to their respective inocula. Empty squares indicate samples for which sequencing was not attempted. HS is harbor seal. m is mallard. ES is elephant seal, M is macaque, SL is California sea lion.

Animal ID	tissue	hour(s) post inoculation					
		HS/H3N8	m/H3N8	ES/H1N1	m/H1N1	m/H5N2	m/H7N5
M15	trachea	72	72		72	72	72
	lung						72
M11	trachea	72	72		72		72
SL01	bronchi		72				72, 96
SL06	bronchi	48, 72	72			72	48, 72
SL08	trachea						48
	bronchi			72			72
ES10	bronchi	48, 72					

Table 8. **IAV amino acid changes compared to inocula detected in *ex vivo* bronchi explants from California sea lions and a Northern elephant seal**. Numbers shown correspond to positions in indicated IAV proteins. *Indicates that the mutation was a synonymous change. Empty squares indicate no sequence differences were detected. An X shows samples for which sequencing was not attempted. hpi is hours post inoculation, HS is harbor seal, m is mallard, ES is elephant seal, M is macaque, SL is California sea lion, HA is hemagglutinin, NA is neuraminidase, PA is polymerase acidic protein, PB is polymerase basic protein.

Animal ID	hpi	HS/H3N8 HA	HS/H3N8 NA	ES/H1N1 PA	m/H7N5 HA	m/H7N5 PB2*
SL01	72	X	X	X	V311I	1368
SL01	96		V394I	X		
SL06	48			X		1368
SL06	72		P118S	X		1368
SL08	48	X	X	X		1368
SL08	72	X	X	I359N, M441I		1368
ES10	48	A154T	P118S	X	X	X
ES10	72	A154T	P118S	X	X	X

IAV titer (PFU/explant)

TRACHEA

HS/H3N8 · m/H3N8 · ES/H1N1 · m/H1N1 · m/H5N2 · m/H7N5

BRONCHI

LUNG

hours post inoculation

PRIMARY BRONCHI

HS/H3N8 · m/H3N8 · ES/H1N1 · m/H1N1 · m/H5N2 · m/H7N5

SECONDARY BRONCHI

LUNG

IAV titer (PFU/explant)

hours post inoculation