Deletion of an intronic polypyrimidine tract of porcine *DNAH17* perturbs splicing and causes defective sperm flagella

Adéla Nosková*,[1], Maya Hiltpold*,[1], Fredi Janett[†], Thomas Echtermann[‡], Zih-Hua Fang*, Xaver Sidler[‡], Christin Selige[§], Andreas Hofer[§], Stefan Neuenschwander**, Hubert Pausch*

*Animal Genomics, Institute of Agricultural Sciences, ETH Zürich, 8315 Lindau, Switzerland

[†]Clinic of Reproductive Medicine, Vetsuisse Faculty, University of Zurich, 8057 Zurich, Switzerland

[‡]Division of Swine Medicine, Vetsuisse Faculty, University of Zurich, 8057 Zurich, Switzerland

[§]SUISAG, 6204 Sempach, Switzerland

**Animal Genetics, Institute of Agricultural Science, ETH Zürich, 8092 Zürich, Switzerland

[1]authors contributed equally

AN: adela.Noskova@usys.ethz.ch
MH: maya.hiltpold@usys.ethz.ch
FJ: fredi.janett@uzh.ch
TE: techtermann@vetclinics.uzh.ch
ZHF: zih-hua.fang@usys.ethz.ch
XS: xsidler@vetclinics.uzh.ch
CS: cse@suisag.ch
AH: aho@suisag.ch
SN: stefan.neuenschwander@usys.ethz.ch
HP: hubert.pausch@usys.ethz.ch

Data availability:

- ***DNA and RNA sequencing*** data are available at the European Nucleotide Archive (ENA) of the EMBL at BioProject
  - PRJEB38156 (https://www.ebi.ac.uk/ena/browser/view/PRJEB38156)
  - PRJEB37956 (https://www.ebi.ac.uk/ena/browser/view/PRJEB37956)
  - PRJEB39374 (https://www.ebi.ac.uk/ena/browser/view/PRJEB39374)
  - PRJNA622908 (https://www.ebi.ac.uk/ena/browser/view/PRJNA622908)
  - PRJNA506525 (https://www.ebi.ac.uk/ena/browser/view/PRJNA506525)
- ***Whole-genome genotypes*** in the form of plink binary files and **phased haplotypes** of 6 affected and 100 unaffected Large White boars are available at zenodo: https://doi.org/10.5281/zenodo.4081475
- **Genotypes of candidate causal variants** for 5 affected and 82 unaffected boars are available in a VCF file at zenodo: https://doi.org/10.5281/zenodo.4081475

46   **Running title**: *DNAH17* exon skipping in infertile boars
47
48   **Key words**: Sperm flagella defect, impaired male fertility, polypyrimidine tract, exon
49   skipping, genetic disorder
50

Corresponding author
Prof. Dr. Hubert Pausch
ETH Zurich
Animal Genomics
Institute of Agricultural Sciences (IAS)
EHB E21
Eschikon 27
8315 Lindau, Switzerland
Phone +41 52 354 92 34
hubert.pausch@usys.ethz.ch
www.ag.ethz.ch

ABSTRACT

Artificial insemination in pig (*Sus scrofa domesticus*) breeding involves the
evaluation of the semen quality of breeding boars. Ejaculates that fulfill predefined
quality requirements are processed, diluted and used for inseminations. Within short
time, eight Swiss Large White boars producing immotile sperm that had multiple
morphological abnormalities of the sperm flagella were noticed. The eight boars
were inbred on a common ancestor suggesting that the novel sperm flagella defect is
a recessive trait. Transmission electron microscopy cross-sections revealed that the
immotile sperm had disorganized flagellar axonemes. Haplotype-based association
testing involving microarray-derived genotypes at 41,094 SNPs of six affected and
100 fertile boars yielded strong association ($P=4.22 \times 10^{-15}$) at chromosome 12.
Autozygosity mapping enabled us to pinpoint the causal mutation on a 1.11 Mb
haplotype located between 3,473,632 and 4,587,759 bp. The haplotype carries an
intronic 13-bp deletion (Chr12:3,556,401-3,556,414 bp) that is compatible with
recessive inheritance. The 13-bp deletion excises the polypyrimidine tract upstream
exon 56 of *DNAH17* (XM_021066525.1:c.8510-17_8510-5del) encoding dynein
axonemal heavy chain 17. Transcriptome analysis of the testis of two affected boars
revealed that the loss of the polypyrimidine tract causes exon skipping which results
in the in-frame loss of 89 amino acids from DNAH17. Disruption of DNAH17 impairs
the assembly of the flagellar axoneme and manifests in multiple morphological
abnormalities of the sperm flagella. Direct gene testing may now be implemented to
monitor the defective allele in the Swiss Large White population and prevent the
frequent manifestation of a sterilizing sperm tail disorder in breeding boars.

INTRODUCTION

Artificial insemination is the most frequent method of breeding in pigs. The semen of
breeding boars is collected once or twice per week at semen collection centers. All
ejaculates are macro- and microscopically evaluated. Traits that are routinely
measured in all ejaculates include ejaculate volume, sperm concentration, motility,
and morphology. Only ejaculates that meet the quality requirements for artificial
insemination are processed, diluted and used for inseminations (Colenbrander et al.,
1993),(Broekhuijse et al., 2011),(Holt et al., 1997). Semen quality and insemination

success vary within and between boars due to environmental, permanent

environmental and genetic effects (Marques et al., 2017).

Access to longitudinal data on standardized semen traits for large cohorts of males is

unique to livestock populations (Thibier and Wagner, 2002),(Knox, 2016). Boars

suitable for breeding are selected based on semen quality and genomic predictions

derived from dense SNP microarray-derived genotypes. Dense genotypes and

repeated semen trait measurements for thousands of males are prerequisites to

unravel the genetic architecture of male fertility. The comprehensive genetic analysis

of male fertility is challenging in most species other than livestock. For instance, the

analysis of male fertility in humans typically relies on correlated proxy phenotypes

such as family size and birth rate (Kosova et al., 2012), small cohorts of males (Sato

et al., 2018),(Sato et al., 2020), and observations from a single ejaculate per

individual (Rahban et al., 2019).

Due to the widespread use of artificial insemination in livestock populations, the

semen from individual male breeding animals may be used for thousands of

inseminations. Boar semen quality is positively correlated with insemination success

and litter size (Holt et al., 1997). Semen traits that are measured at semen collection

centers have medium heritability (Marques et al., 2017). Thus, they are well suited

response variables in genome-wide association studies for male reproductive

performance (Diniz et al., 2014),(Hiltpold et al., 2020). Genome-wide association

studies on male fertility carried out in livestock discovered novel phenotype-genotype

associations that improved our biological understanding of mammalian fertilization

(Pausch et al., 2014),(Lamas-Toranzo et al., 2020, p. 95),(Noda et al.,

2020),(Hiltpold et al., 2020),(Ni et al., 2020, p. 19),(Noskova et al., 2020).

Ejaculates from hundreds of boars are evaluated every year at semen collection

centers as a service to the pig breeding industry. This rich resource of semen quality

records facilitates to investigate sporadically occurring sperm defects using case-

control association testing (Sironen et al., 2006),(Sironen et al., 2011),(Noskova et

al., 2020). Once causative variants have been identified, direct gene tests and

genome-based mating strategies may be implemented to avoid the birth of infertile

males. Moreover, the discovery of genes that harbor pathogenic alleles that

compromise male fertility is important to enhance the diagnostic yield of genetic

testing also in species other than livestock (Xavier et al., 2020).

Here, we investigate an autosomal recessive sperm tail defect of Swiss Large White

boars. Using genome-wide association testing, we map the disorder to porcine

chromosome 12. The analysis of genome-wide DNA and RNA sequencing data of

affected boars revealed that the loss of an intronic polypyrimidine tract of porcine

DNAH17 is causal for the morphological abnormalities of the sperm flagella.

MATERIAL AND METHODS

Ethics approval and consent to participate

Semen samples of five breeding boars were collected by laboratory technicians at an

approved semen collection center as part of their regular service to the Swiss pig

breeding industry. Semen samples of three boars were flushed from the epididymis

post mortem. Testes were collected after regular slaughter at an approved

slaughterhouse. Our study was approved by the veterinary office of the Canton of

Zurich (animal experimentation permit ZH 070/20).

Consent for publication

SUISAG, the Swiss pig breeding and competence center provided written consent to

the analyses performed and agreed to publish results and data.

Animals

Eight Swiss Large White boars with a sperm tail defect were considered in our study

(**Table 1**). Five of them were noticed at the semen collection center of SUISAG

because their ejaculates contained immotile spermatozoa that had multiple

morphological abnormalities of the flagella. Because the five boars were healthy and

pedigree analysis indicated that they were inbred on a common ancestor, recessive

inheritance of the sperm tail defect was suspected. A mating between two suspected

carrier animals was performed in the field to assess phenotypic manifestations in

their offspring. The pregnant sow was purchased and maintained at the research

barn of the Division of Swine Medicine, Vetsuisse Faculty, University of Zurich. The

sow gave birth to a litter with eleven piglets (eight females, three males). One of the

male piglets (Boar_1246) died at the age of 200 days due to hemorrhagic bowel

syndrome (Grahofer et al., 2017). The other two male boars (Boar_1249,

Boar_1254) were slaughtered at the age of 17 months at a regular slaughterhouse.

All three male piglets from the mating of suspected carrier animals expressed the
sperm tail defect (see below). Pedigree records were analyzed using the PYPEDAL
software package (Cole, 2007).

Table 1: Experimental design of the study

Boar	AI center	Research station	GWAS cohort	PCR-based genotyping	Whole-genome sequence data	RNA sequencing data	Macro- and microscopic sperm analyses	Eosin-nigrosin staining	TEM sections
Boar_78	X		X		SAMEA6813549		X		
Boar_79	X		X		SAMEA6813550		X		
Boar_80	X		X	X	SAMEA6813551	SAMEA6832284	X		
Boar_81	X		X		SAMEA6813552	SAMEA6832285	X		
Boar_82	X		X		SAMEA6813553		X		
Boar_1246[a]		X	X				X		
Boar_1249		X		X			X	X	X
Boar_1254		X		X			X	X	X

[a] died at the age of 200 days due to hemorrhagic bowel syndrome; sperm were flushed from the
epididymis post mortem.
Details on the eight boars with a sperm tail defect. Genotypes and haplotypes for the GWAS cohort
are available at https://doi.org/10.5281/zenodo.4081475. DNA and RNA sequencing data from the
boars are available at the European Nucleotide Archive (ENA) of the EMBL at the BioProject
PRJEB38156 under the accession numbers listed in the table.

Phenotypes
Semen samples of five breeding boars were macro- and microscopically evaluated
by laboratory technicians as part of the routine service of SUISAG to the Swiss pig
breeding industry. Spermatozoa from three boars (Boar_1246, Boar_1249,
Boar_1254) were flushed from the epididymis post mortem. Semen samples from a
fertile boar were provided by SUISAG.

Sperm concentration, total sperm count and sperm motility were determined with an
IVOS II CASA system (Hamilton Thorne Inc., Beverly, U.S.A.) using Leja 2-chamber
slides (Leja, Nieuw-Vennep, the Netherlands). For morphological examination,
semen was fixed in buffered formol saline solution (Na2HPO4 4.93g, KH2PO4
2.54g, 38% formaldehyde 125ml, NaCl 5.41g, distilled water q.s. 1000ml) and
smears prepared (Hancock, 1957). At least 200 spermatozoa were subsequently

evaluated by phase contrast microscopy using oil immersion (Olympus BX50,

UplanF1 100×/1.30, Olympus, Wallisellen, Switzerland).

Assessment of sperm viability was performed using the eosin-nigrosin staining

method (Blom, 1950). In stained slides at least 200 spermatozoa were evaluated

under oil immersion on a light microscope (Olympus BX50, UplanF1 100×/1.30,

Olympus, Wallisellen).

In order to prepare sperm flushed from the epididymis for transmission electron

microscopy (TEM), the samples were washed twice in phosphate buffered saline

(PBS) and subsequently centrifuged at 300g. A semen sample from an unaffected

boar was first centrifuged at 300g to increase the sperm concentration and then

washed in PBS. The pellet was fixed with equal volume of 6% glutaraldehyde,

resuspended gently, and centrifuged at 6000g. After removing the supernatant, the

sperm were fixed for a second time with 3% glutaraldehyde, and finally pelleted at

6000g. Pellets were washed three times in PBS, post-fixed in 1% osmium tetroxide,

washed in ddH2O, stained in 1% uranyl acetate, dehydrated in graded series of

ethanol (25, 50, 75, 90, and 100%), and embedded in Epoxy resin through

increasing concentrations (25, 50, 75, and 100%) using PELCO Biowave+ tissue

processor, and then cured at 60 °C for 3 days. Embedded blocks were sectioned

using Leica FC6 microtome and a DIATOME diamond knife with 45° angle into

60 nm sections and mounted on Quantifoil copper grids with formvar carbon films.

Sections were post-stained with 2% uranyl acetate followed by lead citrate. Grids

were imaged using FEI Morgagni 268 electron microscope operated at 100 kV at

20 k magnification.

Genotypes and haplotype inference

Genotypes of 9955 Large White boars (including six boars with the sperm tail defect)

and sows were provided by SUISAG. All pigs were genotyped using Illumina

PorcineSNP60 Bead chips that comprised between 62,163 and 68,528 SNPs. We

considered only autosomal SNPs. Physical positions of the SNPs corresponded to

the Sscrofa11.1-assembly of the porcine genome (Warr et al., 2019). Quality control

on the genotypes was carried using the PLINK (version 1.9) software (Chang et al.,

2015). Animals and SNPs with more than 10% missing genotypes were excluded

from subsequent analyses. We removed SNPs with minor allele frequency (MAF)

less than 0.005 and SNPs for which the observed genotype distribution deviated

significantly (P < 0.00001) from Hardy-Weinberg proportions. After quality control,

our dataset contained 9848 pigs and genotypes at 43,254 autosomal SNPs.

Sporadically missing genotypes were imputed and haplotypes were inferred using 12

iterations of the phasing algorithm implemented in the BEAGLE (version 5.0) software

(Browning et al., 2018), assuming an effective population size of 500 while all other

parameters were set to default values.

Genome-wide association testing

Six boars that produced sperm with multiple morphological abnormalities of the

flagella were considered as case group for a genome-wide association study. The

control group consisted of 100 randomly selected boars that produced normal sperm

and were fertile, i.e., each of them sired at least one litter in the Swiss Large White

breeding unit.

We tested the association between the affection status of the boars and 41,094

autosomal SNPs for which the frequency of the minor allele was greater than 5%.

Single marker-based case-control association testing was performed using Fisher's

exact tests of allelic association as implemented in the PLINK (version 1.9) software

(Chang et al., 2015).

A linear haplotype-based association model was implemented as described in

(Venhoranta et al., 2014) and (Hiltpold et al., 2020). We shifted a sliding window

consisting of 25 contiguous SNPs corresponding to an average haplotype length of

1.28 ± 0.65 Mb along the chromosomes in steps of 5 SNPs. Within each of the 8583

sliding windows considered, we tested the association of between 1 and 10

haplotypes that had frequency greater than 5% with the sperm tail defect using the

linear model (Hiltpold et al., 2020) $y = \mu + \sum_{j=1}^{30} a_j PC_j + bHT + e$, where y is a vector

of phenotypes (1-fertile, 2-infertile), μ is the intercept, PC_j are the top 30 principal

components of a genomic relationship matrix built based on genotypes of 43,254

autosomal SNPs using the PLINK software, a and b are effects of the principal

components and the haplotype (HT) tested, respectively, and e is a vector of

residuals that are assumed to be normally distributed. In total, we tested 47,685

haplotypes for association with the sperm tail defect.

SNPs and haplotypes that exceeded the Bonferroni-corrected significance threshold

($P_{SNP} = 1.22 \times 10^{-6}$, $P_{HAPLO} = 1.05 \times 10^{-6}$) were considered as significantly associated.

The inflation factors of the test statistics were calculated using the *estlambda()*-

function of the GENABEL R package (Aulchenko et al., 2007).

Analysis of positional candidate genes

The top association signal revealed a 1.11 Mb segment of extended homozygosity at

chromosome 12. Genes located within the segment of homozygosity were obtained

from the Refseq annotation of the porcine genome (version 106, available at

ftp://ftp.ncbi.nlm.nih.gov/genomes/refseq/vertebrate_mammalian/Sus_scrofa/annotat

ion_releases/106/GCF_000003025.6_Sscrofa11.1/GCF_000003025.6_Sscrofa11.1_

genomic.gff.gz). Using publicly available RNA sequencing data (BioProject

PRJNA506525, sample accession numbers SAMN10462191 – SAMN10462197)

from (Robic et al., 2019), we quantified mRNA abundance of the positional candidate

genes in porcine testes. To this end, we downloaded between 47 and 58 million 2 x

150 bp paired-end sequencing reads that were generated from RNA extracted from

testis tissue of seven pubertal boars from European Piétrain and Landrace

populations. The RNA sequencing reads were pseudo-aligned to an index of the

porcine transcriptome (Refseq version 106) and transcript abundance was quantified

using the KALLISTO software (Bray et al., 2016). We used the R package TXIMPORT

(Soneson et al., 2015) to aggregate transcript abundances to the gene level.

Whole-genome sequencing and sequence variant genotyping

We sequenced five Swiss Large White boars that produced sperm that had multiple

morphological abnormalities of the sperm flagella using 2×150 bp paired-end reads.

Illumina TruSeq DNA PCR-free libraries with 400 bp insert sizes were prepared and

sequenced at an Illumina NovaSeq6000 instrument. We used the FASTP software

(Chen et al., 2018) to remove adapter sequences, poly-G tails and reads that had

phred-scaled quality less than 15 for more than 15% of the bases. Between

81,145,807 and 114,616,836 filtered read pairs were aligned to the Sscrofa11.1

assembly of the porcine genome using the MEM algorithm of the BWA software (Li,

2013) with default parameter settings. We marked duplicates using the PICARD tools

software suite (https://github.com/broadinstitute/picard) and sorted the alignments by

coordinates using SAMBAMBA (Tarasov et al., 2015). The number of reads covering a

position were extracted from the alignments using the MOSDEPTH software (version

0.2.2) with default parameter settings (Pedersen and Quinlan, 2018). The realized

sequencing read depth of the five boars was 10.9-fold and it ranged from 8.9- to

12.5-fold.

Sequence variants (SNPs and Indels) of the five sequenced boars were genotyped

together with 93 pigs from different breeds for which whole-genome sequence data

were available from our in-house database using the multi-sample variant calling

approach of the GENOME ANALYSIS TOOLKIT (GATK, version 4.1.0; (Depristo et al.,

2011)). We filtered the sequence variants by hard-filtering according to best practice

guidelines of the GATK. A detailed description of our reference-guided variant

discovery and filtration approach is described in (Crysnanto et al., 2019). Functional

consequences of polymorphic sites were predicted according to the Refseq

annotation (version 106) of the porcine genome using the VARIANT EFFECT PREDICTOR

software from Ensembl (McLaren et al., 2016) along with the SpliceRegion.pm plugin

(https://github.com/Ensembl/VEP_plugins/blob/release/101/SpliceRegion.pm). We

considered 82 pigs with known pedigree as a control group for the present study.

Identification of candidate causal variants

We considered 14,806 SNPs and 3,856 Indels that were detected within the 1.11 Mb

segment (between 3,473,632 and 4,587,759 bp) of extended homozygosity at

chromosome 12 as positional candidate causal variants. To identify variants

compatible with recessive inheritance of the sperm tail disorder, we applied a filtering

strategy that takes into account flawed genotypes and the under-calling of

heterozygous genotypes due to relatively low sequencing coverage. Specifically, we

screened for alleles that had following frequency:

- ≥ 0.8 in five affected boars (at least 8 out of 10 alleles),

- ≤ 0.05 in 82 control pigs from different breeds (less than 8 alleles).

This filtration resulted in only one compatible variant which was a 13-bp intronic

deletion.

Whole transcriptome sequencing and read alignment

Testes were collected immediately after slaughter at an approved slaughterhouse

from two boars (Boar_80, Boar_81) that were homozygous for the 13-bp deletion

and produced sperm with multiple morphological abnormalities of the flagella. Tissue

samples were frozen in liquid nitrogen and stored at -80°C until RNA extraction.

Paired-end RNA libraries (2 x 150 bp) were prepared using total RNA purified from

testis tissue using the Illumina TruSeq RNA Sample Preparation Kit (Illumina, San
Diego, CA, USA). The libraries were sequenced at an Illumina NovaSeq6000
instrument yielding 49,198,010 and 70,451,148 reads. Quality control on the raw
RNA sequencing reads was performed using the FASTP software using default
parameter settings (see above). The filtered read pairs (48,889,296 and 69,988,376)
were aligned to the Sscrofa11.1 reference sequence and the Refseq gene
annotation (version 106) using the splice-aware read alignment tool STAR (version
2.7.3a) (Dobin et al., 2013). The number of RNA sequencing reads that covered a
position was counted using the MOSDEPTH (see above).

Bioinformatic analysis

Putative branch point sequences within *DNAH17* (XM_021066525.1) intron 55 were
predicted using the BPP algorithm (Zhang et al., 2017). Putative 3' splice acceptor
sites were predicted using the NNSPLICE software tool
(https://www.fruitfly.org/seq_tools/splice.html (Reese et al., 1997)). Domains of
porcine DNAH17 (protein-ID: A0A287AFU3) were retrieved from Uniprot.

Genotyping of the 13-bp deletion

DNA was isolated from either hair roots (living pigs) or spleen (slaughtered pigs)
using DNeasy blood and tissue kit (Qiagen). A PCR test was established consisting
of a FAM-labelled forward primer (Fw-*DNAH17*: 5'-TGAGCATCTTCTTGGCGAGG-
3') and a reverse primer (Re-*DNAH17*: 5'-GCTGTTGATCAGCACCAGGA-3') which
bind to the wild type and mutant alleles, as well as a wild type specific reverse primer
(Re-wt-*DNAH17*: 5'-TCGAGCTAGACGCGAGGG-3'). PCR fragments were analyzed
on a DNA analyzer 3130xl (Applied Bioscience).

Availability of data

Whole-genome sequence data of 87 pigs including five boars with a sperm tail defect
have been deposited at the European Nucleotide Archive (ENA) of the EMBL at
BioProject PRJEB38156, PRJEB37956, PRJEB39374 and PRJNA622908, under
sample accession numbers listed in **Supplemental file S1**. The genotypes of five
affected and 82 unaffected boars at 14,806 SNPs and 3,856 Indels that were
detected within the segment of autozygosity are available from Zenodo
(https://doi.org/10.5281/zenodo.4081475). RNA sequencing data of testicular tissue
samples of two boars homozygous for the 13-bp deletion have been deposited at the

European Nucleotide Archive (ENA) of the EMBL at the BioProject PRJEB38156

under sample accession numbers SAMEA6832284 and SAMEA6832285. RNA

sequencing data of testicular tissue samples of seven pubertal boars are accessible

via the European Nucleotide Archive (ENA) of the EMBL at the BioProject

PRJNA506525 under sample accession numbers SAMN10462191 –

SAMN10462197. The microarray-derived genotypes and haplotypes of cases and

controls, the eigenvectors considered to account for stratification, the R script use to

carry out haplotype-based association testing, and the results from the haplotype-

based association study are available from Zenodo

(https://doi.org/10.5281/zenodo.4081475).

357   **RESULTS**

Phenotypic manifestation of a sperm tail disorder in Large White boars

Five Swiss Large White boars maintained at a semen collection center produced

ejaculates that contained spermatozoa with defective tails. Microscopic semen

analysis revealed multiple morphological abnormalities of the sperm flagella

including rudimentary, short, coiled and irregularly shaped tails (**Figure 1A**).

Proximal cytoplasmic droplets were frequently observed at the junction of sperm

head and tail. Progressively motile sperm were not detected in the ejaculates. Eosin-

nigrosin staining of semen flushed from the epididymis of two affected boars

indicated that 48 and 60% of the sperm were viable (**Supplemental file S2**).

Ejaculate volume and sperm concentration were normal. Apart from producing

sperm with defective tails, the boars were healthy and had no testicular

abnormalities. Due to the absence of motile sperm and the high degree of

abnormalities of the flagella, the boars were not suitable for breeding and were

slaughtered. The sperm tail defect of the boars is commonly referred to as

asthenoteratozoospermia.

Transmission electron microscopy cross-sections of sperm flagella from a fertile boar

revealed the typical axonemal arrangement of nine outer microtubule doublets

surrounding the central pair (9x2 + 2) (**Figure 1D**). In contrast, cross-sections of

sperm that were flushed from the epididymis of two affected boars revealed multiple

ultrastructural abnormalities including completely disorganized axonemes (**Figure**

1C). Central and outer microtubule doublets were absent in some flagella. Some sperm had superfluous but disorganized axonemal structures. The cytoplasmic bags that were detected at the junction between sperm head and tail using light microscopy contained unassembled axonemal components (**Supplemental file S3**).

Figure 1. Characteristics of a recessive sperm tail defect of Large White boars. Representative phase contrast microscopy images of sperm produced from affected (A) and fertile (B) boars. Ejaculates of affected boars contained sperm with multiple morphological abnormalities of the flagella including short, coiled, broken and bent tails. A proximal cytoplasmic droplet was detected between the sperm head and tail at most of the spermatozoa from affected boars. Representative transmission electron microscopy cross-sections of sperm flagella from affected (C) and fertile (D) boars. Cross sections of the sperm flagella revealed the typical arrangement of nine outer microtubule doublets surrounding the central pair (9×2 + 2) in fertile boars (D). Lack of central and outer microtubules doublets and complete axonemal disorganization were frequently detected in cross-sections of the sperm flagella in affected boars. Pedigree of eight boars that produced sperm with defective tails (D). The pedigree contains only obligate mutation carriers. Red color indicates the eight affected boars. The numbers within the boxes correspond to the IDs in **Table 1**. Boars 1246, 1249 and 1254 originate from a mating between two suspected carrier animals. Ovals and boxes represent female and male ancestors, respectively. The grey box indicates a common ancestor (born in 2005) on which all affected boars were inbred. Shapes filled with yellow color represent individuals that were confirmed to carry the top haplotype in heterozygous state using Illumina Porcine60K genotypes.

The boars that produced defective sperm were closely related (**Figure 1E**). Two
affected boars were fullsibs. The average relationship coefficient between the boars
was 0.28 and it ranged from 0.14 to 0.63. Their average coefficient of inbreeding was
0.087, which is slightly higher than in the fertile boars (F=0.06). All affected boars
were inbred on a boar born in 2005. This common ancestor was present in both the
paternal and maternal ancestry of significantly less (N=10, $P_{Fisher's\ exact}$=4.69 x 10^{-6})
fertile boars. Three sires of affected boars were used in artificial insemination. None
of their ejaculates (n=168) contained an anomalous amount of sperm with defective
flagella. These observations are compatible with an autosomal recessive inheritance
of the sperm tail defect.

In order to verify the preseumed recessive inheritance and monitor phenotypic
manifestations in homozygous boars, a mating between two suspected carrier
animals was performed in the field (**Figure 1E**). The sow gave birth to a litter with
eight female and three male piglets (Boar_1246, Boar_1249, Boar_1254). The three
male piglets were maintained at a research barn. Boar_1246 died at the age of 200
days due to hemorrhagic bowel syndrome. Testes and epididymis were collected
post mortem. Testicular abnormalities were not detected. Microscopic analysis of
sperm flushed from the epididymis revealed that the sperm flagella had multiple
morphological abnormalities. Boar_1249 and Boar_1254 were healthy and
developed normal at the research barn. At 17 months, both boars were slaughtered
at a regular slaughterhouse and testes and epididymis were collected. Microscopic
analysis of sperm flushed from the epididymis revealed that both boars produced
sperm with multiple morphological abnormalities of the flagella.

Haplotype-based association testing maps the sperm tail defect to a 1.11 Mb
interval on porcine chromosome 12
Six affected boars (**Table 1**) were genotyped with the Illumina PorcineSNP60
microarray. As a control group, we considered 100 randomly selected fertile boars of
the Swiss Large White breeding population that also had Illumina PorcineSNP60-
derived genotypes. Following quality control, genotypes at 41,094 autosomal SNPs
were used for association testing. Single marker-based case/control-association
testing using Fisher's exact test of allelic association revealed eleven SNPs located
on porcine chromosomes 11, 12 and 13 that exceeded the Bonferroni-corrected
significance threshold (**Figure 2A**). The strongest association signal (P=1.87 x 10^{-9})

resulted from SNP *ASGA0052524* that is located at 2,785,333 bp on porcine

chromosome 12. An inflation factor of 1.55 indicated that the SNP-based association

study was enriched for false positive association signals. Inspection of the top

principal components of the genomic relationship matrix revealed clustering of the

boars with the sperm tail defect (**Supplemental file S4**), suggesting that population

stratification confounded the association analysis, thereby producing spurious

association signals.

Figure 2. A 1.11 Mb segment on porcine chromosome 12 is associated with the sperm tail

defect. Manhattan plot of a SNP- (A) and haplotype-based (B) genome-wide association study. Red

color represents eleven SNPs and twenty-one haplotypes that exceeded the Bonferroni corrected-

significance threshold. Homozygosity mapping in six boars that produced sperm with defective tails

using SNPs located on porcine chromosome 12 (C). Blue and pale blue represent homozygous

genotypes (AA and BB), heterozygous genotypes are displayed in black. The red bar indicates a 1.11

Mb segment of extended homozygosity that was shared in all six affected boars. mRNA abundance

(quantified in transcripts per million) of 15 genes that are located within the segment of extended

homozygosity in testicular tissue of pubertal boars (D). The vertical line represents the standard

deviation estimated across seven samples.

To take population stratification into account, we repeated the association analysis

using a haplotype-based linear model that included 30 principal components of a

genomic relationship matrix as covariates. An inflation factor of 1.15 indicated that

the principal components-based control of population stratification was mostly

successful. Twenty-one haplotypes located between 94,345 and 9,256,260 bp on

chromosome 12 (SSC12) exceeded the Bonferroni-corrected significance threshold

(**Figure 2B**). The strongest associations (P=4.22 x 10^{-15}) were detected for three

overlapping haplotypes located between 2,166,472 and 3,755,956 bp. The top

haplotype occurred at a frequency of 3.5% among the 100 fertile boars. Seven fertile

boars were heterozygous carriers of the top haplotype. The top haplotype was not

detected in the homozygous state in fertile boars. No haplotypes located on

chromosomes other than SSC12 were associated with the sperm tail defect.

The haplotype-based association study peaked at a 1.11 Mb segment of extended

homozygosity (between 3,473,632 and 4,587,759 bp) at porcine chromosome 12

that was identical by descent in all six affected boars, corroborating recessive

inheritance (**Figure 2C**). According to the Refseq annotation of the porcine genome,

the segment of extended autozygosity encompasses 15 genes (**Figure 2D**).

We suspected that the mutation causing the sperm tail defect affects a transcript that

is abundant in the testes. We found expression levels greater than 10 transcripts per

million for four genes (*DNAH17*, *BIRC5*, *SYNGR2*, *SEC14L1*) that were annotated

within the 1.11 Mb segment of extended homozygosity (**Figure 2D**). Among those,

only *DNAH17* encoding the dynein axonemal heavy chain 17, shows a highly testis-

biased expression in humans (https://gtexportal.org/home/), sheep

(http://biogps.org/dataset/BDS_00015/sheep-atlas/) (Clark et al., 2017) and cattle

(http://cattlegeneatlas.roslin.ed.ac.uk/) (L. Fang et al., 2020). Pathogenic alleles in

human and murine orthologs of *DNAH17* have been detected in males that produce

immotile sperm with multiple morphological abnormalities of the flagella (Whitfield et

al., 2019),(Zhang et al., 2020). Thus, we considered *DNAH17* as a compelling

positional and functional candidate gene for a sperm flagella defect in boars.

An intronic 13-bp deletion is associated with the sperm tail defect

We sequenced five affected boars to an average coverage of 10.9-fold using short

paired-end reads. We also considered 82 sequenced pigs from our in-house variant

database that were not affected by the sperm tail disorder. The analysis of

sequencing depth at the proximal region of SSC12 did not reveal large deletions or

duplications that segregate with the haplotype. To identify candidate causal

mutations, we considered 18,662 variants (14,806 SNPs and 3,856 Indels) that were

detected within the 1.11 Mb interval (between 3,473,632 and 4,587,759 bp) of

extended homozygosity. Of the 82 sequenced control animals, 65 also had array-

called genotypes. Three of the 65 pigs that had array- and sequence-called

genotypes carried the top haplotype in heterozygous state, yielding a haplotype

frequency of 2% in the sequenced control animals.

Of 18,662 positional candidate causal variants, a single variant was compatible with

recessive inheritance of the sperm tail defect: a 13-bp deletion at Chr12:3,556,401-

3,556,414 bp (XM_021066525.1:c.8510-17_8510-5del) located in an intron of

DNAH17. The 13-bp location resides within the top window from the haplotype-

based association study. Sequence variant genotyping using *GATK* indicated that

the five sequenced boars with the sperm tail defect were homozygous for the

deletion. Only the three boars from the control group, that carried the top haplotype

in the heterozygous state, also carried the 13-bp deletion in the heterozygous state.

The deletion did not occur in the homozygous state in animals from the control

group. According to the variant classification algorithm of the VEP software, the

intronic 13-bp deletion was annotated as "splice_region_variant & intron_variant". Its

impact on protein function was predicted to be low.

RNA sequencing reveals that the 13-bp deletion causes skipping of DNAH17
exon 56

Closer inspection of the 13-bp deletion revealed that it excises an intronic pyrimidine-

rich sequence containing a continuous stretch of eight pyrimidine bases between 5

and 17 bases upstream of a canonical 3' splice site of *DNAH17* exon 56 (**Figure**

3A). The VEP plugin SpliceRegion.pm indicated that the deletion coincides with a

putative polypyrimidine tract. A branch point consensus sequence (yUnAy) (Gao et

al., 2008) was predicted (z-score: 4.79) 2-6 nucleotides upstream the 13-bp deletion.

The pyrimidine content between the predicted branch point adenine and the splice

acceptor site at the 3' end of intron 55 is 68.4%. Considering that the polypyrimidine

tract is an important cis-acting element for spliceosome assembly in canonical "GT-

AG"-type acceptor splice sites (Coolidge et al., 1997), we suspected that the 13-bp

deletion perturbs pre-mRNA splicing. *In silico* splice site prediction using NNSPLICE

indicated (prediction score 0.46) that the 13-bp deletion likely prevents recognition of

the 3' splice acceptor site at the intron-exon boundary of exon 56, thus possibly

causing exon skipping.

Figure 3: A 13-bp deletion perturbs splicing of *DNAH17*. Genomic sequence at the intron-exon

boundaries of exon 56 of porcine *DNAH17* (XM_021066525.1) (A). The grey background indicates

the sequence of exon 56. Red color highlights 13 nucleotides that are deleted in boars with the sperm

tail defect. Splice acceptor and donor sites are represented in bold-italic type. Violet color indicates

the predicted (z-score: 4.79) branch point sequence (BPS). Expression of DNAH17 (in counts per

million) in testes of two Swiss Large White boars (SAMEA6832284, SAMEA6832285) homozygous

for the 13-bp deletion (B). The number of reads covering a position was divided by the total number of

reads (in million) mapped to transcripts, and averaged across both boars. Blue color indicates the

exon-intron structure of porcine *DNAH17*. Exon 56 is displayed in red color. DNAH17 domains

according to Uniprot (protein-ID: A0A287AFU3) (C). Different colored boxes represent conserved

domains in the stem of dynein heavy chains (DHC), the microtubule-binding region (MT) and six

ATPase domains (ATPases associated with diverse cellular activities, AAA1-AAA6). Skipping of exon

56 truncates 89 amino acids (red color) from AAA4.

Exon 56 of porcine *DNAH17* is regularly expressed in testis tissue of pubertal boars

(**Supplemental file S5**). In order to examine if the loss of the polypyrimidine tract

upstream exon 56 perturbs splicing, we sequenced RNA purified from testis tissue of

two boars that produced defective sperm and were homozygous for the 13-bp

deletion. We aligned 49 and 70 million paired-end (2x150 nt) RNA sequencing reads

to the porcine transcriptome to quantify transcript and exon abundance. An average

of 49 *DNAH17* transcripts per million (TPM) were detected in the two boars

homozygous for the 13-bp deletion. However, no reads aligned to the sequence of

exon 56 corroborating that the 13-bp deletion prevents recognition of the 3' splice

site, thereby causing exon skipping (**Figure 3B**).

The skipping of exon 56 causes an in-frame loss of 89 amino acids (residues 2836-

2924) representing 2% of porcine DNAH17 (XP_020922184.1). DNAH17 contains

six evolutionarily conserved "ATPase associated with a variety of cellular activities

(AAA)"-domains that are linked together as an asymmetric hexameric ring required

to power the beating movement of sperm flagella (Gleave et al., 2014),(Snider et al.,

2008). The 89 amino acids are truncated from AAA4 of DNAH17 (**Figure 3C**).

553   **DISCUSSION**

Eight Swiss Large White boars produced sperm with multiple morphological

abnormalities of the flagella. Considering that all sperm were immotile, the boars

were judged to be infertile and not used for breeding. Because a majority of the

sperm were viable, fertilization might be possible using intracytoplasmic sperm

injection (ICSI) (Nijs et al., 1996),(Ortega et al., 2011). However, success rates after

ICSI may vary considerably in individuals with asthenoteratozoospermia

(Wambergue et al., 2016),(Whitfield et al., 2019). In any case, the application of ICSI

in livestock populations seems unwarranted if the underlying male-factor infertility

follows recessive inheritance, because a defective allele would be transmitted to the

offspring.

Apart from producing sperm with defective tails, the boars were healthy. Diseases

and environmental stressors were less likely to be causal for the defective sperm

flagella (Jurewicz et al., 2009), because the boars were maintained at a semen

collection center with many other boars that produced normal semen. Idiopathic

infertility and poor semen quality in healthy males may result from pathogenic alleles

in genes that are specifically expressed in the male reproductive tract (Pausch et al.,

2014),(Iso-Touru et al., 2019),(Pausch et al., 2016b),(Noskova et al., 2020). The

eight affected boars were inbred on a common ancestor supporting the hypothesis of

a shared genetic etiology. While the presence of common ancestors in the pedigrees

of all affected boars is compatible with recessive inheritance, it does not rule out

other modes of inheritance (Bourneuf et al., 2017). Considering that all affected but

only few fertile boars were inbred on the common ancestor, a recessive mode of

inheritance was likely.

The morphological and ultrastructural flagella defects of the eight Swiss Large White

boars are similar to those observed in Finnish Yorkshire boars that are homozygous

for a loss-of-function allele in *KPL2* (a.k.a. *SPEF2*) (Andersson et al., 2000),(Sironen

et al., 2006). Although cross-breeding between Finnish Yorkshire and Swiss Large

White pigs had not been documented recently, the defective allele could be located

on an ancient shared haplotype (Pausch et al., 2016a),(Schwarzenbacher et al.,

2016). However, the genomic region on chromosome 16 encompassing *SPEF2* is

not associated with the sperm defect of the Swiss Large White boars, indicating

genetic heterogeneity.

SNPs on chromosomes 11, 12 and 13 were significantly associated with the sperm

disorder using Fisher's exact test of allelic association. This result was puzzling

because an oligogenic inheritance of the sperm flagella defect was unlikely.

Phenotypic misclassification and genetic heterogeneity could lead to an inconclusive

association study (Manchia et al., 2013). However, the morphological abnormalities

of the sperm flagella were strikingly similar in the ejaculates of eight boars.

Moreover, pedigree analysis suggested monogenic recessive inheritance. Principal

components analysis and a genomic inflation factor of 1.55 indicated that population

stratification confounded our SNP-based association study (Price et al., 2010).

Although Fisher's exact tests have been widely applied to map binary traits (Balding,

2006), these tests are prone to type I errors when cases and controls differ in their

ancestries. Following the approach from (Hiltpold et al., 2020), we fitted the top

principal components of a genomic relationship matrix as covariates in a haplotype-

based association model to take population stratification into account. A strong

association signal at chromosome 12 remained while all other association signals

disappeared. In agreement with previous studies (Price et al., 2006),(Pausch et al.,

2011),(Hiltpold et al., 2020) and evidenced by a low inflation factor, the principal

components-based correction successfully eliminated spurious signals that arose

due to population stratification. Considering that the most significantly associated

haplotype also encompasses the causal variant for the sperm flagella defect, our

findings corroborate that genome-wide haplotype-based association testing offers a

powerful approach to identify trait-associated regions in stratified mapping cohorts

(Kadri et al., 2014),(Hiltpold et al., 2020),(Pausch et al., 2016a).

The haplotype associated with multiple morphological abnormalities of the sperm

flagella carries a 13-bp deletion in an intron of *DNAH17* encoding dynein axonemal

heavy chain 17. Our results show that DNAH17 is abundant in porcine testis tissue.

Dynein axonemal heavy chains are required for the axonemal assembly and the

beating movement of the flagella (Inaba, 2003),(Ben Khelifa et al., 2014),(Tu et al.,

2019),(Sironen et al., 2020),(Liu et al., 2020),(Whitfield et al., 2019),(Zhang et al.,

2020, p. 17),(Sha et al., 2020). Pathogenic alleles in human and murine *DNAH17*

manifest in sperm with an abnormal mitochondrial sheath and cytoplasmic droplets,

as well as axonemal disorganization including absence of the central pair and

missing peripheral microtubule doublets (Whitfield et al., 2019), (Zhang et al., 2020,

p. 17),(Sha et al., 2020). Boars that are homozygous for a 13-bp deletion in intron 55

of *DNAH17* also produce sperm with multiple morphological and ultrastructural

abnormalities of the flagella. Although annotated as a low impact variant, the intronic

13-bp deletion perturbs pre-mRNA splicing because it disrupts the polypyrimidine

tract in intron 55 of *DNAH17*. The mutant DNAH17 lacks part of the AAA4 domain

which likely compromises axonemal assembly and the beating movement of the

sperm flagellum in the homozygous state. The assembly of the flagellar axoneme is

less disorganized when other domains of DNAH17 are affected by pathogenic

variants (Whitfield et al., 2019),(Zhang et al., 2020),(Sha et al., 2020), likely because

an intact AAA4 domain is essential to physiological DNAH17 function (Snider et al.,

2008). Thus, our findings suggest that the intronic 13-bp deletion is a loss-of-function

allele.

We previously identified mutations at splice donor sites and nearby splice acceptor

sites that manifest in male reproductive disorders due to aberrant splicing (Iso-Touru

et al., 2019),(Hiltpold et al., 2020). Our present study reveals an intronic deletion

from 5 to 17 nucleotides upstream a canonical 3' splice site that perturbs splicing as

the most likely causal variant for a novel porcine sperm flagella defect. The 13-bp

deletion compromises splicing because it excises a polypyrimidine tract upstream

DNAH17 exon 56. Phenotypic consequences arising from polypyrimidine tract

mutations had rarely been described so far (Lefévre et al., 2002),(Sartelet et al.,

2015),(Abramowicz and Gos, 2018). To the best of our knowledge, our study reports

the first phenotype-genotype association for a mutation affecting an intronic

polypyrimidine tract in pigs. We provide evidence that the loss of the polypyrimidine

tract in porcine *DNAH17* intron 55 prevents recognition of the 3' splice site which

leads to the skipping of exon 56, thus causing defective sperm flagella. Sequence

motifs that govern the assembly of the spliceosome may be more distant to 3' splice

sites than the polypyrimidine tract in porcine *DNAH17* intron 55 (Zhang et al., 2017).

However, standard sequence variant annotation tools are largely blind to putative

consequences arising from mutations within intronic motifs. Thus, the systematic

characterization of branch points and polypyrimidine tracts seems warranted to

refine the functional classification of intronic variants.

Our findings enable the monitoring of the 13-bp deletion and unambiguous

identification of carrier and homozygous animals using direct gene testing. While

homozygosity for the 13-bp deletion is easily recognized in artificial insemination

boars using microscopic semen analysis, it may remain undetected in sows and

natural service boars. However, affected natural service boars will be noticed after

few matings due to low fertility. In agreement with previous findings on loss-of-

function alleles in human and murine *DNAH17* (Whitfield et al., 2019),(Zhang et al.,

2020),(Sha et al., 2020), homozygous mutation carriers were healthy. Considering

that the frequency of the 13-bp deletion is low and the majority of sows is

inseminated artificially, economic losses due to unsuccessful breeding with

homozygous boars are negligible in the Swiss Large White population.

The number of recessive conditions detected in livestock populations increases

steadily (Nicholas and Hobbs, 2014). Low effective population size, founder effects

and inbreeding favor the manifestation of recessive conditions. However, recent

advances in genome-wide genotyping and sequencing offer powerful tools for rapid

detection of causal variants underpinning inherited conditions (Bourneuf et al., 2017).

The phenotypic and economic consequences of the 13-bp deletion in *DNAH17* are

less detrimental than mutations that manifest in malformed and non-viable piglets

(Fang et al., 2020). However, the 13-bp deletion becomes a threat to the breeding

population if it attains higher frequency (Pausch et al., 2014),(Hiltpold et al., 2020).

The 13-bp deletion in *DNAH17* adds to a catalogue of sequence variants that cause

male-factor infertility in pigs (Sironen et al., 2011),(Sironen et al., 2006),(Noskova et

al., 2020). It remains an open question how an increasing number of undesired

recessive alleles may be considered appropriately in livestock populations (Cole,

2015),(Upperman et al., 2019). Although the newly identified 13-bp deletion does

neither compromise the welfare of homozygous animals nor result in huge economic

losses, it's frequency should be kept at a low level to avoid the birth of homozygous

boars that are infertile.

ACKNOWLEDGEMENTS

We acknowledge financial support from SUISAG, Micarna SA and the ETH Zürich

Foundation. We thank Dr. Cecilia Bebeacua (ScopeM) for support in transmission

electron microscopy. We are thankful for the excellent technical support provided by

the ETH Zürich technology platforms FGCZ (https://fgcz.ch/), ScopeM

(https://scopem.ethz.ch/), and Genetic Diversity Centre (https://gdc.ethz.ch/),

respectively, for sequencing, microscopy and DNA fragment analysis.

REFERENCES

Abramowicz, A., Gos, M., 2018. Splicing mutations in human genetic disorders: examples,
detection, and confirmation. J Appl Genet 59, 253–268.
https://doi.org/10.1007/s13353-018-0444-7

Andersson, M., Peltoniemi, O., Makinen, A., Sukura, A., Rodriguez-Martinez, H., 2000. The
Hereditary 'Short Tail' Sperm Defect - A New Reproductive Problem in Yorkshire
Boars. Reproduction in Domestic Animals 35, 59–63. https://doi.org/10.1046/j.1439-
0531.2000.00196.x

Aulchenko, Y.S., Ripke, S., Isaacs, A., van Duijn, C.M., 2007. GenABEL: an R library for
genome-wide association analysis. Bioinformatics 23, 1294–1296.
https://doi.org/10.1093/bioinformatics/btm108

Balding, D.J., 2006. A tutorial on statistical methods for population association studies.
Nature Reviews Genetics 7, 781–791. https://doi.org/10.1038/nrg1916

Ben Khelifa, M., Coutton, C., Zouari, R., Karaouzène, T., Rendu, J., Bidart, M., Yassine, S.,
Pierre, V., Delaroche, J., Hennebicq, S., Grunwald, D., Escalier, D., Pernet-Gallay, K.,
Jouk, P.-S., Thierry-Mieg, N., Touré, A., Arnoult, C., Ray, P.F., 2014. Mutations in
DNAH1, which Encodes an Inner Arm Heavy Chain Dynein, Lead to Male Infertility
from Multiple Morphological Abnormalities of the Sperm Flagella. American Journal
of Human Genetics 94, 95. https://doi.org/10.1016/j.ajhg.2013.11.017

Blom, E., 1950. A one-minute live-dead sperm stain by means of eosin-nigrosin. Fertility and
Sterility 1, 176–177.

Bourneuf, E., Otz, P., Pausch, H., Jagannathan, V., Michot, P., Grohs, C., Piton, G.,
Ammermüller, S., Deloche, M.-C., Fritz, S., Leclerc, H., Péchoux, C., Boukadiri, A.,
Hozé, C., Saintilan, R., Créchet, F., Mosca, M., Segelke, D., Guillaume, F., Bouet, S.,
Baur, A., Vasilescu, A., Genestout, L., Thomas, A., Allais-Bonnet, A., Rocha, D., Colle,
M.-A., Klopp, C., Esquerré, D., Wurmser, C., Flisikowski, K., Schwarzenbacher, H.,
Burgstaller, J., Brügmann, M., Dietschi, E., Rudolph, N., Freick, M., Barbey, S., Fayolle,
G., Danchin-Burge, C., Schibler, L., Bed'Hom, B., Hayes, B.J., Daetwyler, H.D., Fries, R.,
Boichard, D., Pin, D., Drögemüller, C., Capitan, A., 2017. Rapid Discovery of De Novo
Deleterious Mutations in Cattle Enhances the Value of Livestock as Model Species.
Scientific Reports 7, 11466. https://doi.org/10.1038/s41598-017-11523-3
Bray, N.L., Pimentel, H., Melsted, P., Pachter, L., 2016. Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol 34, 525–527. https://doi.org/10.1038/nbt.3519
Broekhuijse, M.L.W.J., Feitsma, H., Gadella, B.M., 2011. Field data analysis of boar semen
quality. Reprod. Domest. Anim. 46 Suppl 2, 59–63. https://doi.org/10.1111/j.1439-
0531.2011.01861.x
Browning, B.L., Zhou, Y., Browning, S.R., 2018. A One-Penny Imputed Genome from Next-
Generation Reference Panels. American Journal of Human Genetics 103, 338–348.
https://doi.org/10.1016/j.ajhg.2018.07.015
Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., Lee, J.J., 2015. Second-
generation PLINK: rising to the challenge of larger and richer datasets. Gigascience 4,
7. https://doi.org/10.1186/s13742-015-0047-8
Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. Fastp: An ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34, i884–i890. https://doi.org/10.1093/bioinformatics/bty560
Clark, E.L., Bush, S.J., McCulloch, M.E.B., Farquhar, I.L., Young, R., Lefevre, L., Pridans, C.,
Tsang, H.G., Wu, C., Afrasiabi, C., Watson, M., Whitelaw, C.B., Freeman, T.C.,
Summers, K.M., Archibald, A.L., Hume, D.A., 2017. A high resolution atlas of gene
expression in the domestic sheep (Ovis aries). PLOS Genetics 13, e1006997.
https://doi.org/10.1371/journal.pgen.1006997
Cole, J.B., 2015. A simple strategy for managing many recessive disorders in a dairy cattle
breeding program. Genet Sel Evol 47. https://doi.org/10.1186/s12711-015-0174-9
Cole, J.B., 2007. PyPedal: A computer program for pedigree analysis. Computers and
electronics in agriculture 57:107-113.
Colenbrander, B., Feitsma, H., Grooten, H.J., 1993. Optimizing semen production for
artificial insemination in swine. J. Reprod. Fertil. Suppl. 48, 207–215.
Coolidge, C.J., Seely, R.J., Patton, J.G., 1997. Functional analysis of the polypyrimidine tract
in pre-mRNA splicing. Nucleic Acids Res 25, 888–896.
Crysnanto, D., Wurmser, C., Pausch, H., 2019. Accurate sequence variant genotyping in
cattle using variation-aware genome graphs. Genetics Selection Evolution 51, 21.
https://doi.org/10.1186/s12711-019-0462-x
Depristo, M.A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J.R., Hartl, C., Philippakis,
A.A., Del Angel, G., Rivas, M.A., Hanna, M., McKenna, A., Fennell, T.J., Kernytsky,
A.M., Sivachenko, A.Y., Cibulskis, K., Gabriel, S.B., Altshuler, D., Daly, M.J., 2011. A
framework for variation discovery and genotyping using next-generation DNA
sequencing data. Nature Genetics 43, 491–501. https://doi.org/10.1038/ng.806
Diniz, D.B., Lopes, M.S., Broekhuijse, M.L.W.J., Lopes, P.S., Harlizius, B., Guimarães, S.E.F.,
Duijvesteijn, N., Knol, E.F., Silva, F.F., 2014. A genome-wide association study reveals

a novel candidate gene for sperm motility in pigs. Animal Reproduction Science 151,
201–207. https://doi.org/10.1016/j.anireprosci.2014.10.014
Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.,
Gingeras, T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29,
15–21. https://doi.org/10.1093/bioinformatics/bts635
Fang, L., Cai, W., Liu, S., Canela-Xandri, O., Gao, Y., Jiang, J., Rawlik, K., Li, B., Schroeder, S.G.,
Rosen, B.D., Li, C., Sonstegard, T.S., Alexander, L.J., Tassell, C.P.V., VanRaden, P.M.,
Cole, J.B., Yu, Y., Zhang, S., Tenesa, A., Ma, L., Liu, G.E., 2020. Comprehensive
analyses of 723 transcriptomes enhance genetic and biological interpretations for
complex traits in cattle. Genome Res 30,790-801.
https://doi.org/10.1101/gr.250704.119
Fang, Z.-H., Nosková, A., Crysnanto, D., Neuenschwander, S., Vögeli, P., Pausch, H., 2020. A
63-bp insertion in exon 2 of the porcine KIF21A gene is associated with
arthrogryposis multiplex congenita. Animal Genetics 51, 820–823.
https://doi.org/10.1111/age.12984
Gao, K., Masuda, A., Matsuura, T., Ohno, K., 2008. Human branch point consensus sequence
is yUnAy. Nucleic Acids Res 36, 2257–2267. https://doi.org/10.1093/nar/gkn073
Gleave, E.S., Schmidt, H., Carter, A.P., 2014. A structural analysis of the AAA+ domains in
Saccharomyces cerevisiae cytoplasmic dynein. J Struct Biol 186, 367–375.
https://doi.org/10.1016/j.jsb.2014.03.019
Grahofer, A., Gurtner, C., Nathues, H., 2017. Haemorrhagic bowel syndrome in fattening
pigs. Porcine Health Manag 3. https://doi.org/10.1186/s40813-017-0074-1
Hancock, J.L., 1957. The morphology of boar spermatozoa. J R Microsc Soc 76, 84–97.
https://doi.org/10.1111/j.1365-2818.1956.tb00443.x
Hiltpold, M., Niu, G., Kadri, N.K., Crysnanto, D., Fang, Z.-H., Spengeler, M., Schmitz-Hsu, F.,
Fuerst, C., Schwarzenbacher, H., Seefried, F.R., Seehusen, F., Witschi, U., Schnieke,
A., Fries, R., Bollwein, H., Flisikowski, K., Pausch, H., 2020. Activation of cryptic
splicing in bovine WDR19 is associated with reduced semen quality and male
fertility. PLOS Genetics 16, e1008804. https://doi.org/10.1371/journal.pgen.1008804
Holt, C., Holt, W.V., Moore, H.D., Reed, H.C., Curnock, R.M., 1997. Objectively measured
boar sperm motility parameters correlate with the outcomes of on-farm
inseminations: results of two fertility trials. J. Androl. 18, 312–323.
Inaba, K., 2003. Molecular Architecture of the Sperm Flagella: Molecules for Motility and
Signaling. jzoo 20, 1043–1056. https://doi.org/10.2108/zsj.20.1043
Iso-Touru, T., Wurmser, C., Venhoranta, H., Hiltpold, M., Savolainen, T., Sironen, A., Fischer,
K., Flisikowski, K., Fries, R., Vicente-Carrillo, A., Alvarez-Rodriguez, M., Nagy, S.,
Mutikainen, M., Peippo, J., Taponen, J., Sahana, G., Guldbrandtsen, B., Simonen, H.,
Rodriguez-Martinez, H., Andersson, M., Pausch, H., 2019. A splice donor variant in
CCDC189 is associated with asthenospermia in Nordic Red dairy cattle. BMC
Genomics 20, 286. https://doi.org/10.1186/s12864-019-5628-y
Jurewicz, J., Hanke, W., Radwan, M., Bonde, J.P., 2009. Environmental factors and semen
quality. Int J Occup Med Environ Health 22, 305–329.
https://doi.org/10.2478/v10001-009-0036-1
Kadri, N.K., Sahana, G., Charlier, C., Iso-Touru, T., Guldbrandtsen, B., Karim, L., Nielsen, U.S.,
Panitz, F., Aamand, G.P., Schulman, N., Georges, M., Vilkki, J., Lund, M.S., Druet, T.,
2014. A 660-Kb Deletion with Antagonistic Effects on Fertility and Milk Production
Segregates at High Frequency in Nordic Red Cattle: Additional Evidence for the

Common Occurrence of Balancing Selection in Livestock. PLOS Genetics 10, e1004049. https://doi.org/10.1371/journal.pgen.1004049

Knox, R.V., 2016. Artificial insemination in pigs today. Theriogenology 85, 83–93. https://doi.org/10.1016/j.theriogenology.2015.07.009

Kosova, G., Scott, N.M., Niederberger, C., Prins, G.S., Ober, C., 2012. Genome-wide Association Study Identifies Candidate Genes for Male Fertility Traits in Humans. The American Journal of Human Genetics 90, 950–961. https://doi.org/10.1016/j.ajhg.2012.04.016

Lamas-Toranzo, I., Hamze, J.G., Bianchi, E., Fernández-Fuertes, B., Pérez-Cerezales, S., Laguna-Barraza, R., Fernández-González, R., Lonergan, P., Gutiérrez-Adán, A., Wright, G.J., Jiménez-Movilla, M., Bermejo-Álvarez, P., 2020. TMEM95 is a sperm membrane protein essential for mammalian fertilization. eLife 9, e53913. https://doi.org/10.7554/eLife.53913

Lefévre, S.H., Chauveinc, L., Stoppa-Lyonnet, D., Michon, J., Lumbroso, L., Berthet, P., Frappaz, D., Dutrillaux, B., Chevillard, S., Malfoy, B., 2002. A T to C mutation in the polypyrimidine tract of the exon 9 splicing site of the RB1 gene responsible for low penetrance hereditary retinoblastoma. Journal of Medical Genetics 39, E21. https://doi.org/10.1136/jmg.39.5.e21

Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM, arXiv:13033997.

Liu, C., Miyata, H., Gao, Y., Sha, Y., Tang, S., Xu, Z., Whitfield, M., Patrat, C., Wu, H., Dulioust, E., Tian, S., Shimada, K., Cong, J., Noda, T., Li, H., Morohoshi, A., Cazin, C., Kherraf, Z.-E., Arnoult, C., Jin, L., He, X., Ray, P.F., Cao, Y., Touré, A., Zhang, F., Ikawa, M., 2020. Bi-allelic DNAH8 Variants Lead to Multiple Morphological Abnormalities of the Sperm Flagella and Primary Male Infertility. The American Journal of Human Genetics 107, 330–341. https://doi.org/10.1016/j.ajhg.2020.06.004

Manchia, M., Cullis, J., Turecki, G., Rouleau, G.A., Uher, R., Alda, M., 2013. The Impact of Phenotypic and Genetic Heterogeneity on Results of Genome Wide Association Studies of Complex Diseases. PLoS One 8. https://doi.org/10.1371/journal.pone.0076295

Marques, D.B.D., Lopes, M.S., Broekhuijse, M.L.W.J., Guimarães, S.E.F., Knol, E.F., Bastiaansen, J.W.M., Silva, F.F., Lopes, P.S., 2017. Genetic parameters for semen quality and quantity traits in five pig lines. J. Anim. Sci. 95, 4251–4259. https://doi.org/10.2527/jas2017.1683

McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R.S., Thormann, A., Flicek, P., Cunningham, F., 2016. The Ensembl Variant Effect Predictor. Genome Biology 17:122. https://doi.org/10.1186/s13059-016-0974-4

Ni, X., Wang, J., Lv, M., Liu, C., Zhong, Y., Tian, S., Wu, H., Cheng, H., Gao, Y., Tan, Q., Chen, B., Li, Q., Song, B., Wei, Z., Zhou, P., He, X., Zhang, F., Cao, Y., 2020. A novel homozygous mutation in WDR19 induces disorganization of microtubules in sperm flagella and nonsyndromic asthenoteratospermia. J Assist Reprod Genet 37, 1431–1439. https://doi.org/10.1007/s10815-020-01770-1

Nicholas, F.W., Hobbs, M., 2014. Mutation discovery for Mendelian traits in non-laboratory animals: a review of achievements up to 2012. Anim. Genet. 45, 157–170. https://doi.org/10.1111/age.12103

Nijs, M., Vanderzwalmen, P., Vandamme, B., Segal-Bertin, G., Lejeune, B., Segal, L., van Roosendaal, E., Schoysman, R., 1996. Fertilizing ability of immotile spermatozoa after

intracytoplasmic sperm injection. Hum. Reprod. 11, 2180–2185.
https://doi.org/10.1093/oxfordjournals.humrep.a019073

Noda, T., Lu, Y., Fujihara, Y., Oura, S., Koyano, T., Kobayashi, S., Matzuk, M.M., Ikawa, M.,
2020. Sperm proteins SOF1, TMEM95, and SPACA6 are required for sperm–oocyte
fusion in mice. PNAS 117, 11493–11502. https://doi.org/10.1073/pnas.1922650117

Noskova, A., Wurmser, C., Crysnanto, D., Sironen, A., Uimari, P., Fries, R., Andersson, M.,
Pausch, H., 2020. Deletion of porcine BOLL is associated with defective acrosomes
and subfertility in Yorkshire boars. Animal Genetics, Advanced online,
https://doi.org/10.1111/age.12998

Ortega, C., Verheyen, G., Raick, D., Camus, M., Devroey, P., Tournaye, H., 2011. Absolute
asthenozoospermia and ICSI: what are the options? Hum Reprod Update 17, 684–
692. https://doi.org/10.1093/humupd/dmr018

Pausch, H., Ammermüller, S., Wurmser, C., Hamann, H., Tetens, J., Drögemüller, C., Fries, R.,
2016a. A nonsense mutation in the COL7A1 gene causes epidermolysis bullosa in
Vorderwald cattle. BMC Genet. 17, 149. https://doi.org/10.1186/s12863-016-0458-2

Pausch, H., Flisikowski, K., Jung, S., Emmerling, R., Edel, C., Götz, K.-U., Fries, R., 2011.
Genome-wide association study identifies two major loci affecting calving ease and
growth-related traits in cattle. Genetics 187, 289–297.
https://doi.org/10.1534/genetics.110.124057

Pausch, H., Kölle, S., Wurmser, C., Schwarzenbacher, H., Emmerling, R., Jansen, S.,
Trottmann, M., Fuerst, C., Götz, K.-U., Fries, R., 2014. A Nonsense Mutation in
TMEM95 Encoding a Nondescript Transmembrane Protein Causes Idiopathic Male
Subfertility in Cattle. PLOS Genetics 10, e1004044.
https://doi.org/10.1371/journal.pgen.1004044

Pausch, H., Venhoranta, H., Wurmser, C., Hakala, K., Iso-Touru, T., Sironen, A., Vingborg,
R.K., Lohi, H., Söderquist, L., Fries, R., Andersson, M., 2016b. A frameshift mutation
in ARMC3 is associated with a tail stump sperm defect in Swedish Red (Bos taurus)
cattle. BMC Genet. 17, 49. https://doi.org/10.1186/s12863-016-0356-7

Pedersen, B.S., Quinlan, A.R., 2018. Mosdepth: Quick coverage calculation for genomes and
exomes. Bioinformatics 34, 867–868. https://doi.org/10.1093/bioinformatics/btx699

Price, A.L., Patterson, N.J., Plenge, R.M., Weinblatt, M.E., Shadick, N.A., Reich, D., 2006.
Principal components analysis corrects for stratification in genome-wide association
studies. Nature Genetics 38, 904–909. https://doi.org/10.1038/ng1847

Price, A.L., Zaitlen, N.A., Reich, D., Patterson, N., 2010. New approaches to population
stratification in genome-wide association studies. Nature reviews. Genetics 11, 459.
https://doi.org/10.1038/nrg2813

Rahban, R., Priskorn, L., Senn, A., Stettler, E., Galli, F., Vargas, J., Van den Bergh, M., Fusconi,
A., Garlantezec, R., Jensen, T.K., Multigner, L., Skakkebaek, N.E., Germond, M.,
Jørgensen, N., Nef, S., NICER Working Group, 2019. Semen quality of young men in
Switzerland: a nationwide cross-sectional population-based study. Andrology 7, 818–
826. https://doi.org/10.1111/andr.12645

Reese, M.G., Eeckman, F.H., Kulp, D., Haussler, D., 1997. Improved splice site detection in
Genie. J. Comput. Biol. 4, 311–323. https://doi.org/10.1089/cmb.1997.4.311

Robic, A., Faraut, T., Djebali, S., Weikard, R., Feve, K., Maman, S., Kuehn, C., 2019. Analysis
of pig transcriptomes suggests a global regulation mechanism enabling temporary
bursts of circular RNAs. RNA Biology 16, 1190–1204.
https://doi.org/10.1080/15476286.2019.1621621

Sartelet, A., Li, W., Pailhoux, E., Richard, C., Tamma, N., Karim, L., Fasquelle, C., Druet, T.,
Coppieters, W., Georges, M., Charlier, C., 2015. Genome-wide next-generation DNA
and RNA sequencing reveals a mutation that perturbs splicing of the
phosphatidylinositol glycan anchor biosynthesis class H gene (PIGH) and causes
arthrogryposis in Belgian Blue cattle. BMC Genomics 16, 316.
https://doi.org/10.1186/s12864-015-1528-y
Sato, Y., Tajima, A., Kiguchi, M., Kogusuri, S., Fujii, A., Sato, T., Nozawa, S., Yoshiike, M.,
Mieno, M., Kojo, K., Uchida, M., Tsuchiya, H., Yamasaki, K., Imoto, I., Iwamoto, T.,
2020. Genome-wide association study of semen volume, sperm concentration, testis
size, and plasma inhibin B levels. Journal of Human Genetics 65, 683–691.
https://doi.org/10.1038/s10038-020-0757-3
Sato, Y., Tajima, A., Sato, T., Nozawa, S., Yoshiike, M., Imoto, I., Yamauchi, A., Iwamoto, T.,
2018. Genome-wide association study identifies ERBB4 on 2q34 as a novel locus
associated with sperm motility in Japanese men. J Med Genet 55, 1–7.
https://doi.org/10.1136/jmedgenet-2017-104991
Schwarzenbacher, H., Burgstaller, J., Seefried, F.R., Wurmser, C., Hilbe, M., Jung, S., Fuerst,
C., Dinhopl, N., Weissenböck, H., Fuerst-Waltl, B., Dolezal, M., Winkler, R., Grueter,
O., Bleul, U., Wittek, T., Fries, R., Pausch, H., 2016. A missense mutation in TUBD1 is
associated with high juvenile mortality in Braunvieh and Fleckvieh cattle. BMC
Genomics 17, 400. https://doi.org/10.1186/s12864-016-2742-y
Sha, Y., Wei, X., Ding, L., Mei, L., Huang, X., Lin, S., Su, Z., Kong, L., Zhang, Y., Ji, Z., 2020.
DNAH17 is associated with asthenozoospermia and multiple morphological
abnormalities of sperm flagella. Annals of Human Genetics 84, 271–279.
https://doi.org/10.1111/ahg.12369
Sironen, A., Shoemark, A., Patel, M., Loebinger, M.R., Mitchison, H.M., 2020. Sperm defects
in primary ciliary dyskinesia and related causes of male infertility. Cell. Mol. Life Sci.
77, 2029–2048. https://doi.org/10.1007/s00018-019-03389-7
Sironen, A., Thomsen, B., Andersson, M., Ahola, V., Vilkki, J., 2006. An intronic insertion in
KPL2 results in aberrant splicing and causes the immotile short-tail sperm defect in
the pig. PNAS 103, 5006–5011. https://doi.org/10.1073/pnas.0506318103
Sironen, A., Uimari, P., Venhoranta, H., Andersson, M., Vilkki, J., 2011. An exonic insertion
within Tex14 gene causes spermatogenic arrest in pigs. BMC Genomics 12, 591.
https://doi.org/10.1186/1471-2164-12-591
Snider, J., Thibault, G., Houry, W.A., 2008. The AAA+ superfamily of functionally diverse
proteins. Genome Biol 9, 216. https://doi.org/10.1186/gb-2008-9-4-216
Soneson, C., Love, M.I., Robinson, M.D., 2015. Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. F1000Res 4, 1521.
https://doi.org/10.12688/f1000research.7563.2
Tarasov, A., Vilella, A.J., Cuppen, E., Nijman, I.J., Prins, P., 2015. Sambamba: Fast processing
of NGS alignment formats. Bioinformatics 31, 2032–2034.
https://doi.org/10.1093/bioinformatics/btv098
Thibier, M., Wagner, H.-G., 2002. World statistics for artificial insemination in cattle.
Livestock Production Science 74, 203–212. https://doi.org/10.1016/S0301-
6226(01)00291-3
Tu, C., Nie, H., Meng, L., Yuan, S., He, W., Luo, A., Li, H., Li, W., Du, J., Lu, G., Lin, G., Tan, Y.-
Q., 2019. Identification of DNAH6 mutations in infertile men with multiple

943   morphological abnormalities of the sperm flagella. Scientific Reports 9, 15864.
944   https://doi.org/10.1038/s41598-019-52436-7

945 Upperman, L.R., Kinghorn, B.P., MacNeil, M.D., Van Eenennaam, A.L., 2019. Management of
946   lethal recessive alleles in beef cattle through the use of mate selection software.
947   Genetics Selection Evolution 51, 36. https://doi.org/10.1186/s12711-019-0477-3

948 Venhoranta, H., Pausch, H., Flisikowski, K., Wurmser, C., Taponen, J., Rautala, H., Kind, A.,
949   Schnieke, A., Fries, R., Lohi, H., Andersson, M., 2014. In frame exon skipping in
950   UBE3B is associated with developmental disorders and increased mortality in cattle.
951   BMC Genomics 15:890, 1–9. https://doi.org/10.1186/1471-2164-15-890

952 Wambergue, C., Zouari, R., Fourati Ben Mustapha, S., Martinez, G., Devillard, F., Hennebicq,
953   S., Satre, V., Brouillet, S., Halouani, L., Marrakchi, O., Makni, M., Latrous, H., Kharouf,
954   M., Amblard, F., Arnoult, C., Ray, P.F., Coutton, C., 2016. Patients with multiple
955   morphological abnormalities of the sperm flagella due to DNAH1 mutations have a
956   good prognosis following intracytoplasmic sperm injection. Hum Reprod 31, 1164–
957   1172. https://doi.org/10.1093/humrep/dew083

958 Warr, A., Affara, N., Aken, B., Beiki, H., Bickhart, D.M., Billis, K., Chow, W., Eory, L., Finlayson,
959   H.A., Flicek, P., Girón, C.G., Griffin, D.K., Hall, R., Hannum, G., Hourlier, T., Howe, K.,
960   Hume, D.A., Izuogu, O., Kim, K., Koren, S., Liu, H., Manchanda, N., Martin, F.J.,
961   Nonneman, D.J., O'Connor, R.E., Phillippy, A.M., Rohrer, G.A., Rosen, B.D., Rund, L.A.,
962   Sargent, C.A., Schook, L.B., Schroeder, S.G., Schwartz, A.S., Skinner, B.M., Talbot, R.,
963   Tseng, E., Tuggle, C.K., Watson, M., Smith, T.P.L., Archibald, A.L., 2019. An improved
964   pig reference genome sequence to enable pig genetics and genomics research.
965   bioRxiv 668921. https://doi.org/10.1101/668921

966 Whitfield, M., Thomas, L., Bequignon, E., Schmitt, A., Stouvenel, L., Montantin, G., Tissier, S.,
967   Duquesnoy, P., Copin, B., Chantot, S., Dastot, F., Faucon, C., Barbotin, A.L., Loyens,
968   A., Siffroi, J.-P., Papon, J.-F., Escudier, E., Amselem, S., Mitchell, V., Touré, A.,
969   Legendre, M., 2019. Mutations in DNAH17, Encoding a Sperm-Specific Axonemal
970   Outer Dynein Arm Heavy Chain, Cause Isolated Male Infertility Due to
971   Asthenozoospermia. The American Journal of Human Genetics 105, 198–212.
972   https://doi.org/10.1016/j.ajhg.2019.04.015

973 Xavier, M.J., Salas-Huetos, A., Oud, M.S., Aston, K.I., Veltman, J.A., 2020. Disease gene
974   discovery in male infertility: past, present and future. Hum Genet.
975   https://doi.org/10.1007/s00439-020-02202-x

976 Zhang, B., Ma, H., Khan, T., Ma, A., Li, T., Zhang, H., Gao, J., Zhou, J., Li, Y., Yu, C., Bao, J., Ali,
977   A., Murtaza, G., Yin, H., Gao, Q., Jiang, X., Zhang, F., Liu, C., Khan, I., Zubair, M.,
978   Hussain, H.M.J., Khan, R., Yousaf, A., Yuan, L., Lu, Y., Xu, X., Wang, Y., Tao, Q., Hao,
979   Q., Fang, H., Cheng, H., Zhang, Y., Shi, Q., 2020. A DNAH17 missense variant causes
980   flagella destabilization and asthenozoospermia. J Exp Med 217.
981   https://doi.org/10.1084/jem.20182365

982 Zhang, Q., Fan, X., Wang, Y., Sun, M., Shao, J., Guo, D., 2017. BPP: a sequence-based
983   algorithm for branch point prediction. Bioinformatics 33, 3166–3172.
984   https://doi.org/10.1093/bioinformatics/btx401

 SUPPLEMENTAL FILES

Supplemental file 1

Title: Accession numbers of 87 boars

Format: xlsx

Description: Accession numbers of the sequence data of five affected

(asthenozoospermic) and 82 unaffected (control) boars. The third column indicates

the haplotype status (0 – not carrier, 1 – heterozygous carrier, 2 – homozygous

carrier) of 65 sequenced boars that also had Illumina PorcineSNP60 BeadChip-

called genotypes.

Supplemental file 2

Format: jpg

Title: Sperm viability assessment by eosin-nigrosin staining

Description: Representative light microscopic image of sperm from an affected boar

stained with eosin-nigrosin. The sperm were flushed post mortem from the

epididymis of Boar_1249. Live sperm appear white in color. Dead sperm appear pink

in color because they take up eosin due to loss of membrane integrity.

1005

Supplemental file 3

Title: Disorganized flagellar axonemes in two affected boars

Format: png

Description: Cross-sections of sperm from two affected boars reveal multiple

morphological and ultrastructural abnormalities of the flagella including the absence

of outer or inner doublets, secondary flagella-like structures within the same

membrane, and flagellar components in the cytoplasmic droplet-like structure.

1013

Supplemental file 4

Title: Principal components analysis of cases and controls

Format: png

Description: Scatter plot of the top five eigenvectors of a genomic relationship matrix.

Grey and red color indicates 100 fertile and six asthenoteratozoospermic boars from

the Swiss Large White population. The raw data underlying the plot (genotypes,

haplotypes, eigenvectors) are available at available at

https://doi.org/10.5281/zenodo.4081475.

1022

Supplemental file 5

Title: Expression profile of porcine *DNAH17* in pubertal boars

Format: png

Description: Expression of *DNAH17* quantified using testis RNAseq alignments of

four pubertal boars (SRR2564762, SRR8237161, SRR8237162, SRR8237165). The

number of reads covering a position was extracted from coordinate sorted BAM files

using the MOSDEPTH software, divided by the total number of reads mapped to

transcripts. Blue color indicates the exon-intron structure of porcine *DNAH17*. Exon

56 is displayed in red color. Although exon expression is somewhat variable between

the samples, the expression of exon 56 is evident in all samples. Raw data

underlying the plot are available at the European Nucleotide Archive (ENA) of the

EMBL at the BioProject PRJNA506525.