Quantitative assessment of NCLDV–host interactions predicted by co-occurrence analyses

Lingjie Meng[a], Hisashi Endo[a], Romain Blanc-Mathieu[a,b], Rodrigo Hernández-Velázquez[a,c], Hiroto Kaneko[a], Hiroyuki Ogata[a,#]

Affiliations:

a. Bioinformatics Center, Institute for Chemical Research, Kyoto University, Gokasho, Uji, 611-0011, Japan

b. Laboratoire de Physiologie Cellulaire & Végétale, CEA, Univ. Grenoble Alpes, CNRS, INRA, IRIG, Grenoble, France

c. Max Planck Institute for Marine Microbiology, Celsiusstraße 1, Bremen, Germany

[#]*Corresponding author*:

Hiroyuki Ogata (E-mail: ogata@kuicr.kyoto-u.ac.jp, Phone: +81-774-38-3270)

Abstract

Nucleocytoplasmic DNA viruses (NCLDVs) are highly diverse and abundant in marine environments. However, knowledge of their hosts is limited because only a few NCLDVs have been isolated. By taking advantage of the rapidly increasing metagenomic data, *in silico* host prediction approaches are expected to fill this gap between known virus–host relationships and the true but largely unknown amount of NCLDVs. In this study, we built co-occurrence networks between NCLDVs and eukaryotes using the *Tara* Oceans metagenome and metabarcoding datasets to predict virus–host interactions. Using the positive likelihood ratio to assess the performance of host prediction for NCLDVs, we demonstrated that co-occurrence approaches can increase the odds of predicting true positive relationships four-fold compared with random host predictions in the high-weight region (weight > 0.4). To refine the host predictions from high-dimensional co-occurrence networks, we employed a recently proposed phylogeny-based method, Taxon Interaction Mapper, and showed that Taxon Interaction Mapper further improved the prediction performance eight-fold using weight cut-off filtration (> 0.4). Finally, we inferred virophage and NCLDV networks that further corroborated that co-occurrence approaches are effective for predicting NCLDV hosts in marine environments.

Importance

NCLDVs infect a wide range of eukaryotes. Their life circle is less dependent on hosts than other viruses. However, our understanding of NCLDV–host systems is highly limited because few of these viruses have been isolated. Co-occurrence inference is a possible way to predict virus–host interactions. For the first time, we quantitatively assessed the effectiveness of co-occurrence analyses for NCLDV host prediction. We also improved the performance of these co-occurrence analyses with a phylogeny-guided filtering method and produced a list of candidate hosts for three NCLDV families.

Introduction

Nucleocytoplasmic large DNA viruses (NCLDVs) represent a group of double-stranded DNA viruses that belong to the viral phylum *Nucleocytoviricota*, which was previously referred to as Megavirales (Virus Taxonomy: 2019 Release) (1, 2). NCLDVs usually possess diverse gene repertoires (74 to more than 2000 proteins), large genomes (45 kb to 2.5 Mb), and outsized virions (80 nm to 1.5 μm) (3–5). NCLDVs usually have high functional autonomy and encode components of replication, transcription, and translation systems (3). Recently, a virus that belongs to a new family of NCLDVs called "Medusaviridae" was found to encode five types of histones (6). The existence of metabolically active viral factories and infectious virophages also indicates that the life cycle of NCLDVs is less dependent on the host cells than other viruses (7, 8). To determine what allows these giant viruses to be more independent than other viruses, the first step is to identify their hosts—"Who infects whom?"

NCLDVs are known to infect a broad range of eukaryotes, from unicellular eukaryotes and macroalgae to animals (9). Amoebae are frequently used hosts in co-culture to isolate large NCLDVs (10). However, there is growing evidence, especially in marine systems, that NCLDVs can infect many phytoplankton groups, such as Pelagophyceae, Mamiellophyceae, and Dinophyceae (11, 12). Several other non-photosynthetic eukaryotic lineages, such as Bicoecea and Choanoflagellatea, were also reported as experimentally identified NCLDV hosts in marine environments (13, 14). Small to large marine organisms, including invertebrates and vertebrates, are infected by viruses that belong to the NCLDV family *Iridoviridae* (15, 16). These studies together indicate ubiquitous infectious relationships between NCLDVs and a wide range of marine eukaryotes. However, our understanding of NCLDV–host systems is highly limited because few viruses have been isolated.

The number of viruses and hosts isolated in the laboratory is effectively a drop in the

ocean compared with the amount that truly exists. NCLDVs have been found to be highly

diverse and abundant based on omics data (17, 18). In only a few liters of coastal seawater,

more than 5,000 *Mimiviridae* species were detected; by comparison, only 20 reference

Mimiviridae with known hosts have been well investigated (19). Global marine metagenomic

data have revealed that the richness and phylogenetic diversity of NCLDVs are even higher

than those of an entire prokaryotic domain (20). From biogeographical evidence, it is clear

that these viruses are prevalent in the oceanic environment but have a heterogeneous

community structure across sizes, depths, and biomes (21). Marine metatranscriptomic data

have also shown that NCLDVs are active everywhere in sunlit oceans and may infect hosts

from small piconanoplankton (0.8–5 μm) to large mesoplankton (180–2000 μm) (22).

Apart from quantity, previous studies also demonstrated that NCLDVs have the

potential to infect a greater diversity of hosts than known to date. Gene transfer between

NCLDVs and eukaryotes has indicated that these viruses may be able to infect most major

eukaryotic lineages (23, 24). NCLDVs might have started coevolving with eukaryotes even

before the last eukaryotic common ancestor (LECA) (25). A recent study supported this

hypothesis by showing that some NCLDVs encode viractins (actin-related genes in viruses),

which could have been acquired from proto-eukaryotes and possibly reintroduced the pre-

LECA eukaryotic lineage (26). Together, the findings of high abundance and a long history

of coevolution showcase that what is known about NCLDV hosts is just the tip of the iceberg.

The real host range of these viruses might be concordant with their known diversity.

Therefore, more effort is needed to identify hosts to elucidate the poorly known virus–host

relationships and the largely unknown NCLDV world.

Substantial effort has been made to reveal the interactions between NCLDVs and

their putative hosts. Apart from the classic co-culture method, some other alternative

methods, including high-throughput cell sorting, are also used (10, 14). Because of its high

efficiency, metagenomics has been increasingly used to investigate the host range of

NCLDVs. Genome analyses, such as to identify horizontal gene transfer (HGT), have also

performed well for NCLDV host prediction (23, 24).

Abundance-based co-occurrence analyses have been used for host prediction and are

supposed to be effective because viruses can only thrive in an environment where their hosts

exist. In addition to virus–host relationships, this method was applied to show the association

between NCLDVs and their "parasites" (virophages) (27). However, co-occurrence-based

prediction is controversial for viral host prediction, because the abundance dynamics of

viruses (e.g., *Emiliania huxleyi* and *Heterosigma akashiwo* viruses) and their hosts are

sometimes not concordant (28, 29). Usually, validation with known virus–host relationships

or corroboration with genomic evidence (e.g., HGT) is used to assess network-based

predictions (17, 30). However, the effectiveness of co-occurrence network methods has never

been quantitatively tested for NCLDV host prediction. The current lack of quantitative

assessment hinders the widespread use of this approach. Therefore, methods are needed that

test the accuracy of NCLDV host prediction with co-occurrence networks. Effort is also

needed to improve the performance of co-occurrence-based prediction.

The *Tara* Oceans expedition is a global-scale survey on marine ecosystems that

expands our knowledge of microbial diversity, organismal interactions, and ecological

drivers of community structure (31). The present study used *Tara* Oceans metagenomic and

metabarcoding datasets to predict the virus–host relationships between NCLDVs and

eukaryotes by constructing co-occurrence networks. To quantitatively assess the performance

of network-based host prediction, we employed the positive likelihood ratio (LR+) for

prediction with reference data for known NCLDV–host relationships. The performance of a

recently proposed phylogeny-based enrichment analysis approach, Taxon Interaction Mapper

(TIM) (32), was also assessed. We further examined NCLDV–virophage networks to

investigate the effectiveness of co-occurrence approaches for predicting NCLDV hosts.

Results

NCLDV–eukaryote co-occurrence networks

From five datasets that corresponded to five size fractions, we generated five co-

occurrence networks. Altogether, these networks were composed of 47978 *polB*–V9

associations (edges). Out of these associations, 47296 had positive weights, and 682 had

negative weights (Fig. 2A). For the nodes, 20148 V9 and 5234 *polB* OTUs were involved in

the networks. The associations that involved the family *Mimiviridae* were numerically

dominant (n = 36830) among the different NCLDV families. The second most numerous

family was *Phycodnaviridae*, with 5521 edges with eukaryotes. No other family had more

than 2000 associations with eukaryotes. *Marseilleviridae*, the least numerous family in the

networks, had 132 edges with eukaryotes. Taxonomic annotation of eukaryotic OTUs

indicated that Alveolata, Opisthokonta, Rhizaria, and Stramenopiles were the major four

eukaryotic groups that were connected to NCLDVs (with 21167, 9179, 6521, and 5327

edges, respectively). Three of these eukaryotic groups belong to the SAR supergroup (i.e.,

Stramenopiles, Alveolata, and Rhizaria), which represented 68.81% of the total associations.

Regarding the pairs between viral families and eukaryotic lineages, *Mimiviridae* and

Alveolata showed the largest number of edges (n = 16548). Besides NCLDV–eukaryote

associations, we detected 57495 *polB*–*polB* associations and 234448 V9–V9 associations

(Fig. S2B). We also included environmental parameters in the network inference and

identified 25 pairs of associations between environmental parameters and *polB* OTUs (Table

S3).

The number of NCLDV–eukaryote associations generally decreased with increasing

size fractionation (Fig. S2A). The largest number of *polB*–V9 associations was found in the

0.8–5-μm fraction (n = 10647). However, the 0.8–inf-μm size fraction network was the

largest (n = 10477) for edges with positive weights. With the annotation of major lineages,

the eukaryotic community compositions in the networks varied over different size fractions

(Fig. 2B). In the smallest size fraction (0.8–5 μm) and the large range size fraction (0.8–inf

μm), Marine Alveolate Group II was the eukaryotic lineage with the largest number of

associations with NCLDVs (21.39% and 19.98%, respectively). Dinophyceae was the second

largest group connected to NCLDVs in these two size fractions and showed the largest

number of connections with NCLDVs in the 5–20-μm size fraction network (22.22% of total

interactions). The viral associations with Metazoa and Collodaria increased with increasing

size fractions. In the largest 180–2000-μm size fraction network, Metazoa contributed

39.31% of the total *polB*–V9 edges.

We calculated the degree of nodes (number of connected edges) for each NCLDV

polB OTU (Fig. 3A, B). Largely consistent with the total number of edges, the average

degree of positive associations per *polB* in all size fractions was higher than negative edges

and decreased along with increasing size fractions (2.69, 2.40, 2.25, and 2.10 from 0.8–5 μm

to 180–2000 μm, and 2.76 for 0.8–inf). Most of the *polB* nodes had more than one positive

association (Fig. 3A). Together with the taxonomic annotation of nodes, *polB*–V9

associations in the networks generated with the *Tara* Oceans data revealed their high

dimensionality and complexity.

Network validation

We quantitatively assessed the performance of predicting *polB*–V9 associations using

the positive likelihood ratio (LR+) (Fig. 1). By defining groups of metagenomic PolBs as

described in the Materials and Methods, 932 OTUs were recruited in the validation, and these

sequences contributed 6191 *polB*–V9 associations in the FlashWeave networks (Fig. S3). To

obtain an overall performance, we assessed the pooled associations (by removing

redundancy) from the five co-occurrence networks. LR+ was separately calculated for edges

with positive and negative weights. As shown in Fig. 4A, the LR+ of host prediction for

positive associations was higher than 1 (LR+ = 1 indicates no change in the likelihood of the

condition). The LR+ was generally higher when the cut-off for the FlashWeave weight was

high than when the cut-off was lower, which indicated that condition positive cases are

enriched in the edges with high weights. This result demonstrated that the co-occurrence-

based host prediction of NCLDVs outperformed random prediction (i.e., random inference of

virus–host pairs). In the high-weight regions: 1) weight > 0.6, the LR+ of associations was

higher than 10; 2) weight > 0.4, the LR+ was roughly higher than 4. Nonetheless, the false

discovery rate (FDR) was high (Fig. S4A), which indicated that the predictions contained

numerous virus–host edges that were not considered condition positive. FDR was 96.34%

when weight $\geq$ 0.4. An assessment of the host prediction for negative weight associations was

also carried out. There were no known NCLDV–host pairs found in the negative networks

(Fig. S4B). The analysis of the remaining part of our study was thus conducted for positive

associations.

Comparing the performance between different size fractions indicated that the

networks of small size fractions (including the 0.8–inf-μm size fraction) performed better in

predicting the NCLDV–host relationships (Fig. 4B). The 0.8–inf-μm size fraction had the

highest average LR+ out of the five size fractions (LR+ = 4.97). The LR+ of small size

fractions was generally higher than that of large size fractions, but there were exceptions

between 180–2000 and 20–180 μm. The average LR+ for the associations in the 20–180-μm

size fraction network was even lower than 1, which indicated that most of the predicted

NCLDV–eukaryote pairs in this size fraction are not yet known. Different from the average

results, the high-weight region (> 0.5) showed that the associations of the large size fractions

(5–20 and 180–2000 μm) performed better than those of small size fractions (0.8–5 μm) in

terms of both LR+ and FDR (Fig. S5 A, B).

Comparison of three filtration strategies did not show a consistent pattern in

prediction performance. When we compared the pooled associations from all size fractions,

the networks from the Q1 filtration strategy performed best (Fig. S6A). However, if we only

used the samples in the 0.8–inf-μm size fraction, which was shown to perform best for

NCLDV host prediction, the most stringent Q3 strategy performed best (Fig. S6B). Finally,

we compared the performance of networks inferred by three different methods: FlashWeave-

S (sensitive model), FlashWeave-HE (heterogeneous model), and FastSpar. FlashWeave-

S performed best (Fig. S7). Of note, the LR+ of these three methods were all larger than 1.

Assessment of host prediction improvement

After the phylogeny-guided TIM filtering of the host prediction, we identified 24

eukaryotic taxonomic groups specifically associated with NCLDV (Fig. S8). To compare the

performance of the TIM results with the above raw FlashWeave results, we transferred the

three primary eukaryotic taxonomic ranks into their associated major lineages (Table S4), and

the connections were plotted on a network (Fig. 5A). This network showed that three out of

nine NCLDV families (*Mimiviridae, Phycodnaviridae*, and *Iridoviridae*) had enriched

connections with eukaryotic lineages. Among the network edges, known virus–host pairs

were found, such as Haptophyta–*Mimiviridae*, Mamiellophyceae–*Phycodnaviridae*, and

Metazoa–*Iridoviridae*. The associations in the TIM-filtered results showed a sharp

improvement in performance from the original result with and without an edge weight cut-

off. The average LR+ of TIM-enriched associations was 42.22, which was higher than the

raw FlashWeave associations without a weight cut-off (3.43) and with a weight cut-off of 0.4

(5.20) (Fig. 5B). The FDR dropped to 0.74 from 0.97 (no cut-off) and 0.95 (weight cut-off of

0.4) (Fig. 5C).

It was apparent from the network that *Mimiviridae* had a diverse putative host range

(12 lineages) of algae, protozoans, and metazoans. Metazoa had the most enriched nodes

connected to *Mimiviridae*; additionally, MAST-3, Cryptophyta, Foraminifera, and Ciliophora

had strong relationships with *Mimiviridae*. For *Phycodnaviridae*, there were six eukaryotic

lineages retained after TIM filtration. Among these, Bacillariophyta, "other filosan (part of

filosan Cercozoa)", and Mamiellophyceae had comparatively strong connections. Moreover,

Rhodophyta, Ciliophora, and Dictyochophyceae had links to both *Mimiviridae* and

Phycodnaviridae. There was also a connection between *Iridoviridae* and Metazoa.

Associations between virophages and NCLDVs

Using 6818 NCLDV *polB* OTUs and 195 virophage major capsid proteins (MCPs),

we identified 535 FlashWeave associations (196 and 339 for pico- and femto-size fractions,

respectively) (Fig. 6A). Most of the associations had positive weights (n = 490), whereas

some had negative weights (n = 45). The average number of associations per virophage MCP

was different in two size fractions: 3.2 in femto- and 5.6 in pico-size fractions. The network

revealed that *Mimiviridae* had the largest number of virophage associations in both size

fractions. We also detected 84 positive associations between virophages and

Phycodnaviridae.

The phylogenetic tree defined three main virophage clades, and they were all found to

have many connections to NCLDVs. To investigate significant relationships, Fisher's exact

test was performed between virophage clades and NCLDV families. Families other than

Phycodnaviridae and *Mimiviridae* did not show significant associations. Therefore, we made

a group, "Other NCLDVs," to include all families except *Phycodnaviridae* and *Mimiviridae.*

First, we only used FlashWeave results with a weight > 0.4, as previous results showed that a

FlashWeave weight of 0.4 is a suitable cut-off that produced at least moderate performance

(Fig. 3A). From the femto-size fraction network, we found two significantly enriched

connections (Fig. 6B): one was between virophage group C and *Mimiviridae* ($p = 0.0022$)

and the other was between group A and "Other NCLDVs" ($p = 0.0439$). Another

significantly enriched relationship between virophage group B and *Phycodnaviridae* ($p =$

0.0410) was found in pico-size fractions when we used all associations without edge weight

cut-off.

Finally, we examined HGTs of virophage MCPs in NCLDV genomes. We found two

HGTs of virophage MCPs; both showed links between clade A and *Iridoviridae* (Table S3).

This result was consistent with Fisher's exact test, which revealed a connection between

virophage clade A and "Other NCLDVs", which includes *Iridoviridae*.

Discussion

NCLDVs can infect a wide range of eukaryotes, from unicellular to multicellular

organisms (33). However, we are still far from a comprehensive understanding of their hosts

because few have been isolated. Therefore, to better understand the ecological functions and

evolutionary significance of NCLDVs, reasonable host prediction is needed. To make these

predictions, we constructed global co-occurrence networks based on the marine metagenome

and metabarcoding datasets from 85 stations of the *Tara* Oceans expedition, which cover all

major oceanic provinces across an extensive latitudinal gradient from pole to pole. The edges

(associations) between *polB* and V9 nodes (OTUs) in the networks were generated by

FlashWeave. The networks were huge (Fig. 2A, S2A), suggesting that numerous NCLDVs

interact with eukaryotes in the ocean. This was expected given the high abundance and

diversity of NCLDVs in marine environments (17, 20) and the identification of HGT between

these viruses and diverse eukaryotic lineages (23). The networks were dominated by the

Mimiviridae nodes. This result is consistent with previous reports that *Mimiviridae* is the

most abundant and has the widest array of transcribed genes out of NCLDV families in marine environments (21, 22). *Mimiviridae* was known to infect amoebae, algae, and stramenopiles (3). In our study, these three eukaryotic groups were all found to have numerous associations with *Mimiviridae*. *Phycodnaviridae* has been known to infect many species of aquatic organisms, such as *Emiliania huxleyi* (Haptophyta), *Ectocarpus siliculosus* (Phaeophyceae), *Chlorella heliozoae* (Trebouxiophyceae), and *Ostreococcus tauri* (Mamiellophyceae) (34–36). Correspondingly, plenty of associations of *Phycodnaviridae* were found in the co-occurrence networks. For the eukaryotic nodes, all high taxonomic rank groups, including the SAR supergroup (i.e., Stramenopiles, Alveolata, Rhizaria), Opisthokonta, Archaeplastida, Amoebozoa, Excavata, and other eukaryotes, have associations with NCLDVs. Among these groups, the SAR supergroup contributed the most (~68%) *polB*–V9 associations. However, this is still lower than in other microbial co-occurrence analyses; for example, in aquatic microbial interactions, they dominated ~92% of the total associations (37). A substantially proportion (~32%) of NCLDV–eukaryote interactions were from non-SAR groups, which covered the known NCLDV host range, such as Archaeplastida and Haptophyta. Our result provides a list of candidate eukaryotes for NCLDV host prediction.

However, it is difficult to directly predict NCLDV hosts from constructed networks because of the high degree of associations per *polB* OTU (Fig. 3A). One node connected to multiple edges was expected in the co-occurrence analysis. In previous NCLDV host prediction studies, additional processing was performed to filter the high dimensional associations to predict the meaningful interactions, such as weight cut-off or a combination of different co-occurrence network inference methods (17, 27). Moreover, no previous study quantitively assessed the performance of co-occurrence networks when predicting NCLDV–host relationships. Qualitatively identifying known pairs and detecting HGT (without validation) have been commonly used to assess prediction reliability (17, 30). Therefore, we aimed to 1) quantitatively assess the performance of co-occurrence-based host prediction for NCLDVs and 2) improve the prediction results using filtering methods.

In a previous study of bacteriophage host prediction, ROC curves were used as an assessment metric to compare different prediction methods (38). However, the number of known virus–host pairs of NCLDVs is not sufficient to generate a dataset for ROC assessment. Therefore, in this study, we carried out an alternative method, the LR+, to assess the performance. LR+ is calculated with two relative values, sensitivity and specificity (Fig. 1). The LR+ of co-occurrence-based host predictions for positive associations was higher

than 1 and increased along with increasing cut-off values for the edge weights (Fig. 4A),

which demonstrated that positive prediction results were more likely to be true positives than

those based on random prediction. In high-weight regions (> 0.6 and > 0.4), LR+ values were

larger than 10 and 4, respectively. These LR+ values indicate that FlashWeave can increase

the probability of predicting true positives (39). However, both the true positive rate (<

18.9%) and false positive rate (< 6.37%) were very low. These low rates were from

FlashWeave with a cut-off of alpha < 0.01 (as described in the Materials and Methods),

which excluded a large proportion of the *polB*–V9 pairs from the results. The FDR of co-

occurrence was even higher than 90% (Fig. S4A). Such a high FDR in co-occurrence

networks demonstrates that condition positive connections (i.e., known interactions) are

embedded in an immense pool of condition negative connections. However, these negative

signals can correspond to unidentified (i.e., currently unknown true interactions), indirect, or

false relationships.

We also found that true positive predictions only existed in positive weight

associations, whereas negative weight associations did not contribute to NCLDV–host

detection (Fig. S4B). This result indicates that the abundance dynamics of NCLDVs and their

potential hosts were positively correlated with each other in the analyzed samples, which

were collected at a global scale; this might be because NCLDVs detected in the dataset were

active viruses that replicate locally in their hosts. Similar results were obtained in other co-

occurrence-based host prediction studies (30, 40). However, several experimental studies

showed that the abundance dynamics of NCLDVs and hosts showed a delay in time (28, 29).

It is possible that the global-scale samples did not have sufficiently high resolution to detect

negative correlations (or correlations with a time delay) due to lack of time-resolution (41).

Therefore, further studies, especially those that focus on a high temporal resolution, are

needed to better understand the detailed dynamics of virus–host associations and the capacity

of co-occurrence-based methods for host prediction.

The networks of different size fractions showed different performance patterns in

predicting NCLDV–host relationships (Fig. 4B). Generally, the associations from small-sized

networks performed better than the associations from large-sized fractions. To date, most of

the known NCLDV hosts are small, such as the genera *Prymnesium* (10 µm), *Aureococcus* (2

µm), and *Ostreococcus* (1 µm). Because of this, our assessment method might be biased

toward small size fractions. However, it is also possible that NCLDV infections are more

prevalent in smaller size fractions. Notably, the 0.8–inf-µm size fraction network, which

covered all four individual size fractions, performed best. This might be because NCLDVs

can infect not only small hosts but also hosts from a broad size range.

According to a previous study, trimming of low-abundance OTUs was recommended

to improve the prediction of true interactions (42). In our study, however, we did not achieve

such performance improvement by treating input abundance data with a rigorous filtration

(upper quartile) (Fig. S6). This result might be because the true positive and false positive

rates defined in this study were too low; therefore, the validation may not be sufficiently

sensitive to reflect the change between different abundance trimming strategies. However, it

is also possible that low-abundance NCLDV OTUs are indeed network participants, as was

demonstrated in a study showing that rare cyanobacterial species might play fundamental

roles in blooming (43). Our result also revealed that FlashWeave-S was better than

FlashWeave-HE and FastSpar at predicting NCLDV–host interactions (Fig, S7). The

difference between FlashWeave-HE and FlashWeave-S is that HE mode can remove

structural zeros during network inference. Structural zero is a typical property of

heterogeneous datasets, like *Tara* Oceans datasets, and may lead to false-positive edges (44).

Conversely, our results demonstrated that retaining structural zeros helped improve NCLDV–

eukaryote prediction, which indicates that the "presence–absence" pattern is more

informative than the "more–less" pattern when identifying NCLDV–host relationships. Our

result is consistent with a previous "K-r-strategist" hypothesis: NCLDVs, like mimiviruses,

are K-strategists that decay slowly and can form stable associations with their hosts (45). A

recent report supported these non-"boom and bust" dynamics of prasinoviruses and their

hosts with an experiment-based mathematical model (46). Overall, all these results show that

co-occurrence is an acceptable method for predicting NCLDV–host interactions in the marine

metagenome, and the LR+ is an effective quantitative metric for assessing the performance of

co-occurrence analysis.

Although the results generated by FlashWeave already increased the probability of

prediction, the condition positive interactions were still embedded in many noise edges, as

shown by the high FDR (Fig. S4A, S5B). We used a phylogeny-guided and network-based

association enrichment tool, TIM, to reduce the high dimension of associations and improve

NCLDV host prediction (32). The results showed that NCLDVs had enriched connections

with 14 major eukaryotic lineages, which included 24 taxonomic groups in three different

ranks (order, class, and phylum) (Table S4) (Fig. 5A, S7). Using the LR+ as a metric,

NCLDV host prediction improved 12-fold with TIM filtration (Fig. 5B). FDR dropped below

23% after TIM treatment (Fig. 5C). In TIM-enriched connections, some are known NCLDV–

host pairs, such as *Phycodnaviridae* and Mamiellophyceae, *Mimiviridae* and Haptophyta, and *Iridoviridae* and Metazoa. Some other studies revealed that *Mimiviridae* can exclusively infect diverse putative hosts (23, 47). Our results support this assumption that *Mimiviridae* has connections with 12 eukaryotic lineages out of 14 total lineages. Among the lineages, *Mimiviridae* had the most numerous links to Metazoa. Some mimiviruses (namaoviruses) are known to infect freshwater sturgeon, *Acipenser fulvescens* (48). Metazoans are presumed to be susceptible to mimiviruses, because the choanoflagellates, a group of eukaryotes that is phylogenetically close to metazoans, were recently proven to be the host of a species of *Mimiviridae* (14). Moreover, the TIM result revealed that *Phycodnaviridae* is closely connected to Bacillariophyta, which consists of three NCBI taxonomic groups: Thalassiophysales, Cymbellales, and Bacillariophyceae. Thalassiophysales was shown to have many HGT candidates with a large range of NCLDVs, and Bacillariophyceae also has a significant HGT candidate with phaeoviruses (23). Although Dictyochophyceae itself has not been proven to be a phycodnavirus host, its sister group Pelagophyceae was experimentally identified as an AaV host (49). Additionally, it is interesting to note the connection between Metazoa (Calanoida) and *Iridoviridae*. Calanoida is an order of arthropods commonly found as zooplankton; most of the sizes are 500–2000 μm. The viruses of the family *Iridoviridae* infect many Arthropod species, including insects and crustaceans (16).

Furthermore, we constructed the connections between virophages and NCLDVs. To date, all isolated virophages are only known to infect *Mimiviridae* (50). As expected, *Mimiviridae* was the family with dominant connections to virophages (Fig. 6A). Recently, *in silico* evidence demonstrated that virophages can infect *Phycodnaviridae*, which indicated that the virophage host range might be more extensive than we know (51). Consistently, a relatively large number of virophage OTUs were found to be associated with *Phycodnaviridae* in our study. The enrichment analysis also revealed significant connections between three virophage clades and NCLDV families (Fig. 6B). To support the enrichment analysis, we conducted an HGT analysis because gene transfers have previously been found between Sputnik virophages and giant viruses (52). Our HGT analysis indicated a previously unknown infectious relationship between virophage clade A and *Iridoviridae*. Overall, the results of virophage–NCLDV associations support our previous statement that co-occurrence is an appropriate tool for investigating NCLDV interactions in marine metagenomic data.

Materials and Methods

Metagenomic and metabarcoding data

The microbial metagenomic and eukaryotic metabarcoding data used in this study were previously generated from plankton samples collected by the *Tara* Oceans expedition from 2008 to 2013 (53, 54). Because our research requires paired metagenomic and metabarcoding datasets, we used data derived from the euphotic zone samples, namely those from the surface (SRF) and Deep Chlorophyll Maximum (DCM) layers (55). Type B DNA polymerase (*polB*) was used as the marker gene for NCLDVs. A total of 6818 NCLDV *polB* OTUs were extracted from the metagenomic datasets (i.e., the second version of the Ocean Microbial Reference Gene Catalog, OM-RGC.v2) using the pplacer phylogenetic placement method (56, 57). These *polB* sequences were classified into seven NCLDV families (*Mimiviridae*, *Phycodnaviridae*, *Marseilleviridae*, *Ascoviridae*, *Iridoviridae*, *Asfarviridae*, and *Poxviridae*) and two other giant virus groups ("Medusaviridae" and "Pithoviridae"). We employed the metabarcoding data for eukaryotes, which targets the 18S ribosomal RNA gene hypervariable V9 region (V9) (58). Taxonomic annotation of the eukaryotic metabarcoding data was previously performed by the *Tara* Oceans consortium with an extensive V9_PR2 reference database, which was derived from the original Protist Ribosomal Reference (PR2) database.

Data processing

A relative abundance matrix for the NCLDV *polB* OTUs was extracted from OM-RGC.v2 for the samples derived from the pico-size fractions (0.22–1.6 or 0.22–3.0 μm). We converted the relative abundances of *polB* OTUs back to absolute read counts based on gene length and read length (assumed to be 100 nt). This process was required because small decimal numbers cannot be used by FlashWeave and because relative abundance data suffer from apparent correlations, which reduce the specificity of co-occurrence networks in revealing microbial interactions (42). To build comprehensive interaction networks involving eukaryotes of different sizes, we extracted the V9 read count matrices from the metabarcoding dataset for the following five size fractions: 0.8–5 μm; 5–20 μm and 3–20 μm (hereafter referred to as "5–20 μm" for simplicity); 20–180 μm; 180–2000 μm; and > 0.8 μm (hereafter referred to as 0.8–inf μm). To create the input files for network inference, the *polB* matrix was combined with each of the V9 matrices (corresponding to different size fractions), and only the samples represented by both *polB* and V9 files were placed in new files. In total, samples from 84 *Tara* Oceans stations (a total of 560 samples for two depths and five size fractions) widely distributed across oceans were used in this study (Fig. S1A). Depending on the individual size fractions, 84–127 samples were retained and included in the co-occurrence

analysis (Fig. S1B). Read counts in the newly generated matrices were normalized using

centered log-ratio (*clr*) transformation after adding a pseudo count of one to all matrix

elements because zero cannot be transformed in *clr*. Following *clr* normalization, we filtered

out low-abundance OTUs with a lower quartile (Q1) filtering approach. Specifically, OTUs

were retained in the matrices when their *clr*-normalized abundance was higher than Q1

(among the non-zero counts in the original count matrix prior to the addition of a pseudo

count of one) in at least five samples. Normalization and filtering were separately applied to

polB and V9. The numbers of OTUs in the final matrices are provided in Fig. S1C.

Co-occurrence-based network inference

Network inference was performed using FlashWeave [v0.15.0 (44)]. FlashWeave is a

fast and compositionally robust tool for ecological network inference based on the local-to-

global learning framework. Meta-variables (such as environmental parameters) can be

included in the FlashWeave network to remove potential indirect associations. We used

temperature, salinity, nitrate, phosphate, and silicate concentrations as meta-variables in our

network inferences to determine their correlations with *polB* OTUs. FlashWeave has a

heterogeneous mode (FlashWeave-HE), which helps overcome sample heterogeneity.

However, FlashWeave-HE was not considered appropriate for the *Tara* Oceans data because

it was shown to predict an insufficient number of known planktonic interactions (44).

Therefore, we mainly used FlashWeave-S with default settings except for the FlashWeave

normalization step and comparison between FlashWeave-S and FlashWeave-HE. A threshold

to determine the statistical significance was set to alpha < 0.01. All detected pairwise

associations were assigned a value called "weight" that ranged between −1 and +1. Edges

with weights > 0 or < 0 were referred to as positive and negative associations, respectively.

To compare the performance of FlashWeave-S to other co-occurrence methods, we used

FlashWeave-HE and FastSpar (59). The FlashWeave-HE settings were the same as

FlashWeave-S but with a command "heterogeneous." In FastSpar, we used 50 iterations, 20

excluded iterations, and a threshold of 0.1 to generate associations. To reduce the high

dimensionality of the datasets, upper quartile (Q3) filtered matrices were used for

comparison.

Network validation

We validated the virus–host associations in inferred networks based on a confusion

matrix defined by the known NCLDV–host information (Fig. 1). Briefly, we manually

compiled 69 known virus–host relationships for NCLDVs (Table S1). In the validation
process, eukaryotic taxonomic groups were annotated at the level of the "Major lineages" in
the extensive PR2 database (updated after publication) (58). The "Major lineages" were used
in the present study because 1) the deficiency of known virus–host relationships limited the
use of lower eukaryotic taxonomy ranks, such as genus, for assessment, and 2) these lineages
adequately represented marine eukaryotes by covering the full spectrum of cataloged
eukaryotic V9 diversity at a comparable phylogenetic depth (58). Then, we performed
BLASTp [2.10.1 (60)] searches from the *Tara* Oceans PolB sequences against the NCLDV
reference database to define groups of metagenomic PolBs with a threshold of 65% sequence
identity by retaining only the best hit for each environmental PolB sequence. This threshold
was determined because, by using reference PolB sequences and RefSeq protein sequence
databases, we found that 60–70% of sequence identity could distinguish whether the
NCLDVs infected hosts of the same major lineages; this was mainly tested for
Phycodnaviridae because of the lack of host information for closely related viruses in other
NCLDV families (Table S2). Then, 65% was chosen because it could provide a better LR+
(as described below) than 60% and 70%.

The likelihood ratio was used in for assessment. This approach is commonly used in
diagnostic testing to assess if a test (host prediction in this study) usefully changes the
probability of the existence of condition positive (true positive). In this study, the LR+ was
used because host prediction is a test to discover condition positive states (61). LR+ is
calculated by dividing the true-positive rate (sensitivity) by the false-positive rate (1 –
specificity). If LR+ is close to 1, the performance of the prediction is comparable to a random
prediction. If LR+ >> 1, a positive prediction result is more likely to be a true positive than
that based on random prediction. From the set of detected associations between a given *polB*
OTU and V9 OTUs that belong to a given major eukaryotic lineage, we only kept the best
positive and negative associations (i.e., the edges with the highest absolute weights) to
simplify the prediction scheme. As an auxiliary assessment, the FDR was also calculated by
dividing the number of false positives by the number of positive predictions (Fig. 1).

Assessment of phylogeny-guided filtering for host prediction

We assessed the performance of TIM (https://github.com/RomainBlancMathieu/TIM)
(32) in improving host predictions by the FlashWeave co-occurrence approach. TIM is a
recently proposed tool for assigning the most likely associations based on the assumption that
evolutionarily related viruses tend to infect evolutionarily related hosts. TIM requires a

phylogenetic tree of viruses and a set of connections between viruses and eukaryotes, and

then tests whether a node of the virus tree is enriched with a specific predicted host group

compared with the rest of the tree using Fisher's exact test and Benjamini–Hochberg

adjustment (32).

We pooled network associations using FlashWeave analysis for five size fractions. To

build a concise and credible viral phylogenetic tree, we removed all of the PolB sequences

that were absent in the FlashWeave network associations, and the remaining sequences were

filtered by the amino acid sequence length ($\geq$ 500 aa). Protein alignment was conducted using

MAFFT-*linsi* [version 7.471 (62)], and 18 sequences were manually removed because they

were not well aligned with other PolB sequences. A total of 501 PolB sequences were used to

make a maximum likelihood phylogenetic tree with FastTree [version 2.1.11 (63)]. Then, the

PolB–V9 associations were mapped on the tree to calculate the significance of the enrichment

of specific associations using TIM. TIM provides a list of nodes in the viral tree and

associated NCBI taxonomies (order, class, and phylum) of eukaryotes that show a significant

enrichment in the leaves under the nodes. The TIM result was visualized with iTOL [version

5 (64)]. The TIM result was converted to a network, in which nodes correspond to the major

eukaryotic lineages. The network edge weight was defined by the number of tree nodes in

each viral family subtree enriched with a specific major eukaryotic lineage. The network was

visualized with Cytoscape [version 3.7.1] using prefuse force directed layout (65). To assess

the effectiveness of TIM in improving prediction, we extracted all the associations predicted

by TIM and compared their performance with the associations without TIM filtration.

Virophage–NCLDV associations

We inferred the networks between NCLDVs and virophages. *mcp* was used as the

marker gene of virophages. First, 47 reference MCP amino acid sequences were collected

from public databases and used to build an HMM profile. The HMM profile was used to

search against the amino acid sequences of OM-RGC v2 using HMMER hmmsearch [version

3.3.1] with the threshold of E-value < 1E−90 (66). This threshold was determined by

searching reference sequences against the GenomeNet nr-aa database. The search detected

195 *Tara* Oceans virophage MCP sequences in the OM-RGC database. Together with 47

reference MCPs, a phylogenetic tree of MCP amino acid sequences was built using MAFFT

and FastTree.

We extracted the abundance profiles for the 195 MCP sequences from the pico-

(0.22–1.6 or 0.22–3.0 µm) and femto-size (< 0.22 µm) fractions. We used samples from the

SRF and DCM depths. PolB and MCP abundance profiles were merged into two matrices
corresponding to the two virophage size fractions. Then, network inference was conducted
using the FlashWeave default settings after Q1 filtration. In the MCP phylogenetic tree, three
virophage clades contributed most of the NCLDV connections. Thus, an NCLDV enrichment
analysis for the three clades was carried out using Fisher's exact test, and the *p*-value was
adjusted by the Benjamin–Hochberg method. This approach was the same as TIM, but we did
not use the TIM software because TIM only works for eukaryotic nodes with NCBI
taxonomy annotations.

We used another approach, HGT, to predict the virophage-NCLDV interactions. First,
we generated an NCLDV genome database, which includes 56 reference NCLDV genomes
corresponding to our *polB* dataset and 2074 metagenome-assembled genomes from a
previous study (23). A total of 827548 coding sequences were included in this database.
Then, 195 virophage MCPs from the metagenomic data were BLASTp searched against this
database using an E-value cut-off of 1E−10 (with a minimum query coverage of 50% and a
minimum sequence identity of 50%). If a virophage MCP obtained a hit in the NCLDV
genome database with a lower E-value compared with hits in the MCP database (the hit to
itself was removed), the hit in the NCLDV genome database was considered an HGT
candidate.
561

Acknowledgments

This work was supported by JSPS/KAKENHI (Nos. 18H02279 and 19H05667 to H.O.
and Nos. 19K15895 and 19H04263 to H.E.), Kyoto University Research Coordination Alliance
(funding to H.E.), and the Collaborative Research Program of the Institute for Chemical
Research, Kyoto University (Nos. 2019-30 and 2020-27). Computational time was provided
by the SuperComputer System, Institute for Chemical Research, Kyoto University. We further
thank the *Tara* Oceans consortium, and the people and sponsors who supported *Tara* Oceans.
Tara Oceans (including both the *Tara* Oceans and *Tara* Oceans Polar Circle expeditions) would
not exist without the leadership of the *Tara* Expeditions Foundation and the continuous support
of 23 institutes (https://oceans.taraexpeditions.org). This article is contribution number XXX
of *Tara* Oceans. We thank Mallory Eckstut, PhD, from Edanz Group (https://en-author-
services.edanzgroup.com/ac) for editing a draft of this manuscript.

574
575

References

1.   Colson P, De Lamballerie X, Yutin N, Asgari S, Bigot Y, Bideshi DK, Cheng XW,

Federici BA, Van Etten JL, Koonin E V., La Scola B, Raoult D. 2013. "Megavirales",

a proposed new order for eukaryotic nucleocytoplasmic large DNA viruses. Arch Virol

158:2517–2521.

2.   Koonin E V., Dolja V V., Krupovic M, Varsani A, Wolf YI, Yutin N, Zerbini FM,

Kuhn JH. 2020. Global Organization and Proposed Megataxonomy of the Virus

World. Microbiol Mol Biol Rev 84:e00061-19.

3.   Koonin E V., Yutin N. 2019. Evolution of the Large Nucleocytoplasmic DNA Viruses

of Eukaryotes and Convergent Origins of Viral Gigantism. Advances in Virus

Research 103:167-202.

4.   Legendre M, Alempic JM, Philippe N, Lartigue A, Jeudy S, Poirot O, Ta NT, Nin S,

Couté Y, Abergel C, Claverie JM. 2019. Pandoravirus celtis illustrates the

microevolution processes at work in the giant Pandoraviridae genomes. Front

Microbiol 10:1–11.

5.   Boratto PVM, Oliveira GP, Machado TB, Andrade ACSP, Baudoin J-P, Klose T,

Schulz F, Azza S, Decloquement P, Chabrière E, Colson P, Levasseur A, La Scola B,

Abrahão JS. 2020. Yaravirus: A novel 80-nm virus infecting Acanthamoeba

castellanii. Proc Natl Acad Sci 117:16579–16586.

6.   Yoshikawa G, Blanc-Mathieu R, Song C, Kayama Y, Mochizuki T, Murata K, Ogata

H, Takemura M. 2019. Medusavirus, a Novel Large DNA Virus Discovered from Hot

Spring Water. J Virol 93:1–3.

7.   Forterre P. 2011. Manipulation of cellular syntheses and the nature of viruses: The

virocell concept. Comptes Rendus Chim 14:392–399.

8.   Mougari S, Sahmi-Bounsiar D, Levasseur A, Colson P, Scola B La. 2019. Virophages

of giant viruses: An update at eleven. Viruses 11:1–28.

9.   Koonin E V., Yutin N. 2010. Origin and evolution of eukaryotic large nucleo-

cytoplasmic DNA viruses. Intervirology 53:284–292.

10.   La Scola B, Audic S, Robert C, Jungang L, De Lamballerie X, Drancourt M, Birtles R,

Claverie JM, Raoult D. 2003. A giant virus in amoebae. Science 299:2033.

11.   Nagasaki K, Yamaguchi M. 1997. Isolation of a virus infectious to the harmful bloom

causing microalga Heterosigma akashiwo (Raphidophyceae). Aquat Microb Ecol

13:135–140.

12.   Moniruzzaman M, LeCleir GR, Brown CM, Gobler CJ, Bidle KD, Wilson WH,

Wilhelm SW. 2014. Genome of brown tide virus (AaV), the little giant of the

611   Megaviridae, elucidates NCLDV genome expansion and host-virus coevolution.

612   Virology 466–467:60–70.

613 13. Colson P, Gimenez G, Boyer M, Fournous G, Raoult D. 2011. The giant Cafeteria

614   roenbergensis virus that infects a widespread marine phagocytic protist is a new

615   member of the fourth domain of life. PLoS One 6:13–17.

616 14. Needham DM, Yoshizawa S, Hosaka T, Poirier C, Choi CJ, Hehenberger E, Irwin

617   NAT, Wilken S, Yung CM, Bachy C, Kurihara R, Nakajima Y, Kojima K, Kimura-

618   Someya T, Leonard G, Malmstrom RR, Mende DR, Olson DK, Sudo Y, Sudek S,

619   Richards TA, DeLong EF, Keeling PJ, Santoro AE, Shirouzu M, Iwasaki W, Worden

620   AZ. 2019. A distinct lineage of giant viruses brings a rhodopsin photosystem to

621   unicellular marine predators. Proc Natl Acad Sci U S A 116:20574–20583.

622 15. Williams T. 2008. Natural Invertebrate Hosts of Iridoviruses (Iridoviridae). Neotrop

623   Entomol 37:615–632.

624 16. Mihara T, Nishimura Y, Shimizu Y, Nishiyama H, Yoshikawa G, Uehara H, Hingamp

625   P, Goto S, Ogata H. 2016. Linking virus genomes with host taxonomy. Viruses 8:10–

626   15.

627 17. Hingamp P, Grimsley N, Acinas SG, Clerissi C, Subirana L, Poulain J, Ferrera I,

628   Sarmento H, Villar E, Lima-Mendez G, Faust K, Sunagawa S, Claverie JM, Moreau

629   H, Desdevises Y, Bork P, Raes J, De Vargas C, Karsenti E, Kandels-Lewis S, Jaillon

630   O, Not F, Pesant S, Wincker P, Ogata H. 2013. Exploring nucleo-cytoplasmic large

631   DNA viruses in Tara Oceans microbial metagenomes. ISME J 7:1678–1695.

632 18. Monier A, Claverie JM, Ogata H. 2008. Taxonomic distribution of large DNA viruses

633   in the sea. Genome Biol 9:R106.

634 19. Li Y, Hingamp P, Watai H, Endo H, Yoshida T, Ogata H. 2018. Degenerate PCR

635   primers to reveal the diversity of giant viruses in coastal waters. Viruses 10:496.

636 20. Mihara T, Koyano H, Hingamp P, Grimsley N, Goto S, Ogata H. 2018. Taxon richness

637   of "Megaviridae" exceeds those of bacteria and archaea in the ocean. Microbes

638   Environ 33:162–171.

639 21. Endo H, Blanc-Mathieu R, Li Y, Salazar G, Henry N, Labadie K, de Vargas C,

640   Sullivan MB, Bowler C, Wincker P, Karp-Boss L, Sunagawa S, Ogata H. 2020.

641   Biogeography of marine giant viruses reveals their interplay with eukaryotes and

642   ecological functions. Nat Ecol Evol https://doi.org/10.1038/s41559-020-01288-w.

643 22. Carradec Q, Pelletier E, Da Silva C, Alberti A, Seeleuthner Y, Blanc-Mathieu R,

644   Lima-Mendez G, Rocha F, Tirichine L, Labadie K, Kirilovsky A, Bertrand A, Engelen

S, Madoui MA, Méheust R, Poulain J, Romac S, Richter DJ, Yoshikawa G, Dimier C,

Kandels-Lewis S, Picheral M, Searson S, Acinas SG, Boss E, Follows M, Gorsky G,

Grimsley N, Karp-Boss L, Krzic U, Pesant S, Reynaud EG, Sardet C, Sieracki M,

Speich S, Stemmann L, Velayoudon D, Weissenbach J, Jaillon O, Aury JM, Karsenti

E, Sullivan MB, Sunagawa S, Bork P, Not F, Hingamp P, Raes J, Guidi L, Ogata H,

De Vargas C, Iudicone D, Bowler C, Wincker P. 2018. A global ocean atlas of

eukaryotic genes. Nat Commun 9:373.

23.   Schulz F, Roux S, Paez-Espino D, Jungbluth S, Walsh DA, Denef VJ, McMahon KD,

Konstantinidis KT, Eloe-Fadrosh EA, Kyrpides NC, Woyke T. 2020. Giant virus

diversity and host interactions through global metagenomics. Nature 578:432–436.

24.   Gallot-Lavallée L, Blanc G. 2017. A glimpse of nucleo-cytoplasmic large DNA virus

biodiversity through the eukaryotic genomicswindow. Viruses 9:17.

25.   Guglielmini J, Woo AC, Krupovic M, Forterre P, Gaia M. 2019. Diversification of

giant and large eukaryotic dsDNA viruses predated the origin of modern eukaryotes.

Proc Natl Acad Sci U S A 116:19585–19592.

26.   Cunha V Da, Gaïa M, Ogata H, Jaillon O, Delmont TO, Forterre P. 2020. Giant

viruses encode novel types of actins possibly related to the origin of eukaryotic actin:

the viractins. bioRxiv doi.org/10.1101/2020.06.16.150565.

27.   Roux S, Chan LK, Egan R, Malmstrom RR, McMahon KD, Sullivan MB. 2017.

Ecogenomics of virophages and their giant virus hosts assessed through time series

metagenomics. Nat Commun 8:858.

28.   Tomaru Y, Katanozaka N, Nishida K, Shirai Y, Tarutani K, Yamaguchi M, Nagasaki

K. 2004. Isolation and characterization of two distinct types of HcRNAV, a single-

stranded RNA virus infecting the bivalve-killing microalga Heterocapsa

circularisquama. Aquat Microb Ecol 34:207–218.

29.   Martínez Martínez J, Schroeder DC, Larsen A, Bratbak G, Wilson WH. 2007.

Molecular dynamics of Emiliania huxleyi and cooccurring viruses during two separate

mesocosm studies. Appl Environ Microbiol 73:554–562.

30.   Moniruzzaman M, Wurch LL, Alexander H, Dyhrman ST, Gobler CJ, Wilhelm SW.

2017. Virus-host relationships of marine single-celled eukaryotes resolved from

metatranscriptomics. Nat Commun 8:16054.

31.   Sunagawa S, Acinas SG, Bork P, Bowler C, Eveillard D, Gorsky G, Guidi L, Iudicone

D, Karsenti E, Lombard F, Ogata H, Pesant S, Sullivan MB, Wincker P, de Vargas C.

2020. Tara Oceans: towards global ocean ecosystems biology. Nat Rev Microbiol

18:428–445.

32.   Blanc-Mathieu R, Kaneko H, Endo H, Chaffron S, Hernández-Velázquez R, Nguyen CH, Mamitsuka H, Henry N, De Vargas C, Sullivan MB, Suttle CA, Guidi L, Ogata H. 2019. Viruses of the eukaryotic plankton are predicted to increase carbon export efficiency in the global sunlit ocean. bioRxiv https://doi.org/doi.org/10.1101/710228.

33.   Fischer MG. 2016. Giant viruses come of age. Curr Opin Microbiol 31:50–57.

34.   Delaroque N, Wolf S, Muller DG, Knippers R. 2000. The brown algal virus EsV-1 particle contains a putative hybrid histidine kinase. Virology 273:383–390.

35.   Fitzgerald LA, Graves M V., Li X, Hartigan J, Pfitzner AJP, Hoffart E, Van Etten JL. 2007. Sequence and annotation of the 288-kb ATCV-1 virus that infects an endosymbiotic chlorella strain of the heliozoon Acanthocystis turfacea. Virology 362:350–361.

36.   Clerissi C, Desdevises Y, Grimsley N. 2012. Prasinoviruses of the Marine Green Alga Ostreococcus tauri Are Mainly Species Specific. J Virol 86:4611–4619.

37.   Bjorbækmo MFM, Evenstad A, Røsæg LL, Krabberød AK, Logares R. 2020. The planktonic protist interactome: where do we stand after a century of research? ISME J 14:544–559.

38.   Edwards RA, McNair K, Faust K, Raes J, Dutilh BE. 2016. Computational approaches to predict bacteriophage-host relationships. FEMS Microbiol Rev 40:258–272.

39.   Salkić NN. 2008. Objective assessment of diagnostic tests validity : a short review for clinicians and other mortals . Part II. Acta Med Acad 39–42.

40.   Coutinho FH, Silveira CB, Gregoracci GB, Thompson CC, Edwards RA, Brussaard CPD, Dutilh BE, Thompson FL. 2017. Marine viruses discovered via metagenomics shed light on viral strategies throughout the oceans. Nat Commun 8:1–12.

41.   Martin-Platero AM, Cleary B, Kauffman K, Preheim SP, McGillicuddy DJ, Alm EJ, Polz MF. 2018. High resolution time series reveals cohesive but short-lived communities in coastal plankton. Nat Commun 9:266.

42.   Berry D, Widder S. 2014. Deciphering microbial interactions and detecting keystone species with co-occurrence networks. Front Microbiol 5:1–14.

43.   Xue Y, Chen H, Yang JR, Liu M, Huang B, Yang J. 2018. Distinct patterns and processes of abundant and rare eukaryotic plankton communities following a reservoir cyanobacterial bloom. ISME J 12:2263–2277.

44.   Tackmann J, Matias Rodrigues JF, von Mering C. 2019. Rapid Inference of Direct Interactions in Large-Scale Ecological Networks from Heterogeneous Microbial

Sequencing Data. Cell Syst 9:286-296.e8.

45. Suttle CA. 2007. Marine viruses - Major players in the global ecosystem. Nat Rev

Microbiol 5:801–812.

46. Yau S, Krasovec M, Felipe Benites L, Rombauts S, Groussin M, Vancaester E, Aury

JM, Derelle E, Desdevises Y, Escande ML, Grimsley N, Guy J, Moreau H, Sanchez-

Brosseau S, van de Peer Y, Vandepoele K, Gourbiere S, Piganeau G. 2020. Virus-host

coexistence in phytoplankton through the genomic lens. Sci Adv 6:eaay2587.

47. Claverie JM, Grzela R, Lartigue A, Bernadac A, Nitsche S, Vacelet J, Ogata H,

Abergel C. 2009. Mimivirus and Mimiviridae: Giant viruses with an increasing

number of potential hosts, including corals and sponges. J Invertebr Pathol 101:172–

180.

48. Clouthier SC, Vanwalleghem E, Copeland S, Klassen C, Hobbs G, Nielsen O,

Anderson ED. 2013. A new species of nucleo-cytoplasmic large DNA virus (NCLDV)

associated with mortalities in Manitoba lake sturgeon Acipenser fulvescens. Dis Aquat

Organ 102:195–209.

49. Gastrich MD, Leigh-Bell JA, Gobler CJ, Anderson OR, Wilhelm SW, Bryan M. 2004.

Viruses as potential regulators of regional brown tide blooms caused by the alga,

Aureococcus anophagefferens. Estuaries 27:112–119.

50. Duponchel S, Fischer MG. 2019. Viva lavidaviruses! five features of virophages that

parasitize giant DNA viruses. PLoS Pathog 15:1–7.

51. Xu S, Zhou L, Liang X, Zhou Y, Chen H, Yan S, Wang Y. 2020. Novel Cell-Virus-

Virophage Tripartite Infection Systems Discovered in the Freshwater Lake Dishui

Lake in Shanghai, China. J Virol 94:e00149-20.

52. Sun S, La Scola B, Bowman VD, Ryan CM, Whitelegge JP, Raoult D, Rossmann MG.

2010. Structural Studies of the Sputnik Virophage. J Virol 84:894–897.

53. Gregory AC, Zayed AA, Conceição-Neto N, Temperton B, Bolduc B, Alberti A,

Ardyna M, Arkhipova K, Carmichael M, Cruaud C, Dimier C, Domínguez-Huerta G,

Ferland J, Kandels S, Liu Y, Marec C, Pesant S, Picheral M, Pisarev S, Poulain J,

Tremblay JÉ, Vik D, Acinas SG, Babin M, Bork P, Boss E, Bowler C, Cochrane G, de

Vargas C, Follows M, Gorsky G, Grimsley N, Guidi L, Hingamp P, Iudicone D,

Jaillon O, Kandels-Lewis S, Karp-Boss L, Karsenti E, Not F, Ogata H, Poulton N,

Raes J, Sardet C, Speich S, Stemmann L, Sullivan MB, Sunagawa S, Wincker P,

Culley AI, Dutilh BE, Roux S. 2019. Marine DNA Viral Macro- and Microdiversity

from Pole to Pole. Cell 177:1109-1123.e14.

54.   Pesant S, Not F, Picheral M, Kandels-Lewis S, Le Bescot N, Gorsky G, Iudicone D,

Karsenti E, Speich S, Trouble R, Dimier C, Searson S. 2015. Open science resources

for the discovery and analysis of Tara Oceans data. Sci Data 2:150023.

55.   de Vargas C, Audic S, Henry N, Decelle J, Mahé F, Logares R, Lara E, Berney C, Le

Bescot N, Probert I, Carmichael M, Poulain J, Romac S, Colin S, Aury J-M, Bittner L,

Chaffron S, Dunthorn M, Engelen S, Flegontova O, Guidi L, Horák A, Jaillon O,

Lima-Mendez G, Lukeš J, Malviya S, Morard R, Mulot M, Scalco E, Siano R, Vincent

F, Zingone A, Dimier C, Picheral M, Searson S, Kandels-Lewis S, Acinas SG, Bork P,

Bowler C, Gorsky G, Grimsley N, Hingamp P, Iudicone D, Not F, Ogata H, Pesant S,

Raes J, Sieracki ME, Speich S, Stemmann L, Sunagawa S, Weissenbach J, Wincker P,

Karsenti E. 2015. Ocean plankton. Eukaryotic plankton diversity in the sunlit ocean.

Science 348:1261605.

56.   Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G, Djahanschiri

B, Zeller G, Mende DR, Alberti A, Cornejo-Castillo FM, Costea PI, Cruaud C,

D'Ovidio F, Engelen S, Ferrera I, Gasol JM, Guidi L, Hildebrand F, Kokoszka F,

Lepoivre C, Lima-Mendez G, Poulain J, Poulos BT, Royo-Llonch M, Sarmento H,

Vieira-Silva S, Dimier C, Picheral M, Searson S, Kandels-Lewis S, Boss E, Follows

M, Karp-Boss L, Krzic U, Reynaud EG, Sardet C, Sieracki M, Velayoudon D, Bowler

C, De Vargas C, Gorsky G, Grimsley N, Hingamp P, Iudicone D, Jaillon O, Not F,

Ogata H, Pesant S, Speich S, Stemmann L, Sullivan MB, Weissenbach J, Wincker P,

Karsenti E, Raes J, Acinas SG, Bork P. 2015. Ocean plankton. Structure and function

of the global ocean microbiome. Science 348:1261359.

57.   Matsen FA, Kodner RB, Armbrust EV. 2010. pplacer: linear time maximum-

likelihood and Bayesian phylogenetic placement of sequences onto a fixed reference

tree. BMC Bioinformatics 11:538.

58.   De Vargas C, Engelen S, Hingamp P, Sieracki M, Vargas C, Audic S, Henry N,

Decelle J, Mahé F, Logares R, Lara E, Berney C, Bescot N, Probert I, Carmichael M,

Poulain J, Romac S. 2015. Eukaryotic plankton diversity in the sunlit ocean. Science

348:1261605–1/11.

59.   Watts SC, Ritchie SC, Inouye M, Holt KE. 2019. FastSpar: Rapid and scalable

correlation estimation for compositional data. Bioinformatics 35:1064–1066.

60.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.

2009. BLAST+: Architecture and applications. BMC Bioinformatics 10:1–9.

61.   Yang WT, Parikh JR, Thomas Stavros A, Otto P, Maislin G. 2018. Exploring the

negative likelihood ratio and how it can be used to minimize false-positives in breast
imaging. Am J Roentgenol 210:301–306.

62. Katoh K, Misawa K, Kuma KI, Miyata T. 2002. MAFFT: A novel method for rapid
multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res
30:3059–3066.

63. Price MN, Dehal PS, Arkin AP. 2009. Fasttree: Computing large minimum evolution
trees with profiles instead of a distance matrix. Mol Biol Evol 26:1641–1650.

64. Letunic I, Bork P. 2019. Interactive Tree of Life (iTOL) v4: Recent updates and new
developments. Nucleic Acids Res 47:256–259.

65. Paul Shannon, Andrew Markiel, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,
Schwikowski B, Ideker T. 2003. Cytoscape: A Software Environment for Integrated
Models. Genome Res 13:426.

66. Eddy SR. 1998. Profile hidden Markov models. Bioinformatics 14:755–763.
794

Author contributions

796   LM and HO designed the study. LM performed most of the bioinformatics analysis.
797 HE generated the primary NCLDV data. RBM, RHV, and HK contributed to the
798 bioinformatics analysis. All authors contributed to the writing of the manuscript.
799

Materials & Correspondence

801   Correspondence and material requests should be addressed to HO (email:
802 ogata@kuicr.kyoto-u.ac.jp).
803

Competing financial interests

805   The authors declare no competing financial interests.

Figure 1. Overall workflow for inferring co-occurrence networks and quantitative assessment. This figure shows how the input data (*Tara* Oceans metagenomics and metabarcoding data) were used in this study. The definition of the confusion matrix for quantitative assessment is shown in the table. The LR+ and FDR equations are given at the lower right corner of the plot.

Figure 2. *polB*–V9 co-occurrence network. We performed co-occurrence analysis at the OTU level and constructed the network with pooled ***polB***–V9 associations from five size fraction networks (A). To better display co-occurrence patterns, PolB OTUs were grouped at the family or family-like level, and V9 OTUs were grouped using annotation at high taxonomic ranks. The size of each node indicates the number of OTUs that belong to the group, and the width of each edge indicates the number of associations between two connected groups. Associations with positive weight are shown in red and negative associations are shown in blue. (B) Number of associations connected to NCLDVs for each major eukaryotic lineage in five size fractions. The top 10 lineages were retained, and other lineages were omitted and shown as "others." Size fractions are presented in μm.

Figure 3. Density plots for the degree of NCLDV nodes in co-occurrence networks. The degree of an NCLDV node is given by the associations between this node and eukaryotes in the networks. The amount of NCLDV nodes are given on the top of the density values. (A) Positive degree (number of positive associations per node) for NCLDV nodes in five size fraction networks. (B) Negative degree (number of negative associations per node) for NCLDV nodes in five size fraction networks. Size fractions are presented in μm.

Figure 4. Positive likelihood ratios (LR+) in the NCLDV virus–host validation. (A) General performance of co-occurrence networks is shown with the LR+ calculated with associations pooled from five size fractions networks. To show the relationship between LR+ and FlashWeave association weight, the LR+ values are plotted along with the association weight. (B) Performance of each size fraction network is shown with the violin plot by ggplot2 with a bandwidth of 2. Size fractions are presented in μm.

Figure 5. Prediction of NCLDV virus–host relationships with TIM. (A) Undirected network that shows the relationships between NCLDVs and eukaryotes after TIM filtration. The size of each node indicates the number of predicted interactions of this group. The weight of network edges as defined by the number of tree nodes enriched in each viral family subtree to specific eukaryotic major lineages in the TIM analysis. Known virus–host relationships are highlighted in red, and the pairs found to have horizontal gene transfer are highlighted in yellow (1). (B) Performance of networks on NCLDV host prediction for original FlashWeave results without a weight cut-off, weight cut-off > 0.4, and TIM filtration, plotted by ggplot2 with a bandwidth of 2. (C) FDR of networks for NCLDV host prediction with the original FlashWeave results without a weight cut-off, weight cut-off > 0.4, and TIM filtration.

Figure 6. Associations between virophages and NCLDVs. (A) Number of associations with virophages is shown for seven NCLDV families and two unclassified groups, "Medusaviruses" and "Pithoviruses." Associations in the femto-size fraction network are shown in yellow, and in the pico-size fraction network are shown in green. The number of positive associations is above the zero axis, and the number of negative associations is below the zero axis. (B) Phylogenetic tree was constructed from 195 environmental virophages and 47 reference MCP sequences. The outside layer indicates three major virophage clades. The

865 inner two layers indicate that the virophage OTUs have at least one association with
866 NCLDVs in femto- or pico-size fractions networks.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.