1 **Title: Quantitative assessment of NCLDV–host interactions**
2 **predicted by co-occurrence analyses**
3
4 **Running title: NCLDV–host prediction based on co-occurrence**
5 **analysis**
6

7 Lingjie Meng[a], Hisashi Endo[a], Romain Blanc-Mathieu[a,b], Samuel Chaffron[c,d], Rodrigo
8 Hernández-Velázquez[a,e], Hiroto Kaneko[a], Hiroyuki Ogata[a,#]
9

Affiliations:
a. Bioinformatics Center, Institute for Chemical Research, Kyoto University, Gokasho, Uji,
611-0011, Japan
b. Laboratoire de Physiologie Cellulaire & Végétale, CEA, Univ. Grenoble Alpes, CNRS,
INRA, IRIG, Grenoble, France
c. Université de Nantes, CNRS UMR 6004, LS2N, F-44000 Nantes, France
d. Research Federation (FR2022) Tara Océan GO-SEE, Paris, France
e. Max Planck Institute for Marine Microbiology, Celsiusstraße 1, Bremen, Germany
18

19 [#]*Corresponding author*:
20 Hiroyuki Ogata (E-mail: ogata@kuicr.kyoto-u.ac.jp, Phone: +81-774-38-3270)
21

Abstract

Nucleocytoplasmic DNA viruses (NCLDVs) are highly diverse and abundant in
marine environments. However, knowledge of their hosts is limited because only a few
NCLDVs have been isolated so far. Taking advantage of the recent large-scale marine
metagenomics census, *in silico* host prediction approaches are expected to fill the gap and
further expand our knowledge of virus–host relationships for unknown NCLDVs. In this
study, we built co-occurrence networks of NCLDVs and eukaryotic taxa to predict virus–host
interactions using *Tara* Oceans sequencing data. Using the positive likelihood ratio to assess
the performance of host prediction for NCLDVs, we benchmarked several co-occurrence
approaches and demonstrated an increase in the odds ratio of predicting true positive
relationships four-fold compared with random host predictions. To further refine host
predictions from high-dimensional co-occurrence networks, we developed a phylogeny-
informed filtering method, Taxon Interaction Mapper, and showed it further improved the

prediction performance by twelve-fold. Finally, we inferred virophage – NCLDV networks to corroborate that co-occurrence approaches are effective for predicting interacting partners of NCLDVs in marine environments.

Importance

NCLDVs can infect a wide range of eukaryotes although their life cycle is less dependent on hosts compared with other viruses. However, our understanding of NCLDV–host systems is highly limited because few of these viruses have been isolated so far. Co-occurrence information has been assumed to be useful to predict virus–host interactions. In this study, we quantitatively show the effectiveness of co-occurrence inference for NCLDV host prediction. We also improve the prediction performance with a phylogeny-guided method, which leads to a concise list of candidate host lineages for three NCLDV families. Our results underpin the usage of co-occurrence approach for metagenomic exploration of the ecology of this diverse group of viruses.

Introduction

Nucleocytoplasmic large DNA viruses (NCLDVs) represent a group of double-stranded DNA viruses that belong to the viral phylum *Nucleocytoviricota* ([Virus Taxonomy: 2019 Release](#)), which was previously referred to as Megavirales (1, 2). NCLDVs usually possess diverse gene repertoires (74 to more than 2,000 proteins), large genomes (45 kb to 2.5 Mb), and outsized virions (80 nm to 1.5 μm) (3–5). NCLDVs have high functional autonomy and encode components of replication, transcription, and translation systems (3). Recently, a virus that belongs to a new family of NCLDVs called "Medusaviridae" was found to encode five types of histones (6). The existence of metabolically active viral factories and infectious virophages also indicates that the life cycle of NCLDVs is less dependent on host cells than other viruses (7, 8). To further understand what makes these giant viruses more independent than other viruses, a first crucial step is to identify their hosts — "Who infects whom?".

NCLDVs are known to infect a broad range of eukaryotes, from unicellular eukaryotes and macroalgae to animals (9). Amoebae are frequently used hosts in co-culture to isolate large NCLDVs (10). However, there is growing evidence, especially in marine systems, that NCLDVs can infect many phytoplankton groups, such as Pelagophyceae, Mamiellophyceae, Dinophyceae, and Haptophyte (11–13). Several other non-photosynthetic eukaryotic lineages, such as Bicoecea and Choanoflagellatea, were also reported as

experimentally identified NCLDV hosts in marine environments (14, 15). Small to large marine organisms, including invertebrates and vertebrates, are infected by viruses that belong to the NCLDV family *Iridoviridae* (16, 17). Together these studies indicate ubiquitous infectious relationships between NCLDVs and a wide range of marine eukaryotes. However, our understanding of NCLDV–host systems is very limited because few viruses have been isolated so far.

The number of viruses and hosts isolated in the laboratory represents a very small fraction of existing interactions in the ocean. Indeed, NCLDVs have been found to be highly diverse and abundant based on omics data (18, 19). In only a few liters of coastal seawater, more than 5,000 *Mimiviridae* species were detected; by comparison, only 20 *Mimiviridae* with known hosts have been well investigated (20). Global marine metagenomic data have revealed that the richness and phylogenetic diversity of NCLDVs are even higher than those of an entire prokaryotic domain (21). From biogeographical evidence, it is clear that these viruses are prevalent in the marine environment but have a heterogeneous community structure across sizes, depths, and biomes (22). Marine metatranscriptomic data have also shown that NCLDVs are active everywhere in sunlit oceans and may infect hosts from small piconanoplankton (0.8–5 µm) to large mesoplankton (180–2000 µm) (23).

Previous studies also demonstrated that NCLDVs have the potential to infect a greater diversity of hosts than known to date through gene transfer analyses (24, 25). NCLDVs might have started coevolving with eukaryotes even before the last eukaryotic common ancestor (LECA) (26). A recent study supported this hypothesis by showing that some NCLDVs encode viractins (actin-related genes in viruses), which could have been acquired from proto-eukaryotes and possibly reintroduced in the pre-LECA eukaryotic lineage (27). Together, these findings underline a lack of knowledge about NCLDV biology and host diversity. Therefore, more effort is needed to identify hosts to elucidate the poorly known virus–host relationships and the largely unknown NCLDV world.

Substantial effort has been made to reveal interactions between NCLDVs and their putative hosts. Apart from the co-culture method, other alternative methods, including high-throughput cell sorting, are also being used (10, 15). Metagenomics, which is particularly useful to assess a large fraction of ecosystem diversity, has been increasingly used to investigate NCLDVs host range. Comparative genomics analyses, such as identification of horizontal gene transfer (HGT) predictions, have also performed well for NCLDV host prediction (24, 25).

Abundance-based co-occurrence analyses have been used for host prediction and are
supposed to be effective because viruses can only thrive in an environment where their hosts
exist (18, 28). In addition to virus–host relationships, co-occurrence networks have been used
to predict the association between NCLDVs and their "parasites" (virophages) (29).
However, the co-occurrence-based prediction is also controversial for viral host prediction
since the abundance dynamics of viruses and their hosts (e.g., *Emiliania huxleyi* and
Heterosigma akashiwo viruses) are sometimes not concordant (30, 31). Usually, validation
with known virus–host relationships or corroboration with genomic evidence (e.g., HGT) is
used to assess network-based predictions (18, 28). However, the effectiveness of previous
and novel co-occurrence network methods has never been quantitatively tested for NCLDV
host prediction. The current lack of quantitative assessment hinders the widespread use of
this approach. Therefore, dedicated methods are needed to test the accuracy of NCLDV host
prediction with co-occurrence networks and to improve the performance of co-occurrence-
based predictions.

The *Tara* Oceans expedition is a global-scale survey on marine ecosystems that
expands our knowledge of microbial diversity, organismal interactions, and ecological
drivers of community structure (32). The present study used *Tara* Oceans metagenomic and
metabarcoding datasets to predict virus–host relationships between NCLDVs and eukaryotes
by constructing co-occurrence networks using different methods. To quantitatively assess the
performance of network-based host prediction, we employed the positive likelihood ratio
(LR+) using reference data for known NCLDV–host relationships. We developed a
phylogeny-based enrichment analysis approach, Taxon Interaction Mapper (TIM), to enhance
the performance in detecting positive signals in the intricate inferred networks. TIM has
previously been used in host predictions for DNA and RNA viruses (33), but without a
quantitative assessment on its effectiveness. In this study, we assessed the performance of
TIM as a filter of co-occurrence networks. We examined NCLDV–virophage networks,
which further justify the use of co-occurrence and filtering approaches to identify NCLDV
interaction partners.

Results

NCLDV–eukaryote co-occurrence networks

134   From five datasets that corresponded to five size fractions (Fig. S1), we generated five
135 co-occurrence networks on a global scale (Fig. 1, S2A). Altogether, these networks were

composed of 20,148 V9 and 5,234 *polB* OTUs (nodes) and 47,978 *polB*–V9 associations (edges). Out of these associations, 47,296 had positive weights, and 682 had negative weights (Fig. 2A). The associations that involved the family *Mimiviridae* were numerically dominant (n = 36,830) among the different NCLDV families. The second largest family was *Phycodnaviridae*, with 5,521 edges involving eukaryotes. No other family had more than 2,000 associations with eukaryotes. *Marseilleviridae*, forming the least associations in the networks, had 132 edges with eukaryotes. Taxonomic annotation of eukaryotic OTUs indicated that Alveolata, Opisthokonta, Rhizaria, and Stramenopiles were the major four eukaryotic groups connected to NCLDVs (with 21,167, 9,179, 6,521, and 5,327 edges, respectively). Three of these eukaryotic groups belong to the SAR supergroup (i.e., Stramenopiles, Alveolata, and Rhizaria), which represented 68.81% of the total associations. Regarding the pairs between viral families and eukaryotic lineages, *Mimiviridae* and Alveolata showed the largest number of edges (n = 16,548). Besides NCLDV–eukaryote associations, we detected 57,495 *polB*–*polB* associations and 234,448 V9–V9 associations (Fig. S2B). We also included environmental parameters in the network inference and identified 25 pairs of associations between environmental parameters and *polB* OTUs (Table S1).

The number of NCLDV–eukaryote associations generally decreased with enlarging size fraction (Fig. S2A). The largest number of *polB*–V9 associations were found in the 0.8–5-μm fraction (n = 10,647). Correspondingly, the eukaryotic community in 0.8–5-μm fraction had the greatest diversity (Fig. S3). However, the 0.8–inf-μm size fraction network was the largest (n = 10,477) for edges with positive weights. With the annotation of major lineages, the eukaryotic community compositions in the networks varied across different size fractions (Fig. 2B). In the smallest size fraction (0.8–5 μm) and the large range size fraction (0.8–inf μm), Marine Alveolate Group II was the eukaryotic lineage with the largest number of associations with NCLDVs (21.39% and 19.98%, respectively). Dinophyceae was the second largest group connected to NCLDVs in these two size fractions and showed the largest number of connections with NCLDVs in the 5–20-μm size fraction network (22.22% of total interactions). The viral associations with Metazoa and Collodaria increased with increasing size fractions. In the largest 180–2000-μm size fraction network, Metazoa contributed 39.31% of the total *polB*–V9 edges.

We calculated the degree of nodes (number of connected edges) for each NCLDV *polB* OTU (Fig. 3A, B). Naturally, the average degree of positive associations per *polB* was higher than negative edges in all size fractions and decreased along with increasing size fractions (2.69, 2.40, 2.25, and 2.10 from 0.8–5 μm to 180–2000 μm, and 2.76 for 0.8–inf).

170 Most of the *polB* nodes had more than one positive association (Fig. 3A). Together with the
171 taxonomic annotation of nodes, *polB*–V9 associations in the networks generated with the
172 *Tara* Oceans data revealed their high dimensionality and complexity.
173
Network validation
We quantitatively assessed the performance of predicting *polB*–V9 associations using
the positive likelihood ratio (LR+) (Fig. 1). By defining groups of metagenomic PolBs as
described in the Materials and Methods, 932 OTUs were recruited in the validation, and these
sequences contributed 6191 *polB*–V9 associations in the FlashWeave networks (Fig. S4). To
obtain an overall performance, we assessed the pooled associations (by removing
redundancy) from the five co-occurrence networks. LR+ was separately calculated for edges
with positive and negative weights because they can represent different infectious patterns.
As shown in Fig. 4A, the LR+ of host prediction for positive associations was higher than 1
(LR+ = 1 indicates no change in the likelihood of the condition). The LR+ generally
increased with the cut-off for FlashWeave weights, which indicated that condition positive
cases are enriched in the edges with higher weights. This result demonstrated that the co-
occurrence-based host prediction of NCLDVs outperformed random prediction (i.e., random
inference of virus–host pairs). In high-weight regions: 1) weight > 0.6, the LR+ of
associations was higher than 10; 2) weight > 0.4, the LR+ was roughly higher than 4.
Nonetheless, the false discovery rate (FDR) was high (Fig. S5A), which indicated that the
predictions contained numerous virus–host edges that were not considered condition positive.
FDR was 91.67% and 96.34% when weight is greater than 0.6 and 0.4, respectively. An
assessment of the host prediction for negative weight associations was also carried out. There
were no known NCLDV–host pairs found in the negative networks (Fig. S5B). The analysis
of the remaining part of our study was thus conducted for positive associations.
Comparing the performance between different size fractions indicated that the
networks of small size fractions (including the 0.8–inf-μm size fraction) performed better in
predicting the NCLDV–host relationships (Fig. 4B, S6). The 0.8–inf-μm size fraction had the
highest average LR+ out of the five size fractions (LR+ = 4.97). The LR+ of small size
fractions was generally higher than that of large size fractions, but there were exceptions
between 180–2000 and 20–180 μm. The LR+ of the associations in the 0.8–inf-μm, 0.8–5-μm
and 5–20-μm was greater than 1. Different from the average results, when the weight is
greater than 0.8, the associations of 5–20-μm size fraction had the best performance in terms
of both LR+ and FDR (Fig. S6 A, B).

We also compared abundance filtration strategies using Flashweave-S (sensitive

model) and FlashWeave-HE (heterogeneous model) but did not find a consistent pattern in

prediction performance (Fig. S7). The networks from the Q1 filtration strategy performed

best using Flashweave-S, but Q1 (lower quartile) filtration was not better than Q2 (middle

quartile) for Flashweave-HE inferred networks. Flashweave-S had a better performance than

HE model with any filtration strategy. Finally, we compared the performance of networks

inferred by all three methods: FlashWeave-S, FastSpar and Spearman. Three methods

generated a comparable number of positive associations, but FlashWeave-S made the largest

number of true positive predictions (Fig. S8A). Noteworthy, the LR+ of these three methods

were all larger than 1, however, FlashWeave-S and Spearman performed better than FastSpar

(Fig. S8B).

Assessment of host prediction improvement

Then we used the newly developed phylogeny-guided host prediction tool, TIM, to

filter *polB*–V9 associations, which is based on the assumption that evolutionarily related

viruses tend to infect evolutionarily related hosts (see Materials and Methods). We identified

24 eukaryotic taxonomic groups specifically associated with NCLDVs (Fig. S9). To compare

the performance of the TIM results with the above raw FlashWeave results, we converted the

three primary eukaryotic taxonomic ranks to their associated major lineages (Table S2), and

the associations were plotted as a network (Fig. 5A). This network showed that three out of

nine NCLDV families (*Mimiviridae, Phycodnaviridae*, and *Iridoviridae*) had enriched

connections in specific eukaryotic lineages. Among the network edges, known virus–host

pairs were found, such as Haptophyta–*Mimiviridae*, Mamiellophyceae–*Phycodnaviridae*, and

Metazoa–*Iridoviridae*. The associations in the TIM-filtered results showed a sharp

improvement in performance from the original result with and without an edge weight cut-

off. The average LR+ of TIM-enriched associations was 42.22, which was higher than the

raw FlashWeave associations without a weight cut-off (3.43), with a weight cut-off of 0.4

(5.20), and with a cut-off at 0.668 (14.23) (Fig. 5B, S9A). The FDR dropped from 0.97 (no

cut-off) and 0.95 (weight cut-off of 0.4) to 0.74 (Fig. 5C).

From the network, diverse putative hosts (13 lineages) emerged for *Mimiviridae,*

including algae, protozoans, and metazoans. Metazoa had the most enriched nodes connected

to *Mimiviridae*; additionally, MAST-3,12, Cryptophyta, Foraminifera, and Ciliophora had

strong relationships with *Mimiviridae*. For *Phycodnaviridae*, there were six eukaryotic

lineages retained after TIM filtration. Among these, Bacillariophyta, "other filosan (part of

filosan Cercozoa)”, and Mamiellophyceae had comparatively strong associations. Moreover,

Rhodophyta, Ciliophora, and Dictyochophyceae had links to both *Mimiviridae* and

Phycodnaviridae. There was also a connection between *Iridoviridae* and Metazoa.

Associations between virophages and NCLDVs

Using 6,818 NCLDV *polB* OTUs and 195 virophage major capsid proteins (MCPs),

we identified 535 FlashWeave associations (196 and 339 for pico- and femto-size fractions,

respectively) (Fig. 6A). Most of the associations had positive weights ($n = 490$), whereas

some had negative weights ($n = 45$). The average number of associations per virophage MCP

was different in two size fractions: 3.2 in femto- and 5.6 in pico-size fractions. The network

revealed that *Mimiviridae* had the largest number of virophage associations in both size

fractions. We also detected 84 positive associations between virophages and

Phycodnaviridae.

The phylogenetic tree defined three main virophage clades, and they were all found to

have many connections to NCLDVs. To investigate significant relationships, Fisher's exact

test was performed between virophage clades and NCLDV families. Families other than

Phycodnaviridae and *Mimiviridae* did not show significant associations. Therefore, we made

a group, “Other NCLDVs,” to include all families except *Phycodnaviridae* and *Mimiviridae*.

First, we only used FlashWeave results with a weight > 0.4, as previous results showed that a

FlashWeave weight of 0.4 is a suitable cut-off that produced moderate performance (Fig.

3A). From the femto-size fraction network, we found two significantly enriched connections

(Fig. 6B): one was between virophage group C and *Mimiviridae* ($p = 0.0022$) and the other

was between group A and “Other NCLDVs” ($p = 0.0439$). Another significantly enriched

relationship between virophage group B and *Phycodnaviridae* ($p = 0.0410$) was found in

pico-size fractions when we used all associations without edge weight cut-off.

Finally, we examined HGTs of virophage MCPs in NCLDV genomes. We found two

HGTs of virophage MCPs; both showed links between clade A and *Iridoviridae* (Table S3).

This result was consistent with the Fisher's exact test result, which revealed a connection

between virophage clade A and “Other NCLDVs” including *Iridoviridae*.

Discussion

NCLDVs can infect a wide range of eukaryotes, from unicellular to multicellular

organisms (34). However, we are still far from a comprehensive knowledge of their hosts

because few have been isolated so far. Therefore, better host prediction algorithms are needed

to understand the ecological functions and evolutionary significance of NCLDVs. To make

these predictions, we constructed global ocean co-occurrence networks based on the marine

metagenome and metabarcoding datasets from 85 stations of the *Tara* Oceans expedition,

which cover all major oceanic provinces across an extensive latitudinal gradient from pole to

pole. The edges (associations) between *polB* and V9 nodes (OTUs) in the networks were

generated using FlashWeave. The networks were particularly dense (Fig. 2A, S2A), thus

suggesting that NCLDVs interact with numerous eukaryotes in the ocean. This was expected

given the high abundance and diversity of NCLDVs in marine environments (18, 21) and the

identification of HGT between these viruses and diverse eukaryotic lineages (24). The

networks were dominated by the *Mimiviridae* nodes, which is consistent with previous

reports that *Mimiviridae* is the most abundant and has the widest array of transcribed genes

out of NCLDV families in marine environments (22, 23). *Mimiviridae* was known to infect

amoebae, algae, and stramenopiles (3). In our study, these three eukaryotic groups were all

found to have numerous associations with *Mimiviridae*. *Phycodnaviridae* has been known to

infect many species of aquatic organisms, such as *Emiliania huxleyi* (Haptophyta),

Ectocarpus siliculosus (Phaeophyceae), *Chlorella heliozoae* (Trebouxiophyceae), and

Ostreococcus tauri (Mamiellophyceae) (35–37). Correspondingly, plenty of associations of

Phycodnaviridae were found in the co-occurrence networks. For the eukaryotic nodes, all

high taxonomic rank groups, including the SAR supergroup (i.e., Stramenopiles, Alveolata,

Rhizaria), Opisthokonta, Archaeplastida, Amoebozoa, Excavata, and other eukaryotes, have

associations with NCLDVs. Among these groups, the SAR supergroup contributed the most

(~68%) *polB*–V9 associations. However, this is still lower than in other microbial co-

occurrence analyses; for example, a previous study showed SAR supergroup dominated

~92% of the total aquatic microbial associations (38). A substantial proportion (~32%) of

NCLDV–eukaryote interactions were from non-SAR groups, which covered the known

NCLDV host range, such as Archaeplastida and Haptophyta.

However, it is difficult to accurately predict NCLDV hosts from constructed networks

because of the high degree of associations per *polB* OTU (Fig. 3A). One node connected to

multiple edges was expected in the co-occurrence analysis. In previous NCLDV host

prediction studies, additional processing was performed to filter the high dimensional

associations to predict the meaningful interactions, such as weight cut-off or a combination of

different co-occurrence network inference methods (18, 29). Moreover, no previous study

quantitively assessed the performance of co-occurrence networks when predicting NCLDV–

host relationships. Qualitatively identifying known pairs and detecting HGT (without

validation) have been commonly used to assess prediction reliability (18, 28). Therefore, we

aimed to 1) quantitatively assess the performance of co-occurrence-based host prediction for

NCLDVs and 2) improve the prediction results using filtering methods.

In a previous study of bacteriophage host prediction, ROC curves were used as an

assessment metric to compare different prediction methods (39). However, the number of

known virus–host pairs of NCLDVs is not sufficient to generate a dataset for ROC

assessment. Therefore, in this study, we carried out an alternative method, the LR+, to assess

the performance. LR+ is calculated with two relative values, sensitivity and specificity (Fig.

1). The LR+ of co-occurrence-based host predictions for positive associations was higher

than 1 and increased along with increasing cut-off values for the edge weights (Fig. 4A),

which demonstrated that positive prediction results were more likely to be true positives than

those based on random prediction. In high-weight regions (> 0.6 and > 0.4), LR+ values were

larger than 10 and 4, respectively. These LR+ values indicate that FlashWeave can increase

the probability of predicting true positives (40). However, both the true positive rate

(sensitivity, < 18.9%) and false positive rate (< 6.37%) were very low (Fig. S5C). These low

rates were from FlashWeave with a cut-off of alpha < 0.01, which excluded a large

proportion of the *polB*–V9 pairs from the results. So only about 4000 predictions could be

validated from a set of 6191 *polB*–V9 FlashWeave associations in this study (Fig. S5D, as

described in the Materials and Methods). The FDR of co-occurrence was even higher than

90% (Fig. S5A). Such a high FDR in co-occurrence networks demonstrates that condition

positive connections (i.e., known interactions) are embedded in an immense pool of condition

negative connections. However, these negative signals can correspond to either unidentified

(i.e., currently unknown true interactions), indirect, or false relationships.

We also found that true positive predictions only existed in positive weight

associations, whereas negative weight associations did not contribute to NCLDV–host

detection (Fig. S5B). This result indicates that the abundance dynamics of NCLDVs and their

potential hosts were positively correlated with each other in the analyzed samples, which

were collected at a global scale; this might be because NCLDVs detected in the dataset were

active viruses that replicate locally in their hosts. Similar results were obtained in other co-

occurrence-based host prediction studies (28, 41). However, several experimental studies

showed that the abundance dynamics of NCLDVs and hosts showed a delay in time (30, 31).

It is possible that the global-scale samples did not have sufficiently high resolution to detect

negative correlations (or correlations with a time delay) due to lack of time-resolution (42).

Therefore, further studies, especially those that focus on a high temporal resolution, are

needed to better understand the detailed dynamics of virus–host associations and the capacity
of co-occurrence-based methods for host prediction.

The networks of different size fractions showed different performance patterns in
predicting NCLDV–host relationships (Fig. 4B). This pattern is not dependent on the
diversity of eukaryotic communities (Fig. S3A, B). Generally, small-sized fractions (0.8–5-
µm and 5–20-µm) networks performed better than large-sized fractions (20-180-µm and 180-
2000-µm) networks. This result is not dependent on the diversity of eukaryotic communities.
To date, most of the known NCLDV hosts are small, such as the genera *Micromonas*,
Aureococcus, and *Ostreococcus* are within the range of 0.8–5-µm, and *Prymnesium*,
Heterosigma, and *Heterocapsa* are within the range of 5–20-µm. Because of this, our
assessment method might be biased toward small size fractions as smaller organisms tend to
be more abundant in the environment (43). However, it is also possible that NCLDV
infections are more prevalent in smaller size fractions. Notably, the 0.8–inf-µm size fraction
network, which covered all four individual size fractions, performed best. This might be
because NCLDVs can infect not only small hosts but also hosts from a broad size range.

Trimming of low-abundance OTUs was recommended to improve the prediction of
true interactions and was often used in co-occurrence studies (44, 45). In our study, however,
we did not achieve such performance improvement by treating input abundance data with a
rigorous filtration (upper quartile) (Fig. S7). This result might be because the true positive
and false positive rates defined in this study were too low; therefore, the validation may not
be sufficiently sensitive to reflect the change between different abundance trimming
strategies. However, it is also possible that low-abundance NCLDV OTUs are indeed
network participants, as was demonstrated in a study showing that rare cyanobacterial species
might play fundamental roles in blooming (46). Our result also revealed that FlashWeave-S
was better than FlashWeave-HE at predicting NCLDV–host interactions (Fig. S7). The
difference between FlashWeave-HE and FlashWeave-S is that HE mode can remove
structural zeros during network inference. Structural zero is a typical property of
heterogeneous datasets, like *Tara* Oceans datasets, and may lead to false-positive edges (47).
Conversely, our results suggested that retaining structural zeros did not negatively influence
the result, which indicates that the "presence–absence" pattern is as informative as the
"more–less" pattern when identifying NCLDV–host relationships. This result is consistent
with a previous "K-r-strategist" hypothesis: some NCLDVs, like mimiviruses, are K-
strategists that decay slowly and can form stable associations with their hosts (48, 49). A
recent report supported these non-"boom and bust" dynamics of prasinoviruses and their

hosts with an experiment-based mathematical model (50). Overall, our results support co-
occurrence networks as a useful method for predicting NCLDV–host interactions in marine
metagenomes, and likelihood ratios as useful quantitative metrics for assessing the
performance of co-occurrence analysis for viral host predictions.
Although the results generated by FlashWeave already improved the accuracy of
predictions, the condition positive interactions were still embedded in many noise edges, as
shown by a very high FDR (Fig. S5A, S6B). To overcome this situation, we developed TIM
to reduce the high dimension of associations and improve NCLDV host prediction (33). The
results showed that NCLDVs had enriched connections with 15 major eukaryotic lineages,
which included 24 taxonomic groups in three different ranks (order, class, and phylum)
(Table S2) (Fig. 5A, S7). Using the LR+ as a prediction diagnostic metric, NCLDV host
prediction improved 12-fold with TIM filtration (Fig. 5B). FDR dropped below 23% after
TIM treatment (Fig. 5C). In TIM-enriched connections, some are known NCLDV–host pairs,
such as *Phycodnaviridae* and Mamiellophyceae, *Mimiviridae* and Haptophyta, and
Iridoviridae and Metazoa. Some other studies revealed that *Mimiviridae* could exclusively
infect diverse putative hosts (24, 51). Our results support the assumption that *Mimiviridae* has
connections with 13 eukaryotic lineages out of 15 total lineages. Among these lineages,
Mimiviridae had the most numerous links to Metazoa. Some mimiviruses (namaoviruses) are
known to infect freshwater sturgeon, *Acipenser fulvescens* (52). Metazoans are presumed to
be susceptible to mimiviruses, because the choanoflagellates, a group of eukaryotes that is
phylogenetically close to metazoans, were recently identified to be the host of a species of
Mimiviridae (15). Moreover, the TIM result revealed that *Phycodnaviridae* is closely
connected to Bacillariophyta, which consists of three NCBI taxonomic groups:
Thalassiophysales, Cymbellales, and Bacillariophyceae. Thalassiophysales was shown to
have many HGT candidates with a large range of NCLDVs, and Bacillariophyceae also has a
significant HGT candidate with phaeoviruses (24). Although Dictyochophyceae itself has not
been proven to be a phycodnavirus host, its sister group Pelagophyceae was experimentally
identified as an AaV host (53). Additionally, it is interesting to note the connection between
Metazoa (Calanoida) and *Iridoviridae*. Calanoida is an order of arthropods commonly found
as zooplankton; most of the sizes are 500–2000 μm. The viruses of the family *Iridoviridae*
infect many Arthropod species, including insects and crustaceans (17).
Furthermore, we also inferred associations between virophages and NCLDVs. To
date, all isolated virophages are only known to infect *Mimiviridae* (54). As expected,
Mimiviridae was the family with dominant connections to virophages (Fig. 6A). Recently, *in*

silico evidence demonstrated that virophages can infect *Phycodnaviridae*, which indicated that the virophage host range might be larger than we know (55). In support of this hypothesis, a relatively large number of virophage OTUs were found to be associated with *Phycodnaviridae* in our study. The enrichment analysis also revealed significant connections between three virophage clades and NCLDV families (Fig. 6B). To support the enrichment analysis, we conducted an HGT analysis because gene transfers have previously been found between Sputnik virophages and giant viruses (56). Our HGT analysis indicated a previously undescribed infectious relationship between virophage clade A and *Iridoviridae*. Overall, the results of virophage–NCLDV associations support our previous statement that co-occurrence networks inference and analysis are appropriate for investigating NCLDV interactions in marine metagenomic data.

Materials and Methods

Metagenomic and metabarcoding data

The microbial metagenomic and eukaryotic metabarcoding data used in this study were previously generated from plankton samples collected by the *Tara* Oceans expedition from 2008 to 2013 (57, 58). Because our research requires paired metagenomic and metabarcoding datasets, we used data derived from the euphotic zone samples, namely those from the surface (SRF) and Deep Chlorophyll Maximum (DCM) layers (59). Type B DNA polymerase (*polB*) was used as the marker gene for NCLDVs. A total of 6818 NCLDV *polB* OTUs were extracted from the metagenomic datasets (i.e., the second version of the Ocean Microbial Reference Gene Catalog, OM-RGC.v2) using the pplacer phylogenetic placement method (ML tree) (22, 60, 61). These *polB* sequences were classified into seven NCLDV families (*Mimiviridae*, *Phycodnaviridae*, *Marseilleviridae*, *Ascoviridae*, *Iridoviridae*, *Asfarviridae*, and *Poxviridae*) and two other giant virus groups ("Medusaviridae" and "Pithoviridae"). For eukaryotes, we employed used the metabarcoding data for eukaryotes, which targeting the 18S ribosomal RNA gene hypervariable V9 region (V9) (62). Taxonomic annotation of the eukaryotic metabarcoding data was previously performed by the *Tara* Oceans consortium using an extensive V9_PR2 reference database (59), which was derived from the original Protist Ribosomal Reference (PR2) database (63). The diversity index of eukaryotic communities was calculated using the package "vegan" (64). Processed frequency data are available from GenomeNet (ftp://ftp.genome.jp/pub/db/community/tara/Cooccurrence).

Data processing

A relative abundance matrix for the NCLDV *polB* OTUs was extracted from OM-RGC.v2 for the samples derived from the pico-size fractions (0.22–1.6 or 0.22–3.0 μm). We converted the relative abundances of *polB* OTUs back to absolute read counts based on gene length and read length (assumed to be 100 nt). This process was required because small decimal numbers cannot be used by FlashWeave and because relative abundance data suffer from apparent correlations, which reduce the specificity of co-occurrence networks in revealing microbial interactions (44). To build comprehensive interaction networks involving eukaryotes of different sizes, we extracted the V9 read count matrices from the metabarcoding dataset for the following five size fractions: 0.8–5 μm; 5–20 μm and 3–20 μm (hereafter referred to as "5–20 μm" for simplicity); 20–180 μm; 180–2000 μm; and > 0.8 μm (hereafter referred to as 0.8–inf μm). To create the input files for network inference, the *polB* matrix was combined with each of the V9 matrices (corresponding to different size fractions), and only the samples represented by both *polB* and V9 files were placed in new files. In total, samples from 84 *Tara* Oceans stations (a total of 560 samples for two depths and five size fractions) widely distributed across oceans were used in this study (Fig. S1A). Depending on the individual size fractions, 84–127 samples were retained and included in the co-occurrence analysis (Fig. S1B). Read counts in the newly generated matrices were normalized using centered log-ratio (*clr*) transformation after adding a pseudo count of one to all matrix elements because zero cannot be transformed in *clr*. Following *clr* normalization, we filtered out low-abundance OTUs with a lower quartile (Q1) filtering approach. Specifically, OTUs were retained in the matrices when their *clr*-normalized abundance was higher than Q1 (among the non-zero counts in the original count matrix prior to the addition of a pseudo count of one) in at least five samples. Normalization and filtering were separately applied to *polB* and V9. The numbers of OTUs in the final matrices are provided in Fig. S1C.

Co-occurrence-based network inference

Network inference was performed using FlashWeave [v0.15.0 (47)]. FlashWeave is a fast and compositionally robust tool for ecological network inference based on the local-to-global learning framework. Meta-variables (such as environmental parameters) can be included in the FlashWeave network to remove potential indirect associations. We used temperature, salinity, nitrate, phosphate, and silicate concentrations as meta-variables in our network inferences to determine their correlations with *polB* OTUs. FlashWeave provides a heterogeneous mode (FlashWeave-HE), which helps overcome sample heterogeneity.

However, FlashWeave-HE may not be appropriate for the *Tara* Oceans data because it was

shown to predict an insufficient number of known planktonic interactions (47). Therefore, we

mainly used FlashWeave-S with default settings except for the FlashWeave normalization

step and comparison between FlashWeave-S and FlashWeave-HE. A threshold to determine

the statistical significance was set to alpha < 0.01. All detected pairwise associations were

assigned a value called "weight" that ranged between −1 and +1. Edges with weights > 0 or <

0 were referred to as positive and negative associations, respectively. To compare the

performance of FlashWeave-S to other co-occurrence methods, we used FlashWeave-HE,

Spearman, and FastSpar (65). The FlashWeave-HE settings were the same as FlashWeave-S

but with a command "heterogeneous". For Spearman, we used stats.spearmanr in package

"Scipy" (66). In FastSpar, we used 50 iterations, 20 excluded iterations, and a threshold of

0.1 to generate associations. To reduce the high dimensionality of the datasets, upper quartile

(Q3) filtered matrices were used for comparison among FlashWeave-S, Spearman, and

FastSpar.

Network validation

We validated the virus–host associations in inferred networks based on a confusion

matrix defined by the known NCLDV–host information (Fig. 1). Briefly, we manually

compiled 69 known virus–host relationships for NCLDVs (Table S4). In the validation

process, eukaryotic taxonomic groups were annotated at the level of the "Major lineages" in

the extensive PR2 database (updated after publication) (62). The "Major lineages" were used

in the present study because 1) the deficiency of known virus–host relationships limited the

use of lower eukaryotic taxonomy ranks, such as genus, for assessment, and 2) these lineages

adequately represented marine eukaryotes by covering the full spectrum of cataloged

eukaryotic V9 diversity at a comparable phylogenetic depth (62). Then, we performed

BLASTp [2.10.1 (67)] searches from the *Tara* Oceans PolB sequences against the NCLDV

reference database to define groups of metagenomic PolBs with a threshold of 65% sequence

identity by retaining only the best hit for each environmental PolB sequence. This threshold

was determined because, by using reference PolB sequences and RefSeq protein sequence

databases, we found that 60–70% of sequence identity could distinguish whether the

NCLDVs infected hosts of the same major lineages; this was mainly tested for

Phycodnaviridae because of the lack of host information for closely related viruses in other

NCLDV families (Table S5). Then, 65% was chosen because it could provide a better LR+

(as described below) than 60% and 70%.

The positive likelihood ratio was used in for assessment to estimate the predictions

accuracy. This approach is commonly used in diagnostic testing to assess if a test (host

prediction in this study) usefully changes the probability of the existence of condition

positive (true positive). In this study, the LR+ was used because host prediction is a test to

discover condition positive states (68). LR+ is calculated by dividing the true-positive rate

(sensitivity) by the false-positive rate (1 –specificity). If LR+ is close to 1, the performance of

the prediction is comparable to a random prediction. If LR+ >> 1, a positive prediction result

is more likely to be a true positive than that based on random prediction. From the set of

detected associations between a given *polB* OTU and V9 OTUs that belong to a given major

eukaryotic lineage, we only kept the best positive and negative associations (i.e., the edges

with the highest absolute weights) to simplify the prediction scheme. As an auxiliary

assessment, the FDR was also calculated by dividing the number of false positives by the

number of positive predictions (Fig. 1). For the comparison among five size fractions, we

only used the abundance in the overlap samples of 0.8–5 μm, 5–20 μm, 20–180 μm, and

180–2000 μm sizes. So the number of samples in five size fractions is comparable (84, 88,

88, 88, 88), which could reduce the bias that may influence the topology of networks (69).

Phylogeny-guided filtering of host predictions and its assessment

We developed Taxon Interaction Mapper (TIM) to improve host predictions by co-

occurrence approaches (33). TIM assumes that evolutionarily related viruses tend to infect

evolutionarily related hosts (17, 70), and extract the most likely virus-host associations from

the co-occurrence networks. TIM requires a phylogenetic tree of viruses (based on marker

genes) and a set of connections between viruses and eukaryotes (co-occurrence edges), and

then tests whether leaves (i.e., viral OTUs) under a node of the virus tree is enriched with a

specific predicted host group compared with the rest of the tree using Fisher's exact test and

Benjamini–Hochberg adjustment (Fig. S9A) (33). TIM is available from

https://github.com/RomainBlancMathieu/TIM.

We pooled network associations using FlashWeave analysis for five size fractions. To

build a concise and credible viral phylogenetic tree, we removed all of the PolB sequences

that were absent in the FlashWeave network associations, and the remaining sequences were

filtered by the amino acid sequence length (≥ 500 aa). Protein alignment was conducted using

MAFFT-*linsi* [version 7.471 (71)], and 18 sequences were manually removed because they

were not well aligned with other PolB sequences. A total of 501 PolB sequences were used to

make a maximum likelihood phylogenetic tree with FastTree [version 2.1.11 (72)]. Then, the

PolB–V9 associations were mapped on the tree to calculate the significance of the enrichment

of specific associations using TIM. TIM provides a list of nodes in the viral tree and

associated NCBI taxonomies (order, class, and phylum) of eukaryotes that show significant

enrichment in the leaves under the nodes. The TIM result was visualized with iTOL [version

5 (73)]. The TIM result was converted to a network, in which nodes correspond to the major

eukaryotic lineages. The network edge weight was defined by the number of tree nodes in

each viral family subtree enriched with a specific major eukaryotic lineage. The network was

visualized with Cytoscape [version 3.7.1] using prefuse force directed layout (74). To assess

the effectiveness of TIM in improving prediction, we extracted all the associations predicted

by TIM and compared their performance with the raw and weight cut-off results.

Virophage–NCLDV associations

We inferred the networks between NCLDVs and virophages using. *mcp* was used as

the marker gene for virophages. First, 47 reference MCP amino acid sequences were

collected from public databases and used to build an HMM profile. The HMM profile was

used to search against the amino acid sequences of OM-RGC v2 using HMMER hmmsearch

[version 3.3.1] with the threshold of E-value $< 1E-90$ (75). This threshold was determined by

searching reference sequences against the GenomeNet nr-aa database. The search detected

195 *Tara* Oceans virophage MCP sequences in the OM-RGC database. Together with 47

reference MCPs, a phylogenetic tree of MCP amino acid sequences was built using MAFFT

and FastTree.

We extracted the abundance profiles for the 195 MCP sequences from the pico-

(0.22–1.6 or 0.22–3.0 µm) and femto-size (< 0.22 µm) fractions. We used samples from the

SRF and DCM depths. PolB and MCP abundance profiles were merged into two matrices

corresponding to the two virophage size fractions. Then, network inference was conducted

using the FlashWeave default settings after Q1 filtration. In the MCP phylogenetic tree, three

virophage clades contributed most of the NCLDV connections. Thus, an NCLDV enrichment

analysis for the three clades was carried out using Fisher's exact test, and the *p*-value was

adjusted by the Benjamin–Hochberg method. This approach was the same as TIM, but we did

not use the TIM software because the current version of TIM requires inputs of eukaryotic

nodes with NCBI taxonomy annotations.

We used another approach, HGT, to predict the virophage-NCLDV interactions. First,

we generated an NCLDV genome database, which includes 56 reference NCLDV genomes

corresponding to our *polB* dataset and 2,074 metagenome-assembled genomes from a

previous study (24). A total of 827,548 coding sequences were included in this database.

Then, 195 virophage MCPs from the metagenomic data were BLASTp searched against this

database using an E-value cut-off of 1E−10 (with a minimum query coverage of 50% and a

minimum sequence identity of 50%). If a virophage MCP obtained a hit in the NCLDV

genome database with a lower E-value compared with hits in the MCP database (the hit to

itself was removed), the hit in the NCLDV genome database was considered an HGT

candidate.

Acknowledgments

This work was supported by JSPS/KAKENHI (Nos. 18H02279 and 19H05667 to H.O.

and Nos. 19K15895 and 19H04263 to H.E.), Kyoto University Research Coordination Alliance

(funding to H.E.), and the Collaborative Research Program of the Institute for Chemical

Research, Kyoto University (Nos. 2019-30 and 2020-27). Computational time was provided

by the SuperComputer System, Institute for Chemical Research, Kyoto University. We further

thank the *Tara* Oceans consortium, and the people and sponsors who supported *Tara* Oceans.

Tara Oceans (including both the *Tara* Oceans and *Tara* Oceans Polar Circle expeditions) would

not exist without the leadership of the *Tara* Expeditions Foundation and the continuous support

of 23 institutes (https://oceans.taraexpeditions.org). This article is contribution number XXX

of *Tara* Oceans. We thank Mallory Eckstut, PhD, from Edanz Group (https://en-author-

services.edanzgroup.com/ac) for editing a draft of this manuscript.

Figure 1. Overall workflow for inferring co-occurrence networks and quantitative assessment. This figure shows how the input data (*Tara* Oceans metagenomics and metabarcoding data) were used in this study. The definition of the confusion matrix for quantitative assessment is shown in the table. The LR+ and FDR equations are given at the lower right corner of the plot.

616
617
Figure 2. *polB*–V9 co-occurrence network. We performed co-occurrence analysis at the
OTU level and constructed the network with pooled *polB* –V9 associations from five size
fraction networks (A). To better display co-occurrence patterns, PolB OTUs were grouped at
the family or family-like level, and V9 OTUs were grouped using annotation at high
taxonomic ranks. The size of each node indicates the number of OTUs that belong to the
group, and the width of each edge indicates the number of associations between two
connected groups. Associations with positive weight are shown in red and negative
associations are shown in blue. (B) Number of associations connected to NCLDVs for each
major eukaryotic lineage in five size fractions. The top 10 lineages were retained, and other
lineages were omitted and shown as "others." Size fractions are presented in μm.

Figure 3. Density plots for the degree of NCLDV nodes in co-occurrence networks. The degree of an NCLDV node is given by the associations between this node and eukaryotes in the networks. The amount of NCLDV nodes are given on the top of the density values. (A) Positive degree (number of positive associations per node) for NCLDV nodes in five size fraction networks. (B) Negative degree (number of negative associations per node) for NCLDV nodes in five size fraction networks. Size fractions are presented in μm. NCLDV nodes with degree = 1 and degree > 1 are separated using a vertical line, and the number of nodes is given.

654
655
656
Figure 4. Positive likelihood ratios (LR+) in the NCLDV virus–host validation. (A)
General performance of co-occurrence networks is shown with the LR+ calculated with
associations pooled from five size fractions networks. To show the relationship between LR+
and FlashWeave association weight, the LR+ values are plotted along with the association
weight. (B) Performance of each size fraction network is shown with the violin plot by
ggplot2 with a bandwidth of 2. Size fractions are presented in μm.

678
679
680
Figure 5. Prediction of NCLDV virus–host relationships with TIM. (A) Undirected
network that shows the relationships between NCLDVs and eukaryotes after TIM filtration.
The size of each node indicates the number of predicted interactions of this group. The
weight of network edges as defined by the number of tree nodes enriched in each viral family
subtree to specific eukaryotic major lineages in the TIM analysis. Known virus–host
relationships are highlighted in red, and the pairs found to have horizontal gene transfer are
highlighted in yellow (1). (B) Performance of networks on NCLDV host prediction for
original FlashWeave results without a weight cut-off, weight cut-off > 0.4, and TIM
filtration, plotted by ggplot2 with a bandwidth of 2. (C) FDR of networks for NCLDV host
prediction with the original FlashWeave results without a weight cut-off, weight cut-off >
0.4, and TIM filtration.
692
693

Figure 6. Associations between virophages and NCLDVs. (A) Number of associations with virophages is shown for seven NCLDV families and two unclassified groups, "Medusaviruses" and "Pithoviruses." Associations in the femto-size fraction network are shown in yellow, and in the pico-size fraction network are shown in green. The number of positive associations is above the zero axis, and the number of negative associations is below the zero axis. (B) Phylogenetic tree was constructed from 195 environmental virophages and 47 reference MCP sequences. The outside layer indicates three major virophage clades. The inner two layers indicate that the virophage OTUs have at least one association with NCLDVs in femto- or pico-size fractions networks.

References

1. Colson P, De Lamballerie X, Yutin N, Asgari S, Bigot Y, Bideshi DK, Cheng XW, Federici BA, Van Etten JL, Koonin E V., La Scola B, Raoult D. 2013. "Megavirales", a proposed new order for eukaryotic nucleocytoplasmic large DNA viruses. Arch Virol 158:2517–2521.

2. Koonin E V., Dolja V V., Krupovic M, Varsani A, Wolf YI, Yutin N, Zerbini FM, Kuhn JH. 2020. Global Organization and Proposed Megataxonomy of the Virus World. Microbiol Mol Biol Rev 84:e00061-19.

3.   Koonin E V., Yutin N. 2019. Evolution of the Large Nucleocytoplasmic DNA Viruses
of Eukaryotes and Convergent Origins of Viral Gigantism. Advances in Virus
Research 103:167-202.

4.   Legendre M, Alempic JM, Philippe N, Lartigue A, Jeudy S, Poirot O, Ta NT, Nin S,
Couté Y, Abergel C, Claverie JM. 2019. Pandoravirus celtis illustrates the
microevolution processes at work in the giant Pandoraviridae genomes. Front
Microbiol 10:1–11.

5.   Boratto PVM, Oliveira GP, Machado TB, Andrade ACSP, Baudoin J-P, Klose T,
Schulz F, Azza S, Decloquement P, Chabrière E, Colson P, Levasseur A, La Scola B,
Abrahão JS. 2020. Yaravirus: A novel 80-nm virus infecting Acanthamoeba
castellanii. Proc Natl Acad Sci 117:16579–16586.

6.   Yoshikawa G, Blanc-Mathieu R, Song C, Kayama Y, Mochizuki T, Murata K, Ogata
H, Takemura M. 2019. Medusavirus, a Novel Large DNA Virus Discovered from Hot
Spring Water. J Virol 93:1–3.

7.   Forterre P. 2011. Manipulation of cellular syntheses and the nature of viruses: The
virocell concept. Comptes Rendus Chim 14:392–399.

8.   Mougari S, Sahmi-Bounsiar D, Levasseur A, Colson P, Scola B La. 2019. Virophages
of giant viruses: An update at eleven. Viruses 11:1–28.

9.   Koonin E V., Yutin N. 2010. Origin and evolution of eukaryotic large nucleo-
cytoplasmic DNA viruses. Intervirology 53:284–292.

10.   La Scola B, Audic S, Robert C, Jungang L, De Lamballerie X, Drancourt M, Birtles R,
Claverie JM, Raoult D. 2003. A giant virus in amoebae. Science 299:2033.

11.   Nagasaki K, Yamaguchi M. 1997. Isolation of a virus infectious to the harmful bloom
causing microalga Heterosigma akashiwo (Raphidophyceae). Aquat Microb Ecol
13:135–140.

12.   Moniruzzaman M, LeCleir GR, Brown CM, Gobler CJ, Bidle KD, Wilson WH,
Wilhelm SW. 2014. Genome of brown tide virus (AaV), the little giant of the
Megaviridae, elucidates NCLDV genome expansion and host-virus coevolution.
Virology 466–467:60–70.

13.   Wilson WH, Schroeder DC, Allen MJ, Holden MTG, Parkhill J, Barrell BG, Churcher
C, Hamlin N, Mungall K, Norbertczak H, Quail MA, Price C, Rabbinowitsch E,
Walker D, Craigon M, Roy D, Ghazal P. 2005. Complete genome sequence and lytic
phase transcription profile of a Coccolithovirus. Science 309:1090–1092.

14.   Colson P, Gimenez G, Boyer M, Fournous G, Raoult D. 2011. The giant Cafeteria

750   roenbergensis virus that infects a widespread marine phagocytic protist is a new

751   member of the fourth domain of life. PLoS One 6:13–17.

752   15.   Needham DM, Yoshizawa S, Hosaka T, Poirier C, Choi CJ, Hehenberger E, Irwin

753   NAT, Wilken S, Yung CM, Bachy C, Kurihara R, Nakajima Y, Kojima K, Kimura-

754   Someya T, Leonard G, Malmstrom RR, Mende DR, Olson DK, Sudo Y, Sudek S,

755   Richards TA, DeLong EF, Keeling PJ, Santoro AE, Shirouzu M, Iwasaki W, Worden

756   AZ. 2019. A distinct lineage of giant viruses brings a rhodopsin photosystem to

757   unicellular marine predators. Proc Natl Acad Sci U S A 116:20574–20583.

758   16.   Williams T. 2008. Natural Invertebrate Hosts of Iridoviruses (Iridoviridae). Neotrop

759   Entomol 37:615–632.

760   17.   Mihara T, Nishimura Y, Shimizu Y, Nishiyama H, Yoshikawa G, Uehara H, Hingamp

761   P, Goto S, Ogata H. 2016. Linking virus genomes with host taxonomy. Viruses 8:10–

762   15.

763   18.   Hingamp P, Grimsley N, Acinas SG, Clerissi C, Subirana L, Poulain J, Ferrera I,

764   Sarmento H, Villar E, Lima-Mendez G, Faust K, Sunagawa S, Claverie JM, Moreau

765   H, Desdevises Y, Bork P, Raes J, De Vargas C, Karsenti E, Kandels-Lewis S, Jaillon

766   O, Not F, Pesant S, Wincker P, Ogata H. 2013. Exploring nucleo-cytoplasmic large

767   DNA viruses in Tara Oceans microbial metagenomes. ISME J 7:1678–1695.

768   19.   Monier A, Claverie JM, Ogata H. 2008. Taxonomic distribution of large DNA viruses

769   in the sea. Genome Biol 9:R106.

770   20.   Li Y, Hingamp P, Watai H, Endo H, Yoshida T, Ogata H. 2018. Degenerate PCR

771   primers to reveal the diversity of giant viruses in coastal waters. Viruses 10:496.

772   21.   Mihara T, Koyano H, Hingamp P, Grimsley N, Goto S, Ogata H. 2018. Taxon richness

773   of "Megaviridae" exceeds those of bacteria and archaea in the ocean. Microbes

774   Environ 33:162–171.

775   22.   Endo H, Blanc-Mathieu R, Li Y, Salazar G, Henry N, Labadie K, de Vargas C,

776   Sullivan MB, Bowler C, Wincker P, Karp-Boss L, Sunagawa S, Ogata H. 2020.

777   Biogeography of marine giant viruses reveals their interplay with eukaryotes and

778   ecological functions. Nat Ecol Evol https://doi.org/10.1038/s41559-020-01288-w.

779   23.   Carradec Q, Pelletier E, Da Silva C, Alberti A, Seeleuthner Y, Blanc-Mathieu R,

780   Lima-Mendez G, Rocha F, Tirichine L, Labadie K, Kirilovsky A, Bertrand A, Engelen

781   S, Madoui MA, Méheust R, Poulain J, Romac S, Richter DJ, Yoshikawa G, Dimier C,

782   Kandels-Lewis S, Picheral M, Searson S, Acinas SG, Boss E, Follows M, Gorsky G,

783   Grimsley N, Karp-Boss L, Krzic U, Pesant S, Reynaud EG, Sardet C, Sieracki M,

Speich S, Stemmann L, Velayoudon D, Weissenbach J, Jaillon O, Aury JM, Karsenti

E, Sullivan MB, Sunagawa S, Bork P, Not F, Hingamp P, Raes J, Guidi L, Ogata H,

De Vargas C, Iudicone D, Bowler C, Wincker P. 2018. A global ocean atlas of

eukaryotic genes. Nat Commun 9:373.

24. Schulz F, Roux S, Paez-Espino D, Jungbluth S, Walsh DA, Denef VJ, McMahon KD,

Konstantinidis KT, Eloe-Fadrosh EA, Kyrpides NC, Woyke T. 2020. Giant virus

diversity and host interactions through global metagenomics. Nature 578:432–436.

25. Gallot-Lavallée L, Blanc G. 2017. A glimpse of nucleo-cytoplasmic large DNA virus

biodiversity through the eukaryotic genomicswindow. Viruses 9:17.

26. Guglielmini J, Woo AC, Krupovic M, Forterre P, Gaia M. 2019. Diversification of

giant and large eukaryotic dsDNA viruses predated the origin of modern eukaryotes.

Proc Natl Acad Sci U S A 116:19585–19592.

27. Cunha V Da, Gaïa M, Ogata H, Jaillon O, Delmont TO, Forterre P. 2020. Giant

viruses encode novel types of actins possibly related to the origin of eukaryotic actin:

the viractins. bioRxiv doi.org/10.1101/2020.06.16.150565.

28. Moniruzzaman M, Wurch LL, Alexander H, Dyhrman ST, Gobler CJ, Wilhelm SW.

2017. Virus-host relationships of marine single-celled eukaryotes resolved from

metatranscriptomics. Nat Commun 8:16054.

29. Roux S, Chan LK, Egan R, Malmstrom RR, McMahon KD, Sullivan MB. 2017.

Ecogenomics of virophages and their giant virus hosts assessed through time series

metagenomics. Nat Commun 8:858.

30. Tomaru Y, Katanozaka N, Nishida K, Shirai Y, Tarutani K, Yamaguchi M, Nagasaki

K. 2004. Isolation and characterization of two distinct types of HcRNAV, a single-

stranded RNA virus infecting the bivalve-killing microalga Heterocapsa

circularisquama. Aquat Microb Ecol 34:207–218.

31. Martínez Martínez J, Schroeder DC, Larsen A, Bratbak G, Wilson WH. 2007.

Molecular dynamics of Emiliania huxleyi and cooccurring viruses during two separate

mesocosm studies. Appl Environ Microbiol 73:554–562.

32. Sunagawa S, Acinas SG, Bork P, Bowler C, Eveillard D, Gorsky G, Guidi L, Iudicone

D, Karsenti E, Lombard F, Ogata H, Pesant S, Sullivan MB, Wincker P, de Vargas C.

2020. Tara Oceans: towards global ocean ecosystems biology. Nat Rev Microbiol

18:428–445.

33. Kaneko H, Blanc-Mathieu R, Endo H, Chaffron S, Delmont TO, Gaia M, Henry N,

Hernández-Velázquez R, Nguyen CH, Mamitsuka H, Forterre P, Jaillon O, De Vargas

C, Sullivan MB, Suttle CA, Guidi L, Ogata H. 2019. Eukaryotic virus composition can

predict the efficiency of carbon export in the global ocean. bioRxiv

https://doi.org/doi.org/10.1101/710228.

34. Fischer MG. 2016. Giant viruses come of age. Curr Opin Microbiol 31:50–57.

35. Delaroque N, Wolf S, Muller DG, Knippers R. 2000. The brown algal virus EsV-1

particle contains a putative hybrid histidine kinase. Virology 273:383–390.

36. Fitzgerald LA, Graves M V., Li X, Hartigan J, Pfitzner AJP, Hoffart E, Van Etten JL.

2007. Sequence and annotation of the 288-kb ATCV-1 virus that infects an

endosymbiotic chlorella strain of the heliozoon Acanthocystis turfacea. Virology

362:350–361.

37. Clerissi C, Desdevises Y, Grimsley N. 2012. Prasinoviruses of the Marine Green Alga

Ostreococcus tauri Are Mainly Species Specific. J Virol 86:4611–4619.

38. Bjorbækmo MFM, Evenstad A, Røsæg LL, Krabberød AK, Logares R. 2020. The

planktonic protist interactome: where do we stand after a century of research? ISME J

14:544–559.

39. Edwards RA, McNair K, Faust K, Raes J, Dutilh BE. 2016. Computational approaches

to predict bacteriophage-host relationships. FEMS Microbiol Rev 40:258–272.

40. Salkić NN. 2008. Objective assessment of diagnostic tests validity : a short review for

clinicians and other mortals . Part II. Acta Med Acad 39–42.

41. Coutinho FH, Silveira CB, Gregoracci GB, Thompson CC, Edwards RA, Brussaard

CPD, Dutilh BE, Thompson FL. 2017. Marine viruses discovered via metagenomics

shed light on viral strategies throughout the oceans. Nat Commun 8:1–12.

42. Martin-Platero AM, Cleary B, Kauffman K, Preheim SP, McGillicuddy DJ, Alm EJ,

Polz MF. 2018. High resolution time series reveals cohesive but short-lived

communities in coastal plankton. Nat Commun 9:266.

43. Huete-Ortega M, Cermeño P, Calvo-Díaz A, Marañón E. 2012. Isometric size-scaling

of metabolic rate and the size abundance distribution of phytoplankton. Proc R Soc B

Biol Sci 279:1815–1823.

44. Berry D, Widder S. 2014. Deciphering microbial interactions and detecting keystone

species with co-occurrence networks. Front Microbiol 5:1–14.

45. Kurtz ZD, Müller CL, Miraldi ER, Littman DR, Blaser MJ, Bonneau RA. 2015. Sparse

and Compositionally Robust Inference of Microbial Ecological Networks. PLoS

Comput Biol 11:1–25.

46. Xue Y, Chen H, Yang JR, Liu M, Huang B, Yang J. 2018. Distinct patterns and

processes of abundant and rare eukaryotic plankton communities following a reservoir
cyanobacterial bloom. ISME J 12:2263–2277.

47. Tackmann J, Matias Rodrigues JF, von Mering C. 2019. Rapid Inference of Direct
Interactions in Large-Scale Ecological Networks from Heterogeneous Microbial
Sequencing Data. Cell Syst 9:286-296.e8.

48. Suttle CA. 2007. Marine viruses - Major players in the global ecosystem. Nat Rev
Microbiol 5:801–812.

49. Blanc-Mathieu R, Dahle H, Hofgaard A, David B, Kalinowski J, Ogata H, Sandaa R-
A. 2020. The genome of a persistent giant algal virus encodes an unprecedented
number of genes involved in energy metabolism. bioRxiv
doi.org/10.1101/2020.07.30.228163.

50. Yau S, Krasovec M, Felipe Benites L, Rombauts S, Groussin M, Vancaester E, Aury
JM, Derelle E, Desdevises Y, Escande ML, Grimsley N, Guy J, Moreau H, Sanchez-
Brosseau S, van de Peer Y, Vandepoele K, Gourbiere S, Piganeau G. 2020. Virus-host
coexistence in phytoplankton through the genomic lens. Sci Adv 6:eaay2587.

51. Claverie JM, Grzela R, Lartigue A, Bernadac A, Nitsche S, Vacelet J, Ogata H,
Abergel C. 2009. Mimivirus and Mimiviridae: Giant viruses with an increasing
number of potential hosts, including corals and sponges. J Invertebr Pathol 101:172–
180.

52. Clouthier SC, Vanwalleghem E, Copeland S, Klassen C, Hobbs G, Nielsen O,
Anderson ED. 2013. A new species of nucleo-cytoplasmic large DNA virus (NCLDV)
associated with mortalities in Manitoba lake sturgeon Acipenser fulvescens. Dis Aquat
Organ 102:195–209.

53. Gastrich MD, Leigh-Bell JA, Gobler CJ, Anderson OR, Wilhelm SW, Bryan M. 2004.
Viruses as potential regulators of regional brown tide blooms caused by the alga,
Aureococcus anophagefferens. Estuaries 27:112–119.

54. Duponchel S, Fischer MG. 2019. Viva lavidaviruses! five features of virophages that
parasitize giant DNA viruses. PLoS Pathog 15:1–7.

55. Xu S, Zhou L, Liang X, Zhou Y, Chen H, Yan S, Wang Y. 2020. Novel Cell-Virus-
Virophage Tripartite Infection Systems Discovered in the Freshwater Lake Dishui
Lake in Shanghai, China. J Virol 94:e00149-20.

56. Sun S, La Scola B, Bowman VD, Ryan CM, Whitelegge JP, Raoult D, Rossmann MG.
2010. Structural Studies of the Sputnik Virophage. J Virol 84:894–897.

57. Gregory AC, Zayed AA, Conceição-Neto N, Temperton B, Bolduc B, Alberti A,

Ardyna M, Arkhipova K, Carmichael M, Cruaud C, Dimier C, Domínguez-Huerta G,

Ferland J, Kandels S, Liu Y, Marec C, Pesant S, Picheral M, Pisarev S, Poulain J,

Tremblay JÉ, Vik D, Acinas SG, Babin M, Bork P, Boss E, Bowler C, Cochrane G, de

Vargas C, Follows M, Gorsky G, Grimsley N, Guidi L, Hingamp P, Iudicone D,

Jaillon O, Kandels-Lewis S, Karp-Boss L, Karsenti E, Not F, Ogata H, Poulton N,

Raes J, Sardet C, Speich S, Stemmann L, Sullivan MB, Sunagawa S, Wincker P,

Culley AI, Dutilh BE, Roux S. 2019. Marine DNA Viral Macro- and Microdiversity

from Pole to Pole. Cell 177:1109-1123.e14.

58.   Pesant S, Not F, Picheral M, Kandels-Lewis S, Le Bescot N, Gorsky G, Iudicone D,

Karsenti E, Speich S, Trouble R, Dimier C, Searson S. 2015. Open science resources

for the discovery and analysis of Tara Oceans data. Sci Data 2:150023.

59.   de Vargas C, Audic S, Henry N, Decelle J, Mahé F, Logares R, Lara E, Berney C, Le

Bescot N, Probert I, Carmichael M, Poulain J, Romac S, Colin S, Aury J-M, Bittner L,

Chaffron S, Dunthorn M, Engelen S, Flegontova O, Guidi L, Horák A, Jaillon O,

Lima-Mendez G, Lukeš J, Malviya S, Morard R, Mulot M, Scalco E, Siano R, Vincent

F, Zingone A, Dimier C, Picheral M, Searson S, Kandels-Lewis S, Acinas SG, Bork P,

Bowler C, Gorsky G, Grimsley N, Hingamp P, Iudicone D, Not F, Ogata H, Pesant S,

Raes J, Sieracki ME, Speich S, Stemmann L, Sunagawa S, Weissenbach J, Wincker P,

Karsenti E. 2015. Ocean plankton. Eukaryotic plankton diversity in the sunlit ocean.

Science 348:1261605.

60.   Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G, Djahanschiri

B, Zeller G, Mende DR, Alberti A, Cornejo-Castillo FM, Costea PI, Cruaud C,

D'Ovidio F, Engelen S, Ferrera I, Gasol JM, Guidi L, Hildebrand F, Kokoszka F,

Lepoivre C, Lima-Mendez G, Poulain J, Poulos BT, Royo-Llonch M, Sarmento H,

Vieira-Silva S, Dimier C, Picheral M, Searson S, Kandels-Lewis S, Boss E, Follows

M, Karp-Boss L, Krzic U, Reynaud EG, Sardet C, Sieracki M, Velayoudon D, Bowler

C, De Vargas C, Gorsky G, Grimsley N, Hingamp P, Iudicone D, Jaillon O, Not F,

Ogata H, Pesant S, Speich S, Stemmann L, Sullivan MB, Weissenbach J, Wincker P,

Karsenti E, Raes J, Acinas SG, Bork P. 2015. Ocean plankton. Structure and function

of the global ocean microbiome. Science 348:1261359.

61.   Matsen FA, Kodner RB, Armbrust EV. 2010. pplacer: linear time maximum-

likelihood and Bayesian phylogenetic placement of sequences onto a fixed reference

tree. BMC Bioinformatics 11:538.

62.   De Vargas C, Engelen S, Hingamp P, Sieracki M, Vargas C, Audic S, Henry N,

920   Decelle J, Mahé F, Logares R, Lara E, Berney C, Bescot N, Probert I, Carmichael M,

921   Poulain J, Romac S. 2015. Eukaryotic plankton diversity in the sunlit ocean. Science

922   103:167–202.

923 63. Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, Boutte C, Burgaud G, De

924   Vargas C, Decelle J, Del Campo J, Dolan JR, Dunthorn M, Edvardsen B, Holzmann

925   M, Kooistra WHCF, Lara E, Le Bescot N, Logares R, Mahé F, Massana R, Montresor

926   M, Morard R, Not F, Pawlowski J, Probert I, Sauvadet AL, Siano R, Stoeck T, Vaulot

927   D, Zimmermann P, Christen R. 2013. The Protist Ribosomal Reference database

928   (PR2): A catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with

929   curated taxonomy. Nucleic Acids Res 41:597–604.

930 64. Oksanen J, Kindt R, Legendre P, O'Hara B, Simpson GL, Solymos PM, Stevens

931   MHH, & Wagner H. 2008. The vegan package. Community Ecol Packag 190.

932 65. Watts SC, Ritchie SC, Inouye M, Holt KE. 2019. FastSpar: Rapid and scalable

933   correlation estimation for compositional data. Bioinformatics 35:1064–1066.

934 66. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D,

935   Burovski E, Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M, Wilson J,

936   Millman KJ, Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E, Carey CJ, Polat İ,

937   Feng Y, Moore EW, VanderPlas J, Laxalde D, Perktold J, Cimrman R, Henriksen I,

938   Quintero EA, Harris CR, Archibald AM, Ribeiro AH, Pedregosa F, van Mulbregt P,

939   Vijaykumar A, Bardelli A Pietro, Rothberg A, Hilboll A, Kloeckner A, Scopatz A, Lee

940   A, Rokem A, Woods CN, Fulton C, Masson C, Häggström C, Fitzgerald C, Nicholson

941   DA, Hagen DR, Pasechnik D V., Olivetti E, Martin E, Wieser E, Silva F, Lenders F,

942   Wilhelm F, Young G, Price GA, Ingold GL, Allen GE, Lee GR, Audren H, Probst I,

943   Dietrich JP, Silterra J, Webber JT, Slavič J, Nothman J, Buchner J, Kulick J,

944   Schönberger JL, de Miranda Cardoso JV, Reimer J, Harrington J, Rodríguez JLC,

945   Nunez-Iglesias J, Kuczynski J, Tritz K, Thoma M, Newville M, Kümmerer M,

946   Bolingbroke M, Tartre M, Pak M, Smith NJ, Nowaczyk N, Shebanov N, Pavlyk O,

947   Brodtkorb PA, Lee P, McGibbon RT, Feldbauer R, Lewis S, Tygier S, Sievert S,

948   Vigna S, Peterson S, More S, Pudlik T, Oshima T, Pingel TJ, Robitaille TP, Spura T,

949   Jones TR, Cera T, Leslie T, Zito T, Krauss T, Upadhyay U, Halchenko YO, Vázquez-

950   Baeza Y. 2020. SciPy 1.0: fundamental algorithms for scientific computing in Python.

951   Nat Methods 17:261–272.

952 67. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.

953   2009. BLAST+: Architecture and applications. BMC Bioinformatics 10:1–9.

68. Yang WT, Parikh JR, Thomas Stavros A, Otto P, Maislin G. 2018. Exploring the
negative likelihood ratio and how it can be used to minimize false-positives in breast
imaging. Am J Roentgenol 210:301–306.

69. Faust K, Lima-Mendez G, Lerat JS, Sathirapongsasuti JF, Knight R, Huttenhower C,
Lenaerts T, Raes J. 2015. Cross-biome comparison of microbial association networks.
Front Microbiol 6:1–13.

70. Roux S, Adriaenssens EM, Dutilh BE, Koonin E V., Kropinski AM, Krupovic M,
Kuhn JH, Lavigne R, Brister JR, Varsani A, Amid C, Aziz RK, Bordenstein SR, Bork
P, Breitbart M, Cochrane GR, Daly RA, Desnues C, Duhaime MB, Emerson JB,
Enault F, Fuhrman JA, Hingamp P, Hugenholtz P, Hurwitz BL, Ivanova NN, Labonté
JM, Lee KB, Malmstrom RR, Martinez-Garcia M, Mizrachi IK, Ogata H, Páez-Espino
D, Petit MA, Putonti C, Rattei T, Reyes A, Rodriguez-Valera F, Rosario K, Schriml L,
Schulz F, Steward GF, Sullivan MB, Sunagawa S, Suttle CA, Temperton B, Tringe
SG, Thurber RV, Webster NS, Whiteson KL, Wilhelm SW, Wommack KE, Woyke T,
Wrighton KC, Yilmaz P, Yoshida T, Young MJ, Yutin N, Allen LZ, Kyrpides NC,
Eloe-Fadrosh EA. 2019. Minimum information about an uncultivated virus genome
(MIUVIG). Nat Biotechnol 37:29–37.

71. Katoh K, Misawa K, Kuma KI, Miyata T. 2002. MAFFT: A novel method for rapid
multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res
30:3059–3066.

72. Price MN, Dehal PS, Arkin AP. 2009. Fasttree: Computing large minimum evolution
trees with profiles instead of a distance matrix. Mol Biol Evol 26:1641–1650.

73. Letunic I, Bork P. 2019. Interactive Tree of Life (iTOL) v4: Recent updates and new
developments. Nucleic Acids Res 47:256–259.

74. Paul Shannon, Andrew Markiel, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,
Schwikowski B, Ideker T. 2003. Cytoscape: A Software Environment for Integrated
Models. Genome Res 13:426.

75. Eddy SR. 1998. Profile hidden Markov models. Bioinformatics 14:755–763.

Author contributions

984 LM and HO designed the study. LM performed most of the bioinformatics analysis.
985 HE generated the primary NCLDV data. RBM, SC, RHV, and HK contributed to the
986 bioinformatics analysis. All authors contributed to the writing of the manuscript.
987

Materials & Correspondence

Correspondence and material requests should be addressed to HO (email: ogata@kuicr.kyoto-u.ac.jp).

Competing financial interests

The authors declare no competing financial interests.