Beyond broadband: towards a spectral decomposition of EEG microstates

Victor Férat [1], Martin Seeber [1], Christoph M. Michel [1,2], Tomas Ros [1,2]

1 Functional Brain Mapping Laboratory, Department of Basic Neurosciences, Campus Biotech, University of Geneva, Geneva, Switzerland

2 Centre for Biomedical Imaging (CIBM) Lausanne-Geneva, Geneva, Switzerland

Credit authorship contribution statement

Victor Férat: Conceptualization, Formal analysis, Methodology, Visualization, Writing - original draft, Writing - review & editing.

Martin Seeber: Writing - review & editing

Christoph Michel: Writing - review & editing

Tomas Ros: Writing - Conceptualization, Formal analysis, Methodology review & editing.

Data availability statement

The data and code supporting our results are openly available in the OSF repository and can be found at: https://osf.io/bfsge/

Other 3rd party software and datasets used in this study can be found by following citations in the paper.

Acknowledgments

The study was supported by the Swiss National Science Foundation (NCCR Synapsy grant No. 51NF40 – 185897 and grant No. 320030_184677) to CMM.

We also would like to thank the Mind-Body-Emotion group at the Max Planck Institute for Human Cognitive and Brain Sciences for all the work they have done to make their dataset public.

Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Abstract: Originally applied to alpha oscillations in the 1970s, MS analysis has since been used to decompose mainly broadband EEG signals (e.g. 1-40 Hz). We hypothesized that MS decomposition within separate, narrow frequency bands could provide more fine-grained information for capturing the spatio-temporal complexity of multichannel EEG. In this study using a large open-access dataset (n=203), we decomposed EEG recordings into 4 classical frequency bands (delta, theta, alpha, beta) in order to compare their individual MS segmentations using mutual information as well as traditional MS measures (e.g. mean duration, time coverage). Firstly, we confirmed that MS topographies were spatially equivalent across all frequencies, matching the canonical broadband maps (A, B, C, D). Interestingly however, we observed strong informational independence of MS temporal sequences between spectral bands, together with significant divergence in traditional MS measures. For example, relative to broadband, alpha/beta band dynamics displayed greater time coverage of maps A & B, while map D was more prevalent in delta/theta bands. Moreover, by using a frequency-specificMS taxonomy (e.g. θA, αC), we were able to predict the eyes-open vs closed- behavioural state significantly better using alpha-band MS features compared with broadband ones (80% vs 73% accuracy). Overall, our findings demonstrate the value and validity of spectrally-specific MS analyses, which may prove useful for identifying new neural mechanisms in fundamental research and/or for biomarker discovery in clinical populations.

Keywords: EEG ; microstates; topography; Resting state ; eyes closed ; eyes open ; alpha; mutual information

1   Introduction

Multi-channel Electroencephalography (EEG) is a long-established tool for exploring the human brain's spatio-temporal activities. Microstate (MS) analysis (Michel and Koenig 2018) , first introduced by Lehmann (Lehmann 1971) in 1971, takes advantage of EEG's high temporal resolution to segment EEG signals into short successive periods of time characterized by metastable scalp topographies. Initially applied to narrow-band alpha oscillations (8 -12 Hz)(Lehmann 1971), microstate analysis is nowadays usually performed on broadband EEG signals (1 – 40Hz) (Michel and Koenig 2018; Zanesco et al. 2020). Historically, only a limited number of studies (Javed et al. 2019; Merrin et al. 1990; Musaeus et al. 2020) have focused on applying MS analysis to the traditional frequencies associated with cortical oscillations (e.g. delta, theta, alpha, beta etc.). For example, in the 1990's, Merrin et al (Merrin et al. 1990) were the first to report on a significant difference in MS segments between schizophrenic patients and controls specifically in the theta EEG band. On the other hand, more recent work in healthy subjects found that MS dynamics were independent of EEG power fluctuations across the frequency spectrum (Britz, Van De Ville, and Michel 2010), which technically supported the rationale for performing broadband MS analysis. Neuroimaging studies have nevertheless emerged showing that anatomically-distinct cortical regions display different dominant EEG frequencies, with occipito-parietal regions more active in the alpha band, and prefrontal regions being biased more toward delta or theta power (Groppe et al. 2013; Keitel and Gross 2016; Mellem et al. 2017). Moreover, ongoing cortical dynamics have been reported to fluctuate from a local resting/idling alpha oscillatory state to task-specific active mode(s)

dominated by other rhythms (e.g. theta (Ribary, Doesburg, and Ward

2017), gamma (Hipp, Engel, and Siegel 2011)). As a consequence, cortical

regions could combine different frequencies for integrating/segregating

information across large-scale networks, a phenomenon termed

"oscillatory multiplexing" (Akam and Kullmann 2014). Finally, of more

clinical significance, a growing body of work has indicated abnormal EEG

spectral power in distinct frequencies across cortical regions in a variety

of brain disorders (Ros et al. 2014; Schulman et al. 2011). Therefore, given

that different spatial topographies uncovered by MS analysis imply

anatomically-distinct cortical generators (according to the forward-model

of EEG generation (Michel and Koenig 2018)), it is reasonable to

hypothesize that distinct MS topographies may display different spatial

and/or temporal profiles across the frequency spectrum.

To investigate this question as well as gain a deeper understanding of

frequency-specific MS signature(s), we sought to explicitly decompose MS

spatio-temporal dynamics within *discrete, narrow-band frequency bands*

(i.e. delta, theta, alpha, and beta), with the aim of comparing them to the

classical analysis of the broadband signal.

Here, we employed a validated, open-source dataset (Babayan et al.

2019) of resting-state EEG recordings from 203 healthy subjects during

both eyes opened and eyes closed conditions. These were then filtered in

the classical EEG bands (delta: 0-4 Hz, theta: 4-8 Hz, alpha: 8-12 Hz, beta:

15-30 Hz) to obtain band specific signals. These narrow-band signals, in

addition to the broadband (1-30 Hz) signal, were then independently

subjected to standard microstate analysis (Pascual-Marqui, Michel, and

Lehmann 1995). Map topography, mean duration, occurrence, time

coverage, and global explained variance (GEV) were used as quantitative
measures of spatiotemporal microstate dynamics. In summary, and using
spatial correlation analysis, we firstly demonstrate remarkably similar
microstate topographies across frequencies, closely matching the classical
broadband maps. Interestingly, however, we observed strong
informational independence of microstate sequences between
frequencies, in addition to significant differences in established measures
of temporal dynamics (mean duration, occurrence, and time coverage).

In conclusion, our results support a more diverse, frequency-specific
application of microstate analysis compatible with the narrow-band MS
analyses of early pioneers (Lehmann 1971; Merrin et al. 1990). We
anticipate this approach to provide a more fine-grained spectral
information not visible to the standard broadband analysis, for example in
the identification of biomarkers in clinical populations or for
understanding the mechanisms underlying EEG microstates.

2 Methods

2.1 Dataset

EEG recordings were obtained from 203 anonymized participants
enrolled in the Mind-Brain-Body study (Babayan et al. 2019). Detailed
protocol and inclusion criteria are reported in (Babayan et al. 2019). The
overall sample consisted of 227 participants divided into 2 groups: the
younger adults group with participant age ranging between 20 and 35
years (N = 153, 45 females, mean age = 25.1 years, SD = 3.1) and an older
adults group with age ranging between 59 and 77 years (N = 74, 37

females, mean age = 67.6 years, SD = 4.7). Medical and psychological
screening was conducted on all participants at the Day Clinic for Cognitive
Neurology of the University Clinic Leipzig and the Max Planck Institute for
Human and Cognitive and Brain Sciences in order to include only healthy
patients. The study protocol was approved by the ethics committee of the
University of Leipzig (reference 154/13-ff). Data were obtained in
accordance with the Declaration of Helsinki.

145　2.2　Recordings

Resting state EEGs were recorded using 61 scalp electrodes
(ActiCAP, Brain Products GmbH, Gilching, Germany), and one additional
VEOG electrode for recording right eye activity. All electrodes were placed
according to the international standard 10–20 extended localization
system with FCz reference, digitized with a sampling frequency of fs=2500
Hz,an amplitude resolution of 0.1 microV ,and bandpass filtered between
0.015Hz and 1 kHz. The ground was located at the sternum and scalp
electrode impedance was kept below 5 KΩ. Recordings took place in an
electrically shielded and sound-attenuated EEG booth. Here, 60s blocks
alternated between eyes open (EO) and eyes closed (EC) conditions for a
total recording of 16 min (8 blocks EC, 8 blocks EO, starting with EC).
During the EO condition, participants were asked to stay awake while
fixating their eyes on a black cross presented on a white background.

159　2.3　Prepocessing

The prepossessing steps are extensively described in (Babayan et al.
2019), which we summarize below. All EEG recordings were down-

sampled from 2500 to 250 Hz and filtered between 1 and 45Hz (8th order,
Butterworth filter). Blocks sharing the same condition were concatenated
leading to the creation of 2 datasets per subject. After visual inspection,
outlying channels were rejected and EEG segments presenting noise
and/or artefacts were removed (except eye movements and eye blinks
that were kept for further prepossessing). PCA was used to reduce data
dimensionality, by keeping PCs (N≥30) that explain 95% of the total data
variance. Then, independent component analysis (ICA) was performed
using the Infomax (runica) algorithm. Components reflecting eye
movement, eye blink or heartbeat related artefacts were removed.

Before performing microstate analysis, the following additional
prepossessing steps were conducted using MNE-python (Gramfort et al.
2013): missing/bad channels were interpolated using spherical spline
interpolation, the reference was re-projected to average and recordings
were down-sampled to 100Hz. Finally, each recording was filtered into
broadband plus the 5 traditional EEG frequency bands: broadband (1-30
Hz), delta (1-4 Hz), theta (4-8 Hz), alpha (8-12 Hz), beta (15-30 Hz). Filter
design consisted of a two-pass forward and reverse, zero-phase, non-
causal band-pass FIR filter with the following parameters.

• **Broadband (1-30 Hz)**: - Lower passband edge: 1.00 - Lower
transition bandwidth: 1.00 Hz (-12 dB cutoff frequency: 0.50
Hz) - Upper passband edge: 30.00 Hz- Upper transition
bandwidth: 7.50 Hz (-12 dB cutoff frequency: 33.75 Hz) Filter
length: 331 samples (3.310 sec)
• **Delta (1-4 Hz)**: - Lower passband edge: 1.00 - Lower transition
bandwidth: 1.00 Hz (-12 dB cutoff frequency: 0.50 Hz) - Upper

passband edge: 4.00 Hz - Upper transition bandwidth: 2.00 Hz

(-12 dB cutoff frequency: 5.00 Hz) - Filter length: 331 samples

(3.310 sec)

• **Theta (4-8 Hz)**: - Lower passband edge: 4.00 - Lower

transition bandwidth: 2.00 Hz (-12 dB cutoff frequency: 3.00

Hz) - Upper passband edge: 8.00 Hz - Upper transition

bandwidth: 2.00 Hz (-12 dB cutoff frequency: 9.00 Hz) - Filter

length: 165 samples (1.650 sec)

• **Alpha (8-12 Hz)**: - Lower passband edge: 8.00 - Lower

transition bandwidth: 2.00 Hz (-12 dB cutoff frequency: 7.00

Hz) - Upper passband edge: 12.00 Hz - Upper transition

bandwidth: 3.00 Hz (-12 dB cutoff frequency: 13.50 Hz) - Filter

length: 165 samples (1.650 sec)

• **Beta (15-30 Hz)**: - Lower passband edge: 15.00 - Lower

transition bandwidth: 3.75 Hz (-12 dB cutoff frequency: 13.12

Hz) - Upper passband edge: 30.00 Hz - Upper transition

bandwidth: 7.50 Hz (-12 dB cutoff frequency: 33.75 Hz) - Filter

length: 89 samples (0.890 sec)

For all filters, a Hamming window with 0.0194 passband ripple and 53

dB stopband attenuation was used to reduce border effects.

2.4   MS segmentation

2.4.1   Segmentation

Microstate segmentation was applied to each combination of

frequency band (broadband, delta, theta, alpha, beta) x behavioural

condition (EO, EC) leading to the computation of 10 optimal clusters using

the methodology described below. First, local maxima of the Global Field

Power (GFP) known to represent portions of EEG data with highest signal

to noise ratio (Koenig and Brandeis 2016), were extracted from each

individual recording. Then, 20 epochs of 500 time points randomly drawn

from the previous extraction were submitted to a modified k-means

cluster analysis using the free academic software Cartool (Brunet, Murray,

and Michel 2011). For each number of cluster centers K ranging from 1 to

12, 50 k-means initialisations were applied to each epoch. The

initialisation with highest global explained variance (GEV) was selected

and kept for further processing. A meta-criterion (Bréchet et al. 2019) was

used to choose the optimal number of cluster centers k for each epoch.

Individual optimal clusters were then merged within conditions and

within frequencies to form 10 groups of 4060 clusters. Each group was

then randomly re-sampled into 100 epochs of 5000 time points, and

submitted to the same clustering algorithm (50 initialisations, with meta

criterion selection), leading to the extraction of 100 optimal clusters per

group. Finally, these 100 clusters were submitted to the modified K means

clustering algorithm to extract, for each number of cluster centroids k, a

set of maps which best represent the spatiotemporal variance of

frequency specific EEG data within each condition.

2.4.2   Selection of "common" MS maps

Given that we found high spatial correlations between MS maps across all

frequencies and EO/EC conditions, we fitted the broadband maps directly

to all the frequency bands in order to have a common reference. This may

be considered a heuristic approach for the sake of simplicity. An

alternative approach we explored was to perform subject-level (i.e. 1[st]

level) clustering on all data concatenated *within-subject* (across

frequencies, and/or conditions), followed by group-level (i.e. 2nd-level)
clustering. We found this to once again produce identical maps to the
broadband decomposition. This method could theoretically be used to find
the most "common" clusters across different datasets, in case of variable
k-means outputs (e.g. visually similar MS maps at different k-values). Since
it is beyond the scope of this paper, we leave it to future studies to validate
this method more rigorously.

2.4.3   Back-fitting of MS Maps

The common topographic maps selected above were then assigned
to every time point from all individual recordings using the traditional MS
back-fitting method (Van De Ville, Britz, and Michel 2010). First, the
spatial correlation was computed between every timepoint and map.
Using the so called 'winner takes all' algorithm, each timepoint was
labelled according to the map with which it shared the highest absolute
spatial correlation. Timepoints were labelled as "non-assigned" when the
absolute spatial correlation was below $r < 0.5$ threshold. To ensure
temporal continuity of MS segmentation, a smoothing step (Brunet et al.
2011; Pascual-Marqui et al. 1995) was applied. Finally, segments with
duration shorter than 3 samples (30ms) were assigned to neighbouring
segments using the following rule: the segment was split into two parts,
where each part was assigned to the neighbouring segment with the
higher spatial correlation. With backfitting completed, we extracted 4
spatiotemporal parameters for each microstate map, namely:

• **Global explained variance** (Gev) described as the sum of
264   variances of the original recording explained by the considered

microstate map weighted by the Global Field Power at each

moment in time. Units are percentages (%) between 0 and 1.

• **Mean duration** (MeanDurs), defined as the mean temporal

duration of segments assigned to each MS map. Units are in

seconds (s).

• **Time coverage** (TimeCov) is the ratio of time frames assigned

to each MS map relative to the total number of time frames from

the recording. Results are Units are percentages (%) between

0 and 1.

274   **2.5   Adjusted Mutual information score**

Scikit-learn (Pedregosa et al. 2011) implementation of the adjusted

mutual information score (AMI) (Vinh, Epps, and Bailey 2010) was used

to quantify the mutual information (MI) shared between different MS

temporal segmentations, whilst simultaneously accounting for random

overlap due to chance. This metric, bounded between 0 and 1, is used to

evaluate the statistical (in)dependence of two variables. In our case, AMI

is estimated between the symbolic sequences of two different microstate

segmentations (e.g. ABDCADB vs ABDBDAC). A high score (approaching

1) indicates that the two segmentations agree on the temporal order of all

labels while a low score (approaching 0) indicates that the segmentations'

labels are not temporally aligned. We selected the corrected version of

this metric in order to control for the impact of differences in label

distribution due to chance (for example differences in overall time

coverage between labels).

2.6 Statistics

Statistical analyses were performed on the 4 main spatiotemporal parameters (Global explained variance, Mean spatial correlation, Mean duration, Time coverage). Tests were conducted using a two sided permutation test for equality of means on paired samples (same subject, either between condition, either between frequencies) under the H0 hypothesis that both frequency (i.e. condition) share the same mean against the alternative H1 that the distributions come from two different populations. P-values were estimated by simulated random sampling with 10000 replications. As a large number of statistical tests were carried out without specific pre-planned hypotheses (Armstrong 2014), P values were corrected for multiple comparisons using the Bonferroni method. Corrected P-values are reported in the Results section, as well as the observed means (m) of both samples together with observed standard deviations. Effect sizes are reported as the standardised difference of means using Cohen's d (d).

2.7 Neurobehavioral prediction models

2.7.1 Model

Linear Support Vector Classification (SVC) with 'l2' norm penalization and squared hinge loss function were used to discriminate EO vs EC states using MS parameters pertaining to broad- as well as narrow- band EEG activity. Models were trained using 15 frequency specific features correspond to the 3 spatiotemporal MS metrics (Gev, MeanDurs, TimeCov) of each maps (A,B,C,D,C') of a given frequency. Band

specific prediction models were fitted with standardized features (after
removing their respective mean and scaling them to unit variance).

2.7.2   Evaluation

As suggested by Bouckaer et al. (Bouckaert n.d.),   10 times
repeated 10 fold cross validation tests were used to assess classification
results. For each of the 100 runs, 9 folds of 20 subject's features each were
used to train the model while 1 fold of 20 subject's features was used to
evaluate the 3 diagnostic metrics:

• **Accuracy**: defined as number of the number of correctly predicted
samples out of all the testing set. **The operating characteristic**
(ROC) which the plot of true positive rate as a function of the false
positive rate. It is used to illustrate the classification trade-off for
different discrimination threshold.
• **Area under the curve (AUC)** defined as the area under the ROC
curve: it is an aggregate measure of performance for all possible
classification thresholds. AUC values are in the range of 0 to 1. A
model with 100% error in its predictions has an AUC of 0.0. If all
its predictions are correct, its AUC is 1.0.

95% confidence intervals were evaluated on each of the 3 metric
distributions with 10 degrees of freedom.

Statistical comparison between models were conducted on both
accuracy and AUC using a two- tailed permutation test for equality of
means on paired samples under the H0 hypothesis that the alpha-band (8-

12 Hz) had a greater mean of the classification metric than the broadband

(1-30 Hz). P-values were estimated by simulated random sampling with

10000 replications.

3 Results

3.1 Spatial Similarity of Microstate Maps

Figure 1: *Spatial correlation between MS topographies across behavioural conditions.*

Global cluster centroids of each frequency band within eyes-open (EO) or eyes-closed (EC) condition. Note that map polarity inversion is ignored in the classical analysis of spontaneous EEG.

Figure 1 illustrates the topographic results of MS segmentations in the different conditions and frequency bands. After visual inspection of optimal clusters at different cluster numbers (k), we identified that a value of k=5 revealed five MS topographies that were similar across all EEG bands and behavioural conditions, consistent with recent findings from our laboratory (Bréchet et al. 2019, 2020; D'Croz-Baron et al. 2019). MS maps were designated in line with the canonical prototypes from the literature and their respective symbols, featuring a left-right orientation (A), a right-left orientation (B), an anterior-posterior orientation (C), fronto-central maximum (D) and occipito-central (C')maximum.

Given the additional frequency dimension, we labelled the MS maps
firstly according to the Greek letters traditionally used for narrow-band
EEG (i.e. δ, θ, α, β) and then the Latin alphabet for the canonical map
symbols (i.e. A, B, C, D) . For example, αA denoted the left-right diagonal
map from the alpha band (α) segmentation, and δC the anterior-posterior
map from the delta band (δ) segmentation. The broadband segmentation
was designated with the prefix 'bb'.

Figure 2: *Spatial correlation between MS topographies across*
frequencies.
Spatial correlation of cluster centers of each sub-frequency bands compared
to broadband for eyes opened (EO) and eyes closed (EC) condition.

As shown in **Figure 2**, when comparing topographies between
broadband and each narrow-band (i.e. the diagonal entries in the
correlation matrix), all spatial correlations were $r > 0.98$. Consequently,

we fitted the broadband maps directly to all the frequency bands in order

to have a common reference.

We similarly observed common MS maps when comparing broadband

topographies between EO and EC conditions (**Suppl Figure 1**), with all

intraclass spatial correlations exceeding $r > 0.98$, thus providing

justification for comparing microstate parameters between behavioural

conditions while fitting condition specific broadband maps.

3.2   Mutual Information of Microstate Sequences

Figure 3: Adjusted mutual information of MS symbolic sequences
a) Mean adjusted mutual information (AMI) is depicted, for each behavioural condition (EC: eyes-closed, EO: eyes-open) across n = 203 subjects.
b) Mean adjusted mutual information of MS symbolic sequences between all

broadband and narrow-band combinations. Mean (n = 203 subjects)

adjusted mutual information (AMI) for all frequency pairs.

Briefly, Adjusted Mutual Information (AMI, bounded between 0 and 1)

is an index of how similar two separate MS segmentations are, by

estimating the degree of shared information (i.e. the number of time

points assigned with the same MS) between their symbolic sequences (e.g.

ABCD vs ABDA). The 'adjusted' aspect ensures the measure is unbiased for

symbolic overlap(s) due to chance (Vinh et al. 2010). Higher AMI

(approaching 1) indicates nearly identical MS temporal sequences, while

lower AMI (approaching 0) indicates temporally independent sequences.

(I.e. low overlap)

As shown in **Figures 3**, the AMI between broadband and narrow-band

segmentations in the EO condition showed a value of s = 0.06 for delta, s =

0.03 for theta, s = 0.05 for alpha, and s = 0.01 for beta. These values are

surprisingly low and we can conclude that the broadband segmentation is

comparatively independent of the narrow-band one. Similar conclusions

of temporal independence can be made by examining the AMI between the

narrow-bands *themselves*, with a maximum AMI value between theta and

alpha bands (EO: s = 0.006, EC: s = 0.009), and a minimum AMI value of s

= 0.001 for non-adjacent EEG bands (delta-alpha, delta-beta, theta-beta)

As a sanity check, when inspecting the EO vs EC transition, the shared

information with broadband decreased for the delta band (s = 0.03) but

increased for the alpha band (s = 0.12). The latter is in line with

expectations, as alpha oscillations are known to increase considerably

during eye closure, which would amplify their contribution to the

broadband signal and consequently their shared dynamics.

3.3 Between-frequency comparison of classical MS measures

Here we tested for significant differences between broadband and narrow-band filtered EEG in the classical MS measures: Global explained variance (Gev), Time Coverage (TimeCov), and Mean Duration (MeanDurs). This was done by conducting paired t-tests between broadband (bb) and respective narrow-band (delta to beta) MS measures across all n=203 subjects. For each MS measure and MS map, results were visualised using heat-plots as the narrow-band *absolute difference* from the broadband mean value. Here, a red/blue background indicated significant positive/negative differences at $p < 0.05$, while a white colour indicated non-significant differences at $p > 0.05$. Exact p-values and effect sizes are reported in Table 2 of the Supplementary Results.

Global Explained Variance (Gev)

Figure 4: *Microstate map differences in Global Explained Variance*
(Gev) between broadband and narrow-band filtered EEG, for eyes open (EO)
and eyes closed (EC) conditions. The first row (in grey) represents the mean
Gev of the broadband segmentation. Coloured rows represent the mean Gev
difference between the narrowband (δ, θ, α, β) and broadband
segmentations, for each microstate (A – C'). Significant differences have
red/blue backgrounds, while non-significant ones have a white/grey
background.

Figure 4 illustrates the percent differences in global explained
variance (Gev) between each narrow-band vs broadband, demonstrating
the presence of specific "fingerprints" between frequencies (rows) or MS

maps (columns). , MS segmentation of **delta** band activity showed

significantly higher GEV across most MS maps in both EO (+16 %) and EC

(+ 16 %) conditions, compared to the broadband segmentation. A similar

but less pronounced effect was found for the theta segmentation,

suggesting that low-frequency EEG fluctuations in the 1-8 Hz (delta-theta)

range may be accounted for more parsimoniously using the 5 canonical

MS maps than 1-30 Hz (broadband) activity.

The profile of the **alpha** band segmentation depended on

behavioural condition. During EC, microstate C had significantly increased

Gev (+ 12%) relative to broadband, indicating that distinct MS

topographies have stronger behavioural specificity at narrow-band

frequencies (see section *Application to Behavioural Classification*).

Conversely, during EO, microstates A (+ 2%) and B (+ 4%), expressed

significantly more Gev compared to broadband.

Conversely**, or the beta** band, Gev was significantly greater compared to

broadband for maps A (EO: + 2%, EC+ 3%) and B (EO: + 2%, EC: + 2%)

while it was significantly reduced for maps D (EO: - 2%, EC: - 1%) and C'

(EO: - 3%, EC: - 3%) . Microstate C explained less variance compared to

broadband but the comparison was significant only for EC (- 2%).

3.3.1   Time Coverage (TimeCov)

Figure 5: Microstate map differences in Time Coverage (TimeCov)
between broadband and narrow-band filtered EEG, for eyes open (EO) and
eyes closed (EC) conditions. The first row (in grey) represents the mean
TimeCov of the broadband segmentation. Coloured rows represent the mean
TimeCov difference between the narrowband (δ, θ, α, β) and broadband
segmentations, for each microstate (A – C'). Significant differences have
red/blue backgrounds, while non-significant ones have a white/grey
background.

During MS segmentation, each time-point is assigned to only *one*

of the five canonical MS topographies through a winner-takes-all process

(i.e. the map with the highest spatial correlation with that time-point

wins). Time Coverage (TimeCov) refers to the *average* prevalence of each

MS map over the whole EEG recording **(Figure 5)**, expressed as a

percentage (i.e. number of time-points assigned to a particular MS map

divided by the total number of time-points).

Comparing the **delta** band segmentation to broadband, TimeCov was

reduced for maps A (EO: - 3%, EC -0%), B (EO: - 2%, EC ns) , and C (EO:

5% , EC: - 6 %)   , while it was increased for maps D (EO: + 5%, EC: + 4%)

and C '(EO: +4, EC: +2%).

Small differences were generally observed between **theta** and broadband

MS distributions, with the notable exception of a significant increase in

map D (EO: + 5%, EC: + 7%) and decrease in map C (EO: - 2%, EC: - 4%),

consistently found across behavioural states. For the **alpha** band,

TimeCov was particularly greater compared to broadband for maps A (EO:

+ 2%, EC ns) and B (EO: + 4%, EC: + 1%)

Finally, comparing the **beta** band to broadband, TimeCov was

increased for maps A (EO: +4%, EC: +5% , $p < 0.05$) , B (EO: +4% , EC:

+4%, $p < 0.05$) ,while it was reduced for maps D (EO: -4%, EC: -2%, $p <$

0.05) and C '(EO: -6%, EC: -6%, $p < 0.05$)

3.3.2 Mean Duration (Mean Durs)

Figure 6: Microstate map differences in Mean Duration (MeanDurs) between broadband and narrow-band filtered EEG, for eyes open (EO) and eyes closed (EC) conditions. The first row (in grey) represents the mean MeanDurs of the broadband segmentation. Coloured rows represent the mean MeanDurs difference between the narrowband (δ, θ, α, β) and broadband segmentations, for each microstate (A – C'). Significant differences have red/blue backgrounds, while non-significant ones have a white/grey background.

While analysing MS mean duration (MeanDurs) we first observed that MS

were significantly shorter in **delta** compared to broadband across all

maps: map A (EO: -14ms, EC: -13ms), map B (EO: -14ms, EC: -14ms) , map

C(EO:-15ms, EC: _-17ms, map D (EO: -7ms, EC: -8ms) and map C' (EO: -

8ms, EC: -11ms).On the contrary **theta** segments were significantly longer

than broadband ones for map A (EO: +6ms, EC: +11), map B (EO: +6ms,

EC: +11ms) and map D(EO: -7ms, EC: -8ms). Similar but non-significant

increases were also observed for maps C and C'.

Notably, all MS maps had significantly increased duration in **alpha** band

compared to broadband: map A (EO: +57ms, EC: +70ms), map A (EO:

+62ms, EC: +70ms) map C (EO: +71ms, EC: +150ms) map D (EO: +42ms,

EC: +72ms) and map C'(EO: +57ms, EC: +63ms).

Less pronounced and less consistent differences were found for the **beta**

band, which revealed a single significant increase of map A duration in the

EC condition (EO: ns, EC: +9ms).

3.4 Within-frequency comparison of classical MS measures during eyes-open (EO) versus eyes-closed (EC)

*Figure 7: EO versus EC comparison using frequency specific microstate parameters. a) Mean global explained variance (Gev (%)),and b) mean time coverage (time coverage (%)) for each microstate (A – C') within each frequency band (broadband, delta, theta, alpha, beta) for both eyes closed condition (EC, blue) and eyes opened condition (EO, red). Significance values are indicated from paired permutation test on mean between conditions: no star: 0.05 > p, ∗: p < 0.05, * color indicates the condition with highest value.*

520   Here, we directly compared EO vs EC conditions within each frequency

521   band, and only relevant cases where narrow-band measures were salient

522   compared to the broadband analysis are reported. Full results are

523   documented in Table 3 and Suppl Figure 2 of the Supplementary Results.

524   3.4.1   Global Explained Variance (Gev)

Across frequencies, map C' explained more variance during EC than

during EO for all bands (bb: d = 0.59, p<.05 | δ: d = 0.43, p<.05 | θ: d =

0.35, p<.05 | α: d = 0.35, p<.05 | β d = 0.67, ns) while map D expressed

less variance(bb: d = -0.14, ns | δ: d = -0.07, ns | θ: d -0.29, p<.05 | α: d = -

0.42, p<.05 | β d = -0.73, p<.05). In contrast, the other MS maps had band-

specific modulations: during EC map C explained relatively less variance

in alpha (d = -1.0, p<.05) and broadband (d = -0.7, p<0.5),but expressed

more variance in delta (d = 0.20, p<.05) and beta (d = 0.21, p<.05). In

addition, statistically significant effects were detected in the narrow-

bands which were not evident in the broadband. For example, : the Gev

of map B decreased from EC to EO (d = -0.31, p < 0.05) in in the beta band

while no significant effect was found in broadband for this map.

537   3.4.2   Time coverage (TimeCov),

Map C was relatively more prevalent in EC vs EO in alpha (d = -0.71, *p <*

.05) and broadband (d = -0.48, *p* < .05), but the effect size was significantly

stronger in the alpha band. In contrast, the opposite effect was observed

in delta (d = 0.22, *p* < .05) and beta (d = 0.42, *p* < .05) bands, which showed

increased coverage of map C during EO compared to EC. Diverging effects

of time coverage in narrow-bands compared to broadband appeared

across a number of MS maps and frequency bands, during the EO vs EC

transition: βA –where TimeCov decreased prevalence (d = -0.20, $p < .05$),
bbA and βC –TimeCov increased, respectively (d = 0.23, $p < .05$) and (d =
0.21, $p <0 .05$). , while bbC TimeCov decreased (d = -0.71, $p < .05$). Where
non-significant differences between EO and EC conditions were found for
bbB coverage, *βB (d =-0.17, p<.05) and* ΘB (d = -0.35, p<.05) were
significantly increased in the EC condition while αB (d = 0.57, p < .05) was
decreased.

All in all, the EO vs EC- comparison revealed significant differences
between the broadband and narrow-band segmentations, which were
frequently map- and/or band- specific.

3.5 Statistical classification of EO vs EC behavioural states

Figure 8**: **Classifying EO vs EC states using classical microstate
measures. *Binary classification performance, computed as the area under*
the curve (AUC) of the receiver-operating characteristic (ROC), for 2
support-vector machine models. Input features of the first model (red) were
broadband MS parameters (see text) while the second model (blue) was
evaluated on alpha-band MS parameters. Shaded areas represent 95% CI.
Mean AUC ± standard deviation is reported in the legend.

As a proof-of-concept, we tested the applicability of band-specific
microstates in the context of behavioural prediction, i.e. the binary
discrimination of EO versus EC states using machine learning. Leveraging

the well-known effect of alpha-band modulation during eye
opening/closure (Barry et al. 2007) we hypothesised that alpha-band
specific MS parameters would be stronger and more specific predictors of
the EO/EC transition than those derived from the broadband
segmentation. As we did not have any MS map specific hypotheses, we
selected as features the basic set of classical MS measures from our study:
5 maps (A, B, C, D, C') x 3 variables (Gev, TimeCov, MeanDurs). Then, using
these 15 features, we applied a linear support vector machine (SVM) to
classify EO vs EC recordings across all 203 subjects, separately for alpha-
band and broadband models.

Following 10 fold cross-validation, we observed higher accuracy (i.e.
sensitivity) for the SVM model tested with alpha-band MS parameters (80
± 5%) than the one tested with broadband MS parameters (73 ± 6%). The
superiority of the alpha-band model was reinforced by a separate analysis
of the area under the curve (AUC) of the receiver-operating characteristic
(ROC), which also incorporates the false-positive rate (i.e. specificity).
Here alpha-band (AUC 87±5%) outperformed broadband (77±7%) by a
full 10%. Finally, these differences were statistically significant for both
accuracy (Cohen's d=1.75, p<0 .001) and AUC (Cohen's d=1.28, p <0 .001)
demonstrating that a narrow-band MS segmentation may provide higher
sensitivity/specificity in differentiating behavioural state relative to a
broadband one.
588

4 Discussion

Historically, the first microstate (MS) analysis was applied by Lehmann and colleagues to narrowband (alpha) oscillations (Lehmann 1971), yet this "frequency-specific" approach appears to have been overlooked during the last decades of MS research in favour of decomposing broadband EEG signals (e.g. 2-40 Hz)(Michel and Koenig 2018; Pascual-Marqui et al. 1995). Hence, the present study specifically explored the MS characteristics of narrow-band EEG signals, their quantitative interrelationship, and whether they provide any novel information compared to course-grained broadband dynamics. This was done by first filtering the broadband EEG signal into several narrow-band frequencies (delta, theta, alpha, and beta), with the goal of comparing MS symbolic sequences and classical measures (explained variance, mean duration, time coverage) between them, as well as across different behavioural conditions (eyes open (EO) vs eyes closed (EC)).

4.1 The spatial dimension: microstate topographies

We first investigated whether analogous MS scalp topographies would be produced by segmenting broadband versus narrow-band EEG signals (including the alpha band (Milz et al. 2017)). Interestingly, we observed highly similar MS topographies (with minimum spatial correlations of $r > 0.98$) across all investigated broad- and narrow- band frequencies (broadband, delta to beta), as well as between EO/EC conditions. This is compatible with recent work by Brechet and colleagues (Bréchet et al. 2020), who observed that states of sleep and wake

exhibited significantly different spectral content (e.g. delta vs beta power)
but very similar MS maps. Moreover, these maps corresponded to the
canonical (broadband) topographies previously described in the
literature (Custo et al. 2017; Michel and Koenig 2018). It is therefore
tempting to assume that identical neuronal sources are involved in
generating the same topographies across frequencies. However, although
different maps imply different generators (forward problem), same
topographies do not necessarily imply identical generators (inverse
problem). Due to the ill-posed nature of EEG signals (constructive and
destructive electromagnetic fields), similar scalp potentials can still be
generated by different underlying brain mechanisms (Helmholtz 1853).
Hence, although we cannot unequivocally conclude that MS maps across
the EEG spectrum are generated by the same brain sources operate, this
would be the most probable and parsimonious interpretation. Moreover,
we must juxtapose our findings with work from other groups (Musaeus et
al. 2020) which applied a similar approach but didn't necessarily find the
same topographies across the EEG spectrum. From a methodological point
of view, it should be kept in mind that narrow-band MS analysis does not
per se require similar topographies between frequencies. In this case,
although cross frequency comparisons would not be possible due to
dissimilar maps, it would remain valid to study and quantify
spatiotemporal MS parameters within each frequency band separately, for
example, in the service of clinical biomarker discovery (Merrin et al.
1990). Reassuringly, the MS maps of our study replicate the ones derived
from independent work utilising the same EEG dataset (Zanesco et al.
2020), further supporting the reproducibility of MS analysis despite
methodological variations between studies (e.g. absence of resampling).

4.2 The temporal dimension: mutual information of microstate sequences

Milz and colleagues (Milz et al. 2017) recently proposed that alpha oscillations were the major component driving microstate dynamics. In general, adjusted mutual information (AMI) analyses reported in our work reveal low values (near or below 0.1) of information shared between the narrow-band segmentations, including alpha, and that of the broadband decomposition. However, consistent with the work of Milz and colleagues (Milz et al. 2017), the alpha-band during eyes-closed (EC) did indeed have the highest shared information with broadband (around 0.125). Importantly however, this relationship did not necessarily hold during eyes-open (delta being highest). This indicates specific narrow-band contribution(s) to broadband dynamics heavily depend on behavioural state. Moreover, if narrow-band(s) topographies were directly responsible for the origin of the spatial distribution of the broadband signal, one would expect much higher AMI values (at least 0.5) than those, we observed. In view of the results presented, it would be inaccurate to claim that alpha band, or any other narrow-band as the dominant source of broadband topographies.

In contrast, our results appear to support the ideas of Croce and colleagues (Croce et al. 2020), who suggested that broadband MS dynamics could not be extrapolated from one or a subset of EEG frequency bands. It remains unclear how the interaction of several narrow-band-components leads to a substantially different broadband MS decomposition. We speculate that this might stem from the fact that i) different narrow band signals could

cancel each other at specific time points and ii) microstate assignment is
non-linear given the winner-takes all approach.

Lastly and most intriguingly, no significant informational interrelations
were found between the narrow-band topographical dynamics
themselves (e.g. delta vs beta, theta vs alpha), indicating that each EEG
band appears to have has its own independent dynamics. This may not be
surprising, considering that spontaneous EEG oscillations have been
reported to dynamically switch from a resting signatures (e.g. alpha) to
task-specific active mode(s) dominated by theta (Ribary et al. 2017), beta
(Fernández et al. 1995) or gamma activities (Hipp et al. 2011)). In this
context, our observations of spatiotemporal independence between
narrow-band EEG components support the operation of "oscillatory
multiplexing" (Akam and Kullmann 2014) mechanisms in the cortex,
whereby brain regions could combine different frequencies for
integrating/segregating information across large-scale networks (Le Van
Quyen 2011)

4.3   Classical microstate measures: explained variance, time coverage, and mean duration

682
683

Global Explained Variance (Gev)

Overall, MS segmentations of low-frequency bands (delta, theta)
explained more of the global topographical variance compared to the
classical broadband segmentaion. Since Gev is normalised by and
therefore independent of global field power, this suggests that the
goodness-of-fit (i.e Gev) of the five-map MS model is clearly greater in the

delta/theta bands compared to broadband EEG. In higher frequencies,

effects are more map-specific, as for example we observed an increase in

Gev for the diagonal maps (A,B) and a decrease for midline (D,C') ones..

Time Coverage (TimeCov)

Different patterns of MS temporal coverage were observed according to

frequency. The delta and theta bands seem to exhibit greater prevalence

of map D and decreased incidence of maps A, B, and C. This is remarkable

insofar map D shares an intriguing overlap with the topography of the

well-known frontal-midline theta-rhythm (fm-theta) (Scheeringa et al.

2008; Töllner et al. 2017). Conversely, beta band dynamics seem to favour

the more frequent appearance of maps A and B in lieu of maps D and C'.

The appearance of alpha-band MS maps, on the other hand, appear to be

more state-dependent, as might be expected given the well-known

expression of the posterior alpha rhythm during eyes-closed. Accordingly,

there was a strong behavioural dissociation particularly for map C, which

displayed a relatively greater temporal prevalence during EC compared to

EO.

The presence of such "spectral fingerprints" suggests the existence of an

affinity between distinct MS topographies and EEG bands suggests that

different cortical generators (i.e. topographies) may be activated

preferentially in certain frequencies (Groppe et al. 2013; Keitel and Gross

2016; Mellem et al. 2017).,.

Mean Duration (MeanDurs)

Microstates are defined as short periods of time during which the scalp
electric field remains quasi-stable. Traditional microstate analysis does
not suggest specific frequency filtering, thus resulting in various
broadband filter settings across studies (Michel and Koenig 2018). Our
findings demonstrate that temporally stable states (around 80ms or
longer) are present within all classical EEG narrow-bands (i.e. delta to
beta). It is established that such spatiotemporal structures do not appear
for randomly shuffled EEG (Wackermann et al. 1993). For most EEG
narrow-bands, mean MS durations were usually in the same range as the
typically reported 70–120 ms, but often longer. For example, the average
MS duration in the eyes open state was around 150 ms for the alpha-band,
compared to 90 ms for broadband. It will be therefore interesting for
future studies to examine the mechanistic links between the aggregated
dynamics of the broadband microstates and those of frequency-specific
modes.

4.4   Discriminating between behavioural states:
eyes open versus eyes closed

Within each EEG narrow-band, between 8 (for theta) and 14 (for alpha)
of the classical MS measures were found to be statistically significant. In
comparison, classical MS measures derived from broadband EEG had a
higher effect size for only one parameter (broadband microstate B mean
duration). For all other 14 parameters, at least one narrow-band
component showed a relatively stronger effect size.

Hence, the addition of the frequency dimension has the primary benefit
of increasing both the number and the specificity of potential neural

markers that could aid clinical prognosis or provide insight into brain
mechanisms. We therefore conclude that the extra frequency dimension
in itself leads to a more fine-grained decomposition of multichannel and
multiplex EEG signals than the standard broadband analysis.

This is directly supported by our behavioural classification results,
where we utilised alpha-band vs broadband MS parameters to predict EO
vs EC states using the recordings of all 203 subjects. The significant boost
in overall accuracy (alpha-band: 80% vs broadband: 73%) and area under
the ROC curve nicely demonstrate that frequency-specific MS measures
may provide higher behavioural predictive power than those derived
from the broadband analysis. Although beyond the scope of this paper, we
expect that this approach to provide advantages in other contexts, such as
event-related potential (ERP) analyses or for discriminating between
different clinical populations.

Interestingly, we found that in a few cases the narrow-band effects
were opposite in directionality to the broadband results. Thus, limiting
the analysis to only the latter could lead to incomplete (or even incorrect)
interpretations of underlying brain dynamics.

5   Potential Limitations and Future Work

The current study may technically be considered exploratory, given the
large number of tests that were carried out and in the absence of well-
defined hypotheses, However, we carried out Bonferroni correction for all
tests which may be considered the most conservative method for
controlling for multiple comparisons. Several studies have thus far

proposed explanations for the origins of broadband MS topographies
(Britz et al. 2010). We feel it is still too early to make analogies or
speculations between these results and those of the narrow-band
dynamics. However, we believe that the application of the methodology
proposed here may lead to valuable insights in order to more fully
understand the underlying spectral tapestry of EEG microstates.

6   Conclusion

Ultimately, we report a number of important and novel findings
between the classical broadband MS analysis, generally performed in the
EEG field, and its application to more narrow frequency bands relevant to
cortical oscillatory activities. In a nutshell, it appears that each canonical
EEG frequency band possesses its own independent spatiotemporal
dynamics, whilst the relative prevalence of MS topographies themselves
differ across frequencies. Moreover, it appears that broadband dynamics
cannot be appropriately explained by the simple summation of individual
narrow-band frequency components.

Analysis of narrow-band MS parameters revealed spatial and temporal
characteristics that both converged and diverged from broadband MS
findings. At the very least, our results indicate that narrow-band analysis
is justified as complementary to the usual broadband MS analysis. A
narrow-band decomposition into frequencies more specific for cortical
oscillatory activity could not only advance and consolidate findings in
clinical disorders e.g. (Merrin et al. 1990) (Musaeus et al. 2020), but also
enable a better understanding of the organization and functioning of
large-scale brain systems..

Data and Code Availability Statement

The data and code supporting our results are openly available in the OSF repository and can be found at: https://osf.io/bfsge/ (DOI 10.17605/OSF.IO/BFSGE).

Other 3rd party software and datasets used in this study can be found by following citations in the paper.

References

Akam, Thomas, and Dimitri M. Kullmann. 2014. 'Oscillatory Multiplexing of Population Codes for Selective Communication in the Mammalian Brain'. *Nature Reviews Neuroscience* 15(2):111–22. doi: 10.1038/nrn3668.

Armstrong, Richard A. 2014. 'When to Use the B Onferroni Correction'. *Ophthalmic and Physiological Optics* 34(5):502–8.

Babayan, Anahit, Miray Erbey, Deniz Kumral, Janis D. Reinelt, Andrea MF Reiter, Josefin Röbbig, H. Lina Schaare, Marie Uhlig, Alfred Anwander, Pierre-Louis Bazin, and others. 2019. 'A Mind-Brain-Body Dataset of MRI, EEG, Cognition, Emotion, and Peripheral Physiology in Young and Old Adults'. *Scientific Data* 6:180308.

Barry, Robert J., Adam R. Clarke, Stuart J. Johnstone, Christopher A. Magee, and Jacqueline A. Rushby. 2007. 'EEG Differences between Eyes-Closed and Eyes-Open Resting Conditions'. *Clinical Neurophysiology* 118(12):2765–73. doi: 10.1016/j.clinph.2007.07.028.

Bouckaert, Remco R. n.d. 'Choosing between Two Learning Algorithms Based on Calibrated Tests'. 8.

Bréchet, Lucie, Denis Brunet, Gwénaël Birot, Rolf Gruetter, Christoph M. Michel, and João Jorge. 2019. 'Capturing the Spatiotemporal

Dynamics of Self-Generated, Task-Initiated Thoughts with EEG
and FMRI'. *Neuroimage* 194:82–92.

Bréchet, Lucie, Denis Brunet, Lampros Perogamvros, Giulio Tononi, and
Christoph M. Michel. 2020. 'EEG Microstates of Dreams'. *Scientific*
Reports 10(1):17069. doi: 10.1038/s41598-020-74075-z.

Britz, Juliane, Dimitri Van De Ville, and Christoph M. Michel. 2010. 'BOLD
Correlates of EEG Topography Reveal Rapid Resting-State
Network Dynamics'. *NeuroImage* 52(4):1162–70. doi:
10.1016/j.neuroimage.2010.02.052.

Brunet, Denis, Micah M. Murray, and Christoph M. Michel. 2011.
'Spatiotemporal Analysis of Multichannel EEG: CARTOOL'.
Computational Intelligence and Neuroscience 2011.

Croce, Pierpaolo, Angelica Quercia, Sergio Costa, and Filippo Zappasodi.
2020. 'EEG Microstates Associated with Intra-and Inter-Subject
Alpha Variability'. *Scientific Reports* 10(1):1–11.

Custo, Anna, Dimitri Van De Ville, William M. Wells, Miralena I. Tomescu,
Denis Brunet, and Christoph M. Michel. 2017.
'Electroencephalographic Resting-State Networks: Source
Localization of Microstates'. *Brain Connectivity* 7(10):671–82.

D'Croz-Baron, David F., Mary Baker, Christoph M. Michel, and Tanja Karp.
2019. 'EEG Microstates Analysis in Young Adults With Autism
Spectrum Disorder During Resting-State'. *Frontiers in Human*
Neuroscience 13:173. doi: 10.3389/fnhum.2019.00173.

Fernández, Thalía, Thalía Harmony, Mario Rodríguez, Jorge Bernal, Juan
Silva, Alfonso Reyes, and Erzsébet Marosi. 1995. 'EEG Activation
Patterns during the Performance of Tasks Involving Different
Components of Mental Calculation'. *Electroencephalography and*
Clinical Neurophysiology 94(3):175–82. doi: 10.1016/0013-
4694(94)00262-J.

Gramfort, Alexandre, Martin Luessi, Eric Larson, Denis A. Engemann,
Daniel Strohmeier, Christian Brodbeck, Roman Goj, Mainak Jas,

Teon Brooks, Lauri Parkkonen, and others. 2013. 'MEG and EEG
Data Analysis with MNE-Python'. *Frontiers in Neuroscience* 7:267.

Groppe, David M., Stephan Bickel, Corey J. Keller, Sanjay K. Jain, Sean T.
Hwang, Cynthia Harden, and Ashesh D. Mehta. 2013. 'Dominant
Frequencies of Resting Human Brain Activity as Measured by the
Electrocorticogram'.   *NeuroImage*   79:223–33.   doi:
10.1016/j.neuroimage.2013.04.044.

Helmholtz, H. von. 1853. 'Ueber Einige Gesetze Der Vertheilung
Elektrischer Strome in Korperlichen Leitern, Mit Anwendung Auf
Die Thierisch-Elektrischen Versuche (Schluss.)'. *Annalen Der*
Physik 165(7):353–77.

Hipp, Joerg F., Andreas K. Engel, and Markus Siegel. 2011. 'Oscillatory
Synchronization in Large-Scale Cortical Networks Predicts
Perception'.   *Neuron*   69(2):387–96.   doi:
10.1016/j.neuron.2010.12.027.

Javed, Ehtasham, Pierpaolo Croce, Filippo Zappasodi, and Cosimo Del
Gratta. 2019. 'Hilbert Spectral Analysis of EEG Data Reveals
Spectral Dynamics Associated with Microstates'. *Journal of*
Neuroscience Methods 325:108317.

Keitel, Anne, and Joachim Gross. 2016. 'Individual Human Brain Areas Can
Be Identified from Their Characteristic Spectral Activation
Fingerprints'. *PLoS Biology* 14(6):e1002498.

Koenig, Thomas, and Daniel Brandeis. 2016. 'Inappropriate Assumptions
about EEG State Changes and Their Impact on the Quantification
of EEG State Dynamics'. *Neuroimage* 125:1104–6.

Le Van Quyen, Michel. 2011. 'The Brainweb of Cross-Scale Interactions'.
New   Ideas   in   Psychology   29(2):57–63.   doi:
10.1016/j.newideapsych.2010.11.001.

Lehmann, D. 1971. 'Multichannel Topography of Human Alpha EEG Fields'.
Electroencephalography and Clinical Neurophysiology 31(5):439–
49.

Mellem, Monika S., Sophie Wohltjen, Stephen J. Gotts, Avniel Singh
Ghuman, and Alex Martin. 2017. 'Intrinsic Frequency Biases and
Profiles across Human Cortex'. *Journal of Neurophysiology*
118(5):2853–64.

Merrin, Edward L., Patricia Meek, Thomas C. Floyd, and Enoch Callaway
III. 1990. 'Topographic Segmentation of Waking EEG in
Medication-Free Schizophrenic Patients'. *International Journal of*
Psychophysiology 9(3):231–36.

Michel, Christoph M., and Thomas Koenig. 2018. 'EEG Microstates as a Tool
for Studying the Temporal Dynamics of Whole-Brain Neuronal
Networks: A Review'. *NeuroImage* 180:577–93. doi:
10.1016/j.neuroimage.2017.11.062.

Milz, Patricia, Roberto D. Pascual-Marqui, Peter Achermann, Kieko Kochi,
and Pascal L. Faber. 2017. 'The EEG Microstate Topography Is
Predominantly Determined by Intracortical Sources in the Alpha
Band'. *Neuroimage* 162:353–61.

Musaeus, Christian S., Knut Engedal, Peter Høgh, Vesna Jelic, Arjun R.
Khanna, Troels Wesenberg Kjær, Morten Mørup, Mala Naik, Anne-
Rita Oeksengaard, Emiliano Santarnecchi, and others. 2020.
'Changes in the Left Temporal Microstate Are a Sign of Cognitive
Decline in Patients with Alzheimer's Disease'. *Brain and Behavior*
e01630.

Pascual-Marqui, Roberto D., Christoph M. Michel, and Dietrich Lehmann.
1995. 'Segmentation of Brain Electrical Activity into Microstates:
Model Estimation and Validation'. *IEEE Transactions on*
Biomedical Engineering 42(7):658–65.

Pedregosa, F., G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M.
Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A.
Passos, D. Cournapeau, M. Brucher, M. Perrot, and E. Duchesnay.
2011. 'Scikit-Learn: Machine Learning in Python'. *Journal of*
Machine Learning Research 12:2825–30.

Ribary, Urs, S. M. Doesburg, and L. M. Ward. 2017. 'Unified Principles of
Thalamo-Cortical Processing: The Neural Switch'. *Biomedical*
Engineering Letters 7(3):229–35. doi: 10.1007/s13534-017-
0033-4.

Ros, Tomas, Bernard J. Baars, Ruth A. Lanius, and Patrik Vuilleumier. 2014.
'Tuning Pathological Brain Oscillations with Neurofeedback: A
Systems Neuroscience Framework'. *Frontiers in Human*
Neuroscience 8. doi: 10.3389/fnhum.2014.01008.

Scheeringa, René, Marcel C. M. Bastiaansen, Karl Magnus Petersson,
Robert Oostenveld, David G. Norris, and Peter Hagoort. 2008.
'Frontal Theta EEG Activity Correlates Negatively with the Default
Mode Network in Resting State'. *International Journal of*
Psychophysiology 67(3):242–51. doi:
10.1016/j.ijpsycho.2007.05.017.

Schulman, Joshua J., Robert Cancro, Sandlin Lowe, Feng Lu, Kerry D.
Walton, and Rodolfo R. Llinás. 2011. 'Imaging of Thalamocortical
Dysrhythmia in Neuropsychiatry'. *Frontiers in Human*
Neuroscience 5. doi: 10.3389/fnhum.2011.00069.

Töllner, Thomas, Yijun Wang, Scott Makeig, Hermann J. Müller, Tzyy-Ping
Jung, and Klaus Gramann. 2017. 'Two Independent Frontal Midline
Theta Oscillations during Conflict Detection and Adaptation in a
Simon-Type Manual Reaching Task'. *The Journal of Neuroscience*
37(9):2504–15. doi: 10.1523/JNEUROSCI.1752-16.2017.

Van De Ville, D., J. Britz, and C. M. Michel. 2010. 'EEG Microstate Sequences
in Healthy Humans at Rest Reveal Scale-Free Dynamics'.
Proceedings of the National Academy of Sciences 107(42):18179–
84. doi: 10.1073/pnas.1007841107.

Vinh, Nguyen Xuan, Julien Epps, and James Bailey. 2010. 'Information
Theoretic Measures for Clusterings Comparison: Variants,
Properties, Normalization and Correction for Chance'. *The Journal*
of Machine Learning Research 11:2837–54.

Wackermann, J., D. Lehmann, CM Michel, and WK Strik. 1993. 'Adaptive
Segmentation of Spontaneous EEG Map Series into Spatially
Defined Microstates'. *International Journal of Psychophysiology*
14(3):269–83.

Zanesco, Anthony P., Brandon G. King, Alea C. Skwara, and Clifford D.
Saron. 2020. 'Within and Between-Person Correlates of the
Temporal Dynamics of Resting EEG Microstates'. *NeuroImage*
211:116631. doi: 10.1016/j.neuroimage.2020.116631.

946