Non-phylogenetic identification of co-evolving genes for reconstructing the archaeal Tree of Life

L. Thibério Rangel[1], Shannon M. Soucy[2], João C. Setubal[3], Johann Peter Gogarten[4, 5], Gregory P. Fournier[1]

[1] Department of Earth, Atmospheric & Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA, USA

[2] Department of Biomedical Data Science, Geisel School of Medicine, Dartmouth College, Hanover, NH, USA

[3] Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, São Paulo, Brasil

[4] Department of Molecular and Cell Biology, University of Connecticut, Storrs, CT, USA

[5] Institute for Systems Genomics, University of Connecticut, Storrs, CT, USA

Corresponding author: L. Thibério Rangel, lthiberiol@gmail.com

Abstract

Assessing the phylogenetic compatibility between individual gene families is a crucial and often computationally demanding step in many phylogenomics analyses. Here we describe the Evolutionary Similarity Index (I_{ES}) to assess shared evolution between gene families using a weighted Orthogonal Distance Regression applied to sequence distances. This approach allows for straightforward pairing of paralogs between co-evolving gene families without resorting to multiple tests, or *a priori* assumptions of molecular interactions between protein products from assessed genes. The utilization of pairwise distance matrices, while less informative than phylogenetic trees, circumvents error-prone comparisons between trees whose topologies are inherently uncertain. Analyses of simulated gene family evolution datasets showed that I_{ES} was more accurate and less susceptible to noise than popular tree-based methods (Robinson-Foulds and geodesic distance) for assessing evolutionary signal compatibility, since it bypasses phylogenetic reconstruction and its inherent uncertainty. Applying I_{ES} to a real dataset of 1,322 genes from 42 archaeal genomes identified eight major clusters of gene families with compatible evolutionary trends. Four of these clusters included genes with a taxonomic distribution across all archaeal phyla, while other clusters included a subset of taxa that do not map to generally accepted archaeal clades, indicating possible shared horizontal transfers by clustered gene families. We identify one strongly connected set of 62 genes from the same cluster, occurring as both single-copy and multiple homologs per genome, with compatible phylogenetic reconstructions closely matching previously published species trees for Archaea. An I_{ES} implementation is available at https://github.com/lthiberiol/evolSimIndex.

Introduction

Phylogenies reconstructed from single genes are known to poorly reflect the underlying history of whole genomes, as the detectable phylogenetic signal from an isolated locus cannot be extrapolated to represent whole genomes (Dagan and Martin 2006; Bapteste et al. 2009; Koonin et al. 2009). To ameliorate this effect, it has become common practice to estimate species' evolutionary histories by concatenating multiple sequence alignments of genes from the core genome, i.e., single copy and widely conserved across sampled genomes, which greatly increases the number of sites available for phylogenetic inference. The preference towards using core genome sequences is due their expected resistance to horizontal gene transfer (HGT) (Thomas and Nielsen 2005; Sorek et al. 2007; Popa and Dagan 2011); however, despite the lower frequency of HGT among some gene families, it has been shown that horizontal exchange also affects genes from the core genome. In fact, the slow substitution rate and corresponding high sequence conservation of the core genome may even favor HGT, permitting

increases in neutral and nearly-neutral HGT at the genus and species levels (Papke and Gogarten 2012;

Shapiro et al. 2012).

Given this context, it is clear that more rigorous methods are needed to identify genes best

reflecting the underlying vertical evolutionary signal in a group of species; such methods should seek to

maximize the compatibility between evolutionary trends of the chosen gene families in order to provide a

more robust basis for phylogenomic reconstruction. Many strategies have been proposed to assess

similarities between the phylogenetic signals obtained by individual gene trees - e.g., Robinson-Foulds

bipartition compatibility (RF) (Robinson and Foulds 1981) and geodesic distance (D_{geo}) (Kimmel and

Sethian 1998; Kupczok et al. 2008; Owen and Provan 2011) - as well as other methods that assess

similarities between phylogenetic profiles (Pellegrini et al. 1999; Vert 2002; Barker and Pagel 2005; Liu

et al. 2018). The majority of tree-based methods are based on straightforward comparisons between tree

topologies (Kunin et al. 2005; Leigh et al. 2008; Puigbò et al. 2009; Mirarab et al. 2014; Gori et al. 2016).

However, while an intuitive solution, comparisons between tree topologies require phylogenetic trees of

all assessed gene families to be accurately reconstructed, adding a substantial computational cost to an

already computationally demanding task. Furthermore, the vastness of tree space, combined with the

inherent uncertainty of phylogenetic reconstruction, provides multiple sources of errors in tree-based

evolutionary similarity assessments. Another method to assess the evolutionary compatibility of genes is

based on similarities between patterns of presence and absence (phylogenetic profiles) of such genes

among genomes of interest . While recent implementations displayed substantial improvements (Liu et al.

2018) when compared to the early ones, reliance on an initial reference tree hamper general applicability.

Phylogenetic profile-based methods also do not assess divergencies between sequences of homologous

genes, which limits the resolution of their results.

Accounting for uncertainty-based variations in tree topology (i.e., bipartition support) further

increases the computational burden and decreases the resolution of the evaluated phylogenetic signal

(e.g., collapsing low support bipartitions or weighing them based on support). A proposed solution to

bypass the computational cost of tree similarity assessments is Pearson's correlation coefficient (*r*)

between evolutionary distance matrices (Goh et al. 2000; Pazos and Valencia 2001; Novichkov et al.

2004; Rangel et al. 2019). Unlike tree-based comparisons, methods based on Pearson's *r* enable simple

implementations to detect similar evolutionary signals between gene families with histories complicated

by multiple homologs within genomes by estimating correlation coefficients using all possible pairings of

paralogs between gene families (Gertz et al. 2003; Ramani and Marcotte 2003). Despite its application in

protein-protein interaction studies, the sensitivity of Pearson's *r* to noise in evolutionary distances and the

granularity of its estimates have yet to be compared to those of tree-based metrics. Direct Coupling

Analysis (DCA) has also been used to pair gene copies between possibly co-evolving gene families

(Gueudré et al. 2016), but despite positive results the assumption that protein products of co-evolving

genes must be directly interacting hampers its general applicability.

New approaches

Given the shortcomings pointed out above, we propose the Evolutionary Similarity Index (I_{ES}) as

a metric for similarities between evolutionary histories based on weighted Orthogonal Distance

Regression (wODR) between evolutionary pairwise distance matrices. We show that evolutionary

similarity estimates from wODR display a linear correlation with performed stepwise perturbations

simulated tree topologies. More common tree-based evolutionary similarity estimates, such as RF and

D_{geo}, tend to overestimate the impact of simulated topology changes, and consequently are significantly

more susceptible to errors in evolutionary history reconstruction. As a case study of this new method, we

assessed evolutionary similarities across 1,322 archaeal gene families and detected significant

evolutionary incompatibilities between conserved single-copy genes, as well as a clear central

evolutionary tendency involving 62 gene families that occur as both single and multiple-copies across

genomes.

Methodology

Orthogonal Distance Regression (ODR) is an errors-in-variables regression method that accounts

for measurement errors in both explanatory and response variables (Boggs et al. 1987), instead of

attributing all errors in the expected values exclusively to the response variable, as performed by Ordinary

Least Squares (OLS). While OLS regressions seek to minimize the sum of squared residuals of the

response variable, ODR minimizes the sum of squared residuals from each data point obtained by the

combination of explanatory and response variables. Novichkov et al. (Novichkov et al. 2004) assessed the

compatibility between the evolutionary history of genes with a reference genomic evolutionary history

using Pearson's *r* and estimates of an OLS regression's intercept. The latter extra step when compared to

other implementations using Pearson's *r* (Ramani and Marcotte 2003; Izarzugaza et al. 2008; Gueudré et

al. 2016) is required to infer that datapoints not fitting a regression line through zero are caused by HGT.

The approach proposed by Novichkov et al. requires two key assumptions that restrict the general

applicability of evolutionary assessments of empirical datasets: 1) there must exist a reference history to

which gene histories are compared; and 2) there are no errors in reference distances between genomes.

The approach described here is based on ODR. Its modelling of errors within both assessed

variables decreases the necessity of comparing gene family pairwise distances against a well-established

reference. Where *a priori* there is no clear separation between explanatory and response variables, errors-

in-variables approaches (e.g., ODR) are better suited to compare pairwise evolutionary distances between

two gene families. Independently weighing datapoints based on their residuals from an initial regression

line provides a framework less susceptible to underestimating overall evolutionary similarities due to few

homologs with strong signal incompatibility. Our implementation uses a wODR model with regression

line through the origin by setting the Y-axis intercept to zero, which avoids overfitting the linear

regression model to the detriment of coherent evolutionary assumptions.

Algorithm explanation

Our I_{ES} implementation performs an initial wODR with pairwise comparisons including all gene

family representatives within each genome. For each genome, all pairs containing one homolog from each

assessed gene family are evaluated, and unique pairs that minimize the sum of squared residuals are

reported. As exemplified in Fig. 1, a hypothetical $gene_1$ occurs exclusively as single copy across 10

genomes (Fig. 1, $tree_1$), while $gene_2$ has an extra copy within genome J (Fig. 1, $tree_2$). To identify which

copy of $gene_2$ in J (i.e., j_1 or j_2) better represents their shared evolution we compare $gene_1$ pairwise

distances involving j with $gene_2$ pairwise distances involving j_1 and j_2. Consequently, to do that we must

duplicate J's rows and columns in $matrix_1$ to match $matrix_2$ dimensions (Fig. 1, $matrix_1$). The scatter plot

in Fig. 1 highlights pairwise distances involving j_1 in blue and j_2 in red, and as shown by the fitted wODR

regression, $gene_2$ pairwise distances involving j_1 fits better to the expected linear association between

$matrix_1$ and $matrix_2$ than pairwise distances involving j_2. The smallest sum of residuals obtained by the j_1

homolog of $gene_2$ correctly pairs it with J's $gene_1$ homolog, while j_2's $gene_2$ homolog is likely a product

of HGT from a shared common ancestor of A and B. When both gene families occur in multiples within

the same genome, all pairs of unique loci are reported. Once best matching genes from each gene family

are paired, or if both occur exclusively as single copy, a final wODR is performed using paired homologs

from each gene family and Y-axis intercept equal to zero. wODR is performed through the SciPy

(Virtanen et al. 2020) API of ODRPACK (Boggs et al. 1989). Initial weights of pairwise distance are

estimated as the inverse of residuals obtained from geometric distance regression with intercept equal to

zero and slope equal to s_Y/s_X, where s_Y and s_X are standard deviations from the regressed distance

matrices. Our method's capability to automatically pair copies of duplicated genes who best reflect the

shared history between two gene families vastly expands the scope of datasets fit to general evaluation of

evolutionary signal compatibilities. The presence of multiple gene copies within a genome constitutes a

key bottleneck to methods commonly used to assess the similarity of evolutionary histories. Tree-based

evolutionary distance assessment algorithms are not generally capable of pairing genes between two gene

families when at least one family contains multiple gene copies within genomes (Stamatakis 2006;

Nguyen et al. 2015; Gori et al. 2016; Huerta-Cepas et al. 2016). While Pearson r implementations either

rely on multiple tests (Gertz et al. 2003; Ramani and Marcotte 2003; Izarzugaza et al. 2008) or on

predicting structural interaction between gene products (Gueudré et al. 2016).

Fig. 1 Steps for I_{ES} estimation between gene families containing multiple gene copies. *tree₁* and *tree₂* are phylogenetic trees of two hypothetical gene families, *gene₁* and *gene₂*, respectively. *matrix₁* and *matrix₂* contain pairwise evolutionary distances between taxa from their respective gene families. The red arrows in *matrix₁* highlight the duplication of pairwise distances involving the *j* homolog of *gene₁* necessary to match dimensions of the two matrices. The wODR scatterplot displays the linear relationships between distances from both gene families, and highlights distances related to the *j₁* homolog of *gene₂* in blue and related to the *j₂* homolog in red. Arrows also highlight pairwise distances homologs in genomes *J* and *I* from both gene families.

Given that regression models only account for data points equally represented in both assessed variables, gene losses and duplications are not directly accounted for when comparing evolutionary histories through wODR. To incorporate unequal genomic occurrence between gene families to our proposed measurement of evolutionary similarity, the wODR Coefficient of Determination, i.e. R^2, is adjusted by the Bray-Curtis Index (I_{BC}). I_{BC} is defined as $1 - D_{BC}$, where is the D_{BC} is Bray-Curtis Dissimilarity (Bray and Curtis 1957) calculated from absolute genome counts in each gene family. From hereon we will refer to the wODR $R^2 \times I_{BC}$ product as I_{ES}. Supplementary Fig. S12 demonstrates wODR R^2 overestimation caused by the decrease in taxa overlap between two gene families, as well as the importance of I_{BC} adjustment. One random taxon was removed from each of two simulated gene families whose evolutions diverge by five Subtree Prune and Regraft (SPR) transformations. As the set of taxa

used during the regression becomes unrepresentative of underlying evolutionary processes, estimates

based on wODR R^2 tend to artificially increase.

Continuing with the example depicted in Fig. 1, despite *gene1* and *gene2* identical genomic

occurrence, their copy numbers diverge within genome *J*, which as mentioned before, arose from a

horizontal exchange of *gene2*. To reflect this difference in evolutionary events within gene family

histories in the proposed I_{ES}, the resulting wODR $R^2 = 1$ is adjusted using an $I_{BC} = 0.95$.

Statistics and data analysis

Pandas Python library (McKinney 2010) was used to manipulate pairwise distance matrices and

for generating condensed versions of the matrices submitted to wODR model. Effect size (*f*) hypothesis

tests of differences between distributions were obtained using Common Language statistics (McGraw and

Wong 1992), and p-value correction for multiple tests was performed using False Discovery Rate

implementation in StatsModels Python library (Seabold and Perktold 2010).

Enrichment of gene families within sets of genomes were assessed using hypergeometric tests

and p-values corrected with Benjamini-Hochberg's False Discovery Rate and expressed as q-values.

Data Simulation

We constructed ten simulated datasets, each one containing 50 trees generated from stepwise

random SPR transformations. These datasets were designed to represent sets of trees with similar

topologies reflecting a shared evolutionary history, perturbed by both phylogenetic noise and HGT. Each

dataset contains one initial random rooted tree with 50 taxa generated by ETE3 (Huerta-Cepas et al.

2016). To obtain the remaining 49 trees, the initial tree (*tree_1*) undergoes a series of 49 consecutive SPR

transformations in such a way that *tree_1* differs from *tree_2*, *tree_3*, and *tree_n* by 1, 2, and *n* SPR

transformations, respectively. At each SPR the branch leading to the regrafted clade undergoes two

transformations to simulate changes in substitution rate after an HGT event. The first transformation

multiplies the branch length by a random uniform variable ranging from 0 to 1, simulating at which point

during the branch's history the transfer occurred. The second transformation multiplies by a random

gamma distributed variable ($\alpha = \beta = 100$), simulating changes in substitution rates in the recipient clade

after said transfer. All simulated trees are available in Supplementary Material.

All simulated trees were also used to generate sequence simulations using INDELible (Fletcher

and Yang 2009) (Supplementary Material). Phylogenetic trees and pairwise distance matrices were

reconstructed using IQTree (Nguyen et al. 2015) using the LG+G model.

Archaeal empirical dataset

Complete genome sequences of 42 Archaea from the Euryarchaeota phylum and from TACK,

DPANN, and Asgardarchaeota groups were downloaded from NCBI GenBank (Supplementary Table

S1). Other candidate phyla known from metagenomic as well as remaining members of the DPANN

group were not included, as their expected phylogenetic relationships are not as well understood.

Clustering of homologous proteins was performed using the orthoMCL (Li et al. 2003) implementation

available in the GET_HOMOLOGUES package (Contreras-Moreira and Vinuesa 2013; Vinuesa and

Contreras-Moreira 2015). Archaea were selected as the test dataset since the evolutionary relationships

between some major groups are well-established, while others remain contested. Furthermore, many sets

of archaeal metabolic genes have a strong phyletic dependence (e.g., methanogenesis among

Euryarchaeota (Borrel et al. 2013)), therefore facilitating a clear assessment of similarities between

evolutionary signals of genes at different phylogenetic distances. Evolutionary similarity comparisons

were restricted to homologous groups present in at least 10 genomes.

Pairwise maximum likelihood distances between homologous proteins were generated using

IQTree under the LG+G evolutionary model. Phylogenetic trees from clusters of gene families with

compatible evolutionary signal (CES) and extended core genome (i.e., single copy and present in at least

35 out of the 42 sampled Archaea genomes) were reconstructed from concatenated multiple sequence

alignments using the LG+C60+F+G and individual partitions corresponding to each concatenated gene.

Enrichment of gene functions among CES clusters were performed using StringDB API

(Szklarczyk et al. 2019). For each genome, homologs from CES gene families were submitted

independently for enrichment assessment. Retrieved protein annotations are available in the

Supplementary Material.

Geodesic and Robinson-Foulds distance calculations

D_{geo} between single copy gene families (both simulated and real datasets) were calculated using

the treeCl Python package (Gori et al. 2016). RF distances between single copy gene families were

calculated by ETE3.

Results and Discussion

Simulated dataset

Evolutionary histories between simulated gene families were compared to each other using three

distinct metrics: RF, D_{geo}, and I_{ES}. Results reported by all three approaches successfully identified the

monotonic increase in SPR operations from a starting tree (Fig. 2 and Supplementary Fig. S1).

Measurements obtained from RF and D_{geo} approaches, however, frequently overestimated the impact of

SPR transformations between two gene families, leading to a fast saturation of dissimilarities between

evolutionary histories (Fig. 2B and Fig. 2C). The dissimilarity saturations detected by RF and D_{geo}

measurements occur as they fail to identify the decreasing similarity between two trees separated by more

than 15 to 20 SPR transformations, or even 10 SPR in some replicates (simulation replicate #1,

Supplementary Fig. S1). Both of these approaches rely on the proportion of compatible bipartitions

shared by two trees, which is very susceptible to small changes at deep bipartitions, where changing one

single leaf can potentially create fully incompatible bipartition tables.

Fig. 2 Scatter plot of evolutionary similarity metrics against number of SPR transformations between simulated gene families

from all ten replicates. Solid black lines are estimated from OLS regressions between number of SPR transformations and

evolutionary similarity metrics. All three scatter plots display the number of SPR transformations between two trees in the X-

axis, while varying the evolution similarity metric displayed in the Y-axis. A) displays wODR R^2 between distance matrices of

simulated gene families in the Y-axis, B) displays geodesic distances between trees reconstructed from each simulated alignment,

and C) displays RF distances estimated from the same trees.

In contrast, I_{ES} estimates are less susceptible to overestimating the impact of shifts in topology of

an underlying tree, displaying a robust linear correlation with the number of SPR transformations between

gene families ($\bar{r} = -0.87$, Fig. 2A). The lower level of information assessed by I_{ES}, pairwise distance

matrices instead of dichotomic trees, is less susceptible to dissimilarity saturation, corresponding to a

more linear relation between expected and observed changes in evolutionary histories. Furthermore, I_{ES}

is much more efficient, computationally. Tree reconstruction of the alignment simulated from *tree_1* of

the first simulation replicate (50 taxa and 500 sites with no indels) under LG+G model by IQTree took 80

seconds in a single thread, while the computation of the pairwise distance matrix for the same alignment

took 1.713 seconds. Both computations were performed on a 3 GHz Intel Xeon W processor. The

difference in computing time of almost 50x, without bipartition support assessment, shows another,

practical advantage for assessing evolutionary similarity through I_{ES} in large datasets.

Robustness assessment between approaches

The dichotomic pattern in a cladogram is extremely susceptible to uncertainties in phylogenetic

reconstruction, combined with the vast tree space available for 50 taxa, causing noise-induced topological

variations to be not directly distinguishable from real deviations in evolutionary history (Szöllosi et al.

2013). The simpler information used to estimate I_{ES} (i.e., pairwise Maximum Likelihood distances) is less

prone to such uncertainty, as it bypasses forming hypotheses about the evolutionary relationships between

taxa. This assumption is corroborated by pairwise comparisons within bootstrap replicates, where I_{ES}

correctly detected replicates as such, i.e., virtually identical to each other, while RF and D_{geo} measures

failed to identify the common nature of bootstrap replicates. In addition to its accurate predictions, I_{ES}

consistently displayed very little variance within its estimates between bootstrap replicates.

Each alignment of simulated sequences was used to generate 10 bootstrap replicates. Pairwise

comparisons between 10 bootstrap replicates summed up to 45 comparisons within a single alignment.

Given that we simulated a total of 500 alignments, we assessed 22,500 pairwise comparisons between

bootstrap replicates across all simulated datasets. I_{ES} values correctly identified bootstrap replicates as

sharing virtually identical evolutionary histories, $\overline{I_{ES}} = 0.96$, and did so very consistently (CV = 1.39%,

where CV stands for Coefficient of Variation). Despite successfully identifying increasing evolutionary

changes between simulated trees, RF distances inconsistently predicted similarities between histories of

bootstrapped trees ($\overline{RF} = 18\%$ and CV = 57.45%), as variations in bootstrapped alignments caused small

perturbations in reconstructed tree topologies, and subsequent underestimation of evolutionary similarity

between bootstrap replicates. While D_{geo} estimates are more likely to overestimate small differences

between trees than RF, and consequently are more prone to saturation, geodesic distance estimates

displayed substantially less variation than RF ($\overline{D_{geo}} = 1.25$ and CV = 14.97%). D_{geo} cannot be directly

normalized or translated to a proportion of incompatibility between trees.

Evolutionary similarities within archaeal gene families

In order to test I_{ES} performance when estimating shared phylogenetic signal in an empirical set of

gene families, we evaluated 1,322 families of homologous proteins assembled from annotated CDSs

extracted from 42 archaeal genomes (Supplementary Table S1). This empirical dataset contains conserved

and accessory gene families with different sizes due to gene losses, duplications, and transfers.

I_{ES} was estimated for all pairwise combinations of gene families present in at least 10 genomes,

with 2,142 out of 748,712 comparisons having I_{ES} values of at least 0.7. Pairs of gene families with an

$I_{ES} \geq 0.7$ were added as nodes to a weighted network with its estimated I_{ES} value as an edge connecting

both gene families. In total 419 unique archaeal gene families were added to the network, while the

remaining 903 gene families did not display any $I_{ES} \geq 0.7$ with other gene families. A 0.7 threshold was

selected based on early applications of Pearson's *r* to evolutionary distances (Goh et al. 2000; Pazos and

Valencia 2001). The resulting evolutionary similarity network (Fig. 3) is heavily imbalanced, with just

11% of nodes involved in 50% of network edges. The majority of gene families (68%) did not display any

I_{ES} above the 0.7 threshold with other gene families, suggesting a general incompatibility between

evolutionary signals, or lack thereof to detect its compatibility with others. However, the high edge

concentration within just a few nodes suggests a strong central signal (Puigbò et al. 2009) present among

few gene families, from which the evolutionary trajectories of others have diverged. Similarities between

evolutionary signals, as estimated by I_{ES}, are strongly associated with genomic linkage ($p = 6.18e^{-75}$

and $f = 0.86$). Gene families frequently occurring in each other's genomic vicinity (i.e., fewer than

10,000 bp apart in at least 21 genomes) displayed significantly greater I_{ES} relative to pairs of gene

families that were further apart (i.e., more than 100,000 bp apart in at least 21 genomes) (Fig. 4a).

Fig. 3 Compatible evolutionary signal network, each node represents a gene family, and edges connecting nodes represent shared

evolutionary history ($I_{ES} \geq 0.7$). Nodes in the same colors are identified as undergoing similar evolutionary trends by Louvain

community detection. Triangular nodes represent single-copy genes, and circular ones are gene families containing gene

duplications. Clusters of CES gene families with less than ten members are not represented.

Fig. 4 A) distributions of I_{ES} between pairs of genes within 10,000 bp of each other in blue and between pairs of genes apart by at least 100,000 bp in orange. Neighboring gene pairs displayed significantly more similar evolutionary signals than non-neighboring gene pairs. B) ratio between the proportion of gene pairs intra and inter CES clusters, Y-axis, occurring within genomic windows, X-axis. 100 window sizes were assessed ranging from 1,000 bp to 1,000,000 bp.

Clusters of gene families with compatible evolutionary signal

Evolutionary trends shared across gene families were assessed using Louvain community detection (Blondel et al. 2008), reporting 41 CES clusters, of which 25 comprise only two gene families and eight major clusters contain ten or more similarly evolving gene families (Fig. 6). Although the assumption of compatibility between evolutionary signals based on I_{ES} estimates and the clustering process are agnostic of specific shared evolutionary events or their causes, each of these clusters is expected to comprise gene families sharing common evolutionary trends and paths. These shared evolutionary trends are corroborated by the very pronounced association between genomic linkage and estimated compatibility between evolutionary signals. Across small nucleotide distances between loci, linkage is a strong predictor of CES relations between genes, but its predictive power rapidly decreases as the number of nucleotides between two given loci increase (Fig. 4b), displaying a linear log-log relationship (Supplementary Fig. S2). Comparisons between intra- and inter-cluster genomic linkage showed that the proportion of CES genes within 1,000 bp of each other is three times the proportion of non-CES genes within the same window. Increasing the surveilled genomic window decreases the difference between proportions; within a 10,000 bp window, the ratio of CES genes is reduced to 1.8 the ratio of non-CES, and at a 100,000 bp window this difference in proportions falls to 1.2 (Fig. 4b).

Among the eight CES clusters with ten or more gene families, four are comprised of mostly core genes, and four are composed of mostly accessory genes (Fig. 6). The four CES clusters of core genes

(cluster#2, cluster#3, cluster#4, and cluster#5) are promising candidates for reconstructing the average

phylogenetic signal present within sampled Archaea. These four core CES clusters are composed of 102

extended core genes (single copy and present in at least 35 genomes) and 146 broadly distributed gene

families present on average in 33 genomes, both as single and multiple copies. CES clusters of accessory

genes (cluster#0, cluster#1, cluster#8, and cluster#15 in Fig. 6) include specific archaeal clades, but do

not map to well-established phylogenetic relationships; rather, they show polyphyletic gene distributions,

likely caused by HGTs and/or gene losses shared by CES gene families. For example, cluster#0 is well

represented amongst Euryarchaeota and hyperthermophilic TACK; cluster#15 comprises gene families

with shared evolutionary trends mainly occurring within Crenarachaeota and hyperthermophilic

Euryarchaeota; CES accessory gene families in cluster#1 and cluster#8 display congruent signals tying

methanogenic Euryarchaeota with Thaumarchaeota and Asgardarchaeota, respectively. Besides the eight

CES clusters with ten or more gene families, the CES network community detection yielded 34 other

clusters containing between two and nine gene families (Supplementary Material). These 34 CES clusters

contain a total of 88 gene families, whose degree centralities are much smaller than the 331 gene families

within the eight major CES clusters (averages of 1.36 and 17.79, respectively).

Fig. 5 Heatmap of enriched KEGG Pathways, columns, within CES clusters of gene families, rows. Shades of red represent the

proportion of genomes with detected KEGG Pathway enrichment within its homologs of CES gene families. Columns and rows

were clustered using complete linkage and correlation coefficients. KEGG Pathways enriched in less than 10% of genomes in

which CES genes occur are not reported. Cluster#15 did not report significant enrichment of KEGG Pathways.

CDSs from 21 out of 42 sampled genomes have functional annotation available in StringDB (Supplementary Material), and through its API we identified annotated KEGG Pathways enriched within homologs of CES gene families from each genome. In the dendrogram and heatmap depicted in Fig. 5 we clearly identify two sets of opposing CES clusters of gene families: accessories (top three rows) and core (bottom four rows), and their associations with genetic information processing and metabolism KEGG Pathways (indicated by column color in the top row). All four CES clusters of core gene families are enriched with KEGG Pathways related to genetic information processing (e.g., Ribosome, DNA replication, and Aminoacyl-tRNA biosynthesis). CES clusters of accessory gene families on the other hand tend to be enriched with KEGG Pathways related to metabolism (e.g., Methane metabolism, Microbial metabolism in diverse environments, and Biosynthesis of antibiotics in Fig. 5). It is also important to emphasize the opposite pattern of enrichment and depletion of KEGG Pathways between clusters of core and accessory genes. KEGG Pathways related to metabolism display minor enrichment signal within CES clusters of core gene families, and KEGG Pathways related to genetic information processing are not enriched within clusters of accessory genes (Fig. 5). CES clusters of accessory genes comprised within cluster#1, whose occurrence is restricted to methanogenic Euryarchaeota and Thaumarchaeota, are enriched for methane metabolism within six genomes. Similarly, gene families from cluster#8, restricted to methanogenic Euryarchaeota and Asgardarchaeota, are also enriched in methane metabolism in five genomes (Fig. 5).

Compatible evolutionary signal clusters and possible vertical evolutionary signals

Phylogenies generated from extended core genomes are generally used as reasonable proxies of the species-tree phylogeny, given the assumption that these genes are less likely to undergo HGT between distantly related groups. However, an extended core phylogeny may not represent the species tree for several reasons, including systematic biases in phylogenetic reconstruction due to shared compositional bias, or strong biases in HGT partners among sets of genes. Nevertheless, the extended core tree can still be used as an adequate representation of the consensus evolutionary signal detected in the sampled archaeal genomes, the closest thing we have to the simple "null hypothesis" of a shared history due to vertical inheritance. The 102 genes composing the extended core genome are not equally distributed across CES clusters (Fig. 3), e.g., cluster#4 contains the greatest number of extended core genes, 44 out of 62 gene families, followed by cluster#3 with 27 among its 111 gene families. The split of the extended core genome into four distinct major CES clusters (Fig. 3) suggests differing sets of HGTs among core genes, creating conflicting evolutionary histories between genes from different clusters (Fig. 7). Closeness centrality measures ($\tilde{C} = 0.56$) and node strength corrected by cluster size ($\tilde{S} = 0.19$) suggest that cluster#4 gene families share stronger and more cohesive evolutionary trends than gene families from other clusters (Supplemental Fig S3). Therefore, cluster#4 contains the set of genes that may be best able

to approximate the average evolutionary signal detected among sampled Archaea as well the most

compatible to each other. Cluster #4's evolutionary history is also the most similar to that inferred from

the extended core genome (Fig. 6 and Fig. 7).

While binning genes with distinct evolutionary histories permits phylogenetic reconstructions less

likely to be subject to spurious signals arising from the averaging of conflicting evolutionary signals, the

resulting phylogenies remain susceptible to phylogenetic reconstruction artefacts. For example, since I_{ES}

is not estimated from phylogenies but from pairwise evolutionary distances, we do not expect it to be

subject to Long Branch Attraction (LBA) artefacts. Nevertheless, phylogenies reconstructed from sets of

genes with high I_{ES} between each other are as susceptible to LBA as any other dataset. By clustering gene

families with high I_{ES} we are able to minimize conflicting evolutionary signals between concatenated

genes, improving the robustness of the phylogenetic signal. Despite its robustness to phylogenetic

artifacts, I_{ES} estimates are still affected by sampling biases. The overrepresentation of specific taxonomic

groups can lead to underestimating deviations in the evolutionary history of less represented groups.

LBA is a frequently invoked in discussions of archaeal phylogeny, specifically with regard to the

phyletic status of the DPANN group (Brochier-Armanet et al. 2011; Raymann et al. 2014; Petitjean et al.

2015; Williams et al. 2015; Feng et al. 2019). Regardless of the set of genes used for phylogenetic

reconstruction, extended core genome or any of the four CES clusters of core genes, all resulting trees

depicted a well-supported DPANN clade composed of Nanohaloarchaea archaeon, *Ca.* Woesearchaeota

archaeon, *Nanoarchaeum equitans*, and *Ca.* Diapherotrites archaeon. To assess the impact of LBA in

reconstructing DPANN, we generated phylogenies from the CES clusters of core genes while using a

single DPANN taxon at a time. Each DPANN taxon showed distinct evolutionary trends across all CES

clusters of genes, suggesting both a more complex extended core gene history for these genomes, and that

the initial monophyletic grouping of DPANN in each case was, in fact, artifactual (Supplementary

Material).

Cluster#4 phylogenies individually testing the position of each DPANN taxa placed

Nanohaloarchaea sister to Halobacteria, with Nanohaloarchaea+Halobacteria being sister to

Methanomicrobia. Cluster#3 also reported Nanohaloarchaea sister to Halobacteria, but with both nested

within Methanomicrobia, rendering this group paraphyletic. This placement for Nanohaloarchaea has

been previously proposed by Narasingarao et al. 2012; Zhaxybayeva et al. 2013; Petitjean et al. 2015; and

Feng et al. 2019. The uncertain placement of Nanoarchaea has also been topic of investigation (Huber et

al. 2002; Brochier et al. 2005). Interestingly, each CES cluster recovered a different placement for

Nanoarchaea: sister to Euryarchaeota (cluster#2), sister to Korarchaeota+Crenarchaeota (cluster#3), sister

to Korarchaeota (cluster#4), and sister to Thermococcales (cluster#5). One of the most accepted

Nanoarchaea placements is as sister to Thermococccales (Brochier et al. 2005; Urbonavičius et al. 2007; Dutilh et al. 2014), which in our analyses was recovered only by cluster#5 (Supplementary Material).

While our tests further support that the monophyly of DPANN is likely due to LBA, we did not detect a significant LBA effect for Woesearchaeota and Diapherotrites. Except for Diapherotrites placing between Class I and II methanogens in cluster#2, phylogenies from all four clusters proposed both taxa grouping together as sister to Euryarchaeota, assuming an archaeal root between TACK+Asgard and Euryarchaeota (Supplementary Material). The disparate placements of DPANN members within trees from CES clusters also suggests that, in addition to LBA, the DPANN clade from the extended core genome phylogeny is further impacted by the heterogeneity of the phylogenetic signal. This may not only produce a "signal averaging" effect favoring a monophyletic DPANN deeper in the archaeal tree, but may also be a contributing factor to the LBA artifact itself. Heterogeneity among combined phylogenetic signals is likely to increase the estimated branch length, as the incorrect assumption of a single underlying phylogeny will lead to more homoplastic sites.

Assuming that a given set of genomes constitutes a monophyletic clade, it is also reasonable to expect a certain number of gene families to be overly represented within the clade and not readily available to genomes outside the clade. Regardless of the driving force behind the enrichment of gene families within a clade, inheritance from a common ancestor or biased HGT (Andam et al. 2010; Andam and Gogarten 2011), we identified 80 gene families enriched within TACK genomes and 111 within Euryarchaeota ($q \leq 0.05$). In contrast to the well accepted TACK and Euryarchaeota clades, we did not detect any number of genes enriched within the four sampled DPANN genomes, providing phylogenetic independent evidence against its monophyly. Complementarily, and in support of a Nanohaloarchaea+Halobacteria clade, we identified 78 genes present in Nanohaloarchaea archaeon enriched within the three Nanohaloarchaea and Halobacteria genomes. When compared to well accepted clades containing a similar number of sampled genomes to DPANN, 456 gene families are enriched within the five genomes within Thaumarchaeota. While the differing degrees of physiological, metabolic, and genetic diversity within these groups certainly influence the number of shared gene families, it remains striking that this particular signal of shared ancestry is conspicuously lacking in DPANN.

Common and distinct evolutionary trends between CES clusters

Among CES clusters of core gene families, cluster#4 and cluster#5 are most evenly represented across archaeal groups, while cluster#2 and cluster#3 are poorly distributed among DPANN (Fig. 6). All four CES clusters of core gene families have low frequency within Thaumarchaeota archaeon SCGC AB-539-E09, and only gene families from cluster#2 and cluster#4 are present in any substance in Thermoplasmatales archaeon SCGC AB-539-N05. All four clusters display very similar overall phylogenies calculated from concatenations of genes within each cluster, varying mainly within the

organization of Euryarchaeota (Fig. 6 and Supplementary Material). All four CES clusters of core genes reconstructed the monophyly of Euryarchaeota, with the exception of cluster#2, which placed *Pyrococcus furiosus*, *Thermococcus kodakarensis*, *Methanocaldococcus jannaschii*, *Methanothermobacter thermautotrophicus*, and *Methanopyruus kandleri* together as sister to Asgardarchaeota+TACK. Only cluster#4 recovered the monophyly of Methanomicrobia as sister to Halobacteria, with the other three CES clusters placing Halobacteria within Methanomicrobia.

All four core CES clusters robustly identified Asgardarchaeota as sister to TACK (Fig. 6), with small variation in the Asgardarchaeota phylogeny, and cluster#5 placed Korarchaeota at the base of the TACK super-phylum. When assessing all-versus-all I_{ES} between CES clusters of core genes, the evolutionary signal detected from cluster#4 is the least dissimilar to the other three (Fig. 7). This shortest path from cluster#4's evolutionary trajectory to others suggests that cluster#4 best approximates the average archaeal evolutionary history (Fig. 7). In general, the overall high I_{ES} estimates between core CES clusters suggest that despite composing distinct clusters, gene histories between clusters are generally congruent, with deviations reflecting small divergences potentially representing genes with specific sets of reticulate histories.

Phylogenetic trees obtained from accessory gene families in cluster#0, cluster#1, cluster#8, and cluster#15 reconstructed all represented archaeal phyla as monophyletic (except for *P. furiosus* in Euryarchaeota in cluster#0, Supplementary Fig. S5), suggesting a shared common origin of accessory genes from each CES cluster by all genomes from the same phylum. Although the monophyly of archaeal phyla within trees of CES clusters of accessory genes does not permit an accurate prediction of the directionality of possible inter-phyla HGTs, intra-phylum distances congruent to the supposed vertical inheritance signal can be used to evaluate inter-phylum distances under a wODR model (Supplemental Fig. S6, S8, S10, and S11). When compared to pairwise distances expected from vertical inheritance, inter-phylum distances that are significantly shorter than estimates obtained from intra-phylum distances may be attributed to HGT acquisition by one of the phyla in question. For each CES cluster of accessory genes, we assessed wODR of its pairwise distances against the vertical evolution estimated from cluster#4.

When comparing pairwise distances obtained from cluster#1 against cluster#4, distances between Euryarchaeota and Thaumarchaeota are consistently placed below the estimated regression line (Supplementary Fig. S6 and S7). This suggests that cluster#1 genes were horizontally transferred between ancestors of both phyla, causing shorter evolutionary distances between phyla than expected if their homologs diverged exclusively by vertical inheritance.

Inter-phyla distances between Euryarchaeota and Crenarchaeota obtained from cluster#0 fit the evolutionary rate expected using intra-phylum distances for this CES cluster (Supplementary Fig. S8),

suggesting that homologs from both phyla were vertically inherited from a common ancestor. On the

other hand, cluster#0 inter-phyla distances involving Thaumarchaeota (Crenarchaeota to Thaumarchaeota

and Euryarchaeota to Thaumarchaeota) are shorter than expected from the wODR using intra-phylum

distances (Supplementary Fig. S8) and display significantly greater residuals than distances between

Crenarchaeota and Euryarchaeota (Supplementary Fig. S9). The absence of cluster#0 genes among

Asgardarchaeota and Korarchaeota and the short inter-phyla distances to Thaumarchaeota homologs

suggest an extensive loss among missing clades and horizontal acquisition by the thaumarchaeal ancestor

from either crenarchaeal or euryarchaeal donors.

Despite the occurrence of accessory genes from cluster#1 and cluster#8 in methanogenic

Euryarchaeota (Fig. 6) and the enrichment of methane metabolism pathways (Fig. 5), evolutionary

histories of both CES clusters are not related (Fig. 3). Gene families in CES cluster#8 did not display

$I_{ES} \geq 0.7$ outside its own cluster, constituting a separate connected component in the CES network

depicted in Fig. 3. That said, cluster#8 gene families display shorter Euryarchaeota-Asgardarchaeota

distances when compared to cluster#4 distances, but unlike cluster#0 and cluster#1, intra-

Asgardarchaeota and intra-Euryarchaeota pairwise distances are not mutually compatible under a single

linear regression (Supplemental Fig. S10). The lack of a strong wODR anchor in the form of intra-phyla

distances suggests a more complex horizontal exchange history of cluster#8 genes, possibly involving

intra-phylum HGTs, which we cannot accurately assess with the dataset used in this study. CES

cluster#15 of accessory genes is well distributed among Crenarchaeota, and its intra-phylum pairwise

distances are congruent to cluster#4 distances, but their patchy occurrence among Euryarchaeota and

Korarchaeota (Fig. 6) does not permit a confident assessment of this cluster's evolutionary history

(Supplementary Fig. S5).

Fig. 6 Phylogenetic tree of Archaea reconstructed from 62 genes within CES cluster#4. The phylogeny was obtained using

LG+F+G+C60 evolutionary model from IQTree and each gene had its parameters independently estimated according to

parameter "-sp". Bipartition supports were estimated using UFBoot and aLRT, each with 1,000 replicates, and bipartitions well

supported by both methods are colored in green (UFBoot ≥ 95% and aLRT ≥ 80%), while bipartitions well supported by a

single method are colored in yellow. Red dotted lines indicate Nanohaloarchaea (1) and Nanoarchaea (2 and 3) placements

reconstructed within phylogenies containing a single DPANN genome at a time. Despite the lack of outgroups to Archaea within

our sample the tree is rooted in DPANN for the sake of visualization. The associated heatmap reflects the representation of gene

families within CES clusters amongst archaeal genomes.

3 *Fig. 7* Scatter plots of pairwise evolutionary distances reconstructed from each widely distributed gene family versus each other

4 in blue. And in red, scatter plots of pairwise evolutionary distances reconstructed from each widely distributed gene family versus

5 pairwise evolutionary distances reconstructed from 102 extended core gene families. Similarities between evolutionary histories

6 of pairs of CES clusters, and between CES clusters and extended core genes were estimated by I_{ES}.

Conclusions

We have presented I_{ES} , a new, robust, and efficient method to determine gene families with

compatible evolutionary histories that are good candidates to be used in phylogenetic tree reconstructions

of a set of organisms. The distance regression basis of our proposed method does not require hypotheses

regarding evolutionary relationships between extant and ancestral taxa represented by the branching

pattern of phylogenetic trees. Besides significant gains in accuracy of its estimates and computing

efficiency, I_{ES} introduces a new and robust approach to pair members of gene families that best represent

their shared evolutionary trends. The strong association between genomic linkage and I_{ES} estimates

within archaeal genomes constitutes independent evidence of the ability of I_{ES} to recover shared

evolutionary histories within empirical datasets.

One major consequence of shared evolutionary trends by gene families is that the set of genomes

in which a given gene family occurs should be similar to genomes in which genes of other families with

compatible signals occur. Despite similar performances of Pearson's r and wODR R^2 in detecting these

trends, I_{ES} achieves the same result in a more efficient way. The utilization of wODR also imparts more

robust statistical support not directly available to previous Pearson's r implementations, whereas the

assessment of pairwise distances between taxa provides robustness in the presence of artefacts associated

to phylogenetic inference (e.g., Long Branch Attraction). The ability to assess residuals of each datapoint

independently also allows for evaluations of specific homologs, a useful tool for HGT detection. I_{ES} can

thus be incorporated into phylogenomics pipelines and used to guide the selection of gene families for

more accurate and robust species-tree inference, as well as the detection of meaningful clusters of gene

families evolved in shared, yet reticulate, patterns.

By assessing similarities of evolutionary signal between archaeal gene families using I_{ES} we were

able to detect several clusters of shared, distinct gene histories. Phylogenetic reconstruction using

concatenated sequences from each of the four major CES clusters of core genes confirmed these distinct

evolutionary histories. The phylogeny resulting from CES cluster#4, in particular, recovers a species tree

hypothesis consistent with that proposed in several other studies, while using a more empirically-

supported selection of gene families that does not presuppose vertical inheritance, an improved alternative

to analyses limited to conserved single-copy core genes.

Besides its impact in improving the reconstruction of the Archaeal Tree of Life, CES clusters

obtained from I_{ES} provided key evidence about horizontal exchange between phyla of functionally related

genes (Supplementary Fig. S5). For example, given the almost exclusive occurrence of genes from CES

cluster#1 among methanogenic Euryarchaeota and Thaumarchaeota, tree-based approaches are not

capable to report the possible HGT between both phyla. Separately, intra- and inter-phyla distances

obtained from CES cluster#1 are strongly correlated to distances described in CES cluster#4, however the

significant placement of inter-phyla distances bellow the wODR line strongly suggests an HGT between

ancestors of both phyla.

The method we used to analyze the archaeal gene sets is general and can thus be applied to any

other set of genomes. Furthermore, the I_{ES} implementation described provides a straightforward

framework for future improvements and a possible alternative to phylogenetic reconciliation approaches

to identify HGT events, as described in Supplementary Figs. S5-S11.

Acknowledgements

This work was supported by Simons Foundation Collaboration on the Origins of Life Award #339603 and NSF Integrated Earth Systems Program Award #1615426 to GPF. SMS was supported through Geisel School of Medicine at Dartmouth's Center for Quantitative Biology through a grant from the National Institute of General Medical Sciences of the National Institutes of Health under Award Number P20GM130454. JCS was funded in part by a CNPq senior researcher fellowship.

References

Andam CP, Gogarten JP. 2011. Biased gene transfer and its implications for the concept of lineage. Biol. Direct [Internet] 6:47. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3191353&tool=pmcentrez&rendertype=abstract

Andam CP, Williams D, Gogarten JP. 2010. Biased gene transfer mimics patterns created through shared ancestry. Proc. Natl. Acad. Sci. U. S. A. [Internet] 107:10679–10684. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2890805&tool=pmcentrez&rendertype=abstract

Bapteste E, O'Malley M a, Beiko RG, Ereshefsky M, Gogarten JP, Franklin-Hall L, Lapointe F-J, Dupré J, Dagan T, Boucher Y, et al. 2009. Prokaryotic evolution and the tree of life are two different things. Biol. Direct [Internet] 4:34. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2761302&tool=pmcentrez&rendertype=abstract

Barker D, Pagel M. 2005. Predicting Functional Gene Links from Phylogenetic-Statistical Analyses of Whole Genomes.Murray D, editor. PLoS Comput. Biol. [Internet] 1:e3. Available from: https://dx.plos.org/10.1371/journal.pcbi.0010003

Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E. 2008. Fast unfolding of communities in large networks. J. Stat. Mech. Theory Exp. [Internet] 2008:P10008. Available from: http://stacks.iop.org/1742-5468/2008/i=10/a=P10008?key=crossref.46968f6ec61eb8f907a760be1c5ace52

Boggs PT, Byrd RH, Schnabel RB. 1987. A Stable and Efficient Algorithm for Nonlinear Orthogonal Distance Regression. SIAM J. Sci. Stat. Comput. [Internet] 8:1052–1078. Available from: http://epubs.siam.org/doi/10.1137/0908085

Boggs PT, Donaldson JR, Byrd R h., Schnabel RB. 1989. Algorithm 676: ODRPACK: software for weighted orthogonal distance regression. ACM Trans. Math. Softw. [Internet] 15:348–364.

Available from: http://dl.acm.org/doi/10.1145/76909.76913

Borrel G, O'Toole PW, Harris HMB, Peyret P, Brugère J-F, Gribaldo S. 2013. Phylogenomic Data
Support a Seventh Order of Methylotrophic Methanogens and Provide Insights into the Evolution of
Methanogenesis. Genome Biol. Evol. [Internet] 5:1769–1780. Available from:
https://academic.oup.com/gbe/article-lookup/doi/10.1093/gbe/evt128

Bray JR, Curtis JT. 1957. An Ordination of the Upland Forest Communities of Southern Wisconsin. Ecol.
Monogr. [Internet] 27:325–349. Available from: http://doi.wiley.com/10.2307/1942268

Brochier-Armanet C, Forterre P, Gribaldo S. 2011. Phylogeny and evolution of the Archaea: one hundred
genomes later. Curr. Opin. Microbiol. [Internet] 14:274–281. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1369527411000634

Brochier C, Gribaldo S, Zivanovic Y, Confalonieri F, Forterre P. 2005. Nanoarchaea: representatives of a
novel archaeal phylum or a fast-evolving euryarchaeal lineage related to Thermococcales? Genome
Biol. [Internet] 6:R42. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15892870

Contreras-Moreira B, Vinuesa P. 2013. GET_HOMOLOGUES, a versatile software package for scalable
and robust microbial pangenome analysis. Appl. Environ. Microbiol. [Internet] 79:7696–7701.
Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3837814&tool=pmcentrez&rendertype=
abstract

Dagan T, Martin W. 2006. The tree of one percent. Genome Biol. 7.

Dutilh BE, Cassman N, McNair K, Sanchez SE, Silva GGZ, Boling L, Barr JJ, Speth DR, Seguritan V,
Aziz RK, et al. 2014. A highly abundant bacteriophage discovered in the unknown sequences of
human faecal metagenomes. Nat. Commun. [Internet] 5:1–11. Available from:
http://www.nature.com/doifinder/10.1038/ncomms5498

Feng Y, Neri U, Gosselin S, Louyakis AS, Papke RT, Gophna U, Gogarten JP. 2019. Reconstructing the
evolutionary origins of extreme halophilic Archaeal lineages. bioRxiv [Internet]:2019.12.19.883488.
Available from: http://biorxiv.org/content/early/2019/12/20/2019.12.19.883488.abstract

Fletcher W, Yang Z. 2009. INDELible: A flexible simulator of biological sequence evolution. Mol. Biol.
Evol. 26:1879–1888.

Gertz J, Elfond G, Shustrova A, Weisinger M, Pellegrini M, Cokus S, Rothschild B. 2003. Inferring
protein interactions from phylogenetic distance matrices. Bioinformatics 19:2039–2045.

Goh CS, Bogan AA, Joachimiak M, Walther D, Cohen FE. 2000. Co-evolution of proteins with their
interaction partners. J. Mol. Biol. 299:283–293.

Gori K, Suchan T, Alvarez N, Goldman N, Dessimoz C. 2016. Clustering Genes of Common
Evolutionary History. Mol. Biol. Evol. [Internet] 33:1590–1605. Available from:

http://mbe.oxfordjournals.org/content/early/2016/02/17/molbev.msw038.short?rss=1

Gueudré T, Baldassi C, Zamparo M, Weigt M, Pagnani A. 2016. Simultaneous identification of specifically interacting paralogs and interprotein contacts by direct coupling analysis. Proc. Natl. Acad. Sci. U. S. A. [Internet] 113:12186–12191. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27729520

Huber H, Hohn MJ, Rachel R, Fuchs T, Wimmer VC, Stetter KO. 2002. A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. Nature [Internet] 417:63–67. Available from: http://www.pnas.org/lookup/doi/10.1073/pnas.1905179116

Huerta-Cepas J, Serra F, Bork P. 2016. ETE 3: Reconstruction, Analysis, and Visualization of Phylogenomic Data. Mol. Biol. Evol. [Internet] 33:1635–1638. Available from: https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/msw046

Izarzugaza JMG, Juan D, Pons C, Pazos F, Valencia A. 2008. Enhancing the prediction of protein pairings between interacting families using orthology information. BMC Bioinformatics [Internet] 9:35. Available from: http://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-9-35

Kimmel R, Sethian J a. 1998. Computing geodesic paths on manifolds. Proc. Natl. Acad. Sci. U. S. A. [Internet] 95:8431–8435. Available from: http://www.ncbi.nlm.nih.gov/pubmed/9671694

Koonin E V, Wolf YI, Puigbò P. 2009. The phylogenetic forest and the quest for the elusive tree of life. Cold Spring Harb. Symp. Quant. Biol. [Internet] 74:205–213. Available from: http://www.ncbi.nlm.nih.gov/pubmed/19687142

Kunin V, Goldovsky L, Darzentas N, Ouzounis CA. 2005. The net of life: reconstructing the microbial phylogenetic network. Genome Res. [Internet] 15:954–959. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15965028

Kupczok A, von Haeseler A, Klaere S. 2008. An exact algorithm for the geodesic distance between phylogenetic trees. J. Comput. Biol. [Internet] 15:577–591. Available from: http://www.ncbi.nlm.nih.gov/pubmed/18631022

Leigh JW, Susko E, Baumgartner M, Roger AJ. 2008. Testing Congruence in Phylogenomic Analysis. Syst. Biol. [Internet] 57:104–115. Available from: http://sysbio.oxfordjournals.org/cgi/doi/10.1080/10635150801910436

Li L, Stoeckert CJ, Roos DS. 2003. OrthoMCL: identification of ortholog groups for eukaryotic genomes. Genome Res. [Internet] 13:2178–2189. Available from: http://www.ncbi.nlm.nih.gov/pubmed/12952885

Liu C, Wright B, Allen-Vercoe E, Gu H, Beiko R. 2018. Phylogenetic Clustering of Genes Reveals Shared Evolutionary Trajectories and Putative Gene Functions.Dagan T, editor. Genome Biol. Evol. [Internet] 10:2255–2265. Available from: https://academic.oup.com/gbe/article/10/9/2255/5076814

McGraw KO, Wong SP. 1992. A common language effect size statistic. Psychol. Bull. [Internet] 111:361–365. Available from: http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-2909.111.2.361

McKinney W. 2010. Data Structures for Statistical Computing in Python. In: p. 56–61. Available from: https://conference.scipy.org/proceedings/scipy2010/mckinney.html

Mirarab S, Bayzid MS, Bossau B, Warnow T. 2014. Statistical binning improves species tree estimation in the presence of gene tree heterogeneity. Science (80-.). [Internet] 346. Available from: http://dx.doi.org/10.1126/science.1250463

Narasingarao P, Podell S, Ugalde JA, Brochier-Armanet C, Emerson JB, Brocks JJ, Heidelberg KB, Banfield JF, Allen EE. 2012. De novo metagenomic assembly reveals abundant novel major lineage of Archaea in hypersaline microbial communities. ISME J. [Internet] 6:81–93. Available from: http://www.nature.com/articles/ismej201178

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol. Biol. Evol. [Internet] 32:268–274. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25371430

Novichkov PS, Omelchenko M V, Gelfand MS, Mironov A a, Wolf YI, Koonin E V. 2004. Genome-wide molecular clock and horizontal gene ransfer in bacterial evolution. J. Bacteriol. [Internet] 186:6575–6585. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=516599&tool=pmcentrez&rendertype=abstract

Owen M, Provan JS. 2011. A Fast Algorithm for Computing Geodesic Distances in Tree Space. IEEE/ACM Trans. Comput. Biol. Bioinforma. [Internet] 8:2–13. Available from: http://ieeexplore.ieee.org/document/5396323/

Papke RT, Gogarten JP. 2012. Ecology. How bacterial lineages emerge. Science [Internet] 336:45–46. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22491845

Pazos F, Valencia A. 2001. Similarity of phylogenetic trees as indicator of protein-protein interaction. Protein Eng. 14:609–614.

Pellegrini M, Marcotte EM, Thompson MJ, Eisenberg D, Yeates TO. 1999. Assigning protein functions by comparative genome analysis: protein phylogenetic profiles. Proc. Natl. Acad. Sci. U. S. A. [Internet] 96:4285–4288. Available from: http://www.ncbi.nlm.nih.gov/pubmed/10200254

Petitjean C, Deschamps P, López-García P, Moreira D. 2015. Rooting the Domain Archaea by Phylogenomic Analysis Supports the Foundation of the New Kingdom Proteoarchaeota. Genome Biol. Evol. [Internet] 7:191–204. Available from: https://academic.oup.com/gbe/article-lookup/doi/10.1093/gbe/evu274

Popa O, Dagan T. 2011. Trends and barriers to lateral gene transfer in prokaryotes. Curr. Opin. Microbiol.

[Internet] 14:615–623. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21856213

Puigbò P, Wolf YI, Koonin E V. 2009. Search for a "Tree of Life" in the thicket of the phylogenetic forest. J. Biol. [Internet] 8:59. Available from: http://www.ncbi.nlm.nih.gov/pubmed/19594957

Ramani AK, Marcotte EM. 2003. Exploiting the co-evolution of interacting proteins to discover interaction specificity. J. Mol. Biol. 327:273–284.

Rangel LT, Marden J, Colston S, Setubal JC, Graf J, Gogarten JP. 2019. Identification and characterization of putative Aeromonas spp. T3SS effectors.Antunes LCM, editor. PLoS One [Internet] 14:e0214035. Available from: https://dx.plos.org/10.1371/journal.pone.0214035

Raymann K, Forterre P, Brochier-Armanet C, Gribaldo S. 2014. Global Phylogenomic Analysis Disentangles the Complex Evolutionary History of DNA Replication in Archaea. Genome Biol. Evol. [Internet] 6:192–212. Available from: https://academic.oup.com/gbe/article-lookup/doi/10.1093/gbe/evu004

Robinson DF, Foulds LR. 1981. Comparison of phylogenetic trees. Math. Biosci. [Internet] 53:131–147. Available from: http://linkinghub.elsevier.com/retrieve/pii/0025556481900432

Seabold S, Perktold J. 2010. statsmodels: Econometric and statistical modeling with python. In: 9th Python in Science Conference.

Shapiro BJ, Friedman J, Cordero OX, Preheim SP, Timberlake SC, Szabó G, Polz MF, Alm EJ. 2012. Population genomics of early events in the ecological differentiation of bacteria. Science [Internet] 336:48–51. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3337212&tool=pmcentrez&rendertype=abstract

Sorek R, Zhu Y, Creevey C, Francino M. 2007. Genome-wide experimental determination of barriers to horizontal gene transfer. Science (80-.). [Internet] 318:1449–1452. Available from: http://www.sciencemag.org/content/318/5855/1449.short

Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. Bioinformatics [Internet] 22:2688–2690. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16928733

Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, Simonovic M, Doncheva NT, Morris JH, Bork P, et al. 2019. STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. Nucleic Acids Res. [Internet] 47:D607–D613. Available from: https://academic.oup.com/nar/article/47/D1/D607/5198476

Szöllosi GJ, Rosikiewicz W, Boussau B, Tannier E, Daubin V. 2013. Efficient exploration of the space of reconciled gene trees. Syst. Biol. 62:901–912.

Thomas CM, Nielsen KM. 2005. Mechanisms of, and barriers to, horizontal gene transfer between bacteria. Nat. Rev. Microbiol. [Internet] 3:711–721. Available from: http://dx.doi.org/10.1038/nrmicro1234

Urbonavičius J, Auxilien S, Walbott H, Trachana K, Golinelli-Pimpaneau B, Brochier-Armanet C, Grosjean H. 2007. Acquisition of a bacterial RumA-type tRNA(uracil-54, C5)-methyltransferase by Archaea through an ancient horizontal gene transfer. Mol. Microbiol. [Internet] 67:323–335. Available from: http://doi.wiley.com/10.1111/j.1365-2958.2007.06047.x

Vert J-P. 2002. A tree kernel to analyse phylogenetic profiles. Bioinformatics [Internet] 18 Suppl 1:S276-84. Available from: http://www.ncbi.nlm.nih.gov/pubmed/12169557

Vinuesa P, Contreras-Moreira B. 2015. Robust Identification of Orthologues and Paralogues for Microbial Pan-Genomics Using GET_HOMOLOGUES: A Case Study of pIncA/C Plasmids. In: p. 203–232. Available from: http://link.springer.com/10.1007/978-1-4939-1720-4_14

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser W, Bright J, et al. 2020. SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat. Methods [Internet] 17:261–272. Available from: http://www.nature.com/articles/s41592-019-0686-2

Williams TA, Heaps SE, Cherlin S, Nye TMW, Boys RJ, Embley TM. 2015. New substitution models for rooting phylogenetic trees. Philos. Trans. R. Soc. B Biol. Sci. [Internet] 370:20140336. Available from: https://royalsocietypublishing.org/doi/10.1098/rstb.2014.0336

Zhaxybayeva O, Stepanauskas R, Mohan NR, Papke RT. 2013. Cell sorting analysis of geographically separated hypersaline environments. Extremophiles [Internet] 17:265–275. Available from: http://link.springer.com/10.1007/s00792-013-0514-z