Multiple neuronal networks coordinate *Hydra* mechanosensory behavior

Krishna N. Badhiwala[1], Abby S. Primack[2], Celina E. Juliano[2], Jacob T. Robinson*[1,3,4]

1. Department of Bioengineering, Rice University, 6100 Main Street, Houston, Texas 77005
2. Department of Molecular and Cellular Biology, University of California, Davis, CA, USA.
3. Department of Electrical and Computer Engineering, Rice University, 6100 Main Street, Houston, Texas 77005
4. Department of Neuroscience, Baylor College of Medicine, One Baylor Plaza, Houston, Texas, 77030

*Correspondence to: jtrobinson@rice.edu.

Abstract

Hydra vulgaris is an emerging model organism for neuroscience due to its small size, transparency, genetic tractability, and regenerative nervous system; however, fundamental properties of its sensorimotor behaviors remain unknown. Here, we use microfluidic devices combined with fluorescent calcium imaging and surgical resectioning to study how the diffuse nervous system coordinates *Hydra*'s mechanosensory response. Mechanical stimuli cause animals to contract, and we find this response relies on at least two distinct networks of neurons in the oral and aboral regions of the animal. Different activity patterns arise in these networks depending on whether the animal is contracting spontaneously or contracting in response to mechanical stimulation. Together, these findings improve our understanding of how *Hydra*'s

diffuse nervous system coordinates sensorimotor behaviors. These insights help reveal how

sensory information is processed in an animal with a diffuse, radially symmetric neural

architecture unlike the dense, bilaterally symmetric nervous systems found in most model

organisms.

Introduction

Discovering the fundamental principles of neural activity and behaviors requires studying the

nervous systems of diverse organisms. Animals have evolved different neural structures like the

nerve net (e.g. *Hydra*), nerve cords and ganglia (e.g. *C. elegans*, *Aplysia*, planaria), and brain

(e.g. *Drosophila*, zebrafish, rodents, and primates). Despite the vastly different structures, many

behaviors are conserved across species, including sensorimotor responses[1–6] and sleep[7–15]. By

comparing neural circuits that support similar behaviors despite different architectures, we can

discover organizational principles of neural circuits that reflect millions of years of evolutionary

pressure. While there are many potential organisms that would support this type of comparative

neuroscience, only a small group of animals have the qualities to support laboratory

experiments: short generation span, ease of breeding and manipulation in laboratory conditions,

small and compact size, optical transparency, and a well-developed genetic toolkit with a

complete spatial and molecular map of the nervous system.

Transparent, millimeter-sized animals in particular offer a number of advantages for

neuroscientists because it is possible to image neural activity throughout the entire nervous

system using genetically-encoded calcium or voltage-sensitive fluorescent proteins[16–25]. In

addition, some millimeter-sized animals are compatible with microfluidic devices for precise

environmental control and microscopy techniques that offer cellular-resolution functional

imaging of the entire nervous system. These properties, combined with genetic tractability,

provide a powerful way of revealing neuronal dynamics across the entire nervous system (not

just a small region) during behaviors. For instance, whole nervous system imaging of confined

or freely moving animals has revealed the neuronal dynamics underlying locomotion[19,26] and

sensory motivated global state transitions in *C. elegans*[27,28], responses to noxious odor and

visuomotor behaviors in zebrafish[19–24], responses to light and odor in *Drosophila*[18,29], and

neuronal ensembles correlated with basal behaviors and response to light in *Hydra*[30,31].

Hydra is unique among the small, transparent organisms discussed above due to its

regenerative ability and highly dynamic nervous system. While most small, transparent model

systems (like *C. elegans* or zebrafish larvae) suffer permanent behavioral deficits from the loss

of one or a few neurons[32–36], *Hydra* can completely recover from a significant neuronal loss to

regain normal contractile behavior in as little as ~ 48 hours[37–39]. This radially symmetric

freshwater cnidarian has a nervous system composed of two diffuse networks of neurons, one

embedded in the endoderm and another embedded in the ectoderm[40,41]. While *Hydra's* diffuse

nerve net is highly dynamic with continuous cellular turnover and migration[42,43], regions with

increased neuron density resembling nerve rings have a comparatively lower neuronal turnover

(Figure 1a)[44–47]. One of these regions is in the oral end in the apex above the ring of tentacles

("hypostomal nerve ring"), and another is in the aboral end in the foot ("peduncle nerve ring")

(Figure 1a). Recent single-cell RNA sequencing has provided a complete molecular and spatial

map of the *Hydra* nervous system, including identification of unique cell-type specific biomarkers

to generate new transgenic models[48]. This existing molecular and spatial map of the nervous

system suggests there is no overlap in the neuronal cell types that make up the hypostomal and

peduncle nerve rings, and the distribution of the cell types varies along the length of the body

(Figure 1b). Finally, the demonstration of microfluidic and transgenic tools combined with

Hydra's dynamic yet 'simple' neural architecture has enabled observations of basal and sensory

motivated behaviors in the regenerating nervous system[31].

To better establish *Hydra* as a model organism for comparative neuroscience, it is critical to

understand their basic sensorimotor behaviors, such as response to touch. While it is well

documented that *Hydra* contract when mechanically agitated or poked with a pipette[49–53], we

found no quantitative reports of how this behavior depends on stimulus intensity or is mediated

by neural activity. Although significant insights in *Hydra* behavior have been made over the last

several decades with simple methodologies and manual observations, including that the

tentacles and/or the hypostome are needed for mechanosensory response, these experiments

lack quantitative characterization of neuronal or behavioral response. Forceps-induced touch

allows local stimulation, but it is difficult to control the force applied manually[54,55]. While

stimulation with mechanical agitation allows control over stimulus intensity, these observations

are limited to changes in body lengths[52].

Here, we use whole animal functional imaging combined with resection studies to discover that

despite the apparently diffuse nerve net in *Hydra*, these animals process sensorimotor

responses in specialized regional networks. To study the mechanosensory response, we first

developed a microfluidic system to apply a local mechanical stimulus and quantify *Hydra's*

behavioral and neural response. We then measured these responses in the absence of select

regions of the body and found at least one of the neuron-rich regions, the hypostome (oral) or

the peduncle (aboral), is required to coordinate spontaneous contractions, though the oral

network plays a more significant role. We found a significant reduction in the mechanosensory

response with the removal of the hypostome, the region where sensory information is likely

processed. These sensorimotor experiments combined with whole-animal neural and

epitheliomuscular imaging reveal that *Hydra* is capable of receiving sensory information along

the body column; however, the oral region is necessary for coordinating the motor response.

Results

***Hydra's* mechanosensory response is dependent on stimulus intensity**

To better understand sensory information processing in *Hydra*, we developed a double-layer

microfluidic system that can apply a local mechanical stimulus while we image the response of

the entire nervous system using fluorescence microscopy (Figure 2, Video 1). This local

mechanical stimulation is made possible by push-down microfluidic valves (Figure 2a) that

deliver mechanical stimuli to a portion of the *Hydra* body with precise temporal and spatial

control (see "Methods"). For all experiments, we pressurized a valve (400 μm diameter) that

was directly above the animal (for 1 sec every 31 sec, see "Methods") to stimulate the body

column while simultaneously performing functional calcium imaging (Figure 2b). We selected

the middle of the body for stimulation region to help ensure that we stimulated roughly the same

region of the animal throughout each experiment. This choice was based on the observation

that the body column region was relatively stationary, whereas the oral and aboral extremities

had large displacements during body contractions and elongations.

Experiments showed that this stimulation paradigm delivered a local mechanical stimulation with

most of the mechanical force localized to a radius of approximately 250 μm around the

microfluidic valve. To measure the locality of this stimulus we performed an experiment using

transgenic *Hydra* (nGreen)[48] expressing GFP pan-neuronally (and in neural progenitors) and

tracked the position and fluorescence intensity (GFP) from individual neurons during mechanical

stimulation (N = 222 neurons over 1 min). When the microfluidic valve was pressurized to

deliver mechanical stimulation, we found significantly increased average cellular (or tissue)

displacement (p < 0.001). Further analysis of the cellular movements showed the spatial

distribution of mechanical force from the stimulation was primarily experienced by the neurons

directly under the valve (Figure 2 – Figure Supplement 1). Specifically, we found that the tissue

directly below the valve was compressed (z direction) when mechanically stimulated - the

neurons directly under the valve had a small magnitude of lateral (x-y direction) displacement.

The tissue bordering the valve was stretched away from the valve center - the neurons in the

neighboring regions around the valve had the largest lateral displacement. This lateral

displacement decreased for neurons that were farther from the center of the valve. Neurons

more than 750 μm from the microfluidic valve center showed a negligible displacement of less

than 5 μm (95% CI lower bound = 5.8 μm), which is ~550% less than the displacement of

neurons bordering the valve.

Having established our method to provide local mechanical stimuli, we characterized *Hydra's*

sensitivity to local touch and the associated neural response. We performed experiments using

transgenic *Hydra* expressing GCaMP6s in neurons[30]. When we delivered mechanical stimuli,

we found bright calcium signals generated by a small number of neurons in the hypostome and

body column and a striking co-activation of many neurons in the ectodermal peduncle nerve ring

(Figure 2b, Video 2, 3). This nerve ring activity appeared as either a single bright calcium spike

(or "contraction pulse") or a volley of bright calcium spikes (or "contraction burst"). We also

found that the calcium-sensitive fluorescence averaged over a region of interest (ROI)

surrounding the peduncle faithfully represented the contraction pulses and bursts measured

from individual neurons (Figure 2 - Figure Supplement 2). When we analyzed single neuron

calcium dynamics from this peduncle nerve ring, we found extremely high correlated activity as

previously reported for contractions pulses and bursts (Figure 2 - Figure Supplement 2)[30,31].

Given the similarity of these data between large and small ROIs, we chose to use the peduncle

ROI to measure neuronal contraction bursts and pulses because it does not require single-

neuron tracking, which significantly increased the throughput of our data analysis. We further

confirmed that this signal is not the result of motion artifacts by measuring fluorescence from

Hydra (nGreen) that express GFP pan-neuronally using a similar ROI. In that case we did not

see the strong fluorescence signals associated with contraction pulses and bursts (Figure 2 –

Figure Supplement 3e-f). Body length proved to be an unreliable quantification of contractions

due to the stimulation artifacts (Figure 2 - Figure Supplement 3d); however, we were able to

accurately measure muscle activity associated with contractions by imaging calcium spikes in

the epithelial muscle cells (Figure 2 - Figure Supplement 4, 5 and Video 4, 5). Based on these

experiments, we define *Hydra's* "mechanosensory response" as calcium spikes in neural activity

from the peduncle ROI and the associated calcium spikes in the epithelial muscles from the

whole body if they occur within one second of mechanical stimulation onset (Figure 2c, d and

Figure 2 – Figure Supplement 6).

Using the neuronal fluorescence calcium imaging described above, we found that the probability

of the mechanosensory response depends on the intensity of the stimulus, which is consistent

with many psychometric functions (Figure 2e). *Hydra* were five times more likely to contract

within one second of receiving a strong mechanical stimulus than during a random one second

interval without a stimulus (stimulus valve pressure 20 and 25 psi; response probability = 0.60 ±

0.06 and 0.77 ± 0.08, mean ± SEM, respectively; no stimulus valve pressure = 0 psi; response

probability = 0.11 ± 0.01, mean ± SEM; Figure 2e, h). During mild stimuli, there was a slight

increase (~ 2x) in response probability above the spontaneous activity, although this increase

was not statistically significant compared to spontaneous contraction bursts or pulses (valve

pressure 10 and 15 psi; response probability = 0.23 ± 0.01 and 0.24 ± 0.03, mean ± SEM,

respectively; Figure 2e, h). We found that *Hydra* did not respond to a weak mechanical stimulus

that corresponded to a valve pressure of 5 psi (response probability = 0.09 ± 0.02, mean ±

SEM; Figure 2e, h).

Further analysis of the calcium activity pattern revealed that the single contraction pulses

(calcium spikes in the peduncle neurons) were more frequent when we repeated mechanical

stimulation every 31 seconds for one hour. While spontaneous contraction pulses or bursts

were observed roughly once every minute in microfluidics, when we stimulated *Hydra* with a

strong mechanical stimuli, we found that the frequency nearly matched the 31 sec between

stimuli (Interval between spontaneous contraction bursts or pulses = 72.98 sec ± 4.58, mean ±

SEM; 66.68 ± 5.73 sec median ± SE; Interval between stimulated contraction bursts or pulses =

20 psi, 38.28 ± 1.77 sec, mean ± SEM; 31.14 ± 2.21 sec, median ± SE; 25 psi, 32.20 ± 1.04

sec, mean ± SEM; 31.14 ± 1.30 sec, median ± SE; Figure 2f,h). While the majority of

spontaneous calcium spikes formed bursts, stimulated calcium spikes were roughly three times

more likely to be a single contraction pulses (Percentage of spontaneous spiking activity that is

a single contraction pulse = 0psi, 16.87 ± 5.38 % mean ± SEM; Percentage of stimulated

spiking activity that is a single contraction pulse = 20 psi, 56.84 ± 2.48%; 25psi, 62.50 ± 5.23 %,

mean ± SEM; Figure 2g, h).

***Hydra* sensitivity to mechanical stimuli is lowest near the aboral end**

Because of the diffuse neural architecture of *Hydra*, we expected each patch of *Hydra* tissue to

be equally responsive to mechanical stimuli. However, when we stimulated transgenic *Hydra*

expressing GCaMP6s (N = 3 whole animals, stimuliated 40 times per region with 22 psi) at three

different regions along their body (oral, middle body and aboral, stimulated 40 times at each

region with 22psi), we found the aboral end of *Hydra* to be less sensitive than the center of the

body (aboral region response probability = 0.1 ± 0.025; mid-body region response probability =

0.42 ± 0.025; mean ± SEM; $p < 0.01$; Figure 2 - Supplement figure 7a). Epitheliomuscular

calcium imaging (N = 8 whole animals expressing GCaMP7b in ectodermal epitheliomuscular

cells, stimulated 40 times in body column region with 22psi) also showed that the sensitivity to

mechanical stimulation generally decreases towards the aboral end of the *Hydra* (Figure 2 –

Supplement figure 7b). Furthermore, we found that the difference in sensitivity along the body

was not an artifact due to differently sized *Hydra* experiencing different pressures from the

microfluidic valves. We observed no statistically significant trend between animal size and

response probability (Figure 2 – Supplement figure 7c). These findings, combined with the

transcriptional analysis and in-situ hybridizations that indicate higher density of sensory neurons

in the oral half of the *Hydra*[48], suggest the oral end may be more sensitive to mechanical stimuli.

Unlike other organisms that have unique motor responses like reversals or acceleration

depending on the location of mechanical stimuli (e.g. *C. elegans*)[56,57], we observed that the

same motor program was initiated regardless of where on the body the animal was touched.

The only difference we observed was the fact that the response probability depended on where

along the oral and aboral axis we delivered the mechanical stimulus.

***Hydra*'s mechanosensory response is mediated by electrically-coupled cells**

Based on the latency of the mechanosensory response, we hypothesized that sensorimotor

information is transmitted by electrical activity in the *Hydra* and not by passive calcium diffusion

through ectodermal cells. This hypothesis is supported by the fact that aneural *Hydra,* which no

longer spontaneously contract in absence of stimuli, are capable of aversive contractile

response to touch, however, their responses are slow and require strong mechanical

stimulus[54,55]. To further test our hypothesis, we created a transgenic *Hydra* strain that expresses

the calcium indicator GCaMP7b (under the EF1α promoter, see "Methods") in the ectodermal

epitheliomuscular cells. With this transgenic line we measured contraction pulses and

contraction bursts by averaging calcium activity in all epitheliomuscular cells (Figure 2 – Figure

Supplement 3, 5). We hypothesized that if the mechanosensory response was primarily

mediated by calcium diffusion through epitheliomuscular cells, we would expect to see

propagation of calcium activity from the site of the stimulation. However, this was not the case;

we observed fast propagation of calcium activity throughout the entire ectoderm. Imaging

calcium activity in the peduncle neurons showed that changes in neural activity correlated with

the observed changes in epithelial muscle cells with increasing stimulus intensity (Figure 2 and

Figure 2 – Figure Supplement 5). This suggests that the mechanosensory response is indeed

mediated by the electrically-coupled cells. Although *Hydra*'s behavioral responses are slow

compared to other invertebrates, the 0.5 sec average response time in our data could not be

explained by calcium diffusion alone, which would take ~100 sec to travel the average distance

of ~0.5-1 mm between the stimulation site and the peduncle or hypostome (Figure 2 – Figure

Supplement 8a). Our data show the calcium signals from the peduncle neurons or ectodermal

muscles start increasing within ~0.1-0.2 sec following stimulation onset, reaching peak

fluorescence at ~0.5-0.6 sec regardless of the stimulus intensity (Figure 2 – Figure Supplement

8b-e). We found no difference in the calcium response times between neural and muscle

calcium imaging, though this could be influenced by the dynamics of the calcium indicator,

which typically cannot give information about latencies less than 50 ms[17].

Aboral neurons are not necessary for mechanosensory response in *Hydra*

Having quantitatively established *Hydra's* sensorimotor response to mechanical stimuli, we next

asked if specific regions of the nervous system play primary roles in mediating this response.

Patch-clamp electrophysiology of individual neurons in *Hydra* has thus far been unsuccessful

despite attempts by many research groups, and minimally invasive, cell-type specific

neuromodulation techniques such as optogenetics have yet to be developed for *Hydra*.

However, the animal's regenerative abilities allow us to resect large portions of tissue without

killing the animal, thus we can borrow from the tradition of lesioning brain regions to study their

functions[52,58–61].

Because the *Hydra* body plan is radially symmetric with cell types primarily varying along the

oral-aboral axis of the body column (Figure 1), we chose to make axial cuts across the body

column to remove select neuronal populations from the animal. Our rationale was that these

resections would remove entire or nearly entire groups of neuronal cell types. We then allowed

~6-12 hours for the animal to recover. This recovery time helps to reduce the confounding

contributions from initial tissue regeneration and allows animals to recover enough to tolerate

microfluidic immobilization. This period is long enough to allow the wounds to close, the

molecular response to injury to be completed[62,63], and the initial molecular events of

regeneration to start; but it is *not* long enough for the animal to regenerate lost neurons, which

takes approximately 30-72 hours (Figure 3 – Figure Supplement 1)[37,61,64,65]. We confirmed that

6-12 hours after resection the animals indeed showed loss of specific neuronal cell types by

measuring the expression levels of subtype-specific neuronal markers via qPCR (Figure 3 -

Figure Supplement 2). To limit the stress from microfluidic immobilization that could exacerbate

the resection wounds and affect activity, we shortened the duration of these experiments for the

majority of the animals (40 min total, 20 min of no stimulation, 20 min of stimulation - valve on 1

sec at 22 psi or 0 psi, off 30 sec). Only three animals per each condition (stimulated and non-

stimulated, and 5 different resections) were experimented on with a longer duration protocol as

used previously (100 min total, 20 min no stimulation, 60 min of stimulation - valve on 1 sec at

22 psi or 0 psi, off 30 sec, 20 min no stimulation; Figure 3 - Figure Supplement 3).

We began resection studies by removing the peduncle and basal disk to create a "footless"

Hydra. We hypothesized that aboral neurons may be important for coordinating and enhancing

body contractions (Figure 3). We based this hypothesis on the fact that aboral neuron activity

has a strong correlation with body contractions[30,31]. In addition, the neuropeptide Hym-176C has

been shown to induce ectodermal muscle contractions and is selectively expressed in the

ectodermal peduncle neurons[48,66–68]. Finally, the presence of gap junction protein innexin-2 in

aboral neurons could facilitate fast electrical conductions that allows these neurons to fire

synchronously[48,55]. This could be necessary for enhancing neuromuscular signaling for body

contractions. Because "footless" *Hydra* lacked peduncle neurons that we had used previously to

measure contraction pulses and bursts (Figure 3 - Supplement Figure 3d-f, and Video 6-9), we

performed these experiments using a transgenic *Hydra* line expressing GCaMP7b in the

ectodermal epitheliomuscular cells, which allowed us to measure contraction pulses and bursts

by averaging calcium activity in all the epitheliomuscular cells (using whole frame ROI which is

more robust to motion artifacts than peduncle ROI; Figure 2 - Figure Supplement 3, Figure 3a,b,

Figure 3 – Figure Supplement 3).

Because neurons in the foot fire synchronously with body contractions, we expected "footless"

animals to show significant changes in contraction behaviors (calcium spiking activity) to be

significantly affected by their removal[30,31,69], but this was not what we observed. Surprisingly, our

experiments with "footless" animals showed that the aboral nerve ring was not required to

regulate spontaneous contraction bursts or pulses or mechanosensory responses. After we

removed the peduncle network in *Hydra*, we found the increase in contraction burst or pulse

activity with stimuli (or mechanosensory response) in "footless" individuals was similar to the

increase in activity observed in whole individuals ("Footless" N = 8 animals stimulated 20 min,

Cohen's d = 1.89, Cliff's delta = 0.81, p < 0.01); Whole N = 8 animals stimulated 20 min,

Cohen's d = 1.80, Cliff's delta = 1.00, p < 0.01; Fig 3b,c). Furthermore, we found no significant

difference in either the spontaneous contraction probability or the mechanosensory response

probability of "footless" animals and whole animals ("Footless" N = 3 animals not stimulated,

spontaneous contraction probability = 0.13 ± 0.01, mean ± SEM; "Footless" N = 3 animals

stimulated for 60 min, mechanosensory response probability = 0.88 ± 0.05, mean ± SEM)

(Whole N = 3 animals not stimulated, spontaneous contraction probability = 0.15 ± 0.01, mean ±

SEM; Whole N = 3 animals stimulated for 60 min, mechanosensory response probability = 0.73

± 0.07, mean ± SEM; Figure 3 - Figure Supplement 3b,c, Video 4, 5,10, and 11).

Oral neurons play a major role in mechanosensory response in *Hydra*

While the mechanosensory response in *Hydra* remained unaffected with the removal of the

aboral nerve ring, we found that removal of the hypostome and tentacles (or "headless" *Hydra*)

resulted in significant changes in both the mechanosensory response and spontaneous

contraction bursts or pulses. When we measured the mechanosensory response in "headless"

Hydra, we found that the animals still responded to mechanical stimulation with a significant

increase in their contraction bursts or pulses, however, they did so with a lower magnitude

compared to whole and "footless" individuals ("Headless" N = 8 animals stimulated for 20 min, p

< 0.01, Cohen's d = 1.37, Cliff's delta = 1.00; Figure 3b, c). Specifically, the "headless" *Hydra*

responded with (> 2x) lower probability compared to whole animals, and they also showed a (>

3x) lower probability of spontaneous contraction bursts and pulses ("Headless" N = 3 animals

stimulated for 60 min, mechanosensory response probability = 0.29 ± 0.02, mean ± SEM;

"Headless" N = 3 animals not stimulated, spontaneous contraction probability = 0.05 ± 0.01,

mean ± SEM; Figure 3 - Figure Supplement 3b, c, Video 12 and 13).

To verify the reduction in mechanical response probability was specific to removing the oral

network of neurons, and not simply the result of injury, we performed experiments with animals

that we cut longitudinally along the body axis ("bisected" *Hydra*) to remove approximately the

same amount of tissue while preserving the neuronal subtypes in both the oral and aboral

networks. We found that these longitudinally "bisected" animals showed mechanosensory

responses that were not significantly different from that of the whole, "headless", or "footless"

animals ("Bisected" N = 3 animals stimulated for 60 min, mechanosensory response probability

= 0.54 ± 0.14, mean ± SEM; Figure 3 - Figure Supplement 3b, c). However, the magnitude of

increase in contraction bursts or pulses activity due to stimulation in "bisected" individuals,

though lower than whole and "footless" individuals, was larger than "headless" individuals

("Bisected" N = 8 animals stimulated for 20 min, $p < 0.001$, Cohen's d = 2.32, Cliff's delta = 1.0).

This suggests that our observations in "headless" *Hydra* indeed depended upon the types of

neurons removed during the headless resection and not simply an injury response. We also

found that these longitudinally "bisected" animals had a lower probability of spontaneous

contraction bursts or pulses than the whole animals but higher than "headless" animals. This

suggests that the entire network needs to be intact for normal contraction bursts or pulses

activity, and the loss of roughly half the network leads to some reduction in contractile activity

("Bisected" N = 3 animals not stimulated, spontaneous contraction probability = 0.08 ± 0.02,

mean ± SEM; Figure 3 - Figure Supplement 3b, c Video 14 and 15).

We next asked if the body column alone is sufficient to mediate the mechanosensory response.

To answer this question, we completely removed both oral and aboral regions. In "body column"

animals, we found significant reduction in both the mechanosensory response and spontaneous

contraction bursts and pulses relative to whole animals. The "body column" animals had a

mechanosensory response probability that was not different from the mechanosensory

response in "headless" animals, while the spontaneous contraction bursts and pulses probability

was lower compared to that of "headless" animals ("Body column" N = 3 animals stimulated for

60 min, mechanosensory response probability = 0.19 ± 0.08; "Body column" N = 3 animals not

stimulated, spontaneous contraction probability = 0.02 ± 0.01 mean ± SEM; Figure 3 - Figure

Supplement 3b,c). Although we found a significant increase in contraction bursts and pulses

with stimulation as compared to spontaneous contraction bursts and pulses activity in "body

column" individuals similar to all resections, the magnitude of the increase was the lowest in

"body column" animals (even lower than "headless" animals) ("Body column" N = 6 animals

stimulated for 20 min, p < 0.05, Cohen's d = 1.10, Cliff's delta = 0.50; Figure 3b, c, Video 16 and

17). Moreover, we did not observe significant increases in contraction bursts and pulses over

the same time period (comparing activity from 0 - 20 min with activity from 20 - 40 min, see

"Methods") in non-stimulated animals, suggesting the higher probability of contraction bursts

and pulses was in fact due to mechanical stimulation (Figure 3 – Figure Supplement 4). Thus,

based on the comparison between the probability of spontaneous contraction bursts and pulses

and mechanosensory response in all resections, we found that the "body column" animals had a

weak response to touch despite their slightly increased contraction bursts and pulses probability

with mechanical stimulation.

378
Oral and aboral networks show different patterns of activity during spontaneous

contractions compared to mechanically stimulated contractions

To identify how the activity of neurons in the oral and aboral networks coordinate spontaneous

and stimulated responses, we manually tracked the calcium activity of several neurons in the

oral and aboral regions (20 minutes no stimulation, 10 minutes stimulation 1 sec every 31 sec, n

= 3 *Hydra*; (Figure 4a, b, Figure 4 – Figure Supplement 1 and 2). We found there were at least

two independent networks of neurons based on a correlation analysis (Figure 4c). Specifically,

we time-aligned the calcium activity with either spontaneous contractions or mechanical

stimulation events to classify these groups of neurons based on their activity (Figure. 4d, e, g).

One group of correlated neurons found throughout the entire body showed averaged calcium

activity less than one second after a mechanical stimulus and spontaneous activity that is

consistent with previously reported Contraction Burst (CB) neurons[30]. We plot calcium dynamics

of these CB neurons as shades of blue in Figure. 4 (and Figure 4 - Figure Supplement 1 and 2).

These neurons show bursts of activity that are synchronized with muscle contractions and show

calcium activity that is highly correlated with the average peduncle ROI. In addition to these CB

neurons, we found other groups of correlated neurons with average calcium activity that is

independent of the CB network. One group showed a distinctive pattern of activity following

mechanical stimulation, but no distinctive activity associated with spontaneous contractions.

Specifically, this group of neurons near the oral end responded approximately 10 seconds after

mechanical stimulation (Figure. 4e, g). We found these putative "mechanically responsive (MR)

neurons" in all three Hydra we analyzed and plot their calcium dynamics as shades of red in

Figure. 4 (and Figure 4 - Figure Supplement 1 and 2). The fact that these MR neurons do not

show activity associated with spontaneous contractions clearly indicates that they are not a part

of the CB network, but rather these two distinct networks (CB and MR) are involved in the

Hydra's response to mechanical stimulation. We also note that these MR neurons do not fire

periodically as would be expected for the Rhythmic Potential (RP) network[30]. In addition to CB

and MR neurons, in Hydra 2 (Figure 4 - Figure Supplement 1) and 3 (Figure 4 - Figure

Supplement 2) we also found neurons that were not associated with either the MR network or

the CB network. These neurons we labeled as "unspecified groups" do not appear to be a part

of the CB or RP networks previously characterized nor the MR neurons we identify here.

These data suggest that there are at least two separate pathways involved in the

mechanosensory response. The first involves the CB neurons and muscle contractions. The

second network involving the MR neurons responds more slowly, with a latency of several

seconds. Because the activity of the MR neurons occurs after the contraction, their role in the

behavioral response remains unclear.

Discussion

Our experiments with "footless", "headless", "bisected", and "body column" animals show that

mechanosensory and spontaneous behaviors are regulated by neural ensembles that are

localized to select regions of the animal; however, some properties of the mechanosensory

response may be evenly distributed throughout the body. We demonstrate that localized touch

produces an increased calcium activity in both peduncle neurons (Figure 2) and ectodermal

epitheliomuscular cells (Figure 3), which is associated with body contractions (Figure 2 - Figure

Supplement 4)[31]. We identified at least two neuronal networks (MR and CB network) with

distinct neuronal activities mediating the stimulated responses (Figure 4), where the CB network

of neurons show fast calcium responses, and the MR neurons show slower calcium response. It

is possible that the MR neurons, primarily found in the oral end (N = 3 *Hydra*), may consist of

ec4 neurons, which are the only neuronal subtype in the oral end with an unknown function. The

other orally located ectodermal neuron populations, ec1B and ec3C, are suggested to belong to

the CB and RP1 circuits, respectively[48]. *Hydra's* responsiveness (i.e., probability of response to

a mechanical stimuli) depends on the stimulus intensity (Figure 2). Interestingly, we found that

the "headless" and "footless" animals can still respond to mechanical stimuli despite missing an

entire regional neuronal network; however, "headless" animals show reduced responsiveness

(Figure 3 and Figure 3 - Figure Supplement 3). The "body column" animals missing both

regional neuronal networks have the most dramatically reduced responsiveness, suggesting

that these two neuronal networks work together and play compensatory roles in mediating the

mechanosensory response.

Surprisingly, although the activity of the peduncle nerve ring is strongly associated with

spontaneous contractions, these neurons are not necessary for body contraction and response

to mechanical stimulation. This raises the question of what role the peduncle nerve ring plays in

Hydra behavior. One possible explanation is that this nerve ring coordinates body contractions

by enhancing the neural signal to ectodermal cells. An additional 4-second reduction in body

contraction duration in "body column" animals compared to headless animals supports the idea

that the peduncle nerve ring is also involved in coordination of contractile behavior (Figure 3 –

Figure Supplement 5). Furthermore, calcium activity propagates from the foot to the hypostome

in whole animals during body contractions[70], supporting a hypothesis that the peduncle network

of neurons may be motor neurons.

Based on these observations, we propose a simple model for sensorimotor information flow in

Hydra where we consider the hypostome as an integration point where sensory and motor

information converge. The information is then communicated to the peduncle where it is

amplified for coordinated whole body control. There may be sufficient functional redundancy

between the hypostome and peduncle regions such that removal or damage to one of them is

well tolerated in *Hydra*. Moreover, the diffuse network in the body column may retain minimal

processing capabilities needed for weak sensorimotor responses. The fast and slow calcium

responses from the CB and MR neurons, respectively, indeed support the hypothesis that *Hydra*

have multiple, separate pathways for behavioral response.

Overall, the quantitative characterization of *Hydra's* sensorimotor responses reported here help

to build the foundation for a more comprehensive investigation of information processing in

Hydra—an animal with clear advantages to supplement commonly studied model organisms in

neuroscience. Important next steps include developing mechanistic models to describe sensory

information processing that supports our results. Although our experiments describe general

information flow in *Hydra*, having a cellular level control of neuronal activity would be a clear

advantage for revealing the function of neuronal cell types as well as their functional

connectivity in the sensorimotor circuits. We expect additional future work with optogenetic

manipulation of specific neuronal subtypes combined with fast, volumetric and ratiometric

imaging techniques will provide a more comprehensive approach for characterizing this

sensorimotor processing in *Hydra*. Recognizing that calcium fluorescence imaging is limited in

its ability measure single spikes[71], we also expect other activity sensors (such as voltage

indicators) will reveal what role the MR neurons play in behavioral response. Building upon the

work reported here, one can then interrogate the roles of these regional networks with multiple

sensory modalities, such as light and heat, to answer questions about how diffuse nervous

systems may be capable of centralized information processing and multisensory integration.

These studies may help reveal a comprehensive model for how internal states and external

stimuli shape the behavioral repertoire in an organism with a highly dynamic neural architecture.

Materials and Methods

***Hydra* strains and maintenance**

Hydra were raised in *Hydra* medium at 18°C in a light-cycled (12 hr: 12 hr; light: dark) incubator

and fed with an excess of freshly hatched *Artemia nauplii* (Brine Shrimp Direct, Ogden, UT,

#BSEP8Z) three times a week (protocol adapted from Steele lab). All experiments were

performed at room temperature with animals starved for 2 days.

The transgenic line nGreen, kindly provided by Rob Steele, was generated by microinjecting

embryos with a plasmid containing the *Hydra* actin promoter driving GFP expression[48]. The

transgenic strains expressing GCaMP6s under the actin promoter in neurons and in ectodermal

epitheliomuscles (Addgene plasmid: #102558) were developed by microinjections of the

embryos by Christophe Dupre in the Yuste lab (Columbia University)[57]. The transgenic strain

expressing GCaMP7b in ectodermal epitheliomuscular cells was co-developed by Juliano Lab

(University of California, Davis) and Robinson Lab (Rice University). Briefly, the plasmid with

codon optimized GCaMP7b under the EF1a promoter was constructed by GenScript

(www.GenScript.com). Injections were performed as previously described[72] with the following

modifications: 1) injection solution was prepared by mixing 1 µL 0.5% phenol red (Sigma P0290-

100ML) with 6 µL plasmid DNA solution prior to centrifugation, and 2) embryos were fertilized

for 1-2 hours prior to injection. Plasmid promoters were cloned in expression vector pHyVec2

(Addgene plasmid: # 34790) using restriction sites NsiI. Plasmids were prepared by Maxiprep

(Qiagen, Valencia, CA) and eluted in RNase-free water. A plasmid DNA solution of 1.4 µg/uL

was injected into embryos using an Eppendorf FemtoJet 4x and Eppendorf InjectMan NI 2

microinjector (Eppendorf; Hamburg, Germany) under a Leica M165 C scope (Leica

Microscopes, Inc; Buffalo Grove, Il). Viable hatchlings with mosaic expression were propagated

by asexual reproduction, and asexual buds were screened and selected for increasing amounts

of transgenic tissue until a line was established with uniform expression in the ectodermal

epithelial cells.

Fluorescence imaging of *Hydra* nerve net

Distribution of neurons in the *Hydra* nerve net was fluorescently imaged with transgenic *Hydra*

vulgaris expressing GFP (nGreen) in neurons and neuronal progenitors (Figure 1a)[48]. *Hydra*

was anesthetized with 0.05% chloretone and immobilized in a ~160 μm tall microfluidic

chamber[31]. High-resolution fluorescence imaging was performed using a confocal microscope

(Nikon TI Eclipse) and 10x (0.45NA) objective, where the *Hydra* was imaged at a single plane

with multiple fields of views stitched together to obtain an image of the whole animal (Figure 1a).

Hydra resections

Hydra were placed in a petri dish and covered with enough *Hydra* medium to prevent

desiccation. When *Hydra* were relaxed and stationary, resections were performed with a single

incision with a scalpel across the body. We referenced published images of in-situ

hybridizations of different cell types and spatial expression patterns to guide the location of

incisions. For "footless" *Hydra,* an axial cut above the peduncle removed ~one third of the lower

body, which included the peduncle and the basal disk. For "headless" *Hydra,* an axial cut below

the hypostome removed ~one third of the upper body, including the tentacles and the

hypostome. For "bisected" *Hydra*, a transverse cut starting from the tip of the hypostome to the

basal disk was made along the midline of the body. For "body column" *Hydra*, an axial cut

above the peduncle removed the lower body followed by another axial cut below the hypostome

to remove the upper body region. This preparation resulted in an open tube body. *Hydra* were

stored in an 18°C incubator after the excisions and until beginning the experiments. *Hydra* can

seal open wounds within ~3-4 hours and repopulate the neuronal population to regain

functionality in 30-72 hours (Figure 3 – Figure Supplement 1, see "Imaging regeneration of

peduncle network"). To allow *Hydra* time to recover but not regain the functionality of lost

neuronal cell types, we performed experiments 6-12 hours post amputation.

Imaging regeneration of peduncle network

Imaging regeneration of peduncle network

Transgenic *Hydra* (GcaMP6s, neurons) were axially cut in the middle of the body column to

generate an oral and aboral halves (Figure 3 – Figure Supplement 1a). The oral half was

immobilized between two coverslips with a ~110 µm PDMS spacer. Calcium fluorescence was

conducted for 20 min every 2 hr on Nikon TI Eclipse inverted microscope with 20% excitation

light from Sola engine and GFP filter cube. We captured frames at ~10Hz (100ms exposures)

with Andor Zyla 4.2 sCMOS camera with NIS software. We used 4x (0.2 N.A) objective for wide-

field imaging to fit the entire *Hydra* in the field of view reduce the likelihood of *Hydra* migrating

out of the imaging frame. *Hydra* were exposed to blue excitation light for 20 min during imaging

and remained under dark conditions for 100 min between subsequent imaging timepoints. One

animal was imaged for ~20 hours starting with 1 hr post resection (Figure 3 – Figure

Supplement 1b). *Hydra* had a visibly open wound at the first imaging point but was not

detectable after 3 hr. There were no visibly active neurons in the regenerating aboral end,

indicating that the resected peduncle neurons had not regenerated. Another animal was imaged

for ~30 hours starting with 37 hr post resection (Figure 3 – Figure Supplement 1c). At the first

imaging time point (t = 37 hr post resection), there were few neurons in the peduncle region that

were active during body contractions, and these groups of neurons resembled a nerve ring as

early as 41 hr post resection. Although the peduncle nerve ring seemed to have formed at this

point, it appeared to not be as densely populated qualitatively as observed in whole animals.

Microfluidic device fabrication

The mechanical stimulation devices are double-layer microfluidic devices with push down valves

custom-designed with CAD software (L-edit) and fabricated using standard photo- and soft-

lithography techniques. All master molds were fabricated with transparency photomasks and

SU-8 2075 (MicroChem). The master mold for the bottom *Hydra* layer (circular chambers, 3 mm

diameter) was fabricated with the height of ~ 105-110 μm thick pattern (photoresist spun at

300rpm for 20 sec, 2100 rpm for 30 sec). The master mold for the top valve layer (9 individual

circular valves, 3 x 3 arrangement, 400 μm diameter each) was fabricated with height of ~110

μm thick (photoresist spun at 300rpm for 20sec, 2100 rpm for 30 sec). Polydimethylsiloxane

(PDMS) Sylgard 184 was used to cast microfluidic devices from the master molds. The bottom

Hydra layer (10:1 PDMS spun at 300rpm for 40 sec ~3 hr post mixing; cured for 12 min at 60°C)

was bonded to the valve layer (~4 mm thick, 10:1 PDMS; cured for ~40 min at 60°C with holes

punched for inlet ports) with oxygen plasma treatment (Harrick Plasma, 330 mTorr for 30 sec)

and baked for at least 10 min at 60°C. *Hydra* Insertion ports were hole-punched through the two

layers for *Hydra* layer with 1.5 mm biopsy punches, and the devices were permanently bonded

(O2 plasma treatment, 330 mTorr for 30 sec) to 500 μm thick fused silica wafer (University

Wafers) and baked for at least 1 hr at 60°C. The design files for the photomasks and step-by-

step fabrication protocols will be available on www.open*Hydra*.org (under Resource

hub/Microfluidics).

Hydra were immobilized and removed from the microfluidic device using syringes to apply

alternating positive and negative pressures as previously reported[31]. The microfluidic devices

were reused after cleaning similarly to the protocol previously reported. Briefly, the devices were

flushed with deionized water, sonicated (at least 10 min), boiled in deionized water (160°C for 1

hr), and oven dried overnight.

After repeated use, the PDMS stiffness can change and affect the valve deflection and the

actual force experienced by *Hydra* through the PDMS membrane. Additionally, uncontrollable

conditions during the device fabrication process can also lead to small differences between

devices. As a result, all data for Figure 2 was taken with a single device and the response curve

was used to calibrate (identify pressure that yielded ~ 60% response probability equivalent to

20-22 psi stimulus intensity) new devices.

Microfluidic mechanical stimulation

We used compressed air to inflate the microfluidic valves. For temporal control over valve

(on/off) dynamics, we used a USB-based controller for 24 solenoid pneumatic valves (Rafael

Gómez-Sjöberg, Microfluidics Lab, Lawrence Berkeley National Laboratory, Berkeley, CA

94720) and a custom-built Matlab GUI (available at www.open*Hydra*.org) that allowed setting

the stimulation parameters, such as the duration of valve 'on' (1 sec), duration of valve 'off' (30

sec), and the duration of stimulation (60 min, 119 cycles of valve 'on' and 'off'), and pre, post-

stimulation acclimation/control period (20 min). We used a pressure regulator to manually

control the air pressure into the valve manifold. In summary, we set the stimulation pressure

with a pressure gauge to regulate the flow of air into the valve manifold. This valve manifold was

controlled with a USB valve controller that allowed us to programmatically inflate the valve (turn

it 'on') with pressurized air with custom stimulation parameters.

To test sensory motor response to mechanical stimuli, we used pressurized air to inflate the

push-down valve and cause it to press down on the *Hydra* immobilized in the bottom layer.

Each experimental condition had at least three *Hydra* each. For a given condition, replication

experiments were conducted on different days. After an animal was immobilized inside the

Hydra chamber, we selected one valve (from the nine valves over the entire chamber) that was

directly above the mid-body column region to deliver stimuli. Although *Hydra* were free to move,

we did not observe large displacement most of the time and, as a result, the same valve

remained in contact with the animal throughout the stimulation period.

We adjusted the air pressure using a pressure regulator for each experiment, and the valves

were inflated using a USB valve controller (see above). The full-length stimulation experiment

consisted of 20 min of no stimulation (control/acclimation) followed by 60 min of stimulation

period (except habituation experiment where the stimulation period was 120 min), where valves

were pulsed with constant pressure (0 (control), 5, 10, 15, 20, 22 or 25 psi) for 1 sec every 31

sec, then another 20 min of no stimulation (control/acclimation). Shorter stimulation experiment

(used for whole-animal muscle imaging of various resections) consisted of 20 min of no

stimulation (acclimation/control) followed by 20 min of stimulation period (valves pulsed for 1

sec every 31 sec with a constant pressure of 0 or 22psi). We chose a 20 min initial control

period based on the high sensitivity to abrupt changes in light intensities (especially to blue

wavelengths used for excitation of GCaMP) in *Hydra* which leads to increased contractile

activity for 2-5 min. Even with the stimulus repeated for two hours at a constant inter-stimulus

interval, we found no obvious evidence of sensitization, habituation, or stimulus entrainment in

Hydra (Figure 2 – Figure Supplement 9). As a result, we chose not to randomize the inter-

stimulus interval.

Distribution of mechanical forces

We characterized the distribution of force exerted by a microfluidic valve by quantifying the

movements of neurons due to mechanical stimulation. We performed fluorescence imaging in

transgenic *Hydra* (nGreen) expressing GFP in neurons for ~8 min. We captured 8,000 frames at

~16Hz (50 msec exposures) with 4x objective (0.2 N.A.) and Andor Zyla 4.2 sCMOS camera

with 2x2 binning (1024x1024 frame size) using MicroManager. *Hydra* was stimulated in the

middle of the body column (valve pulsed for 1 sec every 31 sec, 5 times).

To quantify the movement of neurons and tissue throughout the *Hydra* body, we tracked a total

of 222 neurons that were visible for 1 min capturing the first two stimulation trials. We performed

semi-automated tracking with TrackMate plugin (ImageJ/(Fiji)[72–74]and manually corrected the

tracks where neurons were misidentified. From these tracks, we calculated the displacement of

each of the neurons between each frame (~50 msec). The average cellular displacement (0.4

µm per frame, 50 msec) was calculated by averaging the cellular displacements from all frames

when the valve was not pressurized. We found a significantly increased displacement (6.2 um, p

< 0.001) just after the valve was pressurized (and after the valve was depressurized 1 sec

later). We then generated a vector map of neuronal displacements by calculating the change in

position of each of the neurons in the frame just after the valve was pressurized for the first

stimulation trial (Figure 2 - Figure Supplement 1a). We also plotted the cellular displacement for

each of the neurons and the location of those neurons relative to the center of the valve. We

averaged the highest three displacements in 50 µm radial band increments from the valve

center to quantify how far the mechanical forces extended. This was a more conservative

measurement as the neurons in different tissue layers (endodermal and ectodermal layers

furthest from the valve) may have experienced different forces. By taking the average of the

highest three displacements in 50 µm radial bands increments from the valve center, we

identified the majority of the (shear) force was experienced by neurons bordering the valve in

250 µm radius.

Fluorescence intensity from GFP *Hydra*

To prove that the fluorescence intensity changes we observed with calcium imaging are due to

calcium activity and not motion artifacts from body contractions, we compared the average

fluorescence intensities from three different transgenic *Hydra* lines: 1) expressing GFP pan-

neuronally (nGreen line), 2) expressing GCaMP6s pan-neuronally, and 3) expressing GCaMP7b

in ectodermal epitheliomuscular cells (Figure 2 - Figure Supplement 3). For each *Hydra*, we

compared the average fluorescence intensities from three different regions of interest (ROIs)

that included the peduncle region, whole frame (for whole body) region, and valve region. We

also measured the body length by taking the major axis of an ellipse fitted along the oral-aboral

axis of the body column (from apex of the hypostome to the basal disc) after binarizing the

fluorescence image.

For this analysis, we used fluorescence imaging from transgenic *Hydra* expressing GFP pan-

neuronally (nGreen) performed in "Distribution of mechanical forces". *Hydra* was stimulated in

the middle of the body column (valve pulsed for 1 sec every 31 sec, 5 times). We used calcium

imaging from transgenic *Hydra* (expressing either GCaMP6s pan-neuronally or GCaMP7b in

ectodermal epitheliomuscular cells) stimulated in the middle of the body column (valve pulsed

for 1 sec every 31 sec, 120 times). The average fluorescence intensity and body length traces

were time-aligned to the onset of mechanical stimulation to show 1) changes in average

fluorescence intensity in peduncle ROI and the whole frame ROI are due to calcium activity

(increase in fluorescence from in GCaMP lines) and not motion artifact (no change in

fluorescence from GFP line) and 2) increase or decrease in body length or average

fluorescence from valve ROI are affected by stimulation artifacts.

Average calcium fluorescence from large and small ROIs in the peduncle

We performed fluorescence imaging in transgenic *Hydra* expressing GCaMP6s in neurons for

~1 min during spontaneous behaviors (*Hydra* was not stimulated). We captured frames at

~16Hz (50 msec exposures) with a 10x objective (0.45 N.A.) and Andor Zyla 4.2 sCMOS

camera with 3x3 binning (682x682 frame size) using MicroManager.

Because calcium fluorescence decreases when neurons are inactive, tracking multiple neurons

in a highly deformable region is a challenge. Nonetheless, we tracked 9 neurons from the

peduncle nerve ring that were visible (enough to track) for 1 min. We performed semi-

automated tracking with TrackMate plugin (ImageJ/(Fiji)[72–74] and manually corrected the tracks

where neurons were misidentified. We also used a large peduncle ROI, similar to how we used

an ROI for quantifying neural response to stimulation in all other experiments. We calculated the

average calcium fluorescence trace from small ROIs for individual neurons and large ROI for

the peduncle region. Fluorescence intensity was normalized by calculating $\Delta F/F$, where $\Delta F = F-$

F_0 and F_0 is the minimum fluorescence from prior time points. The large peduncle ROI had

highly correlated calcium fluorescence activity with smaller ROIs for individual neurons which

are known to belong to the contraction burst circuit (Figure 2 - Figure Supplement 2). As a

result, for all other experiments we use large peduncle ROI when measuring the neuronal

activity.

Whole-animal imaging of neural activity at different stimulus intensities

We characterized the neural response to repeated local mechanical stimuli at 0 (control), 5, 10,

15, 20 and 25 psi pressure with animals that expressed GCaMP6s in neurons (Figure 2). Each

pressure condition was experimented with three animals on different days using the same

device and stimulation protocol (valve pulsed for 1 sec every 31 sec for 60 min) to generate the

pressure-response curves (total of 18 animals). Calcium fluorescence imaging for all

experiments was conducted for the 100 min duration of the stimulation protocol (see Mechanical

Stimulation subsection) on Nikon TI Eclipse inverted microscope with 20% excitation light from

Sola engine and GFP filter cube. We captured 100,000 frames at ~16Hz (50 msec exposures)

with Andor Zyla 4.2 sCMOS camera with 4x4 binning using MicroManager. For imaging neural

activity, 12-bit low noise camera dynamic range was used. To synchronize the stimulation onset

times with imaging, we use used a data acquisition device (LabJack U3) to record the TTL

frame out signal (fire-any, pin #2) from the Zyla and the valve on/off timestamps from the valve

controller.

We used 4x (0.2 N.A.) objective for wide-field imaging to fit the entire *Hydra* in the field of view

to reduce the likelihood of *Hydra* migrating out of the imaging frame. There are neurons in the

oral region (hypostome) that are co-active with aboral region (peduncle) neurons during body

contractions; however, neurons in the hypostome appeared to be much sparser than those in

peduncle (Video 18). Discerning these hypostomal network neurons required higher

magnification which significantly reduced the field of view such that considerable amount of

nervous tissue could move in and out of the imaging plane (z-plane) or frame (xy-plane), making

it difficult to obtain reliable calcium fluorescence time-series.

Whole-animal imaging of neural activity with different stimulation regions

To map the sensitivity of different body parts to mechanical stimuli, we performed calcium

imaging of multiple animals (N = 3, whole animals expressing GCaMP6s in neurons) using

imaging settings previously described with modified experimental protocol. Based on the range

animal sizes (1-2.5 mm) and the size of the valve (400 um), we stimulated *Hydra* body at three

different regions along the oral-aboral axis: 1) the oral end (upper third of the body), 2) the

aboral end (lower third of the body), and 3) the third near the mid-body column. Each animal

was stimulated at three different locations (20 min no stimulation, stimulation region #1 - 1 sec

every 31 sec for 20 min, ~2 min no stimulation, stimulation region #2 - 1 sec every 31 sec for 20

min, ~2 min no stimulation, stimulation region #3 - 1 sec every 31 sec for 20 min using 22 psi).

We randomized the order in which the three different body regions were stimulated in each of

the three animals to avoid any stimulus entrainment artifacts. We then analyzed the peduncle

nerve ring activity in response to mechanical stimulation of different body regions as details in

"Analysis of calcium activity" and "Analysis of mechanosensory response".

Whole-animal imaging of neural activity in resected animals

For experiments that involved body lesions (Figure 3 – Figure Supplement 3), the same

experiment protocol and imaging settings as previously described (see "whole animal imaging of

neural activity at different stimulus intensities", 20 min no stimulation, stimulation 1 sec every 31

sec for 60 min, 20 min no stimulation) were used with either 0 psi (control) or 20 - 22 psi (~ 60%

response probability). Each condition group had three animals (total of 18 animals). The

"headless" and "bisected" animals were prepared with the appropriate body regions removed as

described previously (*"Hydra* resections"). Due to difficulty in tracking neurons and weak

GCaMP expression in body column neurons, we were unable to quantify responses from

"footless" and "body column" *Hydra* and thus excluded them from experiments.

Simultaneous electrophysiology and calcium imaging of ectodermal epitheliomuscular

cells

Electrical activity from the epitheliomuscular cells was measured simultaneously with calcium

imaging of the ectodermal epitheliomuscular cells in transgenic *Hydra* using a nano-SPEARs

device previously reported (Figure 2 – Figure Supplement 4)[31]. Briefly, transgenic *Hydra*

expressing GCaMP6s in the ectodermal epitheliomuscular cells starved for at least 48 hr were

used to measure the activity of the muscles (10 fps, 30 min, 4x objective with 15% light

intensity). An inverted microscope with GFP filter and Andor Zyla 4.2 were used for capturing

images. All electrical data was obtained with an Intan Technologies RHD2132 unipolar input

amplifier (http://intantech.com) at a sampling rate of 1KHz, low frequency cutoff and DSP filter of

0.1 Hz and high frequency cutoff of 7.5 KHz. From the calcium activity traces, we identified 30

sec intervals of either high amplitude activity or low amplitude activity to perform cross-

correlation analysis. The high and low amplitude activity regions were manually identified with a

threshold of 20% of the highest peak in the calcium activity. The high amplitude activity region

occurred during contraction bursts. The low amplitude activity region occurred during tentacle

contractions for muscular activity imaging. Both the Intan amplifier and the Zyla were triggered

with the same TTL signal. However, to account for any offset in the timing of the electrical and

optical data, we measured the maximum of the cross-correlogram in a 50 ms (approximately

one duty cycle of the trigger signal) window rather than the cross-correlation at zero offset to

generate the correlation maps. For correlation map, each frame was down sampled to 64 x 64

pixels, and the fluorescence trace for the downsampled pixels across the 30 sec intervals was

cross-correlated with electrical activity during the same 30 sec interval. The intensity of color in

the correlation map was used to indicate correlation values.

Whole-animal imaging of epitheliomuscular activity at different stimulus intensities

We characterized the muscle response to mechanical stimuli using animals that expressed

GCaMP7b under the EF1a promoter in ectodermal epitheliomuscular cells (Figure 3). We

measured contraction pulses and contraction bursts by averaging calcium activity in all the

epitheliomuscular cells (Figure 3a). The correlation between peduncle nerve ring activity and

muscle contractions has been previously established based on simultaneous electrophysiology

and neuronal[31] or ectodermal epitheliomuscular cells calcium imaging (Figure 2 – Figure

Supplement 4). The imaging protocol for epitheliomuscular cells was similar to the imaging of

neural activity (see "whole animal imaging of neural activity at different stimulus intensities"),

except a 16-bit camera dynamic range was used to avoid saturating the sensor.

We first developed a partial psychometric curve for calcium activity of epitheliomuscular cells to

confirm the dependence of response on stimulus intensity. Three animals were imaged for 20

min without stimulation and then stimulated at three different pressures (20 min at 15psi, 20 min

at 20 psi, 20 min at 25psi; or in reverse order), and the epitheliomuscular response curve was

used to identify the stimulus intensity (~ 22psi) which yielded ~ 60% response probability

(Figure 2 - Figure Supplement 5). This stimulation intensity was also selected for stimulation of

N = 8 whole animals (40 min of imaging; 20 min of no stimulation, 20 of stim for 1 sec every 31

sec with 22 psi) and animals with different body regions removed.

Whole-animal imaging of epitheliomuscular activity in resected animals

The various body regions were removed as described previously (*"Hydra* resections") to

prepare "headless", "footless", "bisected" and "body column" *Hydra*. Note, this transgenic line of

Hydra expressing GCaMP7b in the ectodermal cells showed some signs of deficit. They were

particularly sensitive to being handled and were more likely to dissociate after ~30 min in the

chambers during long-term microfluidic immobilization and fluorescence imaging. The resected

Hydra were especially difficult to image for the entire 100 min without any cell dissociation. This

could be due to the specific promoter used for driving GCaMP expression or that phototoxicity is

higher when there is a high expression of GCaMP in a larger number of cells. Because the

lesioned animals were more likely to be damaged during microfluidic immobilization, the

mechanical stimulation protocol (see Mechanical Stimulation subsection) was shortened to total

of 40 min of imaging with 20 min of no stimulation followed by 20 min of repeated stimulation (1

sec every 31 sec for 20 min). For three animals per each resection (total of 30 animals), we

used the full-length protocol (100 min of imaging: 20 min of no stimulation, 60 min of stimulation

every 31 sec, and 20 min of no stimulation) to obtain higher statistical power for quantifying

response probability (Figure 3 – Figure supplement 3). A total of 6-8 animals were stimulated for

each resection condition (whole, N = 6 not-stimulated, N = 8 stimulated; "footless", N = 4, not-

stimulated, N = 8 stimulated; "headless", N = 5 not-stimulated, N = 8 stimulated; "bisected", N =

5 not-stimulated, N = 8 stimulated; "body column" N = 3 stimulated, N = 6 stimulated; total 61

animals, Figure 3; 30 of which were experimented with long (100 min) stimulation protocol

Figure 3 – Figure supplement 3).

Analysis of calcium activity

To analyze neural response, we used the average fluorescence from the peduncle region.

Tracking individual neural responses was difficult due to high deformability of the body and lack

of fluorescence markers when neurons were not active; as a result, we looked at the

synchronous firing activity of the neurons in the peduncle region, which is known to have high

correlation with body contractions. To analyze epitheliomuscular response, we used average

fluorescence from the whole animal to obtain the fluorescence signal over time and analyzed

these signals similarly to neural response. Briefly, using ImageJ (Fiji)[72,73] we used a ROI over

the peduncle region or the whole *Hydra* to obtain fluorescence signal over time for neuronal or

epitheliomuscular calcium activity, respectively.

From the fluorescence signal, we detected the large calcium spikes as individual contractions

(contraction pulses or bursts) using MATLAB peak finding algorithm. We generated a raster plot

of calcium spiking activity time-aligned with the stimulus where each row represented one

stimulation trial with 15 sec before and after stimulation onset. These raster plots were used to

calculate the probability for spontaneous contraction and mechanosensory response.

For contractile behavior analysis, we then annotated the calcium spikes. Single calcium spikes

were labeled as single pulse contraction. A volley of calcium spikes was labeled as a

contraction burst. Both of these led to behavioral contractions, thus time between contractions

was calculated as the time between the end of a contraction event (single pulse or burst) and

the start of the next one (as shown in Figure 3 – Figure Supplement 5). Percent of contractions

that are single pulse contractions was calculated as the fraction behavioral contractions that

were single calcium spikes (not bursts). Contraction duration was used to indicate the amount of

time contraction behavior lasted (time from rise in fluorescence signal to return to baseline, as

shown in Figure 3 – Figure Supplement 5)

Analysis of mechanosensory response
859

To obtain the response probability (whether an animal had a contraction pulse—either a single

pulse or a pulse from a burst) at the time of stimulation (when valve was pressurized), we

generated a raster plot of fluorescence spikes time-aligned to stimulus onset and superimposed

for each 30 sec intervals (time between stimulus). We used an hour of activity (t = 20-80 min) to

generate the raster plot and calculate response probability. We defined the 1 sec window while

the valve was pressed as the response window for each trial (Figure 2 – Figure Supplement 6).

We then calculated the fraction of all trials (119 trials over 60 min or 40 trials over 20 min) per

animal that had at least one fluorescence spike ("contraction pulse") in the 1 sec response

window following stimulus onset to obtain the contraction probability. Extraction of raw

fluorescence for ROIs was performed with ImageJ (Fiji)[72,73] and post-processing analysis was

performed with MATLAB (using peak-finding algorithm to detect spikes). A one-way ANOVA

with Bonferroni correction was used for statistical analysis.

For animals with shorter experiment duration, the mechanosensory response probability in

stimulated animals was calculated over the 20 min segment (time = 20-40 min) with a fraction of

all trials (40 trials over 20 min) that had at least one fluorescence spike. A one-way ANOVA with

Bonferroni correction was used for statistical analysis when comparing multiple conditions. A

paired t-test was used to compare the difference between probability of spontaneous

contraction and mechanosensory probability in the same animals for each of the conditions. The

effect size (magnitude of difference) was measured with Cohen's d and Cliff's delta.

Analysis of spontaneous contraction
882

For non-stimulated animals, fluorescence activity from one hour of activity (time = 20-80 min)

was converted into a raster plot with multiple (119 trials over 60 min) 30 sec long intervals (to

match the stimulation interval in stimulated animals). We obtained "stimulation times" using a

DAQ to record the signal from valve controller except the air pressure was set to 0 psi. The

spontaneous contraction probability was calculated by taking the average of response

probability from a random 1 sec interval (Figure 3 and Figure 3 – Figure Supplement 3). Briefly,

by sliding a 1 sec window by ~0.3 sec over the x axis (30 sec of stimulation interval) on the

raster plot (pooled from all three animals for Figure 4a, per individual stimulated animal for

Figure 3 – Figure Supplement 3), we generated the distribution of the fraction of trials (119 trials

x 3 animals over 60 min) with at least one contraction event in a random 1 sec response window

(Figure 3 – Figure Supplement 3). The distributions were compared with a Kruskal-Wallis test.

For animals with shorter experiment duration, spontaneous contraction probability in stimulated

animals was calculated similar to non-stimulated animals above, except the raster plot was

generated with the first 20 min of fluorescence activity (time = 0-20 min) when no stimulation

was applied. Briefly, we generated a distribution of random probabilities by sliding a 1 sec

window over the x-axis on the raster plot and used the distribution mean as the spontaneous

contraction probability to compare with mechanosensory response probability from the same

animal with a paired t-test (Figure 3). For non-stimulated animals with shorter experiment

duration, spontaneous contraction probability was also calculated over the second 20 min

interval (t = 20-40 min) to confirm the increase in contraction probability during mechanical

stimulation was in fact due to stimuli and not just from temporal variation in spontaneous

contractions (Figure 3 – Figure Supplement 4). Note, these experiments were performed with

Hydra constrained to ~110um thick microfluidic chambers. Although animals are able to behave

under such confinements, the range of behavioral motifs and rates may be altered due to

compression.

Neuron subtype gene expression analysis with RT-qPCR
911

Resected *Hydra* were prepared as described above, with 12 polyps per biological replicate

(except 6 whole polyps for control) and a total of two biological replicates per treatment.

Approximately 12 hrs post resections, tissue was frozen in 1 mL Trizol at -80°C until use. RNA

was isolated using the Zymogen RNA Clean and Concentrator kit (Zymogen #R1017) with an

in-column Zymogen DNAse I digestion (Zymogen #E1010) following the manufacturer's

protocol. cDNA was synthesized using 1 μg of purified RNA and Promega M-MLV RNase H

Minus Point Mutant Reverse Transcriptase (Promega, Madison, WI M3682) using the

manufacturer's protocol for oligo dT-primed synthesis. cDNA synthesis was validated via PCR,

and all cDNA samples were diluted 1:10 in nuclease-free water for use in qPCR experiments.

Each sample was run in three technical replicates per gene in a 10 uL qPCR reaction using Bio-

Rad SsoAdvanced universal SYBR green master mix (Bio-Rad, Hercules, CA 1725271).

Samples were run on a CFX96 Touch Real-Time PCR Detection System (Bio-Rad 1855195).

Data were analyzed with the $2^{-\Delta\Delta Ct}$ method using the 'tidyverse' package in R[75,76].

Briefly, Cq values from technical replicates were pooled for subsequent analyses. *rp49* was

used as an internal control to calculate ΔCq values after first being found to give similar results

across all treatments when compared to a second control gene, *actin.* All results were

normalized to the "whole" animal samples. A template-free water control was performed for all

primer sets to ensure contamination-free reactions. All qPCR primers (Table 1) were validated

via a 10-fold serial dilution standard curve to have a binding efficiency over 90%.

Table 1: qPCR primers for neuronal subtypes

Subtype	transcriptID	F sequence	R sequence
en1	t29955aep	GCC GCA GTA TCA TCA TAC AAA TC	CCA TAA ACC ACA CAT CGC ATA AA
en2	t20666aep	CTT CTT GCT TCT ATC CTC GTT CT	TAC CTC AAG TAA ATT ATC GGT CTC G
en3	t33579aep	CGT TGG TAT GAC TAT AAT CGT TGT TAT G	AGG ATA CAT CAC CCA CCA AAT C
ec1A	t22316aep	GCC TTT CTT TAT CTC GGG TAT CT	ACC TCC CAT GAG TAG CTG TA
ec1B	t20807aep	AAG ATC TAC GAC GTC ATA TCA ATC A	TCA TGC CCT TAT TAC CCT CTT G
ec2	t7411aep	GCG CCT TGT AAC TAT GGT CTT A	TTC GTA GAA CAT TGT CAT CTT CCT
ec3A	t9620aep	CGG TGC TGC TCC TAA TTC AA	CTC CGG TGC ACT GAT TTA TAG G
ec3B	t7664aep	TGT TTC AAA TGC AGA CGA AGA TG	GCG TGT TTA TTT GCC TGG AC
ec3C	t19558aep	CGG TTA GAT ACA CTG CGG TTA G	TAC GTG CCG TTC TTC GTT T
ec4A	t33899aep	GGC TTT AAT CGT TGT AGC TCT TG	CTT GCT ATC TTC TGA CAA GTG ATT G
ec5	t6329aep	CCA ACA ATG GTC GAA TGA AGA AA	ATC GCC AGG TTT GTA TCC TTT A
RP49	t6797aep	GCC AAA CTG GAG AAA ACC TAA AG	TCA GGC ATA AGA TGA CGT GTC
Actin	t11116aep	CGC CCT CGT AGT TGA TAA TGG	AAT CCT TCT GTC CCA TAC CAA C

935
936
937
938   **Whole animal imaging with single neuron resolution**

939
940   We performed fluorescence imaging in transgenic *Hydra (N = 3)* expressing GCaMP6s in

941   neurons for ~30 min (30 min of imaging: 20 min of no stimulation, 10 min of stimulation every 31

942   sec). We captured frames at ~16Hz (50 msec exposures) with a 10x objective (0.45 N.A.) and

943   Andor Zyla 4.2 sCMOS camera with 3x3 binning (682x682 frame size) using MicroManager.

944
Because calcium fluorescence intensity is low (almost indiscernible from the autofluorescence

of *Hydra*) when neurons are inactive, tracking multiple neurons in a highly deformable region

without static nuclear fluorescence (such as RFP) is a challenge. Nonetheless, we manually

tracked 5-6 neurons throughout the animal body that were visible (enough baseline

fluorescence to track even when the neurons were inactive) for the entire duration of the

imaging. We performed semi-automated tracking with TrackMate plugin (ImageJ/(Fiji)[72–74] to

track individual neurons with small circular ROIs (~5 μm radius). We manually tracked

(interpolating the current ROI location based on the past and future locations) the neurons when

the fluorescence levels were dimmer than the background autofluorescence or when neurons

were misidentified by TrackMate. We also used a large peduncle ROI, similar to how we used

an ROI for quantifying neural response to stimulation in all other experiments. We calculated the

average calcium fluorescence traces from small ROIs for individual neurons and large ROI for

the peduncle region. Each of the fluorescence traces were corrected by calculating $\Delta F/F_0$,

where F_0 was the mean fluorescence intensity of the trace.

We performed the correlation analysis (MATLAB) of the calcium fluorescence time-series for the

individual neuron ROIs and peduncle ROI to identify groups of neurons with correlated activity.

We calculated the correlation coefficients for the individual neurons and the peduncle ROI by

correlating the entire 30 min calcium fluorescence trace for each of the neurons. These

correlation coefficients were shown to be statistically significant then randomly reshuffled

calcium fluorescence. We divided each of the neuronal calcium time-series into 30 blocks.

These individual blocks were randomly recombined to generate reshuffled time-series for each

of the neurons to calculate correlation coefficients. This random reshuffling of each neuron was

repeated 1000 times to obtain the mean correlation coefficients for each of the neuron pairs

(Figure 4 – Figure Supplement 3d-f). We then calculated the z-score $(x-\mu)/\sigma$, where x is the

correlation coefficients of the original raw time-series, μ is the mean correlation coefficients of

the time-series randomly reshuffled 1000 times and σ is the standard deviation of the

correlation coefficients of the time-series randomly reshuffled (Figure 4 – Figure Supplement 3g-

i). To identify the neuronal clusters, we used a hierarchical clustering algorithm (MATLAB

function 'linkage' with Euclidian distance of correlation coefficients) on the correlation

coefficients. We then used leaf order from the linkage tree to sort the neurons and generate the

correlation heatmap.

We calculated the average fluorescence traces from the time-aligned calcium fluorescence to

either spontaneous contractions or stimulus events to identify the roles of these neurons. For

spontaneous contractions, the calcium fluorescence (t = 0-20 min) was time-aligned to the onset

of contraction bursts or pulses from the peduncle ROI. For stimulated contractions, the calcium

fluorescence (t = 20-30 min) was time-aligned to the onset of stimulus recorded with a DAQ

(see "Whole-animal imaging of neural activity at different stimulus intensities"). We further

confirmed the different neuronal clusters identified by performing correlation analysis on all

neurons from all three Hydra by calculating the correlation coefficients for the average

fluorescence traces from the time-aligned calcium fluorescence to stimulus events (Figure 4f.g).

References

989   1.   Clark, D. A., Freifeld, L. & Clandinin, T. R. Mapping and Cracking Sensorimotor Circuits
990   in Genetic Model Organisms. *Neuron* (2013). doi:10.1016/j.neuron.2013.05.006

991   2.   Ghosh, D. D., Nitabach, M. N., Zhang, Y. & Harris, G. Multisensory integration in C.
992   elegans. *Current Opinion in Neurobiology* (2017). doi:10.1016/j.conb.2017.01.005

993   3.   Kaplan, H. S., Nichols, A. L. A. & Zimmer, M. Sensorimotor integration in Caenorhabditis
994   elegans: A reappraisal towards dynamic and distributed computations. *Philos. Trans. R.*
995   *Soc. B Biol. Sci.* (2018). doi:10.1098/rstb.2017.0371

996   4.   Ahrens, M. B. *et al.* Brain-wide neuronal dynamics during motor adaptation in zebrafish.
997   *Nature* (2012). doi:10.1038/nature11057

998   5.   Chen, X. *et al.* Brain-wide Organization of Neuronal Activity and Convergent
999   Sensorimotor Transformations in Larval Zebrafish. *Neuron* (2018).
1000   doi:10.1016/j.neuron.2018.09.042

1001   6.   Haesemeyer, M., Robson, D. N., Li, J. M., Schier, A. F. & Engert, F. A Brain-wide Circuit
1002   Model of Heat-Evoked Swimming Behavior in Larval Zebrafish. *Neuron* (2018).
1003   doi:10.1016/j.neuron.2018.04.013

1004   7.   Hill, A. J., Mansfield, R., Lopez, J. M. N. G., Raizen, D. M. & Buskirk, C. Van. Cellular
1005   Stress Induces a Protective Sleep-like State in C. elegans. *Curr. Biol.* 1–7 (2014).
1006   doi:10.1016/j.cub.2014.08.040

1007   8.   Raizen, D. M. *et al.* Lethargus is a Caenorhabditis elegans sleep-like state. *Nature* **451**,
1008   569–72 (2008).

1009   9.   Artiushin, G. & Sehgal, A. The Drosophila circuitry of sleep–wake regulation. *Curr. Opin.*
1010   *Neurobiol.* **44**, 243–250 (2017).

1011   10.   Guo, F., Holla, M., Díaz, M. M. & Rosbash, M. A Circadian Output Circuit Controls Sleep-
1012   Wake Arousal in Drosophila. *Neuron* **100**, 624-635.e4 (2018).

11. Kayser, M. S. & Biron, D. Sleep and development in genetically tractable model

organisms. *Genetics* (2016). doi:10.1534/genetics.116.189589

12. Gandhi, A. V., Mosser, E. A., Oikonomou, G. & Prober, D. A. Melatonin Is required for the

circadian regulation of sleep. *Neuron* (2015). doi:10.1016/j.neuron.2015.02.016

13. Zhdanova, I. V., Wang, S. Y., Leclair, O. U. & Danilova, N. P. Melatonin promotes sleep-

like state in zebrafish. *Brain Res.* (2001). doi:10.1016/S0006-8993(01)02444-1

14. Zimmerman, J. E., Naidoo, N., Raizen, D. M. & Pack, A. I. Conservation of sleep: insights

from non-mammalian model systems. *Trends in Neurosciences* (2008).

doi:10.1016/j.tins.2008.05.001

15. Kanaya, H. J. *et al.* A sleep-like state in *Hydra* unravels conserved sleep mechanisms

during the evolutionary development of the central nervous system. *Sci. Adv.* **6**,

eabb9415 (2020).

16. Broussard, G. J., Liang, R. & Tian, L. Monitoring activity in neural circuits with genetically

encoded indicators. *Front. Mol. Neurosci.* **7**, 97 (2014).

17. Chen, T.-W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*

499, 295–300 (2013).

18. Lemon, W. C. *et al.* Whole-central nervous system functional imaging in larval

Drosophila. *Nat. Commun.* **6**, 7924 (2015).

19. Prevedel, R. *et al.* Simultaneous whole- animal 3D imaging of neuronal activity using

light-field microscopy. *Nat. Methods* **11**, 727–730 (2014).

20. Ahrens, M. B., Orger, M. B., Robson, D. N., Li, J. M. & Keller, P. J. Whole-brain functional

imaging at cellular resolution using light-sheet microscopy. *Nat Methods* **10**, 413–420

(2013).

21. Cong, L. *et al.* Rapid whole brain imaging of neural activity in freely behaving larval

zebrafish (Danio rerio). *Elife* **6**, e28158 (2017).

22. Kim, D. H. *et al.* Pan-neuronal calcium imaging with cellular resolution in freely swimming

zebrafish. *Nat. Methods* **14**, 1107–1114 (2017).

23. Portugues, R., Feierstein, C. E., Engert, F. & Orger, M. B. Whole-brain activity maps

reveal stereotyped, distributed networks for visuomotor behavior. *Neuron* **81**, 1328–1343

(2014).

24. Vladimirov, N. *et al.* Light-sheet functional imaging in fictively behaving zebrafish. *Nat.*

Methods **11**, 883 (2014).

25. Gonzales, D. L., Badhiwala, K. N., Avants, B. W. & Robinson, J. T. Bioelectronics for

Millimeter-Sized Model Organisms. *iScience* **23**, 100917 (2020).

26. Nguyen, J. P. *et al.* Whole-brain calcium imaging with cellular resolution in freely

behaving Caenorhabditis elegans. *Proc. Natl. Acad. Sci.* **113**, E1074–E1081 (2016).

27. Nichols, A. L. A., Eichler, T., Latham, R. & Zimmer, M. A global brain state underlies C.

elegans sleep behavior. *Science (80-.).* **356**, eaam6851 (2017).

28. Gonzales, D. L., Zhou, J., Fan, B. & Robinson, J. T. A microfluidic-induced C. elegans

sleep state. *Nat. Commun.* **10**, 1–13 (2019).

29. Aimon, S. *et al. Fast near-whole-brain imaging in adult drosophila during responses to*

stimuli and behavior. PLoS Biology **17**, (2019).

30. Dupre, C. & Yuste, R. Non-overlapping Neural Networks in Hydra vulgaris.

Curr. Biol. **27**, 1085–1097 (2017).

31. Badhiwala, K. N., Gonzales, D. L., Vercosa, D. G., Avants, B. W. & Robinson, J. T.

Microfluidics for electrophysiology, imaging, and behavioral analysis of: Hydra. *Lab Chip*

18, 2523–2539 (2018).

32. Bargmann, C. I. & Horvitz, H. R. Chemosensory neurons with overlapping functions direct

chemotaxis to multiple chemicals in C. elegans. *Neuron* **7**, 729–742 (1991).

33. Bargmann, C. I. & Avery, L. Laser Killing of Cells in Caenorhabditis elegans. *Methods*

Cell Biol. **48**, 225–250 (1995).

34. El Bejjani, R. & Hammarlund, M. Neural regeneration in Caenorhabditis elegans. *Annu.*

Rev. Genet. **46**, 499–513 (2012).

35.   Kroehne, V., Freudenreich, D., Hans, S., Kaslin, J. & Brand, M. Regeneration of the adult

1067   zebrafish brain from neurogenic radial glia-type progenitors. *Development* **138**, 4831–

1068   4841 (2011).

1069   36.   Hecker, A., Schulze, W., Oster, J., Richter, D. O. & Schuster, S. Removing a single

1070   neuron in a vertebrate brain forever abolishes an essential behavior. *Proc. Natl. Acad.*

1071   *Sci. U. S. A.* **117**, 3254–3260 (2020).

1072   37.   Itayama, T. & Sawada, Y. Development of electrical activity in regenerating aggregates of

1073   hydra cells. *J. Exp. Zool.* **273**, 519–26 (1995).

1074   38.   Soriano, J., Rüdiger, S., Pullarkat, P. & Ott, A. Mechanogenetic coupling of Hydra

1075   symmetry breaking and driven turing instability model. *Biophys. J.* **96**, 1649–1660 (2009).

1076   39.   Gierer, A. *et al.* Regeneration of Hydra from Reaggregated Cells. *Nat. New Biol.* **239**, 98–

1077   101 (1972).

1078   40.   Burnett, A. L. & Diehl, N. A. The nervous system of hydra. I. Types, distribution and origin

1079   of nerve elements. *J. Exp. Zool.* **157**, 217–226 (1964).

1080   41.   Lfntz, T. L. & Barrnett, R. J. Fine structure of the nervous system. *Am. Zool.* (1965).

1081   42.   Bode, H. R., Heimfeld, S., Koizumi, O., LIttlefield, C. L. & Yaross, M. S. Maintenance and

1082   regeneration of the nerve net in hydra. *Integr. Comp. Biol.* **28**, 1053–1063 (1988).

1083   43.   Campbell, R. D. Tissue dynamics of steady state growth in hydra littoralis. III. Behavior of

1084   specific cell types during tissue movements. *J. Exp. Zool.* **164**, 379–391 (1967).

1085   44.   Bode, H. *et al.* Quantitative analysis of cell types during growth and morphogenesis in

1086   Hydra. *Wilhelm Roux Arch. f r Entwicklungsmechanik der Org.* **171**, 269–285 (1973).

1087   45.   Epp, L. & Tardent, P. The distribution of nerve cells inHydra attenuata Pall. *Wilhelm*

1088   *Roux's Arch. Dev. Biol.* **185**, 185–193 (1978).

1089   46.   Koizumi, O. *et al.* Nerve ring of the hypostome in Hydra. I. Its structure, development, and

1090   maintenance. *J. Comp. Neurol.* **326**, 7–21 (1992).

47.   Hufnagel, L. A. & Kass-Simon, G. The two nerve rings of the hypostomal nervous system

of Hydra vulgaris---an immunohistochemical analysis. *Cell Tissue Res.* **366**, 1–15 (2016).

48.   Siebert, S. *et al.* Stem cell differentiation trajectories in Hydra resolved at single-cell

resolution. *Science (80-.).* **365**, eaav9314 (2019).

49.   Mast, S. O. REACTIONS TO TEMPERATURE CHANGES IN SPIRILLUM, HYDRA, AND

FRESH-WATER PLANARIANS. *Am. J. Physiol. Content* **10**, 165–190 (1903).

50.   Wagner, G. Memoirs: On Some Movements and Reactions of Hydra. *J. Cell Sci.* **s2-48**,

585–622 (1905).

51.   Rushforth, N. B. Inhibition of contraction responses of hydra. *Integr. Comp. Biol.* **5**, 505–

513 (1965).

52.   Rushforth, N. B. ., Burnett, A. L. . & Maynard, R. Behavior in Hydra : Contraction

Responses of Hydra pirardi to Mechanical and Light Stimuli. *Am. Assoc. Adv. Sci.* **139**,

760–761 (1963).

53.   Rushforth, N. B. & Burke, D. S. Behavioral and electrophysiological studies of Hydra. II.

Pacemaker activity of isolated tentacles. **140**, 502–519 (1971).

54.   CAMPBELL, R. D., JOSEPHSON, R. K., SCHWAB, W. E. & RUSHFORTH, N. B.

Excitability of nerve-free hydra. *Nature* **262**, 388–390 (1976).

55.   Takaku, Y. *et al.* Innexin gap junctions in nerve cells coordinate spontaneous contractile

behavior in Hydra polyps. *Sci. Rep.* **4**, 1–8 (2014).

56.   Chalfie, M. *et al.* The neural circuit for touch sensitivity in Caenorhabditis elegans. *J.*

Neurosci. **5**, 956–64 (1985).

57.   Wicks, S. R., Roehrig, C. J. & Rankin, C. H. A dynamic network simulation of the

nematode tap withdrawal circuit: Predictions concerning synaptic function using

behavioral criteria. *J. Neurosci.* **16**, 4017–4031 (1996).

58.   Vaidya, A. R., Pujara, M. S., Petrides, M., Murray, E. A. & Fellows, L. K. Lesion Studies in

Contemporary Neuroscience. *Trends Cogn. Sci.* **23**, 653–671 (2019).

59.   Krug, K., Daniel Salzman, C. & Waddell, S. Understanding the brain by controlling neural

1118   activity. *Philos. Trans. R. Soc. B Biol. Sci.* **370**, 1–4 (2015).

1119   60.   PASSANO, L. M. & McCULLOUGH, C. B. Co-ordinating systems and behaviour in hydra.

1120   I. Pacemaker system of the periodic contractions. *J. Exp. Biol. 41 643-664, 1964* 643–

1121   664 (1964).

1122   61.   Pierobon, P. Regional modulation of the response to glutathione in Hydra vulgaris. *J.*

1123   *Exp. Biol.* **218**, 2226–32 (2015).

1124   62.   Cazet, J. & Juliano, C. Oral Regeneration Is the Default Pathway Triggered by Injury in

1125   Hydra. *bioRxiv* 2020.07.06.189811 (2020). doi:10.1101/2020.07.06.189811

1126   63.   Tursch, A., Bartsch, N. & Holstein, T. W. MAPK signaling links the injury response to

1127   Wnt-regulated patterning in Hydra regeneration. *bioRxiv*

1128   2020.07.06.189795 (2020). doi:10.1101/2020.07.06.189795

1129   64.   Bode, H. R. Head regeneration in Hydra. *Dev. Dyn.* **226**, 225–236 (2003).

1130   65.   Goel, T., Wang, R., Martin, S., Lanphear, E. & Collins, E.-M. S. Linalool acts as a fast and

1131   reversible anesthetic in Hydra. *PLoS One* **14**, e0224221 (2019).

1132   66.   Yum, S. *et al.* A Novel Neuropeptide, Hym-176, Induces Contraction of the Ectodermal

1133   Muscle in Hydra. *Biochem. Biophys. Res. Commun.* **248**, 584–590 (1998).

1134   67.   Noro, Y. *et al.* Regionalized nervous system in Hydra and the mechanism of its

1135   development. *Gene Expr. Patterns* (2019). doi:10.1016/J.GEP.2019.01.003

1136   68.   Klimovich, A. *et al.* Prototypical pacemaker neurons interact with the resident microbiota.

1137   *Proc. Natl. Acad. Sci. U. S. A.* **117**, 17854–17863 (2020).

1138   69.   Shimizu, H. & Fujisawa, T. Peduncle of Hydra and the heart of higher organisms share a

1139   common ancestral origin. *Genesis* **36**, 182–186 (2003).

1140   70.   Szymanski, J. R. & Yuste, R. Mapping the Whole-Body Muscle Activity of Hydra vulgaris.

1141   *Curr. Biol.* **29**, 1807-1817.e3 (2019).

71.   Huang, L. *et al.* Relationship between simultaneously recorded spiking activity and

fluorescence signal in GCaMP6 transgenic mice. *Elife* **10**, e51675 (2021).

72. Rueden, C. T. *et al.* ImageJ2: ImageJ for the next generation of scientific image data.

BMC Bioinformatics **18**, 529 (2017).

73. Schindelin, J. *et al.* Fiji: An open-source platform for biological-image analysis. *Nat.*

Methods **9**, 676–682 (2012).

74. Tinevez, J. Y. *et al.* TrackMate: An open and extensible platform for single-particle

tracking. *Methods* **115**, 80–90 (2017).

75. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time

quantitative PCR and the 2-ΔΔCT method. *Methods* **25**, 402–408 (2001).

76. Core R Team. R: A Language and Environment for Statistical Computing. *R Foundation*

for Statistical Computing **2**, https://www.R--project.org (2020).

1154

1155

1156 **Acknowledgements**

1157

We thank Dr. Rob Steele (UC Irvine), Dr. Christophe Dupre (Harvard University), and Dr. Rafael

Yuste (Columbia University) for sharing transgenic *Hydra*. We also thank Dr. Guillaume Duret

and Dr. Caleb Kemere for useful discussions. The promoter sequence for EF1a for generating

transgenic *Hydra* was provided by Jack Cazet (UC Davis, Juliano lab), and the plasmid with

GCAMP7b was cloned (and codon optimized) by GenScript. The open source USB valve

controller was built based on the design by Rafael Gómez-Sjöberg, Microfluidics Lab, Lawrence

Berkeley National Laboratory, Berkeley, CA 94720. This work is funded by NSF EDGE. K.N.B.

is funded by a training fellowship from the Keck Center of the Gulf Coast Consortia on the NSF

IGERT: Neuroengineering from Cells to Systems 1250104. We also thank the Rice Shared

Equipment Authority and nanofabrication cleanroom facility where the devices were fabricated.

1169 **Competing interests**

1170   The authors declare no competing interests.

1171

Figures

 Figure 1: Distribution of neurons in the *Hydra* nerve net. a) Fluorescent image of *Hydra* nervous system. Green fluorescent

 protein (GFP) is expressed in neurons and neuronal progenitors (nGreen transgenic line[48]). Body anatomy is annotated on the left.

 White arrows indicate the body parts: hypostome, tentacles, peduncle, and basal disk. High neuronal density regions are annotated

 on the right. b) Distribution of neuronal cell types varying longitudinally along the body, with endodermal nerve net cell types in tan

 and ectodermal nerve net cell types in green. Cell types were identified through single cell RNA sequencing[48].

Pressure (psi)	Response probability mean +/- SEM	Time between contractions		Single pulse contractions mean +/- SEM
		mean +/- SEM	median +/- SE	
0	0.11 +/- 0.01	72.98 +/- 4.58 sec	66.68 +/- 5.73 sec	16.87 +/- 5.38 %
5	0.09 +/- 0.02	92.66 +/- 5.87 sec	86.65 +/- 7.35 sec	9.52 +/- 2.27 %
10	0.23 +/- 0.01	63.01 +/- 3.73 sec	54.73 +/- 4.67 sec	29.96 +/- 10.29 %
15	0.24 +/- 0.03	83.60 +/- 6.37 sec	64.73 +/- 7.98 sec	17.30 +/- 6.59 %
20	0.60 +/- 0.06	38.28 +/- 1.77 sec	31.14 +/- 2.21 sec	56.84 +/- 2.48 %
25	0.77 +/- 0.08	32.20 +/- 1.04 sec	31.14 +/- 1.30 sec	62.50 +/- 5.23 %

1181

Figure 2: *Hydra's* neuronal response depends on the mechanical stimulus intensity. a) (left) Side view of double-layer

microfluidic device for mechanical stimulation. (right) Device with pressurized valve. *Hydra* is immobilized in the bottom *Hydra* layer,

and pressurized air supplied into the valve layer causes the circular membrane (400 μm diameter) to push down on *Hydra*. b) (left)

Brightfield image of *Hydra* immobilized in the bottom layer of the chip and the arrangement of micro-valves on the top layer. Micro-

valve used for stimulation falsely colored with a light blue circle. (right) Fluorescent image of *Hydra* with pan-neuronal expression of

GCaMP6s. White dashed circle marks the peduncle region of interest (ROI) used for quantifying calcium fluorescence changes. c-d)

Representative calcium fluorescence activity in the peduncle region from an animal not stimulated and animal mechanically

stimulated with 25 psi. Black and green dots indicate fluorescence (calcium) spikes. Gray shaded regions indicate stimulus 'on' time

(also "response window"). c) Fluorescence (calcium) trace from *Hydra* not stimulated (top) and stimulated with 25 psi (middle).

Stimulation protocol in gray (bottom trace): 20 min no stimulation, 1 hr of repeated stimulation (1 sec 'on', 30 sec 'off') and 20 min no

stimulation. Stimulus 'on' times indicated with vertical lines. Magnification of 30 sec fluorescence and stimulation protocol trace from

one stimulation trial. d) Raster plot of stimulus time aligned spiking activity from multiple trials superimposed for *Hydra* not stimulated

(top) and stimulated with 25 psi (bottom). e). Mechanosensory response probability, fraction of trials (out of 119 total) that have at

least one calcium spike (also contraction pulse) occurring during the 1 sec response window (gray shaded region) when valve is

pressurized. Large circles indicate average probability from all animals (N=3) combined for each condition. Small circles indicate

probability from a single animal. Significant pairwise comparisons are shown with brackets (one-way ANOVA with post-hoc

Bonferroni correction). f) Time interval between body contractions under each condition. Dashed line represents the time interval

between stimuli (~31 sec). Brackets indicate significant differences in a Kruskal-Wallis test with post-hoc Dunn-Sidak correction. g)

Percent of all body contractions that are a single pulse; brackets show significant pairwise comparisons from a one-way ANOVA

with post-hoc Bonferroni correction. Error bars are standard error of mean (SEM); N = 3 *Hydra* for each condition; * = $p < 0.05$, ** =

$p < 0.01$, ***** = $p < 0.00001$. h) Table summarizing the mechanosensory response probability, time between contractions, and

percent of contractions that are single pulses for each stimulus intensity (mean ± SEM or median ± SE). Source data for the

quantitative characterization of mechanosensory response are available in the Figure 2 – source data 1.

1205

a

Whole *Hydra*   Footless *Hydra*   Headless *Hydra*   Bisected *Hydra*   Body column *Hydra*

1 mm   1 mm   1 mm   1mm   1 mm

b

n = 3 *Hydra*

Spontaneous activity

Mechanosensory response

Time (sec)   Time (sec)   Time (sec)   Time (sec)   Time (sec)

c

n = 8
d = 1.00
d = 1.80
**

n = 8
d = 0.81
d = 1.89
**

n = 8
d = 1.00
d = 1.37
**

n = 8
d = 1.00
d = 2.32
***

n = 6
d = 0.50
d = 1.10
*

Response probability

● Spontaneous activity (0 - 20 min)   ● Mechanosensory response (20 - 40 min)

1206

Figure 3: Oral region is important for mechanosensory response a) Representative images of resection preparations (6-12

hours post-resections) of transgenic *Hydra* during contraction (GCaMP7b, ectodermal epitheliomuscular cells): whole (or control),

footless, headless, bisected and body column animal. Entire frame ROI was used for analysis of the whole-body epithelial calcium

activity. b) Representative raster plot of stimulus time aligned calcium spikes from three animals with multiple trials superimposed.

Each black dot is a peak in calcium fluorescence identified as a contraction pulse. c) Response probability, fraction of trials that

have at least one calcium spike (also contraction pulse) occurring within 1 sec of stimulation onset. Gray dots are the mean

contraction probability during no stimulation (t = 0-20 min) calculated from 1 sec window shifted by ~0.3 sec over 30 sec intervals.

Blue dots are contraction probability during stimulation (t = 20-40 min) calculated from 1 sec response window during valve 'on'.

Light gray lines connect the probabilities for spontaneous contraction and mechanosensory response for each individual. Cartoon

schematics of *Hydra* indicate the resections performed. Excised body regions are outlined with a dashed line and unfilled area.

Color filled body regions indicate the portion of *Hydra* retained for the experiment. P-values from a paired t-test indicated as follows:

n.s. = not significant; * = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$. Source data for the mechanosensory response in resected animals

are available in the Figure 3 – source data 1.

1220

1221

Figure 4: Distinct networks of neurons involved in spontaneous and stimulated behaviors. a) Fluorescent image of

transgenic *Hydra* expressing GCaMP6s pan-neuronally. Individually tracked neurons ROIs are indicated by arrows, and peduncle

ROI is outlined with a white dashed circle. b) Calcium fluorescence traces from single neurons (top 6 traces) and average calcium

fluorescence from peduncle ROI (bottom trace). Mechanically responsive (MR) neurons are shown in shades of red. Contraction

burst (CB) neurons are shown in shades of blue. c) Heat map shows the correlation coefficients of individually tracked neurons and peduncle ROI from *Hydra* 1, with color bar at the bottom. Correlation was computed using the entire 30 min calcium fluorescence from single neurons shown in (b). d) Average calcium fluorescence traces from each of the neurons and peduncle ROI during spontaneous behaviors time-aligned with the onset of spontaneous body contractions. Dashed line indicates the onset of body contraction. e) Average calcium fluorescence traces from each of the neurons and peduncle ROI during stimulated behaviors time-aligned with the onset of mechanical stimulation. Gray shaded rectangle indicates mechanical stimulation. f) Heat map shows the correlation coefficients of g) the average calcium fluorescence traces from each of the neurons and peduncle ROI (traces in black) from three different *Hydra* during stimulated behaviors time-aligned with the onset of mechanical stimulation. Three groups of neurons were identified across the three *Hydra.* CB neurons (traces in blue), MR neurons (traces in red) and unspecified neurons (traces in yellow). a-e) Calcium fluorescence traces and correlation analysis (entire 30 min of activity, 20 min of spontaneous activity and 10 min of stimulated activity) from one representative *Hydra*. f-g) Correlation analysis of the average stimulus-aligned calcium fluorescence (30 sec stimulation interval average from 10 min of stimulated activity) of each neuron pooled multiple *Hydra* (N = 3 *Hydra*) to highlight the three groups of neurons identified: CB, MR and unspecified. Source data for fluorescent calcium activity from single neurons are available in the Figure 4 – source data 1.

Figure Supplements

Figure 2 – Figure Supplement 1: Distribution of mechanical forces. Fluorescent image (grayscale) of transgenic *Hydra* (nGreen) expressing GFP in neurons and neural progenitors. The Oral region is on the left. Aboral//peduncle region is on the right. Arrows (cyan) overlayed indicate the displacement of individual cells immediately before and after the valve is pressurized for stimulation (~50 msec). The lengths of the arrows are enlarged to be visible with the scaled size of the largest displacement ~31 µm as indicated by the legend (lower left). Semi-transparent (yellow) circle indicates the location of the valve (400 µm diameter). b) Displacement of neurons due to stimulation varies with the location of the neurons from the valve center. Each semi-transparent blue dot represents a single neuron (n= 222 neurons). Black dots (connected by black dashed line) represent the average displacement of the highest three displacements within a 50 µm radial band. Microfluidic valve has a radius of 200 µm indicated by the yellow shaded region. Majority of the cumulative displacement (>60 %) is localized to a 250 µm radius bordering the valve indicated by the blue dashed line at 450 µm. Neurons more than 750 µm from the microfluidic valve center have negligible displacement (<6% cumulative displacement) indicated by the green dashed line.

Figure 2 – Figure Supplement 2: Average calcium fluorescence from large peduncle ROI correlated with calcium fluorescence from smaller ROIs for individual peduncle neurons. a) Fluorescent image of transgenic Hydra expressing GCaMP6s pan-neuronally. Dashed while circle indicates the peduncle ROI. Small cyan circles indicate the individual neuron ROIs. Peduncle region is in the top right. The oral end of the Hydra (not shown in the image) is oriented towards the bottom left corner. b) Average calcium fluorescence trace (black) from the large peduncle ROI. Average calcium traces (gray) from individual neurons in the peduncle.

Figure 2 – Figure Supplement 3: Spikes in calcium fluorescence are due to calcium activity not motion artifacts. a) Fluorescence image of transgenic *Hydra* (nGreen) expressing GFP in neurons false colored with hot green colormap. Neurons appear in white. b) Fluorescence image of transgenic *Hydra*

expressing GCaMP6s in neurons false colored with hot green colormap. c) Fluorescence image of

transgenic *Hydra* expressing GCaMP7b in ectodermal epitheliomuscular cells false colored with hot green

colormap. a-c) Body length (white arrow), valve ROI (top circles), peduncle ROI (larger bottom circles)

and whole frame ROI (entire frame square) annotated on all three images for comparison. d) Average

body length trace time-aligned to stimulus onset from GFP *Hydra* (left) and GCaMP *Hydra* (middle, right).

Light gray shaded region indicates when the stimulation is on. e) Average fluorescence trace calculated

from the annotated peduncle ROI from GFP *Hydra* (left) and GCaMP *Hydra* (middle, right). f) Average

fluorescence trace calculated from the whole frame ROI from GFP *Hydra* (left) and GCaMP *Hydra*

(middle, right). g) Average fluorescence trace calculated from the annotated valve ROI from GFP *Hydra*

(left) and GCaMP *Hydra* (middle, right).

1282

1283

Figure 2 – Figure Supplement 4: Simultaneous electrophysiology and calcium imaging of

ectodermal epitheliomuscular cells. (a) Photograph (left) of a microfluidic immobilization chamber filled

with green dye. Black box highlights the recording region in the microfluidic chamber (110 μm tall). False

colored scanning electron micrograph (middle) shows the recording region (blue, photoresist; light grey,

Pt; dark grey, silica) on the nano-SPEAR chip (50 μm tall). Inset shows a zoom-in of the Pt electrode

(light grey) suspended mid-way between the top and bottom of the photoresist sidewall (blue). Brightfield

image (right) shows *Hydra* immobilized in the microfluidic chamber placed on top of the nano-SPEAR chip

with combined 160 μm tall recording region (Reproduced from Figure 2A, Badhiwala et al, 2018[31] by

permission of The Royal Society of Chemistry. This panel is not covered by the CC-BY 4.0 license, and

further reproduction of this panel would need permission from the copyright holder.) b) Simultaneous

electrophysiology and calcium imaging in transgenic *Hydra* (GCaMP6s, ectodermal epitheliomuscular

cells). Top trace shows mean fluorescence ($\Delta F/F$) from ectodermal epitheliomuscular cells (whole frame

ROI) in *Hydra* (GCaMP, ectodermal) (10Hz). Bottom trace shows simultaneously recorded electrical

activity from the *Hydra* (1KHz). High and low activity periods identified based on peak amplitude. Inset:

left box shows correlation during high activity period, contraction bursts. Traces show peaks in

fluorescence (top trace) coinciding with peaks in electrophysiology (bottom trace). A representative

fluorescence image shows high levels of fluorescence thus calcium activity in the entire body during

contraction burst events. Correlation map spatially plotting the correlation coefficient shows the entire

body has calcium activity correlated with electrophysiology. Inset: right box shows correlation during low

activity period, tentacle pulses. Traces show peaks in fluorescence (top trace) coinciding with peaks in

electrophysiology (bottom trace). A representative fluorescence image shows high levels of fluorescence

in the tentacle region during tentacle pulses. Cross correlation map shows the tentacle region has

calcium activity correlated with electrophysiology. Almost all spikes in electrophysiology coincide with

spike in fluorescence from ectodermal epitheliomuscular cells (and behavioral contractions).

1308

1309

Figure 2 – Figure Supplement 5: *Hydra*'s epitheliomuscular response is dependent on the

mechanical stimulus intensity. a) Gray trace (top) is the stimulation protocol. 20 min no stimulation, 20

min of repeated stimulation (1 sec 'on', 30 sec 'off') at 15 psi, 20 min of repeated stimulation (1 sec 'on',

30 sec 'off') at 20 psi, and 20 min of repeated stimulation (1 sec 'on', 30 sec 'off') at 25 psi. Stimulus 'on'

times indicated by vertical lines. Entire frame ROI used for analysis of the whole-body epithelial calcium

activity. Representative calcium fluorescence trace (blue) from ectodermal epitheliomuscles (GCaMP7b)

from animal stimulated with three different stimulus intensities for 20 min each (15, 20 and 25 psi).

Representative calcium fluorescence trace (teal) from ectodermal epitheliomuscles (GCaMP7b) from

animal stimulated with three different stimulus intensities for 20 min each (25, 20 and 15 psi). The

decrease in fluorescence amplitude observed for both increasing in and decreasing stimulus intensity is

due to photobleaching of the calcium indicator. b) Mechanosensory response probability at different

stimulus intensities (N = 3 animals). Response probability, fraction of trials that have at least one calcium

spike (also contraction pulse) occurring within 1 sec of stimulation onset.

1323

1324

Figure 2 – Figure Supplement 6: Mechanosensory response window. a) Different response windows

tested (1 sec in blue; 5 sec in orange; 10 sec in yellow; and 15 sec in purple) (Top). The lengths of the

rectangles correspond to time (sec) in the raster plot below. Stimulus trace in gray. Representative raster

plot of time-aligned contraction pulses from multiple trials superimposed from one animal stimulated at 20

psi every 31 sec for 60 min (Bottom). Each black dot is a spike in calcium fluorescence identified as a

contraction pulse. Stimulus is applied from 0 - 1 sec as indicated by a step in stimulus trace. b) Response

probability, fraction of trials that have at least one calcium spike (also contraction pulse) occurring during

different response windows (1 sec in blue; 5 sec in orange; 10 sec in yellow; and 15 sec in purple within

stimulation onset).

1334

1335

Figure 2 – Figure Supplement 7: Mechanical sensitivity of different body regions in *Hydra*. a)

Response probability of transgenic Hydra (N=3 animals expressing GCaMP6s in neurons) stimulated at

three different body regions: Oral, Mid-body and Aboral. Annotated Hydra below the plot indicates the

three stimulation regions used. b) Response probability of transgenic Hydra (n=8 animal expressing

GCaMP7b in ectodermal epitheliomuscular cells) Response probability is calculated using average

calcium fluorescence from neurons in the peduncle ROI in a) and using average calcium fluorescence

from the ectodermal epitheliomuscular cells from the entire body in b) and c).

1343

1344

Figure 2 – Figure Supplement 8: *Hydra's* mechanosensory response time is faster than passive

calcium diffusion through epitheliomuscular cells a) Distance calcium diffuses passively over given

time (left, black line plot) approximated using passive diffusion equation (right). Light brown shaded

region indicates the range of body length (~ 0.5 - 1 mm) of *Hydra* in microfluidic chambers used for the

experiments. Teal shaded region indicates the average time between stimulus onset and observed spike

in fluorescence (mean response time 0.5 - 1 sec). Yellow shaded region is the theoretical mean calcium

diffusion time calculated assuming passive diffusion. b-e) Mean calcium fluorescence time-aligned to

stimulus and normalized. Stimulus is applied at 0 sec (valve 'on' from 0 - 1 sec). b) Mean calcium activity

from the peduncle network of neurons at different stimulus intensities. c) Mean calcium activity from

ectodermal epitheliomuscular cells at different stimulus intensities. b-c) The line colors correspond to

different stimulus intensities shown at the bottom. d) Mean calcium activity from peduncle network of

neurons in different body resections stimulated at 20 psi (pressure for ~ 60% response probability in

whole animals). e) Mean calcium activity from ectodermal epitheliomuscular cells in different body

resections different body resections stimulated at 22 psi (pressure for ~ 60% response probability in

whole animals). d-e) The line colors correspond to the different resections shown at the bottom with

cartoon schematic of the resected *Hydra* with the same color.

1361

1362

Figure 2 – Figure Supplement 9: Long-term mechanical stimulation a) Representative calcium

fluorescence trace from the peduncle region from an animal (GcaMP6s, neurons) not stimulated and

animal mechanically stimulated with 20 psi. Stimulus protocol in gray, 20 min no stimulation, 120 min of

repeated stimulation (1 sec 'on', 30 sec 'off') at 20 psi followed by no stimulation. b) Spontaneous and

stimulated (mechanosensory response) contraction pulse probabilities for the first and second hour of

stimulation compared for each animal. Gray circles are mean contraction pulse probability during no

stimulation calculated from an average of 1 sec window shifted by ~ 0.3 sec over 30 sec intervals (N = 3).

Blue circles are contraction probability during stimulation calculated from 1 sec response window during

valve 'on' (N = 2). Light gray lines indicate the change in probabilities from hour 1 to hour 2 for each

individual. (paired t-test, n.s. = not significant)

1373

1374

Figure 3 – Figure Supplement 1: Regeneration of the peduncle network. a) Summary schematic of

the experiment performed. Dashed line indicates where the incision was made to remove the lower half of

the transgenic *Hydra* body (GcaMP6s, neurons) to discover when the peduncle neuronal network is

regenerated. b) Fluorescence images of "footless" *Hydra* during body contractions at various timepoints t

= 1 hr, 5 hr and 9 hr after bisection. Pink arrow indicates the open wound visible at t = 1 hr but not at

other timepoints. c) Fluorescence images of another "footless" *Hydra* during body contractions at various

timepoints t = 37 hr, 39 hr, 41 hr. Blue dashed circle indicates the region where the peduncle network is

found. Blue arrows indicate some of the neurons in the peduncle network with clearly well-connected

neurons at t = 41 hr.

1384

1385

Figure 3 - Figure Supplement 2: RT-qPCR analyses of neuron subtype-specific gene expression in

resected animals demonstrates loss of specific neuron subtypes. RT-qPCR was used to test for the

loss of specific neuron subtypes in whole, tube (no head or foot), headless, footless, and bisected *Hydra*

using uniquely expressed biomarkers for each subtype. There is not a specific biomarker for ec3A

(located in the basal disk), so the marker used to test for the presence or absence of this cell type is also

expressed in ec3B (located in the body column). Therefore, expression of the ec3A/B marker gene is

reduced, but not completely lost in animals with resected feet ("tube" and "footless"). However, ec5

expression (located in the peduncle above the basal disk) is completely lost in animals with resected feet,

thus it is clear that the ec3A subtype is completely lost in these animals. Biomarkers for neuron subtypes

located in the head and tentacles (ec1B, ec2, ec3C, ec4) are completely lost in animals with resected

heads ("headless" and "tube"). Data were analyzed with the $2^{-\Delta\Delta Ct}$ method and results were normalized to

the housekeeping gene *RP49* and to expression in whole animals.

1398

1399

Figure 3 – Figure Supplement 3: Mechanosensory response from ectodermal epitheliomuscular

cells and neurons in resected *Hydra*. a) Representative images of resection preparations of transgenic

Hydra (GCaMP7b, ectodermal epitheliomuscular cells: whole (or control), "footless", "headless",

"bisected", and "body column". White dashed square indicates the whole frame ROI used for quantifying

response. b) Spontaneous probability of at least one spiking event (or contraction pulse) occurring during

a random 1 sec window. (1 sec window slide by ~ 0.3 sec across 30 sec stimulation intervals; Kruskal-

Wallis test with post hoc dunn-sidak correction) c) Mechanosensory response probability during 1 sec of

valve on (Response probability, fraction of trials that have at least one calcium spike (also contraction

pulse) occurring within 1 sec of stimulation onset). Large circles indicate average values from all animals

combined for each condition. Small circles indicate probability from individual animals (one-way anova

with post hoc Bonferroni correction). Error bars are standard error of mean (S.E.M.) (N = 3 *Hydra* for each

condition; analysis from 60 min of stimulation which is imaging t = 20 – 80 min; n.s. = not significant, ** =

1412 p < 0.01, **** = p < 0.0001, ***** = p < 0.00001). d) Representative images of resection preparations of

1413 transgenic *Hydra* (GCaMP6s, neurons: whole (or control), "headless", and "bisected". White dashed circle

1414 indicates the peduncle ROI used for quantifying response. e) Spontaneous probability of at least one

1415 spiking event (or contraction pulse) occurring during a random 1 sec window. (1 sec window slide by ~

1416 0.3 sec across 30 sec stimulation intervals; Kruskal-Wallis test with post hoc dunn-sidak correction) f)

1417 Mechanosensory response probability during 1 sec of valve on. Large circles indicate average values

1418 from all animals combined for each condition. (Response probability, fraction of trials that have at least

1419 one calcium spike (also contraction pulse) occurring within 1 sec of stimulation onset.) Small circles

1420 indicate probability from individual animals (one-way anova with post hoc Bonferroni correction). Error

1421 bars are standard error of mean (S.E.M.) (N = 3 *Hydra* for each condition; analysis from 60 min of

1422 stimulation which is imaging t = 20 – 80 min; n.s. = not significant, ** = p < 0.01, **** = p < 0.0001, ***** =

1423 p < 0.00001).

1424

1425

1426 **Figure 3 – Figure Supplement 4: Contraction activity in non-stimulated animals.** Cartoon schematic

1427 of resection preparations of transgenic *Hydra* (GCaMP7b, ectodermal epitheliomuscular cells): whole (or

1428 control), "footless", "headless", "bisected" and "body column" animals. Entire frame ROI used for analysis

1429 of the whole-body epithelial calcium activity. Gray circles are mean contraction probability during t = 0 - 20

1430 min. Blue circles are mean contraction probability during t = 20 - 40 min, (this time corresponds to when

1431 stimulated animals receive mechanical stimuli). Mean probability calculated from 1 sec window shifted by

1432 ~ 0.3 sec over 30 sec intervals. (Response probability, fraction of trials that have at least one calcium

1433 spike (also contraction pulse) occurring within 1 sec of stimulation onset). Pressure in valves = 0 psi. Light

1434 gray lines pair the spontaneous contractions probability from the first 20 min interval with the second 20

1435 min interval from each individual animal. (Whole N = 4, footless N = 3, headless N = 3, bisected N = 5,

1436 body column N = 3 *Hydra*, paired t-test, n.s. = not significant)

1437

1438

Figure 3 – Figure Supplement 5: Hypostome and peduncle nerve rings work together to

coordinate contractile behavior a) Representative fluorescence trace used to calculate time interval

between contractions. b) Time interval between spontaneous contractions in animals with different

resections: whole, "footless", "headless", "bisected" and "body column". c) Time interval between

stimulated contractions in animals with different resections. d) Representative fluorescence trace used to

calculate time interval between contraction pulses. For illustration purpose, only a select few of the time

intervals are shown. e) Time interval between spontaneous contraction pulses in animals with different

resections. f) Time interval between stimulated contraction pulses in animals with different resections. g)

Representative fluorescence trace used to calculate contraction duration. h) Duration of spontaneous

contractions in animals with different resections. i) Duration of stimulated contractions in animals with

different resections. N= 3 *Hydra* (GcaMP7b, Ectodermal epitheliomuscles) per resection. (Kruskal-Wallis

test with dunn-sidak correction, n.s. = not significant, ** = $p < 0.01$, *** = $p < 0.001$, **** = $p < 0.0001$,*****

= $p < 0.00001$,****** = $p < 0.000001$)

1452

1453

Figure 4 – Figure Supplement 1: Single cell correlation analysis. a) Fluorescence image of

transgenic *Hydra* (#2) expressing GCaMP6s pan-neuronally. Individually tracked neuronal ROIs are

indicated with circles. The neurons are numbered in the order their traces are appear in c-e with the same

colors. Peduncle ROI outlined with white dashed line. b) Calcium fluorescence traces from single neurons

(top 6 traces) and average calcium fluorescence from peduncle ROI (bottom trace). Mechanically

responsive (MR) neurons are shown in shades of red. Contraction burst (CB) neurons are shown in

shades of blue. Unspecified neurons are shown in yellow, and their activity do not resemble any of the

previously identified neuronal networks (contraction burst, rhythmic potential or the mechanically

responsive reported here). c) Heat map shows the correlation coefficients of individually tracked neurons

and peduncle ROI. Colorbar is at the bottom. d) Average calcium fluorescence traces from each of the

neurons and peduncle ROI during spontaneous behaviors time-aligned with the onset of spontaneous

body contractions. Dashed line indicates the onset of body contraction. e) Average calcium fluorescence

traces from each of the neurons and peduncle ROI during stimulated behaviors time-aligned with the

onset of mechanical stimulation. Gray shaded rectangle indicates mechanical stimulation.

1468

1469

Figure 4 – Figure Supplement 2: Single cell correlation analysis. a) Fluorescence image of

transgenic *Hydra* (#3) expressing GCaMP6s pan-neuronally. Individually tracked neuronal ROIs are

indicated with circles. The neurons are numbered in the order their traces are appear in c-e with the same

colors. Peduncle ROI outlined with white dashed line. b) Calcium fluorescence traces from single neurons

(top 5 traces) and average calcium fluorescence from peduncle ROI (bottom trace). Mechanically

responsive (MR) neurons are shown in shades of red. Contraction burst (CB) neurons are shown in

shades of blue. Unspecified neurons are shown in yellow, and their activity do not resemble any of the

previously identified neuronal networks (contraction burst, rhythmic potential or the mechanically

responsive reported here). c) Heat map shows the correlation coefficients of individually tracked neurons

and peduncle ROI. Colorbar is at the bottom. d) Average calcium fluorescence traces from each of the

neurons and peduncle ROI during spontaneous behaviors time-aligned with the onset of spontaneous

body contractions. Dashed line indicates the onset of body contraction. e) Average calcium fluorescence

traces from each of the neurons and peduncle ROI during stimulated behaviors time-aligned with the

onset of mechanical stimulation. Gray shaded rectangle indicates mechanical stimulation.

1484

1485

Figure 4 – Figure Supplement 3: Random shuffling of single cell correlation analysis. Individual

neuronal fluorescence time-series were randomly shuffled to show the correlation coefficients were not

due to random chance in all three *Hydra* (each row is a different *Hydra*). Original correlation coefficients

from 10 mins of raw fluorescence time-series (a-c), shuffled correlation coefficients from randomly

reshuffled blocks reconstructing 10 mins of fluorescence time-series (d-e) and z-score of correlation

coefficients calculated based on random reshuffling (g-i). Original correlation coefficients were calculated

from the 10 mins of raw fluorescence activity for each of the neurons during mechanical stimulation.

Shuffled correlation coefficients were calculated by taking the mean of correlation coefficients from

randomly reshuffled fluorescence time-series. Briefly, the raw fluorescence time-series from 10 mins of

mechanical stimulation was divided into 100 blocks, which were then randomly recombined for each of

the neurons to calculated correlation coefficients and this random reshuffling was repeated 1000 times. Z-

score was calculated using the mean and standard deviation of the correlation coefficients calculated

from random reshuffling repeated 1000 times. The heatmap for z-score (g-i) resemble the original

correlation coefficients (a-c) indicating the correlation coefficients of the raw fluorescence time-series

were not a result of random chance. Except for the diagonals which represents autocorrelation, the mean

correlation coefficients resulting from randomly reshuffled fluorescence (e-f) are zero for all neuron pairs

indicating the correlation due to random chance.

1503

1504   **Description of Videos**

1505   **Video 1:**

1506   Microfluidic system to study mechanosensory response in *Hydra*. Dashed blue circle indicates the

1507   microfluidic valve that presses down on *Hydra.* Mechanosensory response from neurons and

1508   epitheliomuscular cells is shown.

1509

1510   **Video 2:**

1511   Spontaneous neural calcium activity in normal animals. Dashed blue circle indicates the ROI used for

1512   calcium trace shown in blue (bottom). (Playback 100x.)

1513

Video 3:

Stimulated neural calcium activity in normal animals. Dashed blue square indicates the ROI used for

calcium trace shown in blue (bottom). Dashed white circle indicates the location of the valve that presses

down on *Hydra* when inflated. Gray trace shows the stimulus protocol, where vertical lines indicate valve

'on' times. Stimulus applied beginning t = ~ 20 min and ends t = ~80 min. Valve is 'on' for 1 sec and 'off'

for 30 sec. (Playback 100x.)

1520

1521   **Video 4:**

1522   Spontaneous ectodermal epitheliomuscular calcium activity in normal animals. Dashed blue square

1523   indicates the ROI (entire frame) used for the calcium trace shown in blue (bottom). (Playback 100x.)

1524

Video 5:

Stimulated ectodermal epitheliomuscular calcium activity in normal animals. Dashed blue square

indicates the ROI (entire frame) used for calcium trace shown in blue (bottom). Dashed white circle

indicates the location of the valve that presses down on *Hydra* when inflated. Gray trace shows the

stimulus protocol, where vertical lines indicate valve 'on' times. Stimulus applied beginning t = ~ 20 min

and ends t = ~80 min. Valve is 'on' for 1 sec and 'off' for 30 sec. (Playback 100x.)

1531

1532

1533 **Video 6:**

1534 Spontaneous neural calcium activity in longitudinally bisected animals. Dashed blue circle indicates the

1535 ROI used for calcium trace shown in blue (bottom). (Playback 100x.)

1536

Video 7:

Stimulated neural calcium activity in longitudinally bisected animals. Dashed blue circle indicates the ROI

used for calcium trace shown in blue (bottom). Dashed white circle indicates the location of the valve that

presses down on *Hydra* when inflated. Gray trace shows the stimulus protocol, where vertical lines

indicate valve 'on' times. Stimulus applied beginning t = ~ 20 min and ends t = ~80 min. Valve is 'on' for 1

sec and 'off' for 30 sec. (Playback 100x.)

1543

1544 **Video 8:**

1545 Spontaneous neural calcium activity in headless animals. Dashed blue circle indicates the ROI used for

1546 calcium trace shown in blue (bottom). Animal is less active without the head and fewer spikes in calcium

1547 activity of peduncle neurons are observed. (Playback 100x.)

1548

Video 9:

Stimulated neural calcium activity in headless animals. Dashed blue circle indicates the ROI used for

calcium trace shown in blue (bottom). Dashed white circle indicates the location of the valve that presses

down on *Hydra* when inflated. Gray trace shows the stimulus protocol, where vertical lines indicate valve

'on' times. Stimulus applied beginning t = ~ 20 min and ends t = ~80 min. Valve is 'on' for 1 sec and 'off'

for 30 sec. Animal is less active without the head and fewer spikes in calcium activity of peduncle

neurons are observed without stimulation. After stimulation begins, more single pulse contractions occur.

(Playback 100x.)

1557

1558 **Video 10:**

1559   Spontaneous ectodermal epitheliomuscular calcium activity in footless animals. Dashed blue square

1560   indicates the ROI (entire frame) used for the calcium trace shown in blue (bottom). (Playback 100x.)

1561

Video 11:

Stimulated ectodermal epitheliomuscular calcium activity in footless animals. Dashed blue square

indicates the ROI (entire frame) used for calcium trace shown in blue (bottom). Dashed white circle

indicates the location of the valve that presses down on *Hydra* when inflated. Gray trace shows the

stimulus protocol, where vertical lines indicate valve 'on' times. Stimulus applied beginning t = ~ 20 min

and ends t = ~80 min. Valve is 'on' for 1 sec and 'off' for 30 sec. (Playback 100x.)

1568

1569   **Video 12:**

1570   Spontaneous ectodermal epitheliomuscular calcium activity in headless animals. Dashed blue square

1571   indicates the ROI (entire frame) used for the calcium trace shown in blue (bottom). Animal is less active

1572   without the head and fewer spikes in calcium activity of the whole body ectodermal epitheliomuscular

1573   cells are observed. (Playback 100x.)

1574

Video 13:

Stimulated ectodermal epitheliomuscular calcium activity in headless animals. Dashed blue square

indicates the ROI (entire frame) used for calcium trace shown in blue (bottom). Dashed white circle

indicates the location of the valve that presses down on *Hydra* when inflated. Gray trace shows the

stimulus protocol, where vertical lines indicate valve 'on' times. Stimulus applied beginning t = ~ 20 min

and ends t = ~80 min. Valve is 'on' for 1 sec and 'off' for 30 sec. Animal is less active without the head

and fewer spikes in calcium activity of epitheliomuscular cells are observed without stimulation. After

stimulation begins, more single pulse contractions occur. (Playback 100x.)

1583

1584   **Video 14:**

1585   Spontaneous ectodermal epitheliomuscular calcium activity in bisected animals. Dashed blue square

1586   indicates the ROI (entire frame) used for the calcium trace shown in blue (bottom). Animal is less active

1587   without the head and fewer spikes in calcium activity of the whole body ectodermal epitheliomuscular

1588   cells are observed. (Playback 100x.)

1589

Video 15:

Stimulated ectodermal epitheliomuscular calcium activity in bisected animals. Dashed blue square

indicates the ROI (entire frame) used for calcium trace shown in blue (bottom). Dashed white circle

indicates the location of the valve that presses down on *Hydra* when inflated. Gray trace shows the

stimulus protocol, where vertical lines indicate valve 'on' times. Stimulus applied beginning t = ~ 20 min

and ends t = ~80 min. Valve is 'on' for 1 sec and 'off' for 30 sec. Animal is less active without the head

and fewer spikes in calcium activity of epitheliomuscular cells are observed without stimulation. After

stimulation begins, more single pulse contractions occur. (Playback 100x.)

1598

1599   **Video 16:**

1600   Spontaneous ectodermal epitheliomuscular calcium activity in body column animals. Dashed blue square

1601   indicates the ROI (entire frame) used for the calcium trace shown in blue (bottom). Animal is less active

1602   without the head and fewer spikes in calcium activity of the whole body ectodermal epitheliomuscular

1603   cells are observed. (Playback 100x.)

1604

Video 17:

Stimulated ectodermal epitheliomuscular calcium activity in body column animals. Dashed blue square

indicates the ROI (entire frame) used for calcium trace shown in blue (bottom). Dashed white circle

indicates the location of the valve that presses down on *Hydra* when inflated. Gray trace shows the

stimulus protocol, where vertical lines indicate valve 'on' times. Stimulus applied beginning t = ~ 20 min

and ends t = ~80 min. Valve is 'on' for 1 sec and 'off' for 30 sec. Animal is less active without the head

and fewer spikes in calcium activity of epitheliomuscular cells are observed without stimulation. After

stimulation begins, more single pulse contractions occur. (Playback 100x.)

1613

Video 18:

Stimulated neural calcium activity in the hypostome and body column of normal animals. Dashed blue

circle indicates the ROI used for calcium trace shown in blue (bottom). Dashed white circle indicates the

location of the valve that presses down on *Hydra* when inflated. Gray trace shows the stimulus protocol,

where vertical lines indicate valve 'on' times. Stimulus applied beginning t = ~ 20 min, Valve is 'on' for 1

sec and 'off' for 30 sec. Imaged with 10x 0.45 N.A. objective (Playback 100x.)

1620

Source data

Figure 2 – source data 1: source file for stimulus dependent mechanosensory response.

The file (Fig2_SourceData.mat) contains all source data used for quantitatively characterizing

mechanosensory response shown in Figure 2. The file contains a struct named 'HydraData' with each

trial in a different row. The columns contain the raw and processed data for each trial: 'StimCondition'

indicates the stimulation pressure (0, 5, 10, 15, 20 or 25 psi) used during the trial; 'RawFluorescenceFoot'

contains the raw fluorescence values from foot/peduncle ROI; 'StimulationTrace' contains the trace of

stimulus delivery where 0 is valve off, 1 is valve on (and stimulation is being applied) per frame;

'Response Probability' is extracted after processing 'RawFluorescenceFoot' with stimulation onset times

from 'StimulationTrace'; 'TimeBetweenContractions' is time in seconds between contraction burst pulses

extracted from 'RawFluorescenceFoot'; 'SinglePulseContractions_Percent' is the percent of body

contractions that occur as single pulses extracted from 'RawFluorescenceFoot'

Figure 3 – source data 1: source file mechanosensory response in resected animals.

The file (Fig3_SourceData.mat) contains all source data used for characterizing mechanosensory

response in resected animals in Figure 3. The file contains a struct named 'HydraData' with each trial in a

different row. The columns contain the raw and processed data for each trial: 'StimCondition' indicates

stimulation condition (w/ or w/o stimulation; body resections - whole, bisected, no foot = Footless, no head

= Headless, no head and foot = BodyColumn); 'RawFluorescenceFoot' contains the raw fluorescence

values from foot/peduncle ROI; 'StimulationTrace' contains the trace of stimulus delivery where 0 is valve

off, 1 is valve on (and stimulation is being applied) per frame; 'Response Probability' is the stimulated

contraction probability extracted after processing 'RawFluorescenceFoot' with stimulation onset times

from 'StimulationTrace'; 'BasalContractionProbability' is spontaneous contraction probability extracted

after processing 'RawFluorescenceFoot' during (t = 0-20 min) the initial acclimation period without

stimulation.

Figure 4 – source data 1: source file for fluorescent calcium activity from single neuron ROIs.

The file (Fig4_SourceData.zip) contains three folders 'Hydra1', 'Hydra2' and 'Hydra3' for the three

different animals with individually tracked single neurons. Each folder contains a file called

'IndividualNeuronTraces.mat' with three different matrices: 'NeuronROI_RawFluorAllFrames' matrix [# of

rows = # of neurons+1, # of columns = # of frames] contain the raw fluorescence intensities individual

neuronal ROIs. Each row is different neuron and the last row in the matrix is the average fluorescence

from peduncle ROI'; 'ROI_xLocAllFrames' matrix [# of rows = # of neurons, # of columns = # of frames]

contains the x position of the center of circular ROIs for individual neurons; 'ROI_yLocAllFrames' matrix [#

of rows = # of neurons, # of columns = # of frames] contains the y position of the center of circular ROIs

for individual neurons