Adult Stem Cell-derived Complete Lung Organoid Models Emulate Lung Disease in COVID-19

Courtney Tindle[1,2]*, MacKenzie Fuller[1,2]*, Ayden Fonseca[1,2]*, Sahar Taheri[3]*, Stella-Rita Ibeawuchi[4], Nathan Beutler[5], Amanraj Claire[1,2], Vanessa Castillo[1], Moises Hernandez[6], Hana Russo[4], Jason Duran[7], Laura E. Crotty Alexander[8], Ann Tipps[4], Grace Lin[4], Patricia A. Thistlethwaite[6], Ranajoy Chattopadhyay[1, 2, 9]†, Thomas F. Rogers[3,10]†, Debashis Sahoo[3, 11-12]†, Pradipta Ghosh[1,2,12,13]†* and Soumita Das[2, 5]†*

[1]Department of Cellular and Molecular Medicine, University of California San Diego.
[2]HUMANOID CoRE, University of California San Diego.
[3]Department of Computer Science and Engineering, Jacobs School of Engineering, University of California San Diego.
[4]Department of Pathology, University of California San Diego.
[5]Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA 92037, USA.
[6]Department of Surgery, University of California San Diego.
[7]Division of Cardiology, Department of Internal Medicine, UC San Diego Medical Center, La Jolla 92037
[8]Pulmonary Critical Care Section, Veterans Affairs (VA) San Diego Healthcare System, La Jolla, California; Division of Pulmonary, Critical Care and Sleep Medicine, Department of Medicine,
[9]Cell Applications, Inc. San Diego, California
[10]Division of Infectious Diseases, Department of Medicine, University of California, San Diego, La Jolla, CA 92037, USA.
[11]Department of Pediatrics, University of California San Diego.
[12]Moores Cancer Center, University of California San Diego.
[13]Medicine, University of California San Diego.

* Equally contributed

† **Corresponding authors**:

- **Ranajoy Chattopadhyay, Ph.D.; Email:** r.chattopadhyay@cellapplications.com
- **Thomas F. Rogers, M.D., Ph.D; Email:** trogers@ucsd.edu
- **Debashis Sahoo, Ph.D.; Email:** dsahoo@ucsd.edu
- **Pradipta Ghosh, M.D.;** Professor, Department of Cellular and Molecular Medicine & Department of Medicine, Director, Institute for Network Medicine Executive Director, UC San Diego HUMANOID Center of Research Excellence (CoRE), University of California, San Diego, 9500 Gilman Drive, George E. Palade Bldg, Rm 232; La Jolla, CA 92093. **Phone**: 858-822-7633. **Email:** prghosh@ucsd.edu
- **Soumita Das, Ph.D.;** Associate Professor, Department of Pathology, Founding Director, UC San Diego HUMANOID Center of Research Excellence (CoRE), University of California, San Diego, 9500 Gilman Drive, George E. Palade Bldg, Rm 256; La Jolla, CA 92093. **Phone**: 858-246-2062: **Email:** sodas@ucsd.edu

Adult Stem Cell-derived Complete Lung Organoid Models Emulate Lung Disease in COVID-19

Courtney Tindle[1,2]*, MacKenzie Fuller[1,2]*, Ayden Fonseca[1,2]*, Sahar Taheri[3]*, Stella-Rita Ibeawuchi[4], Nathan Beutler[5], Amanraj Claire[1,2], Vanessa Castillo[1], Moises Hernandez[6], Hana Russo[4], Jason Duran[7], Laura E. Crotty Alexander[8], Ann Tipps[4], Grace Lin[4], Patricia A. Thistlethwaite[6], Ranajoy Chattopadhyay[1, 2, 9]†, Thomas F. Rogers[3,10]†, Debashis Sahoo[3, 11-12]†, Pradipta Ghosh[1,2,12,13]†* and Soumita Das[2, 5]†*

[1]*Department of Cellular and Molecular Medicine, University of California San Diego.*
[2]*HUMANOID CoRE, University of California San Diego.*
[3]*Department of Computer Science and Engineering, Jacobs School of Engineering, University of California San Diego.*
[4]*Department of Pathology, University of California San Diego.*
[5]*Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA 92037, USA.*
[6]*Division of Cardiothoracic Surgery, University of California San Diego.*
[7]*Division of Cardiology, Department of Internal Medicine, UC San Diego Medical Center, La Jolla 92037*
[8]*Pulmonary Critical Care Section, Veterans Affairs (VA) San Diego Healthcare System, La Jolla, California; Division of Pulmonary, Critical Care and Sleep Medicine, Department of Medicine,*
[9]*Cell Applications, Inc. San Diego, California*
[10]*Division of Infectious Diseases, Department of Medicine, University of California, San Diego, La Jolla, CA 92037, USA.*
[11]*Department of Pediatrics, University of California San Diego.*
[12]*Moores Cancer Center, University of California San Diego.*
[13]*Medicine, University of California San Diego.*

* Equally contributed

† **Corresponding authors**:

- **Ranajoy Chattopadhyay, Ph.D.; Email:** r.chattopadhyay@cellapplications.com
- **Thomas F. Rogers, M.D.; Ph.D; Email:** trogers@ucsd.edu
- **Debashis Sahoo, Ph.D.; Email:** dsahoo@ucsd.edu
- **Pradipta Ghosh, M.D.;** Professor, Department of Cellular and Molecular Medicine & Department of Medicine**,** Director, Institute for Network Medicine Executive Director, UC San Diego HUMANOID Center of Research Excellence (CoRE), University of California, San Diego, 9500 Gilman Drive, George E. Palade Bldg, Rm 232; La Jolla, CA 92093. **Phone**: 858-822-7633. **Email:** prghosh@ucsd.edu
- **Soumita Das, Ph.D.;** Associate Professor, Department of Pathology, Founding Director, UC San Diego HUMANOID Center of Research Excellence (CoRE), University of California, San Diego, 9500 Gilman Drive, George E. Palade Bldg, Rm 256; La Jolla, CA 92093. **Phone**: 858-246-2062: **Email:** sodas@ucsd.edu

GRAPHIC ABSTRACT

HIGHLIGHTS:

- **Human lung organoids with mixed proximodistal epithelia are created**
- **Proximal airway cells are critical for viral infectivity**
- **Distal alveolar cells are important for emulating host response**
- **Both are required for the overzealous response in severe COVID-19**

IN BRIEF: An integrated stem cell-based disease modeling and computational approach demonstrate how both proximal airway epithelium is critical for SARS-CoV-2 infectivity, but distal differentiation of alveolar pneumocytes is critical for simulating the overzealous host response in fatal COVID-19.

Summary (150 words)

SARS-CoV-2, the virus responsible for COVID-19, causes widespread damage in the lungs in the setting of an overzealous immune response whose origin remains unclear. We present a scalable, propagable, personalized, cost-effective adult stem cell-derived human lung organoid model that is complete with both proximal and distal airway epithelia. Monolayers derived from adult lung organoids (ALOs), primary airway cells, or hiPSC-derived alveolar type-II (AT2) pneumocytes were infected with SARS-CoV-2 to create *in vitro* lung models of COVID-19. Infected ALO-monolayers best recapitulated the transcriptomic signatures in diverse cohorts of COVID-19 patient-derived respiratory samples. The airway (proximal) cells were critical for sustained viral infection, whereas distal alveolar differentiation (AT2→AT1) was critical for mounting the overzealous host immune response in fatal disease; ALO monolayers with well-mixed proximodistal airway components recapitulated both. Findings validate a human lung model of COVID-19 , which can be immediately utilized to investigate COVID-19 pathogenesis and vet new therapies and vaccines.

Introduction

SARS-CoV-2, the virus responsible for COVID-19, causes widespread inflammation and injury in the lungs, giving rise to diffuse alveolar damage (DAD) (Arrossi and Farver, 2020; Borczuk et al., 2020; Damiani et al., 2020; Li et al., 2020; Roden et al., 2020), featuring marked infection and viral burden leading to apoptosis of alveolar pneumocytes (Hussman, 2020), along with pulmonary edema (Bratic and Larsson, 2013; Carsana et al., 2020). DAD leads to poor gas exchange and, ultimately, respiratory failure; the latter appears to be the final common mechanism of death in most patients with severe COVID-19 infection. How the virus causes so much damage remains unclear. A particular challenge is to understand the out-of-control immune reaction to the SARS-CoV-2 infection known as a cytokine storm, which has been implicated in many of the deaths from COVID-19. Although rapidly developed pre-clinical animal models have recapitulated some of the pathognomonic aspects of infection, e.g., induction of disease, and transmission, and even viral shedding in the upper and lower respiratory tract, many failed to develop severe clinical symptoms (Lakdawala and Menachery, 2020). Thus, the need for pre-clinical models remains both urgent and unmet.

To address this need, several groups have attempted to develop human pre-clinical COVID-19 lung models, all within the last few months (Duan et al., 2020; Mulay et al., 2020; Salahudeen et al., 2020). While a head-to-head comparison of the key characteristics of each model can be found in **Table 1**, what is particularly noteworthy is that none recapitulate the heterogeneous epithelial cellularity of both proximal and distal airways, i.e., airway epithelia, basal cells, secretory club cells and alveolar pneumocytes. Also noteworthy is that models derived from iPSCs lack propagability and/or cannot be reproducibly generated for biobanking; nor can they be scaled up in cost-effective ways for use in drug screens. Besides the approaches described so far, there are a few more approaches used for modeling COVID-19—(i) 3D organoids from bronchospheres and tracheospheres have been established before (Hild and Jaffe, 2016; Rock et al., 2009; Tadokoro et al., 2016) and are now used in apical-out cultures for infection with SARS-COV-2 (Suzuki et al., 2020); (ii) the most common model used for drug screening is the air-liquid interphase (ALI model) in which pseudo-stratified primary bronchial or small airway epithelial cells are used to recreate the multilayered mucociliary epithelium (Mou et al., 2016; Randell et al., 2011); (iii) several groups have also generated 3D airway models from iPSCs or tissue-resident stem cells (Dye et al., 2015; Ghaedi et al., 2013; Konishi et al., 2016; McCauley et al., 2017; Miller et al., 2019; Wong et al., 2012); (iv) others have generated AT2 cells from iPSCs using closely overlapping protocols of sequential differentiation starting with definitive endoderm, anterior foregut endoderm, and distal alveolar expression (Chen et al., 2017; Gotoh et al., 2014; Huang et al., 2014; Jacob et al., 2017; Jacob et al., 2019; Yamamoto et al., 2017). (v) Finally, long term *in vitro* culture conditions for pseudo-stratified airway epithelium organoids, derived from healthy and diseased adult humans suitable to assess virus infectivity (Sachs et al., 2019; van der Vaart and Clevers, 2020; Zhou et al., 2018) have been pioneered; unfortunately, these airway organoids expressed virtually no lung mesenchyme or alveolar signature. What remains unclear is if any of these models accurately recapitulate the immunopathologic phenotype that is seen in the lungs in COVID-19.

We present a rigorous transdisciplinary approach that systematically assesses an adult lung organoid model that is propagable, personalized and complete with both proximal airway and distal alveolar cell types

against existing models that are incomplete, and we cross-validate them all against COVID-19 patient-derived respiratory samples. Findings surprisingly show that cellular crosstalk between both proximal and distal components are necessary to emulate how SARS-CoV-2 causes diffuse alveolar pneumocyte damage; the proximal airway mounts a sustained viral infection, but it is the distal alveolar pneumocytes that mount the overzealous host response that has been implicated in a fatal disease.

Results

A rationalized approach for creating and validating acute lung injury in COVID-19

To determine which cell types in the lungs might be most readily infected, we began by analyzing a human lung single-cell sequencing dataset (GSE132914) for the levels of expression of angiotensin-converting enzyme-II (ACE2) and Transmembrane Serine Protease 2 (TMPRSS2), the two receptors that have been shown to be the primary sites of entry for the SARS-CoV-2 (Hoffmann et al., 2020). The dataset was queried with widely accepted markers of all the major cell types (see **Table 2**). Alveolar epithelial type 2 (AT2), ciliated and club cells emerged as the cells with the highest expression of both receptors (**Fig 1A; Fig S1A**). These observations are consistent with published studies demonstrating that ACE2 is indeed expressed highest in AT2 and ciliated cells (Jia et al., 2005; Mulay et al., 2020; Zhao et al., 2020). In a cohort of deceased COVID-19 patients, we observed by H&E (**Fig S1B**) that gas-exchanging flattened AT1 pneumocytes are virtually replaced by cuboidal cells that were subsequently confirmed to be AT2-like cells via immunofluorescent staining with the AT2-specific marker, surfactant protein-C (SFTPC; **Fig 1B** *upper panel*; **Fig S1C**; *top*). We also confirmed that club cells express ACE2 (**Fig S1C**; *bottom*), underscoring the importance of preserving these cells in any ideal lung model of COVID-19. When we analyzed the lungs of deceased COVID-19 patients, the presence of SARS-COV-2 in alveolar pneumocytes was also confirmed, as determined by the colocalization of viral nucleocapsid protein with SFTPC (**Fig 1B**; *lower panel;* **Fig S1D**). Immunohistochemistry studies further showed the presence of SARS-COV-2 virus in alveolar pneumocytes and in alveolar immune cells (**Fig S1E**). These findings are consistent with the gathering consensus that alveolar pneumocytes support the interaction between the epithelial cells and inflammatory cells recruited to the lung; via mechanisms that remain unclear, they are generally believed to contribute to the development of acute lung injury and acute respiratory distress syndrome (ARDS), the severe hypoxemic respiratory failure during COVID-19 (Hou et al., 2020; Spagnolo et al., 2020). Because prior work has demonstrated that SARS-CoV-2 infectivity in patient-derived airway cells is highest in the proximal airway epithelium compared to the distal alveolar pneumocytes (AT1 and AT2) (Hou et al., 2020), and yet, it is the AT2 pneumocytes that harbor the virus, and the AT1 pneumocytes that are ultimately destroyed during diffuse alveolar damage, we hypothesized that both proximal airway and distal (alveolar pneumocyte) components might play distinct roles in the respiratory system to mount the so-called viral infectivity and host immune response phases of the clinical symptoms observed in COVID-19 (Chen and Li, 2020).

Because no existing lung model provides such proximodistal cellular representation (**Table 1**), and hence, may not recapitulate with accuracy the clinical phases of COVID-19, we first sought to develop a lung

model that is complete with both proximal and distal airway epithelia using adult stem cells that were isolated from deep lung biopsies. Lung organoids were generated using the protocol outlined in **Fig 1C** and detailed in *Methods*. Organoids grown in 3D cultures were subsequently dissociated into single cells to create 2D-monolayers (either maintained submerged in media or used in ALI model) for SARS-CoV-2 infection, followed by RNA Seq analysis. Primary airway epithelial cells and hiPSC-derived alveolar type-II (AT2) pneumocytes were used as additional models (**Fig 1D**; *left panel*). Each of these transcriptomic datasets was subsequently used to cross-validate our *ex-vivo* lung models of SARS-CoV-2 infection with the human COVID-19 autopsy lung specimens (**Fig 1D**; *right panel*) to objectively vet each model for their ability to accurately recapitulate the gene expression signatures in the patient-derived lungs.

Creation of a lung organoid model, complete with both proximal and distal airway epithelia

Three lung organoid lines were developed from deep lung biopsies obtained from the normal regions of lung lobes surgically resected for lung cancer; both genders, smokers and non-smokers were represented (**Fig S2A; Table 3**). Three different types of media were compared (**Fig S2B**); the composition of these media was inspired either by their ability to support adult-stem cell-derived mixed epithelial cellularity in other organs (like the gastrointestinal tract (Miyoshi and Stappenbeck, 2013; Sato et al., 2009; Sayed et al., 2020), or rationalized based on published growth conditions for proximal and distal airway components (Gotoh et al., 2014; Sachs et al., 2019; van der Vaart and Clevers, 2020). A growth condition that included conditioned media from L-WRN cells which express Wnt3, R-spondin and Noggin, supplemented with recombinant growth factors, which we named as '*lung organoid expansion media*' emerged as superior compared to alveolosphere media-I and II(Jacob et al., 2019; Yamamoto et al., 2017) (details in the methods), based on its ability to consistently and reproducibly support the best morphology and growth characteristics across multiple attempts to isolate organoids from lung tissue samples. Three adult lung organoid lines (ALO1-3) were developed using the expansion media, monitored for their growth characteristics by brightfield microscopy and cultured with similar phenotypes until P10 and beyond (**Fig S2C-D**). The 3D morphology of the lung organoid was also assessed by H&E staining of slices cut from formalin-fixed paraffin-embedded (FFPE) cell blocks of *HistoGel*-embedded ALO1-3 (**Fig S2E**).

To determine if all the 6 major lung epithelial cells (illustrated in **Fig 2A**) are present in the organoids, we analyzed various cell-type markers by qRT-PCR (**Fig 2B-H**). All three ALO lines had a comparable level of AT2 cell surfactant markers (compared against hiPSC-derived AT2 cells as positive control) and a significant amount of AT1, as determined using the marker AQP5. ALOs also contained basal cells (as determined by the marker ITGA6), ciliated cells (as determined by the marker FOXJ1), club cells (as determined by the marker SCGB1A1) and stem cells (as determined by marker p75-NGFR). As expected, the primary human bronchial epithelial cells (NHBE) had significantly higher expression of basal cell markers than the ALO lines (hence, served as a positive control), but they lacked stemness and club cells (hence, served as a negative control).

The presence of all cell types was also confirmed by assessing protein expression of various cell types within organoids grown in 3D cultures. Two different approaches were used—(i) slices cut from FFPE cell blocks of *HistoGel*-embedded ALO lines (**Fig 2I-J**) or (ii) ALO lines grown in 8-well chamber slides were fixed in matrigel (**Fig 2K**), stained, and assessed by confocal microscopy. Such staining not only confirmed the presence of all cell types in each ALO line but also demonstrated the presence of more than one cell type (i.e., mixed cellularity) of proximal (basal-KRT5) and distal (AT1/AT2 markers) within the same organoid structure. For example, AT2 and basal cells, marked by SFTPB and KRT5, respectively, were found in the same 3D-structure (**Fig 2J**, interrupted curved line). Similarly, ciliated cells and goblet cells stained by Ac-Tub and Muc5, respectively, were found to coexist within the same structure (**Fig 2J**, interrupted box; **Fig 2K**, arrow). The presence of heterogeneous cellularity was documented in all three ALO lines (see multiple additional examples in **Fig S3**).

Organoid cellularity resembles tissue sources in 3D cultures but differentiates in 2D cultures

To model respiratory infections such as COVID-19, it is necessary for pathogens to be able to access the apical surface. Many groups have modeled such access by dissociating 3D organoids into single cells and plating them as 2D-monolayers (Duan et al., 2020; Han et al., 2020; Huang et al., 2020; Mulay et al., 2020; Sachs et al., 2019; Zhou et al., 2018). Because the loss of dimensionality can have a major impact on cellular proportions and impact disease-modeling in unpredictable ways, we assessed the impact of the 3D-to-2D conversion on cellularity by RNA seq analyses. Two commonly encountered methods of growth in 2D-monolayers were tested: (i) monolayers polarized on trans-well inserts but submerged in growth media, and ii) monolayers were grown at the air-liquid interface (popularly known as the 'ALI model'(Dvorak et al., 2011; Prytherch et al., 2011)) for 21-days to differentiate into the mucociliary epithelium (see **Fig 3A; Fig S4A-E**). The submerged 2D-monolayers had several regions of organized vacuolated-appearing spots (**Fig S4A, D**; arrow), presumably due to morphogenesis and cellular organization even in 2D. The ALI-monolayers appeared to be progressively hazier with time after air-lift, likely due to the accumulation of secreted mucin (**Fig S4E**), and formed an intact epithelial barrier, as determined by trans-epithelial electrical resistance (TEER) (**Fig S4C**). RNA Seq datasets were analyzed using the same set of cell markers, as we used in **Fig 1A** (listed in **Table 2**). Cell-type deconvolution of our dataset using CIBERSORTx (https://cibersortx.stanford.edu/runcibersortx.php) showed that cellular proportions in the human lung tissues were also relatively well-preserved in organoids grown in 3D over several passages (**Fig 3B**; *left*); both showed a mixed population of simulated alveolar, basal, club, ciliated and goblet cells. When 3D organoids were dissociated and plated as 2D monolayers on transwells, the AT2 signatures were virtually abolished with a concomitant and prominent emergence of AT1 signatures, suggesting that growth in 2D-monolayers favor differentiation of AT2 cells into AT1 cells (Shami and Evans, 2015)(**Fig 3B**; *middle*). A compensatory reduction in proportion was also observed for the club, goblet and ciliated cells. The same organoids, when grown in long-term 2D culture conditions in the ALI model, showed a strikingly opposite pattern; alveolar signatures were almost entirely replaced by a concomitant increase in ciliated and goblet cells (**Fig 3B**; *right*). These findings are consistent with the well-established notion that ALI conditions favor growth as pseudo-

stratified mucociliary epithelium(Dvorak et al., 2011; Prytherch et al., 2011). As an alternative model for use as monolayers for viral infection, we developed hiPSC-derived AT2 cells and alveolospheres (**Fig 3C**), using established protocols (Huang et al., 2020). Because they were grown in the presence of CHIR99021 (an aminopyrimidine derivate that is a selective and potent Wnt agonist) (Abdelwahab et al., 2019; Jacob et al., 2019; Yamamoto et al., 2017), which probably inhibits the AT2→AT1 differentiation, these monolayers were enriched for AT2 and devoid of AT1 cells (**Fig 3D**).

The multicellularity of lung organoid monolayers was also confirmed by immunofluorescence staining and confocal microscopy of the submerged and ALI monolayers, followed by the visualization of cell markers in either max-projected z-stacks (**Fig 3E**; *left*) or orthogonal views of the same (**Fig 3E**; *right*). As expected, markers for the same cell type (i.e., SFTPB and SFTPC, both AT2 markers) colocalize, but markers for different cell types do not. Submerged monolayers showed the prominent presence of both AT1 (AQP5-positive) and AT2 cells. Compared to the submerged monolayers, the ALI model showed a significant increase in the ciliated epithelium (as determined by Ac Tub; compare Ac Tub stained panels in **Fig 3E** with **3F**). This increase was associated with a concomitant decrease in both KRT5-stained basal and SFTPC-positive AT2 cells (**Fig 3F**).

Taken together, the immunofluorescence images are in agreement with the RNA seq dataset; both demonstrate that the short-term submerged monolayer favors distal differentiation (AT2→AT1), whereas the 21-day ALI model favors proximal mucociliary differentiation. It is noteworthy that these distinct differentiation phenotypes originated from the same 3D-organoids despite the seeding of cells in the same basic media composition (i.e., PneumaCult™) prior to switching over to an ALI-maintenance media for the prolonged growth at Air-Liquid interface; the latter is a well-described methodology that promotes differentiation into ciliated and goblet cells (Rayner et al., 2019).

Differentiated 2D-monolayers show that SARS-CoV-2 infectivity is higher in proximal than distal epithelia

Because the lung organoids with complete proximodistal cellularity could be differentiated into either proximal-predominant monolayers in submerged short-term cultures or distal-predominant monolayers in long-term ALI cultures, this provided us with an opportunity to model the respiratory tract and assess the impact of the virus along the entire proximal-to-distal gradient. We first asked how viral infectivity varies in the two models. Because multiple groups have shown the importance of the ciliated airway cells for infectivity (i.e., viral entry, replication and apical release(Hou et al., 2020; Hui et al., 2020; Milewska et al., 2020; Zhu et al., 2020)), as positive controls, we infected monolayers of human airway epithelia (see **Fig S4F-I**). AT2 cells, which express high levels of viral entry receptors ACE2 and TMPRSS2 (**Fig 1A**, **Fig S1A**) have been shown to be proficient in viral entry, but are least amenable to sustained viral release and infectivity(Hou et al., 2020; Hui et al., 2020). To this end, we infected monolayers of hiPSC-derived homogeneous cultures of AT2 cells as secondary controls (see **Fig S4J-L**). Infection was carried out using the Washington strain of SARS-CoV-2, USA-WA1/2020 [BEI Resources NR-52281 (Rogers et al., 2020)]. As expected, the 2D-lung monolayers we generated, both the submerged or the ALI models, were readily infected with SARS-CoV-2 (**Fig S5A**), as determined by the presence of the viral

envelope gene (E-gene; **Fig 3G**); however, the kinetics of viral amplification differed. When expressed as levels of E gene normalized to the peak values in each model (**Fig 3G**), the kinetics of the ALI-monolayer model mirrored that of the primary airway epithelial monolayers; both showed slow beginning (0 – 48 hpi) followed by an exponential increase in E gene levels from 48 to 72 hpi. The submerged monolayer model showed sustained viral infection during the 48-72 hpi window (**Fig S5A;** *left*). In the case of AT2 cells, the 48-72 hpi window was notably missing in monolayers of hiPSC-derived AT2 cells (**Fig 3G**; **Fig S5A;** *right*). When we specifically analyzed the kinetics of viral E gene expression during the late phase (48-72 hpi window), we found that proximal airway models [human Bronchial airway Epi (HBEpC)] were more permissive than distal models [human Small Airway Epi (HSAEpC) and AT2] to viral replication (**Fig S5B**); the ALO monolayers showed intermediate sustained infectivity (albeit with variability). All models showed extensive cell death and detachment by 96 hrs and, hence, were not analyzed. Confocal imaging of infected ALO monolayers with anti-SARS-COV-2 nucleocapsid protein antibody showed that submerged ALO monolayers did indeed show progressive changes during the 48 to 72h window after infection (**Fig 3H**): by 48 hpi we observed the formation of 'reticulovesicular patterns' that are indicative of viral replication within modified host endoplasmic reticulum (Knoops et al., 2008) (**Fig 3H**; *left*), and by 72 hpi we observed focal cytopathic effect (CPE) (Kaye, 2006) such as cell-rounding, detachment, and bursting of virions (**Fig 3H**; *right*, **FigS5C**).

Taken together, these findings show that sustained viral infectivity is best simulated in monolayers that resemble the proximal mucociliary epithelium, i.e., 2D-monolayers of lung organoids grown as ALI models and the primary airway epithelia. Because prior studies conducted in patient-derived airway cells (Hou et al., 2020) mirror what we see in our monolayers, we conclude that proximal airway cells within our mixed-cellular model appear to be sufficient to model viral infectivity in COVID-19.

Differentiated 2D-monolayers show that host immune response is higher in distal than proximal epithelia

Next, we asked if the newly generated lung models accurately recapitulate the host immune response in COVID-19. To this end, we analyzed the infected ALO monolayers (both the submerged and ALI variants) as well as the airway epithelial (HSAEpC) and AT2 monolayers by RNA seq and compared them all against the transcriptome profile of lungs from deceased COVID-19 patients. A publicly available dataset (GSE151764) (Nienhold et al., 2020), comprised of lung transcriptomes from victims deceased either due to non-infectious causes (controls) or due to COVID-19, was first analyzed for differentially expressed genes (**Fig 4A-B**). This cohort was chosen as a test cohort over others because it was the largest one available at the time of this study with appropriate postmortem control samples. Differentially expressed genes showed an immunophenotype that was consistent with what is expected in viral infections (**Fig 4C; Table 4**; **Fig S6**), and showed overrepresentation of pathways such as interferon, immune, and cytokine signaling (**Fig 4D; Table 5**; **Fig S7**). Differentially expressed gene signatures and reactome pathways that were enriched in the test cohort were fairly representative of the host immune response observed in patient-derived respiratory samples in multiple other validation cohorts; the signature derived from the test cohort could consistently classify control (normal) samples from COVID-19-samples (ROC AUC 0.89 to 1.00 across the board; **Fig 4E**). The most notable finding is that the patient-derived

signature was able to perfectly classify the EpCAM-sorted epithelial fractions from the bronchoalveolar lavage fluids of infected and healthy subjects (ROC AUC 1.00; GSE145926-Epithelium (Liao et al., 2020), suggesting that the respiratory epithelium is a major site where the host immune response is detected in COVID-19. When compared to existing organoid models of COVID-19, we found that the patient-derived COVID-19-lung signature was able to perfectly classify infected *vs.* uninfected late passages (> 50) of hiPSC-derived AT1/2 monolayers (GSE155241) (Han et al., 2020) and infected *vs.* uninfected liver and pancreatic organoids (**Fig 4F**). The COVID-19-lung signatures failed to classify commonly used respiratory models, e.g., A549 cells and bronchial organoids, as well as intestinal organoids (**Fig 4F**). A similar analysis on our own lung models revealed that the COVID-19-lung signature was induced in submerged monolayers with distal-predominant AT2→AT1 differentiation, but not in the proximal-predominant ALI model (ROC AUC 1.00 and 0.50, respectively; **Fig 4G**). The ALI model and the small airway epithelia, both models that mimic the airway epithelia (and lack alveolar pneumocytes; see **Fig 3B**), failed to mount the patient-derived immune signatures (**Fig 4H**; *left*). These findings suggested that the presence of alveolar pneumocytes is critical for emulating host response. To our surprise, induction of the COVID-19-lung signature also failed in hiPSC-derived AT2 monolayers (**Fig 4H**; *right*), indicating that AT2 cells are unlikely to be the source of such host response. These findings indicate that both proximal airway and AT2 cells, when alone, are insufficient to induce the host immune response that is encountered in the lungs of COVID-19 patient.

Next, we analyzed the datasets from our ALO monolayers for differentially expressed genes when challenged with SARS-COV-2 (**Fig 5A-B**). Genes and pathways upregulated in the infected lung organoid-derived monolayer models overlapped significantly with those that were upregulated in the COVID-19 lung signature (compare **Fig 4C-D** with **5C-D**, **Table 6-7**; **Fig S8-S9**). That we observed a partial overlap in upregulated genes and no overlaps among downregulated genes between model and disease (COVID-19) (**Fig 5E**) likely reflect the missing stromal and immune components in our organoid monolayers. Regardless of these missing components, the model-derived DEG signature was sufficient to consistently and accurately classify diverse cohorts of patient-derived respiratory samples (ROC AUC ranging from 0.88 to 1.00; **Fig 5F**); the model-derived DEG signature was significantly induced in COVID-19 samples compared to normal controls (**Fig 5G-H**). Most importantly, the model-derived DEG signature was significantly induced in the epithelial cells recovered from bronchoalveolar lavage (**Fig 5I**).

Taken together, these cross-validation studies from disease to model (**Fig 4**) and *vice versa* (**Fig 5**) provide an objective assessment of the match between the host response in COVID-19 lungs and our submerged ALO monolayers. Such a match was not seen in the case of the other models, e.g., the proximal airway-mimic ALI model, HSAEpC monolayer, or hiPSC-derived AT2 models. Because the submerged ALO monolayers contained both proximal airway epithelia (basal cells) and promoted AT2→AT1 differentiation, findings demonstrate that mixed cellular monolayers can mimic the host response in COVID-19. A subtractive analysis revealed that the cell type that is shared between models which showed induction of host response signatures (i.e., ALO submerged monolayers and GSE155241 (Han et al., 2020); **Fig 5F**) but is absent in models that do not show such response (hu Bronchial organoids, small airway epi, ALI-model of ALO) is AT1. We conclude that distal differentiation from AT2→AT1, a complex process that is comprised of distinct intermediates (Choi et al., 2020), is essential for modeling the host immune response in COVID-19.

Both proximal and distal airway epithelia are required to mount the overzealous host response in COVID-19

We next asked which model best simulated the overzealous host immune response that has been widely implicated in fatal COVID-19. To this end, we relied upon a recently described artificial intelligence (AI)-guided definition of the nature of the overzealous response in fatal COVID-19 (Sahoo et al., 2020). Using ACE2 as a seed gene, a 166-gene signature was identified and validated as an invariant immune response that was shared among all respiratory viral pandemics, including COVID-19 (**Fig 6A**). A subset of 20 genes within the 166-gene signature was subsequently identified as a determinant of disease severity/fatality; these 20 genes represented translational arrest, senescence, and apoptosis. These two signatures referred to as ViP (166-gene) and severe ViP (20-gene) signatures, were used as a computational framework to first vet existing SARS-CoV-2 infection models that have been commonly used for therapeutic screens (**Fig 6B-D**). Surprisingly, we found that each model fell short in one way or another. For example, the Vero E6, which is a commonly used cultured cell model, showed a completely opposite response; instead of being induced, both the 166-gene and 20-gene ViP signatures were suppressed in infected Vero E6 monolayers (**Fig 6B**). Similarly, neither ViP signature was induced in the case of SARS-CoV-2 challenged human bronchial organoids(Suzuki et al., 2020) (**Fig 6C**). Finally, in the case of the hiPSC-derived AT1/2 organoids, which recapitulated the COVID-19-lung derived immune signatures (in **Fig 4F**), the 166-gene ViP signature was induced significantly (**Fig 6D**; *top*), but the 20-gene severity signature was not (**Fig 6D**; *bottom*). These findings show that none of the existing models capture the overzealous host immune response that has been implicated in a fatality.

Our lung models showed that both the 166- and 20-gene ViP signatures were induced significantly in the submerged ALO-derived monolayers that had distal differentiation (**Fig 6E;** *left*), but not in the proximal-mimic ALI model (**Fig 6E;** *right*). Neither signatures were induced in monolayers of small airway epithelial cells (**Fig 6F**) or hiPSC-derived AT2 cells (**Fig 6G**).

Taken together with our infectivity analyses, these findings demonstrate that although the proximal airway epithelia and AT2 cells may be infected, and as described by others (Dye et al., 2015; Hou et al., 2020), may be vital for mounting a viral response and for disease transmission, these cells alone cannot mount the overzealous host immune response that is associated with the fatal disease. Similarly, even though the alveolar pneumocytes, AT1 and AT2 cells, are sufficient to mount the host immune response, in the absence of proximal airway components, they too are insufficient to recapitulate the severe ViP signature that is characterized by cellular senescence and apoptosis. However, when both proximal and distal components are present, i.e., basal, ciliated and AT1 cells, the model mimicked the overzealous host immune response in COVID-19 (**Fig 6H**).

Discussion

The most important discovery we report here is the creation of adult lung organoids that are complete with both proximal airway and distal alveolar epithelia; these organoids can not only be stably propagated and expanded in 3D cultures but also used as monolayers of mixed cellularity for modeling viral and host immune responses during respiratory viral pandemics. Furthermore, an objective analysis of this model and other existing SARS-

CoV-2-infected lung models against patient-lung derived transcriptomes showed that the model which most closely emulates the elements of viral infectivity, lung injury, and inflammation in COVID-19 is one that contained both proximal and distal alveolar signatures (**Fig 6H**), whereas, the presence of just one or the other fell short.

There are three important impacts of this work. First, successful modeling of the human lung organoids that are complete with both proximal and distal signatures has not been accomplished before. The multicellularity of the lung has been a daunting challenge that many experts have tried to recreate *in vitro*; in fact, the demand for perfecting such a model has always remained high, not just in the current context of the COVID-19 pandemic but also with the potential of future pandemics. We have provided the evidence that the organoids that were created using our methodology retain proximal and distal cellularity throughout multiple passages and even within the same organoid. Although a systematic design of experiment (DoE) approach (Bukys et al., 2020) was not involved in getting to this desirable goal, a rationalized approach was taken. For example, a Wnt/R-spondin/Noggin-containing conditioned media was used as a source of the so-called 'niche factors' for any organoid growth (Sato and Clevers, 2015). This was supplemented with recombinant FGF7/10; FGF7 is known to help cell proliferation and differentiation and is required for normal branching morphogenesis (Padela et al., 2008), whereas FGF10 helps in cell maturation (Rabata et al., 2020) and in alveolar regeneration upon injury (Yuan et al., 2019). Together, they are likely to have directed the differentiation toward distal lung lineages (hence, the preservation of alveolar signatures). The presence of both distal alveolar and proximal ciliated cells was critical: proximal cells were required to recreate sustained viral infectivity, and the distal alveolar pneumocytes, in particular, the ability of AT2 cells to differentiate into AT1 pneumocytes was essential to recreate the host response. It is possible that the response is mediated by a distinct AT2-lineage population, i.e., damage-associated transient progenitors (DATPs), which arise as intermediates during AT2→AT1 differentiation upon injury-induced alveolar regeneration (Choi et al., 2020). Although somewhat unexpected, the role of AT1 pneumocytes in mounting innate immune responses has been documented before in the context of bacterial pneumonia (Wong and Johnson, 2013; Yamamoto et al., 2012). In work (Huang et al., 2020) that was published during the preparation of this manuscript, authors used long-term ALI models of hiPSC-derived AT2 monolayers (in growth conditions that inhibit AT2→AT1 differentiation, as we did here for our AT2 model) and showed that SARS-CoV-2 induces iAT2-intrinsic cytotoxicity and inflammatory response, but failed to induce type 1 interferon pathways (IFN α/β). It is possible that prolonged culture of iAT2 pneumocytes gives rise to some DATPs but cannot robustly do so in the presence of inhibitors of AT1 differentiation. This (spatially segregated viral and host immune response) is a common theme across many lung infections (including bacterial pneumonia and other viral pandemics (Chan et al., 2013; Hou et al., 2020; Taubenberger and Morens, 2008; Weinheimer et al., 2012) and hence, this mixed cellularity model is appropriate for use in modeling diverse lung infections and respiratory pandemics to come.

Second, among all the established lung models so far, ours features 4 key properties that are desirable whenever disease models are being considered for their use in HTP modes for rapid screening of candidate therapeutics and vaccines —(i) reproducibility, propagability and scalability, (ii) cost-effectiveness, (iii) personalization, and (iv) modularity, with the potential to be used in multi-dimensional complex lung models with

other cell types if/when needed. We showed that the protocol we have optimized supports isolation, expansion and propagability at least up to 12-15 passages (at the time of submission of this work), with documented retention of proximal and distal airway components up to P8 (by RNA seq). Feasibility has also been established for scaling up for use in 384-well HTP assays. We also showed that the protocols for generating lung organoids could be reproduced in both genders and regardless of the donor's smoking status, consistency in outcome and growth characteristics were observed across all isolation attempts. The ALOs are also cost-effective; the need for exclusive reliance on recombinant growth factors was replaced at least in part with conditioned media from a commonly used cell line (L-WRN/ ATCC® CRL-2647 cells). Such media has a batch to batch stable cocktail of Wnt, R-Spondin, and Noggin and has been shown to improve reproducibility in the context of GI organoids in independent laboratories (VanDussen et al., 2019). The use of conditioned media may not only have made our model more cost-effective than others but also likely improved the rigor and reproducibility in lung modeling and research. Because the model is propagable, repeated iPSC-reprogramming (another expensive step) is also eliminated, further cutting costs compared to many other models. As for personalization, our model is derived from adult lung stem cells from deep lung biopsies; each organoid line was established from one patient. By avoiding iPSCs or EPSCs, this model not only captures genetics but also retains organ-specific epigenetic programming in the lung, and hence, potentially additional programming that may occur in disease (such as in the setting of chronic infection, injury, inflammation, somatic mutations, etc.). The ability to replicate donor phenotype and genotype in vitro allows for potential use as pre-clinical human models for Phase '0' clinical trials. As for modularity, by showing that the 3D lung organoids could be used as polarized monolayers on transwells to allow infectious agents to access the apical surface (in this case, SARS-CoV-2), we demonstrate that the organoids have the potential to be reverse-engineered with additional components in a physiologically relevant spatially segregated manner: for example, immune and stromal cells can be placed in the lower chamber to model complex lung diseases that are yet to be modeled and have no cure (e.g., Idiopathic pulmonary fibrosis, etc).

Third, the value of the ALO models is further enhanced due to the availability of companion readouts/ biomarkers (e.g., ViP signatures in the case of respiratory viral pandemics, or monitoring the E gene, or viral shedding, etc.) that can rapidly and objectively vet treatment efficacy based on set therapeutic goals. Of these readouts, the host response, as assessed by ViP signatures, is a key vantage point because an overzealous host response is what is known to cause fatality. Recently, a systematic review of the existing pre-clinical animal models revealed that most of the animal models of COVID-19 recapitulated mild patterns of human COVID-19; no severe illness associated with mortality was observed, suggesting a wide gap between COVID-19 in humans (Spagnolo et al., 2020) and animal models (Ehaideb et al., 2020). The model revealed here, in conjunction with the ViP signatures described earlier (Sahoo et al., 2020), could serve as a pre-clinical model with companion diagnostics to identify drugs that target both the viral and host response in pandemics.

Acknowledgements

This work was supported by the National Institutes for Health (NIH) grants 1R01DK107585-01A1, 3R01DK107585-05S1 (to SD); R01-AI141630, CA100768 and CA160911 (to PG) and R01-AI 155696 (to PG, DS and SD); R00-CA151673 and R01-GM138385 (to DS), R01- HL32225 (to PT), UCOP-R00RG2642 (to SD and PG), UCOP-R01RG3780 (to P.G. and D.S) and a pilot award from the Sanford Stem Cell Clinical Center at UC San Diego Health (P.G, S.D, D.S). L.C.A's salary was supported in part by the VA San Diego Healthcare System. The authors would like to thank Victor Pretorius, Rachel White and Jen Bigbee (Department of Cardiothoracic Surgery, UC San Diego) who assisted with thoracotomies during rapid autopsies. This manuscript includes data generated at the UC San Diego Institute of Genomic Medicine (IGC) using an Illumina NovaSeq 6000 that was purchased with funding from a National Institutes of Health SIG grant (#S10 OD026929).

Author contributions

Role	Authors
Conceptualization; Funding acquisition; Supervision; Project administration	S.D., P.G., D.S.,
Investigation; Methodology; Data curation, Formal Analysis • Experiments on organoids, monolayers (C.T, M.F, A.F) • Experiments on iPSC-derived AT2 cells (M.F, R.C) • Computational analysis (S.T, D.S, P.G) • Data curation and analysis (D.S., P.G., S.D. S.T., M.F., A.F, C.T) • Rapid Autopsy of Lungs (A.C., V.C., J.D., L.C.A,) • Essential resources (patient samples, tissues, etc) (T.F.R, N.B, J.D., L.C.A, A.T, H.R, P.a.T, M.H)	C.T, M.F, A.F, S.T, SR.I, N.B, A.C, V.C, M.H, H.R, J.D, L.C.A, A.T, G.L, P.A.T, R.C, T.F.R, D.S., S.D., P.G.
Funding acquisition	D.S., S.D., P.G.
Software	D.S.
Visualization; Writing – original draft; Writing – review & editing	P.G., S.D.

Competing interests: The authors declare no competing interests.

References

Abdelwahab, E.M.M., Rapp, J., Feller, D., Csongei, V., Pal, S., Bartis, D., Thickett, D.R., and Pongracz, J.E. (2019). Wnt signaling regulates trans-differentiation of stem cell like type 2 alveolar epithelial cells to type 1 epithelial cells. Respir Res *20*, 204.

Arrossi, A.V., and Farver, C. (2020). The pulmonary pathology of COVID-19. Cleve Clin J Med.

Borczuk, A.C., Salvatore, S.P., Seshan, S.V., Patel, S.S., Bussel, J.B., Mostyka, M., Elsoukkary, S., He, B., Del Vecchio, C., Fortarezza, F.*, et al.* (2020). COVID-19 pulmonary pathology: a multi-institutional autopsy cohort from Italy and New York City. Mod Pathol.

Bratic, A., and Larsson, N.G. (2013). The role of mitochondria in aging. J Clin Invest *123*, 951-957.

Bukys, M.A., Mihas, A., Finney, K., Sears, K., Trivedi, D., Wang, Y., Oberholzer, J., and Jensen, J. (2020). High-Dimensional Design-Of-Experiments Extracts Small-Molecule-Only Induction Conditions for Dorsal Pancreatic Endoderm from Pluripotency. iScience *23*, 101346.

Carsana, L., Sonzogni, A., Nasr, A., Rossi, R.S., Pellegrinelli, A., Zerbi, P., Rech, R., Colombo, R., Antinori, S., Corbellino, M.*, et al.* (2020). Pulmonary post-mortem findings in a series of COVID-19 cases from northern Italy: a two-centre descriptive study. Lancet Infect Dis *20*, 1135-1140.

Chan, R.W., Chan, M.C., Agnihothram, S., Chan, L.L., Kuok, D.I., Fong, J.H., Guan, Y., Poon, L.L., Baric, R.S., Nicholls, J.M.*, et al.* (2013). Tropism of and innate immune responses to the novel human betacoronavirus lineage C virus in human ex vivo respiratory organ cultures. J Virol *87*, 6604-6614.

Chen, Y., and Li, L. (2020). SARS-CoV-2: virus dynamics and host response. Lancet Infect Dis *20*, 515-516.

Chen, Y.W., Huang, S.X., de Carvalho, A., Ho, S.H., Islam, M.N., Volpi, S., Notarangelo, L.D., Ciancanelli, M., Casanova, J.L., Bhattacharya, J.*, et al.* (2017). A three-dimensional model of human lung development and disease from pluripotent stem cells. Nat Cell Biol *19*, 542-549.

Choi, J., Park, J.E., Tsagkogeorga, G., Yanagita, M., Koo, B.K., Han, N., and Lee, J.H. (2020). Inflammatory Signals Induce AT2 Cell-Derived Damage-Associated Transient Progenitors that Mediate Alveolar Regeneration. Cell Stem Cell *27*, 366-382 e367.

Damiani, S., Fiorentino, M., De Palma, A., Foschini, M.P., Lazzarotto, T., Gabrielli, L., Viale, P.L., Attard, L., Riefolo, M., and D'Errico, A. (2020). Pathological Post Mortem Findings in Lungs Infected With Sars-Cov 2. J Pathol.

Duan, F., Guo, L., Yang, L., Han, Y., Thakur, A., Nilsson-Payant, B.E., Wang, P., Zhang, Z., Ma, C.Y., Zhou, X.*, et al.* (2020). Modeling COVID-19 with Human Pluripotent Stem Cell-Derived Cells Reveals Synergistic Effects of Anti-inflammatory Macrophages with ACE2 Inhibition Against SARS-CoV-2. Res Sq.

Dvorak, A., Tilley, A.E., Shaykhiev, R., Wang, R., and Crystal, R.G. (2011). Do airway epithelium air-liquid cultures represent the in vivo airway epithelium transcriptome? Am J Respir Cell Mol Biol *44*, 465-473.

Dye, B.R., Hill, D.R., Ferguson, M.A., Tsai, Y.H., Nagy, M.S., Dyal, R., Wells, J.M., Mayhew, C.N., Nattiv, R., Klein, O.D.*, et al.* (2015). In vitro generation of human pluripotent stem cell derived lung organoids. Elife *4*.

Ehaideb, S.N., Abdullah, M.L., Abuyassin, B., and Bouchama, A. (2020). A systematic review uncovers a wide-gap between COVID-19 in humans and animal models MedRxiv.

Ghaedi, M., Calle, E.A., Mendez, J.J., Gard, A.L., Balestrini, J., Booth, A., Bove, P.F., Gui, L., White, E.S., and Niklason, L.E. (2013). Human iPS cell-derived alveolar epithelium repopulates lung extracellular matrix. J Clin Invest *123*, 4950-4962.

Gotoh, S., Ito, I., Nagasaki, T., Yamamoto, Y., Konishi, S., Korogi, Y., Matsumoto, H., Muro, S., Hirai, T., Funato, M.*, et al.* (2014). Generation of alveolar epithelial spheroids via isolated progenitor cells from human pluripotent stem cells. Stem Cell Reports *3*, 394-403.

Han, Y., Yang, L., Duan, X., Duan, F., Nilsson-Payant, B.E., Yaron, T.M., Wang, P., Tang, X., Zhang, T., Zhao, Z.*, et al.* (2020). Identification of Candidate COVID-19 Therapeutics using hPSC-derived Lung Organoids. bioRxiv.

Hild, M., and Jaffe, A.B. (2016). Production of 3-D Airway Organoids From Primary Human Airway Basal Cells and Their Use in High-Throughput Screening. Curr Protoc Stem Cell Biol *37*, IE 9 1-IE 9 15.

Hoffmann, M., Kleine-Weber, H., Schroeder, S., Kruger, N., Herrler, T., Erichsen, S., Schiergens, T.S., Herrler, G., Wu, N.H., Nitsche, A.*, et al.* (2020). SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell *181*, 271-280 e278.

Hou, Y.J., Okuda, K., Edwards, C.E., Martinez, D.R., Asakura, T., Dinnon, K.H., 3rd, Kato, T., Lee, R.E., Yount, B.L., Mascenik, T.M.*, et al.* (2020). SARS-CoV-2 Reverse Genetics Reveals a Variable Infection Gradient in the Respiratory Tract. Cell *182*, 429-446 e414.

Huang, J., Hume, A.J., Abo, K.M., Werder, R.B., Villacorta-Martin, C., Alysandratos, K.D., Beermann, M.L., Simone-Roach, C., Lindstrom-Vautrin, J., Olejnik, J.*, et al.* (2020). SARS-CoV-2 Infection of Pluripotent Stem Cell-Derived Human Lung Alveolar Type 2 Cells Elicits a Rapid Epithelial-Intrinsic Inflammatory Response. Cell Stem Cell.

Huang, S.X., Islam, M.N., O'Neill, J., Hu, Z., Yang, Y.G., Chen, Y.W., Mumau, M., Green, M.D., Vunjak-Novakovic, G., Bhattacharya, J.*, et al.* (2014). Efficient generation of lung and airway epithelial cells from human pluripotent stem cells. Nat Biotechnol *32*, 84-91.

Hui, K.P.Y., Cheung, M.C., Perera, R., Ng, K.C., Bui, C.H.T., Ho, J.C.W., Ng, M.M.T., Kuok, D.I.T., Shih, K.C., Tsao, S.W.*, et al.* (2020). Tropism, replication competence, and innate immune responses of the coronavirus SARS-CoV-2 in human respiratory tract and conjunctiva: an analysis in ex-vivo and in-vitro cultures. Lancet Respir Med *8*, 687-695.

Hussman, J.P. (2020). Cellular and Molecular Pathways of COVID-19 and Potential Points of Therapeutic Intervention. Front Pharmacol *11*, 1169.

Jacob, A., Morley, M., Hawkins, F., McCauley, K.B., Jean, J.C., Heins, H., Na, C.L., Weaver, T.E., Vedaie, M., Hurley, K.*, et al.* (2017). Differentiation of Human Pluripotent Stem Cells into Functional Lung Alveolar Epithelial Cells. Cell Stem Cell *21*, 472-488 e410.

Jacob, A., Vedaie, M., Roberts, D.A., Thomas, D.C., Villacorta-Martin, C., Alysandratos, K.D., Hawkins, F., and Kotton, D.N. (2019). Derivation of self-renewing lung alveolar epithelial type II cells from human pluripotent stem cells. Nat Protoc *14*, 3303-3332.

Jia, H.P., Look, D.C., Shi, L., Hickey, M., Pewe, L., Netland, J., Farzan, M., Wohlford-Lenane, C., Perlman, S., and McCray, P.B., Jr. (2005). ACE2 receptor expression and severe acute respiratory syndrome coronavirus infection depend on differentiation of human airway epithelia. J Virol *79*, 14614-14621.

Kaye, M. (2006). SARS-associated coronavirus replication in cell lines. Emerg Infect Dis *12*, 128-133.

Knoops, K., Kikkert, M., Worm, S.H., Zevenhoven-Dobbe, J.C., van der Meer, Y., Koster, A.J., Mommaas, A.M., and Snijder, E.J. (2008). SARS-coronavirus replication is supported by a reticulovesicular network of modified endoplasmic reticulum. PLoS Biol *6*, e226.

Konishi, S., Gotoh, S., Tateishi, K., Yamamoto, Y., Korogi, Y., Nagasaki, T., Matsumoto, H., Muro, S., Hirai, T., Ito, I.*, et al.* (2016). Directed Induction of Functional Multi-ciliated Cells in Proximal Airway Epithelial Spheroids from Human Pluripotent Stem Cells. Stem Cell Reports *6*, 18-25.

Lakdawala, S.S., and Menachery, V.D. (2020). The search for a COVID-19 animal model. Science *368*, 942-943.

Li, Y., Wu, J., Wang, S., Li, X., Zhou, J., Huang, B., Luo, D., Cao, Q., Chen, Y., Chen, S.*, et al.* (2020). Progression to Fibrosing Diffuse Alveolar Damage in a Series of 30 Minimally Invasive Autopsies with COVID-19 Pneumonia in Wuhan, China. Histopathology.

Liao, M., Liu, Y., Yuan, J., Wen, Y., Xu, G., Zhao, J., Cheng, L., Li, J., Wang, X., Wang, F.*, et al.* (2020). Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19. Nat Med *26*, 842-844.

McCauley, K.B., Hawkins, F., Serra, M., Thomas, D.C., Jacob, A., and Kotton, D.N. (2017). Efficient Derivation of Functional Human Airway Epithelium from Pluripotent Stem Cells via Temporal Regulation of Wnt Signaling. Cell Stem Cell *20*, 844-857 e846.

Milewska, A., Kula-Pacurar, A., Wadas, J., Suder, A., Szczepanski, A., Dabrowska, A., Owczarek, K., Marcello, A., Ochman, M., Stacel, T.*, et al.* (2020). Replication of Severe Acute Respiratory Syndrome Coronavirus 2 in Human Respiratory Epithelium. J Virol *94*.

Miller, A.J., Dye, B.R., Ferrer-Torres, D., Hill, D.R., Overeem, A.W., Shea, L.D., and Spence, J.R. (2019). Generation of lung organoids from human pluripotent stem cells in vitro. Nat Protoc *14*, 518-540.

Miyoshi, H., and Stappenbeck, T.S. (2013). In vitro expansion and genetic modification of gastrointestinal stem cells in spheroid culture. Nat Protoc *8*, 2471-2482.

Mou, H., Vinarsky, V., Tata, P.R., Brazauskas, K., Choi, S.H., Crooke, A.K., Zhang, B., Solomon, G.M., Turner, B., Bihler, H.*, et al.* (2016). Dual SMAD Signaling Inhibition Enables Long-Term Expansion of Diverse Epithelial Basal Cells. Cell Stem Cell *19*, 217-231.

Mulay, A., Konda, B., Garcia, G., Yao, C., Beil, S., Sen, C., Purkayastha, A., Kolls, J.K., Pociask, D.A., Pessina, P.*, et al.* (2020). SARS-CoV-2 infection of primary human lung epithelium for COVID-19 modeling and drug discovery. bioRxiv.

Nienhold, R., Ciani, Y., Koelzer, V.H., Tzankov, A., Haslbauer, J.D., Menter, T., Schwab, N., Henkel, M., Frank, A., Zsikla, V.*, et al.* (2020). Two distinct immunopathological profiles in autopsy lungs of COVID-19. MedRxiv.

Padela, S., Yi, M., Cabacungan, J., Shek, S., Belcastro, R., Masood, A., Jankov, R.P., and Tanswell, A.K. (2008). A critical role for fibroblast growth factor-7 during early alveolar formation in the neonatal rat. Pediatr Res *63*, 232-238.

Prytherch, Z., Job, C., Marshall, H., Oreffo, V., Foster, M., and BeruBe, K. (2011). Tissue-Specific stem cell differentiation in an in vitro airway model. Macromol Biosci *11*, 1467-1477.

Rabata, A., Fedr, R., Soucek, K., Hampl, A., and Koledova, Z. (2020). 3D Cell Culture Models Demonstrate a Role for FGF and WNT Signaling in Regulation of Lung Epithelial Cell Fate and Morphogenesis. Front Cell Dev Biol *8*, 574.

Randell, S.H., Fulcher, M.L., O'Neal, W., and Olsen, J.C. (2011). Primary epithelial cell models for cystic fibrosis research. Methods Mol Biol *742*, 285-310.

Rayner, R.E., Makena, P., Prasad, G.L., and Cormet-Boyaka, E. (2019). Optimization of Normal Human Bronchial Epithelial (NHBE) Cell 3D Cultures for in vitro Lung Model Studies. Sci Rep *9*, 500.

Rock, J.R., Onaitis, M.W., Rawlins, E.L., Lu, Y., Clark, C.P., Xue, Y., Randell, S.H., and Hogan, B.L. (2009). Basal cells as stem cells of the mouse trachea and human airway epithelium. Proceedings of the National Academy of Sciences of the United States of America *106*, 12771-12775.

Roden, A.C., Bois, M.C., Johnson, T.F., Aubry, M.C., Alexander, M.P., Hagen, C.E., Lin, P.T., Quinton, R.A., Maleszewski, J.J., and Boland, J.M. (2020). The Spectrum of Histopathologic Findings in Lungs of Patients with Fatal COVID-19 Infection. Arch Pathol Lab Med.

Rogers, T.F., Zhao, F., Huang, D., Beutler, N., Burns, A., He, W.T., Limbo, O., Smith, C., Song, G., Woehl, J.*, et al.* (2020). Isolation of potent SARS-CoV-2 neutralizing antibodies and protection from disease in a small animal model. Science *369*, 956-963.

Sachs, N., Papaspyropoulos, A., Zomer-van Ommen, D.D., Heo, I., Bottinger, L., Klay, D., Weeber, F., Huelsz-Prince, G., Iakobachvili, N., Amatngalim, G.D.*, et al.* (2019). Long-term expanding human airway organoids for disease modeling. EMBO J *38*.

Sahoo, D., Katkar, G.D., Khandelwal, S., Behroozikhah, M., Claire, A., Castillo, V., Tindle, C., Fuller, M., Taheri, S., Rogers, T.F.*, et al.* (2020). AI-guided discovery of the invariant host response to viral pandemics. BioRxiv.

Salahudeen, A.A., Choi, S.S., Rustagi, A., Zhu, J., de la, O.S., Flynn, R.A., Margalef-Catala, M., Santos, A.J.M., Ju, J., Batish, A.*, et al.* (2020). Progenitor identification and SARS-CoV-2 infection in long-term human distal lung organoid cultures. bioRxiv.

Sato, T., and Clevers, H. (2015). SnapShot: Growing Organoids from Stem Cells. Cell *161*, 1700-1700 e1701.

Sato, T., Vries, R.G., Snippert, H.J., van de Wetering, M., Barker, N., Stange, D.E., van Es, J.H., Abo, A., Kujala, P., Peters, P.J.*, et al.* (2009). Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. Nature *459*, 262-265.

Sayed, I.M., Suarez, K., Lim, E., Singh, S., Pereira, M., Ibeawuchi, S.R., Katkar, G., Dunkel, Y., Mittal, Y., Chattopadhyay, R.*, et al.* (2020). Host engulfment pathway controls inflammation in inflammatory bowel disease. FEBS J.

Shami, S.G., and Evans, M.J. (2015). Kinetics of Pulmonary Cells. In book: Comparative Biology of the Normal Lung.

Spagnolo, P., Balestro, E., Aliberti, S., Cocconcelli, E., Biondini, D., Casa, G.D., Sverzellati, N., and Maher, T.M. (2020). Pulmonary fibrosis secondary to COVID-19: a call to arms? Lancet Respir Med *8*, 750-752.

Suzuki, T., Itoh, Y., Sakai, Y., Saito, A., Okuzaki, D., Motooka, D., Minami, S., Kobayashi, T., Yamamoto, T., Okamoto, T.*, et al.* (2020). Generation of human bronchial organoids for SARS-CoV-2 research. BioRxiv.

Tadokoro, T., Gao, X., Hong, C.C., Hotten, D., and Hogan, B.L. (2016). BMP signaling and cellular dynamics during regeneration of airway epithelium from basal progenitors. Development *143*, 764-773.

Taubenberger, J.K., and Morens, D.M. (2008). The pathology of influenza virus infections. Annu Rev Pathol *3*, 499-522.

van der Vaart, J., and Clevers, H. (2020). Airway organoids as models of human disease. J Intern Med.

VanDussen, K.L., Sonnek, N.M., and Stappenbeck, T.S. (2019). L-WRN conditioned medium for gastrointestinal epithelial stem cell culture shows replicable batch-to-batch activity levels across multiple research teams. Stem Cell Res *37*, 101430.

Weinheimer, V.K., Becher, A., Tonnies, M., Holland, G., Knepper, J., Bauer, T.T., Schneider, P., Neudecker, J., Ruckert, J.C., Szymanski, K.*, et al.* (2012). Influenza A viruses target type II pneumocytes in the human lung. J Infect Dis *206*, 1685-1694.

Wong, A.P., Bear, C.E., Chin, S., Pasceri, P., Thompson, T.O., Huan, L.J., Ratjen, F., Ellis, J., and Rossant, J. (2012). Directed differentiation of human pluripotent stem cells into mature airway epithelia expressing functional CFTR protein. Nat Biotechnol *30*, 876-882.

Wong, M.H., and Johnson, M.D. (2013). Differential response of primary alveolar type I and type II cells to LPS stimulation. PLoS One *8*, e55545.

Yamamoto, K., Ferrari, J.D., Cao, Y., Ramirez, M.I., Jones, M.R., Quinton, L.J., and Mizgerd, J.P. (2012). Type I alveolar epithelial cells mount innate immune responses during pneumococcal pneumonia. J Immunol *189*, 2450-2459.

Yamamoto, Y., Gotoh, S., Korogi, Y., Seki, M., Konishi, S., Ikeo, S., Sone, N., Nagasaki, T., Matsumoto, H., Muro, S.*, et al.* (2017). Long-term expansion of alveolar stem cells derived from human iPS cells in organoids. Nat Methods *14*, 1097-1106.

Yuan, T., Volckaert, T., Redente, E.F., Hopkins, S., Klinkhammer, K., Wasnick, R., Chao, C.M., Yuan, J., Zhang, J.S., Yao, C.*, et al.* (2019). FGF10-FGFR2B Signaling Generates Basal Cells and Drives Alveolar Epithelial Regeneration by Bronchial Epithelial Stem Cells after Lung Injury. Stem Cell Reports *12*, 1041-1055.

Zhao, Y., Zhao, Z., Wang, Y., Zhou, Y., Ma, Y., and Zuo, W. (2020). Single-Cell RNA Expression Profiling of ACE2, the Receptor of SARS-CoV-2. Am J Respir Crit Care Med *202*, 756-759.

Zhou, J., Li, C., Sachs, N., Chiu, M.C., Wong, B.H., Chu, H., Poon, V.K., Wang, D., Zhao, X., Wen, L.*, et al.* (2018). Differentiated human airway organoids to assess infectivity of emerging influenza virus. Proceedings of the National Academy of Sciences of the United States of America *115*, 6822-6827.

Zhu, N., Wang, W., Liu, Z., Liang, C., Wang, W., Ye, F., Huang, B., Zhao, L., Wang, H., Zhou, W.*, et al.* (2020). Morphogenesis and cytopathic effect of SARS-CoV-2 infection in human airway epithelial cells. Nat Commun *11*, 3910.

<div style="text-align: center;">**FIGURE LEGENDS**</div>

Figure 1. A rationalized approach to building and validating human pre-clinical models of COVID-19. A. Whisker plots display relative levels of ACE2 expression in various cell types in the normal human lung. The cell types were annotated within a publicly available single-cell sequencing dataset (GSE132914) using genes listed in Table S1. *p* values were analyzed by one-way Anova and Tukey post hoc test. **B.** FFPE sections of the human lung from normal and deceased COVID-19 patients were stained for SFTPC, alone or in combination with nucleocapsid protein and analyzed by confocal immunofluorescence. Representative images are shown. Scale bar = 20 µm. **C.** Schematic showing key steps generating an adult stem cell-derived, propagable, lung organoid model, complete with proximal and distal airway components for modeling COVID-19-in-a-dish. **D.** A transcriptome-based approach is used for cross-validation of *in vitro* lung models of SARS-CoV-2 infection (left) versus the human disease, COVID-19 (right), looking for a match in gene expression signatures.

Figure 2. Adult stem cell-derived lung organoids are propagatable models with both proximal and distal airway components. **A.** Schematic lists the various markers used here for qPCR and immunofluorescence to confirm the presence of all cell types in the 3D lung organoids here and in 2D monolayers later (in Fig 3). **B-H.** Bar graphs display the relative abundance of various cell type markers (normalized to 18S) in adult lung organoids (ALO), compared to the airway (NHBE) and/or alveolar (AT2) control cells, as appropriate. *p* values were analyzed by one-way Anova. Error bars denote S.E.M; n = 3-6 datasets from 3 independent ALOs and representing early and late passages. See also **Supplementary Figure 3** for individual ALOs. **I-J**. H&E-stained cell blocks were prepared using HistoGel (I). Slides were stained for the indicated markers and visualized by confocal immunofluorescence microscopy. Representative images are shown in J. Scale bar = 50 µm. **K**. 3D organoids grown in 8-well chamber slides were fixed, immunostained and visualized by confocal microscopy as in J. Scale bar = 50 µm. See also **Supplementary Figure 2**.

Figure 3. **Monolayers derived from lung organoids differentiate into proximal and distal airway components**. **A-B**. Samples collected at various steps of lung organoid isolation and expansion in culture, and from the two types of monolayers prepared using the lung organoids were analyzed by RNA seq and the datasets were compared for % cellular composition. Schematic in A shows the steps, and bar plots in B show the relative proportion of various lung cell types. GLL = general lung lineage; VEM = viral entry markers (ACE2 and TMPRSS2). **C-D**. hiPSC-derived AT2 cells and alveolospheres (C) were plated as monolayers and analyzed by RNA seq. Bar plots in D show % cellular composition. **E-F**. Submerged ALO monolayers in transwells (E) or monolayers were grown as ALI models (F) were fixed and stained for the indicated markers and visualized by confocal immunofluorescence microscopy. The representative max projected Z-stack images (left) and the corresponding orthogonal images (right) are displayed. Arrows indicate AT2 cells. Arrowhead = Club cells. Scale bar = 20 µm. **G.** Monolayers of ALO, hiPSC-derived AT2 cells, and other alternative models (see **Supplementary Figure 4**) were infected or not with SARS-CoV-2 and analyzed for infectivity by qPCR (targeted amplification of viral envelope, E gene (G). **H**. Monolayers of ALO1-3 were challenged with SARS-CoV-2 for indicated time points

prior to fixation and staining for KRT5, SARS-COV2 viral nucleocapsid protein and DAPI and visualized by confocal microscopy. Representative images are shown, displaying reticulovesicular network patterns and various cytopathic effects. Scale bar = 15 μm. See also **Supplementary Figure 5** for infectivity observed in alternative models.

Figure 4. **Gene expression patterns in the lungs of patients with COVID-19 is recapitulated in lung organoid monolayers infected with SARS-CoV-2**. **A-C**. Publicly available RNA Seq datasets (GSE151764) of lung autopsies from patients who were deceased due to COVID-19 or non-infectious causes (healthy normal control) were analyzed for differential expression of genes (B), and displayed as a heatmap labeled with selected genes in C. See also **Supplementary Figure 6** for a heatmap with all genes labeled. **D**. Reactome-pathway analysis shows the major pathways up- or downregulated in COVID-19 lungs. See also **Supplementary Figure 7** for visualization as hierarchical reacfoam. **E**. Bar plots display the ability of the differentially expressed genes in the test cohort (GSE151764) to classify human COVID-19 respiratory samples from four other independent cohorts. **F**. Bar plots display the ability of the differentially expressed genes in the test cohort (GSE151764) to classify published *in vitro* models for SARS-CoV-2 infection where RNA Seq datasets were either generated in this work or publicly available. **G-H.** Bar (top) and violin (bottom) plots compare the relative accuracy of disease modeling in four *in vitro* models used in the current work, as determined by the induction of COVID-19 lung signatures in each model. G; monolayer (left) and ALI models (right) prepared using adult lung organoids (ALOs). H; Primary human small airway epithelium (left) and hiPSC –derived AT2-monolayers (right).

Figure 5. **Genes and pathways induced in the SARS-CoV-2-infected lung organoid monolayers are induced also in the lungs of COVID-19 patients. A-C**. Adult lung organoid monolayers infected or not with SARS-CoV-2 were analyzed by RNA seq and differential expression analysis. Differentially expressed genes (B) are displayed as a heatmap labeled with selected genes in C. See also **Supplementary Figure 8** for a heatmap with all genes labeled. **D**. Reactome-pathway analysis shows the major pathways upregulated in SARS-CoV-2-infected lung organoid monolayers. See also **Supplementary Figure 9** for visualization as hierarchical reactome. **E**. A Venn diagram showing overlaps in DEGs between model (current work; 5B) and disease (COVID-19 lung dataset, GSE151764; Fig 4). **F**. Bar plots display the ability of the DEGs in infected lung monolayers to classify human normal vs COVID-19 respiratory samples from five independent cohorts. **G-I**. Bar (top) and violin (bottom) plots compare the accuracy of disease modeling in three publicly available human lung datasets, as determined by the significant induction of the DEGs that were identified in the SARS-CoV-2-challenged monolayers.

Figure 6. Both proximal and distal airway components are required to model the overzealous host response in COVID-19. A. Schematic summarizing the immune signatures identified based on ACE2-equivalent gene induction observed invariably in any respiratory viral pandemic. The 166-gene ViP signature captures the cytokine storm in COVID-19, whereas the 20-gene subset severe ViP signature is indicative of disease severity/fatality. **B-D**. Publicly available RNA Seq datasets from commonly used lung models, Vero E6 (B),

human bronchial organoids (C) and hPSC-derived AT1/2 cell-predominant lung organoids are classified using the 166-gene ViP signature (top row) and 20-gene severity signature (bottom row). **E-G**. RNA Seq datasets generated in this work using either human small airway epithelial cells (E), adult lung organoids as submerged or ALI models (left and right, respectively, in F) and hiPSC-derived AT2 cells (G) were analyzed and visualized as in B-D. **H**. Summary of findings in this work, its relationship to the observed clinical phases in COVID-19, and key aspects of modeling the same.

FIGURES

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

INVENTORY OF SUPPLEMENTARY MATERIALS

- **DETAILED METHODS**
- **SUPPLEMENTARY TABLES**
- **SUPPLEMENTARY FIGURES AND LEGENDS**
- **BIBLIOGRAPHY**

Detailed Methods

- **Key Resource Table**
- **Contact for Reagent and Resource Sharing**
- **Experimental Model and Subject Details**
  - Human lung organoids
  - Human iPSC(hiPSC)-derived Type II-alveolar pneumocytes
- **Method Details**
  - Human subjects for lung tissue source
  - Organoid isolation, expansion and culture
  - Preparation of monolayer
  - Preparation of air-liquid interface (ALI) monolayers
  - RNA Isolation
  - Quantitative (q)-PCR
  - Immunofluorescence
  - RNA Sequencing and data analysis
- **Quantification and Statistical Analysis**
  - Statistical Analysis
  - Replications
- **Data and Software Availability**

Key Resource Table:

MATERIALS & REAGENTS

ANTIBODIES USED FOR IMMUNOFLUORESCENCE

Name	Manufacturer	Catalog number	Dilution factor
ACE2	Santa Cruz	390851	1:100
ACE2	Biolegend	375802	1:50
Acetylated α-Tubulin	Santa Cruz	23950	1:500
AQP5	Santa Cruz	514022	1:100
CC10	Santa Cruz	365992	1:100
DAPI	Invitrogen	D1306	1:500
KRT5	Abcam	52635	1:100
MUC5AC	Abcam	229451	1:150
Na+/K+ ATPase	Abcam	76020	1:400
SARS-CoV Nucleoprotein	Sino Biological	40143-MM05	1:250
SFTPB	Santa Cruz	133143	1:100
SFTPC	Abcam	90716	1:150
anti-Rat IgG Secondary Antibody, Alexa Flour 594	Invitrogen	A-11007	1:500
anti-Rabbit IgG Secondary Antibody, Alexa Fluor 594	Invitrogen	A-11012	1:500
anti-Mouse IgG Secondary Antibody, Alexa Fluor 488	Invitrogen	A-11011	1:500

ANTIBODIES USED FOR IHC

SARS Spike (SARS-CoV2)	Abcam	273433	1:250
SARS-CoV Nucleoprotein	Sino Biological	40143-MM05	1:500

INSTRUMENTS

Countess II Automated Cell Counter	Thermo Fisher Scientific	AMQAX1000	
Epithelial Volt-Ohm (TEER) Meter	Millipore	MERS00002	
Leica DMI4000B (Automated Inverted Microscope)	Leica Microsystems	DMI4000B	
Leica TCS SPE Confocal	Leica Microsystems	TCS SPE	
Power Pressure Cooker XL	Tristar Products		
Canon Rebel XS DLSR	Canon		
MiniAmp Plus Thermal Cycler	Applied Biosystems	A37835	
QuantStudio5	Applied Biosystems	A28140	
Light Microscope (brightfield images)	Carl Zeiss LLC	Axio Observer, Inverted; 491917-0001-000	
Fisherbrand™ 150 Handheld Homogenizer	Fisher Scientific	15340168	

SOFTWARE

ImageJ	https://imagej.nih.gov/ij/index.html		
GraphPad Prism	https://www.graphpad.com/scientific-software/prism/		
LAS AF Software	https://www.leica-microsystems.com/products/microscope-software/p/leica-las-x-ls/		
QuantStudio Design & Analysis Software	https://www.thermofisher.com/us/en/home/global/forms/life-science/quantstudio-3-5-software.html		
CIBERSORTx	https://cibersortx.stanford.edu/runcibersortx.php		

ENZYMES, CHEMICALS, AND REAGENTS

Zinc Formalin	Fisher Scientific	23-313096	
Xylene	VWR	XX0060-4	
Hematoxylin	Sigma-Aldrich Inc	MHS1	
Ethanol	Koptec	UN1170	
Sodium Citrate	Sigma-Aldrich	W302600	
DAB (10x)	Thermo Fisher	1855920	1:10
Stable Peroxidase substrate buffer (10x)	Thermo Fisher	34062	1:10
3% Hydrogen Peroxide	Target	245-07-3628	
Horse Serum	Vector Labs	30022	
Anti- rabbit IgG	Vector Laboratories	MP-7401	
Paraformaldehyde 16% Solution, EM Grade	Electron Microscopy Sciences	15710	
100% Methanol	Supelco	MX0485	
Glycine	Fisher Scientific	BP381-5	
Bovine Serum Albumin	Sigma-Aldrich	A9647-100G	
Triton-X 100	Sigma-Aldrich	X100-500ML	
Prolong Glass	Invitrogen	P36984	
Nail Polish (Rapid Dry)	Electron Microscopy Sciences	72180	
Gill Modified Hematoxylin (Solution II)	Millipore Sigma	65066-85	
Histogel	Thermo Scientific	HG4000012	
TrypLE Select	Thermo Scientific	12563-011	
Advanced DMEM/F-12	Thermo Scientific	12634-010	
HEPES Buffer	Life Technologies	15630080	
Glutamax	Thermo Scientific	35050-061	
Penicillin-Streptomycin	Thermo Scientific	15140-122	
Collagenase Type I	Thermo Scientific	17100-017	
Matrigel	Corning	354234	
B-27	Thermo Scientific	17504044	
N-acetyl-L-cysteine	Sigma-Aldrich	A9165	
Nicotinamide	Sigma-Aldrich	N0636	
FGF-7 (KGF)	PeproTech	100-19-50ug	
FGF10	PeproTech	100-26-50ug	
A-83-01	Bio-Techne Sales Corp.	2939/50	
SB202190	Sigma-Aldrich	S7067-25MG	
Y-27632	R&D Systems	1254/50	
DPBS	Thermo Scientific	14190-144	

Ultrapure Water	Invitrogen	10977-015	
EDTA	Thermo Scientific	AM9260G	
Hydrocortisone	STEMCELL Technologies	7925	
Heparin	Sigma Aldrich	H3149	
Pneumacult Ex-Plus Medium	STEMCELL Technologies	5040	
PneumaCult ALI Medium	STEMCELL Technologies	5001	
Goat serum	Vector Laboratories	MP-7401	
Fetal Bovine Serum	Sigma-Aldrich	F2442-500ML	
Animal Component-Free Cell Dissociation Kit	STEMCELL Technologies	5426	
Red Blood Cell Lysis Buffer	Invitrogen	00-4333-57	
Cell Recovery Solution	Corning	354253	
Quick-RNA MicroPrep Kit	Zymo Research	R1051	
Quick-RNA MiniPrep Kit	Zymo Research	R1054	
Ethyl alcohol, pure	Sigma-Aldrich	E7023	
TRI Reagent	Zymo Research	R2050-1-200	
2x SYBR Green qPCR Master Mix	Bimake	B21203	
qScript cDNA SuperMix	Quanta Biosciences	95048	
Applied Biosystems TaqMan Fast Advanced Master Mix	Thermo Scientific	4444557	
18S, Hs99999901_s1	Thermo Scientific	4331182	
E_Sarbeco_F1 Forward Primer	IDT	10006888	
E_Sarbeco_R2 Reverse Primer	IDT	10006890	
E_Sarbeco_P1 Probe	IDT	10006892	

OTHER

6-well Tissue Culture Plate	Genesee Scientific	25-105	
12-well Tissue Culture Plate	CytoOne	CC7682-7512	
Transwell Inserts (6.5 mm, 0.4 um pore size)	Corning	3470	
Cell Scraper	Millipore Sigma	C5981-100EA	
Microscope Cover Glass (#1 Thickness) 24 x 50 mm	VWR	16004-098	
Microscope Cover Glass (#1 Thickness) 25 mm diameter	Chemglass Life Sciences	CLS-1760-025	
Millicell EZ Slide 8-Well Chamber	Millipore Sigma	PEZGS0816	
Optical Lens Cleaner	Fisher Scientific	22-143974	
Countess Cell Counting Chamber Slides	Invitrogen	C10312	
Trypan Blue Stain	Invitrogen	T10282	
70 um Cell Strainer	Thermo Fisher Scientific	22-363-548	
100 um Cell Strainer	Corning	352360	
RNase Away	Thermo Fisher Scientific	14-375-35	
Noyes Spring Scissors - Angled	Fine Science Tools	15013-12	

Detailed Methods

Collection of human lung specimens for organoid isolation

To generate adult healthy lung organoids, fresh biopsy bites were prospectively collected after surgical resection of the lung by the cardiothoracic surgeon. Before collection of the lung specimens, each tissue was sent to a gross anatomy room where a pathologist cataloged the area of focus, and the extra specimens were routed to the research lab in Human Transport Media (HTM, Advanced DMEM/F-12, 10 mM HEPES, 1X Glutamax, 1X penicillin-streptomycin, 5 μM Y-27632) for cell isolation. Deidentified lung tissues obtained during surgical resection, that were deemed excess by clinical pathologists, were collected using an approved human research protocol (IRB# 101590; PI: Thistlethwaite). Isolation and biobanking of organoids from these lung tissues were carried out using an approved human research protocol (IRB# 190105: PI Ghosh and Das) that covers human subject research at the UC San Diego HUMANOID Center of Research Excellence (CoRE). For all the deidentified human subjects, information including age, gender, and previous history of the disease, was collected from the chart following the rules of HIPAA and described in the Table.

A portion of the same lung tissue specimen was fixed in 10% Zinc-Formalin for at least 24hrs followed by submersion in 70% EtOH until embedding in FFPE blocks.

Autopsy procedures for lung tissue collection from COVID-19 positive human subjects

The lung specimens from COVID-19 positive human subjects were collected through autopsy (the study was IRB Exempt). All donations to this trial were obtained after telephone consent followed by written email confirmation by the next of kin/power of attorney per California state law (no in-person visitation could be allowed into our COVID-19 ICU during the pandemic). The team member followed the CDC guidelines for COVID19 and the autopsy procedures ((CAP), 2020; (CDC), 2020)). Lung specimens were collected in 10% Zinc-formalin and stored for 72 hrs before processing for histology. Patient characteristics are listed in the Table.

Autopsy #2 was a standard autopsy performed by anatomical pathology in the BSL3 autopsy suite. The patient expired and his family consented for autopsy. After 48 hrs, the lungs were removed and immersion fixed whole in 10% formalin for 48 hrs and then processed further. Lungs were only partially fixed at this time (about 50% fixed in thicker segments) and were sectioned into small 2-4 cm chunks and immersed in 10% formalin for further investigation.

Autopsy #4 and #5 were collected from rapid post-mortem lung biopsies. The procedure was performed in the Jacobs Medical Center ICU (all of the ICU rooms have a pressure-negative environment, with air exhausted through HEPA filters [Biosafety Level 3 (BSL3)] for isolation of SARS-CoV-2 virus). Biopsies were performed 2-4 hrs after patient expiration. The ventilator was shut off to reduce the aerosolization of viral particles at least 1 hr after the loss of pulse and before sample collection. Every team member had personal protective equipment in accordance with the University policies for procedures on patients with COVID-19 (N95 mask + surgical mask, hairnet, full face shield, surgical gowns, double surgical gloves, booties). Lung biopsies

were obtained after L-thoracotomy in the 5th intercostal space by the cardiothoracic surgery team. Samples were taken from the left upper lobe (LUL) and left lower lobe (LLL) and then sectioned further.

Isolation and culture of human whole lung-derived organoids

A previously published protocol was modified to isolate lung organoids from 3 human subjects(Sachs et al., 2019; Zhou et al., 2018). Briefly, normal human lung specimens were washed with PBS/4X penicillin-streptomycin and minced with surgical scissors. Tissue fragments were resuspended in 10 mL of wash buffer (Advanced DMEM/F-12, 10 mM HEPES, 1X Glutamax, 1X penicillin-streptomycin) containing 2 mg/ml Collagenase Type I (Thermo Fisher, USA) and incubated at 37°C for approximately 1 hr. During incubation, tissue pieces were sheared every 10 min with a 10 mL serological pipette and examined under a light microscope to monitor the progress of digestion. When 80-100% of single cells were released from connective tissue, the digestion buffer was neutralized with 10 mL wash buffer with added 2% Fetal Bovine Serum; the suspension was passed through a 100-μm cell strainer and centrifuged at 200 rcf. Remaining erythrocytes were lysed in 2 ml red blood cell lysis buffer (Invitrogen) at room temperature for 5 min, followed by the addition of 10 mL of wash buffer and centrifugation at 200 rcf. Cell pellets were resuspended in cold Matrigel (Corning, USA) and seeded in 25 μl droplets on a 12 well tissue culture plate. The plate was inverted and incubated at 37°C for 10 min to allow complete polymerization of the Matrigel before the addition of 1 mL Lung Expansion Medium per well. Lung expansion media was prepared by modifying the GI-organoid media (50% conditioned media, prepared from L-WRN cells with Wnt3a, R-spondin, and Noggin, ATCC-CRL-3276™) (Ghosh et al., 2020; Sayed et al., 2020a; Sayed et al., 2020b; Sayed et al., 2020c) with a proprietary cocktail from the HUMANOID CoRE containing B27, TGF-β receptor inhibitor, antioxidants, p38 MAPK inhibitor, FGF 7, FGF 10 and ROCK inhibitor. The lung expansion media was compared to alveolosphere media I (IMDM and F12 as the basal medium with B27, low concentration of KGF, Monothioglycerol, GSK3 inhibitor, Ascorbic acid, Dexamethasone, IBMX, cAMP and ROCK inhibitor) and II (F12 as the basal medium with added $CaCl_2$, B27, low concentration of KGF, GSK3 inhibitor, TGF-β receptor inhibitor Dexamethasone, IBMX, cAMP and ROCK inhibitor) modified from previously published literature (Jacob et al., 2019; Yamamoto et al., 2017). Neither alvelosphere media contain any added Wnt3a, R-spondin, and Noggin. The composition of these media was developed either by fundamentals of adult-stem cell-derived mixed epithelial cellularity in other organs (like the gastrointestinal tract (Miyoshi and Stappenbeck, 2013; Sato et al., 2009; Sayed et al., 2020c), or rationalized based on published growth conditions for proximal and distal airway components(Gotoh et al., 2014; Sachs et al., 2019; van der Vaart and Clevers, 2020). Organoids were maintained in a humidified incubator at 37°C/5% CO2, with a complete media change performed every 3 days. After the organoids reached confluency between 7-10 days, organoids were collected in PBS/0.5 mM EDTA and centrifuged at 200 rcf for 5 min. Organoids were dissociated in 1 mL trypLE Select (Gibco, USA) per well at 37°C for 4-5 min and mechanically sheared. Wash buffer was added at a 1:5, trypLE to wash buffer ratio. The cell suspension was subsequently centrifuged, resuspended in Matrigel, and seeded at a 1:5 ratio. Lung organoids were biobanked and passage 3-8 cells were used for experiments. Subculture was performed every 7-10 days.

The preparation of Lung organoid-derived monolayers

Lung-organoid-derived monolayers were prepared using a modified protocol of GI-organoid-derived monolayers (Ghosh et al., 2020; Sayed et al., 2020a; Sayed et al., 2020b; Sayed et al., 2020c). Briefly, transwell inserts (6.5 mm diameter, 0.4 um pore size, Corning) were coated in Matrigel diluted in cold PBS at a 1:40 ratio and incubated for 1hr at room temperature. Confluent organoids were collected in PBS/EDTA on day 7 and dissociated into single cells in trypLE for 6-7 min at 37°C. Following enzymatic digestion, the cell suspension was mechanically sheared through vigorous pipetting with a 1000 μl pipette and neutralized with wash buffer. The suspension was centrifuged, resuspended in Pneumacult Ex-Plus Medium (StemCell, Canada), and passed through a 70-μm cell strainer. The coating solution was aspirated, and cells were seeded onto the apical membrane at 1.8E5 cells per transwell with 200 μl PneumaCult Ex-Plus media. 700 μl of PneumaCult Ex-Plus was added to the basal chamber. Cells were cultured over the course of 2-4 days. A media change of both the apical and basal chambers was performed every 24 hrs.

Air-Liquid Interface (ALI) Model of Lung organoids

Organoids were dissociated into single cells and expanded in T-75 flasks in PneumaCult Ex-Plus Medium until confluency was reached. Cells were dissociated in ACF Enzymatic Dissociation Solution (StemCell, Canada) for 6-7 min at 37°C and neutralized in equal volume ACF Enzyme Inhibition Solution (StemCell, Canada). Cells were seeded in the apical chamber of transwells at 3.3E4 cells per transwell in 200 μL of PneumaCult Ex-Plus Medium. 500 μL of PneumaCult Ex-Plus was added to the basal chamber. Media in both chambers was changed every other day until confluency was reached (~4 days). The media was completely removed from the apical chamber, and media in the basal chamber was replaced with ALI Maintenance Medium (StemCell, Canada). The media in the basal chamber was changed every 2 days. The apical chamber was washed with warm PBS every 5-7 days to remove accumulated mucus. Cells were cultured under ALI conditions for 21+ days until they completed differentiation into a pseudostratified mucociliary epithelium. To assess the integrity of the epithelial barrier, Trans-Epithelial Electrical Resistance (TEER) was measured with an Epithelial Volt-Ohm Meter (Millicell, USA). The media was removed from the basal chamber, and wash media was added to both chambers. Cultures were equilibrated to 37°C before TEER values were measured. Final values were expressed as $\Omega \cdot cm^2$ units and were calculated by multiplying the growth area of the membrane by the raw TEER value.

The culture of primary airway epithelial cells and iPSC-derived alveolar epithelial cells

Primary normal human bronchial epithelial cells (NHBE) were obtained from Lonza and grown according to instructions. NHBE cells were cultured in T25 cell culture tissue flasks with PneumaCult-Ex Plus media (StemCell, Canada). Cells were seeded at ~100,000 cells/T25 flask and incubated at 37°C, 5% CO_2. Once cells reached 70–80% confluency, they were dissociated using 0.25% Trypsin in dissociation media and plated in 24 well transwells (Corning). Primary human bronchial epithelial cells (HBEpC) and small airway epithelial cells (HSAEpC) were obtained from Cell Applications Inc. The HBEpC and HSAEpC were cultured in human bronchial/tracheal epithelial cell media and small airway epithelial cell media, respectively, following the instructions of Cell Application.

Human iPSC-derived alveolar epithelial type 2 cells (iHAEpC2) were obtained from Cell Applications Inc. and cultured in growth media (i536K-05, Cell Applications Inc.) according to the manufacturer's instructions. All the primary cells were used within early passages (5-6) to avoid any gradual disintegration of the airway epithelium with columnar epithelial structure and epithelial integrity.

The infection with SARS-Cov2

Lung organoid-derived monolayers or primary airway epithelial cells either in 384 well plates or in transwells were washed twice with antibiotic-free lung wash media. 1E5 PFU of SARS-CoV-2 strain USA-WA1/2020 (BEI Resources NR-52281) in complete DMEM was added to the apical side of the transwell and allowed to incubate for 24, 48, 72 and 96 hrs at 34°C and 5% CO_2. After incubation, the media was removed from the basal side of the transwell. The apical side of the transwells was then washed twice with (antibiotic-free lung wash media) and then twice with PBS. TRIzol™ Reagent (Thermo Fisher 15596026) was added to the well and incubated at 34˚C and 5% CO_2 for 10 min. The TRIzol™ Reagent was removed and stored at -80 ˚C for RNA analysis.

RNA isolation

Organoids and monolayers used for lung cell type studies were lysed using RNA lysis buffer followed by RNA extraction per Zymo Research Quick-RNA MicroPrep Kit instructions. Tissue samples and monolayers in SARS-CoV2 studies were lysed in TRI-Reagent and RNA was extracted using Zymo Research Direct-zol RNA Miniprep.

Quantitative (q)RT-PCR

Organoid and monolayer cell-type gene expression was measured by qRT-PCR using 2x SYBR Green qPCR Master Mix. cDNA was amplified with gene-specific primer/probe set for the lung cell type markers and qScript cDNA SuperMix (5x). qRT-PCR was performed with the Applied Biosystems QuantStudio 5 Real-Time PCR System. Cycling parameters were as follows: 95 °C for 20 s, followed by 40 cycles of 1 s at 95 °C and 20 s at 60 °C. All samples were assayed in triplicate and eukaryotic 18S ribosomal RNA was used as a reference.

Cell types	Marker	Primer Sequence
Basal cells	ITGA6	ITGA6 F 'CGAAACCAAGGTTCTGAGCCC' ITGA6 R 'CTTGGATCTCCACTGAGGCAGT'
Goblet	Muc5AC	Muc5AC F ' GGAACTGTGGGGACAGCTCTT' Muc5AC R ' GTCACATTCCTCAGCGAGGTC'
Cilia	FoxJ1	FoxJ1 F 'ACTCGTATGCCACGCTCATCTG' FoxJ1 R 'GAGACAGGTTGTGGCGGATTGA'
Club cell	SCGB1A1	SCGB1A1 F ' CAAAAGCCCAGAGAAAGCATC' SCGB1A1 R ' CAGTTGGGGATCTTCAGCTTC'
Alveolar type 1	AQP5, PDPN, P2RX4	AQP5 F ' TACGGTGTGGCACCGCTCAATG' AQP5 R ' AGTCAGTGGAGGCGAAGATGCA' PDPN F ' GTGCCGAAGATGATGTGGTGAC' PDPN R ' GGACTGTGCTTTCTGAAGTTGGC' P2RX4 F ' GTGGCGGATTATGTGATACCAGC' P2RX4 R ' CACACAGTGGTCGCATCTGGAA'
Alveolar type 2	ABCA3, SFTPA1, SFTPC, LAMP3	ABCA3 F ' CTTGACAGTCGCAGAGCACCTT' ABCA3 R ' CTCCGTGAGTTCCACTTGTCCT' SFTPA1 F ' CACCTGGAGAAATGCCATGTCC' SFTPA1 R ' AAGTCGTGGAGTGTGGCTTGGA' SFTPC F ' GTCCTCATCGTCGTGGTGATTG' SFTPC R ' AGAAGGTGGCAGTGGTAACCAG' LAMP3 F ' TGGGAGCCTATTTGACCGTCTC' LAMP3 R ' GCTGACAACTGGAGGCTCTGTT'

Assessment of SARS-CoV-2 infectivity test

Assessment of SARS-CoV-2 infectivity test was determined by qPCR using TaqMan assays and TaqMan Universal PCR Master Mix as done before(Corman et al., 2020; Lamers et al., 2020). cDNA was amplified with gene-specific primer/probe set for the E gene and QPCR was performed with the Applied Biosystems QuantStudio 3 Real-Time PCR System. The specific TaqMan primer/probe set for E gene are as follows: Forward 5'-ACAGGTACGTTAATAGTTAATAGCGT-3', (IDT, Cat# 10006888); Reverse 5'-ATATTGCAGCAGTACGCACACA-3'; Probe 5'-FAM-ACACTAGCCATCCTTACTGCGCTTCG-BBQ-3' and 18S rRNA. Cycling parameters were as follows: 95 °C for 20 s, followed by 40 cycles of 1 s at 95 °C and 20 s at 60 °C. All samples were assayed in triplicate and gene eukaryotic 18S ribosomal RNA was used as a reference.

Immunofluorescence

Organoids and Lung organoid-Derived Monolayers were fixed by either: (1) 4% PFA at room temperature for 30 min and quenched with 30 mM glycine for 5 min, (2) ice-cold 100% Methanol at -20°C for 20 min, (3) ice-cold 100% Methanol on ice for 20 min. Subsequently, samples were permeabilized and blocked for 2 hrs using an in-house blocking buffer (2 mg/mL BSA and 0.1% Triton X-100 in PBS); as described previously(Lopez-Sanchez et al., 2014). Primary antibodies were diluted in blocking buffer and allowed to incubate overnight at 4°C; Secondary antibodies were diluted in blocking buffer and allowed to incubate for 2 hrs in the dark. Antibody dilutions are listed in the Supplementary Key Resource Table. ProLong Glass was used as a mounting medium. #1 Thick Coverslips were applied to slides and sealed. Samples were stored at 4°C until imaged. FFPE embedded Organoid and Lung Tissue sections underwent antigen retrieval as previously described in methods for Immunohistochemistry staining. After antigen retrieval and cooling in DI water; samples were permeabilized and blocked in blocking buffer and treated as mentioned above for immunofluorescence. Images were acquired at room temperature with Leica TCS SPE confocal and with DMI4000 B microscope using the Leica LAS-AF Software. Images were taken with a 40× oil-immersion objective using 405-,488-, 561-nm laser lines for excitation. Z-stack images were acquired by successive Z-slices of 1μm in the desired confocal channels. Fields of view that were representative and/or of interest were determined by randomly imaging 3 different fields. Z-slices of a Z-stack were overlaid to create maximum intensity projection images; all images were processed using FIJI (Image J) software.

Embedding of Organoids in *HistoGel*

Organoids were seeded on a layer of Matrigel in 6-Well plates and grown for 7-8 days. Once mature, organoids were fixed in 4% PFA at room temperature for 30 min and quenched with 30 mM glycine for 5 min. Organoids were gently washed with PBS and harvested using a cell scraper. Organoids were resuspended in PBS using wide-bore 1000 μL pipette tips. Organoids were stained using Gill's hematoxylin for 5 min for easier FFPE embedding and sectioning visualization. Hematoxylin stained organoids were gently washed in PBS and centrifuged and excess hematoxylin was aspirated. Organoids were resuspended in 65°C histogel and centrifuged at 65°C for 5 min. Histogel embedded organoid pellets were allowed to cool to room temperature

and stored in 70% ethanol at 4°C until ready for FFPE embedding by LJI Histology Core. Successive FFPE embedded organoid sections were cut at a 4 µm thickness and fixed on to microscope slides.

Immunohistochemistry

For SARS CoV- nucleoprotein (np) antigen retrieval, slides were immersed in Tris-EDTA buffer (pH 9.0) and boiled for 10 min at 100°C inside a pressure cooker. Endogenous peroxidase activity was blocked by incubation with 3% H_2O_2 for 10 minutes. To block non-specific protein binding 2.5% goat serum was added. Tissues were then incubated with a rabbit SARS CoV-NP antibody (Sino Biological, See Supplementary Key Resource Table) for 1.5 hrs at room temperature in a humidified chamber and then rinsed with TBS or PBS 3 times, for 5 min each. Sections were incubated with horse anti-rabbit IgG secondary antibodies for 30 min at room temperature and then washed with TBS or PBS 3 times for 5 min each. Sections were incubated with DAB and counterstained with hematoxylin for 30 sec.

RNA Sequencing

RNA sequencing libraries were generated using the Illumina TruSeq Stranded Total RNA Library Prep Gold with TruSeq Unique Dual Indexes (Illumina, San Diego, CA). Samples were processed following manufacturer's instructions, except modifying RNA shear time to five minutes. The resulting libraries were multiplexed and sequenced with 100 basepair (bp) Paired-End (PE100) to a depth of approximately 25-40 million reads per sample on an Illumina NovaSeq 6000 by the Institute of Genomic Medicine (IGM) at the University of California San Diego. Samples were demultiplexed using bcl2fastq v2.20 Conversion Software (Illumina, San Diego, CA). RNASeq data was processed using kallisto (version 0.45.0), and human genome GRCh38 Ensembl version 94 annotation (Homo_sapiens GRCh38.94 chr_patch_hapl_scaff.gtf). Gene-level TPM values and gene annotations were computed using tximport and biomaRt R package. A custom python script was used to organize the data and log reduced using log2(TPM+1). The raw data and processed data are deposited in Gene Expression Omnibus under accession no GSE157055, and GSE157057.

Data Collection and Annotation

Publicly available COVID-19 gene expression databases were downloaded from the National Center for Biotechnology Information (NCBI) Gene Expression Omnibus website (GEO) (Barrett et al., 2005; Barrett et al., 2013; Edgar et al., 2002). If the dataset is not normalized, RMA (Robust Multichip Average)(Irizarry et al., 2003a; Irizarry et al., 2003b) is used for microarrays and TPM (Transcripts Per Millions)(Li and Dewey, 2011; Pachter, 2011) is used for RNASeq data for normalization. We used log2(TPM+1) to compute the final log-reduced expression values for RNASeq data. Accession numbers for these crowdsourced datasets are provided in the figures and manuscript. All of the above datasets were processed using the Hegemon data analysis framework (Dalerba et al., 2011; Dalerba et al., 2016; Volkmer et al., 2012).

Analysis of RNA seq Datasets

DESeq2 (Love et al., 2014) was applied to uninfected and infected samples to identify Up- and Down-regulated genes. A gene signature score is computed using both the Up- and Down- regulated genes which are used to order the sample. To compute the gene signature score, first, the genes present in this list were normalized according to a modified Z-score approach centered around StepMiner threshold (formula = (expr - SThr)/3*stddev). The normalized expression values for every probeset for all the genes were added or subtracted (depending on Up and Down-regulated genes) together to create the final score. The samples were ordered based on the final gene signature score. The Gene signature score is used to classify sample categories and the performance of the multi-class classification is measured by ROC-AUC (Receiver Operating Characteristics Area Under The Curve) values. A color-coded bar plot is combined with a violin plot to visualize the gene signature-based classification. All statistical tests were performed using R version 3.2.3 (2015-12-10). Standard t-tests were performed using python scipy.stats.ttest_ind package (version 0.19.0) with Welch's Two Sample t-test (unpaired, unequal variance (equal_var=False), and unequal sample size) parameters. Multiple hypothesis correction was performed by adjusting *p* values with statsmodels.stats.multitest.multipletests (fdr_bh: Benjamini/Hochberg principles). The results were independently validated with R statistical software (R version 3.6.1; 2019-07-05). Pathway analysis of gene lists was carried out via the Reactome database and algorithm (Fabregat et al., 2018). Reactome identifies signaling and metabolic molecules and organizes their relations into biological pathways and processes. Violin, Swarm and Bubble plots are created using python seaborn package version 0.10.1.

Single Cell RNA Seq data analysis

Single Cell RNASeq data from GSE145926 was downloaded from GEO in the HDF5 Feature Barcode Matrix Format. The filtered barcode data matrix was processed using Seurat v3 R package(Stuart et al., 2019) and scanpy python package(Wolf et al., 2018). Pseudo bulk analysis of GSE145926 data was performed by adding counts from the different cell subtypes and normalized using log2(CPM+1). Epithelial cells were identified using SFTPA1, SFTPB, AGER, AQP4, SFTPC, SCGB3A2, KRT5, CYP2F1, CCDC153, and TPPP3 genes using SCINA algorithm(Zhang et al., 2019). Pseudo bulk datasets were prepared by adding counts from the selected cells and normalized using log (CPM+1).

Assessment of cell-type proportions

CIBERSORTx (https://cibersortx.stanford.edu/runcibersortx.php) was used for cell-type deconvolution of our dataset (which was normalized by CPM). As reference samples, we first used the Single Cell RNASeq dataset (GSE132914) from Gene Expression Omnibus (GEO). Next, we analyzed the bulk RNA seq datasets for the identification of cell types of interest using relevant gene markers (see **Table 2**): AT1 cells (PDPN, AQP5, P2RX4, TIMP3, SERPINE1), AT2 cells (SFTPA1, SFTPB, SFTPC, SFTPD, SCGB1A1, ABCA3, LAMP3), BASAL cells (CD44, KRT5, KRT13, KRT14, CKAP4, NGFR, ITGA6), CLUB cells (SCGB1A1, SCGB3A2, SFTPA1, SFTPB, SFTPD, ITGA6, CYP2F1), GOBLET cells (CDX2, MUC5AC, MUC5B, TFF3), Ciliated cells (ACTG2, TUBB4A, FOXA3, FOXJ1, SNTN), Generic Lung Lineage cells (GJA1, TTF1, EPCAM) and cells for Viral Entry Markers

(ACE2,TMPRSS2) were identified using SCINA algorithm. Then, normalized pseudo counts were obtained with the CPM normalization method. The cell-type signature matrix was derived from the Single Cell RNASeq dataset, cell-types, and gene markers of interest. It was constructed by taking an average from gene expression levels for each of the markers across each cell type.

Statistical analysis

All experiments were repeated at least three times, and results were presented either as one representative experiment or as average ± S.E.M. Statistical significance between datasets with three or more experimental groups was determined using one-way ANOVA including a Tukey's test for multiple comparisons. For all tests, a P-value of 0.05 was used as the cutoff to determine significance (*P < 0.05, **P < 0.01, ***P < 0.001, and ****P < 0.0001). All experiments were repeated a least three times, and P-values are indicated in each figure. All statistical analyses were performed using GraphPad prism 6.1. A part of the statistical tests was performed using R version 3.2.3 (2015-12-10). Standard t-tests were performed using python scipy.stats.ttest_ind package (version 0.19.0).

Table 1. A comparison of current versus existing lung organoid models available for modeling COVID-19

Author	Source of stem cells	Propagability	Cell types						SARS-COV-2 infection	Demonstrated reproducibility using more than one patient	Cost-effective (use of conditioned media)	Notes
			AT1	AT2	Club	Basal	Ciliated	Goblet				
Zhou et al PMID: 29891677	Small pieces of normal lung tissue adjacent to the diseased tissue from patients undergoing surgical resection for clinical conditions.	✓ Long term culture >1 y	✗	✗	✓	✓	✓	✓	✗ Infection with H1N1 pandemic Influenza virus	—	—	Proximal differentiation (PD) of human Adult Stem Cell-derived airway organoid (AO) culture. Differentiation conditions (PneumaCult-ALI medium) increase ciliated cells. Serine proteases known to be important for productive viral infection, were elevated after PD.
Sachs et al PMID: 30643021	Generation of normal and tumor organoids from resected surplus lung tissue of patients with lung cancers.	✓ long term culture for over 1 year	✗	✗	✓	✓	✓	— Not clearly mentioned	✗	—	—	airway organoid (AO) expressed **no mesenchyme or alveolar** transcripts. Strongly enriched for bulk lung and small airway epithelial signature limited to basal, club, and ciliated cells. Withdrawal of R-spondin terminated AO expansion after 3–4 passages similar to withdrawal of FGFs
Duan et al PMID: 32839764	hPSC derived lung cells and macrophages	✗	✓	✓	✓	✓	✓	✓ Low	✓ SARS-CoV-2 infection mediated damage onset by macrophages.	—	—	Co-culture of lung cells and macrophages. Protocol followed enables alveolar differentiation process, although described presence of almost all lung cell types.
Salahudeen et al BioRxiv https://doi.org/1 0.1101/2020.07. 27.212076	Cells sorted from human peripheral lung tissues.	— Distal Lung organoid with possibility of long-term culture	From diff of AT2	✓	After diff of basal cells	✓		—	✓ Infection and presence of dsRNA and nucleocapsid	—	—	No RNA seq of infected samples to compare with COVID. Differentiation to different cell types. SARS CoV2 infection in apical-out organoids (not polarized monolayers). The combination of EGF and the Noggin was optimal, without any additional growth-promoting effects of either WNT3A or R-spondin
Han et al BioRxiv doi: https://doi.o rg/10.1101/202 0.05.05.079095	hPSC-derived lung organoids	— Organoids were generated by 50 days of differentiation	✓	✓	—	—	—	—	✓ SARS-CoV-2 and SARS-CoV-2-Pseudo-Entry Viruses.	—	—	AT1, AT2, stromal cells, low number of pulmonary neuroendocrine cells, proliferating cells, and airway epithelial cells were reported. Mostly AT2 based ACE2 receptor was used for virus infection. High throughput screen using hPSC-derived lung organoids identified FDA-approved drug candidates, including imatinib and mycophenolic acid, as inhibitors of SARS-CoV-2 entry.
Youk et al BioRxiv doi: https://doi.org/1 0.1101/2020.07 .10.194498	Adult alveolar stem cells isolated from distal lung parenchymal tissues by collagenase, dispase and sorting	— Multiple passages upto 10 months	From AT2 Lost in higher passages	✓	✗	✗	✗	✗	✓ In the organoid form	—	—	Single cell transcriptomic profiling identified 2 clusters and type I interferon signal pathway are highly elevated at 3 dpi
Mulay et al doi.org/10.1101 /2020.06.29.17 4623	a) Alv organoids with distal lung epithelial cells with lung fibroblast cells	—	✗	✓	✗	✗	✗	✗	✓ In the organoid form	✓	—	Infection of AT2 cells trigger apoptosis that may contribute to alveolar injury. Alteration of innate immune response genes from AT2 cells
	b) Proximal airway ALI with heterogenous cells	—	✗	✗	✓	✓	✓	✓	✓	✓	—	Infection of ciliated and goblet cells. Two separate models for SARS-CoV2 infection
Huang J PMID: 32979316	iPSC derived AT2 cell ALI model	—	✓	✓	✗	✗	✗	✗	✓	—	—	Bulk RNA seq after day 1 and day 4 infection. The infection induces rapid inflammatory responses.
Tindle et al [Current study]	Deep lung tissue sections surgically obtained from patients undergoing lobe resections for lung cancers.	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	RNA Seq and cross-validation of COVID-19 model. Single model with all the cells types and infection of SARS-CoV2 in the 2D form with Apical accessibility that close to physiologic state.

Table 2: Markers used to identify various cell types in the lung

CELL TYPE	MARKERS
AT1	AQP5*, PDPN*, Carboxypeptidase M, CAV-1, CAV-2, HTI56, HOPX, P2RX4*, Na+/K+ ATPase, TIMP3*, SEMA3F PDPN* AQP5* P2RX4* TIMP3* SERPINE*
AT2	ABCA3*, CC10 (SCGB1A1*), CD44v6, Cx32, gp600, ICAM-1, KL-6, LAMP3*, MUC1, SFTPA1*, SFTPB*, SFTPC*, SFTPD*, SERPINE1
CLUB	CC10 (SCGB1A1*), CYP2F2*, ITAG6*, SCGB3A2*, SFTPA1*, SFTPB*, SFTPD*
GOBLET	CDX-2*, MUC5AC*, MUC5B*, TFF3*, UEA1+
CILIATED	ACT (ACTG2*), BTub4 (TUBB4A*), FOXA3*, FOXJ1*, SNTN*
BASAL	CD44v6 (CD44*), ITGA6*, KRT5*, KRT13*, KRT14*, p63 (CKAP4*), p75 (NGFR*)
GENERIC LUNG LINEAGE	Cx43 (GJA1*), TTF-1 (TTF1*; Greatest in AT2 & Club), EpCAM (EPCAM*)
VIRAL ENTRY MARKERS	ACE2*, TMPRSS2*

Asterisk (*) denotes markers used for single cell gating (**Fig 1A**)
Marker used in this work for Immunofluorescence (IF)
Marker used in this work for qPCR
Markers used in both IF and qPCR
Obscure markers (Not a lot of research relative to lung)

Table 3. Characteristics of patients enrolled into this study for obtaining lung tissues to serve as source of stem cells to generate lung organoids

Name	Date of surgery	Age	Sex	Smoking History	Reason for surgery	Histology
ALO1	4/17/2020	64	Male	Current, chronic smoker. Packs/day: 0.50. Years: 53. Pack years: 26.5	Lung Carcinoma	Invasive Squamous Cell Carcinoma, non-keratinizing
ALO2	4/17/2020	59	Male	Non-smoker	Lung Carcinoma	Invasive Adenocarcinoma
ALO3	7/7/2020	46	Female	Non-smoker	Left Lower Lobe Nodule	Invasive Adenocarcinoma

Table 4: Upregulated genes and pathways: Healthy vs COVID-19 lung ([GSE151764](GSE151764))

Genes:

BRCA2	XAGE1B	CDK1	SNAI2	CXCL11
CYBB	CCR5	GBP1	IFITM1	IFI27
KRT5	CCR2	HLA-G	GZMB	IFI35
C1QB	ALOX15B	IDO1	CD163	TDO2
FCGR1A	CMKLR1	ISG20	CD38	GZMA
IL10	MX1	LAG3	BST2	OAS3
IL6	TNFRSF17	MAD2L1	BUB1	POU2AF1
CD44	CCR1	CXCL9	CCL20	CXCL13
CD276	CXCR3	MKI67	CCNB2	GNLY
DMBT1	SLAMF8	IFIT2	TNFSF18	IFIT3
DDX58	IL21	IFIT1	ISG15	TOP2A
TNFAIP8	FOXM1	CXCL10	CDKN3	LILRB1
LAMP3	IFIH1	IRF4	C1QA	HERC6
KIAA0101	IFI6	PSMB9	OAS1	TNFSF13B
MELK	PDCD1LG2	CCL18	OAS2	IFI44L
				STAT1

Pathways:

Name	pValue	FDR
Interferon Signaling	1.1102230246251565e-16	1.1102230246251565e-14
Interferon alpha/beta signaling	1.1102230246251565e-16	1.1102230246251565e-14
Cytokine Signaling in Immune system	1.1102230246251565e-16	1.1102230246251565e-14
Immune System	1.1102230246251565e-16	1.1102230246251565e-14
Interleukin-10 signaling	9.845457782375888e-13	7.87636622590071e-11
Interferon gamma signaling	9.256706512417168e-12	6.109426298195331e-10
Chemokine receptors bind chemokines	1.0827139185209944e-10	6.171469335569668e-9
Signaling by Interleukins	6.813459352272844e-9	3.406729676136422e-7
Insulin-like Growth Factor-2 mRNA Binding Proteins (IGF2BPs/IMPs/VICKZs) bind RNA	1.268436958801189e-7	0.00000055811226187252320
Antiviral mechanism by IFN-stimulated genes	0.00000019330583493903393	0.00007732233397561572
CD163 mediating an anti-inflammatory response	0.000007798676168513374	0.0002807523420664815
OAS antiviral response	0.000010208709973369423	0.00033688742912119096
Peptide ligand-binding receptors	0.000017140576871854662	0.0005142173061556399
Interleukin-4 and Interleukin-13 signaling	0.00010149486613275638	0.0028418562517171786
Cyclin A/B1/B2 associated events during G2/M transition	0.00018878164653901575	0.004908322810014409
G0 and Early G1	0.00036071218383249004	0.009017804595812251
Interleukin-6 signaling	0.0004656678443889506	0.010710360420945864
ISG15 antiviral mechanism	0.0008313991987842773	0.017459383174469822
Regulation of APC/C activators between G1/S and early anaphase	0.0008313991987842773	0.017459383174469822
Polo-like kinase mediated events	0.0011105065129394243	0.022210130258788485
APC/C-mediated degradation of cell cycle proteins	0.0013081035810655894	0.02354586445918061
Regulation of mitotic cell cycle	0.0013081035810655894	0.02354586445918061
G2/M DNA replication checkpoint	0.001750156331829622	0.029752657641103575
Class A/1 (Rhodopsin-like receptors)	0.0023550630454979293	0.035376667816378604
Interleukin-6 family signaling	0.002358444521091907	0.035376667816378604
TNFs bind their physiological receptors	0.002358444521091907	0.035376667816378604

Table 5: Downregulated genes and pathways: Healthy vs COVID-19 lung (GSE151764)

Genes:

CX3CR1	JAML	KLRB1	GRAP2	CD226
ARG1	CX3CR1	LY9	MMP9	CD160
MPO	HLA-DQB2	CCL17	RORC	FOXP3
IL2	TNFRSF9	CCL22	CCR4	CRTAM
BCL2	CXCR5	TCF7	IRS1	CCR6
CA4	CD1C	CXCR4	ITK	CEACAM8
IGF1R	CD69	CD83	KLRG1	PTGS2

Pathways:

Name	pValue	FDR
Chemokine receptors bind chemokines	2.8463564838432376e-11	4.981123846725666e-9
Immune System	1.2524759007703778e-10	1.0896540336702287e-8
Interleukin-4 and Interleukin-13 signaling	2.820398359304477e-9	1.6358310483965965e-7
RUNX1 and FOXP3 control the development of regulatory T lymphocytes (Tregs)	4.3109720904155324e-7	0.0000185371799887679
Peptide ligand-binding receptors	6.709444980712576e-7	0.00002348305743249014
Signaling by Interleukins	0.0000015036584932737185	0.000043606096304937836
Cytokine Signaling in Immune system	0.000026065058545632347	0.0006516264636408087
Dectin-1 mediated noncanonical NF-kB signaling	0.00008640543214522012	0.0018145140750496225
Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell	0.00010833886747929622	0.002058438482106628
Class A/1 (Rhodopsin-like receptors)	0.0001833048828232542	0.0031161830079953212
Interleukin-10 signaling	0.0002366961933546463	0.0035504429003196947
RUNX3 Regulates Immune Response and Cell Migration	0.0005791814112650062	0.0077471849338778265
Extra-nuclear estrogen signaling	0.0005959373026059867	0.0077471849338778265
BH3-only proteins associate with and inactivate anti-apoptotic BCL-2 members	0.000699254752329681	0.008391057027956172
CLEC7A (Dectin-1) signaling	0.000822803514531345	0.009050838659844795
Generation of second messenger molecules	0.0011719919081142338	0.011719919081142338
Innate Immune System	0.0016764040917021106	0.015723603669013264
GPCR ligand binding	0.0017470670743348071	0.015723603669013264
Adaptive Immune System	0.002059835990545711	0.018538523914911398
Estrogen-dependent nuclear events downstream of ESR-membrane signaling	0.004670055830237274	0.03736044664189819
C-type lectin receptors (CLRs)	0.005458044949850915	0.04366435959880732
Transcriptional regulation by RUNX3	0.008124332598870021	0.05687032819209015
BMAL1:CLOCK,NPAS2 activates circadian gene expression	0.009518272708583853	0.06662790896008697
ESR-mediated signaling	0.012073762373888686	0.0845163366172208
Transcriptional regulation by RUNX1	0.012881563708187715	0.08786708746501692
TCR signaling	0.01464451457750282	0.08786708746501692

Table 6: Upregulated genes and pathways: Uninfected vs Infected (48 hpi) lung organoid monolayers

Genes:

IFI35	EPSTI1	AMIGO2	IFITM2
SLC4A11	CMPK2	WARS1	FAAP100
APOL1	OASL	IFI27	ISG15
OAS3	IFI44L	CD14	SLC35F6
IFIT3	IFI44	SAMD9L	
IFIT2	PARP9	SRP9P1	

Pathways:

Name	pValue	FDR
Interferon Signaling	1.1102230246251565e-16	4.218847493575595e-15
Interferon alpha/beta signaling	1.1102230246251565e-16	4.218847493575595e-15
Cytokine Signaling in Immune system	1.1540801647669241e-10	2.8852004119173102e-9
Immune System	0.0000025401148793235596	0.00004826218270714833
OAS antiviral response	0.0004764545663257236	0.007146818494885854
Antiviral mechanism by IFN-stimulated genes	0.0010333472605504879	0.012400167126605854
Interferon gamma signaling	0.0018896946191467867	0.020786640810614654
Transfer of LPS from LBP carrier to CD14	0.006318772245420878	0.056868950208787905
TRIF-mediated programmed cell death	0.020912675855586538	0.16560733288455465
MyD88 deficiency (TLR2/4)	0.03733748270572135	0.16560733288455465
IRAK2 mediated activation of TAK1 complex upon TLR7/8 or 9 stimulation	0.03733748270572135	0.16560733288455465
TRAF6-mediated induction of TAK1 complex within TLR4 complex	0.03937173812023953	0.16560733288455465
IRAK4 deficiency (TLR2/4)	0.03937173812023953	0.16560733288455465
Activation of IRF3/IRF7 mediated by TBK1/IKK epsilon	0.04140183322113866	0.16560733288455465
Caspase activation via Death Receptors in the presence of ligand	0.04140183322113866	0.16560733288455465
IKK complex recruitment mediated by RIP1	0.04948077475965562	0.18550132647663609

Table 7: Downregulated genes and pathways: Uninfected vs Infected (48 hpi) lung organoid monolayers

AC093392.1	ARHGAP19	HLA-V	RN7SL718P
MT-TV	AC138969.3	AC016766.1	

Pathways:

Name	pValue	FDR
rRNA processing in the mitochondrion	0.01892731245535506	0.08366120772817287
tRNA processing in the mitochondrion	0.021271491049123403	0.08366120772817287
Mitochondrial translation termination	0.043991554455812154	0.08366120772817287
Mitochondrial translation elongation	0.043991554455812154	0.08366120772817287
Mitochondrial translation initiation	0.04490921761868061	0.08366120772817287
Mitochondrial translation	0.04765767844112356	0.08366120772817287

SUPPLEMENTARY FIGURES AND LEGENDS

Supplementary Figure 1. Alveolar type II pneumocyte hyperplasia is a pathognomonic feature of lung injury in COVID-19.
A. Whisker plots display relative levels of ACE2 expression in various cell types in the normal human lung. The cell types were annotated within a publicly available single-cell sequencing dataset (GSE132914) using genes listed in Table S1. *p*-values were analyzed by one-way Anova and Tukey post hoc test.
B. FFPE sections of the human lung from deceased COVID-19 patients were analyzed by H&E staining. Representative fields are shown. Images on the right are magnified areas indicated with boxes on the left. Arrows = alveolar type II pneumocyte hyperplasia.
C-D. FFPE sections of the human lung from normal and deceased COVID-19 patients were stained for AT2 and club cell markers and either ACE2 or viral nucleocapsid protein and analyzed by confocal immunofluorescence. Representative images are shown. Scale bar = 50 μm.
E. FFPE sections of the human lung from normal and deceased COVID-19 patients were stained for viral nucleocapsid antibody. Representative images are shown. Arrows = infected cells.

Supplementary Figure 2. Lung organoids are reproducibly established from 3 different donors and propagated in each case over 10 passages.
A. Schematic displaying the key demographics of the patients who served as donors of the lung tissue as a source of adult stem cells for the generation of organoids. Three organoid lines were generated, ALO1-3. ALO, adult lung organoids.
B-D. Bright-field microscopy of organoids in 3D culture grown in different media/conditions (B), imaged serially over days (C) and at different passages (D). Scale bar = 100 μm.
E. Serial cuts of *HistoGel*-embedded organoids were analyzed by H&E staining. Scale bar = 50 μm.

Supplementary Figure 3. Adult stem cell-derived lung organoids are propagatable models with both proximal and distal airway components.

A. Schematic lists the various markers used here for qPCR and immunofluorescence to confirm the presence of all cell types in the 3D lung organoids here and in 2D monolayers later (in Fig 3).

B-H. Bar graphs display the relative abundance of various cell type markers (normalized to 18S) in adult lung organoids (ALO), compared to the airway (NHBE) and/or alveolar (AT2) control cells, as appropriate. *p*-values were analyzed by one-way Anova. Error bars denote S.E.M; n = 3-6 datasets.

I. 3D organoids grown in 8-well chamber slides were fixed, immunostained and visualized by confocal microscopy, as in **Fig 2K**. Scale bar = 50 μm.

Supplementary Figure 4. Alternative models of lung epithelial cells are used here for SARS-CoV-2 infection and/or as a control for qPCR.

A-E. Two different types of 2D polarized monolayers are prepared using adult lung organoids. Schematics in A-B show growth as submerged or ALI models. Panel C shows bar graphs with transepithelial resistance (TEER) across ALI monolayers. Brightfield images in D-E show representative fields of monolayers. Scale bar = 100 μm. Arrows = self-organized vacuolar regions were seen in monolayers on submerged transwell.

F-I. Monolayers of primary airway epithelial cells (small airway epi; F-G; bronchial epi; H-I) were visualized by bright field microscopy (H, H) or by fixing, staining and visualizing by confocal microscopy (H, I). Representative images in G and H are presented as maximum projected z-stacks on the left and as an orthogonal view on the right.

J-L. hiPSC-derived AT2 cells, prepared using the i-HAEpC2 cell kit, were grown in monolayers on transwell inserts to form a polarized. Brightfield images are shown in K. Monolayers were fixed and stained for several markers and analyzed by confocal microscopy. Representative images are shown in L. Scale bar = 20 μm.

Supplementary Figure 5. Proof of infectivity.

A. Monolayers of ALO (either transwell submerged models or ALI, *Left*) and monolayers of hiPSC-derived AT2 cells (*Right*) were infected or not with SARS-CoV-2 and analyzed for viral envelope gene (E gene). Bar graphs display the relative expression of E gene in infected ALO monolayers, indicative of viral infection.

B. Line graphs show the change in E gene expression in infected monolayers over 24 h period (from 48 hpi to 72 hpi) where values at 72 hpi are normalized to that at 48 hpi. Data is presented as S.E.M. of 3 independent repeats.

C. Monolayers of ALO1-3 were challenged with SARS-CoV-2 for indicated time points prior to fixation and staining for KRT5 (red) and viral nucleocapsid protein (green) and DAPI (blue; nuclei) and visualized by confocal microscopy. Representative images are shown, displaying various cytopathic effects. Scale bar = 15 μm.

Supplementary Figure 6. Differential expression analysis of RNA Seq datasets from lung autopsies (normal vs. COVID-19).
Publicly available RNA Seq datasets (GSE151764) of lung autopsies from patients who were deceased due to COVID-19 or non-infectious causes (normal lung control) were analyzed for differential expression of genes and displayed as a heatmap.

Differential Expression Analysis of Human Healthy vs COVID-19 lungs

GSE151764

Pathways upregulated in human SARS-CoV-infected lungs

Pathway name	Entities found	Entities Total	Entities ratio	Entities pValue	Entities FDR	Reactions found	Reactions total	Reactions ratio	Species name
Interferon Signaling	43	396	0.027	1.11E-16	1.11E-14	43	66	0.005	Homo sapiens
Interferon alpha/beta signaling	34	186	0.013	1.11E-16	1.11E-14	9	20	0.002	Homo sapiens
Cytokine Signaling in Immune system	65	1,312	0.089	1.11E-16	1.11E-14	136	735	0.057	Homo sapiens
Immune System	74	2,869	0.196	1.11E-16	1.11E-14	245	1,634	0.126	Homo sapiens
Interleukin-10 signaling	13	86	0.006	9.85E-13	7.88E-11	12	15	0.001	Homo sapiens
Interferon gamma signaling	18	252	0.017	9.26E-12	6.11E-10	9	15	0.001	Homo sapiens
Chemokine receptors bind chemokines	10	57	0.004	1.08E-10	6.17E-9	7	19	0.001	Homo sapiens
Signaling by Interleukins	23	639	0.044	6.81E-9	3.41E-7	80	490	0.038	Homo sapiens
Insulin-like Growth Factor-2 mRNA Binding Proteins (IGF2BPs/IMPs/VICKZs) bind RNA	5	13	0.001	1.27E-7	5.58E-6	2	3	0	Homo sapiens
Antiviral mechanism by IFN-stimulated genes	8	94	0.006	1.93E-6	7.73E-5	25	31	0.002	Homo sapiens
CD163 mediating an anti-inflammatory response	4	14	0.001	7.8E-6	2.81E-4	4	9	0.001	Homo sapiens
OAS antiviral response	4	15	0.001	1.02E-5	3.37E-4	9	15	0.001	Homo sapiens
Peptide ligand-binding receptors	10	211	0.014	1.71E-5	5.14E-4	7	76	0.006	Homo sapiens
Interleukin-4 and Interleukin-13 signaling	9	211	0.014	1.01E-4	2.84E-3	8	46	0.004	Homo sapiens
Cyclin A/B1/B2 associated events during G2/M transition	4	32	0.002	1.89E-4	4.91E-3	20	25	0.002	Homo sapiens
G0 and Early G1	4	38	0.003	3.61E-4	9.02E-3	4	27	0.002	Homo sapiens
Interleukin 6 signaling	3	17	0.001	4.66E 4	1.07E 2	18	19	0.001	Homo sapiens
ISG15 antiviral mechanism	5	83	0.006	8.31E-4	1.75E-2	16	16	0.001	Homo sapiens
Regulation of APC/C activators between G1/S and early anaphase	5	83	0.006	8.31E-4	1.75E-2	8	14	0.001	Homo sapiens
Polo-like kinase mediated events	3	23	0.002	1.11E-3	2.22E-2	12	15	0.001	Homo sapiens

Pathways downregulated in human SARS-CoV-infected lungs

Pathway name	Entities found	Entities Total	Entities ratio	Entities pValue	Entities FDR	Reactions found	Reactions total	Reactions ratio	Species name
Chemokine receptors bind chemokines	8	57	0.004	2.85E-11	4.98E-9	7	19	0.001	Homo sapiens
Immune System	31	2,869	0.196	1.25E-10	1.09E-8	117	1,634	0.126	Homo sapiens
Interleukin-4 and Interleukin-13 signaling	10	211	0.014	2.82E-9	1.64E-7	5	46	0.004	Homo sapiens
RUNX1 and FOXP3 control the development of regulatory T lymphocytes (Tregs)	4	17	0.001	4.31E-7	1.85E-5	14	20	0.002	Homo sapiens
Peptide ligand-binding receptors	8	211	0.014	6.71E-7	2.35E-5	7	76	0.006	Homo sapiens
Signaling by Interleukins	12	639	0.044	1.5E-6	4.36E-5	25	490	0.038	Homo sapiens
Cytokine Signaling in Immune system	15	1,312	0.089	2.61E-6	6.52E-5	31	735	0.057	Homo sapiens
Dectin-1 mediated noncanonical NF-kB signaling	4	66	0.004	8.64E-5	1.81E-3	1	9	0.001	Homo sapiens
Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell	7	316	0.022	1.08E-4	2.06E-3	8	43	0.003	Homo sapiens
Class A/1 (Rhodopsin-like receptors)	8	462	0.031	1.83E-4	3.12E-3	7	152	0.012	Homo sapiens
Interleukin-10 signaling	4	86	0.006	2.37E-4	3.55E-3	2	15	0.001	Homo sapiens
RUNX3 Regulates Immune Response and Cell Migration	2	10	0.001	5.8E-4	7.75E-3	2	5	0	Homo sapiens
Extra-nuclear estrogen signaling	4	110	0.007	5.96E-4	7.75E-3	3	38	0.003	Homo sapiens
BH3-only proteins associate with and inactivate anti-apoptotic BCL-2 members	2	11	0.001	7E-4	8.39E-3	3	4	0	Homo sapiens
CLEC7A (Dectin-1) signaling	4	120	0.008	8.23E-4	9.05E-3	1	45	0.003	Homo sapiens
Generation of second messenger molecules	3	59	0.004	1.17E-3	1.17E-2	8	17	0.001	Homo sapiens
Innate Immune System	12	1,329	0.091	1.68E-3	1.57E-2	34	697	0.054	Homo sapiens
GPCR ligand binding	8	652	0.044	1.75E-3	1.57E-2	7	179	0.014	Homo sapiens
Adaptive Immune System	10	999	0.068	2.06E-3	1.85E-2	52	261	0.02	Homo sapiens
Estrogen-dependent nuclear events downstream of ESR-membrane signaling	2	29	0.002	4.67E-3	3.74E-2	1	12	0.001	Homo sapiens

Supplementary Figure 7. Reactome pathway analysis of differentially expressed genes in lung autopsies (normal vs. COVID-19). Reactome-pathway analysis of the differentially expressed genes shows the major pathways upregulated in COVID-19-affected lungs. _Top_: Visualization as flattened (left) and hierarchical (right, insets) reactome. _Bottom_: Visualization of the same data as tables with statistical analysis indicative of the degree of pathway enrichment.

Supplementary Figure 8. Differential expression analysis of RNA Seq datasets from lung organoid monolayers, infected or not, with SARS-CoV-2. Adult lung organoid monolayers infected or not with SARS-CoV-2 were analyzed by RNA seq and differential expression analysis. Differentially expressed genes are displayed as a heatmap.

Differential Expression Analysis of Human Lung Monolayers: Infected vs Uninfected

Pathways upregulated in human SARS-CoV-infected lungs

Pathway name	Entities found	Entities Total	Entities ratio	Entities pValue	Entities FDR	Reactions found	Reactions total	Reactions ratio	Species name
Interferon Signaling	17	396	0.027	1.11E-16	4.22E-15	23	66	0.005	Homo sapiens
Interferon alpha/beta signaling	17	186	0.013	1.11E-16	4.22E-15	2	20	0.002	Homo sapiens
Cytokine Signaling in Immune system	17	1,312	0.089	1.15E-10	2.89E-9	23	735	0.057	Homo sapiens
Immune System	18	2,869	0.196	2.54E-6	4.83E-5	76	1,634	0.126	Homo sapiens
OAS antiviral response	2	15	0.001	4.76E-4	7.15E-3	3	15	0.001	Homo sapiens
Antiviral mechanism by IFN-stimulated genes	3	94	0.006	1.03E-3	1.24E-2	19	31	0.002	Homo sapiens
Interferon gamma signaling	4	252	0.017	1.89E-3	2.08E-2	2	15	0.001	Homo sapiens
Transfer of LPS from LBP carrier to CD14	1	3	0	6.32E-3	5.69E-2	2	2	0	Homo sapiens
TRIF-mediated programmed cell death	1	10	0.001	2.09E-2	1.66E-1	3	3	0	Homo sapiens
MyD88 deficiency (TLR2/4)	1	18	0.001	3.73E-2	1.66E-1	2	2	0	Homo sapiens
IRAK2 mediated activation of TAK1 complex upon TLR7/8 or 9 stimulation	1	18	0.001	3.73E-2	1.66E-1	1	5	0	Homo sapiens
TRAF6-mediated induction of TAK1 complex within TLR4 complex	1	19	0.001	3.94E-2	1.66E-1	6	6	0	Homo sapiens
IRAK4 deficiency (TLR2/4)	1	19	0.001	3.94E-2	1.66E-1	2	2	0	Homo sapiens
Activation of IRF3/IRF7 mediated by TBK1/IKK epsilon	1	20	0.001	4.14E-2	1.66E-1	7	9	0.001	Homo sapiens
Caspase activation via Death Receptors in the presence of ligand	1	20	0.001	4.14E-2	1.66E-1	1	9	0.001	Homo sapiens
IKK complex recruitment mediated by RIP1	1	24	0.002	4.95E-2	1.86E-1	3	3	0	Homo sapiens
Regulation of TLR by endogenous ligand	1	31	0.002	6.35E-2	1.86E-1	1	12	0.001	Homo sapiens
Caspase activation via extrinsic apoptotic signalling pathway	1	32	0.002	6.54E-2	1.86E-1	1	17	0.001	Homo sapiens
Diseases of Immune System	1	34	0.002	6.94E-2	1.86E-1	4	15	0.001	Homo sapiens
Diseases associated with the TLR signaling cascade	1	34	0.002	6.94E-2	1.86E-1	4	15	0.001	Homo sapiens

Pathways downregulated in human SARS-CoV-infected lungs

Pathway name	Entities found	Entities Total	Entities ratio	Entities pValue	Entities FDR	Reactions found	Reactions total	Reactions ratio	Species name
rRNA processing in the mitochondrion	1	40	0.003	1.89E-2	8.37E-2	1	6	0	Homo sapiens
tRNA processing in the mitochondrion	1	45	0.003	2.13E-2	8.37E-2	2	3	0	Homo sapiens
Mitochondrial translation termination	1	94	0.006	4.4E-2	8.37E-2	5	5	0	Homo sapiens
Mitochondrial translation elongation	1	94	0.006	4.4E-2	8.37E-2	5	8	0.001	Homo sapiens
Mitochondrial translation initiation	1	96	0.007	4.49E-2	8.37E-2	1	4	0	Homo sapiens
Mitochondrial translation	1	102	0.007	4.77E-2	8.37E-2	11	17	0.001	Homo sapiens
Rho GTPase cycle	1	144	0.01	6.67E-2	8.37E-2	1	5	0	Homo sapiens
tRNA processing	1	182	0.012	8.37E-2	8.37E-2	2	54	0.004	Homo sapiens
rRNA processing	1	245	0.017	1.11E-1	1.11E-1	1	21	0.002	Homo sapiens
Translation	1	339	0.023	1.51E-1	1.51E-1	11	99	0.008	Homo sapiens
Signaling by Rho GTPases	1	457	0.031	1.99E-1	1.99E-1	1	117	0.009	Homo sapiens
Metabolism of RNA	1	782	0.053	3.18E-1	3.18E-1	2	187	0.014	Homo sapiens
Metabolism of proteins	1	2,356	0.161	7.06E-1	7.06E-1	11	892	0.069	Homo sapiens
Signal Transduction	1	3,396	0.231	8.42E-1	8.42E-1	1	2,386	0.184	Homo sapiens

Supplementary Figure 9. Reactome pathway analysis of differentially expressed genes in lung organoid monolayers infected with SARS-CoV-2.
Reactome-pathway analysis of the differentially expressed genes shows the major pathways upregulated in SARS-CoV-2-infected lung organoid monolayers. _Top_: Visualization as flattened (left) and hierarchical (right, insets) reactfoam. _Bottom_: Visualization of the same data as tables with statistical analysis indicative of the degree of pathway enrichment.

Supplementary Bibliography

(CAP), C.o.A.P. (2020). Amended COVID-19 autopsy guideline statement from the CAP autopsy committee.

(CDC), C.f.D.C.a.P. (2020). Collection and Submission of Post-mortem Specimens from Deceased Persons with Known or Suspected COVID-19 (Interim Guidance). .

Barrett, T., Suzek, T.O., Troup, D.B., Wilhite, S.E., Ngau, W.C., Ledoux, P., Rudnev, D., Lash, A.E., Fujibuchi, W., and Edgar, R. (2005). NCBI GEO: mining millions of expression profiles--database and tools. Nucleic Acids Res *33*, D562-566.

Barrett, T., Wilhite, S.E., Ledoux, P., Evangelista, C., Kim, I.F., Tomashevsky, M., Marshall, K.A., Phillippy, K.H., Sherman, P.M., Holko, M.*, et al.* (2013). NCBI GEO: archive for functional genomics data sets--update. Nucleic Acids Res *41*, D991-995.

Corman, V.M., Landt, O., Kaiser, M., Molenkamp, R., Meijer, A., Chu, D.K., Bleicker, T., Brunink, S., Schneider, J., Schmidt, M.L.*, et al.* (2020). Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. Euro Surveill *25*.

Dalerba, P., Kalisky, T., Sahoo, D., Rajendran, P.S., Rothenberg, M.E., Leyrat, A.A., Sim, S., Okamoto, J., Johnston, D.M., Qian, D.*, et al.* (2011). Single-cell dissection of transcriptional heterogeneity in human colon tumors. Nat Biotechnol *29*, 1120-1127.

Dalerba, P., Sahoo, D., Paik, S., Guo, X., Yothers, G., Song, N., Wilcox-Fogel, N., Forgo, E., Rajendran, P.S., Miranda, S.P.*, et al.* (2016). CDX2 as a Prognostic Biomarker in Stage II and Stage III Colon Cancer. N Engl J Med *374*, 211-222.

Edgar, R., Domrachev, M., and Lash, A.E. (2002). Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Res *30*, 207-210.

Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P., Haw, R., Jassal, B., Korninger, F., May, B.*, et al.* (2018). The Reactome Pathway Knowledgebase. Nucleic Acids Res *46*, D649-D655.

Ghosh, P., Swanson, L., Sayed, I.M., Mittal, Y., Lim, B.B., Ibeawuchi, S.R., Foretz, M., Viollet, B., Sahoo, D., and Das, S. (2020). The stress polarity signaling (SPS) pathway serves as a marker and a target in the leaky gut barrier: implications in aging and cancer. Life Sci Alliance *3*.

Gotoh, S., Ito, I., Nagasaki, T., Yamamoto, Y., Konishi, S., Korogi, Y., Matsumoto, H., Muro, S., Hirai, T., Funato, M.*, et al.* (2014). Generation of alveolar epithelial spheroids via isolated progenitor cells from human pluripotent stem cells. Stem Cell Reports *3*, 394-403.

Irizarry, R.A., Bolstad, B.M., Collin, F., Cope, L.M., Hobbs, B., and Speed, T.P. (2003a). Summaries of Affymetrix GeneChip probe level data. Nucleic Acids Res *31*, e15.

Irizarry, R.A., Hobbs, B., Collin, F., Beazer-Barclay, Y.D., Antonellis, K.J., Scherf, U., and Speed, T.P. (2003b). Exploration, normalization, and summaries of high density oligonucleotide array probe level data. Biostatistics *4*, 249-264.

Jacob, A., Vedaie, M., Roberts, D.A., Thomas, D.C., Villacorta-Martin, C., Alysandratos, K.D., Hawkins, F., and Kotton, D.N. (2019). Derivation of self-renewing lung alveolar epithelial type II cells from human pluripotent stem cells. Nat Protoc *14*, 3303-3332.

Lamers, M.M., Beumer, J., van der Vaart, J., Knoops, K., Puschhof, J., Breugem, T.I., Ravelli, R.B.G., Paul van Schayck, J., Mykytyn, A.Z., Duimel, H.Q.*, et al.* (2020). SARS-CoV-2 productively infects human gut enterocytes. Science *369*, 50-54.

Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics *12*, 323.

Lopez-Sanchez, I., Dunkel, Y., Roh, Y.S., Mittal, Y., De Minicis, S., Muranyi, A., Singh, S., Shanmugam, K., Aroonsakool, N., Murray, F.*, et al.* (2014). GIV/Girdin is a central hub for profibrogenic signalling networks during liver fibrosis. Nat Commun *5*, 4451.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Miyoshi, H., and Stappenbeck, T.S. (2013). In vitro expansion and genetic modification of gastrointestinal stem cells in spheroid culture. Nat Protoc *8*, 2471-2482.

Pachter, L. (2011). Models for transcript quantification from RNA-Seq. In arXiv e-prints.

Sachs, N., Papaspyropoulos, A., Zomer-van Ommen, D.D., Heo, I., Bottinger, L., Klay, D., Weeber, F., Huelsz-Prince, G., Iakobachvili, N., Amatngalim, G.D.*, et al.* (2019). Long-term expanding human airway organoids for disease modeling. EMBO J *38*.

Sato, T., Vries, R.G., Snippert, H.J., van de Wetering, M., Barker, N., Stange, D.E., van Es, J.H., Abo, A., Kujala, P., Peters, P.J.*, et al.* (2009). Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. Nature *459*, 262-265.

Sayed, I.M., Chakraborty, A., Abd El Hafeez, A.A., A., S., Sahan, A.Z., WJM, H., Sahoo, D., Ghosh, P., Hazra, T.K., and Das, S. (2020a). The DNA Glycosylase NEIL2 Suppresses Fusobacterium-Infection-Induced Inflammation and DNA Damage in Colonic Epithelial Cells. Cells *9*.

Sayed, I.M., Sahan, A.Z., Venkova, T., Chakraborty, A., Mukhopadhyay, D., Bimczok, D., Beswick, E.J., Reyes, V.E., Pinchuk, I., Sahoo, D.*, et al.* (2020b). Helicobacter pylori infection downregulates the DNA glycosylase NEIL2, resulting in increased genome damage and inflammation in gastric epithelial cells. J Biol Chem *295*, 11082-11098.

Sayed, I.M., Suarez, K., Lim, E., Singh, S., Pereira, M., Ibeawuchi, S.R., Katkar, G., Dunkel, Y., Mittal, Y., Chattopadhyay, R.*, et al.* (2020c). Host engulfment pathway controls inflammation in inflammatory bowel disease. FEBS J.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell *177*, 1888-1902 e1821.

van der Vaart, J., and Clevers, H. (2020). Airway organoids as models of human disease. J Intern Med.

Volkmer, J.P., Sahoo, D., Chin, R.K., Ho, P.L., Tang, C., Kurtova, A.V., Willingham, S.B., Pazhanisamy, S.K., Contreras-Trujillo, H., Storm, T.A.*, et al.* (2012). Three differentiation states risk-stratify bladder cancer into distinct subtypes. Proc Natl Acad Sci U S A *109*, 2078-2083.

Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene expression data analysis. Genome Biol *19*, 15.

Yamamoto, Y., Gotoh, S., Korogi, Y., Seki, M., Konishi, S., Ikeo, S., Sone, N., Nagasaki, T., Matsumoto, H., Muro, S.*, et al.* (2017). Long-term expansion of alveolar stem cells derived from human iPS cells in organoids. Nat Methods *14*, 1097-1106.

Zhang, Z., Luo, D., Zhong, X., Choi, J.H., Ma, Y., Wang, S., Mahrt, E., Guo, W., Stawiski, E.W., Modrusan, Z.*, et al.* (2019). SCINA: A Semi-Supervised Subtyping Algorithm of Single Cells and Bulk Samples. Genes (Basel) *10*.

Zhou, J., Li, C., Sachs, N., Chiu, M.C., Wong, B.H., Chu, H., Poon, V.K., Wang, D., Zhao, X., Wen, L.*, et al.* (2018). Differentiated human airway organoids to assess infectivity of emerging influenza virus. Proceedings of the National Academy of Sciences of the United States of America *115*, 6822-6827.