1 **Gulf of Mexico blue hole harbors high levels of novel microbial lineages**
2

3 Patin NV[1,2]*, Dietrich ZA[3], Stancil A[4], Quinan M[4], Beckler JS[4], Hall ER[5], Culter J[5], Smith
4 CG[6], Taillefert M[7], Stewart FJ[1,2,8]
5

1. School of Biological Sciences, Georgia Institute of Technology, Atlanta GA 30332
2. Center for Microbial Dynamics and Infection, Georgia Institute of Technology, Atlanta
GA 30332
3. Bowdoin College, Brunswick ME 04011
4. Harbor Branch Oceanographic Institute, Florida Atlantic University, Ft. Pierce FL 34946
5. Mote Marine Laboratory, Sarasota FL 34236
6. U.S. Geological Survey, St. Petersburg Coastal and Marine Science Center, St.
Petersburg FL 33701
7. School of Earth & Atmospheric Sciences, Georgia Institute of Technology, Atlanta GA
30332
8. Department of Microbiology & Immunology, Montana State University, Bozeman MT
59717
18

19 * Corresponding author: nastassia.patin@noaa.gov

21 **Abstract**
22

Exploration of oxygen-depleted marine environments has consistently revealed novel
microbial taxa and metabolic capabilities that expand our understanding of microbial
evolution and ecology. Marine blue holes are shallow karst formations characterized by
low oxygen and high organic matter content. They are logistically challenging to sample,
and thus our understanding of their biogeochemistry and microbial ecology is limited. We
present a metagenomic characterization of Amberjack Hole on the Florida continental
shelf (Gulf of Mexico). Dissolved oxygen became depleted at the hole's rim (32 m water
depth), remained low but detectable in an intermediate hypoxic zone (40-75 m), and then
increased to a secondary peak before falling below detection in the bottom layer (80-110
m), concomitant with increases in nutrients, dissolved iron, and a series of sequentially
more reduced sulfur species. Microbial communities in the bottom layer contained
heretofore undocumented levels of the recently discovered phylum Woesearchaeota (up
to 58% of the community), along with lineages in the bacterial Candidate Phyla Radiation
(CPR). Thirty-one high-quality metagenome-assembled genomes (MAGs) showed
extensive biochemical capabilities for sulfur and nitrogen cycling, as well as for resisting
and respiring arsenic. One uncharacterized gene associated with a CPR lineage
differentiated hypoxic from anoxic zone communities. Overall, microbial communities and
geochemical profiles were stable across two sampling dates in the spring and fall of 2019.

The blue hole habitat is a natural marine laboratory that provides opportunities for sampling taxa with under-characterized but potentially important roles in redox-stratified microbial processes.

Introduction

Blue holes are subsurface caverns found in karst bedrock environments. They formed during climatic periods when low sea levels exposed the bedrock to weathering, and subsequently became submerged as sea levels rose (1). Marine blue holes differ from anchialine blue holes, such as those found in the Bahamas and the Yucatán peninsula, as they do not have freshwater layers and are not exposed to the atmosphere. Anchialine blue holes can be highly stratified, with anoxic and sulfidic bottom waters (2–4) and microbial communities distinct from other marine and freshwater systems (5–8). However, data on true marine blue holes are limited. A recent study on the Yongle Blue Hole, the deepest known marine blue hole with a bottom depth of 300 m, found the water column became anoxic around 100 m with increases in hydrogen sulfide, methane, and dissolved inorganic carbon below that depth (9). This and two additional studies also found that microbial communities in the Yongle Blue Hole were notably different from those of the surrounding pelagic water column, with anoxic layers in the hole dominated by taxa linked to sulfur oxidation and nitrate reduction (9–11). Further, deep blue hole waters may be resistant to mixing with waters outside the hole, especially in regions with limited seasonal variation or water mass intrusion. These observations suggest the potential for blue holes to harbor novel microbial lineages as a consequence of both unique geochemistry and environmental isolation.

Locations of 18 blue holes have been recorded in offshore waters on the west Florida shelf; however, many more may exist but remain undiscovered due to a lack of systematic survey (12) (J. Culter, Mote Marine Laboratory, unpublished data, December 7, 2017). According to anecdotal reports from recreational divers and fishers, the rims of these holes feature dense communities of corals, sponges, and other invertebrates, in contrast with the more barren sandy bottom of the surrounding shelf. Commercially and recreationally valuable fishes also congregate at the rims. The presence of high biomass has led to speculation that blue holes could represent offshore sources of groundwater-derived nutrients transported via extensions of the Floridan aquifer (13). However, unlike for some of the nearshore blue holes (12), no measurements of groundwater discharge have been made in the offshore blue holes. Elevated nutrient levels at these sites have broad relevance for coastal ecology in the Gulf of Mexico, particularly as they may fuel phytoplankton blooms (14,15). Indeed, the west coast of Florida experiences frequent and intense harmful algal blooms (HABs). While numerous nutrient sources have been identified as potential HAB triggers, the relative importance of these sources remains

81 geographically unconstrained and our knowledge of what drives HABs is by no means
82 complete (16–18).
83
Gulf of Mexico blue holes may be chemically stratified and devoid of oxygen. In one of
the only biological studies of these features, Garman et al. (2011) explored Jewfish Sink,
a coastal blue hole near Hudson, Florida. Oxygen concentrations fell from near saturation
at the rim (~2 m water depth) to zero around 20 m water depth and remained below
detection all the way to the bottom of the hole (~64 m water depth), with the anoxic layer
further characterized by pronounced sulfide accumulation with depth. A clone library of
16S rRNA genes from microbial mats in the hole revealed a taxonomically rich community
(338 operational taxonomic units, including 150 bacterial and 188 archaeal taxa), with
sequences closely related to those from low-oxygen habitats including deep-sea
sediments, salt marshes, cold seeps, and whale falls (6). These sequences represent
taxa linked to a range of metabolisms, including dissimilatory sulfur, methane, and
nitrogen cycling (both oxidative and reductive). This study also reported dense and
chemically variable clouds of particulates within the hole; these included iron-sulfide
minerals, suggesting the potential for microbial-metal interactions in the water column.
98
This work, alongside evidence from the South China Sea sites (9), suggests marine blue
holes are biogeochemically complex features and potential hotspots for microbial
diversity. Indeed, oxygen-depleted marine water columns have been rich targets for
exploring novel microbial processes. Within the past five years alone, studies in these
systems have shed new light on microbial processes of arsenic respiration (19), cryptic
sulfur (20) and oxygen cycling (21), low-oxygen adapted nitrification (22) and
denitrification (23), and anaerobic methane oxidation (24). Moreover, oxygen-depleted
waters are expanding worldwide (25), making it critical to understand how oxygen
concentration impacts microbially regulated nutrient and energy budgets, and how these
impacts vary among sites. Most microbes in these systems are uncultivated and
phylogenetically divergent from better-studied relatives. Many of these taxa remain
unclassified beyond the phylum or class level and likely represent lineages uniquely
adapted to low-oxygen conditions. In lieu of cultivation, metagenomic analysis enables
exploration of both the taxonomy and function of microbial players in blue holes and
similar low-oxygen environments.
114
Here, we used metagenomics to describe the microbial ecosystem in a marine blue hole
on the west Florida shelf. To help interpret microbial processes, we also present a
comprehensive electrochemistry-based analysis of redox chemical speciation. Amberjack
(AJ) Hole lies ~50 km west of Sarasota, Florida at a water depth of 32 m. Like Jewfish
Sink, AJ is conical in shape, with a narrow rim (25 m diameter) and a wider floor
(approximately 100 m diameter, as determined by initial reconnaissance dives for this

study). Water depth at the floor ranges from 110 m at the edge to 90 m at the center,
where a debris pile of fine-grained sediment has accumulated. Our knowledge of AJ, like
that of other marine blue holes, is limited and based primarily on exploration by a small
number of technical divers. The hole's shape and depth present a challenge for SCUBA,
as well as for instrumentation or submersibles. During two expeditions in May and
September 2019, we used a combination of technical divers and Niskin bottles to sample
water column microbial communities. These collections spanned oxygenated waters at
the rim, an anoxic but non-sulfidic intermediate layer, and a bottom layer rich in reduced
sulfur compounds. Our results reveal a system that is highly stratified, apparently stable
between timepoints, and phylogenetically diverse, with high representation by
uncultivated and poorly understood taxa. The results suggest marine blue holes may
serve as unique natural laboratories for exploring redox-stratified microbial ecosystems,
while highlighting a need for future biogeochemical exploration of other blue holes and
similar habitats.
135
136 **Results**
137
138 *Sampling scheme*
139
We sampled physical, chemical, and biological features of Amberjack Hole in May and
September 2019. The May data set included one conductivity, temperature, and depth
(CTD) profile with a coupled dissolved oxygen sensor, along with water column samples
for chemical analyses (5 depths) and microbial community sequencing (11 depths,
including 5 from within the hole). These samples were collected via a combination of diver
bottle water exchange, hand-cast Niskin bottles, and automated Niskin sampling on a
rosette (Table S1). The CTD profile was acquired by attaching a combined CTD and
optical dissolved oxygen sensor to an autonomous lander deployed for 24 hours to
measure sediment respiratory processes and fluxes (data not included in this study). The
lander was positioned at 106 m water depth on the slope of the debris pile; the deepest
May water sample was acquired by a diver at this depth.
151
In September, improved sampling design resulted in higher spatial resolution for all water
column parameters. We obtained two CTD profiles and eight samples from within the hole
for microbiome analysis. All September water samples were acquired by hand-cast Niskin
bottles, with the deepest sample from 95 m, the presumed peak of the debris pile. In
contrast to the May sampling, all water sampling in September was performed on the day
of the CTD casts, ensuring that chemical and biological measurements were temporally
coupled. We therefore focus primarily on September results (below), with exceptions
where noted.
160

Water column physical and chemical profiles

Amberjack Hole was highly stratified (Fig. 1). Salinity increased sharply by one PSU between the overlying water column (35.2) and the blue hole water mass (~30 m, 36.2). Within the hole, salinity decreased slightly with depth (maximum difference 0.4 PSU) until 75 m and then increased to ~36 PSU at the bottom. Dissolved oxygen decreased sharply upon entry into the hole, from 100% saturation at the rim (32 m) to less than 5% saturation by 40 m (Fig. 1B). Oxygen then increased gradually to a secondary maximum of 40% at 75 m, before dropping to near 0% (anoxia) below 80 m.

Dissolved inorganic nitrogen (nitrate (NO_3^-) + nitrite (NO_2^-), or NO_x) spiked between 40 and 70 m from 0 to ~12 µM in May and nearly 17 µM in September, while ammonium (NH_4^+) and phosphate (PO_4^{3-}) increased sharply below 80 m (Fig. 1C). Bottom water nutrient concentrations were higher in September than in May, with NH_4^+ reaching nearly 50 µM and PO_4^{3-} exceeding 8 µM. Particulate nutrients (N, P, and carbon (C)) and chlorophyll *a* concentrations were seemingly less variable throughout the water column in May compared to September (Fig. S2), although this pattern may be an artifact of the lower sampling resolution. In September, particulate nutrients and chlorophyll *a* spiked at the hole opening (between 30 and 40 m) with P at 0.1 µM, N at 1.2 µM, C at 11 µM, and chlorophyll *a* at nearly 1.5 µg/L. Particulate nutrients remained low deeper in the water column with the exception of P, which spiked again to 0.14 µM at 80 m. Particulate C and N also increased slightly below 80 m (Fig. S2).

Coinciding with anoxia, ferrous iron ($Fe(II)_d$) increased from 16 nM to 187 nM between 70 and 80 m, respectively. With increasing depth, sequentially more reduced sulfur compounds were observed (Fig. 1D):

a) at 85 m, $S_2O_3^{2-}$ (thiosulfate, sulfur (S) in the +II oxidation state), reached a maximum of 264 µM;

b) at 90 m, S(0), representing either S_8 (i.e. elemental sulfur, S in the 0 oxidation state) or most S within polysulfide (S_x^{2-}, e.g. S_4^{2-} or S_8^{2-}), in filtered (< 0.7 µm) and unfiltered water reached 126 µM and 599 µM, respectively; and

c) at 95 m (deepest sample), combined $\Sigma S(-II)$ (i.e. either as hydrogen sulfide (HS^-), polysulfide (S_x^{2-}), or both) reached a maximum of 61 µM.

Due to analytical uncertainties associated with the voltammetric quantification scheme, the $\Sigma S(-II)$ as presented may contribute redundantly to both (b) and (c) from a sulfur mass balance perspective. Only 6% of the S(-II) signal in the 95 m sample is lost upon filtering through 0.7 µm GFF filters (not shown), compared to 93 -100% losses in the 85 and 80 m samples, respectively, suggesting that the deepest waters are probably enriched in true hydrogen sulfide (HS^-). Indeed, significant HS^- is fluxing from sediments as revealed by benthic flux and pore water measurements (data not shown).

Notably, the zone of S(0) (i.e. elemental sulfur) accumulation between 80 and 90 m was
marked by a spike in turbidity (Fig. 1B). In addition, particulate P was elevated
concomitantly with Fe_d and $Fe(II)_d$ between 70 and 80 m (Fig. 1D and S2). The latter
suggests the presence of particulate or colloidal minerals and associated adsorption sites
(e.g., Fe oxide colloids).

The May sampling resulted in a single data point from a depth of 85 m, which showed
levels of 19.8 ± 2.3 µM $S_2O_3^{2-}$, 93 µM ΣS(-II), and no S(0), similar to the September
measurements from that depth.

Microbial community composition

16S rRNA gene amplicon sequencing yielded 12,692 to 118,397 reads per sample after
filtering for quality (Table S1). Water column microbial communities differed significantly
based on depth grouping (PERMANOVA p=0.016), partitioning into shallow (0-32 m, oxic
zone above the hole), middle (40-70 m, hypoxic zone), and deep (80-106 m, anoxic zone)
groups (Fig. 2A). The shallow group was characterized by the ubiquitous cyanobacteria
Synechococcus and *Prochlorococcus*, as well as several clades of the heterotrophic
alphaproteobacterium SAR11 (particularly clade Ia) (Figure 2B). Cyanobacteria were
absent below 32 m in both May and September. The middle water column featured high
frequencies of *Nitrosopumilus* sp., a member of the ammonia-oxidizing Thaumarchaeota,
as well as members of the sulfur-oxidizing family *Thioglobaceae* (Fig. 2B). Other groups,
composing 1-10% of the community in this depth zone, included Marine Group II and III
Thermoplasmatota (formerly MGII/III Euryarchaeota) and members of the family
Gimesiaceae (phylum Planctomycetes).

The anoxic and sulfidic deep water column was dominated by Woesearchaeota of the
DPANN superphylum. Woesearchaeota represented 33% - 56% of all sequences below
75 m in both May and September and were comprised of 74 sequence variants (SVs),
with two SVs representing between 84% (May 85 m) and 99% (September 90 m) of the
total Woesearchaeotal fraction. Other well-represented groups included *Nitrosopumilus*
sp, *Thioglobaceae* (SUP05 clade), and a member of the phylum Bacteroidota (formerly
Bacteroidetes) associated with hydrothermal vents ("Bacteroidetes VC2.1 Bac22" in the
SILVA database) (Fig. 2B). While Shannon diversity was similar among depth groups,
Simpson diversity was ~50% higher in the deep compared to the middle and shallow
groups (Fig. S3).

Depth patterns in community composition were highly similar between May and
September. The most abundant taxa showed only minor differences in frequency

between these months, with a few exceptions (Fig. 3). Marine Group II Thermoplasmatota
reached 12% relative abundance in the middle hypoxic zone in September while in May
this taxon never exceeded 1%; in contrast, Marine Group III peaked at around 6% in May
and in September only reached about half that level. In May, *Thioglobaceae* frequency
increased steadily below 30 m to peak at >10% of the community in the deep anoxic zone.
In contrast, *Thioglobaceae* was not detected in the anoxic zone in September. Rather,
Arcobacter sp. (represented by a single SV) spiked at 80 m in September to nearly 10%
of the community. In all other samples, this taxon never exceeded 0.5% at any sampled
depth. The 80 m depth was not sampled in May.
250
251 *Metagenomes and MAG taxonomy*
252
We obtained four metagenomes from two depths in both May and September. These
represent communities at 60 m (both months) in the hypoxic (~15% O_2 saturation) and
Nitrosopumilus-dominated zone, and at 106 m (May) and 95 m (September) in the anoxic,
sulfidic, and Woesearchaeota-dominated zone. Sequencing and assembly results are
provided in Table S2. Metagenome binning involving data from all samples yielded 31
high-quality, non-redundant MAGs representing a diverse array of microbial taxa (Table
1). These included eight archaeal MAGs, seven of which were generated from the 60 m
samples, including six belonging to the phylum Thermoplasmatota and one belonging to
Nitrosopumilus (Thaumarchaeota). The deep metagenomes yielded two MAGs classified
as members of the superphylum Patescibacteria, part of the Candidate Phyla Radiation,
one of which contained a 16S rRNA gene SV classified as *Ca*. Uhrbacteria. The only
archaeal MAG from the deep samples was classified as a member of the order
Woesearchaeota. This MAG contained a 16S rRNA gene with 100% nucleotide identity
to the most abundant SV recovered in amplicon sequencing of the bottom water
communities (Table 1, Table S3). Queries against the Genome Taxonomy Database
(GTDB; using MAG single-copy core genes) and SILVA database (using the 16S rRNA
gene SV) classified this MAG as belonging to the order "Woesearchaeia" in the phylum
Nanoarchaeota. However, this classification is outdated, as "Woesearchaeia" has
recently been given the phylum-level designation Woesearchaeota (26). Only nine MAGs
had a reference genome in GTDB that exceeded the minimum alignment fraction (65%);
the remainder had no known close relative (Table 1).
274
Three MAGs representing the highest amplicon frequencies (from all recovered MAGs)
were placed in phylogenies with closely related taxa, confirming their taxonomic
assignments (based on GTDB and SILVA) as members of the Woesearchaeota (Fig. 4),
Nitrosopumilus (Fig. S4), and *Thioglobaceae* (SUP05 clade) (Fig. S5). The two taxa most
closely related to the Woesarchaeotal MAG are both from oxygen minimum zones in the
Arabian Sea, with the next most closely related taxa from an iron-rich terrestrial hot spring

and a freshwater lake (Lake Baikal). The phylogeny shows Woesearchaeotal MAGs from
a wide range of environments interspersed across branches, providing weak support for
clustering according to habitat type (Fig. 4). The AJ *Nitrosopumilus* MAG was most
closely related to a MAG from a cold seep sponge, and relatively distantly removed from
the nearest cultured *Nitrosopumilus* isolate (Fig S4). The AJ SUP05 MAG was most
closely related to a clade of eight *Thioglobacaeae* spp. MAGs from hydrothermal vents,
and next-most closely related to genomes associated with deep-sea invertebrates (Fig.
S5).
289
290 *Functional annotation*
291
Based on analysis of all metagenome contigs (prior to MAG binning), broad functional
categories (KEGG 'subgroup2' level) separated the 60 m samples from both deep
samples (Fig. S6). Among the categories with a 5-fold or greater difference in frequency
between the sample sets was the KEGG category 'ECM-receptor interaction,' which
contained only two annotated genes (Fig. S7). Notably, a single gene (KO number
K25373, annotated as the eukaryotic protein dentin sialophosphoprotein) occurred at
much higher frequency in the deep samples relative to the 60 m samples. BLASTP
queries against the NCBI nr database linked this sequence to hypothetical proteins from
genomes of the candidate phylum Uhrbacteria (CPR), although these proteins shared
only 30% amino acid identity with the gene in our data. This gene was identified in one of
the two MAGs (BH28) classified as Patescibacteria (CPR), to which the Uhrbacteria
belong. Other categories driving the sample separation included ethylbenzene
degradation, bacterial chemotaxis, and o-glycan biosynthesis (Fig. S6).
305
Although they were not among the most differentiating categories, sulfur and nitrogen
metabolism categories also differed in frequency between the depths (Fig. 5). These
included genes for both assimilatory and dissimilatory metabolism. Some of the most
pronounced differences in sulfur metabolism involved the dissimilatory thiosulfate
(*phs*A/*psr*A) and sulfite (*dsr*AB) reductases, both of which were enriched in deep
metagenomes. Sulfur metabolism genes enriched in the 60 m metagenomes also
included *dmd*ABCD, which encode enzymes for the catabolism of
dimethylsulfoniopropionate (DMSP).
314
Blue hole MAGs encoded a diverse suite of metabolisms (Table 2). We detected genes
involved in both reductive and oxidative pathways of dissimilatory sulfur and nitrogen
cycling, arsenic respiration, methylotrophy, and carbon monoxide metabolism, among
others. One MAG (BH25) is one of only two reported members of the Bacteroidota
(formerly Bacteroidetes) phylum potentially capable of dissimilatory sulfur metabolism,
with the other represented by a MAG from a hot spring (27). Most MAGs (23 out of 31)

contained genes involved in arsenic resistance. Genes for arsenic respiration were detected in three MAGs, with two occurrences of arsenite oxidase gene *aio*A (BH16, Actinobacterial order *Microtrichales*, and BH24, alphaproteobacterial order *Rhodospirillales*) and one of arsenate reductase gene *arr*A (BH30, Desulfobacterota taxon *NaphS2*) (Fig. S8). The latter had 82% amino acid identity with a gene from another *NaphS2* strain isolated from anoxic sediment in the North Sea (DSM:14454); this gene is annotated in the FunGene database as *arr*A. Other MAG-affiliated proteins included nitrous oxide reductase (*nos*Z, in four MAGs representing three phyla), carbon monoxide dehydrogenase (*coo*FS, in BH30 (*NaphS2*, phylum Desulfobacterota)), and the ammonia and methane monooxygenases *amo*AB (in BH19, *Nitrosopumilus*) and *pmo*AB (in BH9 (family *Methylomonadaceae*, Gammaproteobacteria) (Table 2). BH9 also contained several other genes involved in C1 metabolism including *mxa*D, *mch*, *mtd*AB, and *fae* (data not shown).

The Woesearchaeotal MAG (BH21) was notable for its small size (679 Kbp, with 67% (CheckM) and 73% (anvi'o) completion values). As with other Woesearchaeotal genomes, genes for several core biosynthetic pathways were missing, including those for glycolysis/gluconeogenesis, the citric acid cycle, and the pentose phosphate pathway. Genes linked to glyoxylate and glycarboxylate metabolism and fructose and mannose metabolism were detected, along with the carbamate kinase *arc*C; however, no full energetic pathway could be reconstructed.

Only one MAG (BH24, order *Rhodospirillales*) contained genes for the full denitrification pathway ($NO_3 \rightarrow N_2$). Other MAGs had the potential to perform individual steps: $NO_3 \rightarrow NO_2$, seven MAGs; $NO_2 \rightarrow NO$, six MAGs; $NO \rightarrow N_2O$, six MAGs; and $N_2O \rightarrow N_2$, three MAGs (Table 2).

Microscopy and cell counts

Microscopy-based cell counts (prokaryotes) ranged from 5×10^6 to 7×10^6 cells/mL within the hole (data available only from September), almost an order of magnitude lower than counts above the hole (30 m sample; Fig. S9). Counts were lowest at 80 m, the depth of the observed turbidity spike.

Discussion

Oxygen-deficient marine water columns are crucial habitats for understanding ecosystem function under oxygen limitation and across gradients of redox substrates, representing conditions that are predicted to expand substantially in the future (25). Moreover, these systems have been a critical resource for discoveries of novel microbial diversity and

unforeseen linkages between chemical cycles (19–24). While in recent years these
discoveries have been facilitated by community DNA, RNA, and protein sequencing, most
oxygen-depleted waters have not yet been characterized, either from an -omics
perspective or via cultivation-dependent methods. This is due partly to the fact that these
systems are challenging to sample and span a gradient of environmental conditions. For
example, the Pacific oxygen minimum zones (OMZs) are anoxic through several
hundreds of meters of the water column, cover hundreds of square kilometers of open
ocean, and are relatively unaffected by processes in the underlying sediment (28).
Microbial communities in these systems, especially those along the peripheries,
presumably have periodic exchange with microbial communities outside the OMZ, for
example via eddy intrusion, storms, or offshore transport (20). In contrast, we describe a
very different oxygen-deficient zone with intense but apparently stable stratification. We
show that the semi-enclosed blue hole environment exhibits a physiochemical profile and
microbial community both similar to but also remarkably distinct from that of other oxygen-
deficient water columns.
376
Amberjack Hole is characterized by an oxygen profile unlike that of most OMZs, with a
secondary oxygen peak around 75 m where dissolved oxygen rose to 43% saturation
before dropping back to zero (Fig. 1). In open ocean OMZs, oxygen profiles are typically
unimodal, with concentrations falling along an upper oxycline, staying hypoxic or anoxic
through a core layer, and then gradually increasing below the core as organic substrates
are depleted with depth and microbial respiration slows (29,30). In AJ, the secondary
oxygen peak at 75 m represents either a decline in net oxygen consumption driven by
decreased microbial respiration, a transport-related phenomenon that affects oxygen
supply, or both. Exchange of water between the offshore aquifer and blue hole water
column could affect oxygen dynamics within the hole. However, the water discharging
from the aquifer would be either saline connate water or saline water traveling long
distances through the aquifer (13,31,32). As such, it is unlikely that oxygen would remain
at levels to increase oxygen to 43% saturation. Alternatively, oxygen may come into the
hole by the sinking of dense (high σ_T, Fig 1B) oxygenated marine waters that intrude onto
the shelf. Such an event could occur episodically during storms but has not been
observed for this system. Turnover via severe cooling of overlying waters, or potentially
upwelling and spilling of water from off the shelf, is presumably rare, and turnover by
storm events unlikely. Indeed, preliminary calculations, assuming only wind-driven mixing
and based on the potential energy anomaly (33), suggest that full turnover of the AJ water
column would require wind speeds of 488 mph blowing for 36 hours in conjunction with 2
m/s surface current velocities in the same direction (Navid Constantinou, Australian
National University, personal communication, February 2020). Thus, the observed
stratification and oxygen profile may be consistent across seasons and result in a stable
bottom layer that is biologically isolated from the surrounding pelagic environment.

AJ Hole is also characterized by a subsurface maximum in NO_3^- (nitrate) and/or NO_2^-
(nitrite) below the main chemocline (Fig. S2), suggesting that dissolved oxygen is also
consumed by nitrification. The decline in NO_x coincides with the second sharp drop in
oxygen concentration around 80 m. This drop is followed by the detection of reduced
inorganic compounds, consistent with decreasing redox free energy expectations as a
function of depth. The presence of $\Sigma S(-II)$ below 80 m suggests active SO_4^{2-} reduction.
Although we detected genes potentially supporting sulfate reduction (e.g., *dsrAB*, *apr*AB),
we cannot rule out the potential that the encoded proteins instead catalyze oxidative sulfur
metabolism. Indeed, preliminary evidence indicates that $\Sigma S(-II)$ was generated at
millimolar concentrations in the sediments based on pore water profiles, and benthic
lander chamber sediment flux measurements in fact show that the sediment-derived $\Sigma S(-$
$II)$ fluxes (on the order of 10 mM m^{-2} day^{-1}) are comprised entirely of acid-volatile,
purgeable hydrogen sulfide, ΣH_2S (Beckler et al., in prep). Thus, it is likely that the
reduced sulfur in the deep water column largely originates from benthic, rather than water
column, sulfate reduction.

Reduced sulfur is clearly a key energy source in the deepest layers of the blue hole. We
detected high concentrations of sulfur in intermediate redox states, notably S(II) (i.e.
thiosulfate, $S_2O_3^{2-}$) and S(0) (elemental sulfur), overlying a pronounced zone of S(-II)
(hydrogen sulfide) below 80 m. Generally, oxidative sulfur metabolism proceeds with
H_2S/HS^- being oxidized to sulfite and sulfate (SO_3^{2-}/SO_4^{2-}) with S(0) and $S_2O_3^{2-}$ produced
as intermediates (34,35), with the relative completion depending on the pH or on the
composition of the microbial community. Here, the peak concentration of $S_2O_3^{2-}$ was
observed at a depth above that of the peak concentration of S(0), suggesting that different
steps in the overall oxidation of $\Sigma S(-II)$ may be performed by various vertically-distributed
niches within a complex cycle. We argue that if a single microbial population was
performing complete H_2S/HS^- oxidation to SO_4^{2-} (with S(0) or S(+II) as intermediates),
these compounds should exhibit similar depth distributions, not stratification.

The zone of peak reduced sulfur concentration (peak concentrations of all measured
sulfur compounds) occurred below 80 m (Fig. 1D) and appeared vertically decoupled from
detectable dissolved oxygen, which fell below detection between 74 and 79 m (Fig. 1B).
This decoupling suggests the anaerobic sulfur oxidation may also proceed with NO_x as a
terminal oxidant, as NO_x remained detectable above 90 m and decreased in
concentration with depth as reduced sulfur concentrations increased (Fig. 1). Although
representing only the deepest, most sulfidic layers of the blue hole, metagenomes
contained diverse genes for oxidizing sequentially reduced sulfur compounds, as well as
high abundances of genes for each step of the denitrification pathway (Fig. 5). Sulfur-
driven denitrification is common in marine OMZs (e.g. (36)), notably in conjunction with

gammaproteobacteria of the SUP05 lineage (e.g. (37)). In some cases, we detected both sulfur oxidation and denitrification genes in the same MAG, including in the SUP05 (*Thioglobaceae*) MAG (BH20). We also recovered MAGs encoding only sulfur oxidation proteins or incomplete denitrification pathways (Table 2), a pattern observed in other low oxygen water columns and suggesting the likely cross-feeding of sulfur and nitrogen cycle intermediates between taxa (e.g., (38,39)). However, the zone of stratification of sulfur intermediates, roughly 70-90 m, overlaps with the zone of oxygen availability (down to 80 m) as measured with the fluorometric sensor, which has a detection limit only in the low micromolar range. Thus, it is likely that the community in this depth range consumes both oxygen and oxidized nitrogen species (and potentially other oxidants) for use in sulfur oxidation, with the relative concentrations of oxidants and sulfur intermediates potentially driving the vertical separation of microbial niches.

Interestingly, the oxic-anoxic transition at 80 m coincided with a sharp spike in turbidity. This spike was not due to an increase in microbial cell counts (Fig. S9). However, at 80 m, we recorded a sharp increase (to 10% of total) in the frequency of 16S rRNA genes assigned to *Arcobacter, a* sulfide-oxidizing bacterium known to produce filamentous sulfur at oxic-anoxic interfaces (40). Between 80 and 90 m, 70-83% of the combined elemental sulfur (S(0)) and polysulfide (ΣS(-II)) signal was lost after 0.7 µm filtration, suggesting that this lost fraction was in fact elemental sulfur, as polysulfide is soluble (41,42). We did not sample at 80 m in May and therefore cannot confirm if the sharp turbidity and *Arcobacter* spikes are stable over time. However, the production of filamentous or other particulate forms of sulfur under microaerophilic conditions is consistent with the observed sulfide and oxygen gradients in both months (Fig. 1, Fig. S1). Alternatively, *Arcobacter* may be using NO_x to oxidize S(-II), as has been shown in denitrifying members of the genus (43).

Other oxidants may also play a role in AJ. Dissolved ferrous iron (Fe(II)) reached a maximum concentration near the turbidity spike at ~80 m, below the zone of peak O_2/NO_x and above the zone of reduced sulfur (Fig. 1D). Further, total dissolved iron (Fe_d), which includes dissolved organically-stabilized Fe(III) and FeS colloids, remained < 300 nM, but increased below 40 m, peaking just above the dynamic reduced sulfur zone (Fig. 1D). These patterns hint at a possible cryptic iron cycle, in which ΣH_2S diffusing upwards is rapidly oxidized by dissolved Fe(III) (44) to form Fe(II), S(0), and ΣS(-II). Dissolved Fe(III) may then be recycled by reoxidation of Fe(II) by O_2, and the S(0)/S(-II) simultaneously oxidized by O_2 or dissolved Fe(III) to form thiosulfate, $S_2O_3^{2-}$. Alternatively, ΣH_2S may first react with Fe(II) to form the detected FeS colloids, which subsequently oxidize to form Fe(II), S(0), and S(-II). Metabolic pathways for iron oxidization (or reduction) are not easy to detect with metagenomic data, as iron metabolism genes play roles in other cellular processes and are therefore suggestive but not diagnostic of iron metabolism (45,46). Nonetheless, additional metagenomic sampling at finer spatial resolution may help

identify a role for microbial activity in blue hole iron cycling. The oxidation of iron, however,
is rapid even abiotically, and the vertical distribution of iron in this system can partially
explain the apparent vertical disconnect between the $\Sigma S(-II)$ and O_2. Indeed, this
phenomenon has been observed in a nearshore Florida hole (6), in which FeS colloids
formed at the interface between 30 and 50 m. AJ appears to be enriched instead in
colloidal S(0), with only nanomolar concentrations of FeS colloids.
487
The pronounced stratification of blue hole chemistry is consistent with differentiation of
microbial communities with depth. In both May and September, microbiome composition
varied distinctly among oxic, hypoxic, and anoxic, sulfidic layers (Fig. 2). Shannon
diversity was similar among all communities, but Simpson diversity was notably higher in
the anoxic depth group (Fig. S3), reflecting the lower evenness of these microbiomes.
Surprisingly, the deepest communities were dominated by a recently described archaeal
lineage, the Woesearchaeota (47). Woesearchaeota have been detected in a wide variety
of biomes, including groundwater, terrestrial and marine sediments, wetlands, deep-sea
hydrothermal vents, and hypersaline lakes (48). However, their relative abundance is
consistently low, at most ~5% of the total microbial community in any given environment,
with the highest proportions observed in freshwater sediments (48) and high-altitude
lakes (49). In contrast, Woesearchaeota comprised at least one third of the blue hole
microbiome between 75 and 106 meters, reaching a maximum of nearly 60% in the anoxic
sulfidic layer in September (Fig. 2, Fig. 3). Remarkably, one 16S rRNA sequence variant
represented up to 97% (and only two SVs represented up to 99%) of all Woesearchaeotal
amplicons in each sample, suggesting low intrapopulation diversity. Dominance (>50% of
the community) by a single strain variant is relatively uncommon in pelagic marine
microbiomes; it is even more rare that such occurrences involve members of the Archaea.
The AJ Woesearchaeotal MAG, which contained the dominant 16S rRNA
Woesearchaeota amplicon sequence, was 679 Kbp and similar in size to other
Woesearchaeotal genomes, which rarely exceed 1 Mbp (47,48). This MAG was most
closely related to MAGs from an OMZ in the Arabian Sea (Fig. 5), suggesting the potential
for a Woesearchaeotal clade specific to a pelagic, marine, low oxygen niche. At around
70% completion, the AJ MAG did not contain genes of metabolic processes common to
marine low oxygen systems, notably dissimilatory sulfur or nitrogen cycling, anaerobic
metabolism, or autotrophy. Indeed, consistent with characterizations of other
Woesearchaeotal genomes (47,48), AJ Woesearchaeota appear to have limited
metabolic capabilities. It is therefore possible that these cells are partnered in a syntrophic
relationship with other microbes on which they rely for energy or nutrients, as has been
previously suggested (47,48).
518
519 The Woesearchaeotal MAG was one of several novel lineages in the AJ water column.
520 From the deepest samples (95 m and 106 m), recovered MAGs included members of the

sulfur-oxidizing SUP05 clade of *Thioglobaceae*, two members of the CPR phylum Patescibacteria, and three members of the phylum Marinimicrobia (SAR406), among others (Table 1). The middle water column (60 m) was also populated by underdescribed taxa, including several lineages of marine Thermoplasmatota and members of the bacterial phyla Planctomycetota and Myxococcota. Only eight out of nineteen MAGs from 60 m had identifiable close representatives in the GTDB, suggesting a high level of taxonomic novelty. Surprisingly, automatic binning did not recover a Thaumarchaeal MAG from the 60 m assembly despite the large fraction of amplicon SVs belonging to the ammonia-oxidizing genus *Nitrosopumilus*. However, a manually binned MAG could be phylogenomically placed in this lineage (Fig. S4) and contained the dominant SV from the amplicon data set, which comprised up to 86% of all *Nitrosopumilus* spp. amplicons. As with the Woesearchaeota, this low level of SV diversity implies population homogeneity. It remains to be determined if this homogeneity is driven by a potential dearth of ecological niches. Indeed, the AJ *Nitrosopumilus* MAG, at 725 Kbp, was estimated to be only 53-63% complete (Table 1, Table S3); its full biochemical potential therefore remains to be characterized. However, in contrast to recently described and putatively heterotrophic Thaumarchaeota (50,51), blue hole *Nitrosopumilus*, like all known members of this genus, contain *amo*CAB encoding ammonia monooxygenase and therefore likely contribute to nitrification in the blue hole.

The 60 m samples also contained six MAGs belonging to the archaeal phylum Thermoplasmatota. One of these (BH15) could only be placed in the family Thalassoarchaeaceae. The others included four (BH4, BH11, BH12, and BH13) from the Marine Group II lineage, which is one of the four major planktonic archaeal groups (52), and one from the less well-characterized Marine Group III lineage (BH3). The Marine Group II have recently been proposed as an order-level lineage, *Candidatus* Poseidoniales, containing two families delineating the current MGIIa and MGIIb clades (53). Most MGIIa members have been identified from the photic zone, while MGIIb sequences are largely limited to depths below 200 m, although there are exceptions (54). Notably, all AJ MGII MAGs belonged to the MGIIb lineage (Table 1). Several studies have identified genes related to low-oxygen metabolism including reduction of sulfate (55–57) and nitrate (53), suggesting these taxa are more likely to be adapted to microaerophilic or anaerobic conditions. We found a gene for thiosulfate reduction to sulfite (*phs*A) in two of the four MGIIb MAGs (Table 2), but no complete pathways for any forms of anaerobic respiration. Nevertheless, the Marine Group II lineage comprised over 25% (five out of nineteen) of the MAGs recovered from 60 m, suggesting this group as an important component to the hypoxic zone community.

Metabolic potential differentiated the middle and deep water column communities (Fig. 5, Fig. S6). Many of these functions could be linked to MAGs and represented a range of

anaerobic metabolisms. Genes for dissimilatory sulfide and sulfite oxidation (or potentially
reduction; *dsr*AB, *apr*AB), denitrification (*nar, nir, nor, nos*), and dissimilatory nitrate
reduction to ammonia (DNRA, *nrf*) were enriched in deep anoxic samples, but also
common in the hypoxic zone (60 m) (Table 2, Fig. 5). Other genes linked to MAGs
included *aio*AB and *arr*A, encoding arsenite oxidase and arsenate reductase, respectively
(Fig. S8). Reduction of arsenate to arsenite is thought to be an ancient metabolic pathway,
originating before oxygenation of the Earth's atmosphere and oceans (58). It has recently
been proposed as an important microbial metabolic strategy in oxygen-deficient marine
environments, potentially providing arsenite for use as an energy source by other
microbes (19). Arsenic concentrations in Amberjack have not been measured. However,
the detection of *aio*AB and *arr*A, of which the latter was also detected in unbinned contigs
from the deep co-assembly (Fig. S8), as well as genes for arsenic detoxification (*ars*CM)
in diverse MAGs (including in the *Nitrosopumilus* MAG described above), suggest a role
for arsenic cycling in AJ. Remarkably, one MAG (BH24, *Rhodospirillales*) contained
genes for several of these pathways, representing an unusual repertoire of respiration
strategies (Table S4). Finally, one of the genes with the largest differences in frequency
between 60 m and deep samples encodes an unknown protein previously observed in
genomes of *Candidatus* Uhrbacteria, one of the groups within the CPR/Patescibacteria
(Fig. S7). Two MAGs from this group, which is largely affiliated with freshwater and
subsurface aquifer environments, were recovered from the deep samples, and may play
as-yet undetermined ecological roles in this unusual microbiome. This hypothesis is
supported by the low sequence identity of this protein to similar ones in NCBI.

Conclusions

The Amberjack Hole water column is potentially highly stable, with complete turnover
(driven either by wind or the sinking of cold water masses) unlikely. Thus, AJ
microbiomes, particularly those in the deepest layers, potentially have limited connectivity
to other marine communities. The extent to which such isolation explains the observed
unique community composition or drives divergence of individual microbial lineages
remains uncertain. Undoubtedly, the vertical transport, transformation, and stratification
of redox active elements - notably sulfur, nitrogen, and iron - also play a significant role
in structuring this unusual microbiome. The presumed stability of this community, its high
metabolic diversity, and its dominance by understudied microbial taxa, highlight AJ as a
model for detecting novel biogeochemical processes under low oxygen. Marine blue
holes in general are potentially valuable sites for the study of microbial diversification and
linked elemental cycling.

Methods

Sampling scheme

Sampling was conducted May 15-17 2019 and September 19 2019 using the vessels *R/V* William R. Mote and *R/V* Eugenie Clark. Amberjack Hole is located approximately 50 kilometers west of Sarasota, Florida, at 27.28748 N, -83.16139 W. One component of the research involved the deployment of an autonomous benthic lander to the bottom of the hole for 24-48 hours at a time for benthic electrochemistry measurements; these data are not included in this paper. Technical SCUBA divers who guided the lander into and out of the hole also collected water samples (see details below).

In both May and September, all water samples used for nutrient measurements and microbial DNA preservation were filtered immediately on board. Microbial DNA samples were stored on ice until the return to Mote Marine Laboratory, where they were stored at -20°C. Subsequent nutrient analyses described below were conducted at Mote except where noted.

Water column physical parameters and dissolved oxygen

During the May sampling, an EXO2 sonde (YSI Inc/Xylem Inc, Yellow Springs, Ohio) was attached to the autonomous lander deployed for 24 hours to the bottom of the hole. The sensor provided extensive data from the surface (1-2 m) and the bottom position (106 m) but low-resolution readings from the water column due to rapid ascent and descent velocities. Dissolved oxygen, temperature, and salinity data were retrieved from the instrument and plotted in R.

During the fall sampling, an integrated SBE-19plus V2 CTD with an SBE43 DO sensor, WET Labs FLNTUrt chlorophyll and turbidity sensor, Satlantic cosine PAR sensor, and an SBE18 pH sensor (Sea-Bird Electronics Inc) was lowered twice by hand at a rate that provided high-resolution dissolved oxygen, turbidity, salinity, and density data on September 19 2019. CTD casts were performed the same day as water was sampled for microbial DNA and nutrients. Data were retrieved and plotted in R as above.

SBE data were processed using SBE Data Processing software after compensating for sensor thermal mass, sensor alignment, timing offset (the delay associated with pumped sensors, i.e. conductivity), and changes in instrument velocity (due to ship heave). Data were binned (~0.2 m increments) to smooth data variability. Processed and derived data included temperature (°C), potential temperature (°C), salinity (psu), water density (σ_T in kg/m^3), chlorophyll *a* (µg/L based on relative fluorescence), turbidity (NTU), dissolved oxygen (mg/L), O_2 saturation (%), pH, PAR, and Brunt-Väisälä frequency (N =

stratification index). Manufacturer recommendations for calibration and service for all
sensors were followed.

Water column electrochemical measurements

Water samples (~10 m vertical resolution) were subject to solid-state Hg/Au voltammetric
analyses (59) for measurement of the redox environment, i.e. O_2, Mn(+II), organic-Fe(III)
complexes, Fe(II), S(+II) (in the form of $S_2O_3^-$), S(0), S_x(-II), and ΣH_2S (S^{2-}, HS^-, H_2S).
Briefly, samples collected from the water column at 10 m intervals were carefully
transferred via a Tygon (formula 2375) transfer tube line into LDPE bottles while filling
from bottom to top to minimize atmospheric oxygen contaminations. Sample bottles
remained sealed until analyses and were stored at 4ºC. Upon return to the lab (within 4-
6 hours of collection), 20 mL of each sample was carefully pipetted into an
electrochemical cell (Analytical Instrument Systems, Inc.) holding a custom fabricated
Hg/Au amalgam microelectrode working electrode (60), a Pt counter electrode, and a
fritted Ag/AgCl reference electrode (with 3 M KCl electrolyte solution). N_2 was gently
blown over the top of the solution to minimize mixing with the atmosphere. Within 5
minutes, samples were subject to a series of anodic square wave voltammograms
(ASWVs), cathodic square wave voltammograms (CSWVs), and linear sweep
voltammograms (LSVs) to quantify the above redox analytes (59,61). To analytically
distinguish between S(0) and S(-II), which both react at the same potential at an Hg
electrode (42), CSWV measurements were repeated after acidification with HCl (final pH
< 3) and N_2 sparging of the solution for two minutes to remove free ΣH_2S. This same
electrochemical speciation experiment was conducted a second time in separate aliquots
that were filtered through 0.7 µm GFF filters directly in the N_2-degassed electrochemical
cell (this filter size was selected for future organic carbon analyses).

Nutrient, iron, and DIC measurements

Water samples were collected for nutrients and included chlorophyll *a*, dissolved
ammonium (NH_4^+), dissolved NO_x (nitrate (NO_3) + nitrite (NO_2)), dissolved
orthophosphates (PO_4^{3-}), dissolved total and ferrous iron (Fe), dissolved inorganic carbon
(DIC) and particulate carbon (C), nitrogen (N), and phosphorous (P). All samples were
collected and immediately filtered once shipboard.

Chlorophyll *a* was measured according to EPA method 445.0 (62). Briefly, samples were
filtered through a glass fiber filter until clogging, then stored in the dark until analyses.
Prior to analyses, filters were sonicated in 90% acetone to extract chlorophyll from algal
cells and then centrifuged for clarification. The fluorescence of the clarified extract was
then measured using a fluorometer (Turner 10-AU Fluorometer) with special narrow

bandpass filters at an excitation wavelength of 436 nm and an emission wavelength of
680 nm. Analytical quality assurance followed standard laboratory practices assessing
precision and accuracy using sample replicates, container blanks, duplicate and spiked
analyses with results meeting acceptable levels of precision and accuracy.
684
Samples for dissolved NH_4^+ and NO_x were filtered through Pall Supor (PES) 450 47 mm
(0.45 μm pore size) membrane filters while samples for particulates were filtered through
pre-combusted (450°C, 3h) 47 mm (0.7 μm pore size) Whatman GF/F filters. For
particulate carbon and nitrogen, 200 mL was filtered per sample and filters were rinsed
with acidified (10% HCl) filtered seawater to remove inorganic carbon. For particulate P,
500 mL was filtered. Samples were then stored on ice until their return to the lab and
analyzed within 48 hours (dissolved NH_4^+ and NO_x) or within 28 days (particulates).
Analyses for dissolved nutrients followed colorimetric, segmented flow, autoanalyzer
techniques on an AA3 with method reference and method detection limits (MDL) as
follows: dissolved NH_4^+ ((63); 0.07 μM), dissolved NO_x ((64); 0.07 μM). Particulate
phosphorus was analyzed according to (65,66) with in-house modifications for analysis
on a segmented flow analyzer and MDLs of 0.03 μM. For particulate carbon and nitrogen,
samples were analyzed on a Thermo FlashEA1 1112 Elemental Analyzer with MDLs of
0.2 and 0.1 μM, respectively.
699
700 To measure DIC, samples were immediately poisoned with $HgCl_2$ and stored until
701 analyses. DIC was analyzed (Apollo AS-C6 DIC Analyzer) following methods by (67).
702 Accuracy and precision of the instrument was regularly monitored using Certified
703 Reference Materials for Seawater CO_2 Measurements (Dickson Laboratory, Scripps
704 Institution of Oceanography, San Diego, CA; Batch #181 and 186).
705
706 Soluble orthophosphates (PO_4^{3-}) were measured spectrophotometrically using the
707 molybdate-blue technique (68).
708
Finally, the speciation of iron was obtained by measuring Fe(II) by the ferrozine assay
(69) in filtered samples before (dissolved Fe(II)) and after (total dissolved Fe) reduction
by hydroxylamine (0.2 M) using a long waveguide spectrophotometric flow cell. Analytical
quality assurance follows standard laboratory practices assessing precision and accuracy
using sample replicates, container blanks, duplicate and spiked analyses with results
meeting acceptable levels of precision and accuracy.
715
716 *Microbial DNA sampling and preservation*
717
Water column samples were collected on May 15 and 17 2019 and September 19 2019.
In May, 800 mL water was collected by divers from depths of 46 m, 61 m, and 106 m

inside the hole. Sterilized Nalgene bottles filled with deionized water were taken down by
the divers, opened at the appropriate depth, and DI water was replaced with ambient
seawater. Niskin bottles deployed on the CTD rosette also collected 1.9 L samples from
depths of 8 m, 15 m, and 23 m above the hole. In May and September, hand-cast Niskin
bottles collected 1 L water from 0 m (surface), 30 m, 60 m, and 85 m (May), and from 0
m (surface) to 90 m at 10 m intervals, as well as 85 m and 95 m (September). Samples
were filtered onto 0.22 µm Sterivex (MilliporeSigma) filters with a peristaltic pump and
preserved with approximately 3 mL DNA/RNA stabilization buffer (25 mM sodium citrate,
10 mM EDTA, 5.3 M ammonium sulfate (pH 5.2)) and stored at -80°C or on dry ice until
further processing at Georgia Tech. In September, unfiltered water from the following
depths was also preserved (1.4 mL water, 150 µL PBS-buffered formaldehyde) for nucleic
acid staining and microscopy and kept frozen at -20ºC until processing: 30 m, 50 m, 60
m, 70 m, 80 m, 85 m, 90 m, 95 m.
733
734   *Sequencing library preparation*
735
DNA was extracted from each Sterivex cartridge using a custom protocol as described in
(70), except for the September surface (0 m) sample which was lost during sample transit
to Georgia Tech. Briefly, cells were lysed by flushing out RNA stabilizing buffer and
replacing it with lysis buffer (50mM Tris-HCl, 40 mM EDTA, 0.73 M sucrose) and lysozyme
(2 mg in 40 mL of lysis buffer per cartridge), then incubating cartridges for 45 min at 37°C.
Proteinase K was added and cartridges were resealed and incubated for 2 h at 55°C. The
lysate was removed, and the DNA was extracted once with phenol:chloroform:isoamyl
alcohol (25:24:1) and once with chloroform:isoamyl alcohol (24:1). Finally, DNA was
concentrated by spin dialysis using Ultra-4 (100 kDA, Amicon) centrifugal filters. Yield
was assessed using a Qubit 2.0 dsDNA high-sensitivity assay (Invitrogen, Carlsbad, CA).
746
Illumina MiSeq libraries were prepared by amplifying the V4 region of the 16S rRNA gene
using the environmental DNA protocol adapted from (71). Briefly, amplicons were
generated using Platinum® PCR SuperMix (Life Technologies, Carlsbad, CA) with Earth
Microbiome Project primers 515FB and 806RB appended with Illumina-specific adapters.
Template DNA was diluted to approximately 5 ng/µL for all samples and PCRs were
performed in 25-µL reactions using Platinum® PCR SuperMix (Life Technologies) (22 µL),
BSA (Invitrogen) (1 µL), and 0.5 µL each of forward and reverse primer (10 ng/L stock
concentration) with 1 µL template DNA. The thermal cycling protocol consisted of 26
cycles with the following steps: denaturation at 98°C (30 s), followed by 30 cycles of
denaturation at 98°C (5 s), primer annealing at 55°C (5 s) and primer extension at 72°C
(8 s), followed by extension at 72°C for 1 minute. Negative control reactions were run
using 1 µL milliQ water in place of DNA template. Amplicons were analyzed by gel
electrophoresis to verify size (~400 bp, including barcodes and adaptor sequences) and

purified using Diffinity RapidTip2 PCR purification tips (Diffinity Genomics, West Chester,
PA). Amplicons from different samples were pooled at equimolar concentrations and
sequenced using a paired-end Illumina MiSeq 500 cycle kit (2 x 250 bp) with 5% PhiX to
increase read diversity.
764
Metagenomes were generated from four water column samples: two from May (60 m and
106 m) and two from September (60 m and 95 m). Libraries were prepared using the
Illumina Nextera XT DNA library preparation kit (Illumina Inc., San Diego, CA) according
to manufacturer's instruction and run on a Bioanalyzer 2100 instrument (Agilent) using a
high sensitivity DNA chip to determine library insert sizes. An equimolar mixture of the
libraries (final loading concentration of 12 pM) was sequenced on an Illumina MiSeq
instrument (School of Biological Sciences, Georgia Institute of Technology), using a
MiSeq reagent v2 kit for 600 cycles (2 x 300 bp paired end protocol).
773
774　*Amplicon sequence data processing and analysis*
775
Demultiplexed amplicon sequences are available in the Patin FigShare account:
https://figshare.com/projects/Amberjack_Blue_Hole/85013. Raw sequences were run
through the DADA2 algorithm (72) in QIIME2 (73) to assess sequences at sequence
variant resolution, using the following command parameters: --p-trim-left-f 70 --p-trim-left-
r 70 --p-trunc-len-f 150 --p-trunc-len-r 150. Resulting SVs were assigned taxonomy using
the naïve Bayes classifier trained on the Silva 132 database (99% OTUs, 515F/805R
sequence region).
783
The SV table with raw read counts of all water samples were exported from QIIME2 and
transformed in R using the variance stabilizing transformation (vst) in the DESeq2
package (74). Tables were not rarefied to preserve maximum available information.
Metadata including depth and SV taxonomy were imported and combined into a Phyloseq
object (75,76). Alpha and beta diversity analyses were run in DivNet (77) at the SV level.
Shannon and Simpson diversity results were extracted and plotted by depth grouping
using ggplot2 (78). Beta diversity was assessed using the resulting Bray-Curtis distance
matrix in a principal component analysis using the prcomp() function in R and visualized
in a PCA using ggplot2. The distance matrix was also used to test for significant difference
among samples by depth grouping and month using the adonis() function in the R vegan
package (79) with the following command: adonis(formula = bc ~ Depth + Month, data =
metadata, permutations = 999).
796
797　*Metagenomic sequence data processing and analysis*
798

Metagenome sequences are available in the Patin FigShare account:
https://figshare.com/projects/Amberjack_Blue_Hole/85013. Raw sequences from the four
metagenomes (May 60 m, May 106 m, September 60 m, September 95 m) were trimmed
and checked for quality and adapter contamination as described for the amplicons above.
Quality controlled reads were run through MicrobeCensus (80) to generate genome
equivalent (GE) values for each metagenome. To generate more even coverage
distribution across the metagenomes, quality controlled reads were normalized using
BBNorm, part of the open source BBMap package
(https://sourceforge.net/projects/bbmap/).

Six assemblies were generated from the four water column metagenomes. Individual
assemblies of each sample were generated using metaSPAdes (v3.14.0) (81). Co-
assemblies for each depth pair (May 60 m with September 60 m, and May 106 m with
September 95 m) were also generated using MEGAHIT (v1.2.9) (82) using a minimum
contig length of 2500 bp. All assemblies were assessed for quality using metaQUAST
(83) and annotated using Prokka (84) and the KEGG database.

For the latter, open reading frames were generated using Prodigal (85) and clustered
using MeShClust (86) at 90% nucleotide identity. The longest sequence from each cluster
was extracted using a custom Python script and these representative sequences were
run against the KEGG ortholog profile HMM models (KOfams) using KofamScan with the
'prokaryote' database (87). The parameter '-f mapper' was applied to provide only the
most confident annotations (those assigned an individual KO). Orthologies were matched
to their corresponding functions using a parsed version of the 'ko00001.keg' database
text file (https://github.com/edgraham/GhostKoalaParser), which provides a three-tiered
hierarchical categorization of each gene, referred to here as 'Group', 'Subgroup1', and
'Subgroup2'. Sequence coverage of each gene was generated by mapping metagenomic
short reads against each one using Magic-BLAST (88). The Magic-BLAST output was
filtered to include only the best hit for each read and the read counts were normalized by
GE value of the corresponding metagenome.

The normalized counts were used to run hierarchical clustering analyses in Python using
the Seaborn 'clustermap' function. KofamScan outputs were grouped by the three
hierarchical categories to generate heat maps at different levels of categorization. At the
highest categorical level ('Group') the categories 'Human Diseases,' 'Organismal
Systems,' 'Cellular community – eukaryotes,' and 'Brite Hierarchies' were removed before
performing the cluster analysis. The first three groups are not relevant to microbial gene
functions and the fourth provides a different hierarchical categorization scheme for the
same annotations and was thus redundant. All cluster analyses were run using ward
linkage and Euclidean distance methods.

Open reading frames generated from the SPAdes individual assemblies were run through
the Hidden Markov Model search tool described in (89) and available at
https://github.com/ShadeLab/PAPER_Dunivin_meta_arsenic/tree/master/HMM_search
to query for genes involved in arsenic metabolism. HMM outputs were run through the
data_preparation.R script (also available in the GitHub repository) to generate plots
showing the quality distributions of hits to the *aio*A and *arr*A genes. The amino acid
sequence of a potential sequence for the *arr*A gene extracted from BH30 (alignment
fraction 89%, bitscore 237) was also queried against the *arr*A BLAST database from the
same toolkit.

Metagenome-assembled genomes

Genomic bins were generated from all individual assemblies as well as co-assemblies.
For each co-assembly, short reads from both metagenomes were submitted to MaxBin
to leverage contig co-variation patterns across samples. All bins were assessed for quality
using CheckM (90) and anvi'o (v6.1) (91). Bins with a quality score greater 40 were
retained for dereplication, with quality score calculated as completion – 5 x contamination
(CheckM values). Dereplication was performed with dRep (92) with an ANI cutoff of 95%
for secondary clustering and a 20% minimum pairwise overlap between genomes. For
bins belonging to the same secondary cluster (ie, likely representing the same microbial
population), the highest-quality was chosen as a representative. In cases where the
qualities of two MAGs were within 1%, the completeness and redundancy values from
anvi'o were applied to determine the higher quality bin. Further analyses were performed
on these representative bins.

All dereplicated high-quality MAGs were assessed for taxonomy with anvi'o and the
Genome Taxonomy Database toolkit (GTDB-Tk) (93,94). Each MAG was checked for the
presence of rRNA genes using the anvi'o HMM models and one MAG from the 60M
assembly was found to contain a eukaryotic 18S rRNA gene with an 80% match to a
copepod sequence (*Calanus* sp.). This contig was removed from the co-assembly of
origin and the process of binning, quality assessment, and de-replication was repeated.

All MAGs were queried for the presence of 16S rRNA gene amplicons by using the SVs
as a query for a BLASTn analysis against the two co-assembly contigs. MAG contigs
containing 100% identity matches across the entire SV length were identified, and the SV
SILVA-based taxonomy was compared with the whole genome taxonomy assignment
from GTDB.

All MAGs were functionally annotated using both Prokka and KofamScan as described
above, without the read mapping step. The following genes were verified by running
amino acid sequences against the NCBI-nr database using blastp: *pmo*A-*amo*A (K10944;
KEGG), *pmo*B-*amo*B (K10955; KEGG), *pmo*C-*amo*C (K10956; KEGG), K23573 (DSPP,
dentin sialophosphoprotein; KEGG), *pmo*A (Prokka), *pmo*B (Prokka), *aio*B (Prokka),
*ars*C2 (Prokka), *ars*M (Prokka). To confirm the annotations of arsenic respiration and
resistance genes, and to query other genes involved in arsenic metabolism, including the
arsenate reductase gene *arr*A, all MAGs were run through the Hidden Markov Model
search   tool   described   in   (89)   and   available   at
https://github.com/ShadeLab/PAPER_Dunivin_meta_arsenic/tree/master/HMM_search.
888
889 *Manual MAG refinement and generation*
890
Six MAGs contained more than one 16S rRNA gene SV. In cases where the SVs within
a MAG were distantly related to each other, any contigs containing a copy that was
divergent from the genome-wide GTDB-TK taxonomic assessment were removed from
the bin. Quality assessments were repeated on the edited MAG using CheckM and anvi'o,
and if the quality score (as defined above) was over 5 points lower than the original MAG,
the edited MAG was discarded. This happened in only one case (BH22); all other MAGs
were kept in their edited form.
898
One of these edited MAGs was BH28, classified by GTDB as Patescibacteria but
containing an SV classified as *Magnetospiraceae* (Alphaproteobacteria, Rhodospirillales)
by SILVA. Conversely, BH24, which was classified by GTDB as Rhodospirillales,
contained no 16S rRNA genes. The contig containing the *Magnetospiraceae* SV was
removed from the Patescibacteria bin and added to BH24 based on its tetranucleotide
clustering proximity to other contigs in that bin (as seen in the anvi'o interactive interface)
and the matching taxonomy between the MAG and the SV.
906
The dominant *Nitrosopumilus* amplicon SV was not present in any of the MAGs generated
from the 60 M samples. However, it was found on an unbinned contig in the 60M co-
assembly. The anvi'o interactive interface was used to manually generate a bin containing
this sequence, along with a 23S rRNA sequence with 99% identity to *Nitrosopumilus*
catalina, and other contigs of similar tetranucleotide frequency, GC content, and coverage
(BH19). This MAG was assessed for quality using CheckM and anvi'o.
913
914 All MAGs were run through GTDB using the 'classify_wf' (for taxonomic classification)
915 and 'ani_rep' (for closest relative by ANI) commands in GTDB-Tk (93), and through
916 Prokka (84) and KofamScan (87) for functional annotation. Closest relatives were

917   determined using a minimum 65% alignment fraction; if no reference genome exceeded
918   this minimum there was no result.
919
920   All MAG sequences are available in the Patin FigShare account:
921   https://figshare.com/projects/Amberjack_Blue_Hole/85013.
922
923   *Phylogenomic analyses*
924
Phylogenomic trees were generated for the MAGs BH19 (*Nitrosopumilus*), BH20 (SUP05
clade, *Thioglobaceae*), and BH21 (Woesearchaeota). In each case, genomes with the
same or similar taxonomy as the MAG were downloaded from the NCBI Assembly
database and run through anvi'o for quality assessment using anvi-estimate-genome-
completeness. High-quality MAGs (quality score > 40 as defined above) were retained
(Table S4) and single-copy genes were extracted using the anvi'o script anvi-get-
sequences-for-hmm-hits. The two archaeal MAGs (BH19 and BH21) were run against the
'Archaea_76' HMM database and the bacterial MAG (BH20) was run against the
'Bacteria_71' HMM database. Genes occurring in most or all MAGs were concatenated
and aligned (see Table S4 for taxon-specific minimum gene occurrences). The alignment
was run through RAxML (95) with optimization of substitution rates and a GAMMA model
of rate heterogeneity ("PROTGAMMA" substitution model) with 999 bootstraps for a
maximum likelihood phylogeny. The resulting .tre files were uploaded to the interactive
Tree of Life website (96) (itol.embl.de) and labeled with their strain or sample of origin.
939
940   *Cell staining and microscopy*
941
Duplicate samples from the following depths were processed for microscopy: 30 m, 50
m, 70 m, 80 m, and 95 m. Samples preserved in PBS-buffered formaldehyde (1.55 mL
total) were combined with 2.0 mL milliQ water and filtered onto 0.2 um GTBP filters
(Millipore) using vacuum filtration. Filters were dried for approximately 20 minutes and
incubated in the dark on ice with 50 µL DAPI (0.2 µg/mL). Filters were then rinsed in milliQ
water and 100% ethanol, dried for approximately 10 minutes, and placed on a microscope
slide with one drop of Citifluor (Electron Microscopy Sciences). Slides were visualized
with a Zeiss Axio Observer D1 confocal epifluorescence microscope using a DAPI filter
(Zeiss filter set 49). Between twenty and twenty-five photographs were taken in a grid
pattern from each slide and counted in ImageJ (97) using a custom script. The average
value of cell counts from all photos for each sample was plotted in Microsoft Excel.
953
954   **Figure legends**
955

Figure 1. The blue hole water column in September 2019 was highly stratified. (A) Compared to the overlying water, salinity was slightly higher and pH was slightly lower inside the hole (i.e. below 30 m). A coincident dip in salinity and rise in pH was present at 75 m. (B) Dissolved oxygen concentrations varied widely, with both a primary and secondary oxycline. At 80 m, the onset of anoxia immediately below the secondary oxycline coincided with a spike in turbidity. Water density is represented by σ_T, defined as $\rho(S,T)$-1000 kg m^{-3} where $\rho(S,T)$ is the density of a sample of seawater at temperature T and salinity S, measured in kg m^{-3}, at standard atmospheric pressure. (C) Dissolved inorganic carbon (DIC) increased slightly from 20 to 50 m but more intensely between 70 and 90 m, from approximately 2.2 to 2.5 mM. A sharp increase in NO_x (NO_2^- + NO_3^-) between 40 and 50 m was followed by a return to near 0 between 60 and 90 m. Phosphate (PO_4^{3-}) and ammonium (NH_4^+) remained below 1 µm before increasing to 5-6 µm between 70 m and 80 m, respectively. (D) Dissolved ferrous iron (Fe(II)$_d$), and total dissolved iron (Fe$_d$) increased with the transition to anoxia. Sulfur species are presented as follows: $S_2O_3^{2-}$ (thiosulfate, S in the +II oxidation state) peaked between 80 and 90 m. S(0) represents combined dissolved and colloidal elemental sulfur; however, the dissolved fraction may also include a small amount of polysulfide species (i.e. S_x(-II)). Finally, $\sum$S(-II) represents primarily hydrogen sulfide (HS$^-$) but could also be minorly redundant with S(0). All iron and sulfur species increased sharply by 70-85 m, with S(0) representing the largest component of the reduced sulfur pool.

Figure 2. Microbial communities represented three water column depth groupings: shallow (0-32 m), middle (40-70 m), and deep (80-95 m). (A) Principal components analysis shows communities were highly similar within each depth grouping regardless of sampling date. (B) Community composition of representative samples from each of the three depth groupings show both middle and deep water column layers feature high levels (~40% frequency) of a single taxon, the ammonia-oxidizing *Nitrosopumilus* sp. in the middle water column and the Woesearchaeota in the deepest layers. Samples shown here are from September 2019 at 30 m, 40 m, and 85 m.

Figure 3. The relative abundances of the most commonly observed taxa are similar between May (A, B) and September (C, D). Notably, *Nitrosopumilus* sp. and Woesearchaeota dominated the middle and deep water column, respectively. The sulfur-oxidizing SUP05 clade increased continuously with depth in May but decreased in frequency below 70 m in September, while *Arcobacter* sp. spiked sharply at 80 m in September. Sampled depths are represented by points on each line.

Figure 4. Phylogenomic analysis of the Woesearchaeotal MAG and ninety publicly available MAGs from a range of biomes show the AJ population is mostly closely related to Woesearchaeota from other marine water columns.

996
997 Figure 5. Relative abundances of genes involved in sulfur (A) and nitrogen (B) metabolism
998 in each metagenome (May 60 m, May 106 m, September 60 m, September 95 m) show
999 some dissimilatory genes, such as *dsr*AB and *nap*AB, are enriched in the deep water
1000 layer.

1001

1002 **Table Legends**

1003

Table 1. All de-replicated, high-quality metagenome-assembled genomes (MAGs) from
the 60 m and deep co-assemblies, as well as one MAG from the September 95 m
individual assembly. Taxonomy assessed at the whole-genome level and nearest relative
(GTDB) are provided. If applicable, taxonomy according to the associated 16S rRNA gene
(SILVA) is provided with additional higher-order classification in parentheses. Amplicon
frequencies for corresponding SVs are provided for each sample. One MAG contained
two SVs (*, see Table S3) and here the frequencies for the first SV are provided.

1011

1012 Table 2. Taxonomy and functional annotations identified in each MAG. The biochemical
1013 process associated with each gene is provided by headers in bold.

1014

1015 Table S1. Amberjack Hole water column sample metadata and amplicon sequencing
1016 results. Processed reads are those that were quality-filtered, trimmed, and run through
1017 DADA2.

1018

1019 Table S2. Metagenome assembly data including sample source, pre- and post-quality
1020 filtered reads, and assembly statistics for both single and co-assemblies.

1021

1022 Table S3. Metagenome-assembled genomes (MAGs) and their associated samples,
1023 quality metrics, and assigned taxonomy.

1024

1025 Table S4. Metagenome-assembled genomes (MAGs) assessed in the phylogenomic
1026 trees (Fig. 4, Fig. S4, Fig. S5) including the Amberjack MAG of interest and the source of
1027 all other MAGs included in the phylogeny.

1028

1029 **Conflicts of interest**

1030

1031 The authors declare no conflict of interest.

1032

1033 **Funding**

1034

1035 This work was funded by the National Oceanic and Atmospheric Administration Office of
1036 Exploration and Research (grant #NA18OAR0110291).

1037
Acknowledgments

1039
1040 We thank the captains of the *R/V* William R. Mote and *R/V* Eugenie Clark for their
1041 support and professionalism, and staff at Mote Marine Lab for their work in sample
1042 collection and processing. We are grateful to Dr. Navid Constantinou for his help
1043 modeling the wind speeds required for turnover of the water column.
1044
1045

1046 **References**

1047
1.   Mylroie JE, Carew JL, Moore AI. Blue Holes: Definition and Genesis. Carbonates

and Evaporates. 1995;10(2):225–33.

2.   Canganella F, Bianconi G, Kato C, Gonzalez J. Microbial ecology of submerged

marine caves and holes characterized by high levels of hydrogen sulphide. In:

Reviews in Environmental Science and Biotechnology. 2007.

3.   Gischler E, Shinn EA, Oschmann W, Fiebig J, Buster NA. A 1500-Year Holocene

Caribbean Climate Archive from the Blue Hole, Lighthouse Reef, Belize. J Coast

Res. 2008;246:1495–505.

4.   Pohlman JW. The biogeochemistry of anchialine caves: Progress and

possibilities. Hydrobiologia. 2011;677:33–51.

5.   Seymour JR, Humphreys WF, Mitchell JG. Stratification of the microbial

community inhabiting an anchialine sinkhole. Aquat Microb Ecol. 2007;

6.   Garman KM, Rubelmann H, Karlen DJ, Wu T, Garey JR. Comparison of an

inactive submarine spring with an active nearshore anchialine spring in Florida.

Hydrobiologia. 2011;

7.   Gonzalez BC, Iliffe TM, Macalady JL, Schaperdoth I, Kakuk B. Microbial hotspots

in anchialine blue holes: Initial discoveries from the Bahamas. Hydrobiologia.

2011;677(1):149–56.

1066 8. Davis MC, Garey JR. Microbial function and hydrochemistry within a stratified

1067   anchialine sinkhole: A window into coastal aquifer interactions. Water.

1068   2018;10(8):972.

1069 9. Yao P, Wang XC, Bianchi TS, Yang ZS, Fu L, Zhang XH, et al. Carbon Cycling in

1070   the World's Deepest Blue Hole. J Geophys Res Biogeosciences. 2020;

1071 10. He H, Fu L, Liu Q, Fu L, Bi N, Yang Z, et al. Community Structure, Abundance

1072   and Potential Functions of Bacteria and Archaea in the Sansha Yongle Blue Hole,

1073   Xisha, South China Sea. Front Microbiol. 2019;10:2404.

1074 11. He P, Xie L, Zhang X, Li J, Lin X, Pu X, et al. Microbial Diversity and Metabolic

1075   Potential in the Stratified Sansha Yongle Blue Hole in the South China Sea. Sci

1076   Rep. 2020;10:5949.

1077 12. DeWitt D. Submarine Springs and other Karst Features in Offshore Waters of the

1078   Gulf of Mexico and Tampa Bay, Southwest Florida Water Management District.

1079   2003.

1080 13. Hughes JD, Vacher HL, Sanford WE. Three-dimensional flow in the Florida

1081   platform: Theoretical analysis of Kohout convection at its type locality. Geology.

1082   2007;35(7):663–6.

1083 14. Hu C, Muller-Karger FE, Swarzenski PW. Hurricanes, submarine groundwater

1084   discharge, and Florida's red tides. Geophys Res Lett. 2006;33(11):L11601.

1085 15. Smith CG, Swarzenski PW. An investigation of submarine groundwater-borne

1086   nutrient fluxes to the west Florida shelf and recurrent harmful algal blooms. Limnol

1087   Oceanogr. 2012;57(2):471–85.

1088 16. Walsh JJ, Jolliff JK, Darrow BP, Lenes JM, Milroy SP, Remsen A, et al. Red tides

in the Gulf of Mexico: Where, when, and why? J Geophys Res.

2006;111(C11003):1–46.

17. Weisberg RH, Liu Y, Lembke C, Lu C, Hubbard C, Garrett M. The coastal ocean

circulation influence on the 2018 West Florida Shelf *K. brevis* red tide bloom. J

Geophys Res Ocean. 2019;124(4):2501–12.

18. Vargo GA, Heil CA, Fanning KA, Dixon LK, Neely MB, Lester K, et al. Nutrient

availability in support of Karenia brevis blooms on the central West Florida Shelf:

What keeps Karenia blooming? Cont Shelf Res. 2008;28(1):73–98.

19. Saunders JK, Fuchsman CA, McKay C, Rocap G. Complete arsenic-based

respiratory cycle in the marine microbial communities of pelagic oxygen-deficient

zones. Proc Natl Acad Sci U S A. 2019;116(20):9925–30.

20. Callbeck CM, Lavik G, Ferdelman TG, Fuchs B, Gruber-Vodicka HR, Hach PF, et

al. Oxygen minimum zone cryptic sulfur cycling sustained by offshore transport of

key sulfur oxidizing bacteria. Nat Commun. 2018;9:1729.

21. Garcia-Robledo E, Padilla CC, Aldunate M, Stewart FJ, Ulloa O, Paulmier A, et al.

Cryptic oxygen cycling in anoxic marine zones. Proc Natl Acad Sci U S A.

2017;114(31):8319–1824.

22. Sun X, Kop LFM, Lau MCY, Frank J, Jayakumar A, Lücker S, et al. Uncultured

Nitrospina-like species are major nitrite oxidizing bacteria in oxygen minimum

zones. ISME J. 2019;13:2391–402.

23. Tsementzi D, Wu J, Deutsch S, Nath S, Rodriguez-R LM, Burns AS, et al. SAR11

bacteria linked to ocean anoxia and nitrogen loss. Nature. 2016;536:179–83.

24. Thamdrup B, Steinsdóttir HGR, Bertagnolli AD, Padilla CC, Patin N V., Garcia-

Robledo E, et al. Anaerobic methane oxidation is an important sink for methane in

the ocean's largest oxygen minimum zone. Limnol Oceanogr. 2019;64(6):2569–

85.

25. Breitburg D, Levin LA, Oschlies A, Grégoire M, Chavez FP, Conley DJ, et al.

Declining oxygen in the global ocean and coastal waters. Science (80-).

2018;359(6371):eaam7240.

26. Baker BJ, De Anda V, Seitz KW, Dombrowski N, Santoro AE, Lloyd KG. Diversity,

ecology and evolution of Archaea. Nat Microbiol. 2020;5:887–900.

27. Thiel V, Costas AMG, Fortney NW, Martinez JN, Tank M, Roden EE, et al.

"*Candidatus* Thermonerobacter thiotrophicus," a non-phototrophic member of the

Bacteroidetes/Chlorobi with dissimilatory sulfur metabolism in hot spring mat

communities. Front Microbiol. 2019;9:3159.

28. Helly JJ, Levin LA. Global distribution of naturally occurring marine hypoxia on

continental margins. Deep Res Part I Oceanogr Res Pap. 2004;51(9):1159–68.

29. Wyrtki K. The oxygen minima in relation to ocean circulation. Deep Res Oceanogr

Abstr. 1962;9(1–2):11–23.

30. Thamdrup B, Dalsgaard T, Revsbech NP. Widespread functional anoxia in the

oxygen minimum zone of the Eastern South Pacific. Deep Res Part I Oceanogr

Res Pap. 2012;65:36–45.

31. Garman KM, Garey JR. The transition of a freshwater karst aquifer to an anoxic

marine system. Estuaries. 2005;28:686–93.

32. Hughes JD, Vacher HL, Sanford WE. Temporal response of hydraulic head,

temperature, and chloride concentrations to sea-level changes, Floridan aquifer

system, USA. Hydrogeol J. 2009;17(4):793–815.

33.   Simpson JH. The shelf-sea fronts: implications of their existence and behaviour.

Philos Trans R Soc London Ser A, Math Phys Sci. 1981;302(1472):531–46.

34.   Luther GW, Findlay AJ, MacDonald DJ, Owings SM, Hanson TE, Beinart RA, et

al. Thermodynamics and kinetics of sulfide oxidation by oxygen: a look at

inorganically controlled reactions and biologically mediated processes in the

environment. Front Microbiol. 2011;2(62):1–9.

35.   Ghosh W, Dam B. Biochemistry and molecular biology of lithotrophic sulfur

oxidation by taxonomically and ecologically diverse bacteria and archaea. FEMS

Microbiol Rev. 2009;33(6):999–1043.

36.   Wright JJ, Konwar KM, Hallam SJ. Microbial ecology of expanding oxygen

minimum zones. Nat Rev Microbiol. 2012;10:381–94.

37.   Walsh DA, Zaikova E, Howes CG, Song YC, Wright JJ, Tringe SG, et al.

Metagenome of a versatile chemolithoautotroph from expanding oceanic dead

zones. Science (80-). 2009;326(5952):578–82.

38.   Hawley AK, Brewer HM, Norbeck AD, Pasă-Tolić L, Hallam SJ. Metaproteomics

reveals differential modes of metabolic coupling among ubiquitous oxygen

minimum zone microbes. Proc Natl Acad Sci U S A. 2014;111(31):11395–400.

39.   Bertagnolli AD, Stewart FJ. Microbial niches in marine oxygen minimum zones.

Nat Rev Microbiol. 2018;16:723–9.

40.   Wirsen CO, Sievert SM, Cavanaugh CM, Molyneaux SJ, Ahmad A, Taylor LT, et

al. Characterization of an autotrophic sulfide-oxidizing marine Arcobacter sp. that

produces filamentous sulfur. Appl Environ Microbiol. 2002;68(1):316–25.

41.   Rozan TF, Theberge SM, Luther G. Quantifying elemental sulfur (S0), bisulfide

1159   (HS-) and polysulfides (S(x)2-) using a voltammetric method. Anal Chim Acta.

1160   2000;415(1–2):175–84.

1161   42.   Luther GW, Glazer BT, Hohmann L, Popp JI, Taillefert M, Rozan TF, et al. Sulfur

1162   speciation monitored *in situ* with solid state gold amalgam voltammetric

1163   microelectrodes: Polysulfides as a special case in sediments, microbial mats and

1164   hydrothermal vent waters. J Environ Monit. 2001;3:61–6.

1165   43.   Heylen K, Vanparys B, Wittebolle L, Verstraete W, Boon N, De Vos P. Cultivation

1166   of denitrifying bacteria: Optimization of isolation conditions and diversity study.

1167   Appl Environ Microbiol. 2006;72(4):2637–43.

1168   44.   Taillefert M, Bono AB, Luther GW. Reactivity of Freshly Formed Fe(III) in

1169   Synthetic Solutions and (Pore)Waters: Voltammetric Evidence of an Aging

1170   Process. Environ Sci Technol. 2000;34(11):2169–77.

1171   45.   Barco RA, Emerson D, Sylvan JB, Orcutt BN, Jacobson Meyers ME, Ramírez GA,

1172   et al. New insight into microbial iron oxidation as revealed by the proteomic profile

1173   of an obligate iron-oxidizing chemolithoautotroph. Appl Environ Microbiol.

1174   2015;81(17):5927–37.

1175   46.   Garber AI, Nealson KH, Okamoto A, McAllister SM, Chan CS, Barco RA, et al.

1176   FeGenie: A Comprehensive Tool for the Identification of Iron Genes and Iron

1177   Gene Neighborhoods in Genome and Metagenome Assemblies. Front Microbiol.

1178   2020;11:37.

1179   47.   Castelle CJ, Wrighton KC, Thomas BC, Hug LA, Brown CT, Wilkins MJ, et al.

1180   Genomic expansion of domain archaea highlights roles for organisms from new

phyla in anaerobic carbon cycling. Curr Biol. 2015;25(6):690–701.

48. Liu X, Li M, Castelle CJ, Probst AJ, Zhou Z, Pan J, et al. Insights into the ecology,

evolution, and metabolism of the widespread Woesearchaeotal lineages.

Microbiome. 2018;6:102.

49. Ortiz-Alvarez R, Casamayor EO. High occurrence of Pacearchaeota and

Woesearchaeota (Archaea superphylum DPANN) in the surface waters of

oligotrophic high-altitude lakes. Environ Microbiol Rep. 2016;8(2):210–7.

50. Aylward FO, Santoro AE. Heterotrophic Thaumarchaea with Small Genomes Are

Widespread in the Dark Ocean. mSystems. 2020;5(3):e00415-20.

51. Reji L, Francis CA. Metagenome-assembled genomes reveal unique metabolic

adaptations of a basal marine Thaumarchaeota lineage. ISME J. 2020;14:2105–

2115.

52. Santoro AE, Richter RA, Dupont CL. Planktonic Marine Archaea. Ann Rev Mar

Sci. 2019;11:131–58.

53. Rinke C, Rubino F, Messer LF, Youssef N, Parks DH, Chuvochina M, et al. A

phylogenomic and ecological analysis of the globally abundant Marine Group II

archaea (*Ca.* Poseidoniales ord. nov.). ISME J. 2019;13:663–75.

54. Pereira O, Hochart C, Auguet JC, Debroas D, Galand PE. Genomic ecology of

Marine Group II, the most common marine planktonic Archaea across the surface

ocean. Microbiol Open. 2019;8(9):e00852.

55. Martin-Cuadrado AB, Garcia-Heredia I, Moltó AG, López-Úbeda R, Kimes N,

López-García P, et al. A new class of marine Euryarchaeota group II from the

mediterranean deep chlorophyll maximum. ISME J. 2015;9:1619–34.

56. Moreira D, Rodríguez-Valera F, López-García P. Analysis of a genome fragment

of a deep-sea uncultivated Group II euryarchaeote containing 16S rDNA, a

spectinomycin-like operon and several energy metabolism genes. Environ

Microbiol. 2004;6(9):959–69.

57. Martin-Cuadrado AB, Rodriguez-Valera F, Moreira D, Alba JC, Ivars-Martínez E,

Henn MR, et al. Hindsight in the relative abundance, metabolic potential and

genome dynamics of uncultivated marine archaea from comparative

metagenomic analyses of bathypelagic plankton of different oceanic regions.

ISME J. 2008;2:865–86.

58. Sforna MC, Philippot P, Somogyi A, Van Zuilen MA, Medjoubi K, Schoepp-

Cothenet B, et al. Evidence for arsenic metabolism and cycling by

microorganisms 2.7 billion years ago. Nat Geosci. 2014;7:811–5.

59. Luther GW, Glazer BT, Ma S, Trouwborst RE, Moore TS, Metzger E, et al. Use of

voltammetric solid-state (micro)electrodes for studying biogeochemical processes:

Laboratory measurements to real time measurements with an in situ

electrochemical analyzer (ISEA). Mar Chem. 2008;108(3–4):221–35.

60. Brendel PJ, Luther GW. Development of a Gold Amalgam Voltammetric

Microelectrode for the Determination of Dissolved Fe, Mn, O_2, and S(-II) in

Porewaters of Marine and Freshwater Sediments. Environ Sci Technol.

1995;29(3):751–61.

61. Tercier-Waeber M Lou, Taillefert M. Remote in situ voltammetric techniques to

characterize the biogeochemical cycling of trace metals in aquatic systems. J

Environ Monit. 2008;10(1):30–54.

62.   Arar EJ, Collins GB. Method 445.0 In Vitro Determination of Chlorophyll a and

1228   Pheophytin a in Marine and Freshwater Algae by Fluorescence. 1997.

1229   63.   Bran+Luebbe/Seal. Ammonia in water and seawater. Method No G-171-96 Rev

1230   10. 2005;Norderstedt, Germany.

1231   64.   Bran+Luebbe/Seal. Nitrate and nitrite in water and seawater; total nitrogen in

1232   persulfate digests. Method No G-172-96 Rev 10. 2010;Norderstedt, Germany.

1233   65.   Solórzano L, Sharp JH. Determination of total dissolved phosphorus and

1234   particulate phosphorus in natural waters. Limnol Oceanogr. 1980;25(4):754–8.

1235   66.   Solórzano L, Sharp JH. Determination of total dissolved nitrogen in natural

1236   waters. Limnol Oceanogr. 1980;25(4):751–4.

1237   67.   Dickson AG, Sabine CL, Christian JR. Guide to best practices for ocean CO_2

1238   measurements. PICES Spec Publ 3. 2007;

1239   68.   Murphy J, Riley JP. A modified single solution method for the determination of

1240   phosphate in natural waters. Anal Chim Acta. 1962;27:31–6.

1241   69.   Stookey LL. Ferrozine-A New Spectrophotometric Reagent for Iron. Anal Chem.

1242   1970;42(7):779–81.

1243   70.   Padilla CC, Bertagnolli AD, Bristow LA, Sarode N, Glass JB, Thamdrup B, et al.

1244   Metagenomic binning recovers a transcriptionally active gammaproteobacterium

1245   linking methanotrophy to partial denitrification in an anoxic oxygen minimum zone.

1246   Front Mar Sci. 2017;4:23.

1247   71.   Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of

1248   a dual-index sequencing strategy and curation pipeline for analyzing amplicon

1249   sequence data on the MiSeq Illumina sequencing platform. Appl Environ

1250   Microbiol. 2013;79(17):5112–20.

1251   72.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP.

1252   DADA2: High-resolution sample inference from Illumina amplicon data. Nat

1253   Methods. 2016;13:581–3.

1254   73.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.

1255   Reproducible, interactive, scalable and extensible microbiome data science using

1256   QIIME 2. Nat Biotechnol. 2019;37:852–7.

1257   74.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion

1258   for RNA-seq data with DESeq2. Genome Biol. 2014;15:550.

1259   75.   McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive

1260   Analysis and Graphics of Microbiome Census Data. PLoS One.

1261   2013;8(4):e61217.

1262   76.   McMurdie PJ, Holmes S. Waste Not, Want Not: Why Rarefying Microbiome Data

1263   Is Inadmissible. PLoS Comput Biol. 2014;10(4):e1003531.

1264   77.   Willis AD, Martin BD. DivNet: Estimating diversity in networked communities.

1265   bioRxiv. 2018;

1266   78.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-

1267   Verlag; 2016.

1268   79.   Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, Mcglinn D, et al.

1269   vegan: Community Ecology Package [Internet]. Community ecology package.

1270   2019. p. R package version 2.5-5. Available from: https://cran.r-

1271   project.org/package=vegan

80.   Nayfach S, Pollard KS. Average genome size estimation improves comparative

metagenomics and sheds light on the functional ecology of the human

microbiome. Genome Biol. 2015;16:51.

81. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. MetaSPAdes: A new versatile

metagenomic assembler. Genome Res. 2017;27(5):824–34.

82. Li D, Liu CM, Luo R, Sadakane K, Lam TW. MEGAHIT: An ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn

graph. Bioinformatics. 2015;31(10):1674–6.

83. Mikheenko A, Saveliev V, Gurevich A. MetaQUAST: Evaluation of metagenome

assemblies. Bioinformatics. 2016;32(7):1088–90.

84. Seemann T. Prokka: Rapid prokaryotic genome annotation. Bioinformatics.

2014;30(14):2068–9.

85. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:

Prokaryotic gene recognition and translation initiation site identification. BMC

Bioinformatics. 2010;11:119.

86. James BT, Luczak BB, Girgis HZ. MeShClust: an intelligent tool for clustering

DNA sequences. Nucleic Acids Res. 2018;46(14):e83.

87. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al.

KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive

score threshold. Bioinformatics. 2020;

88. Boratyn GM, Thierry-Mieg J, Thierry-Mieg D, Busby B, Madden TL. Magic-

BLAST, an accurate RNA-seq aligner for long and short reads. BMC

Bioinformatics. 2019;

89. Dunivin TK, Yeh SY, Shade A. A global survey of arsenic-related genes in soil

1296   microbiomes. BMC Biol. 2019;17:45.

1297 90. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM:

1298   Assessing the quality of microbial genomes recovered from isolates, single cells,

1299   and metagenomes. Genome Res. 2015;25(7):1043–55.

1300 91. Eren AM, Esen OC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o:

1301   An advanced analysis and visualization platformfor 'omics data. PeerJ.

1302   2015;3:e1319.

1303 92. Olm MR, Brown CT, Brooks B, Banfield JF. DRep: A tool for fast and accurate

1304   genomic comparisons that enables improved genome recovery from

1305   metagenomes through de-replication. ISME J. 2017;11(12):2864–8.

1306 93. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify

1307   genomes with the Genome Taxonomy Database. Bioinformatics.

1308   2019;36(6):1925–7.

1309 94. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA, et

1310   al. A standardized bacterial taxonomy based on genome phylogeny substantially

1311   revises the tree of life. Nat Biotechnol. 2018;36:996–1004.

1312 95. Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and post-

1313   analysis of large phylogenies. Bioinformatics. 2014;30(9):1312–1313.

1314 96. Letunic I, Bork P. Interactive Tree of Life (iTOL) v4: Recent updates and new

1315   developments. Nucleic Acids Res. 2019;47(W1):W256–W259.

1316 97. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al.

1317   Fiji: An open-source platform for biological-image analysis. Nat Methods.

1318   2012;9:676–682.

1319

1320
1321
1322
1323

Figure 1

Figure 2

Figure 3

Legend:
- Woesearchaeota
- *Ca*. Nitrosopumilus
- Synechococcus CC9902
- Prochlorococcus MIT9313
- Thioglobaceae (SUP05)
- SAR11 Clade Ia
- SAR11 Clade II
- Thermoplasmatota Marine Group III
- Thermoplasmatota Marine Group II
- Bacteroidota VC2.1 Bac22
- Arcobacter

Figure 4

Figure 5

A. Sulfur metabolism

B. Nitrogen metabolism