preprint

Place cells in head-fixed mice navigating a floating real-world environment

Mary Ann Go [1*]**, Jake Rogers** [1]**, Giuseppe P. Gava** [1]**, Catherine Davey** [1,2]**,**

Seigfred Prado [1]**, Yu Liu** [1] **and Simon R. Schultz** [1*]

[1]*Department of Bioengineering and Centre for Neurotechnology, Imperial College London, London, United Kingdom*
[2]*Department of Biomedical Engineering, University of Melbourne, Melbourne, Victoria, Australia*

Correspondence*:
Dr. Mary Ann Go, Imperial College London, m.go@imperial.ac.uk
Prof Simon R. Schultz, Imperial College London, s.schultz@imperial.ac.uk

ABSTRACT

The hippocampal place cell system in rodents has provided a major paradigm for the scientific investigation of memory function and dysfunction. Place cells have been observed in area CA1 of the hippocampus of both freely moving animals, and of head-fixed animals navigating in virtual reality environments. However, spatial coding in virtual reality preparations has been observed to be impaired. Here we show that the use of a real-world environment system for head-fixed mice, consisting of a track floating on air, provides some advantages over virtual reality systems for the study of spatial memory. We imaged the hippocampus of head-fixed mice injected with the genetically encoded calcium indicator GCaMP6s while they navigated circularly constrained or open environments on the floating platform. We observed consistent place tuning in a substantial fraction of cells, stable over multiple days, with remapping of place fields when the animal entered a different environment. Spatial information rates were within the range observed in freely moving mice. Manifold analysis indicated that spatial information could be extracted from a low-dimensional subspace of the neural population dynamics. This is the first demonstration of place cells in head-fixed mice navigating on an air-lifted real-world platform, validating its use for the study of brain circuits involved in memory and affected by neurodegenerative disorders.

Keywords: CA1, hippocampus, mouse, multiphoton, two-photon, calcium imaging, behavior, spatial memory

1 INTRODUCTION

Place cells in the hippocampus encode spatial information during navigation, by firing selectively when the animal is in a certain part of its environment (O'Keefe and Dostrovsky, 1971; Muller and Kubie, 1989; Taube, 1995). This location-specific firing requires the encoding and recall of spatial memory, and has been suggested to be the "where" component of episodic memory (Ergorul and Eichenbaum, 2004; Leutgeb et al., 2005; O'Keefe, 2007). As such, it provides a powerful model system for investigating the neural circuit mechanisms underlying learning and memory, as well as the neurological basis of neurodegenerative disorders affecting memory such as Alzheimer's Disease (Cacucci et al., 2008; Mably et al., 2017).

Place cells have been readily recorded using electrophysiological techniques in freely moving mice and
rats. However, additional insight into system function can be gained using single- and multiple-photon
fluorescence imaging techniques, which enable large populations of genetically labelled neurons to be
monitored simultaneously (Peron et al., 2015; Schultz et al., 2016). While calcium fluorescence place
fields in the hippocampus have been imaged using single-photon microendoscopy (Ziv et al., 2013), image
quality, imageable depth of field, optical sectioning and consequent cell separability is much greater with
two- than one-photon microscopy, which is why it has become the gold standard technique for spatially
resolved investigation of cortical circuit function. Two-photon microendoscopy is possible, but is much less
well developed, and optical access to the brain is still inferior to that possible in head-fixed preparations
(Ozbay et al., 2018; Zong et al., 2017). This has spurred the development of solutions for employing rodent
spatial navigation behavioural tasks during head fixation.

One increasingly popular approach is the use of Virtual Reality (VR) platforms. In such systems, a
mouse is typically head-fixed atop a polystyrene ball which is free to rotate. Optical sensors tracking
ball movement are used to update a visual display (Dombeck et al., 2007; Muzzu et al., 2018), allowing
either one- or two-dimensional movement. Place cells have been observed in head-fixed mice navigating in
one-dimensional (i.e. linear track) virtual environments in such an apparatus (Harvey et al., 2009; Dombeck
et al., 2010; Rickgauer et al., 2014). One advantage of these VR systems is that they allow controlled
manipulation of the environment in a way not easily possible in the real world. Moreover, because the
mouse's head is fixed, intracellular recording and two-photon imaging are made feasible. However, there
are several key disadvantages of VR systems, including the lack of translational vestibular input, and the
lack of sensory feedback of modalities that may be more behaviourally salient for rodents than vision
(Ravassard et al., 2015). Two-dimensional place tuning has been shown to be profoundly impaired in
VR spatial navigation (Aghajan et al., 2015), and in addition, the theta rhythm frequency has been found
to be slower in VR environments (Aronov and Tank, 2014). In fact, 2D place tuning has, to date, only
been observed in VR systems where the rodent is suspended in a body jacket attached to a commutator
allowing it to make head movements and to rotate its body through a full 360° (Aronov and Tank, 2014), or
with a complex commutator headplate attachment that allows head movements constrained to horizontal
rotations (Chen et al., 2018, 2019). Neither system allows straightforward extension to two-photon imaging
or intracellular recording.

Recently, a 2D real-world system in which mice are head-fixed while navigating a track floating on air
has been developed (Kislin et al., 2014; Nashaat et al., 2016). The system allows for sensory feedback and
head immobility allows for intracellular recording and two-photon imaging. Until now, the presence of
place cells in such an environment has not been shown. We show here that mice can navigate the floating
track and give the first demonstration that the system allows for place tuning, with spatial information rates
broadly comparable to those observed in head-fixed mice.

2   MATERIALS AND METHODS

2.1   Animals

All experimental procedures were carried out under the Animals (Scientific Procedures) Act 1986 and
according to Home Office and institutional guidelines. Subjects were C57BL/6 mice of age 1 to 9 months
at the time of viral injection. Behavioural data were collected from 6 mice (5 males, 1 female, median age:
8.0 months) for the circular track and from 5 mice (all females, median age: 1.5 months) for the circular
open field. Imaging data were collected from 3 mice (2 males, 1 female, median age: 8.2 months) for the

circular track and from 3 mice (all females, median age: 1.7 months) for the open field. Animals were kept
on a reverse 12-h light, 12-h dark cycle with lights on at 7pm.

2.2   Virus injection and hipppocampal window

Mice were anaesthetised with 1.5-3% isofluorane. Body temperature was monitored with a rectal thermal
probe and kept at 37°C using a heating blanket. Analgesia was administered pre-operatively with Carprofen
(5 mg/kg) and buprenorphine (0.07 mg/kg). A small (∼0.5 mm) craniotomy was made and the virus
AAV1.hSyn1.mRuby2.GSG.P2A.GCaMP6s.WPRSE.SV40 (Addgene 50942, titer 1.9 × 10^{13} vg/ml, ∼50
nL) was injected into the hippocampus (from bregma, in mm: 1.7-1.8 ML, 2.0 AP) 1.5 mm from the dural
surface. The virus contains a green genetically encoded calcium indicator protein (GCaMP6s) and a red
fluorescent protein (mRuby) for cell body localisation. Two weeks post-injection, a hippocampal window
was implanted as described by Dombeck et al. (2010). A circular craniotomy centred on the previously
made injection hole was marked using a 3-mm diameter biopsy punch and the cranial bone was removed
using a dental drill. The cortex above the injection site was aspirated using a 27 gauge needle connected to
a water pump until the fibers of the corpus callosum became visible. A stainless steel cannula (diameter: 3
mm, height: 1.5 mm) with a glass bottom was then pressed down into the tissue and fixed in place using
histoacryl glue. The surrounding skull was roughened using a scalpel blade to make crisscross lines before
a stainless steel headplate (aperture: 8.5 mm) was attached to the skull, centred on the craniotomy, using
histoacryl glue. Exposed skull outside the headplate aperture was covered with dental cement mixed with
black powder paint. Mice were given 5-7 days to recover before behavioural training was started.

2.3   Behavioural training

Approximately one week after the hippocampal window was implanted, the animals were habituated to
the experimenter by handling and were placed under water restriction. Behavioural training started the
following day. Animals were trained to move in the dark either along a circular track (outer diameter: 32.5
cm, width: 5 cm) or in a circular open field (diameter: 32.5 cm), both floating on an air table (Mobile
HomeCage Large, Neurotar). The floating tracks were made of carbon fibre (weight: 100 ± 2.8 g) and
had 4-cm high walls lined with visual (phosphorescent tapes, Gebildet E055 and E068) and tactile cues
(sandpaper, cardboard, foam, bubble wrap). The phosphorescent tapes emitted light at 500 (blue) and
520 (green) nm and glowed for the duration of the imaging session ($\leq$1 hr per track, Supplementary Fig.
1). The floor of the circular open field was also lined with tactile cues (mesh tape). The air table rests
under a two-photon resonant scanning microscope (Scientifica Ltd, Uckfield UK, Fig. 1A) and is fitted
with a magnet-based position tracking system that acquires data (including, among others, Euclidean and
polar mouse coordinates and speed) at 100 Hz. A lick spout was attached to the headplate mount and
automated water delivery was controlled using a peristaltic pump (Campden Instruments). Water rewards
were accompanied by a beep.

Animals were trained twice daily in 45-min sessions. In the circular track, animals were trained with
one circular track in the morning and a second circular track with different visual and tactile cues in the
afternoon. Animals were given a water reward of 4 uL per loop traversed at random locations. In the
circular open field, a 4-uL reward was given every time the animal went from an outer region (outer 6 cm of
open field) to either one of two inner regions (two inner 20.5 cm diameter half circles), or vice versa. Daily
water intake was limited to 1-3 mL and was individually adjusted for each mouse to maintain the target
weight of 85% of the pre-restriction weight. At the start of each training session, two rewards were given to
motivate the mice to lick for water. There was no limit to the number of rewards animals could have during
sessions. If the animal did not reach the target volume for the day during training, the remaining volume

was given at the end of the last training session for the day. Animals were trained for 11-14 sessions in the
circular track and 16-18 sessions in the open field before imaging was started. Mice that did not have good
GCaMP6s expression in the CA1 region were excluded from imaging experiments.

2.4 Two-photon imaging

We used a commercial two-photon resonant scanning microscope (VivoScope, Scientifica) equipped with a
tiltable objective mount and a $16\times$ water-immersion objective (LWD 0.8 NA, Nikon). Ultrasound gel at
50% concentration was used as immersion liquid. GCaMP6s and mRuby were excited at 940 nm with a
Ti:sapphire laser (Mai Tai, Newport). The laser power underneath the objective was 60-166 mW. Images
(512×512 pixels, 330×330 μm or 490×490 μm field of view) were acquired at 30 Hz during imaging
sessions which lasted up to an hour. In remapping experiments, mice were imaged for up to an hour in
one track then moved to another track where they were imaged for up to another hour. SciScan software
(Scientifica) was used for microscope control, and image acquisition was TTL-synchronized to position
tracking and reward timing signals. Light from the phosphorescent tapes on the floating track walls used as
visual cues was insignificant in the green and red imaging channels.

To characterize stability of place fields, animals were imaged in the circular track in multiple sessions
spanning up to 8 days. To locate the same set of cells in later sessions, we used the red channel image from
previous sessions as reference.

2.5 Processing of calcium imaging movies

We used the MATLAB (Mathworks, Natick, MA) implementation of the CaImAn software package
(Giovannucci et al., 2019) for motion correction, automatic identification of regions of interest (ROIs)
and deconvolution of neural activity from fluorescence traces. To remove motion artefact from the
calcium imaging videos, we first did rigid image registration then non-rigid image registration. For
ROI identification, the maximum number of ROIs and the average cell size for a given field of view (FOV)
were estimated by examining representative images in ImageJ. Overlapping ROIs were excluded. Signal
contribution from the surrounding neuropil was removed using the FISSA toolbox (Keemink et al., 2018).
Neural activity was then deconvolved from the neuropil-decontaminated fluorescence traces using the
OASIS algorithm (Friedrich et al., 2017). This produced an event train which preserved both the time and
amplitude of inferred calcium transient events. In some analyses, the presence/absence of calcium transient
events was used; integrating the number of such events over a fixed time window, and dividing by the
length of the window yields the neural event rate (units events/sec). In other analyses, we took into account
the amplitude of events; in the absence of per-cell calibration to true spike counts, we consider the units
of the amplitude of this event train to be (dimensionless) $\Delta F/F$, inherited from the original time series
extracted from each ROI. Integrating the amplitude of such events over a fixed time window and dividing
by the length of the window yields the neural activity rate (units $\Delta F/F.s^{-1}$).

To track cells across multiple imaging sessions, we motion-corrected images from different sessions
using the motion-corrected image from one session as a template. We then temporally concatenated the
videos from different sessions and ran the ROI segmentation algorithm on the concatenated video.

2.6 Data analysis

Inferred neural activity and mouse tracking data were analysed using custom scripts written in MATLAB.
For place field analysis, only time points for which the animal speed exceeded 20 mm/s were included, to
eliminate periods when the animals paused for rewards or for grooming.

For the circular track, mouse position was linearised by converting angular distance to Euclidean distance
using the known circumference of the circular track. We used 2-cm bins and computed neural event rates
by dividing the total number of neural events during occupancy of the bin by the total occupancy time there.
These rate maps were smoothed using a boxcar average over three bins and each map was normalised by
its maximum value. Spatial information (in bits/event) for each cell was computed as described previously
(Skaggs et al., 1992). Neurons were classified as place cells if they met the following criteria: 1) neural
events were present for at least half of the traversals (laps) through the circular track, 2) neural events were
present for at least 5% of the time the mouse spent within the place field and 3) the cell contained spatial
information greater than chance. Chance-level spatial information for a cell was determined by performing
1,000 shuffles of the inferred neural activity and calculating the spatial information rate for each shuffle.
The cell was considered a place cell if its information rate exceeded 99% of the values for the shuffled
data. The location of the place field was defined by the bin location of maximum neural event rate while
the place field size was determined by the number of bins for which the neural event rate was at least
50% of the maximum. We quantified stability of place fields across days by calculating recurrence, field
correlation and place field shift for all possible pairs of sessions separated by N days. Recurrence is the
fraction of place cells in one day that retain the classification after N days. Place field correlation is the
mean correlation of the rate maps of place cells that retained their classification in two sessions N days
apart. Place field shift is the difference in field location of place cells in two sessions at N days interval.

For the circular open field, bins were 2 cm × 2 cm and the rate maps were smoothed with a 2D Gaussian
smoothing kernel with a standard deviation of 1.5. Neurons were classified as place cells if 1) neural events
were present for at least 2% of the time the mouse spent within the place field and 2) the cell contained
spatial information greater than chance. Place field location was obtained by calculating the centroid
location of the normalised rate map above a threshold of 0.5.

In the Results that follow, we describe estimates of the means of distributions by the sample mean ± the
standard error of the mean, unless otherwise indicated. One exception to this is in the reporting of mean
activity rates ($\Delta F/F.s^{-1}$), which are expected to fall (approximately) on a log-normal distribution, thus a
symmetric distribution of errors would be inappropriate even in the limit of large samples. In this case,
therefore, we instead report the 90% confidence interval of the mean.

Neuronal variability analyses were performed by adapting methods from classical visual neuroscience
literature (Tolhurst et al., 1983) to calcium transient amplitude-event trains; instead of measuring how
variable neuronal responses were across repeated presentations of a stimulus, we analyzed how variable
responses were to laps around the same track. For each cell, we accumulated the amplitude of calcium
transient events occurring when the mouse was in each 2-cm bin, dividing it by the amount of time spent in
that spatial bin. Averaging or taking the variance of this quantity across laps gives the mean activity or
variance, respectively, for that spatial bin; as the mouse progresses along the track (e.g. into and out of
a place field), the mean changes, and thus we obtain the relationship between variance and mean of the
activity for that cell.

We used total least squares linear regression to fit a power law model $y = ax^\beta$ to the relationship between
activity variance and mean for each individual cell. The power law exponent β, which can be read off
from the slope of the fitted line, provides useful information about the reliability of neuronal signalling.
An exponent of 1 indicates reliability equivalent to that of a Poisson process; above 1 implies additional
sources of variability.

To study neural population dynamics, we embedded the population calcium traces, recorded during
periods when mouse running speed exceeded 20 mm/s, in a manifold to extract lower-dimensional
population activity patterns. Given the population activity matrix $\mathbf{X}$ of size $N \times T$, where N is the number
of recorded units and T the number of time samples, there are T population vectors $\mathbf{x}_t$ of length N. The aim
of this analysis was to uncover a lower-dimensional embedding $\mathbf{Y}$ ($M \times T$), where $M \ll N$ is the number
of manifold dimensions. To do this, we used Classical Multidimensional Scaling (MDS), after observing
qualitatively and quantitatively (in terms of variance accounted for) better representation (Supplementary
Fig. 2) in comparison to Principal Components Analysis (PCA). We generated a dissimilarity matrix to
account for the difference between population activity vectors $\mathbf{x}_t$ over the entire time course T. We used
the cosine distance metric, which evaluates the angle between any two population activity vectors $\mathbf{x}_t$ at
any two time points. ISOMAP produced a similar embedding (Supplementary Fig. 2). For the purpose of
comparing manifold dimensionality, we defined the dimensionality as the number of manifold components
(MDS eigenvalues and associated eigenvectors) required to explain 90% of the variance in the population
activity.

To decode the mouse's angular position (in either the circular track or open field environment), we applied
an Optimal Linear Estimator (Muzzu et al., 2018), incorporating a varying number of manifold dimensions,
and using 5-fold cross-validation.

3   RESULTS

3.1   Mouse behaviour in floating environment resembles tethered and free behaviour

Head-fixed mice were trained to navigate the floating track system (Fig. 1A) in the dark using operant
conditioning. We designed visual cues on the environment walls using phosphorescent tapes. These were
visible in the dark for the duration of the training session (45 min) and did not add significant weight to
the tracks, which would have increased their rotational inertia and made them harder to control. The outer
diameter (32.5 cm) of the floating track was constrained by the distance from the objective lens to the back
wall of the microscope.

In the circular track, the mice initially displayed minimal activity. They struggled to propel the track
with their feet, tracing circular trajectories that were irregular. Without any intervention, the mice quickly
adapted to moving in smooth circular trajectories (Fig. 1Bi) running up to speeds of 200 mm/s (Fig. 1Ci).
Activity and average speed increased after about 9 sessions (Fig. 1Di, Ei). By the end of the training period,
mice could complete >100 laps in 45 min.

In the circular open field, our goal was for the trajectory of a mouse to completely cover the 2D
environment, an important requirement for place field analysis. We rewarded the mice when they went
from an outer region (outer 6 cm of open field) to either one of two inner regions (inner 20.5 cm diameter
half circles) and vice-versa and found this protocol to be effective in achieving the desired behaviour. The
mice adapted to controlling the circular track with their feet, tracing meandering trajectories that were
sometimes near the wall and sometimes crossed the arena. This behaviour resembles those of freely-moving
(Benjamini et al., 2011; Samson et al., 2015) and tethered (Dupret et al., 2010; Trouche et al., 2016) mice
exploring circular arenas (Fig. 1Bii). Mice sometimes showed preference for peripheral locations in the
open field. Locomotion speeds were similar to the circular track (Fig. 1Cii), although it took longer for
the mice to increase activity and average speed, requiring 15 or so sessions (Fig. 1Dii, Eii). To quantify
coverage, we divided the circular arena into 2 cm $\times$ 2 cm bins. By the end of the training period, mouse
trajectories covered >90% of the open field in 45 min.

                                                                6

Figure 1. Experimental setup and behavioural training. **(A)** Experimental setup. Mouse is head fixed while navigating a floating track under a two-photon microscope (in light grey). Inset: Close-up view of head-fixed mouse on floating track. **(B)** 20-min spatial trajectories of head-fixed mice navigating **(i)** a circular track and **(ii)** a circular open field. **(C)** Distribution of mouse locomotion speeds during the behavioural sessions shown in B. **(D)** Fraction of session spent running in the **(i)** circular track ($N = 10$ mice) and **(ii)** open field ($N = 5$) throughout the training period. **(E)** Progression of running speed in the **(i)** circular track and **(ii)** open field throughout the training period.

Mice were trained in the floating track system after viral injection and hippocampal window implantation.
The similarity of their behaviour to that's observed in tethered and freely-moving mice is validation that
our surgical procedures do not adversely affect behaviour, consistent with previous reports (Dombeck
et al., 2010; Pilz et al., 2016). Greater numbers of animals were available at earlier than later stages of the
experimental pipeline simply because of the requirement to pass successive criteria relating to behavioural

performance, quality of preparation, GCaMP6s expression located in area of interest, and registration of
ROIs over multiple sessions.

3.2   CA1 cells form reliable 1D place fields in a floating circular track

To optically record the activity of CA1 neurons, we injected the adeno-associated virus
AAV1.hSyn1.mRuby2.GSG.P2A.GCaMP6s.WPRSE.SV40 into the hippocampus, removed the overlying
cortex and implanted an imaging window. Two-photon imaging (at 940 nm) through the hippocampal
window three weeks later showed robust expression of both the calcium indicator GCaMP6s and the static
marker mRuby in a large population of CA1 neurons (Fig. 2A). We primarily used the mRuby image to
repeatedly identify the same cells across time.

To verify place coding in CA1 neurons, we acquired two-photon time-series videos of GCaMP6s
fluorescence in the hippocampi of mice running along a circular track. We used an automated algorithm
(Giovannucci et al., 2019) that identifies cells and extracts their activity from the GCaMP6s fluorescence
changes ($\Delta F/F$). We found 53-225 (median: 101) active neurons per FOV (330×330 μm, 6 FOVs in 3
mice) imaged for 12-28 min (see Fig. 2B showing 225 ROIs for the representative image shown in Fig.
2A). We extracted $\Delta F/F$ traces from the ROIs and observed significant calcium transients (Fig. 2C). We
then used the deconvolved neuronal activity as a measure of spiking activity (Fig. 2D).

In total, we analysed 721 cells from 6 imaging areas in 3 mice and found that 12-43% (median: 35%) of
the detected neurons in each imaging area showed location-specific activity characteristic of place cells (Fig.
2E). These cells had well-defined fields of neuronal activity which were apparent with repeated traversals
(laps) of the circular track, though not occurring in every lap (Fig. 2F). Moreover, similar to reports in rats
(Mehta et al., 1997; Lee and Knierim, 2007), we observed in some cells a backward shift in the location of
the place field in later laps (Fig. 2F). Place cells were active in $72.1 \pm 0.6\%$ (mean $\pm$ s.e.m.) of the laps
and their firing fields were homogeneously distributed across the circular track (Fig. 2G-I). These cells
showed a skewed distribution of place field size with an average of 19.7 ± 0.4 cm (Fig. 2J), comparable
to reported field sizes for mice (Chen et al., 2013; Ziv et al., 2013). On average, the information rate of
place cells was 0.90 ± 0.02 bits/event (Fig. 2K), within the range of values reported for freely moving
mice (Chen et al., 2013; Mou et al., 2018; Gonzalez et al., 2019) and for head-fixed mice navigating virtual
linear tracks (Arriaga and Han, 2017).

Imaged hippocampal CA1 neurons produced an average of 0.65 ± 0.01 calcium transient events per
second ($n = 721$ cells) during locomotion around the circular track. This is consistent with previous
observations from freely-moving mice (McHugh et al., 1996), given that many of the calcium transients
we measure likely result from calcium influx due to multiple action potentials. Place cells fired at higher
average rates than non-place-sensitive cells (mean 0.95 ± 0.02 events/sec, $n = 271$ versus 0.46 ± 0.01
events/sec, $n = 450$, respectively; significance level 2×10^{-81}, one-sided Student's t-test). The distribution
of place cell mean neural event rates during the session was skewed towards higher rates (Fig. 3A). The
distribution of mean neural activity rates, however (i.e. the rate of events weighted by the amplitude of
each calcium transient), was apparently more close to being log-normally distributed across cells (Fig. 3A),
with a skew towards lower activity contributed by lower activity non-place cells, consistent with data from
extracellular recordings (Buzsáki and Mizuseki, 2014). While a Kolmogorov-Smirnov test of the overall
distribution of activity rejected the null hypothesis of log-normal distribution due to this low tail ($p = 7$
$\times 10^{-7}$, $n = 721$), the log-normal model could not be rejected for the distribution of place cell activities (p
$= 0.7$, $n = 271$).

Figure 2. Place cell firing during navigation of a circular track. **(A)** Two-photon image from a typical CA1 imaging session. Green: GCaMP6s, red: mRuby. **(B)** 225 segmented ROIs from image in **A**. **(C)** Ca^{2+} transients for 50 (of 225) randomly selected cells from the region shown in **A**. Trace height is normalised to the 99th percentile of the $\Delta F/F$ for each cell. Magenta trace at top shows position of the mouse along the circular track. **(D)** Rastergram showing events detected for all 225 cells in this example, over a longer time period. Blue dots indicate the time of onset of each Ca^{2+} transient, with dot area showing relative event amplitude (normalised for each cell). Magenta trace indicates mouse position. **(E)** Spatial trajectory over 24 minutes of recording (gray line) together with locations of firing (blue dots) for five representative cells from the region shown in **A**. **(F)** Neuronal activity maps for the cells in **E** across repeated laps, obtained by integrating the inferred activity in each spatial bin, and normalising by the maximum for each cell. **(G)** Normalised neural event rate maps for 97 (of 225) cells from the region shown in **A** considered place-sensitive, sorted by place preference (peak location). **(H)** Normalized place field map for 271 place cells pooled from 6 fields of view (FOVs) from 3 mice, sorted by place preference. (I-K) Distributions over all cells in **H** of location of maximal activity **(I)**, place field size **(J)**, and spatial information (calculated via Skaggs approximation) **(K)**. Triangles in this and later plots denote mean values. Mean place field size ($\pm$ s.e.m.): 19.7 ± 0.4 cm, mean spatial information: 0.90 ± 0.02 bits/event.

The reliability of neuronal activity has often been measured in the sensory neuroscience literature (Tolhurst et al., 1983; van Steveninck et al., 1997) by calculating the ratio of the variance to the mean of the number of spikes fired in response to each identical stimulus presentation. We are not aware of a similar analysis in the hippocampal literature (Markus et al., 1995, performed a related but somewhat different analysis), however the degree of control our head-fixed preparation affords makes such an analysis feasible. We measured the mean and variance over repeated laps of the neural activity rate for each cell, as described in Methods. Fig. 3C shows the neural activity variance to mean relationships for a pair of cells (one a place cell, the other not), together with power law fits. The exponent of the power law fit (slope of the straight line on a double-logarithmic plot) captures the intrinsic variability of the cell's response. We found that places cells had systematically lower exponents (i.e. more reliable activity from lap to lap, taking into account the mean level of activity) than did non-place cells (Fig. 3D); mean exponent 1.72 ± 0.01, $n = 269$ for place cells, and 1.94 ± 0.01, $n = 448$ for non-place cells). Although place cells were significantly more reliable from trial to trial than non-place cells, our results suggest that CA1 activity, as measured from calcium fluorescence, is somewhat more variable from trial to trial than neocortical activity (Tolhurst et al., 1983).

3.3   Place cells remap almost completely between arenas

Place cells have the ability to change their activity patterns in different environments, a phenomenon known as global remapping. To determine whether place tuning in our floating track is environment-dependent, we imaged mice in a circular track with visual cues (track A) for up to 24 min then transferred them to a second circular track with distinct visual cues and additional tactile cues (track B) where they were imaged for up to 28 more minutes. We found that upon switching to track B, the location-specific activity of many cells disappeared, indicating remapping (Fig. 4A-C). We analysed 154 cells (from 3 imaging areas in 3 mice) which were active in both tracks. 58 of the cells had place fields in both tracks, only 3% of which retained their place field location upon switching to track B. The neuronal activity patterns in the second track could not be predicted from their neuronal activity in the first track - mean activity correlation between tracks was 0.02 ± 0.02. Moreover, place field locations in track B appeared to be randomly redistributed around the circular track relative to their positions in track A (failure to reject the null hypothesis of a circular uniform distribution of place field shifts, $p = 0.95$, Hodges-Ajne test, $n = 103$; Fig. 4C). Altogether, these results show global remapping as a result of visual and tactile cues providing sufficient sensory context for

Figure 3. CA1 place cells show higher and more reliable activity than non-place cells. **(A)** Histograms of calcium transient detected event rates for place cells (PC) and non-place-sensitive cells (NPC). Sample means for each group indicated by triangles. **(B)** Histograms of activity rates for place and non-place cells. Activity rates are calculated here by integrating all calcium transient events across a lap, weighted by the amplitude of each event, and dividing by the time taken for the mouse to complete the lap. **(C)** The variance of the activity rate (per unit time spent) in each 2-cm spatial bin as the mouse completed the lap plotted against the mean activity rate, for two cells, one place-sensitive and other not. Total least-squares regression fits are shown. This plot is double-logarithmic, and the slope of the linear fit thus indicates the exponent in the relationship between the variability and mean. **(D)** The distribution of variability exponents for place and non-place cells.

mice to identify the tracks as being different environments despite their identical geometry and dimensions.
Similar remapping of place fields in different environments has been previously reported in freely-moving
rodents. (Muller and Kubie, 1987; Anderson and Jeffery, 2003; Arriaga and Han, 2017).

3.4   Dynamic place fields over days

To study the stability of place fields over days, we imaged mice in the circular track multiple times over 8
days. Place cells had consistent location-specific activity within a session but the location was not constant
over days (Fig. 5A). For each mouse, there was a similar number of place cells across sessions and the set
of place fields for each session fully covered the length of the circular track (Fig. 5B). We found a total

Figure 4. Place cell remapping. **(A)** Spatial trajectory over 24 minutes of recording (gray line) and firing locations (blue dots) for three representative cells in a mouse that navigated two circular tracks with distinct visual and tactile cues. Note that spatial activity of cells in track A is not maintained in track B, indicating remapping. **(B)** Normalised neural event rate maps for track A place cells (PCs) during navigation in track A (left) and B (right) sorted according to place field locations in track A. Cells pooled from three imaging areas in three mice. **(C)** Normalised neural event rate maps for track B PCs during navigation in track A (left) and B (right) sorted according to place field locations in track B. **(D)** Shift in place field locations for place cells common to both tracks.

of 803 cells in the FOV across all imaging sessions, of which $88 \pm 2\%$ were active on any day (Fig. 5C), within the range of reported values for freely-moving mice (Gonzalez et al., 2019). On average, $29 \pm 3\%$ of the active cells in each session were place sensitive. Only $4 \pm 1\%$ of all cells were place sensitive in all sessions. If a cell was active on one day, the probability that the cell would be active in a later session (here called recurrence) did not change with time (Fig. 5D) within the time period that we examined (< 8 days). Imaging over a much longer timescale (30 days), Ziv et al. (2013) observed a decrease in the recurrence probability of active cells with time. If a cell was place-sensitive on one day, the probability of the cell being place-sensitive in a later session decreased with time (Fig. 5D). Of the place cells that retained the classification at a later session, the majority retained the place field location. These results are qualitatively similar to previous observations from freely-moving mice (Ziv et al., 2013). The decline in similarity of place fields (i.e. substantial remapping) across our imaging period (Fig. 5B) is comparable to the observed decline in VR studies imaged within a similar time period (Hainmueller and Bartos, 2018) and is more pronounced than the observed decline in freely-moving mice (Ziv et al., 2013; Gonzalez et al., 2019) imaged over a longer time period (10-30 days). Remapping of place fields over time was consistent with a decrease in place field correlations (mean correlation of the rate maps of place cells that retained their classification in two sessions N days apart, Fig. 5F), similar to reports for head-fixed mice in VR (Hainmueller and Bartos, 2018).

3.5 Place fields emerge in a floating open arena

In view of reports that spatial selectivity is impaired in two-dimensional virtual environments (Aghajan et al., 2015), we were interested to see whether 2D place fields could be observed in the floating track

Figure 5. Dynamic place fields. **(A)** Neuronal activity maps across laps in a session and across imaging days for 4 representative cells. **(B)** Normalised place field maps for place cells found on multiple days ordered by maximum location on different days; data from one representative mouse. **(C)** Proportion of all cells found in all imaging sessions that were active on each day and proportion of active cells on that day that were found to be place-sensitive. **(D)** Fraction of cells that were active or place-sensitive in one session which retained that classification at a later session (recurrence) versus number of days interval between the imaging sessions. **(E)** Distribution of place field shifts for 1 to 7 day intervals between imaging sessions. **(F)** Place field correlation for cells that retained their classification in two sessions versus number of days interval between the imaging sessions. (C-E) Data pooled from 4 imaging areas in 3 mice. Where an error bar is missing, the average was taken from a single imaged area.

system. To investigate this, we imaged mice navigating an open circular arena for 20-36 min. We found
44-301 (median: 231) active neurons per imaged FOV (490×490 μm, 4 FOVs in 3 mice, 807 cells in
total). 16-46% (median: 25%) of these cells showed location-specific activity characteristic of place cells
(Fig. 6A-B)). These activity patterns and place field maps are qualitatively similar to real-world systems
(Cacucci et al., 2007; Renaudineau et al., 2009; Ziv et al., 2013) and VR systems that allow rotations of
the head or body (Aronov and Tank, 2014; Chen et al., 2018). Some cells responded to the presence of
the wall of the arena where landmarks were located, firing near them (Fig. 6A), resembling border cells
that have been observed in other regions of the hippocampal formation in rats (Solstad et al., 2008; Lever
et al., 2009; Boccara et al., 2010). Overall, mice spent 62 ± 13% of the time in the outer 6 cm of the
arena and a greater proportion of cells had place fields with centroids located in this outer region than
in the inner region (64% vs 36%) (Fig. 6C). Place cells had a skewed distribution of field size with an
average of 170 ± 9 cm^2 or 21 ± 1% of the environment (Fig. 6D). This is comparable to values reported
for tetrode recordings in freely-moving mice (Cacucci et al., 2008) and in head-fixed mice navigating VR
systems which allow horizontal head rotations (Chen et al., 2018). The distribution of spatial information
was similarly skewed with an average of 0.95 ± 0.04 bits/event (Fig. 6E). This value is within the range
reported for freely-moving mice navigating real-world systems (Cacucci et al., 2008; Renaudineau et al.,
2009; Rochefort et al., 2011) and is greater than that reported for head-fixed mice navigating VR systems
with horizontal head rotation (Chen et al., 2018).

Figure 6. Place tuning for open field behaviour. (**A**) Spatial trajectory over 20-32 minutes of recording (gray line) together with firing locations (blue dots) for 5 representative cells from 3 mice. (**B**) Normalised place field maps for the cells in **A**. (**C-E**) Distributions over 204 cells (pooled from 4 imaging areas in 3 mice) of place field centroid location (**C**, inner: inner circle with 10.25 cm radius, outer: outer 6 cm of open arena), place field size (**D**), and spatial information (**E**). Mean place field size ($\pm$ s.e.m.): 170 ± 9 cm^2, mean spatial information: 0.95 ± 0.04 bits/event.

3.6 Spatial location is accurately encoded in low dimensions of the neural manifold

To study how spatial behaviour on the air-lifted platform is reflected in neural population dynamics, we performed a neural manifold analysis using the MDS technique. Applying this to data recorded from mice running in the circular track revealed distinct recapitulation of the spatial trajectory in the low order components of the neural manifold (typical example shown in Fig. 7A). This result was not specific to MDS, and very similar results were observed using PCA and Isomap (Supplementary Fig. 2). Open field exploration, being less constrained, unsurprisingly revealed much more complex dynamics on the manifold. Colour-coding points (corresponding to patterns of activity across the observed neurons) by angular location around the circular track revealed some clustering of points in manifold localities, spread however throughout the manifold (Fig. 7B; typical example). Colour-coding instead for radial position revealed additional structure at larger scales (Fig. 7C). Variance in the neural population dynamics was accounted for by a smaller number of components in circular track than open-field experiments (Fig. 7D); or equivalently, to account for a fixed amoount of variance required more components (dimensions) for mice exploring an open-field. Considering the number of components required to account for a fixed percentage (90%) of the variance as the "dimensionality" of the manifold ((Stringer et al., 2019)), we examined how

the dimensionality grew with the number of neurons incorporated within an ensemble. This grew at a
faster rate with ensemble size for the circular track than open-field environments (Fig. 7E), suggesting
that the greater complexity of the exploration trajectory was recapitulated in higher-dimensional neural
population dynamics. Finally, we assessed the extent to which spatial information could be decoded from
low-dimensional neural manifolds, by training an Optimal Linear Estimator (OLE) decoding algorithm
on the angular position data. Angular position could be decoded with high accuracy from a very small
number of components in the circular track case (Fig. 7F); in comparison, decoding of angular position in
the open-field task yielded lower fidelity estimates, and required more manifold dimensions.

4 DISCUSSION

We have here provided the first demonstration of hippocampal place tuning in head-fixed mice navigating
a real-world floating track. Our results indicate reliable place fields in a one-dimensional circular track,
comparable to those observed in freely moving (McHugh et al., 1996; Chen et al., 2013) and head-fixed
mice navigating a virtual reality apparatus (Dombeck et al., 2010), with spatial information rates similar to
both (Arriaga and Han, 2017; Mou et al., 2018; Gonzalez et al., 2019). When mice re-entered the same
environment, place fields were recapitulated, with fields dynamically changing over the course of days,
with a rate of reconfiguration similar to that observed in VR recordings (Hainmueller and Bartos, 2018),
but significantly faster than that seen in real-world free exploration (Ziv et al., 2013; Gonzalez et al., 2019).

We also demonstrated the presence of 2D place fields in an open arena, similar to those seen in mice
during free exploration (McHugh et al., 1996; Cacucci et al., 2007; Renaudineau et al., 2009; Ziv et al.,
2013) and VR systems that allow rotations of the head or body (Aronov and Tank, 2014; Chen et al.,
2018), with spatial information rates similar to the former. We initially found this surprising, as the lack of
vestibular sensory input might be expected to impair place cell formation (Stackman et al., 2002; Aghajan
et al., 2015). This suggests that perhaps the presence of multimodal sensory information goes some way
to overcome the lack of vestibular signals in the floating track, i.e. the combination of information from
different sources may be an important aspect of place cell formation, rather than there being a strict
requirement for vestibular cues. We speculate that the lack of vestibular information may likewise affect
the properties of the motion-dependent theta rhythm in the floating track system, similar to observations
from VR (Aronov and Tank, 2014; Ravassard et al., 2015; Chen et al., 2018), although this has yet to be
verified. Many hippocampal researchers are skeptical of the utility of VR for studying spatial navigation,
with Donato and Moser (2016) commenting on the Aghajan et al. (2015) results: "These data cast doubt on
whether the way animals interpret 2D or 3D space can be ever be understood using VR under conditions
of head or body restriction. Strategies that compensate for the loss of synchrony between vestibular
information and the animal's behaviour would be a welcome advance". Our results here suggest that the
availability of a richer, multimodal sensory scene can go some way towards addressing this issue.

The floating track system has several advantages over VR systems. Notably, it allows the expression of 2D
place cells in head-fixed setups without the need for a complex commutator headplate (Chen et al., 2018),
making two-photon imaging straightforward. In addition, although not as dynamically reconfigurable as
pure VR systems, the floating track system employs multimodal sensory stimuli (tactile, auditory, olfactory)
which contribute to the integrated spatial context (Ravassard et al., 2015; Geva-Sagiv et al., 2015), making
it adaptable to a variety of tasks.

Compared to real-world free exploration, place coding in the floating track system does not persist for
multiple days to the same degree, similar to VR systems. Another limitation of the floating track system

Figure 7. Neural manifold analysis reveals mouse spatial trajectories in a low-dimensional subspace of the neural population dynamics. (**A**) Multidimensional Scaling (MDS) analysis of a typical circular track experiment. The first three components shown in three-dimensional projection at left, with 2D projections at right. Each dot reflects the projection of the neural population activity at a given time point; colour code indicates spatial location in degrees of angle around the track. (**B**) MDS depiction of a typical open field experiment. In this case the colour code is as in (A), reflecting angular but not radial location. (**C**) Remapping the colour code to reflect normalised radial location indicates additional manifold structure relating to spatial position. (**D**) The variance explained by inclusion of an increasing number of manifold dimensions for the circular track and open field environments. (**E**) The growth of dimensionality, defined as the number of components needed to capture 90% of the variance, with the number of cells included in the manifold. (**F**) Decoding performance, measured by Pearson correlation coefficient, for circular track and open field environments.

compared to real-world free exploration is that it is constrained to relatively small environments - the

longest track we used in this study was 1 metre in length, although we have designed environments that
work within the current setup that allow tracks as long as 2 metres. Although not as long as the 18 metre
tracks of Brun et al. (2008), this should be well within the ecological regime for mice. The floating track
system has one distinct advantage over real-world behavioural tasks in allowing head-fixation, making it
compatible with a wider range of optical imaging technologies and allowing for straightforward extension
to intracellular recording.

This study focused on CA1 hippocampal place cell representations during exploration. The floating
track system presented here offers advantages over traditional VR systems and free exploration real-world
systems for investigating neural encoding in a variety of brain systems, and may well prove a useful tool for
dissecting neural pathways underpinning cognitive functions based upon spatial representations, such as
spatial working memory tasks (Lalonde, 2002; Dudchenko, 2004) and object location and recognition tasks
(Vogel-Ciernia and Wood, 2014). Floating track behavioural tests may occupy a useful position between
virtual reality paradigms, which allow head-fixation but are generally limited to simplistic unimodal sensory
stimulation, and real-world behavioural tasks.

We found that approximately 88% of the CA1 cells we detected were active (fired at least once) in
any given session (see Fig. 5C). If a cell was active in a given session, then it was around 90% likely
to be active in any subsequent session, with no obvious temporal dependence in a timescale of several
days. Approximately 29% of detected cells were place-sensitive (see section 3.2). One caveat concerning
multiphoton calcium imaging in sparsely firing areas such as hippocampal region CA1 is that accurate
identification of the boundaries of ROIs corresponding to single cells depends upon the cells firing at
least once. In our data analysis here, we used the CaImAn software package (Giovannucci et al., 2019) to
extract ROIs, together with extensive modifications to the data processing pipeline to improve registration
of ROIs across sessions on our dataset, as well as incorporating additional neuropil decontamination. We
have also used ABLE (Reynolds et al., 2017) in place of CaImAn in this pipeline with similar results. In
both of these software tools, cells are registered only if they are active at some point in the set of sessions
imaged. This suggests that we are likely to be under-estimating the total number of neurons in the field
of view, and thus over-estimating the fraction of active and place-tuned cells. It should also be noted that
the non-place-tuned cells we observed may include cells that might show place tuning in a freely moving
(non head-fixed) scenario. We additionally note that comparison of results from multiphoton imaging with
electrophysiological studies can be complicated by both inherent biases in cell selection, as described
above, and by the different observation model due to calcium measurements, which has been suggested to
lead to a lower fraction of responsive neurons, but sharper selectivity (Siegle et al., 2020).

The visual stimulus apparatus we used in this study consisted of phosphorescent tape formed into
patterned structures on the walls of the floating track. This resulted in scotopic illumination conditions,
which correspond to decreased spatial acuity (Umino et al., 2008). However, this is not a fundamental
aspect of the platform, and instead photopic illumination could have been used, together with higher walls
and a cone around the objective lens to exclude light from the microscope emission path.

A further caveat to our study is the difference in the age and gender profiles of the mice imaged in the
circular track and open field tasks, with younger, female mice used in the latter task. While there is data
suggesting no gender effect should be expected (Fritz et al., 2017), we might expect lower spatial selectivity
and reduced place field stability (Yan et al., 2003) in the older mice we imaged in the circular track. We
can, thus, view our results in the circular track as conservative with regard to the tuning and stability that
might be expected in younger mice, while demonstrating the applicability of the platform to the study of
neurodegenerative models which require the use of older mice.

The idea that coding in CA1 and other brain areas is dynamic - i.e. that the ensemble of cells encoding a
given memory evolves over time - is now relatively well-established (Ziv et al., 2013; Rubin et al., 2015;
Hainmueller and Bartos, 2018). Our results suggest that the time constant of this reconfiguration may
depend upon environmental factors and the number and type of sensory cues being integrated, with head-
fixed preparations including virtual-reality and floating track paradigms inducing faster reconfiguration,
while memories formed during freely moving behaviour may induce cell assemblies that are stable for
longer durations, as observed in Ziv et al. (2013). (However, noting the caveat above, it is possible that
greater place field stability would have been observed in younger mice). Interestingly, similar changes
in the specific ensemble of neurons encoding a sensory stimulus have been observed in the olfactory
bulb (Kato et al., 2012), whereas motor cortex seems to show no such effect (Peters et al., 2014). This
underscores the importance of further research on the specific parameters affecting the stability of neural
representations employed in perception, memory and cognitive behaviour, with technology such as that
described here now making practical the longitudinal experiments necessary to address this issue.

The relatively high reproducibility of stimulus conditions in the circular track task allowed us to make use
of a trial-to-trial variability analysis method widely employed in visual neuroscience, which has however
not, to our knowledge, previously been employed to study neural representations in spatial memory. This
led to the interesting result that while place cells in CA1 fire at higher rates than non-place cells, their
activity is in fact more reliable from trial to trial, given the firing rate - i.e. that while neural response
variance increases with the mean level of activity, it does so with a lower exponent for place cells than
for non-place cells, indicating comparative reliability. We speculate that, like spatial information (Cacucci
et al., 2008), the response variability exponent may turn out to be a sensitive indicator of neurodegeneration
under conditions of aberrant excitability (Busche et al., 2008).

As well as examining single cell response properties, we used a manifold learning approach to study the
behaviour of the entire ensemble of recorded cells during spatial exploration. We used the Multidimensional
Scaling (MDS) manifold learning approach, after observing that it performed better than Principal
Components Analysis (PCA) in terms of variance accounted for, however we verified qualitatively that a
similar picture was provided by both PCA and Isomap (see Supplementary Fig. 2), and expect our results
to be robust to any sufficiently comprehensive manifold learning approach. The spatial position of the
mouse in the 1D (circular track) task was visually apparent in the lowest dimensions of the neural manifold;
while this was less immediately obvious (because of the non-cyclical nature of the exploratory trajectory)
in the 2D task, a decoding analysis indicated that for the open field task also, spatial location could be
decoded with some success from a relatively small number of manifold dimensions, although with much
less fidelity than for the circular track. As might be expected given the more constrained nature of the
task, the dimensionality of the neural manifold obtained in mice exploring the circular track was lower
than that observed in mice exploring the open field. The use of manifold learning in conjunction with a
floating real-world environment may provide a useful tool for uncovering fundamental principles governing
encoding for memory, perception and cognition.

In this paper we have validated a platform technology which, by combining multiphoton calcium imaging
in head-fixed mice with floating track implementation of a behavioural task, allows robust and sensitive
readout of memory encoding and retrieval performance from the activity of hippocampal neurons. As well
as enabling advances in cognitive neuroscience, we expect this tool to be of great utility for the study of
mouse models of neurodegenerative disorders, and a powerful tool for aiding the pre-clinical development
of therapeutics for these diseases.

CONFLICT OF INTEREST STATEMENT

500 The authors declare that the research was conducted in the absence of any commercial or financial
501 relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

502 MAG and SRS designed the experiments. MAG worked out the technical details of the experiments. MAG
503 and JR performed the experiments with support from YL. MAG processed the experimental data and
504 performed the analysis with support from CD, GPG, SP and SRS. MAG and SRS wrote the manuscript.
505 MAG and SRS supervised the project. All authors discussed the results and commented on the manuscript.

FUNDING

506 This work was supported by BBSRC award BB/R022437/1, by the EPSRC CDT in Neurotechnology
507 for Life and Health (EPSRC award EP/L016737/1), by the Universities UK Rutherford Fund (award
508 RF-2018-27), and by Mrs Anne Uren and the Michael Uren Foundation.

ACKNOWLEDGMENTS

509 We thank David Dupret, Richard Morris and Leo Khiroug for useful discussions, and Sihao Lu for his
510 assistance with the Matlab adaptation of the FISSA toolbox.

SUPPLEMENTAL DATA

511 The Supplementary Material for this article can be found in the attached Supplementary Material file.

REFERENCES

Aghajan, Z. M., Acharya, L., Moore, J. J., Cushman, J. D., Vuong, C., and Mehta, M. R. (2015). Impaired
spatial selectivity and intact phase precession in two-dimensional reality. *Nat. Neurosci.* 18, 121–128.
doi:10.1038/nn.3884
Anderson, M. I. and Jeffery, K. J. (2003). Heterogeneous modulation of place cell firing by changes in
context. *J. Neurosci.* 23, 8827–8835. doi:10.1523/JNEUROSCI.23-26-08827.2003
Aronov, D. and Tank, D. W. (2014). Engagement of neural circuits underlying 2d spatial navigation in a
rodent virtual reality system. *Neuron* 84, 442–456. doi:10.1016/j.neuron.2014.08.042
Arriaga, M. and Han, E. B. (2017). Dedicated hippocampal inhibitory networks for locomotion and
immobility. *J. Neurosci.* 37, 9222–9238. doi:doi:10.1523/jneurosci.1076-17.2017
Benjamini, Y., Fonio, E., Galili, T., Havkin, G. Z., and Golani, I. (2011). Quantifying the buildup in
extent and complexity of free exploration in mice. *Proc. Natl. Acad. Sci. U.S.A.* 108, 15580–15587.
doi:10.1073/pnas.1014837108
Boccara, C. N., Sargolini, F., Thoresen, V. H., Solstad, T., Witter, M. P., Moser, E. I., et al. (2010). Grid
cells in pre- and parasubiculum. *Nat. Neurosci.* 13, 987–994. doi:https://doi.org/10.1038/nn.2602
Brun, V. H., Solstad, T., Kjelstrup, K. B., Fyhn, M., Witter, M. P., Moser, E. I., et al. (2008). Progressive
increase in grid scale from dorsal to ventral medial entorhinal cortex. *Hippocampus* 18, 1200–1212

Busche, M. A., Eichhoff, G., Adelsberger, H., Abramowski, D., Wiederhold, K.-H., Haass, C., et al. (2008).
Clusters of hyperactive neurons near amyloid plaques in a mouse model of alzheimer's disease. *Science*
321, 1686–1689

Buzsáki, G. and Mizuseki, K. (2014). The log-dynamic brain: how skewed distributions affect network
operations. *Nat. Rev. Neurosci.* 15, 264–278

Cacucci, F., Wills, T. J., Lever, C., Giese, K. P., and O'Keefe, J. (2007). Experience-dependent
increase in CA1 place cell spatial information, but not spatial reproducibility, is dependent on the
autophosphorylation of the isoform of the calcium/calmodulin-dependent protein kinase ii. *J. Neurosci.*
27, 7854 –7859

Cacucci, F., Yi, M., Wills, T. J., Chapman, P., and O'Keefe, J. (2008). Place cell firing correlates with
memory deficits and amyloid plaque burden in tg2576 Alzheimer mouse model. *Proc. Natl. Acad. Sci.*
U.S.A. 105, 7863–7868

Chen, G., King, J. A., Burgess, N., and O'Keefe, J. (2013). How vision and movement combine in the
hippocampal place code. *Proc. Natl. Acad. Sci. U.S.A.* 110, 378–383. doi:10.1073/pnas.1215834110

Chen, G., King, J. A., Lu, Y., Cacucci, F., and Burgess, N. (2018). Spatial cell firing during virtual
navigation of open arenas by head-restrained mice. *eLIFE* 7, e34789. doi:10.7554/eLife.34789

Chen, G., Lu, Y., King, J. A., Cacucci, F., and Burgess, N. (2019). Differential influences of environment
and self-motion on place and grid cell firing. *Nature Communications* 10, 1–11

Dombeck, D. A., Harvey, C. D., Tian, L., Looger, L. L., and Tank, D. W. (2010). Functional imaging of
hippocampal place cells at cellular resolution during virtual navigation. *Nat. Neurosci.* 13, 1433–1440.
doi:10.1038/nn.2648

Dombeck, D. A., Khabbaz, A. N., Collman, F., Adelman, T. L., and Tank, D. W. (2007). Imaging
large-scale neural activity with cellular resolution in awake, mobile mice. *Neuron* 56, 43–57

Donato, F. and Moser, E. I. (2016). A world away from reality. *Nature* 533, 325–326

Dudchenko, P. A. (2004). An overview of the tasks used to test working memory in rodents. *Neuroscience*
& Biobehavioral Reviews 28, 699–709

Dupret, D., O'Neill, J., Pleydell-Bouverie, B., and Csicsvari, J. (2010). The reorganization and reactivation
of hippocampal maps predict spatial memory performance. *Nat. Neurosci.* 13, 995–1002. doi:10.1038/
nn.2599

Ergorul, C. and Eichenbaum, H. (2004). The hippocampus and memory for "what,""where," and "when".
Learning & Memory 11, 397–405

Friedrich, J., Yang, W., Soudry, D., Mu, Y., Ahrens, M. B., Yuste, R., et al. (2017). Multi-scale approaches
for high-speed imaging and analysis of large neural populations. *PLOS Comput. Biol.* 13, e1005685.
doi:10.1371/journal.pcbi.1005685

Fritz, A.-K., Amrein, I., and Wolfer, D. P. (2017). Similar reliability and equivalent performance of female
and male mice in the open field and water-maze place navigation task. In *American Journal of Medical*
Genetics Part C: Seminars in Medical Genetics (Wiley Online Library), vol. 175, 380–391

Geva-Sagiv, M., Las, L., Yovel, Y., and Ulanovsky, N. (2015). Spatial cognition in bats and rats: from
sensory acquisition to multiscale maps and navigation. *Nat. Rev. Neurosci.* 16, 94–108. doi:10.1038/
nrn3888

Giovannucci, A., Friedrich, J., Gunn, P., Kalfon, J., Brown, B. L., Koay, S. A., et al. (2019). CaImAn an
open source tool for scalable calcium imaging data analysis. *eLife* 8, 38173. doi:10.7554/eLife.38173

Gonzalez, W. G., Zhang, H., Harutyunyan, A., and Lois, C. (2019). Persistence of neuronal representations
through time and damage in the hippocampus. *Science* 365, 821–825. doi:DOI:10.1126/science.aav9199

Hainmueller, T. and Bartos, M. (2018). Parallel emergence of stable and dynamic memory engrams in the
hippocampus. *Nature* 558, 292–296. doi:10.1038/s41586-018-0191-2

Harvey, C. D., Collman, F., Dombeck, D. A., and Tank, D. W. (2009). Intracellular dynamics of
hippocampal place cells during virtual navigation. *Nature* 461, 941–946. doi:10.1038/nature08499

Kato, H. K., Chu, M. W., Isaacson, J. S., and Komiyama, T. (2012). Dynamic sensory representations in
the olfactory bulb: modulation by wakefulness and experience. *Neuron* 76, 962–975

Keemink, S. W., Lowe, S. C., Pakan, J. M. P., Dylda, E., van Rossum, M. C. W., and Rochefort, N. L.
(2018). FISSA: A neuropil decontamination toolbox for calcium imaging signals. *Sci. Rep.* 8, 3493.
doi:10.1038/s41598-018-21640-2

Kislin, M., Mugantseva, E., Molotkov, D., Kulesskaya, N., Khirug, S., Kirilkin, I., et al. (2014). Flat-floored
air-lifted platform: a new method for combining behavior with microscopy or electrophysiology on
awake freely moving rodents. *J. Vis. Exp.* 88, e51869. doi:10.3791/51869

Lalonde, R. (2002). The neurobiological basis of spontaneous alternation. *Neuroscience & Biobehavioral*
Reviews 26, 91–104

Lee, I. and Knierim, J. J. (2007). The relationship between the field-shifting phenomenon and
representational coherence of place cells in CA1 and CA3 in a cue-altered environment. *Learn Mem.* 14,
807–815. doi:10.1101/lm.706207

Leutgeb, S., Leutgeb, J. K., Barnes, C. A., Moser, E. I., McNaughton, B. L., and Moser, M.-B. (2005).
Independent codes for spatial and episodic memory in hippocampal neuronal ensembles. *Science* 309,
619–623

Lever, C., Burton, S., Jeewajee, A., O'Keefe, J., and Burgess, N. (2009). Boundary vector cells in the
subiculum of the hippocampal formation. *J. Neurosci.* 29, 9771–9777. doi:10.1523/jneurosci.1319-09.
2009

Mably, A. J., Gereke, B. J., Jones, D. T., and Colgin, L. L. (2017). Impairments in spatial representations
and rhythmic coordination of place cells in the 3xTg mouse model of Alzheimer's disease. *Hippocampus*
27, 378–392

Markus, E. J., Qin, Y.-L., Leonard, B., Skaggs, W. E., McNaughton, B. L., and Barnes, C. A. (1995).
Interactions between location and task affect the spatial and directional firing of hippocampal neurons. *J.*
Neurosci. 15, 7079–7094

McHugh, T. J., Blum, K. I., Tsien, J. Z., Tonegawa, S., and Wilson, M. A. (1996). Impaired hippocampal
representation of space in ca1-specific nmdar1 knockout mice. *Cell* 87, 1339–1349

Mehta, M. R., Barnes, C. A., and Mcnaughton, B. L. (1997). Experience-dependent, asymmetric expansion
of hippocampal place fields. *Proc. Natl. Acad. Sci. U.S.A.* 94, 8918–8921

Mou, X., Cheng, J., Yu, Y. S. W., Kee, S. E., and Ji, D. (2018). Comparing mouse and rat hippocampal
place cell activities and firing sequences in the same environments. *Front. Cell. Neurosci.* 12, 1–13.
doi:10.3389/fncel.2018.00332

Muller, R. U. and Kubie, J. L. (1987). The effects of changes in the environment on the spatial firing of
hippocampal complex-spike cells. *J. Neurosci.* 7, 1951–l968. doi:10.1523/JNEUROSCI.07-07-01951.
1987

Muller, R. U. and Kubie, J. L. (1989). The firing of hippocampal place cells predicts the future position of
freely moving rats. *J. Neurosci.* 9, 4101–10

Muzzu, T., Mitolo, S., Gava, G. P., and Schultz, S. R. (2018). Encoding of locomotion kinematics in the
mouse cerebellum. *PloS ONE* 13, e0203900

Nashaat, M. A., Oraby, H., Sachdev, R. N. S., and Winter, Y. (2016). Air-track: a real-world floating
environment for active sensing in head-fixed mice. *J. Neurophysiol.* 116, 1542–1553. doi:10.1152/jn.
00088.2016

O'Keefe, J. (2007). Hippocampal neurophysiology in the behaving animal. In *The Hippocampus Book*, ed.
P. Anderson (Oxford: Oxford University Press). 475–548

O'Keefe, J. and Dostrovsky, J. (1971). The hippocampus as a spatial map. preliminary evidence from unit
activity in the freely-moving rat. *Brain Res.* 34, 171–175

Ozbay, B. N., Futia, G. L., Ma, M., Bright, V. M., Gopinath, J. T., Hughes, E. G., et al. (2018). Three
dimensional two-photon brain imaging in freely moving mice using a miniature fiber coupled microscope
with active axial-scanning. *Scientific reports* 8, 1–14

Peron, S., Chen, T.-W., and Svoboda, K. (2015). Comprehensive imaging of cortical networks. *Curr. Opin.*
in Neurobiol. 32, 115–123

Peters, A. J., Chen, S. X., and Komiyama, T. (2014). Emergence of reproducible spatiotemporal activity
during motor learning. *Nature* 510, 263–267

Pilz, G.-A., Carta, S., Stäuble, A., Ayaz, A., Jessberger, S., and Helmchen, F. (2016). Functional imaging
of dentate granule cells in the adult mouse hippocampus. *J. Neurosci.* 36, 7407–7414. doi:10.1523/
JNEUROSCI.3065-15.2016

Ravassard, P., Kees, A., Willers, B., Ho, D., Aharon, D., Cushman, J., et al. (2015). Multisensory control
of hippocampal spatiotemporal selectivity. *Sci. Rep.* 340, 16171. doi:10.1126/science.1232655

Renaudineau, S., Poucet, B., Laroche, S., Davis, S., and Save, E. (2009). Impaired long-term stability of
CA1 place cell representation in mice lacking the transcription factor zif268/egr1. *Proc. Natl. Acad. Sci.*
U.S.A. 106, 11771–11775. doi:10.1073/pnas.0900484106

Reynolds, S., Abrahamsson, T., Schuck, R., Sjöström, P. J., Schultz, S. R., and Dragotti, P. L. (2017). Able:
An activity-based level set segmentation algorithm for two-photon calcium imaging data. *Eneuro* 4

Rickgauer, J. P., Deisseroth, K., and Tank, D. W. (2014). Simultaneous cellular-resolution optical
perturbation and imaging of place cell firing fields. *Nat. Neurosci.* 17, 1342–1346. doi:10.1038/nn.3866

Rochefort, C., Arabo, A., André, M., Poucet, B., Save, E., and Rondi-Reig, L. (2011). Cerebellum shapes
hippocampal spatial code. *Science* 334, 385–389. doi:10.1126/science.1207403

Rubin, A., Geva, N., Sheintuch, L., and Ziv, Y. (2015). Hippocampal ensemble dynamics timestamp events
in long-term memory. *Elife* 4, e12247

Samson, A. L., Ju, L., Kim, H. A., Zhang, S. R., Lee, J. A. A., Sturgeon, S. A., et al. (2015). Mousemove:
an open source program for semi-automated analysis of movement and cognitive testing in rodents. *Sci.*
Rep. 5, 16171. doi:10.1038/srep16171

Schultz, S. R., Copeland, C. S., Foust, A. J., Quicke, P., and Schuck, R. (2016). Advances in two-photon
scanning and scanless microscopy technologies for functional neural circuit imaging. *Proc. IEEE* 105,
139–157

Siegle, J. H., Ledochowitsch, P., Jia, X., Millman, D., Ocker, G. K., Caldejon, S., et al. (2020). Reconciling
functional differences in populations of neurons recorded with two-photon imaging and electrophysiology.
bioRxiv

Skaggs, W. E., McNaughton, B. L., and Gothard, K. M. (1992). An information-theoretic approach to
deciphering the hippocampal code. In *NIPS*. 1030–1037

Solstad, T., Boccara, C. N., Kropff, E., Moser, M.-B., and Moser, E. I. (2008). Representation of geometric
borders in the entorhinal cortex. *Science* 322, 1865–1868. doi:10.1126/science.1166466

Stackman, R. W., Clark, A. S., and Taube, J. S. (2002). Hippocampal spatial representations require
vestibular input. *Hippocampus* 12, 291–303. doi:10.1002/hipo.1112

Stringer, C., Pachitariu, M., Steinmetz, N., Carandini, M., and Harris, K. D. (2019). High-dimensional
geometry of population responses in visual cortex. *Nature* 571, 361–365

Taube, J. S. (1995). Place cells recorded in the parasubiculum of freely moving rats. *Hippocampus* 5,
569–583. doi:10.1002/hipo.450050608

Tolhurst, D. J., Movshon, J. A., and Dean, A. F. (1983). The statistical reliability of signals in single
neurons in cat and monkey visual cortex. *Vis. Res.* 23, 775–785

Trouche, S., Perestenko, P. V., van de Ven, G. M., Bratley, C. T., McNamara, C. G., Campo-Urriza, N.,
et al. (2016). Recoding a cocaine-place memory engram to a neutral engram in the hippocampus. *Nat.*
Neurosci. 19, 564–567. doi:10.1038/nn.4250

Umino, Y., Solessio, E., and Barlow, R. B. (2008). Speed, spatial, and temporal tuning of rod and cone
vision in mouse. *J. Neurosci.* 28, 189–198. doi:10.1523/JNEUROSCI.3551-07.2008

van Steveninck, R. R. d. R., Lewen, G. D., Strong, S. P., Koberle, R., and Bialek, W. (1997). Reproducibility
and variability in neural spike trains. *Science* 275, 1805–1808

Vogel-Ciernia, A. and Wood, M. A. (2014). Examining object location and object recognition memory in
mice. *Current protocols in neuroscience* 69, 8–31

Yan, J., Zhang, Y., Roder, J., and McDonald, R. J. (2003). Aging effects on spatial tuning of hippocampal
place cells in mice. *Experimental brain research* 150, 184–193

Ziv, Y., Burns, L. D., Cocker, E. D., Hamel, E. O., Ghosh, K. K., Kitch, L. J., et al. (2013). Long-term
dynamics of CA1 hippocampal place codes. *Nat. Neurosci.* 16, 264

Zong, W., Wu, R., Li, M., Hu, Y., Li, Y., Li, J., et al. (2017). Fast high-resolution miniature two-photon
microscopy for brain imaging in freely behaving mice. *Nature methods* 14, 713–719

Supplementary Material

Figure S1. Power law decay of luminance of phosphorescent tapes coloured (**A**) green (peak 520 nm) and (**B**) blue (peak 500 nm). Dashed lines denote curve fits of the form $y = ax^{\beta}$. The luminance over an imaging period of an hour is well within the sensitivity range for mice, within the scotopic (low-light) range (Umino et al., 2008).

Figure S2. Neural manifolds extracted using (**A**) MDS, (**B**) PCA and (**C**) Isomap are very similar. Example shown here is the same circular track recording shown in the main manuscript. While MDS systematically captures at least as much of the variance as PCA (normally more) for the same number of dimensions, it also shows residual sructure related to mouse spatial position beyond the second dimension, which is not apparent in PCAof Isomap. Isomap provides a very similar depiction to MDS and PCA, although the "variance accounted for" quantification of manifold performance does not directly apply in this case. Our conclusion is that the precise technique used to visualise the manifold is not crucial.