An engineered decoy receptor for SARS-CoV-2 broadly binds protein S sequence variants

Kui K. Chan[1], Timothy J.C. Tan[2], Krishna K. Narayanan[2] and Erik Procko[2]

[1] Orthogonal Biologics, Champaign IL 61821

[2] Department of Biochemistry and Cancer Center at Illinois, University of Illinois, Urbana IL 61801

Correspondence: procko@illinois.edu

ABSTRACT

The spike S of SARS-CoV-2 recognizes ACE2 on the host cell membrane to initiate entry. Soluble decoy receptors, in which the ACE2 ectodomain is engineered to block S with high affinity, potently neutralize infection and, due to close similarity with the natural receptor, hold out the promise of being broadly active against virus variants without opportunity for escape. Here, we directly test this hypothesis. We find an engineered decoy receptor, sACE2$_2$.v2.4, tightly binds S of SARS-associated viruses from humans and bats, despite the ACE2-binding surface being a region of high diversity. Saturation mutagenesis of the receptor-binding domain followed by in vitro selection, with wild type ACE2 and the engineered decoy competing for binding sites, failed to find S mutants that discriminate in favor of the wild type receptor. We conclude that resistance to engineered decoys will be rare and that decoys may be active against future outbreaks of SARS-associated betacoronaviruses.

INTRODUCTION

Zoonotic coronaviruses have crossed over from animal reservoirs multiple times in the past two decades, and it is almost certain that wild animals will continue to be a source of devastating outbreaks. Unlike ubiquitous human coronaviruses responsible for common respiratory illnesses, these zoonotic coronaviruses with pandemic potential cause serious and complex diseases, in part due to their tissue tropisms driven by receptor usage. Severe Acute Respiratory Syndrome Coronaviruses 1 (SARS-CoV-1) and 2 (SARS-CoV-2) engage angiotensin-converting enzyme 2 (ACE2) for cell attachment and entry (*1-7*). ACE2 is a protease responsible for regulating blood volume and pressure that is expressed on the surface of cells in the lung, heart and gastrointestinal tract, among other tissues (*8*, *9*). The ongoing spread of SARS-CoV-2 and the disease it causes, COVID-19, has had a crippling toll on global healthcare systems and economies, and effective treatments and vaccines are urgently needed.

As SARS-CoV-2 becomes endemic in the human population, it has the potential to mutate and undergo genetic drift and recombination. To what extent this will occur as increasing numbers of people are infected and mount counter immune responses is unknown, but already a variant in the viral spike protein S (D614G) has rapidly emerged from multiple independent events and effects S protein stability and dynamics (*10*, *11*). Another S variant (D839Y) became prevalent in Portugal, possibly due to a founder effect (*12*). Coronaviruses have moderate to high mutation rates. For example, 10^{-4} substitutions per year per site occur in HCoV-NL63 (*13*), an alphacoronavirus that also binds ACE2, albeit via a smaller interface that is only partially shared with the RBDs of SARS-associated betacoronaviruses (*14*). Additionally, large changes in coronavirus genomes have frequently occurred in nature from recombination events, especially in bats where co-infection levels can be high (*15*, *16*). Recombination of MERS-CoVs has also been documented in camels (*17*). This will all have profound implications for the current pandemic's trajectory, the potential for future coronavirus pandemics, and whether drug resistance in SARS-CoV-2 becomes prevalent.

The viral spike is a vulnerable target for neutralizing monoclonal antibodies that are progressing
through clinical trials, yet in tissue culture escape mutations in the spike rapidly emerge to all antibodies
tested (*18*). Deep mutagenesis of the isolated receptor-binding domain (RBD) by yeast surface display
has easily identified mutations in S that retain high expression and ACE2 affinity, yet are no longer bound
by monoclonal antibodies and confer resistance (*19*). This has motivated the development of cocktails of
non-competing monoclonals (*18, 20*), inspired by lessons learned from the treatment of HIV-1 and Ebola,
to limit the possibilities for the virus to escape. Notably, drug maker Eli Lilly has a monoclonal
monotherapy (LY-CoV555) in advanced trials (NCT04427501) where the emergence of resistant virus
variants has occurred; the trial has been updated to include an arm with a second monoclonal (LY-
CoV016). However, even the use of monoclonal cocktails does not address future coronavirus spill overs
from wild animals that may be antigenically distinct. Indeed, large screening efforts were required to find
antibodies from recovered SARS-CoV-1 patients that cross-react with SARS-CoV-2 (*21*), indicating
antibodies have confined capacity for interacting with variable epitopes on the spike surface, and are
unlikely to be broad and pan-specific for all SARS-related viruses.

An alternative protein-based antiviral to monoclonal antibodies is to use soluble ACE2 (sACE2) as a
decoy to compete for receptor-binding sites on the viral spike (*6, 22-25*). In principle, the virus has
limited potential to escape sACE2-mediated neutralization without simultaneously decreasing affinity for
the native ACE2 receptor, rendering the virus less virulent. Multiple groups have now engineered sACE2
to create high affinity decoys for SARS-CoV-2 that rival matured monoclonal antibodies and potently
neutralize infection (*25-27*). In our group, deep mutagenesis was used to identify a large number of
mutations in ACE2 that increase affinity for S (*25*). These mutations were dispersed across the interface
and also at distal sites where they are predicted to enhance folding of the virus-recognized conformation.
A combination of three mutations, called sACE2$_2$.v2.4, increases affinity 35-fold and binds SARS-CoV-2
S (K$_D$ 600 pM) with affinity comparable to the best monoclonal antibodies (*25*). Even tighter apparent
affinities are reached through avid binding to trimeric spike expressed on a membrane. Despite
engineering being focused exclusively on SARS-CoV-2 affinity, sACE2$_2$.v2.4 potently neutralized
authentic SARS-CoV-1 and -2 infection in tissue culture, suggesting it's close resemblance to the wild
type receptor confers broad activity against ACE2-utilizing betacoronaviruses generally. Soluble
ACE2$_2$.v2.4 is dimeric and monodisperse without aggregation, catalytically active, highly soluble, stable
after storage at 37°C for days, and well expressed at levels greater than the wild type protein. Due to both
its high activity and favorable properties for manufacture, sACE2$_2$.v2.4 is a genuine drug candidate for
preclinical development.

Engineered, high affinity decoy receptors, while very similar to natural ACE2, nonetheless have
mutations present at or near the interaction surface. There is therefore an opportunity for viral spike
variants to discriminate between an engineered decoy and wild type receptors, providing a route towards
resistance. Here, we show that the engineered decoy sACE2$_2$.v2.4 binds broadly and tightly to the RBDs
of diverse SARS-associated betacoronaviruses that use ACE2 for entry. We further fail to find mutations
within the RBD, which directly contacts ACE2 and is where possible escape mutations will most likely
reside, that redirect specificity towards the wild type receptor. We conclude that resistance to an
engineered decoy receptor will be rare, and sACE2$_2$.v2.4 targets common attributes for affinity to S in
SARS-associated viruses.

RESULTS

An engineered decoy receptor broadly binds RBDs from SARS-associated CoVs with tight affinity

The affinities of the decoy receptor sACE2$_2$.v2.4 were determined for purified RBDs from the S
proteins of five coronaviruses from *Rhinolophus* bat species (isolates LYRa11, Rs4231, Rs7327, Rs4084
and RsSHC014) and two human coronaviruses, SARS-CoV-1 and SARS-CoV-2. These viruses fall

within a common clade of betacoronaviruses that use ACE2 as an entry receptor (*7*). They share close
sequence identity within the RBD core while variation is highest within the functional ACE2 binding site
(Figures 1 and S1), possibly due to a co-evolutionary 'arms race' with polymorphic ACE2 sequences in
ecologically diverse bat species (*28*). Affinity was measured by biolayer interferometry (BLI), with
$sACE2_2$ (a.a. S19-G732) fused at the C-terminus with the Fc moiety of human IgG1 immobilized to the
sensor surface and monomeric 8his-tagged RBD (Figure S2) used as the soluble analyte.   This
arrangement excludes avidity effects, which otherwise cause artificially tight (picomolar) apparent
affinities whenever dimeric $sACE2_2$ in solution is bound to immobilized RBD decorating an interaction
surface. Wild type $sACE2_2$ bound all the RBDs with affinities ranging from 16 nM for SARS-CoV-2 to
91 nM for LYRa11, with median affinity 60 nM (Table 1). The measured affinities for the RBDs of
SARS-CoV-1 and SARS-CoV-2 are comparable to published data (*4, 25, 29-31*). Engineered
$sACE2_2$.v2.4 displayed large increases in affinity for all the RBDs, with K_{DS} ranging from 0.4 nM for
SARS-CoV-2 to 3.5 nM for isolate Rs4231, with median affinity less than 2 nM (Table 1).   The
approximate 35-fold affinity increase of the engineered decoy applies universally to coronaviruses in the
test panel and the molecular basis for affinity enhancement must therefore be grounded in common
attributes of RBD/ACE2 recognition.

Figure 1. SARS-associated coronaviruses have high sequence diversity at the ACE2-binding site.
The RBD of SARS-CoV-2 (PDB 6M17) is colored by diversity between 7 SARS-associated CoV strains
(blue, conserved; red, variable).

Table 1. BLI kinetics for immobilized $sACE2_2$-IgG1 binding to coronavirus RBDs								
CoV strain [a]	Wild type $sACE2_2$-IgG1 [b]				$sACE2_2$.v2.4-IgG1			
	k_{on} ($M^{-1}s^{-1}$)	k_{off} (s^{-1})	K_D (nM)	χ^{2} [c]	k_{on} ($M^{-1}s^{-1}$)	k_{off} (s^{-1})	K_D (nM)	χ^2
LYRa11	8.7×10^5	7.9×10^{-2}	91	0.12	1.4×10^6	2.5×10^{-3}	1.8	0.10
Rs7327	6.4×10^5	4.0×10^{-2}	63	0.25	9.8×10^5	1.8×10^{-3}	1.9	0.11
Rs4231	3.2×10^5	2.2×10^{-2}	69	0.04	4.5×10^5	1.6×10^{-3}	3.5	0.10
Rs4084	2.9×10^5	2.5×10^{-2}	85	0.24	4.8×10^5	1.5×10^{-3}	3.1	0.10
RsSHC014	8.8×10^5	2.6×10^{-2}	29	0.20	1.6×10^6	2.0×10^{-3}	1.3	0.29

SARS-1	6.6×10^3	1.2×10^{-4}	58	0.03	3.0×10^3	5.6×10^{-6}	2.1	0.03
SARS-2	1.4×10^6	8.1×10^{-3}	16	0.25	6.6×10^5	2.8×10^{-4}	0.4	0.09

[a] Purified RBDs at 5 to 7 concentrations were used as the soluble analytes.
[b] IgG1-Fc fused $sACE2_2$ was immobilized to anti-human IgG Fc capture biosensors.
[c] χ^2 values represent the goodness of curve fitting. Acceptable values were considered to be less than 3.

A deep mutational scan of the RBD in the context of full-length S reveals residues in the ACE2 binding site are mutationally tolerant

To explore potential sequence diversity in S of SARS-CoV-2 that may act as a 'reservoir' for drug
resistance, the mutational tolerance of the RBD was evaluated by deep mutagenesis (*32*). Saturation
mutagenesis was focused to the RBD (a.a. C336-L517) of full-length S tagged at the extracellular N-
terminus with a c-myc epitope for detection of surface expression. The spike library, encompassing 3,640
single amino acid substitutions, was transfected in human Expi293F cells under conditions where cells
typically acquire no more than a single sequence variant (*33*, *34*). The culture was incubated with wild
type, 8his-tagged, dimeric $sACE2_2$ at a sub-saturating concentration (2.5 nM). Bound $sACE2_2$-8h and
surface-expressed S were stained with fluorescent antibodies for flow cytometry analysis (Figure 2A).
Compared to cells expressing wild type S, the library was poorly expressed, indicating many mutations
are deleterious for folding and expression. A cell population was clearly discernable expressing S
variants that bind ACE2 with decreased affinity (Figure 2B). After gating for c-myc-positive cells
expressing S, cells with high and low levels of bound $sACE2_2$ were collected by fluorescence-activated
cell sorting (FACS), called the ACE2-High and ACE2-Low populations, respectively (Figure 2C). Both
the expression and $sACE2_2$ binding signals decreased over minutes to hours during sorting, possibly due
to shedding of the S1 subunit. Cells were therefore collected and pooled from three separate FACS
experiments for a combined 8 hours sort time.

Figure 2. FACS selection for variants of S with high or low binding signal to ACE2. (A) Flow
cytometry analysis of Expi293F cells expressing full-length S of SARS-CoV-2 with an N-terminal c-myc
tag. Staining for the myc-epitope is on the x-axis while the detection of bound $sACE2_2$-8h (2.5 nM) is on
the y-axis. S plasmid was diluted 1500-fold by weight with carrier DNA so that cells typically express no
more than one coding variant; under these conditions most cells are negative. **(B)** Flow cytometry of cells
transfected with the RBD single site-saturation mutagenesis (SSM) library shows cells expressing S
variants with reduced $sACE2_2$-8h binding. **(C)** Gating strategy for FACS. S-expressing cells positive for
the c-myc epitope were gated (blue) and the highest ("ACE2-High") and lowest ("ACE2-Low") 20% of
cells with bound $sACE2_2$-8h relative to myc-S expression were collected.

Transcripts in the sorted cells were Illumina sequenced and compared to the naive plasmid library to
determine an enrichment ratio for each amino acid substitution (*35*). Mutations in S that express and bind
ACE2 tightly are selectively enriched in the ACE2-High sort (Figure S3); mutations that express but have
reduced ACE2 binding are selectively enriched in the ACE2-Low sort; and mutations that are poorly

expressed are depleted from both sorted populations. Positional conservation scores were calculated by
averaging the $\log_2$ enrichment ratios for each of the possible amino acids at a residue position. By adding
conservation scores for both the ACE2-High and ACE2-Low sorts we derive a score for surface
expression, which shows that the hydrophobic RBD core is tightly conserved for folding and trafficking
of the viral spike (Figure 3A). By comparison, residues on the exposed RBD surface are mutationally
permissive for S surface expression. This matches the mutational tolerance of proteins generally.

Figure 3. Deep mutagenesis reveals that the ACE2-binding site of SARS-CoV-2 tolerates many
mutations. (A) Positional scores for surface expression are mapped to the structure of the SARS-CoV-2
RBD (PDB 6M17, oriented as in Figure 1). Blue residues in the protein core are highly conserved in the
FACS selection for surface S expression (judged by depletion of mutations from the ACE2-High and
ACE2-Low gates), while surface residues in red tolerate mutations. **(B)** Correlation plot of expression
scores from mutant selection in human cells of full-length S (x-axis) versus the conservation scores
(mean of the $\log_2$ enrichment ratios at a residue position) from mutant selection in the isolated RBD by
yeast display (y-axis). Notable outliers are indicated. **(C)** Conservation scores from the ACE2-High gated
cell population are mapped to the RBD structure, with residues colored from low (blue) to high (red)
mutational tolerance. **(D)** Correlation plot of RBD conservation scores for high ACE2 binding from deep
mutagenesis of S in human cells (x-axis) versus deep mutagenesis of the RBD on the yeast surface
(mean of $\Delta K_{D\ app}$; y-axis).

For tight ACE2 binding (i.e. S variants in the ACE2-High population), conservation increases for
RBD residues at the ACE2 interface, yet mutational tolerance remains high (Figure 3C). The sequence
diversity observed among natural betacoronaviruses, which display high diversity at the ACE2 binding
site, is therefore replicated in the deep mutational scan, which predicts the SARS-CoV-2 spike tolerates
substantial genetic diversity at the receptor-binding site for function. From this accessible sequence
diversity SARS-CoV-2 might feasibly mutate to acquire resistance to monoclonal antibodies or
engineered decoy receptors targeting the ACE2-binding site.

Comparison to a deep mutational scan of the isolated RBD by yeast surface display

Two deep mutational scans have been reported for the isolated RBD displayed on the surface of
yeast (*36*, *37*). We compare our data, from a selection of full-length S expressed in human cells, to the
publicly accessible Starr et al data set (*36*). Important residues within the RBD for surface expression of
full-length spike in human cells are closely correlated with data from yeast surface display of the isolated
RBD (Figure 3B), with the exception of a notable region. The surface of the RBD opposing the ACE2-
binding site (e.g. V362, Y365 and C391) is free to mutate for yeast surface display, but its sequence is
constrained in our experiments; this region of the RBD is buried by connecting structural elements to the
global fold of an S subunit in the closed-down conformation (this is the dominant conformation for S
subunits and is inaccessible to receptor binding) (*2*, *4*, *38*, *39*). We used targeted mutagenesis to
individually test alanine substitutions to all the cysteines in the RBD (Figure S4). We found all cysteine-
to-alanine mutations severely diminish S surface expression in Expi293F cells, including C391A and
C525A on the RBD 'backside' that were neutral in the yeast display scan (*36*). These differences
demonstrate that there are tighter sequence constraints on the RBD in the context of a full spike expressed
at a human cell membrane, yet overall we consider the two data sets to closely agree.

For binding to dimeric sACE2$_2$, we note that interface residues were more tightly conserved in the
Starr et al data set (Figure 3D), possibly a consequence of three differences between the deep mutagenesis
experiments. First, our selections for ACE2 binding of S variants at the plasma membrane appears to
primarily reflect mutational effects on surface expression, which is almost certainly more stringent in
human cells. Yeast permit many poorly folded proteins to leak to the cell surface (*40*). Second, the yeast
selections were conducted at multiple sACE2 concentrations from which apparent K_D changes were
computed (*36*); the Starr et al data in this regard is very comprehensive. Due to the long sort times
required for our human cell libraries where only a small fraction of cells express spike, we sorted at a
single sACE2$_2$ concentration that cannot accurately capture a range of different binding affinities
quantitatively. Third, dimeric sACE2$_2$ may geometrically complement trimeric S densely packed on a
human cell membrane, such that avidity masks the effects of affinity-reducing mutations. Nonetheless,
there is overall agreement that ACE2 binding often persists following mutations to the RBD surface, and
our data simply suggests mutational tolerance may be even greater than that already observed by Starr et
al.

A screen for S variants that preferentially bind wild type ACE2 over the engineered decoy

Having shown that the ACE2-binding site of SARS-CoV-2 protein S tolerates many mutations, we
asked whether mutations might therefore be found that confer resistance to the engineered decoy
sACE2$_2$.v2.4. Resistance mutations are anticipated to lose affinity for sACE2$_2$.v2.4 while maintaining
binding to the wild type receptor, and are most likely to reside in the RBD where physical contacts are
made. Similar reasoning formed the foundation of a deep mutagenesis-based selection of the isolated
RBD by yeast surface display to find escape mutations to monoclonal antibodies, and the results were
predictive of escape mutations in pseudovirus growth selections (*19*).

To address whether escape mutations from the engineered decoy might be found in the RBD, we
repurposed the S protein library for a specificity selection. Cells expressing the library, encoding all
possible substitutions in the RBD, were co-incubated with wild type sACE2 fused to the Fc region of
IgG1 and 8his-tagged sACE2.v2.4 at concentrations where both proteins bind competitively (*25*). It was
immediately apparent from flow cytometry of the Expi293F culture expressing the S library that there
were cells expressing S variants shifted towards preferential binding to sACE2.v2.4, but no significant
population with preferential binding to the wild type receptor (Figures 4A and 4B). Cells expressing S
variants that might preferentially bind sACE2(WT)-IgG1 or sACE2.v2.4 were gated and collected by
FACS (Figure 4C), followed by deep sequencing of S transcripts to determine enrichment ratios. There
was close agreement between two independent replicate experiments (Figures 4D-4G). Most RBD
mutations were depleted following sorting, consistent with deleterious effects on S folding and
expression.

Figure 4. A competition-based selection to identify RBD mutations within S of SARS-CoV-2 that
preferentially bind wild type or engineered ACE2 receptors. (A) Expi293F cells were transfected with
wild type myc-S and incubated with competing sACE2(WT)-IgG1 (25 nM) and sACE2.v2.4-8h (20 nM).
Bound protein was detected by flow cytometry after immuno-staining for the respective epitope tags. **(B)**
As in A, except cells were transfected with the RBD SSM library. A population of cells expressing S
variants with increased specificity towards sACE2.v2.4 is apparent (cells shifted to the upper-left of the
main population). **(C)** Gates used for FACS of cells expressing the RBD SSM library. After excluding
cells without bound protein, the top 20% of cells for bound sACE2.v2.4-8h (magenta gate) and for bound
sACE2(WT)-IgG1 (green gate) were collected. **(D-E)** Agreement between $\log_2$ enrichment ratios from two
independent FACS selections for cells expressing S variants with increased specificity for (D)
sACE2(WT) or (E) sACE2.v2.4. R^2 values are calculated for nonsynonymous mutations (black).
Nonsense mutations are red. **(F-G)** Conservation scores are calculated from the mean of the $\log_2$
enrichment ratios for all nonsynonymous substitutions at a given residue position. Correlation plots show
agreement between conservation scores for two independent selections for cells within the (D)
sACE2(WT) or (E) sACE2.v2.4 specific gates.

Soluble ACE2.v2.4 has three mutations from wild type ACE2: T27Y buried within the RBD
interface, and L79T and N330Y at the interface periphery (Figure 5A). A substantial number of mutations
in the RBD of S were selectively enriched for preferential binding to sACE2.v2.4 (Figure 5B, upper-left
quadrant). While sACE2.v2.4-specificity mutations could be found immediately adjacent to the sites of
engineered mutations in ACE2 (in particular mutations to S-F486 adjacent to ACE2-L79 and S-T500

adjacent to ACE2-N330), major hot spots for sACE2₂.v2.4-specificity mutations were also mapped to
RBD loop 498-506, contacting the region where the ACE2-α1 helix packs against a β-hairpin motif
(Figure 5A). By comparison, there were no hot spots in the RBD for sACE2₂(WT)-specificity mutations.
Indeed, only a small number of mutations were selectively enriched for preferential binding to wild type
receptor (Figure 5B), and the abundance of these putative wild type-specific mutations barely rose above
the expected level of noise in the deep mutagenesis data. In this competition assay, S binding to wild type
sACE2₂ is therefore more sensitive to RBD mutations than S binding to engineered sACE2₂.v2.4.

Figure 5. Mutations within the RBD that confer specificity towards wild type ACE2 are rare. (A) The
SARS-CoV-2 RBD is colored by specificity score (the difference between the conservation scores for cells
collected in the sACE2₂(WT) and sACE2₂.v2.4 specific gates). Residues that are hot spots for mutations
with increased specificity towards sACE2₂(WT) are blue or towards sACE2₂.v2.4 are red. The contacting
surface of ACE2 is shown as a green ribbon, with sites of mutations in sACE2₂.v2.4 labeled and shown
as green spheres. **(B)** Log₂ enrichment ratios for mutations in S expressed by cell populations collected in
the sACE2₂(WT) (x-axis) and sACE2₂.v2.4 (y-axis) specific gates. Data are the mean from two
independent sorting experiments. S mutants in blue were predicted to have increased specificity for
sACE2₂(WT) and were tested by targeted mutagenesis in Figure S5. S mutants in purple were predicted
to have increased specificity for sACE2₂.v2.4 and were tested by targeted mutagenesis in Figure S6.
Other nonsynonymous mutations are black. Nonsense mutations are red. **(C)** Wild type myc-S (blue) and
two variants, N501W (orange) and N501Y (green), were expressed in Expi293F cells and tested by flow
cytometry for binding to sACE2₂(WT)-8h (dashed lines) or sACE2₂.v2.4-8h (solid lines). Minor increases
in specific binding for wild type sACE2₂ are observed.

To determine whether the potential wild type ACE2-specific mutations found by deep mutagenesis
are real as opposed to false predictions due to data noise, we tested 24 mutants of S selectively enriched in
the wild type-specific gate by targeted mutagenesis (blue data points in Figure 5B). Only minor shifts
towards binding wild type sACE2₂ were observed (Figure S5). Two S mutants were investigated further
in sACE2₂ titration experiments, N501W and N501Y, which both retained high receptor binding and
displayed small shifts towards wild type sACE2₂ in the competition experiment. N501 of S is located in

the 498-506 loop and its substitution to large aromatic side chains might alter the loop conformation to
cause steric strain with nearby ACE2 mutation N330Y in sACE2$_2$.v2.4. After titrating the concentrations
of 8his-tagged sACE2$_2$(WT) and sACE2$_2$.v2.4 and measuring bound protein to S-expressing cells by flow
cytometry, it was found S-N501W and S-N501Y do show enhanced specificity for wild type sACE2$_2$, but
the effect is small and sACE2$_2$.v2.4 remains the stronger binder (Figure 5C); these mutations therefore
will not confer resistance in the virus to the engineered decoy. By comparison, multiple independent
escape mutations are readily found in S of SARS-CoV-2 that diminish the efficacy of monoclonal
antibodies by many orders of magnitude (*18, 19*).

Finally, 8 representative mutations to S predicted from the deep mutational scan to increase
specificity towards sACE2$_2$.v2.4 (purple data points in Figure 5B) were cloned and 7 were found to have
large shifts towards preferential sACE2$_2$.v2.4 binding in the competition assay (Figure S6). These S
mutations were Y449K/Q/S, L455G/R/Y and G504K. None of the mutated sites is in direct contact with
an engineered residue on sACE2$_2$.v2.4 and the molecular bases for specificity changes are therefore
ambiguous, but we speculate may involve local conformational perturbations. Validation by targeted
mutagenesis therefore confirms that the selection can successfully find mutations in S with altered
specificity. The inability to find mutations in the RBD that impart high specificity for the wild type
receptor means such mutations are rare or may not even exist, at least within the receptor-binding domain
where direct physical contacts with receptors occur. We cannot exclude mutations elsewhere having
long-range conformational effects. Engineered, soluble decoy receptors therefore live up to their promise
as broad therapeutic candidates against which a virus cannot easily escape.

DISCUSSION

The allure of soluble decoy receptors is that the virus cannot easily mutate to escape neutralization.
Mutations that reduce affinity of the soluble decoy will likely also decrease affinity for the wild type
receptor on host cells, thereby coming at the cost of diminished infectivity and virulence. However, this
hypothesis has not been rigorously tested, and since engineered decoy receptors differ from their wild
type counterparts, even if by just a small number of mutations, it is possible a virus may evolve to
discriminate between the two. Here, we show that an engineered decoy receptor for SARS-CoV-2
broadly binds with low nanomolar K_D to the spikes of SARS-associated betacoronaviruses that use ACE2
for entry, despite high sequence diversity within the ACE2-binding site. Mutations in S of SARS-CoV-2
that confer high specificity for wild type ACE2 were not found in a comprehensive screen of all
substitutions within the RBD. The engineered decoy receptor is therefore broad against zoonotic ACE2-
utilizing coronaviruses that may spill over from animal reservoirs in the future and against variants of
SARS-CoV-2 that may arise as the current COVID-19 pandemic rages on. We argue it is unlikely that
decoy receptors will need to be combined in cocktail formulations, as is required for monoclonal
antibodies or designed miniprotein binders to prevent the rapid emergence of resistance (*18, 41*),
facilitating manufacture and distribution. Our findings give insight into how a potential therapeutic can
achieve breath with a low chance of virus resistance for a family of highly infectious and deadly viruses.

Soluble decoy receptors have proven effective in the clinic, especially for modulating immune
responses. Etanercept (trade name Enbrel; soluble TNF receptor), aflibercept (Eylea; a soluble chimera of
VEGF receptors 1 and 2) and abatacept (Orencia; soluble CTLA-4) are just three examples of soluble
receptors that have profoundly impacted the treatment of human disease (*42*), yet no soluble receptors for
a viral pathogen are approved drugs. There are two main reasons for this. First, the affinity of entry
receptors for viral glycoproteins is often moderate to low, which reduces neutralization potency compared
to affinity-matured monoclonal antibodies. For SARS-CoV-2, we and others have solved this problem by
engineering ACE2 to have picomolar affinity for viral S (*25-27*). Second, virus entry receptors have
endogenous functions for normal physiology and their soluble counterparts may impact this normal
physiology to exert unacceptable toxicity. For example, the entry receptor for human cytomegalovirus is

a growth factor receptor, and growth factor interactions had to be knocked out to make a virus-specific decoy suitable for in vivo administration (*43*). However, ACE2 in this regard is different and its endogenous activity – the catalytic conversion of vasoconstrictive and inflammatory peptides of the renin-angiotensin system – may be of direct benefit for addressing COVID-19 symptoms. During infection, ACE2 activity is dysregulated and the renin-angiotensin system becomes imbalanced, possibly driving aspects of acute-respiratory distress syndrome (ARDS) (*44-46*). Administration of recombinant sACE2 converts angiotensin (Ang) I and II to the protective peptides Ang-(1-9) and Ang-(1-7), respectively, with potential benefits for the pulmonary and cardiovascular systems that include decreased lung elastance, increased blood oxygenation, reduced hypertension and diminished inflammation (*44*, *45*, *47-50*). Soluble, wild type $ACE2_2$ has been developed as a drug for ARDS with an acceptable safety profile in humans (*51*, *52*) and is currently under evaluation in a clinical trial by Apeiron. Promisingly, a first clinical case report found that administration of wild type $sACE2_2$ was correlated with reduced viral burden and improved clinical outcomes in a 45 year old woman with severe COVID-19 (DOI:https://doi.org/10.1016/S2213-2600(20)30418-5). Engineered, high affinity $sACE2_2$ decoys, most likely as fusions with immunoglobulin Fc for increased serum stability (*23*, *50*, *53*), represent next generation therapeutics with dual mechanisms of action: (i) potent virus neutralization due to high affinity blockade of the viral spike and (ii) proteolytic turnover of peptide hormones for direct relief of COVID-19 symptoms.

MATERIALS AND METHODS

Plasmids. Residue numbers for constructs begin from the start methionine as amino acid (a.a.) 1. The cloning of human codon-optimized, mature S from SARS-CoV-2 (GenBank Acc. No. YP_009724390.1; a.a. V16-T1273) into the NheI-XhoI sites of pCEP4 (Invitrogen) with an N-terminal, extracellular c-myc tag is described elsewhere (*25*). Soluble ACE2 (a.a. 1-732 encoding a dimer; wild type or engineered variant $sACE2_2$.v2.4) fused to an 8his purification tag or to human IgG1-Fc (a.a. D221-K447; nG1m1 isoallotype; GenBank KY432415.1) and cloned in to the NheI-XhoI sites of pcDNA3.1(+) (Invitrogen) is also previously described (*25*). The RBDs of SARS-CoV-1 (Urbani isolate; GenBank AAP13441.1; a.a. T320-D518), SARS-CoV-2 (YP_009724390; a.a. T333-K529), LYRa11 (AHX37558.1; a.a. T324-D522), Rs7327 (ATO98218.1; a.a. T321-D519), Rs4231 (ATO98157.1; a.a. T320-D518), Rs4084 (ATO98132.1; a.a. T321-D519) and RsSHC014 (AGZ48806.1; a.a. T321-D519) were cloned with N-terminal influenza HA leader peptides (sequence MKTIIALSYIFCLVFA) and C-terminal 8-his tags (sequence GSGHHHHHHHH) into the NheI-XhoI sites of pcDNA3.1(+). These plasmids are deposited with Addgene under accession numbers 145145 and 161821-161826. Mutations were made by overlap extension PCR and verified by Sanger sequencing.

Tissue Culture. Expi293F cells (ThermoFisher) were grown in Expi293 Expression Medium (ThermoFisher) at 125 rpm, 8 % CO2, 37 °C.

Recombinant Protein Production. Plasmids (500 ng DNA per ml culture) and polyethylenimine (MW 25,000; Polysciences; 3 μg per ml culture) were mixed with OptiMEM (Gibco; 100 μl per ml culture), incubated 20 minutes at room temperature and added to Expi293F cells at a density of 2×10^6 / ml. Transfection Enhancers (ThermoFisher) were added 18-23 h post-transfection. Culture supernatant was harvested 4-6 days later by two centrifugation steps (800 × g for 10 minutes to remove cells and 20,000 × g for 20 minutes to remove debris). IgG1 Fc fused and 8his-tagged proteins were subsequently purified as previously described (*25*) using KANEKA KanCapA 3G Affinity (Pall) and HisPur Ni-NTA (Thermo Scientific) resins, respectively. Eluted proteins from affinity chromatography were then separated on a Superdex 200 Increase 10/300 GL column (GE Healthcare Life Sciences) equilibrated with Dulbecco's phosphate-buffered saline (PBS). Proteins from peak fractions were concentrated using centrifugal ultrafiltration devices (Millipore) to final concentrations of ~1 mg/ml (RBD-8h proteins), ~10 mg/ml ($sACE2_2$-8h proteins) and ~50 mg/ml ($sACE2_2$-IgG1 proteins). Concentrations were determined by

absorbance at 280 nm using calculated extinction coefficients. Reported concentrations for sACE2$_2$ are
based on monomeric subunits. Aliquots were snap frozen in liquid N$_2$ and stored at -80 °C.

Biolayer Interferometry. BLI kinetics were collected on an Octet RED96a and analyzed with a 1:1
binding model (global fit) using instrument software (Molecular Devices). IgG1 Fc-fused sACE2$_2$ (wild
type or engineered variant sACE2$_2$.v2.4) were immobilized at 100 nM for 10 minutes to anti-human IgG
Fc biosensors (Molecular Devices). The assay buffer was 10 mM HEPES pH 7.6, 150 mM NaCl, 3 mM
EDTA, 0.05% polysorbate 20, 0.5% non-fat dry milk (Bio-Rad). Loaded sensors were equilibrated for 30
s in buffer, then dipped in RBD-8h solutions for 60 s to measure association and transferred back to
buffer to measure dissociation over 300 s.

Library Construction, FACS and Illumina Sequencing Analysis. Using plasmid pCEP4-myc-S
encoding tagged, full length S of SARS-CoV-2, saturation mutagenesis was focused to residues C336-
L517 forming the RBD. Degenerate NNK codons were introduced at all RBD positions using overlap
extension PCR as previously described (*54*). Transient transfection conditions were used that typically
provide no more than a single coding variant per cell (*33, 34*). Expi293F cells at 2 × 10^6 / ml were
transfected with a mixture of 1 ng coding plasmid (i.e. library DNA) with 1.5 μg pCEP4-ΔCMV carrier
plasmid (described in (*34*)). The medium was replaced 2 h post-transfection and cells were collected 24 h
post-transfection for FACS. Cells were washed with ice-cold PBS supplemented with 0.2 % bovine serum
albumin (PBS-BSA).

For investigations of binding to wild type sACE2$_2$, the cells expressing the S library were resuspended in
2.5 nM sACE2$_2$(WT)-8h and incubated 30 minutes on ice. Cells were washed twice with PBS-BSA and
then co-stained for 20 minutes with anti-myc Alexa 647 (clone 9B11, 1/250 dilution; Cell Signaling
Technology) and anti-HIS-FITC (chicken polyclonal, 1/100 dilution; Immunology Consultants
Laboratory). Cells were again washed twice before sorting on a BD FACS Aria II at the Roy J. Carver
Biotechnology Center. Dead cells, doublets and debris were excluded by first gating on the main
population by forward/side scattering and then excluding DAPI positive cells. From the myc-S-positive
(Alexa 647) population, the 20 % of cells with the highest and 20% of cells with the lowest anti-HIS-
FITC fluorescence for bound sACE2$_2$(WT)-8h were collected (Figure 2C). Collection tubes were coated
overnight with fetal bovine serum prior to sorting and contained Expi293 Expression Medium. Collected
cell pellets were frozen at -80°C and were pooled across separate sort experiments prior to extraction of
total RNA.

The competition selection was performed similarly, with the exception that cells expressing the S library
were incubated for 30 minutes in a mixture of 20 nM sACE2$_2$.v2.4-8h and 25 nM sACE2$_2$(WT)-IgG1.
After washing twice, bound proteins were stained for 30 minutes with anti-human IgG-APC (clone
HP6017, 1/250 dilution; BioLegend) and anti-HIS-FITC (chicken polyclonal, 1/100 dilution;
Immunology Consultants Laboratory). Cells were washed twice and sorted. After gating for the main
population of viable cells as described above, the 20 % of cells with the highest FITC-relative-to-APC
and highest APC-relative-to-FITC signals were collected (Figure 4C).

Total RNA was extracted from the collected cells using a GeneJET RNA purification kit (Thermo
Scientific). First strand cDNA was synthesized with Accuscript (Agilent) primed with a gene-specific
oligonucleotide. The region of S scanned by saturation mutagenesis was PCR amplified as 3 overlapping
fragments that together span the full RBD sequence. Following a second round of PCR, primers added
adapters for annealing to the Illumina flow cell and sequencing primers, together with barcodes for
experiment identification. The PCR products were sequenced on an Illumina NovaSeq 6000 using a
2×250 nt paired end protocol. Data were analyzed using Enrich (*35*), where the frequencies of S variants
in the transcripts of the sorted populations were compared to their frequencies in the naive plasmid
library. Log$_2$ enrichment ratios for all the individual mutations were calculated and normalized by

subtracting the $\log_2$ enrichment ratio for the wild type sequence across the same PCR-amplified fragment.
Conservation scores at residue positions were calculated by averaging the $\log_2$ enrichment ratios for all
non-synonymous mutations at the residue.

Flow Cytometry Analysis of SARS-CoV-2 S Mutants. Expi293F cells at 2.0×10^6 cells/ml were
transfected with plasmid DNA (300 ng per ml of culture for measuring myc-S surface expression and
sACE2$_2$ competition binding, 500 ng per ml for titration experiments) encoding myc-S variants using
Expifectamine (ThermoFisher) according to the manufacturer's directions. At 24 h post-transfection, cells
were washed with PBS-BSA. To detect surface expressed myc-S, cells were incubated with anti-myc
Alexa 647 (clone 9B11, 1/250 dilution; Cell Signaling Technology) on a rocker at 4°C for 30 minutes. To
measure competitive binding of wild type and engineered receptors, cells were instead incubated with 25
nM sACE2$_2$(WT)-IgG1 and 20 nM sACE2$_2$.v2.4-8h for 30 minutes at 4°C, washed twice, and stained
with anti-human IgG-APC (clone HP6017, 1/250 dilution; BioLegend) and anti-HIS-FITC (chicken
polyclonal, 1/100 dilution; Immunology Consultants Laboratory) secondary antibodies for 20 minutes at
4°C. Finally, in titration experiments, transfected cells were incubated for 30 minutes at 4°C with 1/3
serial dilutions of sACE22(WT)-8h or sACE22.v2.4-8h, followed by two washes and a 30 minute
incubation with anti-myc Alexa 647 (clone 9B11, 1/250 dilution) and anti-HIS-FITC (chicken polyclonal,
1/100 dilution). For all experiments, cells were washed twice prior to analysis on an Accuri C6 Flow
Cytometer (BD Biosciences) and data were processed with FCS Express (De Novo Software).
Quantification of myc-S surface expression is detailed in Figure S4.

Reagent and Data Availability. Plasmids for RBD protein expression are deposited with Addgene
(numbers 145145 and 161821-161826). Illumina sequencing data are deposited in NCBI's Gene
Expression Omnibus (GEO) under series accession number GSE159372.

ACKNOWLEDGEMENTS

429 The Roy J. Carver Biotechnology Center at the University of Illinois assisted with flow cytometry and
430 Illumina sequencing. E.P. designed the study and completed deep mutagenesis. K.K.C. purified proteins,
431 tested BLI kinetics. K.K.C., K.K.N. and T.J.C.T. prepared samples for flow cytometry. This work was in
432 part supported by NIH award R01AI129719 to E.P. The University of Illinois has filed a provisional
433 patent for engineered decoy receptors and E.P. and K.K.C. are co-founders of Orthogonal Biologics, Inc.

REFERENCES

1.   P. Zhou *et al.*, A pneumonia outbreak associated with a new coronavirus of probable bat origin.
Nature. **579**, 270–273 (2020).

2.   A. C. Walls *et al.*, Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein.
Cell (2020), doi:10.1016/j.cell.2020.02.058.

3.   Y. Wan, J. Shang, R. Graham, R. S. Baric, F. Li, Receptor recognition by novel coronavirus from
Wuhan: An analysis based on decade-long structural studies of SARS. *J. Virol.* (2020),
doi:10.1128/JVI.00127-20.

4.   D. Wrapp *et al.*, Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.
Science, eabb2507 (2020).

5.   M. Hoffmann *et al.*, SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked
by a Clinically Proven Protease Inhibitor. *Cell* (2020), doi:10.1016/j.cell.2020.02.052.

6.   W. Li *et al.*, Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus.
447   *Nature*. **426**, 450–454 (2003).

448   7.   M. Letko, A. Marzi, V. Munster, Functional assessment of cell entry and receptor usage for
449   SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol*. **11**, 1860 (2020).

450   8.   L. Samavati, B. D. Uhal, ACE2, Much More Than Just a Receptor for SARS-COV-2. *Front. Cell.*
451   *Infect. Microbiol.* **10**, 752 (2020).

452   9.   F. Jiang *et al.*, Angiotensin-converting enzyme 2 and angiotensin 1–7: novel therapeutic targets.
453   *Nature Reviews Cardiology*. **11**, 413–426 (2014).

454   10.   L. Zhang *et al.*, The D614G mutation in the SARS-CoV-2 spike protein reduces S1 shedding and
455   increases infectivity. *bioRxiv*, 2020.06.12.148726 (2020).

456   11.   B. Korber *et al.*, Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases
457   Infectivity of the COVID-19 Virus. *Cell*. **182**, 812–827.e19 (2020).

458   12.   V. Borges *et al.*, On the track of the D839Y mutation in the SARS-CoV-2 Spike fusion peptide:
459   emergence and geotemporal spread of a highly prevalent variant in Portugal. *medRxiv*,
460   2020.08.10.20171884 (2020).

461   13.   K. Pyrc *et al.*, Mosaic structure of human coronavirus NL63, one thousand years of evolution. *J.*
462   *Mol. Biol.* **364**, 964–973 (2006).

463   14.   K. Wu, W. Li, G. Peng, F. Li, Crystal structure of NL63 respiratory coronavirus receptor-binding
464   domain complexed with its human receptor. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 19970–19974
465   (2009).

466   15.   S. Su *et al.*, Epidemiology, Genetic Recombination, and Pathogenesis of Coronaviruses. *Trends*
467   *Microbiol.* **24**, 490–502 (2016).

468   16.   M. F. Boni *et al.*, Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for
469   the COVID-19 pandemic. *Nat Microbiol* (2020).

470   17.   J. S. M. Sabir *et al.*, Co-circulation of three camel coronavirus species and recombination of
471   MERS-CoVs in Saudi Arabia. *Science*. **351**, 81–84 (2016).

472   18.   A. Baum *et al.*, Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape
473   seen with individual antibodies. *Science*, eabd0831 (2020).

474   19.   A. J. Greaney *et al.*, Complete mapping of mutations to the SARS-CoV-2 spike receptor-binding
475   domain that escape antibody recognition. *bioRxiv*, 2020.09.10.292078 (2020).

476   20.   M. A. Tortorici *et al.*, Ultrapotent human antibodies protect against SARS-CoV-2 challenge via
477   multiple mechanisms. *Science*, eabe3354 (2020).

478   21.   D. Pinto *et al.*, Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
479   antibody. *Nature* (2020), doi:10.1038/s41586-020-2349-y.

480   22.   H. Hofmann *et al.*, Susceptibility to SARS coronavirus S protein-driven infection correlates with
481   expression of angiotensin converting enzyme 2 and infection can be blocked by soluble receptor.

Biochem. Biophys. Res. Commun. **319**, 1216–1221 (2004).

23.   C. Lei *et al.*, Neutralization of SARS-CoV-2 spike pseudotyped virus by recombinant ACE2-Ig.
Nat Commun. **11**, 2070 (2020).

24.   V. Monteil *et al.*, Inhibition of SARS-CoV-2 Infections in Engineered Human Tissues Using
Clinical-Grade Soluble Human ACE2. *Cell* (2020), doi:10.1016/j.cell.2020.04.004.

25.   K. K. Chan *et al.*, Engineering human ACE2 to optimize binding to the spike protein of SARS
coronavirus 2. *Science*. **4**, eabc0870 (2020).

26.   A. Glasgow *et al.*, Engineered ACE2 receptor traps potently neutralize SARS-CoV-2. *bioRxiv*,
2020.07.31.231746 (2020).

27.   Y. Higuchi *et al.*, High affinity modified ACE2 receptors prevent SARS-CoV-2 infection. *bioRxiv*,
2020.09.16.299891 (2020).

28.   H. K. Frank, D. Enard, S. D. Boyd, Exceptional diversity and selection pressure on SARS-CoV
and SARS-CoV-2 host receptor in bats compared to other mammals. *bioRxiv*. **5**, 562 (2020).

29.   J. Shang *et al.*, Structural basis of receptor recognition by SARS-CoV-2. *Nature*. **382**, 1199
(2020).

30.   R. N. Kirchdoerfer *et al.*, Stabilized coronavirus spikes are resistant to conformational changes
induced by receptor recognition or proteolysis. *Sci Rep*. **8**, 15701 (2018).

31.   W. Li *et al.*, Receptor and viral determinants of SARS-coronavirus adaptation to human ACE2.
EMBO J. **24**, 1634–1643 (2005).

32.   D. M. Fowler, S. Fields, Deep mutational scanning: a new style of protein science. *Nat. Methods*.
11, 801–807 (2014).

33.   J. D. Heredia *et al.*, Mapping Interaction Sites on Human Chemokine Receptors by Deep
Mutational Scanning. *J. Immunol.* **200**, ji1800343–3839 (2018).

34.   J. Park *et al.*, Structural architecture of a dimeric class C GPCR based on co-trafficking of sweet
taste receptor subunits. *Journal of Biological Chemistry*. **294**, 4759–4774 (2019).

35.   D. M. Fowler, C. L. Araya, W. Gerard, S. Fields, Enrich: software for analysis of protein function
by enrichment and depletion of variants. *Bioinformatics*. **27**, 3430–3431 (2011).

36.   T. N. Starr *et al.*, Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain Reveals
Constraints on Folding and ACE2 Binding. *Cell*. **182**, 1295–1310.e20 (2020).

37.   T. W. Linsky *et al.*, De novo design of ACE2 protein decoys to neutralize SARS-CoV-2. *bioRxiv*,
2020.08.03.231340 (2020).

38.   Y. Cai *et al.*, Distinct conformational states of SARS-CoV-2 spike protein. *Science*. **369**, 1586
(2020).

39.   H. Yao *et al.*, Molecular architecture of the SARS-CoV-2 virus. *Cell* (2020).

40.   G. J. Rocklin *et al.*, Global analysis of protein folding using massively parallel design, synthesis,
and testing. *Science*. **357**, 168–175 (2017).

41.   L. Cao *et al.*, De novo design of picomolar SARS-CoV-2 miniprotein inhibitors. *Science*,
eabd9909 (2020).

42.   S. S. Usmani *et al.*, THPdb: Database of FDA-approved peptide and protein therapeutics. *PLoS
ONE*. **12**, e0181748 (2017).

43.   J. Park *et al.*, Engineered receptors for human cytomegalovirus that are orthogonal to normal
human biology. *PLoS Pathog.* **16**, e1008647 (2020).

44.   Y. Imai *et al.*, Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature*.
436, 112–116 (2005).

45.   B. Treml *et al.*, Recombinant angiotensin-converting enzyme 2 improves pulmonary blood flow
and oxygenation in lipopolysaccharide-induced lung injury in piglets. *Crit. Care Med.* **38**, 596–
601 (2010).

46.   P. Verdecchia, C. Cavallini, A. Spanevello, F. Angeli, The pivotal link between ACE2 deficiency
and SARS-CoV-2 infection. *Eur J Intern Med*. **76**, 14–20 (2020).

47.   D. Wang *et al.*, Renin-angiotensin-system, a potential pharmacological candidate, in acute
respiratory distress syndrome during mechanical ventilation. *Pulm Pharmacol Ther*. **58**, 101833
(2019).

48.   M. K. Chung *et al.*, SARS-CoV-2 and ACE2: The biology and clinical data settling the ARB and
ACEI controversy. *EBioMedicine*. **58**, 102907–102907 (2020).

49.   J. A. Johnson, J. West, K. B. Maynard, A. R. Hemnes, ACE2 improves right ventricular function
in a pressure overload model. *PLoS ONE*. **6**, e20828 (2011).

50.   P. Liu *et al.*, Novel ACE2-Fc chimeric fusion provides long-lasting hypertension control and
organ protection in mouse models of systemic renin angiotensin system activation. *Kidney Int.* **94**,
114–125 (2018).

51.   M. Haschke *et al.*, Pharmacokinetics and pharmacodynamics of recombinant human angiotensin-
converting enzyme 2 in healthy human subjects. *Clin Pharmacokinet*. **52**, 783–792 (2013).

52.   A. Khan *et al.*, A pilot clinical trial of recombinant human angiotensin-converting enzyme 2 in
acute respiratory distress syndrome. *Crit Care*. **21**, 234 (2017).

53.   N. Iwanaga *et al.*, Novel ACE2-IgG1 fusions with improved activity against SARS-CoV2.
bioRxiv, 2020.06.15.152157 (2020).

54.   E. Procko *et al.*, Computational design of a protein-based enzyme inhibitor. *J. Mol. Biol.* **425**,
3563–3575 (2013).

549