Donor-recipient interactions drive dynamics of horizontal gene transfer via natural competence

Yu-Yu Cheng[1], James M. Papadopoulos[1,3], Tanya Falbel[2], Briana M. Burton[2], and Ophelia S. Venturelli[1,2,3*]

[1]Department of Biochemistry, University of Wisconsin - Madison, WI, United States
[2]Department of Bacteriology, University of Wisconsin - Madison, WI, United States
[3]Department of Chemical & Biological Engineering, University of Wisconsin-Madison

*To whom correspondence should be addressed: venturelli@wisc.edu

ABSTRACT

Horizontal gene transfer (HGT) in microbial communities is shaped by a complex web of abiotic and biotic interactions. We investigate the role of donor-recipient interactions on the dynamics of HGT via natural competence in a synthetic community. We show that the donor can play an active role in the gene transfer process that depends on the source of the transferred DNA. The efficiency of plasmid gene transfer depends on the abundance of the donor strain, whereas donor growth impairment augments the efficiency of chromosomal gene transfer. We show that antibiotic stress can either diminish or enhance the efficiency of gene transfer. A dynamic computational model of the system captures the impact of donor growth perturbations on gene transfer efficiency. Our results suggest that anti-HGT strategies targeting the donor strain can lead to opposing effects on the rate of HGT that depend on the microbial interaction network and source of transferred DNA.

INTRODUCTION

Horizontal gene transfer (HGT) is a major mechanism of genetic variation in microbial communities that enables the acquisition of new functional capabilities (*1*). Horizontally acquired sequences can provide a selective advantage by facilitating evolutionary adaptation to changing environments (*2*). Conjugation and natural transformation are prevalent processes that enable HGT in bacterial communities. Conjugation involves cell-to-cell contact between a donor and recipient cell and thus the live donor cell actively participates in the HGT process (*3*). By contrast, extracellular DNA (eDNA) can be acquired by a recipient cell that has activated the natural competence pathway in the absence of living donor cells (*4*). The capability for natural competence is widespread across Gram-positive and Gram-negative bacteria (*5*) and is also a common trait among bacterial pathogens (*6*). For example, inter-species or intra-strain gene transfer (*7–9*) via natural transformation is implicated in the ability of *Streptococcus pneumonia* (*S. pneumonia*) to adapt and persist on a human host. In *Bacillus subtilis* (*B. subtilis)*, natural competence is a major driver of genome diversity, which has enabled this species to secure niches in animal, plant, soil and marine environments (*10*).

The molecular and ecological factors that determine the rate of eDNA release or transfer of DNA between donor and recipient in microbial communities are not well understood. The current paradigm for natural competence requires the presence of eDNA and recipient cells that have activated the natural competence program. However, interactions between living donor and recipient have been shown to influence HGT frequencies in microbial communities. Specific naturally competent species can exploit predation to enhance DNA release from intact or recently killed donor cells. For example, *Acinetobacter baumannii*, *Vibrio cholerae* and *S. pneumoniae* can play an active role in gene transfer via predation using type-VI secretion systems or bacteriocins to enhance DNA release (*11–13*). Further, spatial proximity was a major factor for gene transfer between specific donors and recipients using conjugation-independent mechanisms. For instance, spatial proximity was shown to enhance intra-strain gene transfer in *B. subtilis*, *Porphyromonas gingivalis* and *Pseudomonas stutzeri*, as well as inter-species gene transfer between the donor *Escherichia coli* (*E. coli*) and recipient *Vibrio* species (*14–17*). However, there are many unresolved questions about the role of donor-recipient interactions on gene transfer via natural competence in microbial communities.

The source DNA for natural transformation can be derived from the donor cell chromosome or plasmids harbored by the donor, as opposed to conjugation which is primarily selective for mobile plasmids. Plasmids or chromosome as source molecules for gene transfer exhibit differences in copy number (*18*, *19*), stability in the extracellular environment (*20*), release profiles as a function of growth stage (*21–23*) and molecular mechanisms of DNA uptake by the recipient cell (*5*). In addition to these properties, multi-copy plasmids can impose a substantial metabolic burden by sequestering intracellular cellular resources, leading to substantial changes

in metabolic and stress response activities, cell morphology and growth rate (*24–26*). However,
the effects of plasmid metabolic burden on rates of HGT in microbial communities has not been
investigated.
To address these gaps, we investigate gene transfer dynamics in a synthetic consortium
composed of the genetically manipulable donor and recipient species *E. coli* and *B. subtilis*. Using
a systems-level approach, we demonstrate that donor-recipient interactions and the source of the
transferred sequences (plasmid or chromosome) are major factors influencing the efficiency of
HGT and its dependence on environmental factors. We find that the presence of the *E. coli* donor
substantially enhances plasmid transfer compared to purified DNA, whereas the reciprocal trend
is observed for chromosomal transfer. We construct dynamic computational models of species
growth and HGT via natural competence to further dissect the interactions between donor and
recipient in the system. To test the model predictions, we selectively modulate the growth of the
donor by administering antibiotics, introducing plasmids with a range of metabolic costs or tuning
the rate of cell lysis. Our data show that the abundance of the donor is positively correlated with
plasmid HGT efficiency, whereas substantial donor growth impairment is required for transfer of
chromosomal sequences, consistent with the model predictions. Antibiotic stress can either
enhance or reduce the transfer efficiency of plasmid derived genes. In sum, our results indicate
environmental perturbations that impact the growth of the donor in microbial communities can
differentially impact horizontal gene transfer rates derived from plasmid or chromosomal origins.
70
71 **RESULTS**
72
Temporal characterization of inter-species gene transfer in a synthetic community
To study the factors influencing the frequency of HGT in microbial communities, we constructed
a synthetic community composed of two distantly related species *E. coli* (donor) and *B. subtilis*
(recipient). An integration cassette comprised of 500 bp sequences homologous to the *B. subtilis*
chromosome flanking a spectinomycin resistance gene (*specR*) was introduced into an *E. coli*
plasmid or an erythromycin resistance gene (*ermR*) was integrated onto the *E. coli* chromosome.
To characterize the dynamics of gene transfer, we performed time-series measurements of
absolute species abundance, integration cassette extracellular DNA (eDNA) concentration and *B.*
subtilis transformation frequency in the presence of purified DNA (100 ng mL^{-1} plasmid or
chromosomal DNA) or living *E. coli* harboring the plasmid (*Ec-P*) or chromosomal cassette (*Ec-*
C) in monoculture or co-culture (**Figs. 1 and 2 (A and E)**). Transformation frequency or HGT
efficiency was defined as the fraction of *B. subtilis* colony forming units (CFU) that were resistant
to the antibiotic gene contained in the integration cassette divided by the total number of *B. subtilis*
CFU. To enable control of competence activity, a second copy of the master regulator *comK* was
introduced onto the *B. subtilis* chromosome under control of a xylose-inducible promoter (*27*).
B. subtilis exhibited a lower growth rate and carrying capacity in the presence of *E. coli*,
whereas the presence of *B. subtilis* did not impact the growth of *E. coli*, indicating a negative inter-
species interaction from *E. coli* to *B. subtilis* (**Fig. 2, B and F**). Although the *E. coli* growth
response was not altered in the presence of *B. subtilis*, the eDNA concentration was substantially
higher in the co-culture than in the *E. coli* monoculture (**Fig. 2, C and G**). Notably, the efficiency
of plasmid transfer was significantly enhanced across all time points by up to ~480-fold at 8 hr in
the co-culture compared to *B. subtilis* monoculture supplemented with purified plasmid DNA,
demonstrating that inter-species interactions were a critical determinant of HGT efficiency (**Fig.**
2D). Corroborating this result, the HGT efficiency of a co-culture containing native *comK B. subtilis*
PY79 (WT *B. subtilis*) and *Ec-P* was substantially higher than the WT *B. subtilis* monoculture
exposed to purified plasmid DNA, indicating that the observed increase in HGT efficiency was not
attributed to synthetic control of the competence pathway (**fig. S1A**). In addition, *E. coli* negatively
impacted WT *B. subtilis* growth by a smaller magnitude compared to engineered *B. subtilis*,
suggesting that the xylose-induced competence-dependent growth arrest may further reduce the

competitive ability of *B. subtilis* in the community (**Fig. 2 (B and F) and fig. S1B**). By contrast,
transformants were not detected at any time point in the *Ec-C* co-culture and a low transformation
frequency was observed in the *B. subtilis* monoculture in the presence of purified chromosomal
DNA, indicating that the efficiency of chromosomal transfer was reduced in the presence of the
donor species (**Fig. 2H**).

The critical role of the *E. coli* donor in enhancing plasmid transfer efficiency suggested
that plasmid transfer may involve mechanisms akin to conjugation. Since natural competence
involves eDNA released from the donor, we tested whether eDNA was required for inter-species
gene transfer by introducing a DNA degrading enzyme (DNase I) into the co-culture. In the
presence of DNase I, no transformants were detected in the *Ec-P* and *B. subtilis* co-culture,
corroborating that HGT occurred via natural transformation (**fig. S2A**).

Since spatial proximity between donor and recipient was shown to enhance gene transfer
by natural competence in other systems (*11*, *12*, *14*, *17*), we next sought to characterize the effect
of spatial proximity in the *Ec-P* and *B. subtilis* co-culture. *Ec-P* and *B. subtilis* were physically
separated by a 0.4 µm filter, which prevented cell-to-cell contact while allowing exchange of
diffusible molecules (**fig. S2B**). The HGT efficiency of the physically separated co-culture
condition was 25-fold lower compared to the well-mixed condition in the absence of the filter,
indicating that spatial proximity between donor and recipient enhanced HGT efficiency.

We next used fluorescence microscopy to visualize the spatial distribution of *E. coli* and
B. subtilis cells in the mixed culture. The RFP-labeled *E. coli* and GFP-labeled *B. subtilis* were in
close spatial proximity after 3 hr (**fig. S2C**). To quantify the spatial distribution of *B. subtilis* in the
presence and absence of *E. coli*, each image was partitioned into regions of interest (ROI)
containing single or groups of cells in close spatial proximity. Our results showed that 99% of *B.*
subtilis (at least 1000 cells) were found in close proximity to *E. coli* and the number of cells within
ROI containing both species was higher on average compared to the number of cells within ROI
containing a single species (**fig. S2D**). In addition, the average number of *E. coli* cells within *E.*
coli only ROI was higher than the number of *B. subtilis* in *B. subtilis* only ROI (**fig. S2D**). These
data suggest that *E. coli* can self-aggregate in the presence or absence of *B. subtilis*, whereas
aggregation of *B. subtilis* occurs more frequently in the presence of *E. coli*. In sum, these data
suggest that spatial proximity and cell-to-cell contact between donor and recipient may influence
plasmid transfer efficiency via natural competence.

The *B. subtilis* competence developmental program is regulated by a complex and
dynamic regulatory network that activates the pathway in a subpopulation of cells in response to
specific extracellular signals (*28*). Therefore, we sought to test whether the presence of *E. coli*
impacted the activation response of the competence pathway. To quantify the expression of the
competence pathway, we performed time-series flow cytometry measurements of *B. subtilis*
harboring a *comK* promoter (P_{comK}) fused to GFP in the presence and absence of *Ec-P* labeled
with constitutive RFP. In the presence and absence of *Ec-P*, the fraction of cells expressing GFP
(GFP-ON state) increased and then decreased over time, consistent the previously characterized
transient activation of the pathway in a sub-population of cells (**fig. S3**) (*28*). These data suggest
that the observed enhancement of plasmid transfer efficiency in the presence of *Ec-P* was not
attributed to changes in the activation response of competence pathway across the population.

We next explored whether the differences in HGT efficiencies in the co-culture compared
to monoculture conditions could be explained by variations in eDNA concentration. The eDNA
concentrations observed in the co-cultures and monocultures (100 ng mL^{-1} purified DNA) were in
the saturated regime of the eDNA-transformation frequency dose-response curve ($\sim 10^2$-10^3 ng
mL^{-1} for plasmid and $\sim 10^0$-10^2 ng mL^{-1} for chromosome) (horizontal lines in **Fig. 2I**), suggesting
that eDNA concentration was not a limiting factor for gene transfer. After accounting for
differences in gene copy numbers, the chromosome and plasmid transformation frequency were
similar for the *B. subtilis* monoculture exposed to purified DNA, with a moderate and statistically

significant increase in efficiency for chromosome than plasmid derived genes (**Fig. 2J**). These
data suggest the majority of the differences in the efficiency of chromosomal and plasmid gene
transfer in the *B. subtilis* monoculture condition could be explained variation in gene copy number,
as opposed to other properties of the DNA including topology (e.g. circular, linear or
strandedness), size, stability, or sequence. By ruling out *B. subtilis* competence activity, eDNA
concentration and properties as major determinants of transformation frequency in the synthetic
community, we next investigated how the inter-species interaction between *E. coli* and *B. subtilis*
influenced HGT dynamics.

Dynamic computational modeling of community dynamics and HGT
To interrogate the mechanisms that influence gene transfer, we developed a dynamic
computational model that represents species growth modified by inter-species interactions, eDNA
release and HGT (**Supplementary Materials**). The model captures the temporal changes in live
E. coli (*E*), dead *E. coli* (*Ed*), *B. subtilis* (*B*), eDNA (*G*), and transformed *B. subtilis* (*Bt*) (**Fig. 3A**).
We represent species growth dynamics and microbial interactions using terms from the Lotka-
Volterra model (*29*). To account for the time-delays in eDNA release relative to *E. coli* growth (**Fig.
2, B, C, F, and G**), *E* is converted into *Ed* at a constant rate and *Ed* releases *G* as a function of *B*
and *Ed.* Our data showed that the extracellular concentration of plasmid increased in the *B.*
subtilis monoculture condition, whereas the extracellular chromosome concentration displayed
the reciprocal temporal trend (**Fig. 2, C and G**). To capture these temporal trends, we assume
that *G* can be produced at a rate proportional to *B* and degraded at a rate proportional to *B* or *E*.
In addition, we assume the transformation frequency of *B* depends on *E* and the presence of *G*,
to represent the observed differences in transformation frequency in co-culture and monoculture
conditions (**Fig. 2, D and H**).

The model was fit to the time-series measurements of CFU, eDNA, and transformation
frequency and was able to recapitulate the temporal changes across conditions (**Fig. 2, B, C, D,**
F, G, and H). To further understand the differences in molecular mechanisms between the *Ec-P*
and *Ec-C* co-cultures, we computed the fold-change of each inferred parameter between the two
models (**Fig. 3A**). The inter-species interaction coefficient α_{eb} representing the impact of *B* on the
growth rate of *E* was substantially larger in magnitude in the *Ec-P* model compared to the *Ec-C*
model, suggesting that plasmid metabolic burden can alter the magnitude of inter-species
competition by reducing the fitness of *E. coli,* which in turn enhances the competitive ability of *B.*
subtilis. In addition, the chromosome degradation rate γ_g was significantly larger than plasmid,
consistent with the enhanced persistence of plasmid than chromosomal DNA over time in the
environment (*20*). To investigate the contribution of each parameter to the efficiency of HGT, each
parameter was perturbed by $\pm 20\%$ and the transformation frequency, defined as the ratio of *Bt*
to the sum of *B* and *Bt*, was determined at 6 hr (**Fig. 3B**). Notably, increasing the death rate of
Ec-P or *Ec-C* (γ_e) has opposing effects on the HGT efficiency in the model.

To test the model predicted effects of *E. coli* death rate on HGT efficiency, we introduced
an IPTG-inducible phage φX174 lysis gene *E* into *Ec-P* and *Ec-C* to program the donor lysis rate
(**Fig. 3, C and E**). The eDNA concentration increased substantially as a function of IPTG,
demonstrating that enhanced expression of lysis gene *E* triggered donor lysis and eDNA release
(**fig. S4**). In both communities, the abundance of *B. subtilis* and *E. coli* displayed opposing trends
as a function of IPTG, consistent with relationship between γ_e and species abundance in the
models (**Fig. 3, C, D, E and F**). Notably, the transformation frequency in the *B. subtilis* and *Ec-P*
co-culture exhibited a decreasing trend with IPTG concentration, consistent with the relationship
between γ_e and transformation frequency in the plasmid model (**Fig. 3, C and D**). The reciprocal
pattern was observed for the *Ec-C* community wherein transformants were only detected at high
IPTG concentrations, reflecting a moderate increase in transformation frequency as a function of
γ_e in the chromosome model (**Fig. 3, E and F**). These data demonstrated that substantial growth

inhibition of *Ec-C* was required to observe transformants above the detection limit. Inhibition of
Ec-C growth in the presence of high IPTG concentrations enhanced both eDNA concentration
and *B. subtilis* abundance, suggesting that both of these factors were critical for efficient
chromosomal transfer (**Fig. 3 (E and F) and fig. S4B**).
To further test the predictions of the models, we varied both the initial proportion and total
density of *B. subtilis* and *Ec-P* or *Ec-C* in the co-cultures and measured transformation frequency
at 6 hr. Our model captured the trends across the different conditions for the *Ec-P* co-culture,
including an enhanced number of *B. subtilis* transformants in conditions with high initial species
densities (**fig. S5A**). The measured transformation frequency for the *Ec-C* co-culture was below
the level of detection across the majority of conditions, reflecting the model prediction of a low
number of *B. subtilis* transformants across all conditions (**fig. S5B**). In sum, our model accurately
forecasted the differential roles of donor cell lysis rates in the *Ec-P* and *Ec-C* co-cultures, as well
as the effects of initial species densities on HGT efficiency.
The eDNA release rates (γ_e and λ_{d1}) are larger for *Ec-P* than *Ec-C*, suggesting that *E. coli*
lysis is enhanced in the presence of the plasmid (**Fig. 3A and table S1**). To test this prediction,
we varied the degree of plasmid-mediated metabolic burden by introducing a set of plasmids with
different copy numbers and sizes into *Ec-C* (**Fig. 3G**). Corroborating the plasmid metabolic burden
effect, the *E. coli* doubling time was correlated with the predicted number of plasmid DNA base
pairs per cell (**fig. S6**). The extracellular chromosome concentration increased with plasmid
metabolic burden and the donor strains harboring pSC101* and ColE1 exhibited a higher lysis
rate in the presence of *B. subtilis* compared to monoculture (**Fig. 3G**). Notably, transformants
were not observed in the presence of any of the *Ec-C* donor strains except the strain harboring
the plasmid with the highest metabolic cost (pBB275), demonstrating that plasmid metabolic
burden could substantially enhance the efficiency of chromosomal HGT in microbial communities
(**Fig. 3H**). In sum, the abundance of the recipient, lysis rate of the donor and plasmid metabolic
burden enhanced the efficiency of chromosomal transfer in the synthetic microbial community.
227
Impact of antibiotics that selectively target donor growth on horizontal gene transfer
Antibiotics have been shown to impact the frequencies of HGT via induction of conjugation and
competence pathways (*30–32*). To investigate how antibiotics impact gene transfer dynamics in
the synthetic community, we quantified the temporal change in eDNA release, absolute species
abundance and HGT efficiency in the co-culture in response to the antibiotics streptomycin
(bactericidal) or chloramphenicol (bacteriostatic). We used sub-lethal antibiotic concentrations for
B. subtilis to study how selective inhibition of the donor growth rate impacts HGT efficiency (**fig.
S7**). In response to streptomycin, the growth of *Ec-P* and *Ec-C* were impaired, whereas the growth
of *B. subtilis* was enhanced compared to the no antibiotic control, consistent with an attenuation
of the inhibitory interaction impacting *B. subtilis* (**Figs. 3A, 4 (A, B, E, and F), and table S1**). The
eDNA concentration was higher in the *Ec-P* and *Ec-C* co-cultures in the presence of streptomycin,
indicating that streptomycin enhanced the lysis rate of *E. coli* (**Fig. 4, C and G**). Although the
concentration of eDNA was augmented in the presence of streptomycin, the transformation
frequency in the *Ec-P* co-culture was significantly reduced compared to the no antibiotic control
(**Fig. 4D**), corroborating the key role of *Ec-P* abundance on HGT efficiency (**Fig. 3, C and D**). By
contrast, the transformation frequency in the *Ec-C* co-culture in the presence of streptomycin was
enhanced compared to the no antibiotic condition, consistent the major contribution of the
abundance of the recipient to chromosomal transfer (**Figs. 3E and 4H**). In the presence of
streptomycin, *Ec-P* failed to grow in the community, whereas *Ec-C* persisted over time, indicating
that plasmid metabolic burden not only reduced *E. coli* growth and enhanced lysis, but also
magnified antibiotic sensitivity (**Figs. 4, A and E**).
Exposure to sub-lethal concentration of chloramphenicol had opposing effects on the
plasmid and chromosomal temporal release profiles, while consistently reducing the growth rate
of *E. coli* (**Fig. 4, A, C, E, and G**). These temporal trends are consistent with the differential impact

of sub-lethal chloramphenicol concentrations on the stringent response, which leads to enhanced
replication of plasmids regulated by relaxed control while inhibiting chromosomal replication (*33*).
The presence of chloramphenicol had differential impacts on HGT efficiency in the *Ec-C* or *Ec-P*
co-cultures by enhancing or not altering gene transfer, respectively (**Fig. 4, D and H**). In sum, our
results showed that both the cold-shock antibiotic streptomycin or heat-shock antibiotic
chloramphenicol consistently enhanced chromosome transfer even though the rate of eDNA
release exhibited opposing temporal responses (higher and lower donor lysis rate in the presence
of streptomycin or chloramphenicol, respectively than the no antibiotic condition) (**Fig. 4H**) (*34*).
However, streptomycin and chloramphenicol exhibited differential effects on plasmid gene
transfer by either reducing or not altering HGT efficiency (**Fig. 4D**). Therefore, the mechanism of
action of the antibiotic combined with the origin of the transferred DNA molecule dictated the role
of the antibiotic on inter-species gene transfer in the synthetic community.
264
Investigating key molecular and ecological factors shaping gene transfer
We next sought to understand the relationships between different measured variables of the
system across all experiments to provide insight into the factors shaping inter-species plasmid or
chromosome gene transfer. The abundance of *Ec-C* and *B. subtilis* were negatively correlated,
consistent with the inferred negative inter-species interaction α_{be} of *E* on the growth rate of *B* in
the *Ec-C* model (**Fig. 3A, fig. S8A, and table S1**). The plasmid transfer frequency was positively
correlated with *E. coli* abundance (Pearson correlation coefficient *r* = 0.78, *p*-value = 9.37e-7),
consistent with our model's prediction (**Fig. 5 (A and B)**). By contrast, the *Ec-C* abundance was
not correlated to transformation frequency, mirroring the model prediction (**Fig. 5, C and D**).
Indeed, the efficiency of plasmid transfer was maximized for high *E. coli* abundance, whereas
chromosome transfer displayed the opposite relationship with *E. coli* abundance and required
strong inhibition of *E. coli* growth to observe transformants (**Fig. 5, E and G**). The Pearson
correlations between plasmid or chromosome eDNA concentration and transformation frequency
were not statistically significant, suggesting that eDNA was not limiting for HGT in these conditions
(**fig. S8, B and C**).
Inhibition of *E. coli* growth by antibiotic stress and programmed cell lysis resulted in an
increase in *B. subtilis* abundance by weakening the strong negative interaction impacting *B.*
subtilis (**Figs. 3 (C and E) and 4 (B and F)**). The abundance of *Ec-P* or *Ec-C* and the product of
B. subtilis abundance and time-lagged eDNA concentration were negatively correlated (Pearson
correlation coefficient *r* = -0.61, *p*-value = 0.00163 for plasmid and *r* = -0.89, *p*-value = 2.94e-11
for chromosome) (**Fig. 5, E and G**), reflecting the trend in the models (**Fig. 5, F and H**). The
strongest negative correlation between *E. coli* abundance and chromosome eDNA concentration
occurred for eDNA concentration measured four hours earlier, suggesting a time delay between
cell death and eDNA release (**fig. S8D**). In sum, our results demonstrate that the abundance of
the *E. coli* donor was a key determinant of efficient plasmid transfer. However, our data suggests
efficient chromosome transfer required both sufficient eDNA concentration and *B. subtilis*
abundance in the community (**Fig. 5G**). Therefore, our results suggest the strong negative
interaction from *Ec-C* to *B. subtilis* is the major factor that impeded chromosomal HGT in the
synthetic community and thus environmental perturbations that alleviated microbial competition
and increased donor cell lysis rates substantially enhanced chromosomal transfer.
295
296
DISCUSSION
The intracellular network and environmental signals that control the expression of the natural
competence pathway in *B. subtilis* in monoculture conditions have been extensively characterized
(*35, 36*). The current paradigm for gene transfer via natural competence in *B. subtilis* is that the
recipient drives control of this process by activating competence gene expression in a fraction of
the population in response to specific environmental signals (*37*). However, *B. subtilis* has evolved

in diverse microbial communities shaped by a complex and dynamic web of abiotic and biotic
interactions that impact the rates of HGT_(*10*).   Reduced complexity consortia composed of
genetically manipulable species allow precise control of environmental, ecological and
intracellular network factors and are thus powerful systems to dissect the variables shaping the
rate of HGT.
Using a detailed and quantitative analysis of a synthetic microbial community, our results
challenge the current paradigm of recipient driven control of gene transfer via natural competence
by demonstrating that the donor strain *E. coli* was a major driver of plasmid transfer as opposed
to the concentration of eDNA (*14*). Notably, efficient chromosomal HGT required strong growth
inhibition of the donor, which in turn increased the abundance of *B. subtilis* by impairing the
strength of microbial competition and eDNA concentration. Therefore, our results demonstrate
that the source of the transferred DNA determined the effect of the donor strain on the rates of
HGT in the community. The emergence of antibiotic resistance pathogens via HGT has
accelerated in recent years and is an imminent threat to global public health (*6*). Selective
targeting of donor strains harboring antibiotic resistance genes has been proposed as a strategy
to reduce the propagation of these genes through microbial communities (*38*, *39*). However, our
results indicate that altering the abundance and lysis rates of donor strains could either potentiate
or reduce the rate of HGT via natural competence depending on whether the antibiotic resistance
genes are derived from plasmid or chromosome origin.
Specific antibiotics have been shown to induce competence gene expression or enhance
transformation frequencies in *S. pneumoniae*, *Legionella pneumophila* and *B. subtilis*
monocultures due to activation of stress responses or variation in gene copy numbers (*14*, *31*, *32*,
40). However, the effect of an antibiotic on the efficiency of gene transfer depends on both the
specific mechanism of action and the effective concentration. Using the synthetic community, we
found that the bactericidal antibiotic streptomycin substantially reduced inter-species plasmid
transfer, whereas the bacteriostatic antibiotic chloramphenicol did not impact the efficiency of this
process. By contrast, both antibiotics enhanced inter-species transfer of the chromosomal derived
genes. Therefore, in a microbial community context, antibiotic stress could either diminish or
enhance inter-species gene transfer, which has implications for the design of antibiotic treatments.
Taken together, these results demonstrate that the plasmid or chromosome origin of transferred
DNA sequences is a major determinant of the impact of antibiotic stress on gene transfer due to
potential differences in the molecular mechanisms of HGT. Based on these results, the use of
selective chemical inhibitors that target natural competence activities in the recipient and growth
of the donor strain could be a promising strategy to reduce the rate of horizontal gene transfer of
plasmid derived genes in microbial communities_(*41*). By contrast, the rate of HGT of
chromosomal derived genes could be reduced by exploiting microbial competition to exclude the
recipient while promoting the growth of the donor.
Our results demonstrated that spatial proximity of *E. coli* and *B. subtilis* substantially
enhanced plasmid transfer. Consistent with this result, cell-to-cell contact and spatial proximity
have been shown to play an important role in inter-species and intra-strain gene transfer via
natural competence in several other microbial communities (*11*, *12*, *14*, *17*). The molecular
mechanisms driving these physical associations are not understood, with the exception of gene
transfer mediated by the contact-dependent type VI secretion system (T6SS). *E. coli* has been
shown to self-aggregate and adhere to different species including yeast and mammalian cells
(*42–44*), suggesting that this property could influence HGT efficiency in the synthetic community.
Future work could investigate the molecular mechanisms and genetic determinants of cell-to-cell
contact in the HGT process via natural competence in microbial communities.
We showed that plasmid metabolic burden reduced the growth rate of the donor and
enhanced cell lysis, leading to an increase in the rate of eDNA release and chromosomal HGT.
Future work will investigate if plasmid metabolic burden leads to higher lysis and gene transfer
efficiencies across diverse bacterial species. If this phenomenon is generalizable to other species,

the frequency of transferring genetic information encoded on the chromosome to other species
via natural competence may be higher for strains that harbor metabolically costly plasmids.
Therefore, prediction of potential hub donor species in HGT networks could leverage
metagenomics linking plasmids to host genomes (*45*). In addition, the effect of plasmid metabolic
burden on cell lysis presents potential challenges for biocontainment of engineered DNA
sequences in the environment (*46*).
The reduced complexity of the synthetic community enabled a detailed and quantitative
investigation of molecular and ecological factors influencing gene transfer dynamics. However,
this community does not capture the taxonomic diversity of natural microbiomes. In natural
communities, the donor and/or recipient taxa are largely unknown as well as the effects of
environmental perturbations such as antibiotic stress on HGT events. Horizontal gene transfer
events can be identified in natural communities *in situ* via time-series analysis of strain-level
genome sequences using high-throughput chromosomal confirmation capture (Hi-C) (*47*). Future
work could couple such multiplexed measurements of HGT within microbiomes with network
inference techniques to decipher the complex web of interactions between abiotic and biotic
factors that drive HGT. A deeper and quantitative understanding of the ecological and molecular
factors shaping HGT efficiency could be exploited to design anti-HGT microbiome interventions
to reduce the propagation of antibiotic resistance and virulence genes in microbiomes.
372

MATERIALS AND METHODS
Strain and plasmid construction
The strain background for *Ec-P* and *Ec-C* was *E. coli* MG1655. For flow cytometry analysis, RFP
driven by the P_{J23100} constitutive promoter was introduced onto the *Ec-P* genome downstream of
caiE gene using CRISPR gene editing techniques (*48*). *Ec-P* was transformed with the plasmid
pBB275 which contains a spectinomycin resistance gene (*specR*) flanked by two 500 bp *ycgO*
sequences homologous to the *B. subtilis* PY79 genome. To construct *Ec-C*, an erythromycin
resistance gene (*ermR*) and GFP driven by the P_{veg} constitutive promoter flanked by two 500 bp
yvbJ sequences homologous to *B. subtilis* PY79 were integrated onto the *E. coli* genome
(downstream of *caiE* gene). For programmable *E. coli* lysis, the plasmid pYC01 harboring the
φX174 lysis gene *E* driven by an IPTG-inducible promoter was used to control the donor lysis rate.
The plasmid pYC01 was constructed from pCSaE500 (Addgene #53182) by replacing the lysis
gene promoter with $P_{A1lacO-1}$ promoter (*49*). To characterize plasmid metabolic burden, ColE1
(pBbE2k-RFP), p15A (pBbA2a-RFP), and pSC101* (pBbS2c-RFP) were derived from the
BglBrick library were individually transformed into *Ec-C* using standard chemical transformation
techniques (*50*). To increase the transformation efficiency of *B. subtilis* PY79 in LB medium, a
xylose-inducible *comK* was integrated into *amyE* locus (*27*). To monitor the competence activity,
P_{comK}-*gfp* was introduced into *lacA* locus. To image the *B. subtilis* under microscope, $P_{hyperspank}$-
gfp (*51*) was integrated into *lacA* locus. All DNA oligos used for cloning were synthesized by
Integrated DNA Technologies (IDT). The list of plasmids and strains used in this study can be
found in **table S3**. The estimated copy numbers per cell for pSC101*, p15A, ColE1 and pBB275
were 3-4, 20-30, 50-70 and 50-70, respectively (*49*). The plasmid pBB275 was ~1.5 kb larger in
size than ColE1 and shared the same origin of replication. To normalize for differences in DNA
copy numbers, the eDNA copy number was calculated by NEBioCalculator with pBB275 size
4927 bp and *E. coli* MG1655 genome size 4639675 bp.
398

Bacterial culturing experiments
For bacterial culturing experiments, cells were first inoculated from a -80°C glycerol stock into 4
mL Lennox LB media (Sigma-Aldrich) containing antibiotics for selection and cultured at 37°C for
12 hours. The OD600 was measured by NanoDrop One (Thermo Fisher Scientific) and the cells
were diluted to an OD600 of 0.1 in 10 mL LB supplemented with 50 mM xylose (Thermo Fisher

Scientific) in 14 mL Falcon™ Round-Bottom Tubes (Thermo Fisher Scientific) and cultured at
37°C with shaking. For the *B. subtilis* monoculture experiments, the media was supplemented
with 100 ng mL^{-1} purified *E. coli* plasmid or genomic DNA. Plasmid and genomic DNA were
purified using a Plasmid Miniprep Kit (Qiagen) and DNeasy Blood & Tissue Kit (Qiagen). Samples
were collected every two hours for measuring colony forming units (CFU), extracellular DNA, and
flow cytometry. CFU measurements were performed by diluting the cell culture into phosphate-
buffered saline (PBS) buffer (Sigma-Aldrich) and plating onto LB agar plates containing 100 μg
ml^{-1} spectinomycin (GoldBio) for *Ec-P* or 1 μg mL^{-1} erythromycin (Sigma-Aldrich) and 25 μg/mL
lincomycin (VWR) for *Ec-C*. In the *Ec-P* co-culture, *B. subtilis* were selected for on LB agar plates
containing 5 μg mL^{-1} chloramphenicol (Sigma-Aldrich), 1 μg mL^{-1} erythromycin, and 25 μg mL^{-1}
lincomycin. In the *Ec-C* co-culture, *B. subtilis* were selected for on LB agar plates containing 5 μg
mL^{-1} chloramphenicol and 10 μg mL^{-1} kanamycin (Sigma-Aldrich). *B. subtilis* transformed with
plasmid pBB275 was selected on LB agar plates containing 5 μg mL^{-1} chloramphenicol, 1 μg mL^{-1}
erythromycin, 25 μg mL^{-1} lincomycin, and 100 μg mL^{-1} spectinomycin. *B. subtilis* transformed
with *Ec-C* chromosomal DNA was selected on LB agar plates containing 5 μg mL^{-1}
chloramphenicol, 10 μg/mL kanamycin, 1 μg mL^{-1} erythromycin, and 25 μg mL^{-1} lincomycin. Agar
plates were incubated at 37°C overnight and colonies were counted after 24 hours.
Transformation frequency was defined as the ratio of the number of *B. subtilis* CFU per mL to the
total number of *B. subtilis* CFU per mL. The detection limit of transformants was 1 CFU mL^{-1}
transformed *B. subtilis* divided by the total *B. subtilis* CFU mL^{-1}.
To test if the horizontal gene transfer occurred via natural transformation, 2 units mL^{-1}
DNase I (Thermo Fisher Scientific) was added to the culture. According to the manufacturer, the
quantity of 1 unit of DNase I can degrades 1 μg of plasmid DNA in 10 min at 37°C. Therefore, we
assumed that 2 units mL^{-1} DNase I can fully degrade 100 ng mL^{-1} plasmid administered to the *B.
subtilis* monoculture and extracellular plasmid released in co-culture the with *Ec-P*.
429
Quantitative real-time PCR measurements of extracellular DNA
To quantify eDNA in the cell culture, an aliquot of the cell culture was centrifuged at 5000xg for 5
min and the supernatant was filtered using a 0.2 μm Whatman Puradisc Polyethersulfone Syringe
Filter (GE Healthcare). To measure eDNA concentration by qPCR, the filtered supernatant was
added to a PCR reaction containing the SsoAdvanced™ Universal Probes Supermix (Bio-Rad)
and TaqMan primers with probes 5'FAM/ZEN/3'IABkFQ (Integrated DNA Technologies). The
TaqMan primers were designed to amplify a 129bp region of the *specR* gene on the plasmid
pBB275 in *Ec-P*, and a 145bp region of the *ermR* on the *Ec-C* genome. The plasmid and genomic
DNA standards were prepared as follows: plasmid was purified from *E. coli* DH5α using the
Plasmid Miniprep Kit (Qiagen) and genomic DNA was extracted from *Ec-C* using DNeasy Blood
& Tissue Kit (Qiagen). The extracted DNA was quantified by Quant-iT dsDNA Assay Kit (Thermo
Fisher Scientific) and then serially diluted to span a broad range of concentrations. The qPCR
reactions were run on the CFX Connect Real-Time PCR Detection System (Bio-Rad). The eDNA
concentrations were determined by interpolating the standard curve mapping DNA concentration
to Cq values.
445
Flow cytometer measurements of B. subtilis competence activity
To characterize competence activity in *B. subtilis*, a modified *B. subtilis* PY79 strain harboring
P$_{comK}$-*gfp* was co-cultured with *Ec-P* constitutively expressing RFP driven by a P$_{J23100}$ promoter.
Fluorescence was measured on a LSRFortessa X-20 Flow Cytometer (BD Biosciences). A blue
(488 nm) laser was used to quantify GFP and emission was detected using a 530/30 nm filter and
a yellow-green (561 nm) laser was used to excite RFP and emission was detected using a 610/20
nm filter. The GFP and RFP fluorescence distribution was displayed using the FlowJo software.
The RFP-OFF and RFP-ON cells were separated at 0 hr, but they merged after 2 hours (**fig. S3B)**.

To quantify the fraction of GFP-ON cells, a threshold in GFP (5000 au) was selected to separate
the OFF and ON states of the bimodal fluorescence distribution of the *B. subtilis* monoculture
in the presence of 50 mM xylose and 100 ng mL^{-1} purified plasmid DNA (**fig. S3A**) (*52*).
However, the choice of the threshold does not affect the qualitative trend between monoculture
and co-culture conditions. Data analysis was performed using custom MATLAB scripts.
459
Plate reader measurement of bacterial growth with antibiotics
To determine the sub-lethal concentration for *B. subtilis* and effect on *E. coli* growth, the overnight
culture of *Ec-P* and *B. subtilis* (with chloramphenicol resistance gene) was diluted to OD600 equal
to 0.1 in LB media with different concentrations of streptomycin (Sigma-Aldrich) or
chloramphenicol (Sigma-Aldrich). The cell cultures were incubated in a 96-well plate (VWR,
Catalog Number 82050-744) with shaking and sealed with the Breathe-Easy Adhesive Microplate
Seals (Thermo Fisher Scientific) in a TECAN Spark 10M Multimode Microplate Reader and
OD600 measurements were performed every five minutes.
468
Transwell experiment
To test if spatial proximity between donor and recipient enhances plasmid-mediated gene transfer,
cells were grown in 6-well Nunc Cell Culture Plate (Thermo Fisher Scientific). *Ec-P* and *B. subtilis*
or 100 ng mL^{-1} purified plasmid DNA (pBB275) and *B. subtilis* were separated by a 0.4 µm filter.
The plate was sealed with Breathe-Easy Adhesive Microplate Seals (Thermo Fisher Scientific)
and shaken incubated at 37°C with shaking. After 6 hr, samples from the two compartments
separated by the filter were plated on LB agar plates containing the appropriate antibiotics and
colonies were counted after 24 hr incubation at 37°C. Our controls showed that 0.01% of *E. coli*
were observed in the *B. subtilis* compartment at 6 hr, demonstrating that the filter functioned as a
physical barrier to cell diffusion between compartments.
479
Fluorescent microscopy imaging of bacteria
To image fluorescently labeled *E. coli* and *B. subtilis* in the co-culture, 4 µL of the cell cultures
were transferred to the glass slide after 3 hr incubation with shaking at 3hr at 37°C. To immobilize
the cells for imaging, 5 µL of 0.1 % (w/v) Poly-L-lysine (Millipore Sigma) was spread evenly on
the glass slide. Pipetting of the cultures was minimized to prevent the disruption of the physically
associated cells. Single cells were imaged by Nikon Eclipse Ti Microscope with 40X magnification.
Fluorescence was imaged using the following filters (Chroma): GFP: 470 nm/40 nm (ex),
525/50 nm (em) or RFP: 560 nm/40 nm (ex), 630/70 nm (em). GFP, RFP, and phase-contrast
images were taken from multiple spots on the glass slide. At least 1000 *B. subtilis* and *E. coli* cells
from multiple images were counted by ilastik (*53*) to quantify the degree of spatial proximity within
regions of interest (ROI).
491
Statistical analysis
We used an unpaired *t*-test to determine if there is a significant difference between the means of
two groups in **Figure 2J, 3C,** and **fig. S3C**. We used a Mann-Whitney U Test to determine if two
samples are derived from the same population in **fig. S2D**. Statistically different groups are
denoted by * *p*-value < 0.05, ** if *p*-value < 0.01, or *** if *p*-value < 0.001. The Pearson correlation
coefficient, *p*-value, and least-squares line were calculated to quantify the correlation between
two variables in **Figure 5 (A, C, E, and G)**, the inset in **fig. S6,** and **fig. S8**.

ACKNOWLEDGEMENTS
We would like to thank Jason Zuke and Zhengyi Chen for the assistance with the experiments,
and Victor Zavala, Joshua Pulsipher, and Ryan Clark for helpful discussions about the

computational model. This work was supported by the Defense Advanced Research Projects Agency (DARPA) Grant HR0011-19-2-0002.

AUTHOR CONTRIBUTIONS

Y.Y.C., B.M.B. and O.S.V. conceived the research. Y.Y.C. performed the experiments. O.S.V., B.M.B. and Y.Y.C. analyzed the data. O.S.V. and Y.Y.C. wrote the manuscript. Y.Y.C. and O.S.V. developed the model. J.M.P. constructed the *Ec-C* strain. O.S.V. secured the funding.

COMPETING INTERESTS

The authors declare no competing financial interests.

REFERENCES

1. S. M. Soucy, J. Huang, J. P. Gogarten, Horizontal gene transfer: Building the web of life. *Nat. Rev. Genet.* **16**, 472–482 (2015).

2. J. Ropars, R. C. Rodríguez de la Vega, M. López-Villavicencio, J. Gouzy, E. Sallet, É. Dumas, S. Lacoste, R. Debuchy, J. Dupont, A. Branca, T. Giraud, Adaptive horizontal gene transfers between multiple cheese-associated fungi. *Curr. Biol.* **25**, 2562–2569 (2015).

3. M. Clarke, L. Maddera, R. L. Harris, P. M. Silverman, F-pili dynamics by live-cell imaging. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 17978–17981 (2008).

4. S. Overballe-Petersen, K. Harms, L. A. A. Orlando, J. V. M. Mayar, S. Rasmussen, T. W. Dahl, M. T. Rosing, A. M. Poole, T. Sicheritz-Ponten, S. Brunak, S. Inselmann, J. de Vries, W. Wackernagel, O. G. Pybus, R. Nielsen, P. J. Johnsen, K. M. Nielsen, E. Willerslev, Bacterial natural transformation by highly fragmented and damaged DNA. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 19860–19865 (2013).

5. C. Johnston, B. Martin, G. Fichant, P. Polard, J.-P. Claverys, Bacterial transformation: distribution, shared mechanisms and divergent control. *Nat. Rev. Microbiol.* **12**, 181–196 (2014).

6. N. A. Lerminiaux, A. D. S. Cameron, Horizontal transfer of antibiotic resistance genes in clinical environments. *Can. J. Microbiol.* **65**, 34–44 (2019).

7. A. Jensen, O. Valdórsson, N. Frimodt-Møller, S. Hollingshead, M. Kilian, Commensal *Streptococci* serve as a reservoir for β-lactam resistance genes in *Streptococcus pneumoniae*. *Antimicrob. Agents Chemother.* **59**, 3529–3540 (2015).

8. J. Sauerbier, P. Maurer, M. Rieger, R. Hakenbeck, *Streptococcus pneumoniae* R6 interspecies transformation: genetic analysis of penicillin resistance determinants and genome-wide recombination events. *Mol. Microbiol.* **86**, 692–706 (2012).

9. R. Hakenbeck, T. Grebe, D. Zähner, J. B. Stock, β-lactam resistance in *Streptococcus pneumoniae*: penicillin-binding proteins and non-penicillin-binding proteins. *Mol. Microbiol.* **33**, 673–678 (1999).

10. P. H. Brito, B. Chevreux, C. R. Serra, G. Schyns, A. O. Henriques, J. B. Pereira-Leal, Genetic competence drives genome diversity in *Bacillus subtilis*. *Genome Biol. Evol.* **10**, 108–124 (2018).

11. R. M. Cooper, L. Tsimring, J. Hasty, Inter-species population dynamics enhance microbial horizontal gene transfer and spread of antibiotic resistance. *Elife*. **6**, e25950 (2017).

12. S. Borgeaud, L. C. Metzger, T. Scrignari, M. Blokesch, The type VI secretion system of *Vibrio cholerae* fosters horizontal gene transfer. *Science*. **347**, 63–67 (2015).

13. W.-Y. Wholey, T. J. Kochan, D. N. Storck, S. Dawid, Coordinated bacteriocin expression and competence in *Streptococcus pneumoniae* contributes to genetic adaptation through neighbor predation. *PLoS Pathog.* **12**, e1005413 (2016).

14. X. Zhang, T. Jin, L. Deng, C. Wang, Y. Zhang, X. Chen, Stress-induced, highly efficient,

donor cell-dependent cell-to-cell natural transformation in *Bacillus subtilis*. *J. Bacteriol.*
200, e00267-18 (2018).

15.   G. D. Tribble, T. W. Rigney, D. H. V. Dao, C. T. Wong, J. E. Kerr, B. E. Taylor, S. Pacha,
H. B. Kaplan, Natural competence is a major mechanism for horizontal DNA transfer in
the oral pathogen *Porphyromonas gingivalis*. *MBio*. **3**, e00231-11 (2012).

16.   G. J. Stewart, C. A. Carlson, J. L. Ingraham, Evidence for an active role of donor cells in
natural transformation of *Pseudomonas stutzeri*. *J. Bacteriol.* **156**, 30–35 (1983).

17.   J. H. Paul, J. M. Thurmond, M. E. Frischer, J. P. Cannon, Intergeneric natural plasmid
transformation between *E. coli* and a marine *Vibrio* species. *Mol. Ecol.* **1**, 37–46 (1992).

18.   R. Frame, J. O. Bishop, The number of sex-factors per chromosome in *Escherichia coli*.
Biochem. J. **121**, 93–103 (1971).

19.   C. L. Lee, D. S. W. Ow, S. K. W. Oh, Quantitative real-time polymerase chain reaction for
determination of plasmid copy number in bacteria. *J. Microbiol. Methods*. **65**, 258–267
(2006).

20.   D. Mao, Y. Luo, J. Mathieu, Q. Wang, L. Feng, Q. Mu, C. Feng, P. J. J. Alvarez,
Persistence of extracellular DNA in river sediment facilitates antibiotic resistance gene
propagation. *Environ. Sci. Technol.* **48**, 71–78 (2014).

21.   K. Matsui, N. Ishii, Z. Kawabata, Release of extracellular transformable plasmid DNA
from *Escherichia coli* cocultivated with algae. *Appl. Environ. Microbiol.* **69**, 2399–2404
(2003).

22.   M. G. Lorenz, D. Gerjets, W. Wackernagel, Release of transforming plasmid and
chromosomal DNA from two cultured soil bacteria. *Arch. Microbiol.* **156**, 319–326 (1991).

23.   A. L. Ibáñez de Aldecoa, O. Zafra, J. E. González-Pastor, Mechanisms and regulation of
extracellular DNA release and its biological roles in microbial communities. *Front.*
Microbiol. **8**, 1390 (2017).

24.   J. C. Diaz Ricci, M. E. Hernández, Plasmid effects on *Escherichia coli* metabolism. *Crit.*
Rev. Biotechnol. **20**, 79–108 (2000).

25.   H. Ingmer, C. Miller, S. N. Cohen, The RepA protein of plasmid pSC101 controls
Escherichia coli cell division through the SOS response. *Mol. Microbiol.* **42**, 519–526
(2001).

26.   M. E. Patient, D. K. Summers, ColE1 multimer formation triggers inhibition of *Escherichia*
coli cell division. *Mol. Microbiol.* **9**, 1089–1095 (1993).

27.   X.-Z. Zhang, Y.-H. P. Zhang, Simple, fast and high-efficiency transformation system for
directed evolution of cellulase in *Bacillus subtilis*. *Microb. Biotechnol.* **4**, 98–105 (2011).

28.   N. Mirouze, Y. Desai, A. Raj, D. Dubnau, Spo0A~P Imposes a Temporal Gate for the
Bimodal Expression of Competence in *Bacillus subtilis*. *PLoS Genet.* **8**, e1002586 (2012).

29.   O. S. Venturelli, A. V Carr, G. Fisher, R. H. Hsu, R. Lau, B. P. Bowen, S. Hromada, T.
Northen, A. P. Arkin, Deciphering microbial interactions in synthetic human gut
microbiome communities. *Mol. Syst. Biol.* **14**, e8157 (2018).

30.   R. T. Jeters, G.-R. Wang, K. Moon, N. B. Shoemaker, A. A. Salyers, Tetracycline-
associated transcriptional regulation of transfer genes of the *Bacteroides* conjugative
transposon CTnDOT. *J. Bacteriol.* **191**, 6374–6382 (2009).

31.   M. Prudhomme, L. Attaiech, G. Sanchez, B. Martin, J.-P. Claverys, Antibiotic stress
induces genetic transformability in the human pathogen *Streptoccus pneumoniae*.
Science. **313**, 89–92 (2006).

32.   X. Charpentier, E. Kay, D. Schneider, H. A. Shuman, Antibiotics and UV radiation induce
competence for natural transformation in *Legionella pneumophila*. *J. Bacteriol.* **193**,
1114–1121 (2011).

33.   A. Lim, E. Lee, E. Yip, H. Lam, K. Hui, Effects of the stringent response on pBR322
plasmid copy number in *Escherichia coli* strains. *J. Exp. Microbiol. Immunol.* **9**, 86–91
(2006).

34. M. Au, J. Ip, I. Tao, M. A. Zhang, Evaluation of stringency elicited by antibiotics on
plasmid copy number of pBR322 per genome in DH5α *Escherichia coli* cells. *J. Exp.*
Microbiol. Immunol. **13**, 83–88 (2009).
35. R. Rahmer, K. M. Heravi, J. Altenbuchner, Construction of a super-competent *Bacillus*
subtilis 168 using the P$_{mtlA}$-*comKS* inducible cassette. *Front. Microbiol.* **6**, 1431 (2015).
36. P. Tortosa, L. Logsdon, B. Kraigher, Y. Itoh, I. Mandic-Mulec, D. Dubnau, Specificity and
genetic polymorphism of the *Bacillus* competence quorum-sensing system. *J. Bacteriol.*
183, 451–460 (2001).
37. J.-P. Claverys, M. Prudhomme, B. Martin, Induction of Competence Regulons as a
General Response to Stress in Gram-Positive Bacteria. *Annu. Rev. Microbiol.* **60**, 451-
475 (2006).
38. R. López-igual, J. Bernal-bayard, A. Rodríguez-patón, J. Ghigo, D. Mazel, Engineered
toxin-intein antimicrobials can selectively target and kill antibiotic-resistant bacteria in
mixed populations. *Nat. Biotechnol.* **37**, 755–760 (2019).
39. D. Bikard, C. W. Euler, W. Jiang, P. M. Nussenzweig, G. W. Goldberg, X. Duportet, V. A.
Fischetti, L. A. Marraffini, Exploiting CRISPR-Cas nucleases to produce sequence-
specific antimicrobials. *Nat. Biotechnol.* **32**, 1146–1150 (2014).
40. J. Slager, M. Kjos, L. Attaiech, J.-W. Veening, Antibiotic-induced replication stress
triggers bacterial competence by increasing gene dosage near the origin. *Cell*. **157**, 395–
406 (2014).
41. A. Domenech, A. R. Brochado, V. Sender, K. Hentrich, B. Henriques-Normark, A. Typas,
J.-W. Veening, Proton motive force disruptors block bacterial competence and horizontal
gene transfer. *Cell Host Microbe*. **27**, 544–555 (2020).
42. A. K. Møller, M. P. Leatham, T. Conway, P. J. M. Nuijten, L. A. M. de Haan, K. A.
Krogfelt, P. S. Cohen, An *Escherichia coli* MG1655 lipopolysaccharide deep-rough core
mutant grows and survives in mouse cecal mucus but fails to colonize the mouse large
intestine. *Infect. Immun.* **71**, 2142–2152 (2003).
43. G. P. Dubey, S. Ben-Yehuda, Intercellular nanotubes mediate bacterial communication.
Cell. **144**, 590–600 (2011).
44. K. Jann, G. Schmidt, E. Blumenstock, K. Vosbeck, *Escherichia coli* adhesion to
Saccharomyces cerevisiae and mammalian cells: role of piliation and surface
hydrophobicity. *Infect. Immun.* **32**, 484–489 (1981).
45. Y. Suzuki, S. Nishijima, Y. Furuta, J. Yoshimura, W. Suda, K. Oshima, M. Hattori, S.
Morishita, Long-read metagenomic exploration of extrachromosomal mobile genetic
elements in the human gut. *Microbiome*. **7**, 119 (2019).
46. T. Blazejewski, H.-I. Ho, H. H. Wang, Synthetic sequence entanglement augments
stability and containment of genetic information in cells. *Science.* **365**, 595–598 (2019).
47. E. Yaffe, D. A. Relman, Tracking microbial evolution in the human gut using Hi-C reveals
extensive horizontal gene transfer, persistence and adaptation. *Nat. Microbiol.* **5**, 343–
353 (2020).
48. R. G. Egbert, H. S. Rishi, B. A. Adler, D. M. McCormick, E. Toro, R. T. Gill, A. P. Arkin, A
versatile platform strain for high-fidelity multiplex genome editing. *Nucleic Acids Res.* **47**,
3244–3256 (2019).
49. R. Lutz, H. Bujard, Independent and tight regulation of transcriptional units in *Escherichia*
coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. *Nucleic Acids Res.*
25, 1203–1210 (1997).
50. T. S. Lee, R. A. Krupa, F. Zhang, M. Hajimorad, W. J. Holtz, N. Prasad, S. K. Lee, J. D.
Keasling, BglBrick vectors and datasheets: A synthetic biology platform for gene
expression. *J. Biol. Eng.* **5**, 12 (2011).
51. W. Overkamp, K. Beilharz, R. D. O. Weme, A. Solopova, H. Karsens, Á. T. Kovács, J.
Kok, O. P. Kuipers, J.-W. Veening, Benchmarking various green fluorescent protein

variants in *Bacillus subtilis*, *Streptococcus pneumoniae*, and *Lactococcus lactis* for live
cell imaging. *Appl. Environ. Microbiol.* **79**, 6481–6490 (2013).
52.   B. J. Haijema, J. Hahn, J. Haynes, D. Dubnau, A ComGA-dependent checkpoint limits
growth during the escape from competence. *Mol. Microbiol.* **40**, 52–64 (2001).
53.   S. Berg, D. Kutra, T. Kroeger, C. N. Straehle, B. X. Kausler, C. Haubold, M. Schiegg, J.
Ales, T. Beier, M. Rudy, K. Eren, J. I. Cervantes, B. Xu, F. Beuttenmueller, A. Wolny, C.
Zhang, U. Koethe, F. A. Hamprecht, A. Kreshuk, ilastik: interactive machine learning for
(bio)image analysis. *Nat. Methods.* **16**, 1226–1232 (2019).
54.   O. Fridman, A. Goldberg, I. Ronin, N. Shoresh, N. Q. Balaban, Optimization of lag time
underlies antibiotic tolerance in evolved bacterial populations. *Nature.* **513**, 418–421
(2014).
667
668

Figure 1. Horizontal gene transfer from *E. coli* to *B. subtilis* in the synthetic bacterial consortium. *E. coli* donor has an integration cassette that is composed of an antibiotic resistance gene (AbR) flanked by two homologous sequences (HS) of *B. subtilis*. The integration cassette is either on the plasmid (*Ec-P*) or genome (*Ec-C*). Competent *B. subtilis* can take up DNA and the AbR can be integrated into *B. subtilis* genome via homologous recombination. To characterize the gene transfer process, *E. coli* (i), *B. subtilis* (ii), transformed *B. subtilis* (iii), and eDNA (iv) were measured over time.

Figure 2. Temporal characterization of gene transfer in the synthetic microbial community.
(A) Schematic of experimental design to characterize the temporal changes in plasmid gene transfer in the *B. subtilis* (blue) monoculture in the presence of purified plasmid DNA (100 ng mL^{-1}) or co-culture of *Ec-P* (purple) and *B. subtilis*. Extracellular plasmid release was characterized in the *Ec-P* monoculture. **(B)** Time-series measurements of *Ec-P* and *B. subtilis* CFU mL^{-1} in the monocultures (dashed) and co-culture (solid). Data points represent experimental measurements and lines denote model fits to the data. **(C)** Time-series measurements of extracellular plasmid DNA in the *Ec-P* and *B. subtilis* monocultures (dashed) or co-culture (solid). Data points represent experimental measurements and lines denote model fits to the data. **(D)** Time-series measurements of plasmid transformation frequency for the *B. subtilis* monoculture (dashed) in the presence of purified plasmid DNA 100 ng mL^{-1}) or co-culture with *Ec-P* (solid). Data points represent experimental measurements and lines denote model fits to the data. **(E)** Schematic of experimental design to characterize the temporal changes in chromosomal gene transfer in the *B. subtilis* (blue) monoculture supplemented with purified genomic *Ec-C* DNA (100 ng mL^{-1}) or co-culture of *B. subtilis* and *Ec-C* (purple). **(F)** Time-series measurements of *Ec-C* and *B. subtilis* CFU mL^{-1} in monoculture (dashed) and co-culture (solid). Data points represent experimental measurements and lines denote model fits to the data. **(G)** Time-series measurements of extracellular chromosomal DNA in the *Ec-C* and *B. subtilis* monocultures (dashed) and co-culture (solid). Data points and lines denote experimental measurements and model fits, respectively. **(H)** Time-series measurements of chromosome transformation frequency in the *B. subtilis* monoculture supplemented with purified genomic *Ec-C* DNA (100 ng mL^{-1}) or the co-culture (no transformants observed). Data points and lines represent experimental measurements and model fits to the data, respectively. **(I)** Relationship between purified plasmid or chromosome concentration and transformation frequency in the *B. subtilis* monoculture at 6 hr. Horizontal lines indicate the range of extracellular plasmid and chromosome concentrations measured in the *B. subtilis* monoculture with purified DNA (100 ng mL^{-1}) and co-cultures with *Ec-P* or *Ec-C* (**Fig. 2, C and G**). Data points and lines denote experimental measurements experimental measurements and the average of two biological replicates. **(J)** Relationship between eDNA copy number per mL of purified plasmid or chromosome and transformation frequency at 6 hr in the *B. subtilis* monoculture. All conditions were supplemented with 50 mM xylose.

Figure 3. Model prediction on the role of donor in plasmid-mediated and chromosome-mediated gene transfer. (A) Schematic of the key interactions in the dynamic computational model of growth and horizontal gene transfer. The model species include *E. coli* (*E*), dead *E. coli* (*E_d*), *B. subtilis* (*B*), eDNA (*G*), and transformed *B. subtilis* (*B_t*). The line width and color of each edge are proportional to the fold change of each parameter in the plasmid to chromosome model. **(B)** Relationship between each parameter and the fold change of the predicted transformation

frequency in the plasmid or chromosome model at 6 hr for 20% increase or decrease in the parameter value compared to the estimated parameter set. **(C)** Schematic of a co-culture experiment of *Ec-P* harboring an IPTG-inducible lysis gene and *B. subtilis* (left). Relationship between IPTG concentration and plasmid transformation frequency or CFU mL^{-1} (right) at 6 hr in the co-culture. Transformation frequency exhibited a decreasing trend with IPTG concentration (*p*-value < 0.01 (**) or *p*-value <0.001 (***) using an unpaired *t*-test. **(D)** *Ec-P* model prediction of the concentration (CFU mL^{-1}) of *E* and *B*, or plasmid transformation frequency at 6 hr as a function of the fold change in the *Ec-P* death rate γ_e compared to the estimated parameter value. **(E)** Schematic of a co-culture experiment of *Ec-C* harboring an IPTG-inducible lysis gene and *B. subtilis* (left). Relationship between IPTG concentration and chromosome transformation frequency or CFU mL^{-1} (right) at 6 hr in the co-culture. Transformants were detected at higher IPTG concentrations. Shaded regions denote the detection limits for transformation frequency. **(F)** *Ec-C* model prediction of the concentration (CFU mL^{-1}) of *E* and *B*, or chromosome transformation frequency at 6 hr as a function of the fold change in the *Ec-C* death rate γ_e compared to the estimated parameter value. We assume that the detection limit of transformants in the model is 1 CFU mL^{-1} B_t divided by the sum of B_t and *B* CFU mL^{-1}. **(G)** Schematic of experimental design (left) of co-cultures composed of *Ec-C* harboring a single plasmid with variable origin of replication and *B. subtilis* (solid) or *Ec-C* monoculture (dashed). Time-series measurements of extracellular chromosome concentration in the *Ec-C* monocultures or co-cultures. **(H)** Chromosome transformation frequency at 4 hr in each co-culture of *Ec-C* harboring different plasmids and *B. subtilis*. Shaded area indicates the transformation frequency detection limit. Lines and bars in **Fig. 3, C, E, G, and H** represent the average of 2-3 biological replicates. All conditions were supplemented with 50 mM xylose.

Figure 4. Impact of antibiotics on the dynamics of plasmid and chromosomal gene transfer in the synthetic microbial community. (A) Time-series measurements of *Ec-P* CFU mL^{-1} in co-culture with *B. subtilis* in the presence of streptomycin (10 µg mL^{-1}), chloramphenicol (5 µg mL^{-1}) or no antibiotic. Data points and lines represent experimental measurements and growth model fits to the data. Inset: Inferred doubling times based on the growth models. **(B)** Time-series measurements of *B. subtilis* CFU mL^{-1} in the co-culture with *Ec-P* in the presence of streptomycin (10 µg mL^{-1}), chloramphenicol (5 µg mL^{-1}) or no antibiotic. Data points and lines represent experimental measurements and growth model fits to the data. Inset: Inferred doubling time based on the growth models. **(C)** Time-series measurements of extracellular plasmid DNA in the *Ec-P* and *B. subtilis* co-culture. Data points and lines represent experimental measurements and averages of biological replicates, respectively. **(D)** Time-series measurements of plasmid transformation frequency in the *Ec-P* and *B. subtilis* co-cultures. Data points and lines denote experimental measurements and mean of biological replicates, respectively. **(E)** Time-series measurements of *Ec-C* CFU mL^{-1} in co-culture with *B. subtilis* in the presence of streptomycin (10 µg mL^{-1}), chloramphenicol (5 µg mL^{-1}) or no antibiotic. Data points and lines represent experimental measurements and growth model fits to the data. Inset: Inferred doubling times based on the growth models. **(F)** Time-series measurements of *B. subtilis* CFU mL^{-1} in the co-culture with *Ec-C* in the presence of streptomycin (10 µg mL^{-1}), chloramphenicol (5 µg mL^{-1}) or no antibiotic. Data points and lines represent experimental measurements and growth model fits to the data. Inset: Inferred doubling time based on the growth models. **(G)** Time-series measurements of extracellular chromosome DNA in the co-culture with *Ec-C*. Data points and lines denote experimental measurements and mean of biological replicates, respectively. **(H)** Time-series measurements of chromosome transformation frequency in the *Ec-C* and *B. subtilis* co-culture. Data points and lines denote experimental measurements and mean of biological replicates, respectively. All conditions were supplemented with 50 mM xylose.

Figure 5. Global trends in transformation frequency, *E. coli* CFU, *B. subtilis* CFU, and eDNA concentration across different experimental datasets. (A) Scatter plot of *Ec-P* CFU mL^{-1} and plasmid transformation frequency (Pearson correlation coefficient *r* = 0.78, *p*-value = 9.37e-7). The size of each data point is proportional to the eDNA concentration. **(B)** Scatter plot of plasmid model prediction of the relationship between *E* concentration and transformation frequency for a broad range of *E* death rates γ_e. Data points connected by separate lines denote different values of γ_e. Gray arrow indicates the direction of time. **(C)** Scatter plot of *Ec-C* CFU mL^{-1} and chromosome transformation frequency. The size of the data point is proportional to the eDNA concentration. **(D)** Scatter plot of chromosome model prediction of the relationship between *E* concentration and transformation frequency for a broad range of *E* death rates γ_e. Data points connected by separate lines denote different values of γ_e. Gray arrow indicates the direction of time. **(E)** Scatter plot of *Ec-P* CFU mL^{-1} and the product of *B. subtilis* CFU mL^{-1} and time-lagged eDNA concentration (Pearson correlation coefficient *r* = -0.61, *p*-value = 0.00163). The size of each datapoint is proportional the transformation frequency. **(F)** Scatter plot of the plasmid model prediction of the relationship between *E* and the product of *B* and time-lagged *G* concentration across a broad range of *E* death rates γ_e. Data points connected by separate lines denote different values of γ_e. The size of each data point is proportional to the transformation frequency. Gray arrow indicates the direction of time. **(G)** Scatter plot of *Ec-C* CFU mL^{-1} and the product of *B. subtilis* CFU mL^{-1} and time-lagged extracellular chromosome concentration (Pearson correlation coefficient *r* = -0.89, *p*-value = 2.94e-11). The size of each data point is proportional to the transformation frequency. Data points with red outlines denote measurements where no *B. subtilis* transformants were detected and the size of these data points are proportional to the detection limit. **(H)** Scatter plot of the chromosome model prediction of the relationship between *E* and the product of *B* and time-lagged *G* concentration across a broad range of *E* death rates γ_e. Data points connected by separate lines denote different values of γ_e. The size of each data point is proportional to the transformation frequency. The gray arrow indicates the direction of time. All conditions were supplemented with 50 mM xylose.