Expanding diversity of Asgard archaea and the elusive ancestry of eukaryotes

Yang Liu[1†], Kira S. Makarova[2†], Wen-Cong Huang[1†], Yuri I. Wolf[2], Anastasia Nikolskaya[2], Xinxu

Zhang[1], Mingwei Cai[1], Cui-Jing Zhang[1], Wei Xu[3], Zhuhua Luo[3], Lei Cheng[4], Eugene V. Koonin[2*], Meng

Li[1*]

1 Shenzhen Key Laboratory of Marine Microbiome Engineering, Institute for Advanced Study, Shenzhen

University, Shenzhen, Guangdong, 518060, P. R. China

2 National Center for Biotechnology Information, National Library of Medicine, National Institutes of

Health, Bethesda, Maryland 20894, USA

3 State Key Laboratory Breeding Base of Marine Genetic Resources, Key Laboratory of Marine Genetic

Resources, Fujian Key Laboratory of Marine Genetic Resources, Third Institute of Oceanography, State

Oceanic Administration, Xiamen 361005, P. R. China

4 Key Laboratory of Development and Application of Rural Renewable Energy, Biogas Institute of

Ministry of Agriculture, Chengdu 610041, P.R. China

† These authors contributed equally to this work.

*Authors for correspondence: koonin@ncbi.nlm.nih.gov or limeng848@szu.edu.cn

Running title: Asgard archaea genomics

Keywords:

Abstract

Comparative analysis of 162 (nearly) complete genomes of Asgard archaea, including 75 not reported previously, substantially expands the phylogenetic and metabolic diversity of the Asgard superphylum, with six additional phyla proposed. Phylogenetic analysis does not strongly support origin of eukaryotes from within Asgard, leaning instead towards a three-domain topology, with eukaryotes branching outside archaea. Comprehensive protein domain analysis in the 162 Asgard genomes results in a major expansion of the set of eukaryote signature proteins (ESPs). The Asgard ESPs show variable phyletic distributions and domain architectures, suggestive of dynamic evolution via horizontal gene transfer (HGT), gene loss, gene duplication and domain shuffling. The results appear best compatible with the origin of the conserved core of eukaryote genes from an unknown ancestral lineage deep within or outside the extant archaeal diversity. Such hypothetical ancestors would accumulate components of the mobile archaeal 'eukaryome' via extensive HGT, eventually, giving rise to eukaryote-like cells.

Introduction

The Asgard archaea are a recently discovered archaeal superphylum that is rapidly expanding, thanks to metagenomic sequencing (*1–5*). The Asgard genomes encode a diverse repertoire of eukaryotic signature proteins (ESPs) that far exceeds the diversity of ESPs in other archaea. The Asgard ESPs are particularly enriched in proteins involved in membrane trafficking, vesicle formation and transport, cytoskeleton formation and the ubiquitin network, suggesting that these archaea possess a eukaryote-type cytoskeleton and an intracellular membrane system (*2*).

The discovery of the Asgard archaea rekindled the decades old but still unresolved fundamental debate on the evolutionary relationship between eukaryotes and archaea that has shaped around the '2-domain (2D) versus 3-domain (3D) tree of life' theme (*6–8*). The central question is whether the eukaryotic nuclear lineage evolved from a common ancestor shared with archaea, as in the 3D tree, or from within the archaea, as in the 2D tree. The discovery and phylogenomic analysis of Asgard archaea yielded strong evidence in support of the 2D tree, in which eukaryotes appeared to share common ancestry with one of the Asgard lineages, Heimdallarchaeota (*1*, *2*, *5*). However, the debate is not over as arguments have been made for the 3D topology, in particular, based on the phylogenetic analysis of RNA polymerases, some of the most highly conserved, universal proteins (*9*, *10*).

Molecular phylogenetic methods alone might be insufficient to resolve the ancient ancestral relationship between archaea and eukarya. To arrive at a compelling solution, supporting biological evidence is crucial (*11*), because, for example, the transition from archaeal, ether-linked membrane lipids to eukaryotic, ester-linked lipids that constitute eukaryotic (and bacterial) membranes (*12*) and the apparent lack of the phagocytosis capacity in Asgard archaea (*13*) are major problems for scenarios of the origin of eukaryotes from Asgard or any other archaeal lineage. Some biochemical evidence indicates that Asgard archaea possess an actin cytoskeleton regulated by accessory proteins, such as profilins and gelsolins, and the endosomal sorting complex required for transport machinery (ESCRT) that can be predicted to function similarly to the eukaryotic counterparts (*14–16*). Generally, however, the biology of Asgard archaea remains poorly characterized, in large part, because of their recalcitrance to growth in culture (*17*). To date, only one Asgard archaeon, *Candidatus* Prometheoarchaeum syntrophicum strain MK-D1, has been isolated and grown in culture (*17*). This organism has been reported to form extracellular protrusions that are involved in its interaction with syntrophic bacteria, but no visible organelle-like structure and, apparently, little intracellular complexity.

The only complete, closed genome of an Asgard archaeon also comes from *Candidatus* P. syntrophicum strain MK-D1 (*17*) whereas all other genome sequences were obtained by binning multiple metagenomics

contigs. Furthermore, these genome sequences represent but a small fraction of the Asgard diversity that

has been revealed by 16S rRNA sequencing (*18*, *19*).An additional challenge to the study of the

relationship between archaea and eukaryotes is that identification and analysis of the archaeal ESPs are

non-trivial tasks due to the high sequence divergence of many if not most of these proteins. At present,

the most efficient, realistic approach to the study of Asgard archaea and their eukaryotic connections

involves obtaining high quality genome sequences and analyzing them using the most powerful and

robust of the available computational methods.

Here we describe metagenomic mining of the expanding diversity of the superphylum Asgard, including

the identification of six additional phylum-level lineages that thrive in a wide variety of ecosystems and

are inferred to possess versatile metabolic capacities. We show that these uncultivated Asgard groups

carry a broad repertoire of ESPs many of which have not been reported previously. Our in depth

phylogenomic analysis of these genomes provides insights into the evolution of Asgard archaea but calls

into question the origin of eukaryotes from within Asgard.

Results

Reconstruction of Asgard archaeal genomes from metagenomics data

We reconstructed 75 metagenome-assembled genomes (MAGs) from the Asgard superphylum that have

not been reported previously. These MAGs were recovered from various water depths of the Yap trench,

intertidal mangrove sediments of Mai Po Nature Reserve (Hong Kong, China) and Futian Mangrove

Nature Reserve (Shenzhen, China), seagrass sediments of Swan Lake Nature Reserve (Rongcheng,

China) and petroleum samples of Shengli oilfield (Shandong, China) (Supplementary Table 1,

Supplementary Figure 1). For all analyses described here, these 75 MAGs were combined with 87

publicly available genomes, resulting in a set of 162 Asgard genomes. The 75 genomes reconstructed here

were, on average, 82% complete and showed evidence of low contamination of about 3%, on average

(Supplementary Figure 2).

Classification of Asgard genes into clusters of orthologs

The previous analyses of Asgard genomes detected a large fraction of "dark matter" genes (*20*). For

example, in the recently published complete genome of *Candidatus* Prometheoarchaeum syntrophicum,

45% of the proteins are annotated as "hypothetical". We made an effort to improve the annotation of

Asgard genomes by investigating this dark matter in greater depth, and developing a dedicated platform

for Asgard comparative genomics. To this end, we constructed Asgard Clusters of Orthologous Genes

(asCOGs) and used the most sensitive available methods of sequence analysis to annotate additional

Asgard proteins, attempting, in particular, to expand the catalogue of Asgard homologs of ESPs (see

Materials and Methods for details).

Preliminary clustering by sequence similarity and analysis of the protein cluster representation across the

genomes identified the set of 76 most complete Asgard MAGs (46 genomes available previously and 30

ones reported here) that cover most of the group diversity (Supplementary Table 1). The first version of

the asCOGs presented here consists of 14,704 orthologous protein families built for this 76-genome set.

The asCOGs cover from 72% to 98% (92% on average) of the proteins in these 76 genomes (additional

data file 1). Many asCOGs include individual domains of large, multidomain proteins.

The gene commonality plot for the asCOGs shows an abrupt drop at the right end, which reflects a

surprising deficit of nearly universal genes (Fig. 1). Such shape of the gene commonality curve appears

anomalous compared to other major groups of archaea or bacteria with many sequenced genomes (*21*).

For example, in the case of the TACK superphylum of archaea, for which the number of genomes

available is similar to that for Asgard, with a comparable level of diversity, the commonality plot shows

no drop at the right end, but instead, presents a clear uptick, which corresponds to the core of genes

represented in (almost) all genomes (Fig. 1). Apparently, most of the Asgard genomes remain incomplete,

such that conserved genes were missed randomly. Currently, there are only three gene families that are

present in all Asgard MAGs, namely, a Zn-ribbon domain, a Threonyl-tRNA synthetase and an

aminotransferase (additional data file 1).

We employed the asCOG profiles to annotate the remaining 86 Asgard MAGs, including those that were

sequenced in the later stages of this work (Supplementary Table 1). On average, 89% of the proteins

encoded in these genomes were covered by asCOGs (Supplementary Table 1). Thus, the asCOGs

database appears to be an efficient tool for annotation and comparative genomic analysis of Asgard

MAGs and complete genomes.

Expanding the phylogenetic diversity of Asgard archaea

Phylogenetic analysis of the Asgard MAGs based on a concatenated alignment of 209 core asCOGs (see

Methods and additional data file 2) placed many of the genomes reported here into the previously

delineated major Asgard lineages (Fig. 2a, Supplementary Table 1, additional data file 2), namely,

Thorarchaeota (n=20), Lokiarchaeota (n=18), Hermodarchaeota (n=9), Gerdarchaeota (n=3),

Helarchaeota (n=2), and Odinarchaeota (n=1). Additionally, we identified 6 previously unknown major

Asgard lineages that appear to be strong candidates to become additional phyla (Fig. 2a and b,

Supplementary Table 1; see also Taxonomic description of new taxa in the Supplementary Information

and additional data file 1). A clade formed by As_085 and As_075 is a deeply branching sister group to

the previously recognized Heimdallarchaeota(*2*). Furthermore, our phylogenetic analysis supported the

further split of "Heimdallarchaeota" into 4 phylum-level lineages according to the branch length in the

concatenated phylogeny (see Materials and Methods; see also Taxonomic description of new taxa in the

Supplementary Information and additional data file 1). The putative phyla within the old

Heimdallarchaeota included the previously defined Gerdarchaeota (*4*), and three additional phyla that

could be represented by As_002 (LC2) , As_003 (LC3) and AB_125 (As_001), respectively. Another 3

previously undescribed lineages were related, respectively, to Hel-, Loki-, Odin- and Thorarchaeota.

Specifically, As_181, As_178 and As_183 formed a clade that was deeply rooted at the Hel-Loki-Odin-

Thor clade; As_129 and As_130 formed a sister group to Odinarchaeota; and a lineage represented by

As_086 was a sister group to Thorarchaeota. These results were buttressed by the 16S rRNA gene

phylogeny, comparisons of the mean amino acid identity and 16S rRNA sequence identity (Fig. 2b,

Supplementary Figure 3, Supplementary Figure 4, Supplementary Table 2 and Supplementary Table 3).

We propose the name Wukongarchaeota after Wukong, a Chinese legendary figure who caused havoc in

the heavenly palace, for the putative phylum represented by MAGs As_085 and As_075 (*Candidatus*

Wukongarchaeum yapensis), and names of Asgard deities in the Norse mythology for the other 5

proposed phyla: (1) Hodarchaeota, after Hod, the god of darkness, for MAG As_027 (*Candidatus*

Hodarchaeum mangrove); (2) Kariarchaeota, after Kari, the god of the North wind, for MAG As_030

(*Candidatus* Kariarchaeum pelagius); (3) Borrarchaeota after Borr, the creator god and father of Odin, for

MAG As_133 (*Candidatus* Borrarchaeum yapensis); (4) Baldrarchaeota, after Baldr, the god of light and

brother of Thor, for MAG As_130 (*Candidatus* Baldrarchaeum yapensis); (5) and Hermodarchaeota after

Hermod, the messenger of the gods, son of Odin and brother of Baldr, for MAG As_086 (*Candidatus*

Hermodarchaeum yapensis) (Fig. 2a). For details, see Taxonomic Description of new taxa in the

Supplementary Information.

The gene content of Asgard MAGs agrees well with the phylogenetic structure of the group. The phyletic

patterns of the asCOG form clusters that generally correspond to the clades identified by phylogenetic

analysis (Fig. 3a), suggesting that gene gain and loss within Asgard archaea largely proceeded in a clock-

like manner and/or that horizontal gene exchange preferentially occurred between genomes within the

same clade.

Phylogenomic analysis and positions of Asgard archaea and eukaryotes in the tree of life

The outcome of phylogenetic reconstruction, especially, when deep branchings are involved, such as

those that are relevant for the 3D vs 2D conundrum, depends on the phylogenetic methods employed, the

selection of genes for phylogeny construction and, perhaps most dramatically, on the species sampling

(*22–24*). In many cases, initially uncertain positions of lineages in a tree settle over time once more

representatives of the groups in question and their relatives become available.

In our analysis of the universal phylogeny, we aimed to make the species set for phylogenetic

reconstruction as broadly representative as possible, while keeping its size manageable, to allow the use

of powerful phylogenetic methods. The tree was constructed from alignments of conserved proteins of

162 Asgard archaea, 286 other archaea, 98 bacteria and 72 eukaryotes (see Supplementary Material and

Methods for details of the procedure including the selection of a representative species set and

Supplementary Table 4). Members of 30 families of (nearly) universal proteins that appear to have

evolved without much HGT and have been previously employed for the reconstruction of the tree of life

(*25*) were used to generate a concatenated alignment of 7411 positions, after removing low information

content positions (Supplementary Table 4). For the phylogenetic reconstruction, we used the IQ-tree

program with several phylogenetic models (see Methods and Supplementary Table 5 for details).

Surprisingly, the resulting trees had the 3D topology, with high support values for all key bifurcations

(Fig. 2c, additional data file 2).

A full investigation of the effects of different factors, in particular, the marker gene selection, on the tree

topology is beyond the scope of this work. Nonetheless, we addressed the possibility that the 3D topology

resulted from the model used for the tree reconstruction and/or the species selection. To this end, we

constructed 100 trees from the same alignment by randomly sampling 5 representatives of Asgard

archaea, other archaea, bacteria and eukaryotes each. For these smaller sets of species, the best model

identified by Williams et al. (LG+C60+G4+F) could be employed, resulting in 50 3D and 50 2D trees

(Supplementary Table 5, additional data file 2). Because IQ-tree identified this model as over-specified

for such a small alignment, we also tested a more restricted model (LG+C20+G4+F), obtaining 58 3D and

42 2D trees for the same set of 100 samples (Supplementary Table 5, additional data file 2).

The results of our phylogenetic analysis indicate that: 1) species sampling substantially affects the tree

topology; 2) even the set of most highly conserved genes that appear to be minimally prone to HGT,

yields conflicting signals for different species sets. Additional markers, less highly conserved and more

prone to HGT, are unlikely to improve phylogenetic resolution and might cause systematic error. Notably,

the topology of our complete phylogenetic tree (Fig. 2a) within the archaeal clade is mostly consistent

with the tree obtained in a preliminary analysis of a larger set of archaeal genomes and a larger marker

gene set (*26*). Taking into account these observations and the fact that we used the largest set of Asgard

archaea and other archaea compared to all previous phylogenetic analyses, the appearance of the 3D

topology in our tree indicates that the origin of eukaryotes from within Asgard cannot be considered a

settled issue. Various factors affecting the tree topology, including further increased species

representation, particularly, of Asgard and the deep branches of the TACK superphylum, such as

Bathyarchaeota and Korarchaeota, remain to be explored in order to definitively resolve the evolutionary

relationship between archaea and eukaryotes.

The core gene set of Asgard archaea

We next analyzed the core set of conserved Asgard genes which we arbitrarily defined as all asCOGs that

are present at least in one third of the MAGs in each of the 12 phylum-level lineages, with the mean

representation across lineages >75%. Under these criteria, the Asgard core includes 378 asCOGs

(Supplementary Table 6). As expected, most of these protein families, 293 (77%), are universal (present

in bacteria, other archaea and eukaryotes), 62 (16%) are represented in other archaea and eukaryotes, but

not in bacteria, 15 (4%) are found in other archaea and bacteria, but not in eukaryotes, 7 (2%) are archaea-

specific, and only 1 (0.003%) is shared exclusively with eukaryotes (Supplementary Figure 5). Most of

the core asCOGs show comparable levels of similarity to homologs from two or all three domains of life.

The second largest fraction of the core asCOGs shows substantially greater sequence similarity (at least,

25% higher similarity score) to homologous proteins from archaea than to those from eukaryotes and/or

bacteria (Supplementary Table 6). Compared with the 219 genes that comprise the pan-archaeal core (*27*),

the Asgard core set lacks 12 genes, each of which, however, is present in some subset of the Asgard

genomes. These include three genes of diphthamide biosynthesis and 2 ribosomal proteins, L40E and

L37E. The intricate evolutionary history of gene encoding translation elongation factors and enzymes of

diphthamide biosynthesis in Asgard has been analyzed previously (*28*). Also of note is the displacement

of the typical archaeal glyceraldehyde-3-phosphate dehydrogenase (type II) by a bacterial one (type I) in

most of the Asgard genomes (cog.001204, additional data file 1).

Functional distribution of the core asCOGs is shown in Fig. 3b (also see additional data file 1). For

comparison, we also derived an extended gene core for the TACK superphylum, using similar criteria (at

least 50% in each of the 6 lineages and 75% of the genomes overall, Fig 3b). For at least half of the

Asgard core genes, across most functional classes, there were no orthologs in the TACK core. The most

pronounced differences were found, as expected, in the category U (intracellular trafficking, secretion,

and vesicular transport). In Asgard archaea, this category includes 19 core genes compared with 7 genes

in TACK; 13 of these genes are specific to the Asgard archaea and include components of ESCRT I and

II, 3 distinct Roadblock/longin families, 2 distinct families of small GTPases, and a few other genes

implicated in related processes (Supplementary Table 6).

We compared the protein annotation obtained using asCOGs with the available annotation of '*Candidatus*

Prometheoarchaeum syntrophicum' and found that using asCOGs allowed at least a general functional

prediction for 649 of the 1756 (37%) 'hypothetical proteins' in this organism, the only one in Asgard with

a closed genome. We also identified 139 proteins, in addition to the 80 described originally, that can be

considered Eukaryotic Signature Proteins, or ESPs (see next section).

Eukaryotic features of Asgard archaea gleaned from genome analysis

The enrichment of Asgard proteomes with homologs of eukaryote signature proteins (ESPs), such as

ESCRTs, components of protein sorting complexes including coat proteins, complete ubiquitin

machinery, actins and actin-binding proteins gelsolins and profilins, might be the strongest argument in

support of a direct evolutionary relationship between Asgard archaea and eukaryotes (*2*, *29*). However,

the definition of ESPs is fuzzy because many of these proteins, in addition to their occurrence in Asgard,

are either scattered among several other archaeal genomes, often, from diverse groups (*30*), or consist of

promiscuous domains that are common in archaea, bacteria and eukaryotes, such as WD40 (after the

conserved terminal amino acids of the repeat units, also known as beta-transducin repeats), LRR

(Leucine-Rich Repeats), TPR (TetratricoPeptide Repeats), HEAT (Huntingtin-EF3-protein phosphatase

2A-TOR1) and other, largely, repetitive domains (*31*, *32*). Furthermore, the sequences of some ESPs have

diverged to the extent that they become hardly detectable with standard computational methods. Our

computational strategy for delineating an extensive yet robust ESP set is described under Materials and

Methods. The ESP set we identified contained 505 asCOGs, including 238 that were not closely similar

(E-value=10^{-10}, length coverage 75%) to those previously described by Zaremba-Niedzwiedzka et al.

(*2*)(Supplementary Table 7). In a general agreement with previous observations, the majority of these

ESPs, 329 of the 505, belonged to the 'Intracellular trafficking, secretion, and vesicular transport' (U)

functional class, followed by 'Posttranslational modification, protein turnover, chaperones' (O), with 101

asCOGs (Supplementary Table 7). Among the asCOGs in the U class, 130 were Roadblock/LC7

superfamily proteins, including longins, sybindin and profilins, and 94 were small GTPases of several

families, such as RagA-like, Arf-like and Rab-like ones, as discussed previously (*33*).

The phyletic patterns of ESP asCOGs in Asgard archaea are extremely patchy and largely lineage-specific

(Fig. 4), indicating that most of the proteins in this set are not uniformly conserved throughout Asgard

evolution, but rather, are prone to frequent HGT, gene losses and duplications. These evolutionary

processes are correlated in prokaryotes, resulting in the overall picture of highly dynamic evolution (*34*).

Even the most highly conserved ESP asCOG are missing in some Asgard lineages but show multiple

duplications in others (Fig. 4 and Supplementary Table 7). Surprisingly, many gaps in the ESPs

distribution were detected in the Heimdallarchaeota that include the suspected ancestors of eukaryotes.

Characteristically, many ESPs are multidomain proteins, with 37% assigned to more than one asCOG,

compared to 17% among non-ESP proteins (Supplementary Table 7). Some multidomain ESPs in Asgard

archaea have the same domain organizations as their homologs in eukaryotes, but these are a minority and

typically contain only two domains. Examples include the fusion of two EAP30/Vps37 domains (*35*), and

Vps23 and E2 domains (*35*) in ESCRT complexes, multiple Rag family GTPases, in which longin domain

is fused to the GTPase domain, and several others. By contrast, most of the domain architectures of the

multidomain ESP proteins were not detected in eukaryotes and often are found only in a narrow subset of

Asgard archaea, suggesting extensive domain shuffling during Asgard evolution (Fig. 5a). For example,

we identified many proteins containing a fusion of Vps28/Vps23 from ESCRT I complex (*35*) with C-

terminal domains of several homologous subunits of adaptin and COPI coatomer complexes (*36*, *37*), and

E3 UFM1-protein ligase 1, which is involved in the UFM1 ubiquitin pathway (*38*) (Fig. 5a). Generally, a

protein with such a combination of domains can be predicted to be involved in ubiquitin-dependent

membrane remodeling but, because its domain architecture is unique, the precise function cannot be

inferred.

The majority of the ESP genes of Asgard archaea do not belong to conserved genomic neighborhoods, but

several such putative operons were detected. Perhaps, the most notable one is the ESCRT neighborhood

which includes genes coding for subunits of ESCRT I, II and III, and often, components of the ubiquitin

system (*2*), suggesting an ancient link between the two systems that persists in eukaryotes (*35*). We

predicted another operon that is conserved in most Asgard archaea and consists of genes encoding a

LAMTOR1-like protein of the Roadblock superfamily, a Rab-like small GTPase, and a protein containing

the DENN (differentially expressed in normal and neoplastic cells) domain that so far has been identified

only in eukaryotes (Fig. 5b). Two proteins consisting of a DENN domain fused to longin are subunits of

the folliculin (FLCN) complex that is conserved in eukaryotes. The FLCN complex is the sensor of amino

acid starvation interacting with Rag GTPase and Ragulator lysosomal complex, and a key component of

the mTORC1 pathway, the central regulator of cell growth in eukaryotes (*39*). Some Heimdallarchaea

encode several proteins with the exact same domain organization as FLCN (Fig. 5b). Ragulator is a

complex that consists of 5 subunits, each containing the Roadblock domain. In Asgard archaea, however,

the GTPase present in the operon is from a family that is distinct from the Rag GTPases, which interact

with both FLCN and Ragulator complexes in eukaryotes, despite the fact that Rag family GTPases are

abundant in Asgards (*33*) (Supplementary Table 7). Nevertheless, this conserved module of Asgard

proteins is a strong candidate to function as a guanine nucleotide exchange factor for Rab and Rag

GTPases, analogously to the eukaryotic FLCN. In eukaryotes, the DENN domain is present in many

proteins with different domain architectures that interact with different partners and perform a variety of

functions (*40*, *41*). The Asgard archaea also encode other DENN domain proteins, and the respective

genes form expanded families of paralogs in Loki, Hel and Heimdall lineages, again, with domain

architectures distinct from those in eukaryotes (Fig. 5b) (*42*).

Prompted by the identification of a FLCN-like complex, we searched for other components of the

mTORC1 regulatory pathway in Asgard archaea. The GATOR1 complex that consists of three subunits,

Depdc5, Nprl2, and Nprl3, is another amino acid starvation sensor that is involved in this pathway in

eukaryotes (*43*). Nitrogen permease regulators 2 and 3 (NPRL2 and NPRL3) are homologous GATOR1

subunits that contain a longin domain and a small NPRL2-specific C-terminal domain (*43*). We identified

a protein family with this domain organization in most Thor MAGs and a few Loki MAGs. Several other

ESP asCOGs include proteins with high similarity to the longin domain of NPRL2. Additionally, we

identified many fusions of the NPRL2-like longin domain with various domains related to prokaryotic

two-component signal transduction system (Fig. 5c). Considering the absence of a homolog of

phosphatidylinositol 3-kinase, the catalytic domain of the mTOR protein, it seems likely that, in Asgard

archaea, the key growth regulation pathway remains centered at typical prokaryotic two-component signal

transduction systems whereas at least some of the regulators and sensors in this pathway are "eukaryotic".

The abundance of NPRL2-like longin domains in Asgard archaea implies that the link between this

domain and amino acid starvation regulation emerged at the onset of Asgard evolution if not earlier.

Diverse metabolic repertoires, ancestral metabolism of Asgard archaea, and syntrophic evolution

Examination of the distribution of the asCOGs among the 12 Asgard archaeal phyla showed that the

metabolic pathway repertoire was conserved among the MAGs of each phylum but differed between the

phyla (Fig. 3a). Three distinct lifestyles were predicted by the asCOG analysis for different major

branches of Asgard archaea, namely, anaerobic heterotrophy, facultative aerobic heterotrophy, and

chemolithotrophy (Fig. 6, Supplementary Figure 9). For the last Asgard archaeal common ancestor

(LAsCA), a mixotrophic life style, including both production and consumption of H_2, can be inferred

from parsimony considerations (Fig. 6, Supplementary Table 8; see Materials and Methods for further

details). Loki-, Thor-, Hermod-, Baldr- and Borrarchaeota encode all enzymes for the complete (archaeal)

Wood-Ljungdahl pathway (WLP) and are predicted to oxidize organic substrates, likely, by using the

reverse WLP, given the lack of enzymes for oxidation of inorganic compounds (e.g., hydrogen,

sulfur/sulfide and nitrogen/ammonia). The genomes of these five Asgard phyla encode homologues of

membrane-bound respiratory H_2-evolving Group 4 [NiFe] hydrogenase (Supplementary Figure 6) and/or

cytosolic cofactor-coupled bidirectional Group 3 [NiFe] hydrogenase (44) (Supplementary Figure 7).

Phylogenetic analysis of both group 3 and group 4 [NiFe] hydrogenases showed that Asgard archaea form

distinct clades well separated from the functionally characterized hydrogenases, hampering the prediction

of their specific functions in Asgard archaea. The functionally characterized group 4 [NiFe] hydrogenases

in the Thermococci are involved in the fermentation of organic substrates to H_2, acetate and carbon

dioxide (45, 46). The presence of group 3 [NiFe] hydrogenases suggests that these Asgard archaea cannot

use H_2 as an electron donor because they lack the enzyme complex coupling H_2 oxidation to membrane

potential generation. Thus, in these organisms, bifurcate electrons from H_2 are likely to be used to support

the fermentation of organic substrates exclusively (45–47).

Both Wukongarchaeota genomes (As_075 and As_085) encode a bona fide membrane-bound Group 1k

[NiFe] hydrogenase that could mediate hydrogenotrophic respiration using heterodisulfide as the terminal

electron acceptor (48, 49) (Fig. 6, Supplementary Figure 8, Supplementary Figure 10). The group 1k

[NiFe] hydrogenase is exclusively found in methanogens of the order Methanosarcinales (Euryarchaoeta)

(50), and it is the first discovery of the group 1 [NiFe] hydrogenase in the Asgard archaea.

Wukongarchaeota also encode all enzymes for a complete WLP and a putative ADP-dependent acetyl-

CoA synthetase for acetate synthesis. Unlike all other Asgard archaea, Wukongarchaeota lack genes for

citrate cycle and beta-oxidation. Thus, Wukongarchaeota appear to be obligate chemolithotrophic

acetogens. The genomes of Wukongarchaeota were discovered only in seawater of the euphotic zone of

the Yap trench (0 m and 125 m). Dissolved H_2 concentration is known to be the highest in surface

seawater, where the active microbial fermentation, compared to deep sea (51), could produce sufficient

amounts of hydrogen for the growth of Wukongarchaeota. Hodarchaeota, Gerdarchaeota, Kariarchaeota,

and Heimdallarchaeota share a common ancestor with Wukongarchaeota (Fig. 6). However, genome

analysis implies different lifestyles for these organisms. Hod-, Gerd- and Kariarchaeota encode various

electron transport chain components, including heme/copper-type cytochrome/quinol oxidase, nitrate

reductase, and NADH dehydrogenase, most likely, allowing the use of oxygen and nitrate as electron

acceptors during aerobic and anaerobic respiration, respectively (44). In addition, Hod-, Gerd- and

Heimdallarchaeota encode phosphoadenosine phosphosulfate (PAPS) reductase and adenylylsulfate

kinase for sulfate reduction, enabling the use of sulfate as electron acceptor during anaerobic respiration.

Gerd-, Heimdall-, and Hodarchaeota are only found in coastal and deep-sea sedimentary environments,

whereas Kariarchaeota were found also in marine water. The versatile predicted metabolic capacities of

these groups suggest that Hod-, Gerd- and Kariarchaeota might occupy both anoxic and oxic niches. In

contrast, Heimdallarchaeota appear to be able to thrive only in anoxic environments.

In the widely considered syntrophy scenarios (*52*), eukaryogenesis has been proposed to involve

metabolic symbiosis (syntrophy) between an archaeon and one or two bacterial partners which, in the

original hydrogen/syntrophy hypothesis, were postulated to donate H_2 for methane or hydrogen sulfide

production by the consortium (*53*, *54*). The syntrophic scenarios were boosted by the discovery of

apparent syntrophy between *Candidatus* P. syntrophicum and Deltaproteobacteria which led to the

proposal of the Etangle-Engulf-Endogenize (E^3) model of eukaryogenesis (*17*, *55*). Reconstruction of the

Lokiarchaeon metabolism has suggested that this organism was hydrogen-dependent, in accord with the

hydrogen-syntrophic scenarios (*54*). In contrast, subsequent analysis of the metabolic potentials of 4

Asgard phyla has led to the inference that these organisms were primarily organoheterotrophic and H_2-

producing, the 'reverse flow model' model of protoeukaryote energy metabolism that involves electron or

hydrogen flow from an Asgard archaeon to the alphaproteobacterial ancestor of mitochondria, in the

opposite direction from that in the original hydrogen-syntrophy hypotheses (*44*). Here, we discovered a

deeply branching Asgard group, Wukongarchaeota, which appears to include obligate hydrogenotrophic

acetogens, suggesting the possibility of the LAsCA being a hydrogen-dependent autotroph (Fig. 6). This

finding suggests that LAsCA both produced and consumed H_2. Thus, depending on the exact relationship

between Asgard archaea and eukaryotes that remains to be elucidated, our findings could be compatible

with different syntrophic scenarios that postulate H_2 transfer from bacteria to the archaeal symbiont or in

the opposite direction.

Conclusions

The Asgard archaea that were discovered only 5 years ago as a result of the painstaking assembly of

several Loki Castle metagenomes have grown into a highly diverse archaeal superphylum. The most

remarkable feature of the Asgards is their apparent evolutionary affinity with eukaryotes that has been

buttressed by two independent lines of evidence: phylogenetic analysis of highly conserved genes and

detection of multiple ESPs that are absent or far less common in other archaea. The 75 MAGs added here

substantially expand the phylogenetic and metabolic diversity of the Asgard superphylum. The extended

set of Asgard genomes provides for a phylogeny based on a far more representative species sampling than

available previously and a substantially expanded ESP analysis employing powerful computational

methods. This extended analysis reveals 6 putative additional Asgard phyla but does not immediately

clarify the Asgard-eukaryote relationship. Indeed, phylogenetic analysis of conserved genes in an

expanded set of archaea, bacteria and eukaryotes yields conflicting 3D and 2D signals, with an

unexpected preference for the 3D topology. Thus, the conclusion that eukaryotes emerged from within

Asgard archaea, in particular, from the Heimdall lineage, appears to be premature. Further phylogenomic

study with an even broader representation of diverse archaeal lineages as well as, possibly, even more

sophisticated evolutionary models are required to clarify the relationships between archaea and

eukaryotes.

Our analysis of Asgard genomes substantially expanded the set of ESPs encoded by this group of archaea

and revealed numerous, complex domain architectures of these proteins. These results further emphasize

the excess of ESPs in Asgards compared to other archaea and provide additional support to the conclusion

that most of the Asgard ESPs are involved in membrane remodeling and intracellular trafficking.

However, in parallel with the phylogenomic results, detailed analysis of the ESPs reveals a complex

picture. Most of the multidomain Asgard ESPs possess domain architectures distinct from typical

eukaryotic ones and some of these arrangements include signature prokaryotic domains, suggesting

substantial functional differences from the respective eukaryotic systems. Furthermore, virtually all the

ESPs show patchy distributions in Asgard and other archaea, indicative of a history of extensive HGT,

gene losses and paralogous family expansion. All these findings seem to be best compatible with the

model of a dispersed, dynamic archaeal 'eukaryome' (*30*) that widely spreads among archaea via HGT, so

far reaching the highest ESP density in the Asgard archaea.

The results of this work cannot rule out the possibility of the emergence of eukaryotes from within the

Asgard but seem to be better compatible with a different evolutionary scenario under which the conserved

core of eukaryote genes involved in informational processes originates from an as yet unknown ancestor

group that might be a deep archaeal branch or could lie outside the presently characterized archaeal

diversity. These hypothetical ancestral forms might have accumulated components of the mobile archaeal

'eukaryome' to an even greater extent than the Asgards archaea, eventually, giving rise to eukaryote-like

cells, likely, via a form of syntrophy with one or more bacterial partners. Combined genomic and

(undoubtedly, far more challenging) biological study of diverse archaea is essential for further advancing

our understanding of eukaryogenesis.

Author contributions

ML, EVK, KSM and YL initiated the study;; YL performed metagenomic assembly, binning, metabolism
analysis; KSM, AN and YIW performed comparative genomic analysis; YL, KSM, YIW and WCH
performed phylogenetic analysis; KSM and YIW constructed asCOGs; KSM, YIW, YL, ML, and EVK
analyzed the data; YL, KSM, WCH, EVK, and ML wrote the manuscript that was read, edited and
approved by all authors.

Acknowledgements

ML and YL are supported by National Natural Science Foundation of China (Grant No. 91851105,
31970105 and 31700430), the Key Project of Department of Education of Guangdong Province
(No.2017KZDXM071), and the Shenzhen Science and Technology Program (Grant no.
JCYJ20170818091727570 and KQTD20180412181334790). KSM, YIW, SAS and EVK are supported
by the Intramural Research Program of the National Institutes of Health of the USA (National Library of
Medicine).

Competing interests

The authors declare no competing interests

References

1.	A. Spang, J. H. Saw, S. L. Jørgensen, K. Zaremba-Niedzwiedzka, J. Martijn, A. E. Lind, R. van Eijk, C. Schleper, L. Guy, T. J. G. Ettema, Complex archaea that bridge the gap between prokaryotes and eukaryotes. *Nature*. **521**, 173–179 (2015).

2.	K. Zaremba-Niedzwiedzka, E. F. Caceres, J. H. Saw, D. B ckstr m, L. Juzokaite, E. Vancaester, K. W. Seitz, K. Anantharaman, P. Starnawski, K. U. Kjeldsen, M. B. Stott, T. Nunoura, J. F. Banfield, A. Schramm, B. J. Baker, A. Spang, T. J. G. Ettema, Asgard archaea illuminate the origin of eukaryotic cellular complexity. *Nature*. **541**, 353–358 (2017).

3.	F. MacLeod, G. S. Kindler, H. L. Wong, R. Chen, B. P. Burns, Asgard archaea: Diversity, function, and evolutionary implications in a range of microbiomes. *AIMS Microbiol*. **5**, 48–61 (2019).

4.	M. Cai, Y. Liu, X. Yin, Z. Zhou, M. W. Friedrich, T. Richter-Heitmann, R. Nimzyk, A. Kulkarni, X. Wang, W. Li, J. Pan, Y. Yang, J.-D. Gu, M. Li, Diverse Asgard archaea including the novel phylum Gerdarchaeota participate in organic matter degradation. *Sci. China Life Sci*. **63**, 886–897 (2020).

5.	T. A. Williams, C. J. Cox, P. G. Foster, G. J. Szöllősi, T. M. Embley, Phylogenomics provides robust support for a two-domains tree of life. *Nat Ecol Evol* (2019), doi:10.1038/s41559-019-1040-x.

6.	T. A. Williams, P. G. Foster, C. J. Cox, T. M. Embley, An archaeal origin of eukaryotes supports only two primary domains of life. *Nature*. **504**, 231–236 (2013).

7.	C. J. Cox, P. G. Foster, R. P. Hirt, S. R. Harris, T. M. Embley, The archaebacterial origin of eukaryotes. *Proc Natl Acad Sci U S A*. **105**, 20356–20361 (2008).

8.	N. Yutin, K. S. Makarova, S. L. Mekhedov, Y. I. Wolf, E. V. Koonin, The deep archaeal roots of eukaryotes. *Mol. Biol. Evol.* **25**, 1619–1630 (2008).

9.	V. Da Cunha, M. Gaia, D. Gadelle, A. Nasir, P. Forterre, Lokiarchaea are close relatives of Euryarchaeota, not bridging the gap between prokaryotes and eukaryotes. *PLoS Genet*. **13**, e1006810 (2017).

10.	V. D. Cunha, M. Gaia, A. Nasir, P. Forterre, Asgard archaea do not close the debate about the universal tree of life topology. *PLoS Genet*. **14**, e1007215 (2018).

11.	P. Forterre, The universal tree of life: an update. *Front. Microbiol*. **6**, 717 (2015).

12.	J. Lombard, P. López-García, D. Moreira, The early evolution of lipid membranes and the three domains of life. *Nat. Rev. Microbiol*. **10**, 507–515 (2012).

13.	J. A. Burns, A. A. Pittis, E. Kim, Gene-based predictive models of trophic modes suggest Asgard archaea are not phagocytotic. *Nat. Ecol. Evol*. **6**, a016006 (2018).

14.   C. Akıl, R. C. Robinson, Genomes of Asgard archaea encode profilins that regulate actin.
Nature. **562**, 439–443 (2018).

15.   C. Akıl, L. T. Tran, M. Orhant-Prioux, Y. Baskaran, E. Manser, L. Blanchoin, R. C.
Robinson, Insights into the evolution of regulated actin dynamics via characterization of
primitive gelsolin/cofilin proteins from Asgard archaea. *Proc Natl Acad Sci U S A*. **117**, 19904–
19913 (2020).

16.   Z. Lu, T. Fu, T. Li, Y. Liu, S. Zhang, J. Li, J. Dai, E. V. Koonin, G. Li, H. Chu, M. Li,
Coevolution of Eukaryote-like Vps4 and ESCRT-III Subunits in the Asgard Archaea. *mBio*. **11**,
e00417-20, /mbio/11/3/mBio.00417-20.atom (2020).

17.   H. Imachi, M. K. Nobu, N. Nakahara, Y. Morono, M. Ogawara, Y. Takaki, Y. Takano,
K. Uematsu, T. Ikuta, M. Ito, Y. Matsui, M. Miyazaki, K. Murata, Y. Saito, S. Sakai, C. Song, E.
Tasumi, Y. Yamanaka, T. Yamaguchi, Y. Kamagata, H. Tamaki, K. Takai, Isolation of an
archaeon at the prokaryote-eukaryote interface. *Nature*. **577**, 519–525 (2020).

18.   S. M. Karst, M. S. Dueholm, S. J. McIlroy, R. H. Kirkegaard, P. H. Nielsen, M.
Albertsen, *Nat. Biotechnol.*, in press, doi:10.1038/nbt.4045.

19.   R.-Y. Zhang, B. Zou, Y.-W. Yan, C. O. Jeon, M. Li, M. Cai, Z.-X. Quan, Design of
targeted primers based on 16S rRNA sequences in meta-transcriptomic datasets and
identification of a novel taxonomic group in the Asgard archaea. *BMC microbiology*. **20**, 25
(2020).

20.   K. S. Makarova, Y. I. Wolf, E. V. Koonin, Towards functional characterization of
archaeal genomic dark matter. *Biochem Soc Trans*. **47**, 389–398 (2019).

21.   E. V. Koonin, Y. I. Wolf, Genomics of bacteria and archaea: the emerging dynamic view
of the prokaryotic world. *Nucleic Acids Res*. **36**, 6688–6719 (2008).

22.   A. R. Nabhan, I. N. Sarkar, The impact of taxon sampling on phylogenetic inference: a
review of two decades of controversy. *Brief Bioinform*. **13**, 122–134 (2012).

23.   R. Gouy, D. Baurain, H. Philippe, Rooting the tree of life: the phylogenetic jury is still
out. *Philos Trans R Soc Lond B Biol Sci*. **370**, 20140329 (2015).

24.   J. Felsenstein, *Inferring Phylogenies* (Oxford University Press, Oxford, New York, 2nd
Edition., 2003).

25.   F. D. Ciccarelli, T. Doerks, C. von Mering, C. J. Creevey, B. Snel, P. Bork, Toward
automatic reconstruction of a highly resolved tree of life. *Science (New York, N.Y.)*. **311**, 1283–
1287 (2006).

26.   C. Rinke, M. Chuvochina, A. J. Mussig, P.-A. Chaumeil, D. W. Waite, W. B. Whitman,
D. H. Parks, P. Hugenholtz, *bioRxiv*, in press, doi:10.1101/2020.03.01.972265.

27.   K. S. Makarova, Y. I. Wolf, E. V. Koonin, Archaeal Clusters of Orthologous Genes
(arCOGs): An Update and Application for Analysis of Shared Features between
Thermococcales, Methanococcales, and Methanobacteriales. *Life (Basel)*. **5**, 818–840 (2015).

28.   A. B. Narrowe, A. Spang, C. W. Stairs, E. F. Caceres, B. J. Baker, C. S. Miller, T. J. G.
Ettema, Complex Evolutionary History of Translation Elongation Factor 2 and Diphthamide
Biosynthesis in Archaea and Parabasalids. *Genome Biol Evol*. **10**, 2380–2393 (2018).

29.   L. Eme, A. Spang, J. Lombard, C. W. Stairs, T. J. G. Ettema, Archaea and the origin of
eukaryotes. *Nat. Rev. Microbiol.* **15**, 711–723 (2017).

30.   E. V. Koonin, N. Yutin, The dispersed archaeal eukaryome and the complex archaeal
ancestor of eukaryotes. *Cold Spring Harb Perspect Biol*. **6**, a016188 (2014).

31.   R. L. Tatusov, N. D. Fedorova, J. D. Jackson, A. R. Jacobs, B. Kiryutin, E. V. Koonin, D.
M. Krylov, R. Mazumder, S. L. Mekhedov, A. N. Nikolskaya, B. S. Rao, S. Smirnov, A. V.
Sverdlov, S. Vasudevan, Y. I. Wolf, J. J. Yin, D. A. Natale, The COG database: an updated
version includes eukaryotes. *BMC Bioinformatics*. **4**, 41 (2003).

32.   E. V. Koonin, N. D. Fedorova, J. D. Jackson, A. R. Jacobs, D. M. Krylov, K. S.
Makarova, R. Mazumder, S. L. Mekhedov, A. N. Nikolskaya, B. S. Rao, I. B. Rogozin, S.
Smirnov, A. V. Sorokin, A. V. Sverdlov, S. Vasudevan, Y. I. Wolf, J. J. Yin, D. A. Natale, A
comprehensive evolutionary classification of proteins encoded in complete eukaryotic genomes.
Genome Biol. **5**, R7 (2004).

33.   C. M. Klinger, A. Spang, J. B. Dacks, T. J. G. Ettema, Tracing the Archaeal Origins of
Eukaryotic Membrane-Trafficking System Building Blocks. *Mol Biol Evol*. **33**, 1528–1541
(2016).

34.   P. Puigbò, A. E. Lobkovsky, D. M. Kristensen, Y. I. Wolf, E. V. Koonin, Genomes in
turmoil: quantification of genome dynamics in prokaryote supergenomes. *BMC Biol*. **12**, 66
(2014).

35.   L. Christ, C. Raiborg, E. M. Wenzel, C. Campsteijn, H. Stenmark, Cellular Functions and
Molecular Mechanisms of the ESCRT Membrane-Scission Machinery. *Trends Biochem Sci*. **42**,
42–56 (2017).

36.   N. Gomez-Navarro, E. Miller, Protein sorting at the ER-Golgi interface. *J Cell Biol*. **215**,
769–778 (2016).

37.   K. M. K. Kibria, J. Ferdous, R. Sardar, A. Panda, D. Gupta, A. Mohmmed, P. Malhotra,
A genome-wide analysis of coatomer protein (COP) subunits of apicomplexan parasites and their
evolutionary relationships. *BMC Genomics*. **20**, 98 (2019).

38.   Y. Wei, X. Xu, UFMylation: A Unique & Fashionable Modification for Life. *Genomics
Proteomics Bioinformatics*. **14**, 140–146 (2016).

39.   R. E. Lawrence, S. A. Fromm, Y. Fu, A. L. Yokom, D. J. Kim, A. M. Thelen, L. N.
Young, C.-Y. Lim, A. J. Samelson, J. H. Hurley, R. Zoncu, Structural mechanism of a Rag

GTPase activation checkpoint by the lysosomal folliculin complex. *Science*. **366**, 971–977 (2019).

40. M.-Y. Su, S. A. Fromm, R. Zoncu, J. H. Hurley, Structure of the C9orf72 ARF GAP complex that is haploinsufficient in ALS and FTD. *Nature*. **585**, 251–255 (2020).

41. N. de Martín Garrido, C. H. S. Aylett, Nutrient Signaling and Lysosome Positioning Crosstalk Through a Multifunctional Protein, Folliculin. *Front Cell Dev Biol*. **8**, 108 (2020).

42. A. L. Marat, H. Dokainish, P. S. McPherson, DENN domain proteins: regulators of Rab GTPases. *J Biol Chem*. **286**, 13791–13800 (2011).

43. K. Shen, R. K. Huang, E. J. Brignole, K. J. Condon, M. L. Valenstein, L. Chantranupong, A. Bomaliyamu, A. Choe, C. Hong, Z. Yu, D. M. Sabatini, Architecture of the human GATOR1 and GATOR1-Rag GTPases complexes. *Nature*. **556**, 64–69 (2018).

44. A. Spang, C. W. Stairs, N. Dombrowski, L. Eme, J. Lombard, E. F. Caceres, C. Greening, B. J. Baker, T. J. G. Ettema, Proposal of the reverse flow model for the origin of the eukaryotic cell based on comparative analyses of Asgard archaeal metabolism. *Nat. Microbiol*. **4**, 1138–1148 (2019).

45. H. Yu, C.-H. Wu, G. J. Schut, D. K. Haja, G. Zhao, J. W. Peters, M. W. W. Adams, H. Li, Structure of an Ancient Respiratory System. *Cell*. **173**, 1636-1649.e16 (2018).

46. G. J. Schut, E. S. Boyd, J. W. Peters, M. W. W. Adams, The modular respiratory complexes involved in hydrogen and sulfur metabolism by heterotrophic hyperthermophilic archaea and their evolutionary implications. *FEMS Microbiol Rev*. **37**, 182–203 (2013).

47. F. O. Bryant, M. W. Adams, Characterization of hydrogenase from the hyperthermophilic archaebacterium, Pyrococcus furiosus. *J Biol Chem*. **264**, 5070–5079 (1989).

48. U. Deppenmeier, M. Blaut, B. Schmidt, G. Gottschalk, Purification and properties of a F420-nonreactive, membrane-bound hydrogenase from Methanosarcina strain Gö1. *Arch Microbiol*. **157**, 505–511 (1992).

49. R. K. Thauer, A.-K. Kaster, M. Goenrich, M. Schick, T. Hiromoto, S. Shima, Hydrogenases from methanogenic archaea, nickel, a novel cofactor, and H2 storage. *Annu Rev Biochem*. **79**, 507–536 (2010).

50. D. Søndergaard, C. N. S. Pedersen, C. Greening, HydDB: A web tool for hydrogenase classification and analysis. *Scientific Reports*. **6**, 34212 (2016).

51. R. Conrad, W. Seiler, Methane and hydrogen in seawater (Atlantic Ocean). *Deep Sea Res. Part I Oceanogr. Res. Pap*. **35**, 1903–1917 (1988).

52. P. López-García, D. Moreira, The Syntrophy hypothesis for the origin of eukaryotes revisited. *Nat Microbiol*. **5**, 655–667 (2020).

53. W. Martin, M. Müller, The hydrogen hypothesis for the first eukaryote. *Nature*. **392**, 37–41 (1998).

54.   D. Moreira, P. López-García, Symbiosis Between Methanogenic Archaea and δ-
Proteobacteria as the Origin of Eukaryotes: The Syntrophic Hypothesis. *J Mol Evol*. **47**, 517–530
(1998).

55.   P. López-García, D. Moreira, Cultured Asgard Archaea Shed Light on Eukaryogenesis.
Cell. **181**, 232–235 (2020).

.

Figures

Figure 1. Gene commonality plots for Asgard archaea and the TACK superphylum.

The gene commonality plot showing the number of asCOGs in log scale (Y-axis) that include the given
fraction of analyzed genomes (X-axis). The Asgard plot is compared with the TACK superphylum plot
based on assignment of TACK genomes to arCOGs.

Figure 2. Phylogenetic analysis of Asgard archaea and their relationships with eukaryotes.

(a) Maximum likelihood tree, inferred with IQ-tree and LG+F+R10 model, constructed from
concatenated alignments of the protein sequences from 209 core Asgard Clusters of Orthologs (asCOGs).
Only the 12 phylum-level clades are shown, with species within each clade collapsed. See supplementary
Methods and Supplementary Table 5 for details.

(b) Maximum likelihood tree, inferred with IQ-tree and SYM+R8 model, based on 16S rRNA gene
sequences. Red stars in **(b)** denote MAGs reconstructed in the current study.

(**c**) Phylogenetic tree of bacteria, archaea and eukaryotes, inferred with IQ-tree under LG+R10 model,
constructed from concatenated alignments of the protein sequences of 30 universally conserved genes
(see Material and Methods for details). The tree shows the relationships between the major clades.

The trees are unrooted and are shown in a pseudorooted form for visualization purposes only. The actual
trees and alignments are in Additional data file 2 and list of the trees are provided in the Supplementary
Table 4 and 5.

Figure 3. Phyletic patterns of asCOGs and functional distribution of Asgard core genes.

(a) Classical Multidimensional Scaling analysis of binary presence-absence phyletic patterns for 13,939
asCOGs that are represented in at least two genomes (see Material and Methods for details).

(b) Functional breakdown of Asgard core genes (378 asCOGs) compared with TACK superphylum core
genes (489 arCOGs). The values were normalized as described in Materials and Methods. Functional
classes of genes: J, Translation, ribosomal structure and biogenesis; K, Transcription; L, Replication,

recombination and repair; D, Cell cycle control, cell division, chromosome partitioning; V, Defense

mechanisms; T, Signal transduction mechanisms; M, Cell wall/membrane/envelope biogenesis; N, Cell

motility; U, Intracellular trafficking, secretion, and vesicular transport; O, Posttranslational modification,

protein turnover, chaperones; X, Mobilome: prophages, transposons; C, Energy production and

conversion; G, Carbohydrate transport and metabolism; E, Amino acid transport and metabolism; F,

Nucleotide transport and metabolism; H, Coenzyme transport and metabolism; I, Lipid transport and

metabolism; P, Inorganic ion transport and metabolism; Q, Secondary metabolites biosynthesis, transport

and catabolism; R, General function prediction only; S, Function unknown;

Figure 4. Phyletic patterns of Eukaryotic Signature Proteins (ESPs) encoded in Asgard genomes.

All 505 ESP asCOGs are grouped by distance between binary presence-absence phyletic patterns. The

most highly conserved ESP asCOGs are shown within the red rectangle. Below the plot of the number of

ESP domains in each genome is shown. For details, see Supplementary Table 7.

Figure 5. Domain architectures of selected Asgard ESPs.

(a) ESPs with unique domain architectures. The schematic of each multidomain protein is roughly

proportional to the respective protein length. The identified domains are shown inside the arrows

approximately according to their location and are briefly annotated. Homologous domains are shown by

the same color or pattern.

(b) DENN domain proteins in Asgards. Upper part of the figure above the dashed line shows putative

operon encoding DENN domain proteins. Genes are shown by block arrows with the length proportional

to the size of the corresponding protein. For each protein, the nucleotide contig or genome partition

accession number, Asgard genome ID and lineage are indicated. The part of the figure below the dashed

line shows domain organization of diverse proteins containing DENN domain. Homologous domains are

shown by the same color. The inset on the right side explains identity of the domains

(c) NPRL2-like proteins in Asgards. Designations are the same as in Figure 4a.

Figure 6. Reconstruction and evolution of the key metabolic processes in Asgard archaea.

The schematic phylogeny of Asgard archaea is from Figure 1a.

LAsCA, Last Asgard Common Ancestor; WLP, Wood-Ljungdahl pathway.

Supplementary Materials

Taxonomic description

Materials and Methods

Supplementary References

Tables S1-S8

Figures S1-S10

Additional Data files are available at ftp://ftp.ncbi.nih.gov/pub/wolf/_suppl/asgard20/

Additional data file 1 (Additional_data_file_1.tgz): Complete asCOG data archive

Additional data file 2 (Additional_data_file_2.tgz): Phylogenetic trees and alignments archive

a

b

c

A

B

378
asCOGs

■ Asgard only   ■ Shared   ■ TACK only

A

B

C

Expanding diversity of Asgard archaea and the elusive ancestry of eukaryotes

Yang Liu[1†], Kira S. Makarova[2†], Wen-Cong Huang[1†], Yuri I. Wolf[2], Anastasia Nikolskaya[2], Xinxu

Zhang[1], Mingwei Cai[1], Cui-Jing Zhang[1], Wei Xu[3], Zhuhua Luo[3], Lei Cheng[4], Eugene V. Koonin[2*], Meng

Li[1*]

1 Shenzhen Key Laboratory of Marine Microbiome Engineering, Institute for Advanced Study, Shenzhen

University, Shenzhen, Guangdong, 518060, P. R. China

2 National Center for Biotechnology Information, National Library of Medicine, National Institutes of

Health, Bethesda, Maryland 20894, USA

3 State Key Laboratory Breeding Base of Marine Genetic Resources, Key Laboratory of Marine Genetic

Resources, Fujian Key Laboratory of Marine Genetic Resources, Third Institute of Oceanography, State

Oceanic Administration, Xiamen 361005, P. R. China

4 Key Laboratory of Development and Application of Rural Renewable Energy, Biogas Institute of

Ministry of Agriculture, Chengdu 610041, P.R. China

† These authors contributed equally to this work.

*Authors for correspondence: koonin@ncbi.nlm.nih.gov or limeng848@szu.edu.cn

Supplementary Information

Taxonomic Description of new taxa

'*Candidatus* **Wukongarchaeum**' (Wu.kong.ar.chae'um. N. L. n. Wukong a legendary Chinese figure, also known as Monkey King, who caused havoc in the heavenly palace); N.L. neut. N. *archaeum* (from Gr. adj. archaios ancient) archaeon; N. L. neut. N. Wukongarchaeum.

'*Candidatus* **Wukongarchaeum yapensis**' (yap'ensis N. L. masc. adj. pertaining to Yap trench, which is the geographical position where the first type material of this species was obtained). Type material is the genome designated as As_085 (Yap4.bin4.70) representing '*Candidatus* Wukongarchaeum yapensis'. The genome "As_085" represents a MAG consisting of 2.16 Mbps in 277 contigs with an estimated completeness of 92.52%, an estimated contamination of 4.05%, a 16S and 23S rRNA gene and 14 tRNAs. The MAG recovered from a marine water metagenome (Yap trench, Western Pacific), with an estimated depth of coverage of 31.4, has a GC content of 38%.

'*Candidatus* **Hodarchaeum**' (Hod.ar.chae'um. N. L. n. Hod a son of Odin in Norse mythology); N.L. neut. N. archaeum (from Gr. adj. archaios ancient) archaeon; N. L. neut. N. Hodarchaeum.

'*Candidatus* **Hodarchaeum mangrovi**' (man.gro'vi N.L. fem. n. of a mangrove, referring to the isolation of the type material from mangrove soil). Type material is the genome designated as As_027 (FT2_5_011) representing '*Candidatus* Hodarchaeum mangrovi'. The genome "As_027" represents a MAG consisting of 4.01 Mbps in 348 contigs with an estimated completeness of 93.61%, an estimated contamination of 0.93%, a 23S rRNA gene and 14 tRNAs. The MAG recovered from mangrove sediment metagenomes (Futian Nature Reserve, China), with an estimated depth of coverage of 17.9, has a GC content of 32.9%.

'*Candidatus* **Kariarchaeum**' (Ka.ri.ar.chae'um. N. L. n. Kari the god of wind and brother to Aegir in Norse mythology); N.L. neut. N. *archaeum* (from Gr. adj. archaios ancient) archaeon; N. L. neut. N. Kariarchaeum.

'*Candidatus* **Kariarchaeum pelagius**' (pe.la'gi.us. L. masc. adj. of or belonging to the sea, referring to the isolation of the type material from the Ocean). Type material is the genome designated as As_030 (RS678) representing '*Candidatus* Kariarchaeum pelagius'. The genome "As_030" represents a MAG consisting of 1.41 Mbps in 76 contigs, an estimated completeness of 83.18%, with an estimated contamination of 1.87%, a 23S, 16S and 5S rRNA genes and 18 tRNAs. The MAG recovered from a marine metagenome (Saudi Arabia: Red Sea) has a GC content of 30.11%.

'*Candidatus* **Borrarchaeum**' (Borr.ar.chae'um. N. L. n. Borr a creator god and father of Odin); N.L. neut. N. *archaeum* (from Gr. adj. archaios ancient) archaeon; N. L. neut. N. Borrarchaeum.

'*Candidatus* **Borrarchaeum yapensis**' (yap'ensis N. L. masc. adj. pertaining to Yap trench, which is the geographical position where the first type material of this species was obtained). Type material is the genome designated as As_181 (Yap2000.bin9.141) representing '*Candidatus* Borrarchaeum yapensis'. The genome "As_181" represents a MAG consisting of 3.63 Mbps in 125 contigs, with an estimated completeness of 95.02%, an estimated contamination of 5.61% and 11 tRNAs. The MAG, recovered from a marine water metagenome (Yap trench, Western Pacific) with an estimated depth coverage of 15.04, has a GC content of 37.1%.

'*Candidatus* Baldrarchaeum' (Bal.dr.ar.chae'um. N. L. n. Baldr the god of light and son of Odin and
brorther of Thor in Norse mythology); N.L. neut. N. *archaeum* (from Gr. adj. archaios ancient) archaeon;
N. L. neut. N. Baldrarchaeum.

'*Candidatus* Baldrarchaeum yapensis' (yap'ensis N. L. masc. adj. pertaining to Yap trench, which is the
geographical position where the first type material of this species was obtained). Type material is the
genome designated as As_130 (Yap30.bin9.72) representing '*Candidatus* Baldrarchaeum yapensis'. The
genome "As_130" represents a MAG consisting of 2.27 Mbps in 100 contigs, with an estimated
completeness of 93.93%, an estimated contamination of 3.74%, a 23S and 16S rRNA gene and 15 tRNAs.
The MAG, recovered from a marine water metagenome (Yap trench, Western Pacific) with an estimated
depth coverage of 39.99, has a GC content of 45.95%.

'*Candidatus* Hermodarchaeum' (Her.mod.ar.chae'um. N. L. n. Hermod, messengers of the gods in the
Norse mythology and son of Odin and brother of Baldr in the Norse mythology); N.L. neut. N. *archaeum*
(from Gr. adj. archaios ancient) archaeon; N. L. neut. N. Hermodarchaeum.

'*Candidatus* Hermodarchaeum yapensis' (yap'ensis N. L. masc. adj. pertaining to Yap trench, which is
the geographical position where the first type material of this species was obtained). Type material is the
genome designated as As_086 (Yap4.bin9.105) representing '*Candidatus* Hermodarchaeum yapensis'.
The genome 'As_086' represent a MAG consisting of 2.71 Mbps in 77 contigs, with an estimated
completeness of 92.99%, an estimated contamination of 1.87%, a 23S and 16S rRNA gene and 16 tRNAs.
The MAG, recovered from a marine water metagenome (Yap trench, Western Pacific) with an estimated
depth coverage of 19.24, has a GC content of 44.69%.

Candidatus **Wukongarchaeaceae** (Wu.kong.ar.chae.a.ce'ae. N.L. neut. n. Wukongarchaeum a
(Candidatus) type genus of the family; -aceae ending to denote the family; N.L. fem. pl. n.
Wukongarchaeaceae the Wukongarchaeum family).

The family is delineated based on 209 concatenated Asgard Cluster of Orthologs (AsCOGs) and 16S
rRNA gene phylogeny. The description is the same as that of its sole genus and species. Type genus is
Candidatus Wukongarchaeum.

Candidatus **Wukongarchaeales** (Wu.kong.ar.chae.a'les. N.L. neut. n. Wukongarchaeum a (Candidatus)
type genus of the order; -ales ending to denote the order; N.L fem. pl. n. Wukongarchaeales the
Wukongarchaeum order).

The order is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is Candidatus Wukongarchaeum.

Candidatus **Wukongarchaeia** (Wu.kong.ar.chae'i.a. N.L. neut. n. Wukongarchaeum a (Candidatus) type
genus of the order of the class; -ia ending to denote the class; N.L fem. pl. n. Wukongarchaeia the
Wukongarchaeum class).

The class is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type order *Candidatus* Wukongarchaeales.

Candidatus **Wukongarchaeota** (Wu.kong.ar.chae.o'ta. N.L. neut. n. Wukongarchaeum a (Candidatus)
type genus of the class of the phylum; -ota ending to denote the phylum; N.L neut. pl. n.
Wukongarchaeota the Wukongarchaeum phylum)

The phylum is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type class *Candidatus* Wukongarchaeia.

Candidatus **Hodarchaeaceae** (Hod.ar.chae.a.ce'ae. N.L. neut. n. Hodarchaeum a (Candidatus) type
genus of the family; -aceae ending to denote the family; N.L. fem. pl. n. Hodarchaeaceae the
Hodarchaeum family).

The family is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is *Candidatus* Hodarchaeum.

Candidatus **Hodarchaeales** (Hod.ar.chae.a'les. N.L. neut. n. Hodarchaeum a (Candidatus) type genus of
the order; -ales ending to denote the order; N.L fem. pl. n. Hodarchaeales the Hodarchaeum order).

The order is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is *Candidatus* Hodarchaeum.

Candidatus **Hodarchaeia** (Hod.ar.chae'i.a. N.L. neut. n. Hodarchaeum a (Candidatus) type genus of the
order of the class; -ia ending to denote the class; N.L fem. pl. n. Hodarchaeia the Hodarchaeum class).

The class is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type order *Candidatus* Hodarchaeales.

Candidatus **Hodarchaeota** (Hod.ar.chae.o'ta. N.L. neut. n. Hodarchaeum a (Candidatus) type genus of
the class of the phylum; -ota ending to denote the phylum; N.L neut. pl. n. Hodarchaeota the
Hodarchaeum phylum)

The phylum is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type class *Candidatus* Hodarchaeia.

Candidatus **Hodarchaeaceae** (Hod.ar.chae.a.ce'ae. N.L. neut. n. Hodarchaeum a (Candidatus) type
genus of the family; -aceae ending to denote the family; N.L. fem. pl. n. Hodarchaeaceae the
Hodarchaeum family).

The family is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is *Candidatus* Hodarchaeum.

Candidatus **Hodarchaeales** (Hod.ar.chae.a'les. N.L. neut. n. Hodarchaeum a (Candidatus) type genus of
the order; -ales ending to denote the order; N.L fem. pl. n. Hodarchaeales the Hodarchaeum order).

The order is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is *Candidatus* Hodarchaeum.

Candidatus **Hodarchaeia** (Hod.ar.chae'i.a. N.L. neut. n. Hodarchaeum a (Candidatus) type genus of the
order of the class; -ia ending to denote the class; N.L fem. pl. n. Hodarchaeia the Hodarchaeum class).

The class is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type order *Candidatus* Hodarchaeales.

Candidatus **Hodarchaeota** (Hod.ar.chae.o'ta. N.L. neut. n. Hodarchaeum a (Candidatus) type genus of
the class of the phylum; -ota ending to denote the phylum; N.L neut. pl. n. Hodarchaeota the
Hodarchaeum phylum)

The phylum is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type class *Candidatus* Hodarchaeia.

Candidatus **Kariarchaeaceae** (Ka.ri.ar.chae.a.ce'ae. N.L. neut. n. Kariarchaeum a (Candidatus) type
genus of the family; -aceae ending to denote the family; N.L. fem. pl. n. Kariarchaeaceae the
Kariarchaeum family).

The family is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is *Candidatus* Kariarchaeum.

Candidatus **Kariarchaeales** (Ka.ri.ar.chae.a'les. N.L. neut. n. Kariarchaeum a (Candidatus) type genus of
the order; -ales ending to denote the order; N.L fem. pl. n. Kariarchaeales the Kariarchaeum order).

The order is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is *Candidatus* Kariarchaeum.

Candidatus **Kariarchaeia** (Ka.ri.ar.chae'i.a. N.L. neut. n. Kariarchaeum a (Candidatus) type genus of the
order of the class; -ia ending to denote the class; N.L fem. pl. n. Kariarchaeia the Kariarchaeum class).

The class is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type order *Candidatus* Kariarchaeales.

Candidatus **Kariarchaeota** (Ka.ri.ar.chae.o'ta. N.L. neut. n. Kariarchaeum a (Candidatus) type genus of
the class of the phylum; -ota ending to denote the phylum; N.L neut. pl. n. Kariarchaeota the
Kariarchaeum phylum)

The phylum is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type class *Candidatus* Kariarchaeia.

Candidatus **Borrarchaeaceae** (Borr.ar.chae.a.ce'ae. N.L. neut. n. Borrarchaeum a (Candidatus) type
genus of the family; -aceae ending to denote the family; N.L. fem. pl. n. Borrarchaeaceae the
Borrarchaeum family).

The family is delineated based on 209 concatenated AsCOGs phylogeny. The description is the same as
that of its sole genus and species. Type genus is *Candidatus* Borrarchaeum.

Candidatus **Borrarchaeales** (Borr.ar.chae.a'les. N.L. neut. n. Borrarchaeum a (Candidatus) type genus of
the order; -ales ending to denote the order; N.L fem. pl. n. Borrarchaeales the Borrarchaeum order).

The order is delineated based on 209 concatenated AsCOGs phylogeny. The description is the same as
that of its sole genus and species. Type genus is *Candidatus* Borrarchaeum.

Candidatus **Borrarchaeia** (Borr.ar.chae'i.a. N.L. neut. n. Borrarchaeum a (Candidatus) type genus of the
order of the class; -ia ending to denote the class; N.L fem. pl. n. Borrarchaeia the Borrarchaeum class).

The class is delineated based on 209 concatenated AsCOGs phylogeny. The description is the same as
that of its sole and type order *Candidatus* Borrarchaeales.

Candidatus **Borrarchaeota** (Borr.ar.chae.o'ta. N.L. neut. n. Borrarchaeum a (Candidatus) type genus of
the class of the phylum; -ota ending to denote the phylum; N.L neut. pl. n. Borrarchaeota the
Borrarchaeum phylum)

The phylum is delineated based on 209 concatenated AsCOGs phylogeny. The description is the same as
that of its sole and type class *Candidatus* Borrarchaeia.

Candidatus **Baldrarchaeaceae** (Bal.dr.ar.chae.a.ce'ae. N.L. neut. n. Baldrarchaeum a (Candidatus) type
genus of the family; -aceae ending to denote the family; N.L. fem. pl. n. Baldrarchaeaceae the
Baldrarchaeum family).

The family is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is *Candidatus* Baldrarchaeum.

Candidatus **Baldrarchaeales** (Bal.dr.ar.chae.a'les. N.L. neut. n. Bladrarchaeum a (Candidatus) type
genus of the order; -ales ending to denote the order; N.L fem. pl. n. Baldrarchaeales the Baldrarchaeum
order).

The order is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is *Candidatus* Baldrarchaeum.

Candidatus **Baldrarchaeia** (Bal.dr.ar.chae'i.a. N.L. neut. n. Baldrarchaeum a (Candidatus) type genus of
the order of the class; -ia ending to denote the class; N.L fem. pl. n. Baldrarchaeia the Baldrarchaeum
class).

The class is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type order *Candidatus* Baldrarchaeales.

Candidatus **Baldrarchaeota** (Bal.dr.ar.chae.o'ta. N.L. neut. n. Baldrarchaeum a (Candidatus) type genus
of the class of the phylum; -ota ending to denote the phylum; N.L neut. pl. n. Baldrarchaeota the
Baldrarchaeum phylum)

The phylum is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type class *Candidatus* Baldrarchaeia.

Candidatus **Hermodarchaeaceae** (Her.mod.ar.chae.a.ce'ae. N.L. neut. n. Hermodarchaeum a
(Candidatus) type genus of the family; -aceae ending to denote the family; N.L. fem. pl. n.
Hermodarchaeaceae the Hermodarchaeum family).

The family is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is *Candidatus* Hermodarchaeum.

Candidatus **Hermodarchaeales** (Her.mod.ar.chae.a'les. N.L. neut. n. Hermodarchaeum a (Candidatus)
type genus of the order; -ales ending to denote the order; N.L fem. pl. n. Hermodarchaeales the
Hermodarchaeum order).

The order is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole genus and species. Type genus is *Candidatus* Hermodarchaeum.

Candidatus **Hermodarchaeia** (Her.mod.ar.chae'i.a. N.L. neut. n. Hermodarchaeum a (Candidatus) type
genus of the order of the class; -ia ending to denote the class; N.L fem. pl. n. Hermodarchaeia the
Hermodarchaeum class).

The class is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type order *Candidatus* Hermodarchaeales.

Candidatus **Hermodarchaeota** (Her.mod.ar.chae.o'ta. N.L. neut. n. Hermodarchaeum a (Candidatus)
type genus of the class of the phylum; -ota ending to denote the phylum; N.L neut. pl. n.
Hermodarchaeota the Hermodarchaeum phylum)

The phylum is delineated based on 209 concatenated AsCOGs and 16S rRNA gene phylogeny. The
description is the same as that of its sole and type class *Candidatus* Hermodarchaeia.

Materials and Methods

Sampling collections and DNA sequencing

YT samples were obtained from the Rongcheng Nation Swan Nature Reserve (Rongcheng, China) in November 2018. The sediment cores were collected using columnar samplers at depth intervals of 0–2, 21–26, and 36–41 cm at a seagrass meadow and a non-seagrass–covered site nearby. After collection, bulk sediments were immediately sealed in plastic bags, placed in a pre-cooled icebox, and transported to laboratory within 4 hours. For each sample, DNA was extracted from 10 g sediment using PowerSoil DNA Isolation kit (QIAGEN, Germany), according to the manufacturer's protocol. Following extraction, nucleic acids were sequenced using Illumina HiSeq2500 (Illumina, USA) PE150 by Novogene (Nanjing, China).

MP5 samples were obtained from Mai Po Nature Reserve (Hong Kong, China) in September 2014. Three subsurface sediment samples were collected from a site covering with mangrove forest at depth intervals of 0-2, 10-15 and 20-25 cm. Two subsurface sediment samples were taken at an intertidal mudflat with depths of 0-5 and 13-16 cm. Samples were transported back to laboratory as described for YT metagenomes. DNA was extracted from 5g wet sediment per sample using the PowerSoil DNA Isolation Kit (MO BIO, USA) following the manufacturer's protocol. Metagenomic sequencing data were generated using Illumina HiSeq2500 (Illumina, USA) PE150 by Novogene (Tianjin, China).

FT samples were taken from Futian Nature Reserve (Shenzhen, Guangdong, China) in April 2017. Sediment samples were collected as described for YT samples at depth intervals of 0-2, 6-8, 12-14, 20-22, and 28-30 cm. DNA was extracted from 5g wet sediment per sample using DNeasy PowerSoil kit (Qiagen, Germany) as per manufacturer's instructions. Nucleic acids were sequenced using Illumina HiSeq2000 (Illumina, USA) PE150 by Novogene (Tianjin, China).

The surface sediment sample of the CJE metagenome was collected from Changjiang estuary during a cruise in August 2016. The sample was grabbed from the water bed, sealed immediately in a 50 ml tubes and stored in liquid nitrogen onboard. After transportation to laboratory, 10g wet sediments were used for DNA extraction as per manufacturer's protocol. Nucleic acids were sequenced using Illumina HiSeq2000 (Illumina, USA) PE150 by Novogene (Tianjin, China).

An oil sand sample was collected from Shengli Oilfield (Shandong, China) into bottles and they were transported to laboratory where they were stored at 4 ℃. The sample was used as inoculum to perform enrichment with anaerobic medium in vials as described elsewhere (*1*). After 253d of enrichment, the genomic DNA was extracted as described elsewhere (*2*). Nucleic acids were sequenced Illumina HiSeq2000 (Illumina, USA) PE150 by Novogene (Tianjin, China).

The seawater samples of Yap metagenomes were collected at Yap trench region by CTD SBE911plus (Sea-Bird Electronics, USA) during the 37th Dayang cruise in 2016. 8L of seawater per sample was filtered through a 0.22 μm-mesh membrane filter immediately after recovery onboard. The membrane was then cut into ~0.2 cm^2 pieces with a flame-sterilized scissors and added to a PowerBead Tube (MO BIO, USA) and the subsequent steps were implemented according to the manufacturer's protocol to extract DNA. The DNA per sample was amplified in five separative reactions using REPLI-g Single Cell Kits (Qiagen, Germany) following the manufacturer's protocol, given to the challenging nature of sample retrieval and DNA recovery. The products were pooled together and purified using QIAamp DNA Mini

Kit (Qiagen, Germany) according to the manufacturer's recommendations. Parallel blank controls were
set for sampling, DNA extraction and ampliation with 0.22 µm-mesh membrane filtering Milli-Q water
(18.2 MΩ; Millipore, USA). Nucleic acids were sequenced using HiSeq X Ten (Illumina, USA) PE150.

Metagenomic assembly, binning and gene calling

FT, MP5 and YT metagenomes. These three sets of metagenomes were assembled and binned using the
same method. Raw shotgun metagenomic sequencing reads were trimmed with "read_qc" module from
metaWRAP (v.1.1) (*3*). All clean reads from the same set were pooled together prior to *de novo* assemble
to one co-assembly. Clean reads were sent out to MEGAHIT (v1.1.2) with flag "--presets meta-large" for
co-assembling job (*4*). Sequencing coverage was determined for each assembled scaffold by mapping
reads from each sample to the co-assembly using Bowtie2 (*5*). The binning analysis was carried out 8
times with 8 different combinations of specificity and sensitivity parameters using MetaBAT2(52) ("--
maxP 60 or 95" AND "--minS 60 or 95" AND "--maxEdges 200 or 500") on the assembly with a
minimum length of 2000 bp (*6*). DAS Tool (v1.0) was used as a dereplication and aggregation strategy on
those eight binning results to construct accurate bins (*7*). Manual curation was used for reducing the
genome contamination based on differential coverages, GC contents, and the presence of duplicate genes.

The depth coverage and N50 statistics of 38 Asgard MAGs recovered from YT metagenomes range from
7.72 to 298.86 (median: 21.77) and from 6658 to 381755 bps (median: 26337.5 bps) respectively; for 13
Asgard MAGs recovered from FT metagenomes, the values range from 6.98 to 33.22 (median 16.89) and
from 3889 to 8957 bps (median: 5000 bps), respectively; and for 11 Asgard MAGs recovered from MP5
metagenomes, the values range from 7.06 to 54.42 (median: 10) and from 3898 to 19362 bps (median:
7581 bps) respectively. (**Supplementary Figure 2**).

CJE metagenome. Raw metagenomic shotgun sequencing reads were trimmed using Sickle
(https://github.com/najoshi/sickle) with default settings. The trimmed reads were de novo assembled
using IDBA-UD (v 1.1.1) with the parameters: "-mink 65, -maxk 145, -step 10" (*8*). Sequencing coverage
was determined as described above. The binning analysis were performed with MetaBAT2 12 times, with
12 combinations of specificity and sensitivity parameters (--m 1500, 2000, or 2500" AND "--maxP 85 or
90" AND "--minS 80, 85 and 90") for further refinement (*6*). All binning results were merged and refined
using DAS Tool (v1.0) (*7*).

2 Asgard MAGs recovered from CJE metagenome have a depth coverage of 19.16 and 20.56, and a N50
statistics of 8740 and 5246 bps (**Supplementary Figure 2**).

J65 metagenome. Raw metagenomic shotgun sequencing reads were trimmed with Trimmomatic (v0.38)
(*9*). The clean reads were then fed to SPAdes (v 3.12.0) for *de novo* assembly with the parameters: "--
meta -k 21, 33, 55, 77" (*10*). Sequencing coverage was determined using BBMap (v 38.24) toolkit with
the parameters: "bbmap.sh minid=0.99" (https://github.com/BioInfoTools/BBMap). MetaBAT2 (v 2.12.1)
was used to perform binning analysis with the parameter: "-m 2000" (*6*).

One Asgard MAG recovered from J65 metagenome had a depth coverage of 14.52 and a N50 statistics of
10460 bps (**Supplementary Figure 2**).

Yap metagenome. For each Yap metagenome, raw metagenomic shotgun sequencing reads were trimmed
with Trimmomatic (v0.38) (*9*). Assembly and binning analysis were performed as described for CJE
metagenome for each Yap metagenome.

The depth coverage and N50 statistics of 24 Asgard MAGs recovered from Yap metagenomes range from
7.78 to 82.33 (median: 13.91) and from 5097 to 889102 bps (median: 18155.5 bps) respectively.
(**Supplementary Figure 2**)

A total of 89 Asgard MAGs were reconstructed in this work. Additional 95 Asgard MAGs were
downloaded from public databases (e.g. NCBI FTP site). For all 184 genomes, a uniform gene calling
protocol was applied. Specifically, the completeness, contamination, and strain heterogeneity of the
genomes were estimated by using CheckM (v.1.0.12) (*11*) and DAS Tool under the taxonomic scope of
domain (i.e., Bacteria and Archaea). Protein-coding genes were predicted using Prodigal (v 2.6.3) (*12*)
embedded in Prokka (v 1.13) (*13*). Transfer RNAs (tRNAs) were identified with tRNAscan-SE (v1.23)
using the archaeal tRNA model (*14*). After quality screening, further analysis focused on 162 high quality
Asgard MAGs.

Genome set

311   The Asgard archaea genome set analyzed in this work consisted of 161 Asgard MAGs and one complete
312   Asgard genome (**Supplementary Table 1**). For comparison, selected representative genomes of archaea
313   (296), bacteria (100) and eukaryotes (76) were downloaded from Refseq and Genbank using the NCBI
314   FTP site (ftp://ftp.ncbi.nlm.nih.gov/genomes/all/).

Average amino-acid identity

316   The average amino-acid identity (AAI) across TACK archaeal reference genomes and the 184 Asgard
317   genomes was calculated using compareM (v0.0.23) with the "aai_wf" at default settings
318   (https://github.com/dparks1134/CompareM).

asCOGs construction

Initial clustering of 250,634 proteins encoded in 76 Asgard MAGs was performed using two approaches:
first, footprints of arCOG profiles were obtained by running PSI-BLAST (*15*), initiated with arCOG
alignments, against the set of predicted Asgard proteins. The footprint sequences were extracted and
clustered according to the arCOG best hit. The remaining protein sequences (both full-length proteins and
the sequence fragments outside of the footprints, if longer than 60 aa) were clustered using MMseqs2
(*16*), with similarity threshold of 0.5. Sequences within clusters were aligned using MUSCLE (*17*); the
resulting alignments were passed through several rounds of merging and splitting. The merging phase
involved comparing alignments to each other using HHSEARCH (*18*), finding full-length cluster-to-
cluster matches, merging the sequence sets and re-aligning the new clusters. The splitting phase consisted
of the construction of an approximate phylogenetic tree of the sequences using FastTree (*19*) (gamma-
distributed site rates, WAG evolutionary model) with balanced mid-point tree rooting, identification of
subtrees maximizing the fraction of species (MAGs) representation and minimizing the number of
paralogs, and pruning such subtrees as separate clusters of putative orthologs. Clusters, derived from
arCOGs, were prohibited from merging across distinct arCOGs to prevent distant paralogs from forming
mixed clusters.

Phylogenetic analysis

16S rRNA gene phylogenetic analysis. 16S rRNA gene sequences were identified in 73 genomes of Asgard archaea (26 generated in this study, 47 from public database) using Barrnap (v 0.9) with the "--kingdom arc" option (https://github.com/tseemann/barrnap). These sequences were combined with 46 published 16S rRNA sequences of Asgard archaea, to assess the novelty of the sequences obtained in this work. The novelty of 16S rRNA gene sequences was measured in terms of their sequence identity to previously identified Asgard archaeal 16S rRNA gene and phylogenetic relationships. Specifically, the pairwise sequence identity of two Asgard archaeal 16S rRNA gene sequences (>1300bp) was obtained by first globally aligning the sequences with "Stretcher" in EMBOSS package and then calculating the percent identity excluding gaps (*20*). The Asgard archaeal 16S rRNA gene sequences were aligned with 311 reference sequences from Euryarchaeota (n=231), DPANN (n=22), Korarchaeota (n=1), Crenarchaeota (n=41), Bathyarchaeota (n=2), Thaumarchaeota (n=13) and Aigarchaeota (n=1) using mafft-linsi (v7.471) (*21*) and trimmed with BMGE (v1.12) (settings: -m DNAPAM250:4, -g 0.5) (*22*).The alignment was used for phylogenetic inference with IQ-Tree (v2.0.6) based on SYM+R8 (selected by ModelFinder) to generate a maximum-likelihood tree (*23*).

Asgard phylogeny. A set of asCOGs that were considered most suitable as phylogenetic markers for Asgard archaea was selected using the preliminary classification of the 76 genomes in AsCOGs into previously described lineages: Loki, Thor, Odin, Hel and Heimdall. The following criteria were adopted: the asCOG have to be i) present in at least half of the genomes in all lineages, ii) present in at least 75% among the 76 genomes, iii) the mean number of paralogs per genome not to exceed 1.25. For the 209 asCOGs matching these criteria, the corresponding protein sequences were obtained from the extended set of 162 MAGs and aligned using MUSCLE (*17*); the 'index' paralog to include in the phylogeny was selected for each MAG based on the similarity to the alignment consensus. Alignments were trimmed to exclude columns containing more than 2/3 gap characters and with homogeneity below 0.05 (homogeneity is calculated from the score of the consensus amino acid against the alignment column, compared to the score of the perfect match (*24*) and concatenated, resulting in an alignment of 50,706 characters from 162 sequences (one sequence per MAG). Phylogeny was reconstructed using FastTree (gamma-distributed site rates, WAG evolutionary model) (*19*) and IQ-Tree (LG+F+R10 model, selected by ModelFinder) (*23*), producing very similar tree topologies.

Tree of life. To elucidate the relationships between the Asgard archaea and other major clades of archaea, bacteria and eukaryotes, 30 families of conserved proteins were selected that appear to have evolved mostly vertically (25, 26). The set of 162 MAGs of Asgard archaea was supplemented with 66 TACK archaea and 220 non-TACK archaea (the former, having been described as the closest archaeal relatives of the Asgard archaea (*27*), were sampled more densely), 98 bacteria and 72 eukaryotes. Prokaryotic genomes were sampled from the set of completely sequenced genomes to represent the maximum diversity within their respective clades (briefly, all proteins, encoded by genomes within a groups were clustered at 75% identity level; distances between genomes were estimated from the number of shared proteins within these clusters; UPGMA trees were reconstructed from the distances, and genome sets maximizing the total tree branch length were selected to represent the groups). The set of eukaryotic genomes was manually selected to represent the maximum possible variety of eukaryotic taxa. Genomes, in which more than four markers were missing, were excluded from the bacterial and eukaryotic sets. When multiple paralogs of a marker were present in a genome, preliminary phylogenetic trees were

constructed from protein sequence alignments, and paralogs with the shortest branches were selected to
represent the corresponding genomes in the set. Sequences were aligned using MUSCLE (*17*), and
alignment columns, containing more than 2/3 gap characters or with alignment column homogeneity
below 0.05 were removed (*24*). The resulting concatenated alignments of the 30 markers consisted of
7411 sites. Phylogeny was reconstructed using FastTree (gamma-distributed site rates, WAG evolutionary
model) (*19*) and IQ-Tree (*23*) with three models: LG+R10, selected by IQ-tree ModelFinder as the best
fit, GTR20+F+R10 (following the suggestion of Williams et al. (*28*) to use GTR, we let IQ-tree to select
the best version of the GTR model), and LG+C20+G4+F (again, the mixture model was used following
the suggestion of Williams et al. (*28*); we were unable to use the higher-specified C60 model due to
memory limitations of our hardware and used the C20 model instead).

Ordination of asCOG phyletic patterns using Classical Multidimensional Scaling

Binary asCOG presence-absence patterns were compared between pairs of Asgard MAGs using the
following procedure: first, similarity between asCOG sets $\{A\}$ and $\{B\}$ was calculated as $S_{A,B} =$
$|A \cap B|/\sqrt{|A||B|}$ (the number of shared AsCOGs normalized by the geometric mean of the number of
AsCOGs in the two MAGs); then, the distance between the patterns was calculated as $d_{A,B} = -\ln(S_{A,B})$.
The 162x162 distance matrix was embedded into a 2-dimensional space using Classical Multidimensional
Scaling analysis implemented as the *cmdscale* function in R. The projection retained 89% of the original
datapoint inertia.

Identification and analysis of ESPs

To identify eukaryotic signature proteins (ESPs), several strategies were employed. First, ESPs reported
by Zaremba-Niedzwiedzka et al. (*29*) were mapped to asCOGs using PSI-BLAST (*15*). These asCOGs
were additionally examined case by case using HHpred (*30*) with representative of the respective asCOG
or the respective asCOG alignment used as the query. Second, all asCOGs were mapped to CDD profiles
(*31*) using PSI-BLAST, hits to eukaryote-specific domains were selected. Most of the putative ESP
asCOGs identified in this search, and all with E-value >1e-10, were additionally examined using HHpred,
with a representative of the respective asCOG or the respective asCOG alignment as the query. Third, we
analyzed frequently occurring asCOGs (present in at least 50% of Asgard genomes and in at least 30% of
Heimdall genomes) that were not annotated automatically with the above two approaches using HHpred
with a representative of the respective asCOG or the respective asCOG alignment as the query . Fourth,
most of the putative ESP asCOGs detected with these approaches were used as queries for a PSI-BLAST
search that was run for three iterations (with inclusion threshold E-value=0.0001) against an Asgard only
protein sequence database. Additional unannotated asCOGs with similarity to the (putative) ESPs
identified in this search were further examined using HHpred. Fifth, the genomic neighborhoods or all
ESPs were examined, and proteins encoded by unannotated neighbor genes were analyzed using HHpred
server.

Metabolic pathway reconstruction

The patterns of gene presence-absence in the asCOGs were used to reconstruct the metabolic pathways of
Asgard archaea (Supplementary Table 8). The asCOGs were linked to the KEGG database and to the list
of predicted metabolic enzymes of Asgard archaea reported by Spang et al. (*32*) (Supplementary Table 8).
The classification of [NiFe] hydrogenases was performed by comparing the Asgard proteins of

cog.001539, cog.002254, cog.010021, cog.011939 and cog.012499 to HydDB (*33*). For phylogenetic

analysis, the reference sequences of group 1, 3 and 4 [NiFe] hydrogenases were retrieved from HydDB.

The sequences were filtered using cd-hit with a sequence identity cut-off of 90% prior to adding

orthologous genes of cog.001539, cog.002254, cog.010021, cog.011939 and cog.012499 of Asgard

archaea. All sequences for group 1, group 3 and group 4 [NiFe] hydrogenases were aligned using mafft-

LINSI (*21*) and trimmed with BMGE (-m BLOSUM30 -h 0.6) (*22*). Maximum-likelihood phylogenetic

analyses were performed using IQ-tree (*23*) with the best-fit model (group 1:LG + C60 + R + F, group 3:

LG + C60 + R +F and group 4 LG + C50 + R + F), respectively, according to Bayesian information

criterion (BIC). Support values were estimated using the SH-like approximate-likelihood ratio test and

ultrafast bootstraps, respectively. We adapted a relaxed common denominator approach to determine the

presence of certain pathway in one Asgard phyla (*32*), and combined with maximum parsimony principle

(*34*) to infer the metabolisms of major ancestral forms.

References

1.   L. Cheng, T.-L. Qiu, X. Li, W.-D. Wang, Y. Deng, X.-B. Yin, H. Zhang, Isolation and characterization of Methanoculleus receptaculi sp. nov. from Shengli oil field, China. *FEMS Microbiology Letters*. **285**, 65–71 (2008).

2.   J. Peng, Z. Lü, J. Rui, Y. Lu, Dynamics of the Methanogenic Archaeal Community during Plant Residue Decomposition in an Anoxic Rice Field Soil. *Appl. Environ. Microbiol.* **74**, 2894–2901 (2008).

3.   G. V. Uritskiy, J. DiRuggiero, J. Taylor, MetaWRAP—a flexible pipeline for genome-resolved metagenomic data analysis. *Microbiome*. **6**, 158 (2018).

4.   D. Li, C. M. Liu, R. Luo, K. Sadakane, T. W. Lam, MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*. **31**, 1674–1676 (2015).

5.   B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat Methods*. **9**, 357–359 (2012).

6.   D. D. Kang, F. Li, E. Kirton, A. Thomas, R. Egan, H. An, Z. Wang, MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ*. **7**, e7359 (2019).

7.   C. M. K. Sieber, A. J. Probst, A. Sharrar, B. C. Thomas, M. Hess, S. G. Tringe, J. F. Banfield, Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nat Microbiol*. **3**, 836–843 (2018).

8.   Y. Peng, H. C. M. Leung, S. M. Yiu, F. Y. L. Chin, IDBA-UD: a de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics*. **28**, 1420–1428 (2012).

9.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. **30**, 2114–2120 (2014).

10.   S. Nurk, D. Meleshko, A. Korobeynikov, P. A. Pevzner, metaSPAdes: a new versatile metagenomic assembler. *Genome Res*. **27**, 824–834 (2017).

11.   D. H. Parks, M. Imelfort, C. T. Skennerton, P. Hugenholtz, G. W. Tyson, CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Research*. **25**, 1043–1055 (2015).

12.   D. Hyatt, G.-L. Chen, P. F. LoCascio, M. L. Land, F. W. Larimer, L. J. Hauser, Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*. **11**, 119 (2010).

13.   T. Seemann, Prokka: rapid prokaryotic genome annotation. *Bioinformatics*. **30**, 2068–2069 (2014).

14.   P. P. Chan, T. M. Lowe, tRNAscan-SE: Searching for tRNA genes in genomic sequences. *Methods Mol Biol*. **1962**, 1–14 (2019).

15.   A. A. Schäffer, L. Aravind, T. L. Madden, S. Shavirin, J. L. Spouge, Y. I. Wolf, E. V. Koonin, S.
F. Altschul, Improving the accuracy of PSI-BLAST protein database searches with composition-based
statistics and other refinements. *Nucleic Acids Res*. **29**, 2994–3005 (2001).

16.   M. Steinegger, J. Söding, MMseqs2 enables sensitive protein sequence searching for the analysis
of massive data sets. *Nat Biotechnol*. **35**, 1026–1028 (2017).

17.   R. C. Edgar, MUSCLE: a multiple sequence alignment method with reduced time and space
complexity. *BMC Bioinformatics*. **5**, 113 (2004).

18.   J. Söding, Protein homology detection by HMM-HMM comparison. *Bioinformatics*. **21**, 951–960
(2005).

19.   M. N. Price, P. S. Dehal, A. P. Arkin, FastTree 2--approximately maximum-likelihood trees for
large alignments. *PLoS One*. **5**, e9490 (2010).

20.   P. Rice, I. Longden, A. Bleasby, EMBOSS: the European Molecular Biology Open Software
Suite. *Trends Genet*. **16**, 276–277 (2000).

21.   K. Katoh, D. M. Standley, MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Molecular Biology and Evolution*. **30**, 772–780 (2013).

22.   A. Criscuolo, S. Gribaldo, BMGE (Block Mapping and Gathering with Entropy): a new software
for selection of phylogenetic informative regions from multiple sequence alignments. *BMC Evolutionary*
Biology. **10**, 210 (2010).

23.   L.-T. Nguyen, H. A. Schmidt, A. von Haeseler, B. Q. Minh, IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol*. **32**, 268–274
(2015).

24.   E. Esterman, Y. I. Wolf, R. Kogay, E. V. Koonin, O. Zhaxybayeva, *bioRxiv*, in press,
doi:10.1101/2020.10.08.331884.

25.   F. D. Ciccarelli, T. Doerks, C. von Mering, C. J. Creevey, B. Snel, P. Bork, Toward automatic
reconstruction of a highly resolved tree of life. *Science (New York, N.Y.)*. **311**, 1283–1287 (2006).

26.   P. Puigbò, Y. I. Wolf, E. V. Koonin, Search for a "Tree of Life" in the thicket of the phylogenetic
forest. *Journal of biology*. **8**, 59 (2009).

27.   A. Spang, J. H. Saw, S. L. Jørgensen, K. Zaremba-Niedzwiedzka, J. Martijn, A. E. Lind, R. van
Eijk, C. Schleper, L. Guy, T. J. G. Ettema, Complex archaea that bridge the gap between prokaryotes and
eukaryotes. *Nature*. **521**, 173–179 (2015).

28.   T. A. Williams, C. J. Cox, P. G. Foster, G. J. Szöllősi, T. M. Embley, Phylogenomics provides
robust support for a two-domains tree of life. *Nat Ecol Evol*. **4**, 138–147 (2020).

29.   K. Zaremba-Niedzwiedzka, E. F. Caceres, J. H. Saw, D. B ckstr m, L. Juzokaite, E. Vancaester,
K. W. Seitz, K. Anantharaman, P. Starnawski, K. U. Kjeldsen, M. B. Stott, T. Nunoura, J. F. Banfield, A.

Schramm, B. J. Baker, A. Spang, T. J. G. Ettema, Asgard archaea illuminate the origin of eukaryotic
cellular complexity. *Nature*. **541**, 353–358 (2017).

30.   L. Zimmermann, A. Stephens, S.-Z. Nam, D. Rau, J. Kübler, M. Lozajic, F. Gabler, J. Söding, A.
N. Lupas, V. Alva, A Completely Reimplemented MPI Bioinformatics Toolkit with a New HHpred
Server at its Core. *J Mol Biol*. **430**, 2237–2243 (2018).

31.   M. Yang, M. K. Derbyshire, R. A. Yamashita, A. Marchler-Bauer, NCBI's Conserved Domain
Database and Tools for Protein Domain Analysis. *Curr Protoc Bioinformatics*. **69**, e90 (2020).

32.   A. Spang, C. W. Stairs, N. Dombrowski, L. Eme, J. Lombard, E. F. Caceres, C. Greening, B. J.
Baker, T. J. G. Ettema, Proposal of the reverse flow model for the origin of the eukaryotic cell based on
comparative analyses of Asgard archaeal metabolism. *Nature Microbiology*. **4**, 1138–1148 (2019).

33.   D. Søndergaard, C. N. S. Pedersen, C. Greening, HydDB: A web tool for hydrogenase
classification and analysis. *Scientific Reports*. **6**, 34212 (2016).

34.   D. L. Swofford, W. P. Maddison, Reconstructing ancestral character states under Wagner
parsimony. *Mathematical Biosciences*. **87**, 199–229 (1987).

Data availability

Asgard archaea genomes generated in this study are available in eLMSG (an eLibrary of Microbial Systematics and Genomics, https://www.biosino.org/elmsg/index) under accession numbers from LMSG_G000000521.1 to LMSG_G000000609.1 and NCBI database under the project XXXXX.

Additional Data files are available at ftp://ftp.ncbi.nih.gov/pub/wolf/_suppl/asgard20/

Additional data file 1 (Additional_data_file_1.tgz): Complete asCOG data archive

Additional data file 2 (Additional_data_file_2.tgz): Phylogenetic trees and alignments archive

Supplementary Figures

Supplementary Figure 1 Global distribution of Asgard genomes analyzed in the study. The pie chart shows the proportion of Asgard genomes found in a biotope. Bold letters in the map show the sampling locations.

Supplementary Figure 2 a. Distribution of completeness and contamination for 76 Asgard MAGs assessed by CheckM (v 1.0.12). Distribution of depth coverage (b) and N50 statistics (c) for Asgard MAGs reconstructed in this study. The numbers in parentheses indicate the number of Asgard genomes recovered from a given sampling location. The data for this plot can be found in **Supplementary Table 1.**

Supplementary Figure 3 Comparison of the mean amino-acid identity (AAI) of Asgard and TACK superphyla. a. Shared AAI across Asgard and TACK lineages. Background: comparing all Asgard and TACK lineages included in the analyses but excluding archaea belonging to the same lineages and the six phyla proposed in the current work to investigate the distribution of AAI that defines a phylum. The AAI comparison of (b) Thorarchaeota, (c) Hermodarchaeota (d) Odinarchaeota, (e) Baldrarchaeota, (f) Lokiarchaeota, (g) Helarchaeota, (h) Borrarchaeota, (i) Heimdallarchaeota, (j) Kariarchaeota, (k) Gerdarchaeota, (l) Hodarchaeota and (m) Wukongarchaeota to other Asgad and TACK lineages. The lower and upper hinges of the boxplot correspond to the first and third quartiles. Data beyond the whiskers are shown as individual data points. Number in the parenthesis indicates the number of genomes in the lineages. Data for this plot is included in **Supplementary Table 2.**

Supplementary Figure 4 Comparison of the 16S rRNA gene sequence (>1300 bp) identity of Asgard and TACK lineages. a. Shared 16S rRNA gene sequence identity across Asgard and TACK lineages. Background:comparing all Asgard and TACK lineages included in the analyses but excluding archaea belonging to the same lineages and the six phyla proposed in the current study to investigate the distribution of 16S rRNA gene identity that defines a phylum. 16S rRNA gene identity comparison of (b) Thorarchaeota, (c) Hermodarchaeota (d) Odinarchaeota, (e) Lokiarchaeota, (f) Helarchaeota, (g) Heimdallarchaeota, (h) Kariarchaeota, (i) Gerdarchaeota, (j) Hodarchaeota and (k) Wukongarchaeota to other Asgard and TACK lineages. The lower and upper hinges of the boxplot correspond to the first and third quartiles. Data beyond the whiskers are shown as individual data points. Number in the parenthesis indicates the number of 16S rRNA gene sequences compared in the lineages. Line represents a 16S rRNA gene identity of 75%. Data for this plot could be found in **Supplementary Table 3.**

Supplementary Figure 5 Presence-absence of orthologs of Asgard core genes in other archaea, bacteria and eukaryotes.

Supplementary Figure 6 Phylogenetic analysis of group 4 [NiFe] hydrogenases in the Asgard archaea. The unrooted maximum-likelihood phylogenetic tree was built from an alignment of 425 sequences including 110 sequences of Asgard archaea, with 308 amino-acid positions.

Supplementary Figure 7 Phylogenetic analysis of group 3 [NiFe] hydrogenases in the Asgard archaea. The unrooted maximum-likelihood phylogenetic tree was built from an alignment of 813 sequences including 335 sequences of Asgard archaea, 514 amino-acid positions.

Supplementary Figure 8 Phylogenetic analysis of group 1 [NiFe] hydrogenases in the Asgard archaea.
The unrooted maximum-likelihood phylogenetic tree was built from an alignment of 541 sequences
including 2 sequences of Wukongarchaeaota, with 745 amino-acid positions.

Supplementary Figure 9 A schematic representation of the presence and absence of selected metabolic
features in all (putative) phyla of Asgard archaea.

Supplementary Figure 10 Gene structure of the contig encoding Group 1 [Ni,Fe]-hydrogenase in
Wukongarchaeota. Abbreviations: Ftr, Formylmethanofuran:tetrahydromethanopterin formyltransferase;
FwdD Formylmethanofuran dehydrogenase subunit D; FwdB, Formylmethanofuran dehydrogenase
subunit B; FwdA, Formylmethanofuran dehydrogenase subunit A; FwdC, Formylmethanofuran
dehydrogenase subunit C; HdrC, Heterodisulfide reductase, subunit C; HyaB, Ni,Fe-hydrogenase I large
subunit; HyaA, Ni,Fe-hydrogenase I small subunit.

Supplementary Tables

Supplementary Table 1 Genome information, proposed taxonomy and isolation data

Supplementary Table 2 Mean amino-acid identity values (in %) comparing 66 TACK genomes and 184
Asgard genomes (162 high quality and 22 low-quality)

Supplementary Table 3 The 16S rRNA gene sequence identity (in %) comparing TACK lineages and
Asgard lineages. The identity was calculated using sequences longer than 1300 bps

Supplementary Table 4 Species and phyletic markers used for the tree of life reconstruction

Supplementary Table 5 Data for phylogenetic trees: the trees in the Newick format and the underlying
alignments

Supplementary Table 6 The asCOGs annotation

Supplementary Table 7 Eukaryotic signature proteins in Asgard archaea

Supplementary Table 8 The presence-absence of metabolic enzymes in Asgard archaea.

Biotope
● Coastal sediment ● Hot spring ● Hypersaline lake sediment ● Marine water ● Petroleum seep (Marine)
● Freshwater sediment ● Hydrothermal vent ● Marine sediment ● Petroleum field

Tree scale: 1

various Asgard archaea

group 4c

group 4g

group 4g

group 4h

group 4d

Asgard archaea cluster

group 4i

group 4b

group 4f

group 4a

group 4e

group 4g

various Asgard archaea

Tree scale: 0.1

various Asgard archaea

Lokiarchaeota

group 3c

group 3d

group 3c

group 3a

Asgard archaea cluster

group 3b

various Asgard archaea

Tree scale: 1.25

group 1a

group 1h

group 1g

group 1i

group 1b

group 1j

group 1k

Wukongarchaeota

group 1c

group 1f

group 1d

group 1e