Individual-based simulations of genome evolution with ancestry: the

GᴇɴᴏᴍᴇAᴅᴍɪxR R package

Thijs Janzen[1,2*], Fernando Diaz[3]

[1] Groningen Institute for Evolutionary Life Sciences, University of Groningen, Box

111039700 CC Groningen, The Netherlands

[2] Carl von Ossietzky University, Carl-von-Ossietzky-Str. 9–11, 26111, Oldenburg,

Germany

[3] Department of Entomology, University of Arizona, Tucson, AZ, United States of

America

[*] t.janzen@rug.nl

Running headline: the GᴇɴᴏᴍᴇAᴅᴍɪxR R package

ABSTRACT

1.   Hybridization between populations or species results in a mosaic of the

two parental genomes. Genome admixture has received increasing attention for

its implications in speciation, human evolution, Evolve and Resequencing (E&R)

and genetic mapping. However, a thorough understanding of how local ancestry

changes after admixture, and how selection affects patterns of local ancestry

remains elusive. The complexity of these questions limits analytical treatment,

but these scenarios are specifically suitable for simulation. Currently, there is no

simulation framework that models whole-genome evolution of local ancestry

following admixture.

2.   Here, we present the R package GENOMEADMIXR, which uses an individual-

based model to simulate genomic patterns of local ancestry following admixture

forward in time. GENOMEADMIXR provides user-friendly functions to set up and

analyze simulations under evolutionary scenarios with selection, linkage and

migration.

3.   We show the flexible functionality of the GENOMEADMIXR workflow by

demonstrating 1) how to design an E&R simulation using GENOMEADMIXR and 2)

how to use GENOMEADMIXR to verify analytical expectations following from the

theory of junctions.

4.   GENOMEADMIXR provides a mechanistic approach to explore expected

genome responses to realistic admixture scenarios. With this package, we aim to

aid researchers in testing specific hypotheses based on empirical findings

involving admixing populations.

Keywords: Genome evolution, admixture, ancestry, individual-based modeling,

Evolution and Resequencing.

INTRODUCTION

Genetic exchange has long been recognized as an important driver of genetic diversity, from the recombination of conspecific genomes and the evolution of speciation with migration (Abbott *et al.* 2013) to introgressive hybridization (Janzen *et al.* 2018; Lavretsky *et al.* 2019) and horizontal gene transfer (Keeling & Palmer 2008). With recent advances in sequencing technologies, the view of genetic exchange has now moved from information at specific loci (*e.g.* alleles, genes) to the level of whole genome admixture (Leitwein *et al.* 2018; Lavretsky *et al.* 2019; Chafin & Douglas 2020). The focus is now on understanding how major drivers of evolution (*i.e.* recombination, selection and migration) shape the linear connection of loci along the genome (*i.e.* synteny and linkage) and how these processes evolve under particular scenarios of genome divergence. Understanding these dynamics has helped to elucidate the impact of migration on human evolution (Hellenthal *et al.* 2014; Payseur & Rieseberg 2016), the impact of hybridization on speciation (Schumer *et al.* 2014b, 2018), and helped as well in designing experiments relying on admixture, for instance Evolve and Resequencing (E&R) experiments (Franssen *et al.* 2017; Barghi & Schlötterer 2019; Otte & Schlötterer 2020).

Recombination is one of the main drivers shaping macro-genomic patterns after admixture, where contiguous blocks of ancestry are broken down into smaller blocks over time (Fisher 1954, 1959). As time progresses, genomes transform into macro-genomic mosaics consisting of many small contiguous ancestry blocks. Our understanding of how these mosaics form is relatively robust, but analytical treatments are only known for boundary cases, such as admixture between two distant populations, or for restricted backcrossing

(Fisher 1954, 1959; Stam 1980; Macleod *et al.* 2005; Janzen *et al.* 2018;

Lavretsky *et al.* 2019).

Linkage patterns can also be generated by selection through genetic

hitchhiking or selection of epistatic interactions, which can alter genome-wide

patterns of genetic variation, population dynamics and evolvability (Gerrish *et al.*

2007; Arnold & Kunte 2017; Zhou *et al.* 2017). Thus, macro-genomic patterns

can be driven both by recombination and selection alike. However, a

mathematical treatment of the interaction between selection, recombination and

admixture is currently lacking. In contrast, individual-based simulations are very

suitable to explore these interactions, for instance using packages such as

`MimicrEE2` (Vlachos & Kofler 2018) (written in Java) and `SLiM` (Haller &

Messer 2017, 2018) (written in C++). Unfortunately, these do not explicitly

track local ancestry, and information on haplotypes is often only indirectly

available or requires extensive post-processing. `forqs` (Kessner & Novembre

2014) and `SELAM` (Corbett-Detig & Jones 2016) (both written in C++) remedy

this issue and are specifically focused on tracking local ancestry and inheritance

of ancestry blocks, however these lack ease of use (requiring for instance local

compilation of C++ code, and configuration of input files) and are not easily

applied cross-platform. Here, we present GENOMEADMIXR, a software package

written in the R language. The R language is readily used within biology and can

easily be used cross-platform. GENOMEADMIXR provides routines to simulate

admixture in a host of scenarios (Figure 1), and includes a wide range of routines

available to analyze and visualize simulation results.

MATERIALS and METHODS

Description

GᴇɴᴏᴍᴇAᴅᴍɪxR is written in the R programming language (Team 2020).

Similar to SliM (Haller & Messer 2017, 2018) and SELAM (Corbett-Detig & Jones

2016), GᴇɴᴏᴍᴇAᴅᴍɪxR simulates a population forward in time using a Wright-

Fisher model with non-overlapping generations and a constant population size.

GᴇɴᴏᴍᴇAᴅᴍɪxR simulated diploid individuals that sexually reproduce. For

computational tractability, only one pair of chromosomes is simulated an all

individuals are assumed to be hermaphroditic. Recombination is modeled in the

same way as in for instance SLiM and SELAM (Corbett-Detig & Jones 2016; Haller

& Messer 2018), i.e. as a Poisson process, with the number of crossovers Poisson

distributed with the size in Morgan of the chromosome as rate parameter. The

location of crossovers is drawn from a uniform distribution, without

interference. The simulation tracks the locations of changes in local ancestry

(junctions) (like SELAM) but does not explicitly model nucleotides (as for

instance SLiM).

Usage

A development version can be installed from Github using the "devtools"

package:

```
112  > install.packages("devtools")
113  > devtools::install_github("thijsjanzen/GenomeAdmixR")
```

115 Alternatively, soon a version will be available from CRAN:

```
117  > install.packages("GenomeAdmixR")
```

118   The core of the GENOMEADMIXR package is formed by the simulation

119   function `simulate_admixture` (Table 1). To simulate a simple admixture

120   scenario where two distinct populations hybridize to form a population, which

121   continues to admix for 100 generations, we write:

123   ```
> pop <- simulate_admixture(pop_size = 1000, number_of_founders = 2, total_runtime =
124   100)
```

126     *Molecular markers and statistic estimations*

127     To simulate uncertainty in tracking local ancestry, in line with molecular

128     methods (Schumer *et al.* 2014a; Dennenmoser *et al.* 2019; Lavretsky *et al.* 2019),

129     GENOMEADMIXR includes functionality to track local ancestry at molecular

130     markers located along the genome. Resulting local ancestry information is

131     returned from the function in long format, which facilitates downstream

132     analysis. The number of used ancestral genomes will therefore represent the

133     alternative alleles at marker position. Consequently, using two ancestral

134     genomes would simulate a biallelic model.

135     Continuing along the selected scenario, but tracking 1000 markers along

136     the genome, one would get:

138     ```
> pop <- simulate_admixture(pop_size = 1000, number_of_founders = 2, total_runtime =
139     100, markers = seq(0, 1, length.out = 1000))
```

Based on these markers, a set of genetic diversity statistics can be

calculated, including allele frequency (`calculate_allele_frequencies`,

Table 1), heterozygosity (`calculate_heterozygosity,` Table 1), linkage

disequilibrium   (`calculate_ld`,   Table   1),   Tajima's   D

(`calculate_tajima_d`, Table 1) and F_{ST} (`calculate_fst`, Table 1), as

well as changes of these over generations for better assessing significance of

simulated evolution. Caution should be taken when more than four distinct

ancestries are included in the simulation, because most genetic diversity

statistics assume a maximum nucleotide number of four. For a single population

we can estimate Linkage Disequilibrium (LD), average heterozygosity and

tajima's D:

153
154
155

```
> ld <- calculate_ld(pop)
> het <- calculate_heterozygosity(pop)
> td <- calculate_tajima_d(pop)
```

Visualization

`Simulate_admixture` by default returns the frequencies of all

ancestors at the markers specified at the start and the end of the simulation.

Allele   frequency   changes   can   be   visualized   using

`plot_difference_frequencies_`   (Table   1,   Figure   2)   or

`plot_start_end` (Table 1, Figure 2) and using `plot_frequencies`

(Table 1, Figure 2) the final frequencies are plotted. The functions

`plot_over_time` (Table 1, Figure 2) and `plot_joyplot_frequencies`

(Table 1, Figure 2) provide functionality to show allele frequency changes over

time. All plotting functions return ggplot2 (Wickham 2009) objects, which can be

further customized by the user.

Migration

The function `simulate_admixture_migration` evolves two

independent populations in parallel, given a fraction of exchange of parental

individuals per generation.

```
176 > migr_pop <- simulate_admixture_migration(total_runtime = 1000, pop_size = c(100,
177 100), migration_rate = 0.01)
```

179 Then, we can analyze genetic divergence across these two populations using the

180 F_{ST} statistic:

```
182 > calculate_fst(migr_pop$population_1, migr_pop$population_2)
```

There is functionality included to stop the simulation once a threshold of genetic

differentiation (*i.e. F_{ST}*) between the two populations is reached.

Selection

In the absence of selection, allelic frequencies at haplotype sequences of

ancestral types remain constant, barring drift effects due to finite population

size. `simulate_admixture` provides additional options to impose fitness

benefits upon individuals that contain a marker under selection (where selection

is interpreted as favouring alleles from a specific ancestor). We follow

conventional fitness notation (Crow & Kimura 1970) and denote [w_{aa}, w_{Aa}, w_{AA}]

= [*1, 1+hs, 1+s*], where *w* indicates fitness, *s* indicates the selective benefit of

allele *A*, and *h* indicates the degree of dominance. Parents of offspring in the next

generation are drawn from the current parental population proportional to

fitness and the probability of selecting individual parent j as one of the parents of

an offspring in the next generation is given by:

$$P(j) = \frac{w_j}{\sum_{i=1}^{N} w_i}$$

The selection matrix [w_{aa}, w_{Aa}, w_{AA}] can be provided by the user, allowing for the

implementation of a wide range of potential selective benefits upon receiving a

marker, including overdominance and epistasis. Furthermore, the user is not

restricted to only providing selection on a single marker, and can expand the

selection matrix by adding extra rows for each marker. If multiple loci are under

selection, the fitness of an individual is the product of the finesses at each locus.

An example of adding selection to the previously selected hybridization scenario

is:

```
> s_matrix <- as.matrix(t(c(0.5, 1, 1.1, 1.5, 0)))
```
209   ```
> pop <- simulate_admixture(pop_size = 1000, number_of_founders = 2, total_runtime =
```
210   ```
100, markers = seq(0, 1, length.out = 1000), select_matrix = s_matrix)
```

212   **Demonstration of the package**

213   To demonstrate the functionality of GENOMEADMIXR, we firstly show a

214   series of simulations to compare alternative strategies in Evolution &

215   Resequencing (*E&R*) experiments. Secondly, we show how simulations of

216   GENOMEADMIXR match analytical predictions in junction theory.

218   *E&R experiment*

219   One of the goals in *E&R* experiments is to identify underlying genetic loci driving

220   responses to selection (Burke & Rose 2009; Schlötterer *et al.* 2015). Increased

221   genetic variability in the starting population improves the chances of capturing

222   additional candidate loci, but LD when sampling diverging populations can mask

223   the changes at individual positions (Kessner & Novembre 2015). We compare

224   two different sampling methods to increase genetic variability. *Model 1* starts

225   with a well-mixed population formed from 100 isofemale lines sampled from a

226   single (wild) population. *Model 2* aims to maximize genetic variability by

227   sampling two isofemale lines from different and diverging populations. Using

228   GENOMEADMIXR, we simulate the entire E&R pipeline and explore the impact of

229   both sampling schemes. Figure S1 summarizes the entire simulation setup and R

230   code to replicate our findings can be found in the supplementary material. Figure

231   3 (for a single simulation) indicates that the resolution to detect the marker

232   under selection in the experiment was much higher in *Model 1.* Also, *Model 1*

233   seems to experience less genetic hitchhiking compared to *Model 2*. We

234   performed 10 independent simulations with the same results, and replicates can

235   be used for further analysis and statistical tests. Overall, the obtained results

236   indicate that *Model 1* is for this pipeline the preferred approach (see for a more

237   detailed analysis the Supplementary Material).

239   *Theory of junctions*

240   From the extended theory of junctions (Janzen *et al.* 2018) we know that the

241   expected number of junctions (where a junction delineates the end of one

242   contiguous stretch of genomic content from the same ancestor, and the start of

243   another), depends on the initial heterozygosity H_0. Janzen's extended theory of

244 junctions focuses specifically on the scenario with two ancestors, whereas

245 GENOMEADMIXR allows for an arbitrary number of ancestors. We simulate the

246 accumulation of junctions using GENOMEADMIXR for two population sizes (100

247 and 1000 individuals) and vary the number of ancestors n in [2, 4, 8]. We find

248 that when we average the number of accumulated junctions over 1000

249 replicates, the average number of junctions closely follows that of our analytical

250 expectation (Figure 4).

252 **DISCUSSION**

253   We have presented here a new R package that uses individual-based

254 simulations to study patterns in local ancestry after admixture. We have shown

255 how GENOMEADMIXR can be used within an experimental evolution framework

256 and how results obtained using GENOMEADMIXR are in line with theoretical

257 expectations.

258   We expect GENOMEADMIXR to be applicable in many fields. Firstly, we

259 expect GENOMEADMIXR to allow for improved ease of use of individual-based

260 simulations to study patterns in the formation of contiguous tracts of ancestry,

261 and their breakdown due to recombination. The study of the breakdown of

262 ancestry blocks has recently attracted renewed interest, with studies expanding

263 the theory of junctions (the delineations between two ancestry tracts) towards

264 potential marker effects (Janzen *et al.* 2018), studies utilizing patterns in the

265 distribution of ancestry tracts to infer the onset of hybridization (Ungerer *et al.*

266 1998; Buerkle & Rieseberg 2008; Pool & Nielsen 2009; Gravel 2012; Liang &

267 Nielsen 2014; Corbett-Detig & Nielsen 2017) and the inclusion of ancestral tract

268 information in the analysis of human history (Hellenthal *et al.* 2014; Payseur &

269 Rieseberg 2016). `GenomeAdmixR` complements these analyses by providing a

270 framework to verify findings through simulation.

271   Secondly, we expect GENOMEADMIXR to be an important tool in

272 investigating population genomics and designing *E&R* experiments, where

273 GENOMEADMIXR can be used to test whether the setup of an experiment is

274 sufficient to detect signatures of selection on a background of existing LD. The

275 functionality of the package allows simulating the effect of *i)* previous population

276 dynamics that occurred before sampling (*natural populations*), *ii)* initial

277 parameters when obtaining the starting population for selection (*mixed*

278 *populations*) as well as *iii)* variation in genomic and *selection* parameters. Here,

279 we demonstrate how to use GENOMEADMIXR to estimate the level of resolution

280 obtained when combining inbred lines from different populations in the starting

281 population used in an E&R experiment (Parts *et al.* 2011; Koide *et al.* 2012). We

282 find that although this model narrowed down the haplotype blocks around the

283 selected maker substantially, this still lacks resolution when compared with

284 traditional sampling (Kofler & Schlötterer 2014; Schlötterer *et al.* 2015) from a

285 single population (even after 30 admixing generations). However, the obtained

286 resolution seems deep enough to investigate questions involving variation

287 between evolutionarily independent entities if these are not possible to address

288 by sampling a single population. Haplotype block dynamics have always been

289 important for understanding *E&R* experiments, but have recently been used to

290 explain massive genomic regions responding to selection in these studies

291 (Orozco-Terwengel *et al.* 2012; Turner & Miller 2012; Tobler *et al.* 2013),

292 limiting the ability to differentiate causal variant from hitchhiking regions.

293 Recent methods focus on haplotype block estimations and analyses (Franssen *et*

294 *al.* 2017; Burny *et al.* 2020; Otte & Schlötterer 2020) so much so that the

295 paradigm is shifting from analysis of single nucleotide polymorphisms to

296 selected haplotype blocks (Barghi & Schlötterer 2019).

297   Thirdly, we expect GENOMEADMIXR function as a useful teaching tool,

298 where it can be used to demonstrate the impact of hybridization on linkage

299 patterns, the subsequent interaction between selection and drift and more

300 generally to provide an easy toolkit to explore the interaction between

301 recombination and hybridization.

302   GENOMEADMIXR shares many similarities with it's predecessors SliM

303 (Haller & Messer 2017, 2018) and SELAM (Corbett-Detig & Jones 2016), both

304 powerful admixture simulation programs. However, SLiM is mainly focused on

305 more advanced population demographic scenarios and is not directly suited for

306 tracking ancestry. In contrast, SELAM does track local ancestry, but requires

307 local compilation of C++ code (which can be difficult) and uses complicated input

308 tables to parameterize the simulations. GENOMEADMIXR aims to alleviate both

309 these issues, firstly GENOMEADMIXR is specifically focused on tracking ancestry,

310 and it's admixture, over time. Secondly, because GENOMEADMIXR is an R package,

311 parameterization can be done through the use of the R scripting language, which

312 facilitates sharing of code between platforms, replication of approaches and ease

313 of use.

314   In conclusion, we believe that GENOMEADMIXR has a bright future ahead,

315 with a myriad of potential implementations across different fields of population

316 genetics and molecular ecology.

318 **AUTHOR CONTRIBUTIONS**

319   TJ and FD conceived the model, TJ developed the R code, FD and TJ jointly

320   wrote and revised the manuscript.

324   **DATA AVAILABILITY**

325   A development version of the package can be found at

326   https://github.com/thijsjanzen/GenomeAdmixR, and a stable version will soon

327   be available on CRAN (https://cran.r-project.org/package=GenomeAdmixR).

329   **ACKNOWLEDGEMENTS**

330   We thank Omer Markovitch, Pratik R. Gupte and Cyrus Mallon for helpful

331   comments on an earlier version of the manuscript. The authors indicate that they

332   have no conflicts of interest.

333
334

REFERENCES

Abbott, R., Albach, D., Ansell, S., Arntzen, J.W., Baird, S.J.E., Bierne, N., Boughman, J., Brelsford, a., Buerkle, C. a., Buggs, R., Butlin, R.K., Dieckmann, U., Eroukhmanoff, F., Grill, a., Cahan, S.H., Hermansen, J.S., Hewitt, G., Hudson, a. G., Jiggins, C., Jones, J., Keller, B., Marczewski, T., Mallet, J., Martinez-Rodriguez, P., Möst, M., Mullen, S., Nichols, R., Nolte, a. W., Parisod, C., Pfennig, K., Rice, a. M., Ritchie, M.G., Seifert, B., Smadja, C.M., Stelkens, R., Szymura, J.M., Väinölä, R., Wolf, J.B.W. & Zinner, D. (2013). Hybridization and speciation. *Journal of Evolutionary Biology*, **26**, 229–246.

Arnold, M.L. & Kunte, K. (2017). Adaptive Genetic Exchange: A Tangled History of Admixture and Evolutionary Innovation. *Trends in Ecology & Evolution*, **32**, 601–611.

Barghi, N. & Schlötterer, C. (2019). Shifting the paradigm in evolve and resequence studies: From analysis of single nucleotide polymorphisms to selected haplotype blocks. *Molecular Ecology*, **28**, 521–524.

Buerkle, C.A. & Rieseberg, L.H. (2008). The rate of genome stabilization in homoploid hybrid species. *Evolution*, **62**, 266–275.

Burke, M.K. & Rose, M.R. (2009). Experimental evolution with Drosophila. *Am J Physiol Regul Integr Comp Physiol*.

Burny, C., Nolte, V., Nouhaud, P., Dolezal, M., Schlötterer, C. & Baer, C. (2020). Secondary Evolve and Resequencing: An Experimental Confirmation of Putative Selection Targets without Phenotyping. *Genome Biology and Evolution*, **12**, 151–159.

Chafin, T.K. & Douglas, M.R. (2020). Genome-wide local ancestries discriminate homoploid hybrid speciation from secondary introgression in the red wolf (Canidae : Canis rufus). 1–49.

Corbett-Detig, R. & Jones, M. (2016). SELAM: Simulation of epistasis and local adaptation during admixture with mate choice. *Bioinformatics*, **32**, 3035–3037.

Corbett-Detig, R. & Nielsen, R. (2017). A Hidden Markov Model Approach for Simultaneously Estimating Local Ancestry and Admixture Time Using Next Generation Sequence Data in Samples of Arbitrary Ploidy. *PLoS Genetics*, **13**, 1–40.

Crow, J.F. & Kimura, M. (1970). *An Introduction to Population Genetics Theory*. Harper & Row, New York. Retrieved August 25, 2020, from https://www.cabdirect.org/cabdirect/abstract/19710105376

Dennenmoser, S., Schatz, M.C., Sedlazeck, F.J., Zytnicki, M., Nolte, A.W. & Altmüller, J. (2019). Genome-wide patterns of transposon proliferation in an evolutionary young hybrid fish. *Molecular Ecology*, 0–2.

Fisher, R.A. (1954). A fuller theory of "Junctions" in inbreeding. *Heredity*, **8**, 187–197.

Fisher, R.A. (1959). An algebraically exact examination of junction formation and transmission in parent-offspring inbreeding. *Heredity*, **13**, 179–186.

Franssen, S.U., Barton, N.H. & Schlotterer, C. (2017). Reconstruction of haplotype-blocks selected during experimental evolution. *Molecular Biology and Evolution*, **34**, 174–184.

Gerrish, P.J., Colato, A., Perelson, A.S. & Sniegowski, P.D. (2007). Complete genetic linkage can subvert natural selection. *PNAS*, **104**, 6266–6271.

384   Gravel, S. (2012). Population genetics models of local ancestry. *Genetics*, **191**,
385   607–619.
386   Haller, B.C. & Messer, P.W. (2017). SLiM 2: Flexible, interactive forward genetic
387   simulations. *Molecular Biology and Evolution*, **34**, 230–240.
388   Haller, B.C. & Messer, P.W. (2018). SLiM 3: Forward Genetic Simulations Beyond
389   the Wright–Fisher Model. *Molecular Biology and Evolution*, **36**, 632–637.
390   Hellenthal, G., Busby, G.B.J., Band, G., Wilson, J.F., Capelli, C., Falush, D. & Myers, S.
391   (2014). A Genetic Atlas of Human. *Science*, **343**, 747–751.
392   Janzen, T., Nolte, A.W. & Traulsen, A. (2018). The breakdown of genomic ancestry
393   blocks in hybrid lineages given a finite number of recombination sites.
394   *Evolution*, **72**, 735–750. Retrieved from
395   http://doi.wiley.com/10.1111/evo.13436
396   Keeling, P.J. & Palmer, J.D. (2008). Horizontal gene transfer in eukaryotic
397   evolution. *Nature Reviews Genetics*, **9**, 605–618.
398   Kessner, D. & Novembre, J. (2014). Forqs: Forward-in-time simulation of
399   recombination, quantitative traits and selection. *Bioinformatics*, **30**, 576–
400   577.
401   Kessner, D. & Novembre, J. (2015). Power analysis of artificial selection
402   experiments using efficient whole genome simulation of quantitative traits.
403   *Genetics*, **199**, 991–1005.
404   Kofler, R. & Schlötterer, C. (2014). A guide for the design of evolve and
405   resequencing studies. *Molecular Biology and Evolution*, **31**, 474–483.
406   Koide, T., Goto, T. & Takano-Shimizu, T. (2012). Genomic mixing to elucidate the
407   genetic system of complex traits. *Experimental Animals*, **61**, 503–509.
408   Lavretsky, P., Janzen, T. & McCracken, K.G. (2019). Identifying hybrids & the
409   genomics of hybridization: Mallards & American black ducks of Eastern
410   North America. *Ecology and Evolution*, **9**, 3470–3490.
411   Leitwein, M., Gagnaire, P.A., Desmarais, E., Berrebi, P. & Guinand, B. (2018).
412   Genomic consequences of a recent three-way admixture in supplemented
413   wild brown trout populations revealed by local ancestry tracts. *Molecular
414   Ecology*, **27**, 3466–3483.
415   Liang, M. & Nielsen, R. (2014). The lengths of admixture tracts. *Genetics*, **197**,
416   953–967.
417   Macleod, A.K., Haley, C.S., Woolliams, J.A. & Stam, P. (2005). Marker densities and
418   the mapping of ancestral junctions. *Genetical Research*, **85**, 69–79.
419   Orozco-Terwengel, P., Kapun, M., Nolte, V., Kofler, R., Flatt, T. & Schlãtterer, C.
420   (2012). Adaptation of Drosophila to a novel laboratory environment reveals
421   temporally heterogeneous trajectories of selected alleles. *Molecular Ecology*,
422   **21**, 4931–4941.
423   Otte, K.A. & Schlötterer, C. (2020). Detecting selected haplotype blocks in Evolve
424   and Resequence experiments. *Molecular Ecology Resources*, 1–37.
425   Parts, L., Cubillos, F.A., Warringer, J., Jain, K., Salinas, F., Bumpstead, S.J., Molin, M.,
426   Zia, A., Simpson, J.T., Quail, M.A., Moses, A., Louis, E.J., Durbin, R. & Liti, G.
427   (2011). Revealing the genetic structure of a trait by sequencing a population
428   under selection. 1131–1138.
429   Payseur, B.A. & Rieseberg, L.H. (2016). A genomic perspective on hybridization
430   and speciation. *Molecular Ecology*, **25**, 2337–2360.
431   Pool, J.E. & Nielsen, R. (2009). Inference of historical changes in migration rate
432   from the lengths of migrant tracts. *Genetics*, **181**, 711–719.

433 Schlötterer, C., Kofler, R., Versace, E., Tobler, R. & Franssen, S.U. (2015).
434   Combining experimental evolution with next-generation sequencing: A
435   powerful tool to study adaptation from standing genetic variation. *Heredity*,
436   **114**, 431–440.
437 Schumer, M., Cui, R., Powell, D.L., Dresner, R., Rosenthal, G.G. & Andolfatto, P.
438   (2014a). High-resolution mapping reveals hundreds of genetic
439   incompatibilities in hybridizing fish species. *eLife*, **2014**, 1–21.
440 Schumer, M., Rosenthal, G.G. & Andolfatto, P. (2014b). How common is
441   homoploid hybrid speciation? *Evolution*, **68**, 1553–1560.
442 Schumer, M., Xu, C., Powell, D.L., Durvasula, A., Skov, L., Holland, C., Blazier, C.,
443   Sankararaman, S., Andolfatto, P., Rosenthal, G.G. & Przeworski, M. (2018).
444   Natural selection interacts with recombination to shape the evolution of
445   hybrid genomes. *Science*, **360**, 656–660.
446 Stam, P. (1980). The distribution of the fraction of the genome identical by
447   descent in finite random mating populations. *Genetical Research*, **35**, 131.
448 Team, R.C. (2020). R: A Language and Environment for Statistical Computing.
449 Tobler, R., Franssen, S.U., Kofler, R., Orozco-terwengel, P., Nolte, V., Hermisson, J.
450   & Schlotter, C. (2013). Massive Habitat-Specific Genomic Response in D .
451   melanogaster Populations during Experimental Evolution in Hot and Cold
452   Environments. *Molecular Biology and Evolution*.
453 Turner, T.L. & Miller, P.M. (2012). Investigating natural variation in drosophila
454   courtship song by the evolve and resequence approach. *Genetics*, **191**, 633–
455   642.
456 Ungerer, M.C., Baird, S.J., Pan, J. & Rieseberg, L.H. (1998). Rapid hybrid speciation
457   in wild sunflowers. *Proceedings of the National Academy of Sciences of the
458   United States of America*, **95**, 11757–11762.
459 Vlachos, C. & Kofler, R. (2018). MimicrEE2: Genome-wide forward simulations of
460   Evolve and Resequencing studies. *PLoS Computational Biology*, **14**, 1–10.
461   Retrieved from http://dx.doi.org/10.1371/journal.pcbi.1006413
462 Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis - Hadley
463   Wickham - Google Books*. Retrieved September 9, 2020, from
464   https://books.google.nl/books?hl=nl&lr=&id=XgFkDAAAQBAJ&oi=fnd&pg=
465   PR8&dq=ggplot2&ots=so58bP5WbN&sig=ThS6gEgxaK9XADL_HeG5gDU04
466   Pc#v=onepage&q=ggplot2&f=false
467 Zhou, Y., Qiu, H. & Xu, S. (2017). Modeling Continuous Admixture Using
468   Admixture-Induced Linkage Disequilibrium. *Scientific Reports*, **7**, 1–10.
469
470

471 **Figures**

473 **Figure 1.** Overview of applications of the GENOMEADMIXR package. The
474 GENOMEADMIXR package can be used to **A**) simulate admixture of genomes
475 (rectangular, colored) over time, passed on between individuals (rounded
476 squares) when two populations (dotted lines) meet. **B**) Simulate ongoing genetic
477 exchange under migration, **C**) explore the impact of selection for alleles from the
478 red ancestor in the first section of the genome and **D**) Simulate the formation of
479 Recombinant Inbred Lines, where continued inbreeding generates fully
480 homozygous.
481

484 **Figure 2.** Example visualizations available in GenomeAdmixR. Shown are
485 example plots after an hybridization scenario between two distinct ancestral
486 populations, parameter values used are: population size = 10,000, total runtime
487 = 100. 1000 markers evenly spaced in [0, 1] were used. **A**:
488 `plot_difference_frequencies`, which visualizes the average change in
489 frequency between the start and end of the simulation **B**: plot_start_end, which
490 plots the average frequencies per ancestor, both at the start and the end of the
491 simulation. **C**: plot_frequencies, which plots the average frequency of each
492 ancestor across the genome **D**: plot_over_time, which plots the average
493 frequency of a single marker (here a marker at 50 cM) in the population, over
494 time. **E**: plot_joyplot_frequencies, which visualizes the average frequency over
495 time.

500
501
Figure 3. Summary of results obtained after simulating two scenarios for an E&R experiment using the GENOMEADMIXR package. Genome-wide Δ allele frequency after 5, 10 and 20 generations of selection are compared between *Model 1* and *Model 2*. The models differ in starting populations, with *Model* 1 only founded by 100 isofemales form a single population, and *Model* 2 founded by two isofemales form diverging populations. Selection was simulated for only one location (red point in the lower zoomed panel) and tracked in 300 biallelic markers, while the rest of the genome evolves neutrally.

Figure 4. Accumulation of junctions for different numbers of ancestors (2, 4 and 8 unique ancestors), for a small population (N = 100) and a population of intermediate size (N = 1000). Square dots indicate the mean number of junctions observed across 1000 replicate simulations with the GENOMEADMIXR package. Shaded areas indicate the 95% Confidence Interval across 1000 replicates. The solid lines indicate the analytical prediction following equation 1. Mean simulation dynamics follow the analytical prediction very closely.

519 **Table 1. Overview of available functions in the package**

Simulation	Input	Output	Summary
`simulate_admixture`	- input population (if available) - population size - number of unique ancestors (if no input population was given) - initial frequency of each unique ancestor - number of generations to simulate - size of chromosome in Morgan - selection matrix (see text) - a vector with locations of molecular markers	- Admixed population - tracked allele frequencies through time	Forward simulate a population
`simulate_admixture_migration`	Same input as `simulate_admixture` except: - two input populations (if available) - population sizes for both populations - migration rate - possibility to stop at a critical Fst value	- Two admixed populations - tracked allele frequencies through time	Forward simulate two populations connected via a fixed rate of migration m
Statistical methods			
`calculate_heterozygosity`	- input population - vector with locations of markers	- population average heterozygosity	Calculates average heterozygosity
`calculate_fst`	- two input populations - number of individuals sampled to calculate the statistic - number of markers sampled to calculate the statistic	Fst value	Calculates the Weir & Cockerham Fst for a set of markers
`calculate_LD`	- input population - number of individuals sampled to calculate the statistic - number of markers sampled to calculate the statistic	- pair wise linkage disequilibrium values for all markers - pair wise R^2 values for all markers	Calculates all pairwise LD and R^2 values for a set of markers
`calculate_tajima_D`	- input population - number of individuals sampled to calculate the statistic - vector with locations of markers used	- Tajima's D - π : the average pairwise difference between markers - S (the number of segregating sites) - θ : the expectation of π	Calculates Tajima's D
`calculate_marker_frequency`	- input population - locations of markers	- a table containing the frequency of each ancestor at the provided marker locations.	Estimates the frequency of ancestors at a marker location within the population
`calculate_allele_frequencies`			Estimates the frequency of ancestors in the population for

			sliding windows across the genome.
Visualization			
`plot.individual`	- genome of an individual (result of `simulate_admixture`)	Visualization of ancestry along both chromosomes, with different ancestry represented by different colors.	Plots both chromosomes of an individual, where colors represent ancestors
`plot_chromosome`	- chromosome of an inivindual	Visualization of ancestry along one chromosome, with different ancestry represented by different colors.	Visualize ancestry long a chromosome using different colors for each ancestor.
`plot_joyplot_frequencies`	- tracked frequencies over time (output object of simulate_admixture) - vector of timepoints to include in the plot - selection of ancestors to plot	A so called 'joyplot' or 'ridge plot', which visualizes density distributions over time.	Plots the distribution of genomic frequencies within a region as a joyplot, where the change over time in the distribution is visualized.
`plot_start_end`	- output of `simulate_admixture` - selection of ancestors to plot	Plots both the initial and final frequency of each ancestor along the genome.	Plot the distribution of genomic frequencies at the start and at the end of a simulation together in one plot
`plot_difference_frequencies`			Plots the difference in frequency, compared between the start and the end.
`plot_frequencies`	- output of `simulate_admixture` - a vector with marker locations	Plots the frequency of each ancestor at the end of the simulation	Plot the frequency of each ancestor along the genome
`plot_over_time`	- output of `simulate_admixture` - focal marker to focus on	Plot ancestor frequencies over time.	Plot the frequency of each ancestor at a specific marker, over time.
Utilities			
`save_population`	- population object - file name		Save a population to file
`load_population`	- file name	Population object	Load a population from file

521