Individual-based simulations of genome evolution with ancestry: the

GENOMEADMIXR R package

Thijs Janzen[1,2*], Fernando Diaz[3]

[1] Groningen Institute for Evolutionary Life Sciences, University of Groningen, Box

11103 9700 CC Groningen, The Netherlands

[2] Carl von Ossietzky University, Carl-von-Ossietzky-Str. 9–11, 26111, Oldenburg,

Germany

[3] Department of Entomology, University of Arizona, Tucson, AZ, United States of

America

[*] t.janzen@rug.nl

Running headline: the GENOMEADMIXR R package

ABSTRACT

1.   Hybridization between populations or species results in a mosaic of the

two parental genomes. This and other types of genome admixture have received

increasing attention for their implications in speciation, human evolution, Evolve

and Resequence (E&R) and genetic mapping. However, a thorough

understanding of how local ancestry changes after admixture, and how selection

affects patterns of local ancestry remains elusive. The complexity of these

questions limits analytical treatment, but these scenarios are specifically suitable

for simulation.

2.   Here, we present the R package GENOMEADMIXR, which uses an individual-

based model to simulate genomic patterns following admixture forward in time.

GENOMEADMIXR provides user-friendly functions to set up and analyze

simulations under evolutionary scenarios with selection, linkage and migration.

3.   We show the flexible functionality of the GENOMEADMIXR workflow by

demonstrating 1) how to design an E&R simulation using GENOMEADMIXR and 2)

how to use GENOMEADMIXR to verify analytical expectations following from the

theory of junctions.

4.   GENOMEADMIXR provides a mechanistic approach to explore expected

genome responses to realistic admixture scenarios. With this package, we aim to

aid researchers in testing specific hypotheses based on empirical findings

involving admixing populations.

Keywords: Genome evolution, admixture, ancestry, individual-based modeling,

Evolve and Resequence.

INTRODUCTION

Genetic exchange has long been recognized as an important driver of genetic diversity, from the recombination of conspecific genomes and the evolution of speciation with migration (Abbott *et al.* 2013) to introgressive hybridization (Janzen *et al.* 2018; Lavretsky *et al.* 2019) and horizontal gene transfer (Keeling & Palmer 2008). With recent advances in sequencing technologies, the view of genetic exchange has now moved from information at specific loci (*e.g.* alleles, genes) to the level of whole genome admixture (Leitwein *et al.* 2018; Lavretsky *et al.* 2019; Chafin & Douglas 2020). The focus is now on understanding how major drivers of evolution (*i.e.* recombination, selection and migration) shape the linear connection of loci along the genome (*i.e.* synteny and linkage) and how these processes evolve under particular scenarios of genome divergence. Understanding these dynamics has helped to elucidate the impact of migration on human evolution (Hellenthal *et al.* 2014; Payseur & Rieseberg 2016), the impact of hybridization on speciation (Schumer *et al.* 2014b, 2018), and helped as well in designing experiments relying on admixture, for instance Evolve and Resequence (E&R) experiments (Franssen *et al.* 2017; Barghi & Schlötterer 2019; Otte & Schlötterer 2020).

Recombination is one of the main drivers shaping macro-genomic patterns after admixture, where contiguous blocks of ancestry are broken down into smaller blocks over time (Fisher 1954, 1959). As time progresses, genomes transform into macro-genomic mosaics consisting of many small contiguous ancestry blocks. Our understanding of how these mosaics form is relatively robust, but analytical treatments are only known for boundary cases, such as admixture between two distant populations, or for restricted backcrossing

(Fisher 1954, 1959; Stam 1980; Macleod *et al.* 2005; Janzen *et al.* 2018;

Lavretsky *et al.* 2019).

Linkage Disequilibrium blocks can also be generated by selection through

genetic hitchhiking or selection of epistatic interactions, which can alter genome-

wide patterns of genetic variation, population dynamics and evolvability (Gerrish

et al. 2007; Arnold & Kunte 2017; Zhou *et al.* 2017). Thus, macro-genomic

patterns can be driven both by recombination and selection alike. However, a

mathematical treatment of the interaction between selection, recombination and

admixture is currently lacking. In contrast, individual-based simulations are very

suitable to explore these interactions, for instance using packages such as

MimicrEE2 (Vlachos & Kofler 2018) (written in Java) and SLiM (Haller &

Messer 2017, 2018) (written in C++). Unfortunately, these do not explicitly

track local ancestry, and information on haplotypes is often only indirectly

available or requires extensive post-processing. forqs (Kessner & Novembre

2014) and SELAM (Corbett-Detig & Jones 2016) (both written in C++) remedy

this issue and are specifically focused on tracking local ancestry and inheritance

of ancestry blocks, however these lack ease of use (requiring for instance local

compilation of C++ code, and configuration of input files) and are not easily

applied cross-platform. The R package plmgg (Cottin *et al.* 2020) is easily used

cross-platform, but focuses mainly on plant-like admixture, including selfing.

Here, we present GENOMEADMIXR, a software package written in the R

language. The R language is readily used within biology and can easily be used

cross-platform. GENOMEADMIXR provides routines to simulate admixture in a host

of scenarios (Figure 1), and includes a wide range of routines available to analyze

and visualize simulation results.

MATERIALS and METHODS

Description

MATERIALS and METHODS

Description

The interface of GENOMEADMIXR is written in the R programming language

(Team 2020), with the underlying simulation code using C++, integrated using

Rcpp (Eddelbuettel & Francois 2011). Similar to SliM (Haller & Messer 2017,

2018) and SELAM (Corbett-Detig & Jones 2016), GENOMEADMIXR simulates a

population forward in time usidng a Wright-Fisher model with non-overlapping

generations and a constant population size. GENOMEADMIXR simulates diploid

individuals that sexually reproduce. For computational tractability, only one pair

of chromosomes is simulated, and all individuals are assumed to be

hermaphroditic. Recombination is modeled in the same way as in for instance

SLiM and SELAM (Corbett-Detig & Jones 2016; Haller & Messer 2018), i.e. as a

Poisson process, with the number of crossovers Poisson distributed with the size

of the chromosome in Morgan as rate parameter. The location of crossovers is

drawn from a uniform distribution, without interference.

The package offers two different implementations of this simulation

scheme. Firstly, the user can use the ***Ancestry module*** of GENOMEADMIXR to

perform the simulations using known local ancestry, where the simulation tracks

the locations of changes in local ancestry (like SELAM and forqs) but does not

explicitly model nucleotides or mutation (as mutation between ancestries

complicates the simplifying assumptions of junctions' theory used to propagate

the simulation). Secondly, the ***Sequence module*** can be used to perform

simulations starting with sequencing data. Recombination is provided in cM/Mb

and simulations can therefore be performed on a section rather than the entire

chromosome. When using the *Sequence module*, mutation options are included as

well. Sequencing data can be loaded from VCF or PLINK format (using the

function `read_input_data`). To demonstrate the functionality of the package,

we have included sample data from the *Drosophila melanogaster* Reference Panel

(MacKay *et al.* 2012; Huang *et al.* 2014). The sample data consists of 5000 SNPs

with minimal allele frequency of 0.05, located along the 3R chromosome arm.

Usage

The GENOMEADMIXR package can be installed from CRAN:

124
125　
```
> install.packages(“GenomeAdmixR”)
```
126

The core of GENOMEADMIXR is formed by the function

`simulate_admixture`, customizable with two modules: the ***Ancestry module***

and the ***Sequence module*** (Table 1). Here we show how to implement

simulations using both modules side-by-side. To simulate a simple admixture

scenario where two distinct populations hybridize (e.g. a population of admixed

individuals resulting from a single mating event between two unrelated

individuals, resulting in an exactly 50/50 mixing of ancestral genomes in the first

generation) to form a population, which continues to admix for 100 generations,

we write:

Ancestry module　　　　　　　　***Sequence module***

138　
139　
140　
141　
142　
143　

144　
145　
146　
147　
148　
149　

```
> simulated_population <-
simulate_admixture(module =
ancestry_module(number_of_founders =
2), pop_size = 1000, total_runtime =
100)
```

```
> data("dgrp2.3R.5k.data")
> simulated_population <-
simulate_admixture_data(module =
sequence_module(input_data =
dgrp2.3R.5k.data), pop_size = 1000,
total_runtime = 100)
```

151 *Molecular markers and statistic estimations*

152   GENOMEADMIXR includes the functionality to track molecular markers

located along the genome, in line with molecular methods (Schumer *et al.* 2014a;

Dennenmoser *et al.* 2019; Lavretsky *et al.* 2019). For the *ancestry module*,

markers also serve to simulate the uncertainty in tracking local ancestry due to

limited coverage. The markers are assumed to be on a fixed position (in Morgan

for the *ancestry module*, and in bp for the *sequence module*) and are tracked over

time. Resulting local information is returned from the function in long format,

facilitating downstream analyses. Repeating the previous scenario, but now

tracking 1000 markers at fixed positions along the genome:

Ancestry module        **Sequence module**

```
163  > simulated_population <-              170  > simulated_population <-
164  simulate_admixture(module =           171  simulate_admixture_data(module =
165  ancestry_module(number_of_founders = 2, 172  sequence_module(input_data =
166  markers = seq(0, 1, length.out =      173  dgrp2.3R.5k.data, markers =
167  1000)), pop_size = 1000, total_runtime 174  dgrp2.3R.5k.data$markers[1:1000]),
168  = 100)                                175  pop_size = 1000, total_runtime = 100)
169
176
```

177   Based on these markers, a set of genetic diversity statistics can be

calculated, including allele frequency (`calculate_allele_frequencies`,

Table 1), heterozygosity (`calculate_heterozygosity`, Table 1), linkage

disequilibrium (`calculate_ld`, Table 1), and F_{ST} (F_{ST} is estimated following

Weir and Cockerham (Weir & Cockerham 1984), using the R package

`hierfstat` (Goudet 2005) for the implementation, see `calculate_fst`,

Table 1), as well as changes of these over generations for better assessing

significance of simulated evolution. When using the *ancestry module*, caution

should be taken when using more than four distinct ancestries, because most

genetic diversity statistics assume a maximum nucleotide number of four.

Demonstration of genetic diversity statistics

We estimate LD and average heterozygosity at different points in time.

Our expectation is for LD and heterozygosity to decrease over time. In the code

below, we either use 100 regularly spaced markers in Morgan (*ancestry module*),

or we draw 100 random marker positions from the data (*sequence module*).

193
194 ***Ancestry module***

Sequence module

```
> markers <- seq(from = 0, to = 1, length.out =
100)
> pop_10 <- simulate_admixture(module =
ancestry_module(number_of_founders = 2, markers
markers), total_runtime = 10, pop_size = 1000)

> ld_10 <- calculate_ld(pop_10, markers)
> het_10 <-
calculate_heterozygosity(pop_10$population,
locations = markers)
> pop_100 <- simulate_admixture(module =
ancestry_module(input_population = pop_10, markers
= markers), pop_size = 100, total_runtime = 90)
> ld_100 <- calculate_ld(pop_100, markers)
> het_100 <-
calculate_heterozygosity(pop_100$population,
locations = markers)
```

```
> data("dgrp2.3R.5k.data")
> markers <- sort(sample(dgrp2.3R.5k.data$markers,
100))
> pop_10 <- simulate_admixture_data(module =
sequence_module(input_data = dgrp2.3R.5k.data,
markers = markers), pop_size = 100, total_runtime =
10)
> ld_10 <- calculate_ld(pop_10, markers)
> het_10 <-
calculate_heterozygosity(pop_10$population,
locations = markers)
> pop_100 <- simulate_admixture_data(module =
sequence_module(input_data = pop_10, markers =
markers), pop_size = 100, total_runtime = 90)
> ld_100 <- calculate_ld(pop_100, markers)
> het_100 <-
calculate_heterozygosity(pop_100$population,
locations = markers)
```

The results (Figure 2) show that LD (here plotted as the correlation (R^2)

between loci) initially is relatively high, especially for closely located loci.

However, as time increases, LD disappears. Similarly, initially heterozygosity is

high (as expected), but decreases over time across the genome, until most

markers are fixed. Patterns are similar across the *ancestry* and *sequence*

simulations, although initially, LD is slightly higher for the *sequence* simulations..

Visualization

`simulate_admixture` by default returns the frequencies of all

ancestors at the start and the end of the simulation, if markers are specified.

Allele   frequency   changes   can   be   visualized   using

`plot_difference_frequencies_`   (Table   1,   Figure   3)   or

`plot_start_end` (Table 1, Figure 3) and using `plot_frequencies`

(Table 1, Figure 3) the final frequencies are plotted. The functions

`plot_over_time` (Table 1, Figure 3) and `plot_joyplot_frequencies`

(Table 1, Figure 3) provide functionality to show allele frequency changes over

time. All plotting functions return ggplot2 (Wickham 2009) objects, which can be

further customized by the user.

Migration

`Simulate_admixture` can be extended further by including migration,

where the specific settings for migration can be specified using the function

'migration_settings' (Table 1). Migration is simulated by evolving two

independent populations in parallel, given a fraction migrants each generations.

Ancestry module   ***Sequence module***

```
262   > migr_pop <- simulate_admixture(
263   module = ancestry_module(),
264   total_runtime = 100,
265   migration = migration_settings(pop_size
266   = c(100, 100), migration_rate = 0.01))
267
268
269
```

```
270   > migr_pop <- simulate_admixture
271   (module =
272   sequence_module(input_data_population_1
273   = dgrp2.3R.5k.data,
274   input_data_population_2 =
275   dgrp2.3R.5k.data), total_runtime = 100,
276   migration = migration_settings(
277   pop_size = c(100, 100), migration_rate
278   = 0.01))
```

280   Then, we can analyze genetic divergence across these two populations

281 using the F_{ST} statistic:

283 *Ancestry module*   *Sequence module*

```
284   >  calculate_fst(migr_pop$population_1,
285   migr_pop$population_2)
```

```
286   >  calculate_fst(migr_pop$population_1,
287   migr_pop$population_2)
```

There is functionality included to stop the simulation once a threshold of

genetic differentiation (*i.e. F_{ST}*) between the two populations is reached, which

facilitates scenarios where two related, but genetically different populations are

required.

Selection

`simulate_admixture` provides options to impose fitness benefits

upon individuals that contain a marker under selection (where selection is

interpreted as favouring alleles from a specific ancestor (*ancestry module*) or

SNP (*sequence module*). We follow conventional fitness notation (Crow & Kimura

1970) and denote [w_{aa}, w_{Aa}, w_{AA}] = [*1, 1+hs, 1+s*], where *w* indicates fitness, *s*

indicates the selective benefit of allele *A*, and *h* indicates the degree of

dominance. Parents of offspring in the next generation are drawn from the
parental population proportional to fitness.

The selection matrix [location, w_{aa}, w_{Aa}, w_{AA}, ancestor] includes
information on the selected marker, the different fitness weights, and the origin
of the allele under selection. The weights can be freely provided by the user,
allowing for the implementation of a wide range of potential selective benefits
upon receiving a marker, including overdominance and epistasis.

Furthermore, the user is not restricted to providing selection on a single
marker and can expand the selection matrix by adding extra rows for each
marker. This opens up the possibility for the user to explore polygenic selection
(as for instance for a Quantitative Trait Locus), by specifying several loci with
defined distances, where each locus provides a small fitness benefit. Hence, the
focal trait under selection here is translated directly in its resulting fitness effect.
If multiple loci are under selection, the fitness of an individual is the product of
the finesses at each locus. An example of adding selection on two markers at
locations of 50 and 60 cM to the previously selected hybridization scenario is:
317

318	*Ancestry module*	*Sequence module*

```
319   > s = 0.1
320   > ancestor_under_selection = 0
321   > markers <- c(0.5, 0.6)
322
323   > s_matrix <- matrix(nrow = 2, ncol = 5)
324   > s_matrix[1, ] <- c(markers[1], 1, 1 + 0.5
325   * s, 1 + s, ancestor_under_selection)
326   > s_matrix[2, ] <- c(markers[2], 1, 1 + 0.5
327   * s, 1 + s, ancestor_under_selection)
328
329   > pop <- simulate_admixture(module =
330   ancestry_module(number_of_founders = 2,
331   markers = markers), pop_size = 1000,
332   total_runtime = 100,select_matrix =
333   s_matrix)
```

```
334   > s <- 0.1
335   > allele_under_selection <- 1
336   > markers <-
337   sort(sample(dgrp2.3R.5k.data$markers, 2))
338   > s_matrix <- matrix(nrow = 2, ncol = 5)
339   > s_matrix[1, ] <- c(markers[1], 1, 1 + 0.5
340   * s, 1 + s, allele_under_selection)
341   > s_matrix[2, ] <- c(markers[2], 1, 1 + 0.5
342   * s, 1 + s, allele_under_selection)
343
344   > pop <- simulate_admixture(module =
345   sequence_module(input_data =
346   dgrp2.3R.5k.data, markers = markers ),
347   pop_size = 1000, total_runtime = 100, ,
348   select_matrix = s_matrix)
```

350   Demonstration of the package

To demonstrate the functionality of GENOMEADMIXR, we firstly show a

series of simulations to compare alternative strategies in Evolve & Resequence

(*E&R*) experiments. Secondly, we show how simulations of GENOMEADMIXR match

analytical predictions in junction theory.

E&R experiment

One of the goals in *E&R* experiments is to identify underlying genetic loci

driving responses to selection (Burke & Rose 2009; Schlötterer *et al.* 2015). The

most common sampling method starts with a "*well mixed population*" founded by

individuals sampled from the same natural population (*Model 1*). This method

minimizes Linkage Disequilibrium (*LD*) while trying to compensate for genetic

variability by performing large sampling (Kofler & Schlötterer 2014).

Alternatively, genetic variability can be maximized by sampling from different

populations, but this can generate additional *LD* (Kawecki *et al.* 2012; Kofler &

Schlötterer 2014; Schlötterer *et al.* 2015). However, it has been proposed that

genetic bases of population' or species' differences can be investigated by

investigating mixing lines from different populations or species (Parts *et al.*

2011). This method (*Model 2*) has been used in yeast to fine-tune Quantitative

Trait Locus (*QTL)* studies from phenotypically extreme inbred lines using

experimental selection (Parts *et al.* 2011; Koide *et al.* 2012).

We used GᴇɴᴏᴍᴇAᴅᴍɪxR to simulate the entire *E&R* pipeline and explore

the impact of both sampling schemes (Figure S1 demonstrates the simulation

setup). Figure 4 (for a single simulation) indicates that the resolution to detect

the marker under selection in the experiment was much higher in *Model 1.* Also,

Model 1 seems to experience less genetic hitchhiking compared to *Model 2*. We

performed 10 independent simulations with the same results, and replicates can

be used for further analysis and statistical tests. Overall, the obtained results

indicate that *Model 1* is for this pipeline the preferred approach. We

implemented the mentioned pipeline using both the *ancestry module* (Figure 4a)

and the *sequence module* (Figure 4b, using *D. melanogaster* Reference Panel data

of the 3R arm chromosome) rendering largely congruent results.

We further expanded *Model 1*, using individuals instead of isofemale lines

(*Models 3* and 4) revealing similar trends (See Figure S2).

Theory of junctions

From the extended theory of junctions (Janzen *et al.* 2018) we know that

the expected number of junctions (where a junction delineates the end of one

contiguous stretch of genomic content from the same ancestor, and the start of

another), depends on the initial heterozygosity H_0 (where heterozygosity here

reflects a locus with two alleles stemming from two different ancestors, e.g.

hetero-ancestry). Janzen's extended theory of junctions focuses specifically on

the scenario with two ancestors, whereas GENOMEADMIXR allows for an arbitrary

number of ancestors. We simulate the accumulation of junctions using the

ancestry module of GENOMEADMIXR for two population sizes (100 and 1000

individuals) and vary the number of ancestors n in [2, 4, 8]. We find that when

we average the number of accumulated junctions over 1000 replicates, the

average number of junctions closely follows that of our analytical expectation

(Figure 5).

DISCUSSION

We have presented here a new R package that uses individual-based simulations to study genomic dynamics following admixture. We have shown how both the *ancestry* and *sequence module* can be used for a general approach and more specifically, we have shown how GENOMEADMIXR can be used within an experimental evolution framework and how results obtained using GENOMEADMIXR are in line with theoretical expectations.

We expect GENOMEADMIXR to be applicable in many fields. Firstly, we expect GENOMEADMIXR to allow for improved ease of use of individual-based simulations to study patterns in the formation of contiguous tracts of ancestry, and their breakdown due to recombination. The study of the breakdown of ancestry blocks has recently attracted renewed interest, with studies expanding the theory of junctions towards potential marker effects (Janzen *et al.* 2018), studies utilizing patterns in the distribution of ancestry tracts to infer the onset of hybridization (Ungerer *et al.* 1998; Buerkle & Rieseberg 2008; Pool & Nielsen 2009; Gravel 2012; Liang & Nielsen 2014; Corbett-Detig & Nielsen 2017; Medina *et al.* 2018; Schumer *et al.* 2020; Shchur *et al.* 2020) and the inclusion of ancestral tract information in the analysis of human history (Hellenthal *et al.* 2014; Payseur & Rieseberg 2016). GENOMEADMIXR complements these analyses by providing a framework to verify findings through simulation. Furthermore, preliminary sequencing data can be used to verify findings using the *sequence module*, providing an additional tool to explore the efficacy of ancestry tract information.

Secondly, we expect GENOMEADMIXR to be an important tool in investigating population genomics and designing *E&R* experiments, where

GENOMEADMIXR can be used to test whether the setup of an experiment is

sufficient to detect signatures of selection on a background of existing LD. The

functionality of the package allows simulating the effect of *i)* previous population

dynamics that occurred before sampling (*natural populations*), *ii)* initial

parameters when obtaining the starting population for selection (*mixed*

populations) as well as *iii)* variation in genomic and *selection* parameters. Here,

we have demonstrated how to use GENOMEADMIXR to estimate the level of

resolution obtained when combining inbred lines from different populations in

the starting population of an E&R experiment (Parts *et al.* 2011; Koide *et al.*

2012). We found that although this model narrowed down the haplotype blocks

around the selected maker substantially, this still lacks resolution when

compared with traditional sampling (Kofler & Schlötterer 2014; Schlötterer *et al.*

2015) from a single population (even after 30 admixing generations). However,

the obtained resolution seems deep enough to investigate questions involving

variation between evolutionarily independent entities if these are not possible to

address by sampling a single population. For example, Comeault & Matute

(2018) used this model to study genomic trajectories following species hybridization.

Thirdly, we expect GENOMEADMIXR to function as a useful teaching tool,

where it can be used to demonstrate the impact of hybridization on linkage

patterns, the subsequent interaction between selection and drift and more

generally to provide an easy toolkit to explore the interaction between

recombination and hybridization.

GENOMEADMIXR shares many similarities with its predecessors SLiM (Haller

& Messer 2017, 2018) and SELAM (Corbett-Detig & Jones 2016), both powerful

admixture simulation programs. However, SLiM is mainly focused on more

advanced population demographic scenarios and is not directly suited for

tracking ancestry. In contrast, SELAM tracks local ancestry, but requires local

compilation of C++ code (which can be difficult) and uses complicated input

tables to parameterize the simulations. GENOMEADMIXR aims to alleviate both

these issues, firstly GENOMEADMIXR is specifically focused on tracking ancestry,

and its admixture over time. Secondly, because GENOMEADMIXR is an R package,

parameterization can be done through the use of the R scripting language, which

facilitates sharing of code between platforms, replication of approaches and ease

of use.

By providing both an *ancestry* and *sequence module*, we think that

GENOMEADMIXR provides a very complete package that can be used both to

explore theoretical expectations (*ancestry module*), and verify these expectations

using sequencing data (*sequence module*). Furthermore, because GENOMEADMIXR

readily accepts sequencing data in popular formats, It can be readily

incorporated in existing pipelines. Thus, we expect that GENOMEADMIXR has a

bright future ahead, with a myriad of potential implementations across different

fields of population genetics and molecular ecology.

AUTHOR CONTRIBUTIONS

TJ and FD conceived the model, TJ developed the R code, FD and TJ jointly

wrote and revised the manuscript.

DATA AVAILABILITY

The R package is available via CRAN on [https://cran.r-](https://cran.r-project.org/package=GenomeAdmixR)

[project.org/package=GenomeAdmixR](https://cran.r-project.org/package=GenomeAdmixR). All code used for the paper can be found

in the Supplemental Material.

ACKNOWLEDGEMENTS

We thank Omer Markovitch, Pratik R. Gupte and Cyrus Mallon for helpful

comments on an earlier version of the manuscript. The authors indicate that they

have no conflicts of interest.

490
491

REFERENCES

Abbott, R., Albach, D., Ansell, S., Arntzen, J.W., Baird, S.J.E., Bierne, N., Boughman, J., Brelsford, a., Buerkle, C. a., Buggs, R., Butlin, R.K., Dieckmann, U., Eroukhmanoff, F., Grill, a., Cahan, S.H., Hermansen, J.S., Hewitt, G., Hudson, a. G., Jiggins, C., Jones, J., Keller, B., Marczewski, T., Mallet, J., Martinez-Rodriguez, P., Möst, M., Mullen, S., Nichols, R., Nolte, a. W., Parisod, C., Pfennig, K., Rice, a. M., Ritchie, M.G., Seifert, B., Smadja, C.M., Stelkens, R., Szymura, J.M., Väinölä, R., Wolf, J.B.W. & Zinner, D. (2013). Hybridization and speciation. *Journal of Evolutionary Biology*, **26**, 229–246.

Arnold, M.L. & Kunte, K. (2017). Adaptive Genetic Exchange: A Tangled History of Admixture and Evolutionary Innovation. *Trends in Ecology & Evolution*, **32**, 601–611.

Barghi, N. & Schlötterer, C. (2019). Shifting the paradigm in evolve and resequence studies: From analysis of single nucleotide polymorphisms to selected haplotype blocks. *Molecular Ecology*, **28**, 521–524.

Buerkle, C.A. & Rieseberg, L.H. (2008). The rate of genome stabilization in homoploid hybrid species. *Evolution*, **62**, 266–275.

Burke, M.K. & Rose, M.R. (2009). Experimental evolution with Drosophila. *Am J Physiol Regul Integr Comp Physiol*.

Burny, C., Nolte, V., Nouhaud, P., Dolezal, M., Schlötterer, C. & Baer, C. (2020). Secondary Evolve and Resequencing: An Experimental Confirmation of Putative Selection Targets without Phenotyping. *Genome Biology and Evolution*, **12**, 151–159.

Chafin, T.K. & Douglas, M.R. (2020). Genome-wide local ancestries discriminate homoploid hybrid speciation from secondary introgression in the red wolf (Canidae : Canis rufus). 1–49.

Comeault, A.A. & Matute, D.R. (2018). Genetic divergence and the number of hybridizing species affect the path to homoploid hybrid speciation. *Proceedings of the National Academy of Sciences of the United States of America*, **115**, 9761–9766.

Corbett-Detig, R. & Jones, M. (2016). SELAM: Simulation of epistasis and local adaptation during admixture with mate choice. *Bioinformatics*, **32**, 3035–3037.

Corbett-Detig, R. & Nielsen, R. (2017). A Hidden Markov Model Approach for Simultaneously Estimating Local Ancestry and Admixture Time Using Next Generation Sequence Data in Samples of Arbitrary Ploidy. *PLoS Genetics*, **13**, 1–40.

Cottin, A., Penaud, B., Glaszmann, J.C., Yahiaoui, N. & Gautier, M. (2020). Simulation-based evaluation of three methods for local ancestry deconvolution of non-model crop species genomes. *G3: Genes, Genomes, Genetics*, **10**, 569–579.

Crow, J.F. & Kimura, M. (1970). *An Introduction to Population Genetics Theory*. Harper & Row, New York.

Dennenmoser, S., Schatz, M.C., Sedlazeck, F.J., Zytnicki, M., Nolte, A.W. & Altmüller, J. (2019). Genome-wide patterns of transposon proliferation in an evolutionary young hybrid fish. *Molecular Ecology*, 0–2.

Eddelbuettel, D. & Francois, R. (2011). Seamless R and C++ integration with Rcpp. *Journal of Statistical Software*, **40**, 1–18. Retrieved from

http://www.jstatsoft.org/v40/i08/

Fisher, R.A. (1954). A fuller theory of "Junctions" in inbreeding. *Heredity*, **8**, 187–
197.

Fisher, R.A. (1959). An algebraically exact examination of junction formation and
transmission in parent-offspring inbreeding. *Heredity*, **13**, 179–186.

Franssen, S.U., Barton, N.H. & Schlotterer, C. (2017). Reconstruction of haplotype-
blocks selected during experimental evolution. *Molecular Biology and*
Evolution, **34**, 174–184.

Gerrish, P.J., Colato, A., Perelson, A.S. & Sniegowski, P.D. (2007). Complete genetic
linkage can subvert natural selection. *PNAS*, **104**, 6266–6271.

Goudet, J. (2005). HIERFSTAT , a package for R to compute and test hierarchical F
-statistics. *Molecular Ecology Notes*, **2**, 184–186.

Gravel, S. (2012). Population genetics models of local ancestry. *Genetics*, **191**,
607–619.

Haller, B.C. & Messer, P.W. (2017). SLiM 2: Flexible, interactive forward genetic
simulations. *Molecular Biology and Evolution*, **34**, 230–240.

Haller, B.C. & Messer, P.W. (2018). SLiM 3: Forward Genetic Simulations Beyond
the Wright–Fisher Model. *Molecular Biology and Evolution*, **36**, 632–637.

Hellenthal, G., Busby, G.B.J., Band, G., Wilson, J.F., Capelli, C., Falush, D. & Myers, S.
(2014). A Genetic Atlas of Human. *Science*, **343**, 747–751.

Huang, W., Massouras, A., Inoue, Y., Peiffer, J., Ràmia, M., Tarone, A.M., Turlapati,
L., Zichner, T., Zhu, D., Lyman, R.F., Magwire, M.M., Blankenburg, K., Carbone,
M.A., Chang, K., Ellis, L.L., Fernandez, S., Han, Y., Highnam, G., Hjelmen, C.E.,
Jack, J.R., Javaid, M., Jayaseelan, J., Kalra, D., Lee, S., Lewis, L., Munidasa, M.,
Ongeri, F., Patel, S., Perales, L., Perez, A., Pu, L.L., Rollmann, S.M., Ruth, R.,
Saada, N., Warner, C., Williams, A., Wu, Y.Q., Yamamoto, A., Zhang, Y., Zhu, Y.,
Anholt, R.R.H., Korbel, J.O., Mittelman, D., Muzny, D.M., Gibbs, R.A.,
Barbadilla, A., Johnston, J.S., Stone, E.A., Richards, S., Deplancke, B. & Mackay,
T.F.C. (2014). Natural variation in genome architecture among 205
Drosophila melanogaster Genetic Reference Panel lines. *Genome Research*,
24, 1193–1208.

Janzen, T., Nolte, A.W. & Traulsen, A. (2018). The breakdown of genomic ancestry
blocks in hybrid lineages given a finite number of recombination sites.
Evolution, **72**, 735–750. Retrieved from
http://doi.wiley.com/10.1111/evo.13436

Kawecki, T.J., Lenski, R.E., Ebert, D., Hollis, B., Olivieri, I. & Whitlock, M.C. (2012).
Experimental evolution. *Trends in ecology & evolution*, **27**, 547–60.

Keeling, P.J. & Palmer, J.D. (2008). Horizontal gene transfer in eukaryotic
evolution. *Nature Reviews Genetics*, **9**, 605–618.

Kessner, D. & Novembre, J. (2014). Forqs: Forward-in-time simulation of
recombination, quantitative traits and selection. *Bioinformatics*, **30**, 576–
577.

Kessner, D. & Novembre, J. (2015). Power analysis of artificial selection
experiments using efficient whole genome simulation of quantitative traits.
Genetics, **199**, 991–1005.

Kofler, R. & Schlötterer, C. (2014). A guide for the design of evolve and
resequencing studies. *Molecular Biology and Evolution*, **31**, 474–483.

Koide, T., Goto, T. & Takano-Shimizu, T. (2012). Genomic mixing to elucidate the
genetic system of complex traits. *Experimental Animals*, **61**, 503–509.

Lavretsky, P., Janzen, T. & McCracken, K.G. (2019). Identifying hybrids & the
genomics of hybridization: Mallards & American black ducks of Eastern
North America. *Ecology and Evolution*, **9**, 3470–3490.
Leitwein, M., Gagnaire, P.A., Desmarais, E., Berrebi, P. & Guinand, B. (2018).
Genomic consequences of a recent three-way admixture in supplemented
wild brown trout populations revealed by local ancestry tracts. *Molecular*
Ecology, **27**, 3466–3483.
Liang, M. & Nielsen, R. (2014). The lengths of admixture tracts. *Genetics*, **197**,
953–967.
MacKay, T.F.C., Richards, S., Stone, E.A., Barbadilla, A., Ayroles, J.F., Zhu, D.,
Casillas, S., Han, Y., Magwire, M.M., Cridland, J.M., Richardson, M.F., Anholt,
R.R.H., Barrón, M., Bess, C., Blankenburg, K.P., Carbone, M.A., Castellano, D.,
Chaboub, L., Duncan, L., Harris, Z., Javaid, M., Jayaseelan, J.C., Jhangiani, S.N.,
Jordan, K.W., Lara, F., Lawrence, F., Lee, S.L., Librado, P., Linheiro, R.S.,
Lyman, R.F., MacKey, A.J., Munidasa, M., Muzny, D.M., Nazareth, L., Newsham,
I., Perales, L., Pu, L.L., Qu, C., Ràmia, M., Reid, J.G., Rollmann, S.M., Rozas, J.,
Saada, N., Turlapati, L., Worley, K.C., Wu, Y.Q., Yamamoto, A., Zhu, Y.,
Bergman, C.M., Thornton, K.R., Mittelman, D. & Gibbs, R.A. (2012). The
Drosophila melanogaster Genetic Reference Panel. *Nature*, **482**, 173–178.
Macleod, A.K., Haley, C.S., Woolliams, J.A. & Stam, P. (2005). Marker densities and
the mapping of ancestral junctions. *Genetical Research*, **85**, 69–79.
Medina, P., Thornlow, B., Nielsen, R. & Corbett-Detig, R. (2018). Estimating the
timing of multiple admixture pulses during local ancestry inference.
Genetics, **210**, 1089–1107.
Orozco-Terwengel, P., Kapun, M., Nolte, V., Kofler, R., Flatt, T. & Schlätterer, C.
(2012). Adaptation of Drosophila to a novel laboratory environment reveals
temporally heterogeneous trajectories of selected alleles. *Molecular Ecology*,
21, 4931–4941.
Otte, K.A. & Schlötterer, C. (2020). Detecting selected haplotype blocks in Evolve
and Resequence experiments. *Molecular Ecology Resources*, 1–37.
Parts, L., Cubillos, F.A., Warringer, J., Jain, K., Salinas, F., Bumpstead, S.J., Molin, M.,
Zia, A., Simpson, J.T., Quail, M.A., Moses, A., Louis, E.J., Durbin, R. & Liti, G.
(2011). Revealing the genetic structure of a trait by sequencing a population
under selection. 1131–1138.
Payseur, B.A. & Rieseberg, L.H. (2016). A genomic perspective on hybridization
and speciation. *Molecular Ecology*, **25**, 2337–2360.
Pool, J.E. & Nielsen, R. (2009). Inference of historical changes in migration rate
from the lengths of migrant tracts. *Genetics*, **181**, 711–719.
Schlötterer, C., Kofler, R., Versace, E., Tobler, R. & Franssen, S.U. (2015).
Combining experimental evolution with next-generation sequencing: A
powerful tool to study adaptation from standing genetic variation. *Heredity*,
114, 431–440.
Schumer, M., Cui, R., Powell, D.L., Dresner, R., Rosenthal, G.G. & Andolfatto, P.
(2014a). High-resolution mapping reveals hundreds of genetic
incompatibilities in hybridizing fish species. *eLife*, **2014**, 1–21.
Schumer, M., Powell, D.L. & Corbett-Detig, R. (2020). Versatile simulations of
admixture and accurate local ancestry inference with mixnmatch and
ancestryinfer. *Molecular Ecology Resources*, **20**, 1141–1151.
Schumer, M., Rosenthal, G.G. & Andolfatto, P. (2014b). How common is

homoploid hybrid speciation? *Evolution*, **68**, 1553–1560.

Schumer, M., Xu, C., Powell, D.L., Durvasula, A., Skov, L., Holland, C., Blazier, C.,
Sankararaman, S., Andolfatto, P., Rosenthal, G.G. & Przeworski, M. (2018).
Natural selection interacts with recombination to shape the evolution of
hybrid genomes. *Science*, **360**, 656–660.

Shchur, V., Svedberg, J., Medina, P., Corbett-Detig, R. & Nielsen, R.A.S.M. (2020).
On the distribution of tract lengths during adaptive introgression. *G3: Genes,*
Genomes, Genetics, **10**, 3663–3673.

Stam, P. (1980). The distribution of the fraction of the genome identical by
descent in finite random mating populations. *Genetical Research*, **35**, 131.

Team, R.C. (2020). R: A Language and Environment for Statistical Computing.

Tobler, R., Franssen, S.U., Kofler, R., Orozco-terwengel, P., Nolte, V., Hermisson, J.
& Schlotterer, C. (2013). Massive Habitat-Specific Genomic Response in D .
melanogaster Populations during Experimental Evolution in Hot and Cold
Environments. *Molecular Biology and Evolution*.

Turner, T.L. & Miller, P.M. (2012). Investigating natural variation in drosophila
courtship song by the evolve and resequence approach. *Genetics*, **191**, 633–
642.

Ungerer, M.C., Baird, S.J., Pan, J. & Rieseberg, L.H. (1998). Rapid hybrid speciation
in wild sunflowers. *Proceedings of the National Academy of Sciences of the*
United States of America, **95**, 11757–11762.

Vlachos, C. & Kofler, R. (2018). MimicrEE2: Genome-wide forward simulations of
Evolve and Resequencing studies. *PLoS Computational Biology*, **14**, 1–10.
Retrieved from http://dx.doi.org/10.1371/journal.pcbi.1006413

Weir, B.S. & Cockerham, C.C. (1984). Estimating F-Statistics for the Analysis of
Population Structure Author (s): B . S . Weir and C . Clark Cockerham
Published by⬚: Society for the Study of Evolution Stable URL⬚:
http://www.jstor.org/stable/2408641. *Evolution*, **38**, 1358–1370.

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis - Hadley*
Wickham - Google Books. Retrieved September 9, 2020, from
https://books.google.nl/books?hl=nl&lr=&id=XgFkDAAAQBAJ&oi=fnd&pg=
PR8&dq=ggplot2&ots=so58bP5WbN&sig=ThS6gEgxaK9XADL_HeG5gDU04
Pc#v=onepage&q=ggplot2&f=false

Zhou, Y., Qiu, H. & Xu, S. (2017). Modeling Continuous Admixture Using
Admixture-Induced Linkage Disequilibrium. *Scientific Reports*, **7**, 1–10.

674
675

Figures

Figure 1. Overview of applications of the GᴇɴᴏᴍᴇAᴅᴍɪxR package. The
GᴇɴᴏᴍᴇAᴅᴍɪxR package can be used to **A**) simulate admixture of genomes
(rectangular, colored) over time, passed on between individuals (rounded
squares) when two populations (dotted lines) meet. **B**) Simulate ongoing genetic
exchange under migration, **C**) explore the impact of selection for alleles from the
red ancestor in the first section of the genome and **D**) Simulate the formation of
Recombinant Inbred Lines, where continued inbreeding generates fully
homozygous.
686

Figure 2. Plot showing Linkage Disequilibrium (R^2) (left) and average heterozygosity (right) for a population 10 and 100 generations after admixture, where the initial admixture event involved two unrelated populations. Shown are results using the *ancestry module* (left columns) and the *sequence module* (right columns).

Figure 3. Example visualizations available in GenomeAdmixR. Shown are
example plots after an hybridization scenario using the *ancestry module* between
two distinct ancestral populations, parameter values used are: population size =
10,000, total runtime = 100. 1000 markers evenly spaced in [0, 1] were used. **A**:
`plot_difference_frequencies`, which visualizes the average change in
frequency between the start and end of the simulation **B**: plot_start_end, which
plots the average frequencies per ancestor, both at the start and the end of the
simulation. **C**: plot_frequencies, which plots the average frequency of each
ancestor across the genome **D**: plot_over_time, which plots the average
frequency of a single marker (here a marker at 50 cM) in the population, over
time. **E**: plot_joyplot_frequencies, which visualizes the average frequency over
time.

a) Ancestry module

b) Sequence module

Figure 4. Summary of results obtained after simulating two scenarios for an
E&R experiment using the GENOMEADMIXR package. Genome-wide ⏷⏷allele
frequency after 5, 10 and 20 generations of selection are compared between the
two models. The models differ in their starting populations as they were founded
form a single or diverging populations (*e.g. Models 1* vs *Model 2* respectively).
Model 1, founded by 100 isofemales form a single population; *Model 2*, founded
by two isofemales form diverging populations. Selection was simulated for only
one location (red point in the lower zoomed panel) in a 0.2 Morgan window, and
tracked in 1000 markers, while the rest of the genome evolves neutrally. The
results are shown for simulations using the **a)** *Ancestry module* and, **b)** the
Sequence module starting with data from the *D.* melanogaster Reference Panel.
730

Figure 5. Accumulation of junctions for different numbers of ancestors (2, 4 and 8 unique ancestors), for a small population (N = 100) and a population of intermediate size (N = 1000). Square dots indicate the mean number of junctions observed across 1000 replicate simulations with the GENOMEADMIXR package. Shaded areas indicate the 95% Confidence Interval across 1000 replicates. The solid lines indicate the analytical prediction following equation 1. Mean simulation dynamics follow the analytical prediction very closely.

740   **Table 1. Overview of available functions in the package**

Simulation	Input	Output	Summary
`simulate_admixture`	- chosen module (ancestry module or sequence_module) - population size - number of generations to simulate - migration settings (generated with function migration_settings) - selection matrix (see text) - number of threads	- Admixed population - tracked allele frequencies through time	Forward simulate a population
`ancestry_module`	- input population - number of unique ancestors (if no input population was given) - initial frequency of each unique ancestor - size of chromosome in Morgan a vector with locations of molecular markers	Ancestry module object	
`sequence_module`	- molecular data - initial frequencies - size of chromosome in morgan - recombination rate (if not an entire chromosome is modeled) - a vector with locations of molecular markers - mutation rate - substitution matrix	Sequence module object	
`migration settings`	- migration rate - option to stop at critical FST value - critical FST value - population size vector for both populations - initial frequencies (for ancestry_module simulation) - generations between measure of FST - number of individuals sampled to calculate FST - number of markers used to calculate FST	Migration settings object	
Statistical methods			
`calculate_heterozygosity`	- input population - vector with locations of markers	- population average heterozygosity	Calculates average heterozygosity
`calculate_fst`	- two input populations - number of individuals sampled to calculate the statistic - number of markers sampled to calculate the statistic	Fst value	Calculates the Weir & Cockerham Fst for a set of markers
`calculate_LD`	- input population - number of individuals sampled to calculate the statistic - number of markers sampled to calculate the statistic	- pair wise linkage disequilibrium values for all markers - pair wise R^2 values for all markers	Calculates all pairwise LD and R^2 values for a set of markers
`calculate_marker_frequency`	- input population - locations of markers	- a table containing the frequency of each ancestor at the provided marker locations.	Estimates the frequency of ancestors at a marker location within the population

`calculate_allele_frequencies`			Estimates the frequency of ancestors in the population for sliding windows across the genome.
Visualization			
`plot.individual`	- genome of an individual (result of `simulate_admixture`)	Visualization of ancestry along both chromosomes, with different ancestry represented by different colors.	Plots both chromosomes of an individual, where colors represent ancestors
`plot_chromosome`	- chromosome of an inividual	Visualization of ancestry along one chromosome, with different ancestry represented by different colors.	Visualize ancestry long a chromosome using different colors for each ancestor.
`plot_joyplot_frequencies`	- tracked frequencies over time (output object of simulate_admixture) - vector of timepoints to include in the plot - selection of ancestors to plot	A so called 'joyplot' or 'ridge plot', which visualizes density distributions over time.	Plots the distribution of genomic frequencies within a region as a joyplot, where the change over time in the distribution is visualized.
`plot_start_end`	- output of `simulate_admixture` - selection of ancestors to plot	Plots both the initial and final frequency of each ancestor along the genome.	Plot the distribution of genomic frequencies at the start and at the end of a simulation together in one plot
`plot_difference_frequencies`			Plots the difference in frequency, compared between the start and the end.
`plot frequencies`	- output of `simulate_admixture` - a vector with marker locations	Plots the frequency of each ancestor at the end of the simulation	Plot the frequency of each ancestor along the genome
`plot_over_time`	- output of `simulate_admixture` - focal marker to focus on	Plot ancestor frequencies over time.	Plot the frequency of each ancestor at a specific marker, over time.
Utilities			
`save_population`	- population object - file name		Save a population to file
`load population`	- file name	Population object	Load a population from file
`read_input_data`	- file names - type of data (vcf or plink) - chosen chromosome - number of snps	- genomeadmixr_data object	Read sequencing data from file
`write plink`	- population object - marker locations - file name prefix - recombination rate		Writes output in PLINK format to file

741