Effect of high proportion concentrate dietary on Yak jejunal structure, physiological function

and protein composition during cold season

Jianlei Jia [a,b], Chunnian Liang [a], Xiaoyun Wu [a], Lin Xiong [a], Pengjia Bao [a], Qian Chen [b], Ping

Yan [a*]

a Key Laboratory of Yak Breeding Engineering Gansu Province, Lanzhou, Institute of Husbandry and

Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences, Lanzhou 730050, China;

b Key of laboratory of Plateau Ecology and Agriculture, Qinghai University, Xining 810016, China;

* Corresponding author. Tel: +86-13239649588; EM: pingyanlz@163.com

Abstract:

The current study aimed to investigate the damage of long-term high concentrate diet feeding pattern on Yak jejunal structure, physiological function and protein composition during cold season. Twelve Datong male Yak (Bos grunniens) with the same age from cold season were randomly selected and slaughtered to determine Yak jejunal digestive enzyme activity, morphology and protein composition on different feeding patterns in Tibetan Plateau. The results showed that Yak jejunum digestive enzyme activity and morphology of grazing reared group were better than those in the intensively reared group. A total of 96 differentially expressed proteins were identified by label-free Mass Spectrometry (MS), which could be concluded to two predominant themes: protein structure and inflammatory response. Nine differentially expressed proteins were correlated in Yak jejunum damage in different feeding patterns. According to our study, feeding patterns resulted in the differences in Yak jejunum physiological function, morphology and protein composition. Our results confirmed that feeding of long-term high concentrate diet could damage the jejunum epithelial morphology and function.

Keywords: Feeding pattern (intensively and grazing), Morphology, Physiological function, Protein composition, Yak, Jejunum

Abbreviations: ME, metabolic energy; CP, crude protein; Ca, Calcium; P, Phosphorus; DM, dry matter; AMS, amylases, LPS, lipases; CLS, carboxymethyl cellulase; CLS, carboxymethyl cellulase

1. Introduction

Yak (*Bos grunniens*) mainly lives above an altitude of 3000 meters in the Tibetan plateau and develops a dogged anti-adversity ability to exist, multiply and produce during the long period of natural selection artificial domestication [1]. Yak production is an economic pillar of the Tibet plateau area. The Tibetan plateau is known for its extremely harsh conditions, characterized by high altitude, severe cold, low air oxygen, strong ultraviolet radiation and short forage growing season. Herbage and nutrients are insufficient to support the livestock during the cold period, especially when raised under traditional grazing management [2].

The modern intensive management pattern had gradually been replaced by traditional grazing patterns in the Tibet plateau area [3]. The dietary concentration was more than 75% under modern intensive reared pattern, Yak bodyweight had a faster daily gain, and herdsman had a higher economic benefit. The ruminant needs an intake of 40%-70% dietary roughage to maintain normal rumen function and gastrointestinal microflora environment [4]. However, long-term feeding of high concentrate diet can eventually have a serious impact on animal health, leading to rumen microorganisms' metabolic disorders, such as accumulation of volatile fatty acids (VFAs) and decrease of pH decreased. Fermentable carbohydrate enters the small intestine through the rumen, causes intestinal acidosis, changes the microbial community structure and destroys the integrity of intestinal epithelial morphology and structure [5].

The dietary nutrients are digested in intestinal digestive enzymes and absorbed into the organism through intestinal epithelium to provide energy for animals [6]. The jejunum is an

important organ of nutrition digestion, absorption and metabolism, and a barrier against

harmful substances. It plays an important role in Yak health. Jejunal congenital barriers can

be damaged by external factors, leading to enteric flora disturbance and decreased production

performance. Yak can graze on the pasture without dietary supplementation all the year in

traditional grazing management, and the Yak body can firm a unique intestinal environment

and digestive function to adapt to plateau grazing conditions. However, the modern intensive

management pattern changes Yak diet habit, and high proportion concentrate diet reduces

feedstuff residence time in Yak rumen and improves the amount of jejunum carbohydrate and

jejunum microbial fermentation level, which has a negative effect on the monolayer structure

of jejunum epithelial cells and damages the intestinal barrier [7,8]. This may be an important

physiological mechanism that a high concentrate diet destroys Yak jejunum structure and

barrier. Here, we explore jejunal barrier dysfunction via long-term high concentrate dietary

during cold season. Twelve Datong male Yak with different feeding patterns (intensively and

grazing) during cold season were randomly selected and slaughtered, then determine Yak

jejunal digestive enzyme activity with ELISA method and jejunal morphology with frozen

section method. Using label-free method to identify proteins that specifically expressed with

different feeding pattern, and we analyzed the signal pathway of target proteins. Collectively,

our results shed new light on the damage of long-term high concentrate diet feeding on Yak

jejunum. Meanwhile, this study laid a technical foundation in Yak intensively reared pattern.

2.Material and Methods

2.1 Ethics Statement

All procedures involving the use of animals were approved by the Animal Care

Committee of Lanzhou Institute of Animal Science & Veterinary Pharmaceutics, Chinese

Academy of Agricultural Sciences, China (QHDX-18-04-07-01). Animal slaughtering was

approved by the National Administration of Cattle Slaughtering and Quarantine regulations

(Qinghai, China).

2.2 Study Site

The study was performed at Haibei Tibetan Autonomous Prefecture, Qinghai Province,

China, situated at northeastern Qinghai-Tibetan Plateau. This area is over 3000 m above sea

level and has a dry cold climate.

2.3 Animals and Diets

Twelve Datong male Yaks (4±0.5 years old) were selected under grazing conditions on

natural grasslands in Haibei Tibetan Autonomous Prefecture. All the Yaks were distributed

into two groups. The Yaks of the experimental group were feed in a roofed shed via an

intensively reared pattern, and the Yaks of the control group were feed in grazing pasture

during the experimental period. The experiment lasted for 180 days from 16th November

2018 to 4th May 2019, and the first 10 d was an adjustment period of experimental group

Yak. The following 170 d was for the data collection period.

Based on the determination of the feed intake of the grazing Datong Yak, the

concentrated feed was prepared based on the recommendations for the Chinese Beef Cattle

Raising Standard (NY/T 815-2004). The ingredients composition and nutritional level of

Yaks of the experimental group are presented in Table 1. The roughage was a mixed ration of

maize silage and oat hay in 1:1. The daily dietary of each Yak contained 4 kg concentrate and

2 kg roughage (DM) during the experimental period. The mixed diet nutrient composition

was determined in the Qinghai University laboratory by 'Feed Analysis and Quality Test

Technology' [9].

2.4 Measurement of Samples

Datong male Yaks were stunned by the captive bolt pistol, and the blood was drained at

the end of the experiment. The jejunum was immediately isolated after slaughtering, and

jejunal contents (10 g) were aliquoted into 2 mL sterile tubes and then conserved at -80°C for

digestive enzyme activity analysis. The activity of amylases (AMS), trypsinases (TRS),

lipases (LPS) and carboxymethyl cellulose (CLS) were determined According to Fan's

method [10] using commercially available TaKaRa kits.

The separated jejunum tissue was divided into two parts. One part of jejunum tissue was

fixed in 4% buffered formaldehyde for 72 hours. The samples were dehydrated under

different concentrations of glucose and embedded in a frozen embedding medium. Tissue

sections (5-7 µm) were cut and stained with hematoxylin and eosin (HE), displaying villous

height, villous width, crypt depth, mucosa thickness and muscular thickness, respectively.

Images were acquired using the DP70 software (Olympus, Nagano, Japan) per transverse

section of jejunum tissue. The slides were observed using BX51 microscopy (Olympus,

Nagano, Japan). Measurements were made using Image Proplus5.1 software. Ten

well-oriented and intact crypt-villus units of each slide were measured in triplicate [11].

Another part was aliquoted into a 1.5 mL centrifugal tube when the jejunum surface

contaminants were removed by washing with phosphate-buffered saline solution and

conserved at -80℃ ultracold storage freezer for proteomics and RT-PCR analysis.

Proteomics analysis was conducted by Novogene (China, Beijing). There were 3 replicates in

each group. Jejunum proteins were extracted by lysis buffer method [12]. Digestion of the

protein (250 μg for each sample) was performed according to the FASP procedure. Label-free

Mass Spectrometry (MS) experiments were performed on a Q Exactive mass spectrometer

that was coupled to an Easy nLC system (Thermo Fisher Scientific, MA, USA). The MS data

were analyzed using MaxQuant software and compared with the UniProt Bos taurus

database. Relative quantitative real-time PCR (RT-PCR) was performed to determine the

copy number of target genes. The primers were synthesized at Shanghai Biological

Engineering Ltd., China. The RT-PCR reaction was carried out in a Light Cycler 480

fluorescence quantitative meter (Fritz Hoffmann-La Roche Co. Ltd., Basel, Switzerland).

2.5 Statistical Analysis

Bioinformatics was analyzed via R language toolkit by Novogene (China, Beijing).

Functional annotation and classification of all identified proteins were determined using the

Blast2GO and InterProScan program against the uniport database. Pathway analyses were

extracted using the search pathway tool of the KEGG mapper platform

(http://www.genome.jp/kegg/mapper.html) and BLAST program. Pathway enrichment

statistics were conducted by the Fisher's exact test, and the pathways with a corrected P <

0.05 were defined as the most significant pathways. The STRING program

(http://string-db.org/) for the retrieval of interacting genes/proteins database for the prediction

of the physical and functional interactions was used to analyze the PPIs. The graphical

visualization and analysis of the interaction network were performed in Cytoscape software.

The data were expressed as mean ± standard deviation (SD). Duncan's post hoc test was

used to determine any significant differences between each group. Differences were

considered significant at $P < 0.05$ and extremely significant at $P<0.1$.

3. Results

3.1 Digestive enzyme activity analysis of jejunum

The activity of AMS, LPS, TRS and CLS in different feeding patterns are presented in

Table 2. The data indicated that the AMS and CLS activities of the grazing reared group was

higher than that in the intensively reared group, but there was no significant difference

between the two groups ($P>0.05$); The LPS activity of the intensively reared group was

significantly greater than that in grazing reared group ($P<0.05$). The TRS activity of the

intensively reared group was significantly lower than that in the grazing reared group

($P<0.05$). The Yak jejunum digestive enzyme activity of the grazing reared pattern was

higher than the intensively reared pattern.

3.2 Histological analysis of jejunum morphology

The villous height, villous width, crypt depth, mucosa thickness and muscular thickness

in grazing reared and intensively reared groups are summarized in Table 3. The villous

height, villous width, mucosa thickness and muscular thickness of jejunum were significantly

greater in the grazing reared group than those in the intensively reared group ($P<0.05$).

However, the control group's crypt depth was lower than that in the intensively reared group,

but no significant difference was observed between those two groups ($P>0.05$).

3.3 Changes in proteome profiles during different feeding pattern

    We successfully identified 1208 proteins with Bos taurus database matching of Mascot

and label-free method. Then we applied a manual thresholding approach and a probabilistic

prediction algorithm, yielding 1066 high-confidence candidates. Of these 1066 proteins, 96

differentially expressed proteins were identified between the intensively reared group and

grazing reared group using $P < 0.05$ or a quantitative ratio of > 1.2 or < 0.83 (Table S1).

There were 17 up-regulated and 79 down-regulated differentially expressed proteins in the

grazing reared group compared to that in the intensively reared group (Fig. 1).

3.3 Bioinformatics analyses

To explore biological functions associated with differentially expressed proteins of Yak

jejunum during different feeding patterns, enrichment analysis in the Gene Ontology (GO)

containing cellular components (CC), molecular function (MF) and biological process (BP)

was performed (Fig. 2). The major enriched terms of differentially expressed proteins were

membrane (GO:0016020), cytoplasm (GO:0005737) and mitochondrion (GO: 0005739;

GO:0005740) for cellular components. Other enriched terms were molecular function in

oxidoreductase activity (GO:0016491), cell-cell adherens junction (GO:0005525), adenylate

kinase activity (GO:0004017), protein binding (GO:0019904) and structural molecule activity

(GO:0003954). Apoptosis (GO: 0048468), inflammatory response (GO:0016020), cellular

response to fatty acid (GO:0008152), actin cytoskeleton reorganization (GO:0055114) and

adenylate cyclase-inhibiting G protein-coupled receptor (GO:0046034), which were enriched

in biological processes.

The major pathways associated with differentially expressed proteins of Yak jejunum

during grazing reared group and intensively reared group were identified using KEGG

pathway analysis via KAAS software. The KEGG results showed that out of a total of 71

pathways, 49 were significantly enriched (p-value lower than 0.05).The top 6 of 20

significantly enriched pathways, ranked by significance and percent overlap are tight junction

signaling pathway (ko04530), NF-kappa B signaling pathway (ko04064), MAPK signaling

pathway (ko04010), cytokine-cytokine receptor interaction (ko04060), calcium signaling

pathway (ko04020) and signaling pathways regulating pluripotency of stem cells (ko04550,

Fig. 3). The results of the KEGG pathway analysis matched the results of the GO analysis.

To gain information on interaction network, possible interactions among the 96 proteins

were analyzed online using the Uniport database. As expected, the 96 target proteins

constituted a complex and strong PPi network, and those results of this analysis identified

two predominant themes: protein structure and inflammatory response. To provide further

insights into the biological processes identified by this approach, we took Fishers' test

(Significance A/B Test) for target terms, those terms with the highest number of proteins

from our results show the proteins that result in the identification of an enriched biological

process. For example, the identification of the "Tight junction" (Occludin, Claudin, Zo-1,

ATP1B3 and HKDC1) and "Inflammatory reaction" (GPR41, IL-6, TNF-α and CCL5) terms

results in our list of differentially abundant proteins (Fig. 4).

3.5 Validation of differentially expressed proteins coding genes by RT-PCR

As shown in Fig. 5, the gene expressions of OCLN, CLDN1, TJP1 and HKDC1 were

higher in the grazing reared group than those in the intensively reared group ($P<0.05$) via

RT-PCR, and the ATP1B3 expression had no significant difference between the two groups.

However, the gene expressions of IL6, GPR41, CCL5 and TNF were higher in the intensively

reared group than those in the grazing reared group ($P<0.05$). In summary, the results of

selected differentially expressed proteins coding genes by RT-PCR were the same expression

tendency of the label-free analysis. The RT-PCR assay illustrated that the label-free results

were reliable for further analyses.

4. Discussion

Jejunum is the main organ for dietary nutrition digestion, absorption and metabolism,

and plays an important role in Yak health. The dietary concentrate is more than 75% under

modern intensive management pattern to enhance daily gain and performance. However,

long-term high concentrate diet feeding has a serious impact on animal health, such as jejunal

acidosis [4]. The level of dietary nutrition digestion, absorption and metabolism can be

reflected through digestive enzyme activity. Starch is the main energy source for ruminants.

AMS can hydrolyze starch to glucose, maltose and oligosaccharides, which acts on the

4-c-D-glycosidic bonds of the starch molecule [13]. Fat is the necessary material for energy

storage and nutrition in the animal body, and the animal essential fatty acids are mainly

provided by dietary fat. LPS can digest dietary fat to absorbable nutrients, such as free fatty

acid, glycerin and monoglyceride. TRS can hydrolyze dietary proteins to small molecular

amino acids, and those amino acids are involved in the pathway of intracellular signaling

transmission and apoptosis. It plays a vital role in animal immune response [14]. The ruminant

cellulose-degrading is mainly dependent on the cellulase secreted by microorganisms, and

CLS activity is an important ability to degrade cellulose [15]. In our study, Yak jejunum

digestive enzyme activity can be in a weakened state with long-term high concentrate diet

feeding. We observed that the digestion and absorption of grazing reared patterns were better

than the intensively reared group.

The morphophysiological variations in the jejunal epithelium of ruminants identified the

ability of dietary nutrients digestion and absorption [11]. Changes in the structure of villous

absorptive surface (height, width and crypt) in jejunal epithelial were closely related to the

extent of enterocyte development, which were direct representations of the quality of the

intestinal environment as the indicator of intestinal health [16]. The present study showed that

the ability of dietary nutrients digestion and absorption increased with the increase of villous

absorptive surface, and submucosa and muscular thickness of jejunum were decreased with

long-term high concentrate diet, decreasing the absorption of nutrients in Yak jejunum. When

the high concentrate diet was fed for a long-term, many nutrients were degraded into

short-chain fatty acids (SCFAs) by microorganisms in the rumen, leading to an increase in

the jejunum VFAs and a decrease in the jejunum pH value. VFAs had a strong lipid solubility

when the pH value dropped below the critical value. VFAs can infiltrate into jejunum mucosa

cells to acidify cells, damaging the intestinal barrier and causes an inflammatory response

[17,18]. We found that the significantly decreased resulted in the intensively reared group of

Yak jejunum, such as villous height, villous width, mucosa thickness and muscular thickness.

These results agreed with previous studies in ruminants.

SCFAs in rumen permeate into Yak jejunum, destroy the structural integrity of

morphophysiological, and damage epithelial barrier function [19]. Jejunal mechanical barrier

plays a protective role in jejunum via the normal morphological structure of jejunal mucosa

[20-22]. Tight junction proteins are the important parts of the jejunum mechanical barrier

composed of the transporter protein family, including occludin and claudin, which are mainly

involved in enterocytes proliferation, differentiation, apoptosis and cell bypass permeability

regulation. The damaged mechanical barrier induced jejunal dysfunction causes pathogens to

infect enterocytes, which results in tight junction structural damage, permeability increasing

and inflammatory response [19]. According to recent reports and our study, target proteins can

be concluded to two predominant themes: protein structure (tight junction signaling pathway,

calcium signaling pathway and signaling pathways regulating pluripotency of stem cells) and

inflammatory response (MAPK signaling pathway, chemokine signaling pathway and

cytokine-cytokine receptor interaction). GPR 41, a key activator of the jejunal mechanical

barrier, participates in regulating body inflammatory response and affects the tight junction

signaling pathway [23]. The large concentrate intake can start enterocytes gluconeogenesis

process, activates the tight junction signaling pathway and regulates downstream related cell

apoptosis pathways, decreasing the tight junction proteins expressions, such as occludin,

claudin and Zo-1. Meanwhile, the proliferation of regulatory T cells promotes the

inflammatory response pathway (MAPK) related gene expression to secrete many

anti-inflammatory cytokines (IL-6, TNF-α and CCL5) [24-26]. GPR41 was detected in the

current study. Long-term high concentrate diet feeding changes the normal jejunal metabolic

pathways. Several identified candidate proteins were identified by label-free MS, followed by

qPCR, including occludin, claudin, Zo-1, ATP1B3, HKDC1, IL-6, TNF-α and CCL5.

Compared with the grazing reared group, the expressing quantity of occludin, claudin, Zo-1,

ATP1B3 and HKDC1 in Yak jejunum presented significantly down-regulated in the

intensively reared group. However, IL-6, TNF-α and CCL5 showed a contrary tendency,

indicating that many SCFAs in the jejunum of the high-concentrate diet could damage the

mechanical barrier and reduce the differentially expressed proteins. At the same time, GPR41

was activated to up-regulate the expression and promoted the expression of the

proinflammatory factors. In summary, the results provide evidence of the validity of the

damage of the high concentrate diet on Yak jejunum.

5. Conclusions

We found that feeding pattern results in Yak different jejunum physiological function,

morphology and protein composition. Long-term high concentrate diet feeding can damage

the jejunum epithelial morphology and structure. Therefore, we suggest that the high

concentrate diet is suitable for the short-term fattening pattern of reserved Yak.

Acknowledgments

The authors are grateful to all the participants who took part in this study.

This research was funded by the Agricultural Science and Technology Innovation

Program (CAAS-ASTIP-2014-LIHPS-01), the National Beef Cattle Industry Technology &

System (CARS-38), National Natural Science Foundation of China (31972561).

References

1. Jia Jianlei, Qian Chen, Jing Zhao, et al (2018). Differential Expression of Proteins in

Datong Yak and Chaidamu Yellow Cattle Longissimus lumborum Muscles and Relation to

Meat Water Holding Capacity[J]. Kafkas Univ Vet Fak Derg. 24(5): 691-700.

2. Duane E. Ullrey, Charlotte Kirk Baer and Wilson G (2011). Pond. Encyclopedia of Animal

science[M]. Taylor & Francis Group, LLC. P1123-1124.

3. Long RJ, Ding ML, Guo XH, et al (2008). The yak grazing system on the Qinghai-Tibetan

plateau and its status[J]. Journal of the Australian Rangeland Society. 30(2): 241-246.

4. Huang Xiaodan, Huiyin Tan, André-Denis G Wright, et al (2012). Comparison of

methanogen diversity of yak (Bos grunniens) and cattle (Bos taurus) from the

Qinghai-Tibetan plateau, China[J]. BMC Microbiology. 237(12):1-10.

5. Wang Jinbao, Rui Zhang, Lu Zhang, et al (2015). Histomorphometric Adaptation of Yak

(Bos grunniens) Abomasum to the Qinghai-Tibetan Plateau Environment[J]. Int. J.

Morphol. 33(2):764-776.

6. Cao DX, Wang SL, Hao LZ, et al (2019). Changes of β-carotene and retinol levels and

BCO1 gene and protein expressions in yak tissues at different nutritional seasons[J].

Journal of Animal and Feed Sciences. 28:120-127.

7. Zhang Lihong, Xiong Jiang, Jiakui Li, et al (2020). Characterization of the microbial

community structure in intestinal segments of yak (Bos grunniens) [J]. Anaerobe. 61:

102-115.

8. Moharrery A, Brask M, Weisbjerg MR (2014). The effect of different physical forms of

rapeseed as a fat supplement on the activity of some enzymes in the duodenal chyme of

dairy cows [J]. Journal AnimalFeed Science. 23(4): 293-298.

9. Yang S (1993). Feed Analysis and Quality Test Technology [M]. Beijing Agriculture

Press: Beijing.

10. Fan Xiaohong, Lizhuang Hao, Shujie Liu, et al (2017). Effect of different additions of

canola rapeseed on the activity of digestive enzymes in jejunum and chyme[J]. Chinese

Feed. 18(4):15-23.

11. Ding Baoan, Shuangquan Ma, Stephen R. Madigosky, et al (2018). Seasonal changes of

rumen and intestine morphology of the Qinghai yak (Bos grunniens)[J]. Veterinary World.

11(8): 1135-1138.

12. Jia Jianlei, Qiang Ding, Liping Zhang, et al (2014). Establishment of the optimum

two-dimensional electrophoresis system of ovine ovarian tissue[J]. Genet. Mol. Res. 13(3):

6528-6538.

13. Li Aoyun, Yaping Wang, Jiakui Li, et al (2019). Probiotics isolated from yaks improves

the growth performance, antioxidant activity, and cytokines related to immunity and

inflammation in mice[J]. Microb Cell Fact. 18:112-123.

14. Ding Luming, Ruijun Long, Zhanhuan Shang, et al (2008). Feeding behaviour of yaks on

spring, transitional, summer and winter pasture in the alpine region of the Qinghai–Tibetan

plateau[J]. Applied Animal Behaviour Science. 111(3-4):373-390.

15. Liu Yang, Caiyu Yan, Fujiang Hou, et al (2017). Key sources and seasonal dynamics of

greenhouse gas fluxes from yak grazing systems on the Qinghai-Tibetan Plateau[J].

Scientific Reports. 7:1-11.

16. Guo Wei, Weiwei Wang, Ying Zhang, et al (2020). Characterization of Anaerobic Rumen

Fungal Community Composition in Yak, Tibetan Sheep and Small Tail Han Sheep Grazing

on the Qinghai-Tibetan Plateau[J]. Animals. 10(1):144-156.

17. Xue Dan, Huai Chen, Fang Chen, et al (2016). Analysis of the rumen bacteria and

methanogenic archaea of yak (Bos grunniens) steers grazing on the Qinghai-Tibetan

Plateau[J]. Livestock Science. 188:61-71.

18. Peng YS, M.A.Brown, Wu JP (2008). Fatty Acid Profile in Milk Fat from Qinghai

Plateau Yak at Different Altitudes and Parities[J]. The Professional Animal Scientist. 24(5):

479-487.

19. Peng Along, Yaotian Fan, Hongrong Wang, et al (2020). Effects of Thiamine on

Maintaining Rumen Epithelial Barrier Function in Ruminants[J]. Chinese Journal of

Animal Nutrition. 2020, 32(6):1-6.

20. Xue Chunxu (2017). The impact of high concentrate diet on small intestinal fermentation,

microbial community and epithelial morphology of goats[D]. Nanjing Agricultural

University.

21. Ma Li, Shixiao Xu, Hongjin Liu, et al (2019). Yak rumen microbial diversity at different

forage growth stages of an alpine meadow on the Qinghai-Tibet Plateau[J]. PeerJ. 7:1-27.

22. Li Yang (2019). Effects of subacute rumen acidosis on the absorption of volatile fatty

acids in rumen epithelium of dairy goats and its mechanism research[D]. INNER

MONGOLIA AGRICULTURAL UNIVERSITY.

23. Zhang HM, Wang Y, Chang GJ, et al (2018). Effect of high-concentrate diet on the

inflammatory response mediated by short-chain fatty acids-GPR41/43 in cecum of lactating

goats[J]. Animal Husbandry&Veterinary Medicine. 50(2):61-66.

24. Mamiko Kobayashi, Daisuke Mikami, Junsuke Uwada, et al (2018). A short-chain fatty

acid, propionate, enhances the cytotoxic effect of cisplatin by modulating GPR41 signaling

pathways in HepG2 cells[J]. Oncotarget. 9(59): 31342-31354.

25. Mamiko Kobayashi, DaisukeMikami, HidekiKimura, et al (2017). Short-chain fatty acids,

GPR41 and GPR43 ligands, inhibit TNF-α-induced MCP-1 expression by modulating p38

and JNK signaling pathways in human renal cortical epithelial cells[J]. Biochemical and

Biophysical Research Communications. 486(2):499-505.

26. Li Dinghua, Liu Chiman, Luo Ruibang, et al (2015). MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph[J].

Bioinformatics. 31(10):1674-1676.

Table 1

Ingredients and proximate nutrition content of concentrate

Composition	Content (%)	Nutritional composition	Content
Maize	67.0	ME [b]	3.01 Mcal/kg
Soybean meal	5.0	CP	14.88 %DM
Cottonseed meal	10.0	Ca	0.66 %DM
Wheat bran	14.0	P	0.43 %DM
Salt	0.5		
Sodium Bicarbonate	1.5		
Stone Powder	1.5		
Premix Feed [a]	0.5		

379   [a] The premix provided the following per kg of diets：VA 12000IU, VD 20 00IU, VE 30IU, Cu 12mg, Fe

380   64mg, Mn 56mg, Zn 60mg, I 1.2mg, Se 0.4mg, Co 0.4mg;

381   [b] ME was calculation value (NRC,2007), others were measured value.

382　Table 2

383　The alkaline phosphatase activity of yak jejunum in different feeding pattern

	AMS	LPS	CLS	TRS
grazing group	0.176±0.043	0.218±0.072[a]	4.102±0.835	70.642±7.206[a]
Intensively group	0.121±0.071	0.542±0.053[b]	2.145±0.332	36.109±3.058[b]

384　Note. Different letters represent significant differences (P<0.05), and the following is the same.

385 Table 3

386 The villous height, villous width, crypt depth, mucosa thickness and muscular thickness of yak jejunum

387 in different feeding pattern

	villous height	villous width	crypt depth	mucosa thickness	muscular thickness
grazing group	563.33 ± 11.48^a	149.29 ± 2.67^a	178.84 ± 6.59	808.38 ± 15.52^a	528.91 ± 20.93^a
intensively group	408.31 ± 10.46^b	82.88 ± 5.45^b	194.61 ± 8.37	939.18 ± 14.14^b	751.51 ± 15.14^b

388 Note. Different letters represent significant differences ($P<0.05$), and the following is the same.

420   Fig 1.   Differentially expressed proteins, including 3 replicates with yak jejunum during grazing reared

421   group (A) and intensively reared group (B). The image presents the relative abundance of proteins using

422   different colors, where deeper red represents higher intensity and blue represents lower intensity

442   Fig.2 Enrichment GO terms Analysis

474   Fig.3 Enrichment KEGG pathways Analysis (Fisher's Exact Test, P value < 0.01)

505 Fig.4 Protein-protein interaction networks of the differential abundance proteins of different feeding
506 groups in Yak jejunum based on Cytoscape software. The nodes were proteins from Bos taurus database
507 and the lines were the predicted functional annotations

537 Fig.5 Effects of different feeding pattern on the differentially expressed proteins coding genes in yak
538 jejunum during intensively reared group (A) and grazing reared group (B).

539
540