Jejunal barrier dysfunction of Datong Yak with high proportion concentrate dietary during the cold season

Jianlei *Jia* [1,2], Chunnian *Liang* [1], Xiaoyun *Wu* [1], Lin *Xiong* [1], Pengjia *Bao* [1], Qian *Chen* [2], Ping *Yan* [1]*

* Corresponding author：Ping Yan

Tel: +86-13239649588; E-Mail: pingyanlz@163.com

1 Key Laboratory of Yak Breeding Engineering Gansu Province, Lanzhou, Institute of Husbandry and

 Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences, Lanzhou 730050, China;

2 Key of laboratory of Plateau Ecology and Agriculture, Qinghai University, Xining 810016, China;

ORCID

Jianlei *Jia*   https://orcid.org/0000-0001-5843-1709

Chunnian *Liang*   https://orcid.org/0000-0002-4592-9206

Xiaoyun *Wu*   https://orcid.org/0000-0002-9168-6663

Lin *Xiong*   https://orcid.org/0000-0002-4369-1804

Pengjia *Bao*   https://orcid.org/0000-0003-3596-7804

Qian *Chen*   https://orcid.org/0000-0002-8108-4642

Ping *Yan*   https://orcid.org/0000-0002-4040-7038

Title of the manuscript: High proportion concentrate-fed for Datong Yak Jejunal barrier

dysfunction during the cold season

ABSTRACT

Objective: Ruminant intestine health played a vital role in nutrient absorption and metabolism.

Methods: In present study, twelve Datong male Yak were randomly selected and slaughtered to

determine jejunal digestive enzyme activity, morphology and protein composition between forage-fed

and concentrate-fed during cold season in Tibetan Plateau.

Results: The results showed that Yak jejunum digestive enzyme activity and morphology of forage-fed

group were better than that in concentrate-fed group. A total of 96 differentially expressed proteins were

identified by label-free MS, which could be concluded to two predominant themes: protein structure

and inflammatory response. Nine differentially expression proteins were correlated in yak jejunum

damage in high dietary concentrate.

Conclusion: This fact was confirmed long-term high dietary concentrate feeding could damage the

jejunum epithelial morphology and function.

Keywords: Feeding pattern (forage-fed and concentrate-fed), Morphology, Physiological function,

Protein composition, Yak (Bos grunniens), Jejunum

INTRODUCTION

Yak (Bos grunniens) mainly live above an altitude of 3000 meters in the Tibetan plateau, and develop a dogged anti-adversity ability to exist, multiply and produce during the long period of natural selecting and artificial domestication [1]. Yak feeding is an economic pillar of Tibet plateau area. Tibetan plateau is known as extreme harsh conditions, characterized by high altitude, severe cold, low air oxygen, strong ultraviolet radiation and short forage growing season. Herbage and nutrients are often insufficient to support the livestock during the cold period, especially when raised under traditional grazing management [2]. The saying 'alive in summer, strong in autumn, thin in winter, tired in spring' has been passed down from generation to generation by the local people.

The modern intensive management pattern recently had gradually replaced to traditional grazing pattern in Tibet plateau area [3]. The dietary concentrate was more than 75% under modern intensive reared pattern, Yak bodyweight had a faster daily gain, and herdsman had a higher economic benefit. The ruminant need intake 40%-70% dietary roughage to maintain normal rumen function and gastrointestinal microflora environment [4]. However, long-term high dietary concentrate feeding can eventually have a serious impact on animal health, which could lead to rumen microorganisms metabolic disorder (such as VFAs accumulated and pH decreased), then fermentable carbohydrate entered small intestine through rumen, caused intestinal acidosis, changed the microbial community structure, and destroyed the integrity of intestinal epithelial morphology and structure [5].

The dietary nutrients were digested in intestine digestive enzymes, and absorbed into organism through intestinal epithelium to provide energy for animals [6]. Jejunum was an important organ of nutrition digestion, absorption and metabolism, also an important barrier against harmful substances, played an important role in yak health. The research showed that jejunal congenital barriers was easy to be damaged by the influence of external factors, which leaded to enteric flora disturbance, decrease the production performance. Yak were grazing on the pasture without supplementary dietary all the year in traditional grazing management, and yak body had formed a unique intestinal environment and digestive function to adapt to plateau grazing conditions. However, the modern intensive management pattern changed yak diet habit, high proportion concentrate dietary reduced feedstuff residence time in

yak rumen, improved the amount of jejunum carbohydrate and jejunum microbial fermentation level, which had a negative effect on the jejunum epithelial cells monolayer structure and damage the intestinal barrier [7,8]. This may be an important physiological mechanism that high proportion concentrate dietary destroy yak jejunum structure and barrier. Here, twelve Datong male Yak with different feeding patterns (forage-fed and concentrate-fed) during cold season were randomly selected and slaughtered to determine Yak jejunal digestive enzyme activity, morphology and protein composition. Collectively, our results shed new light on the damage of long-term high dietary concentrate feeding on Yak jejunum. Meanwhile, this study laid a technical foundation in yak organism health associated with dietary structure.

MATERIALS AND METHODS

All procedures involving the use of animals were approved by the Animal Care Committee of Lanzhou Institute of Animal Science & Veterinary Pharmaceutics, Chinese Academy of Agricultural Sciences. Animal slaughtering was approved by the National Administration of Cattle Slaughtering and Quarantine regulations (Qinghai, China).

Study Site

The study was performed at Haibei Tibetan Autonomous Prefecture, Qinghai Province of China, situated at northeastern Qinghai-Tibetan Plateau. This area is over 3000m above sea level and has a dry cold climate.

Animals and Diets

Twelve Datong male Yaks (4±0.5 years old) were selected under grazing conditions on natural grasslands in Haibei Tibetan Autonomous Prefecture. All the Yak were equally distributed into 2 groups. The experimental group Yak were feed in a roofed shed via intensively reared pattern, and the control group Yak were feed in grazing pasture during the experimental period. The experiment lasted 180 days from 16th November, 2018 to 4th May, 2019, and the first 10d was an adjustment period of experimental group Yak and the following 170d was for data collection period.

Based on the determination of the feed intake of the grazing Datong Yak, we prepared the concentrated feed based on the recommendations for Chinese Beef cattle Raising Standard (NY/T 815-2004). The

ingredients composition and nutritional level of experimental group Yak feed was presented in Table

1. The roughage was mixed ration of maize silage and oat hay in 1:1. The daily dietary of each yak

contained with 4kg concentrate and 2kg roughage (DM) during the experimental period. The mixed diet

nutrient composition was determined in Qinghai University laboratory by 'Feed Analysis and Quality

Test Technology' [9].

Measurement of Samples

Datong male yaks were stunned by captive bolt pistol and their blood was drained at the end of

experiment. The jejunum was immediately isolated after slaughtering, and approximately 10g of jejunal

contents were aliquoted into 2mL sterile tubes and then conserved at -80°C ultracold storage freezer for

digestive enzyme activity analysis. The activity of Amylases (AMS), Trypsinases (TRS), Lipases (LPS)

and Carboxymethyl cellulase (CLS) were determined According to Fan's method used commercially

TaKaRa kits [10].

The separated jejunum tissue was divided two parts. One part for histological examination were fixed

in 4% buffered formaldehyde for 72 hours, then the samples were dehydrated under different

concentration of glucose, embedded in frozen embedding medium, and serial 5-7 μm transverse sections

were cut and stained with hematoxylin and eosin (HE), which displayed villous height, villous width,

crypt depth, mucosa thickness and muscular thickness, respectively. Images were acquired using the

DP70 software (Olympus, Nagano, Japan) per transverse section of jejunum tissue. The slides were

observed using BX51 microscopy (Olympus, Nagano, Japan). Measurements were made using Image

Proplus5.1 software. 10 well-oriented and intact crypt-villus units of each slide were measured in

triplicate.

Other part was aliquoted into 1.5mL centrifugal tube, when the jejunum surface contaminants were

removed by washing with phosphate buffered saline solution, and then conserved at-80℃ ultracold

storage freezer for proteomics and qPCR analysis. proteomics analysis was conducted by Novogene

(China, Beijing). There were 3 replicates each group. Jejunum proteins were extracted by lysis buffer

method (7M Urea, 2M Thiourea, 4% CHAPS(w/v), 100 mM DTT, 60 mM Tris-HCl, 0.5% pH 3-10

Ampholine, 20μL proteinase inhibiors) [11]. Digestion of the protein (250 μg for each sample) was

performed according to the FASP procedure described. Label-free Mass Spectrometry (MS)

experiments were performed on a Q Exactive mass spectrometer that was coupled to an Easy nLC

system (Thermo Fisher Scientific, MA, USA). The MS data were analyzed using MaxQuant software

and compared with the UniProt Bos taurus database. Relative quantitative real-time PCR (qPCR) was

performed to determine the copy number of target genes. Shanghai Biological Engineering Ltd.

synthesized the primers. The qPCR reaction was carried out in a Light Cycler 480 fluorescence

quantitative meter (Fritz Hoffmann-La Roche Co. Limited, Basel, Switzerland).

Statistical Analysis

Functional annotation and classification of all identified proteins were determined by using the

Blast2GO program against the uniport database, including cell component, biological process, and

molecular function. Pathway analyses were extracted using the search pathway tool of the KEGG

mapper platform (http://www.genome.jp/kegg/mapper.html). Pathway enrichment statistics were

conducted by Fisher's exact test, and the pathways with a corrected p value < 0.05 were defined as the

most significant pathways. The STRING program (http://string-db.org/) for the retrieval of interacting

genes/proteins database for the prediction of the physical and functional interactions was used to

analyze the PPi. The graphical visualization and analysis of interaction network was performed in

Cytoscape software.

The results of the analysis were given as mean ± SD. Duncan's post hoc test was used to determine any

significant differences between groups. Differences were considered as significant at $P < 0.05$ and as a

trend at $P<0.1$.

RESULTS

Digestive enzyme activity analysis of jejunum

The activity of AMS, LPS, TRS as well as CLS in different feeding patterns were presented in Table 2.

The data indicated that the AMS and CLS activity of forage-fed group was higher than that in

concentrate-fed group, but no significant difference between those two groups ($P>0.05$); The LPS

activity of concentrate-fed group was significantly greater than that in forage-fed group ($P<0.05$); The

TRS activity of concentrate-fed group was significantly lower than that in forage-fed group ($P<0.05$).

In contrast, the Yak jejunum digestive enzyme activity of forage-fed pattern was better than concentrate-

fed pattern.

Histological analysis of jejunum morphology

The villous height, villous width, crypt depth, mucosa thickness and muscular thickness during forage-

fed group and concentrate-fed group were summarized in Table 3. The villous height, villous width,

mucosa thickness and muscular thickness of jejunum were significantly greater in forage-fed group

compared to that in concentrate-fed group ($p<0.05$). however, the crypt depth of control group was

lower than that in concentrate-fed group, but no significant difference between those two groups

($P>0.05$).

Changes in proteome profiles during different feeding pattern

168　The changes proteins of yak jejunum during different feeding pattern were investigated using label-free

169　MS. A total of 96 differentially expressed proteins were identified between concentrate-fed group and

170　forage-fed group as determined using $P < 0.05$ and a quantitative ratio of > 2 or < 0.5. There were 17

171　up-regulated and 79 down-regulated differentially proteins in forage-fed group compared to that in

172　concentrate-fed group (Fig.1).

Bioinformatics analyses

To explore biological functions associated with differentially expressed proteins of yak jejunum during

different feeding pattern, enrichment analysis in the Gene Ontology (GO) contained cellular

components (CC), molecular function (MF) and biological process (BP) was performed (Fig. 2). The

major enriched terms of differentially expressed proteins were membrane (GO:0016020), cytoplasm

(GO:0005737) and mitochondrion (GO:0005739, GO:0005740) for cellular components. Differentially

expressed proteins were enriched of molecular function in oxidoreductase activity (GO:0016491), cell-

cell adherens junction (GO:0005525), adenylate kinase activity (GO:0004017), protein binding

(GO:0019904) and structural molecule activity (GO:0003954). Apoptosis (GO:0048468),

Inflammatory Response (GO:0016020), Cellular response to fatty acid (GO:0008152), Actin

cytoskeleton reorganization (GO:0055114) and Adenylate cyclase-inhibiting G protein-coupled

receptor (GO:0046034) were enriched in biological processes.

The major pathways associated with differentially expressed proteins of yak jejunum during

concentrate-fed group and forage-fed group were identified using KEGG Pathway Analysis via KAAS

software. 96 differentially expressed proteins across 20 pathways were significantly enriched ($p < 0.05$).

The significantly enriched pathways were Tight junction signaling pathway (ko04530), NF-kappa B

signaling pathway (ko04064), MAPK signaling pathway (ko04010), Cytokine-cytokine receptor

interaction (ko04060), Calcium signaling pathway (ko04020), and Signaling pathways regulating

pluripotency of stem cells (ko04550, Fig. 3). The results of KEGG pathway analysis matched the results

of GO analysis.

To explore the information of differentially expressed proteins of yak jejunum during concentrate-fed

group and forage-fed group network, connected proteins among the 96 proteins were analyzed via

STRING program and Cytoscape software. As showed, the results identified two predominant themes:

protein structure (Occluding, Claudin, Zo-1, ATP1B3 and HKDC1) and inflammatory response

(GPR41, IL-6, TNF-α and CCL5) constituted a complex and strong PPi network (Fig. 4).Therefore,

these target proteins could be regarded as important for jejunal structure, physiological function and

protein composition.

Validation of differentially expressed proteins coding genes by qPCR

As shown in Fig.5, the gene expression of OCLN, CLDN1, TJP1 and HKDC1 were higher in forage-

fed group than that in concentrate-fed group ($P<0.05$) via RT-PCR, and the ATP1B3 expression was

no significant difference between two groups, however, the gene expression of IL6, GPR41, CCL5 and

TNF were higher in concentrate-fed group than that in forage-fed group ($P<0.05$). In summary, the

results of selected differentially expressed proteins coding genes by RT-PCR were the same expression

tendency of the label-free analysis. The RT-PCR assay illustrated that the label-free results were reliable

for further analyses.

DISCUSSION

Jejunum was the main organ for dietary nutrition digestion, absorption and metabolism, and played an

important role in yak health. The dietary concentrate was more than 75% under modern intensive

management pattern to enhance daily gain and performance, However, long-term high dietary

concentrate feeding can eventually have a serious impact on animal health, such as jejunal acidosis [12].

The level of dietary nutrition digestion, absorption and metabolism could be reflected through digestive

enzyme activity. Starch was the main energy source for ruminants, and AMS could hydrolyze starch to

glucose, maltose and oligosaccharides, which acted on the 4-c-D-glycosidic bonds of starch molecule

[13]. Fat was the necessary materials of energy storage and nutrition in animal body, and the animal

essential fatty acids were mainly provided by dietary fat. LPS could digest dietary fat to absorbable

nutrition, such as free fatty acid, glycerin and monoglyceride. TRS could hydrolyze dietary proteins to

small molecular amino acids, those amino acids were involved in the pathway of intracellular signaling

transmission and apoptosis, and played an important role in animal immune response [14]. The ruminant

cellulose degrading mainly depended on the cellulase secreted by microorganism, and CLS activity was

an important reflection of ability to degrade cellulose [15]. In our research, yak jejunum digestive enzyme

activity would be in a weakened state with long-term high dietary concentrate feeding, we believed that

the digestion and absorption of forage-fed pattern was better than concentrate-fed group.

The morphophysiological variations in jejunal epithelium of ruminants measured the ability of dietary

nutrients digestion and absorption [16]. Changes in the structure of villous absorptive surface (height,

width and crypt) in jejunal epithelial was closely related with extent of enterocytes development, which

were direct representation of the quality of intestinal environment, and often be used as the indicator of

intestinal health [17]. The present research showed that the ability of dietary nutrients digestion and

absorption increased with increasing villous absorptive surface, and submucosa thickness and muscular

thickness of jejunum were decreased with long-term high dietary concentrate, decreasing the absorption

of nutrients in Yak jejunum. When the high concentrate dietary was be fed for a long-term, a large

number of nutrients were degraded into short-chain fatty acids (SCFAs) by microorganisms in rumen,

which lead to increase the jejunum volatile fatty acids (VFAs), and decrease the jejunum pH value.

VFAs had a strongly lipid solubility, when the pH value dropped below the critical value, VFAs would

infiltrate into jejunum mucosa cells to acidify cells, which damaged the intestinal barrier and caused

the inflammatory response [18]. In our study, we found that the significantly decreased results in

concentrate-fed group of yak jejunum villous height, villous width, mucosa thickness and muscular

thickness. these results agreed with that reported in the ruminant intestine when fed with higher

concentrate feedstuff.

When SCFAs in rumen permeated into Yak jejunum, which would destroy the structural integrity of

morphophysiological and damage epithelial barrier function [19]. Jejunal mechanical barrier played a

mechanical protective role in jejunum via the normal morphological structure of jejunal mucosa [20-22].

Tight junction proteins were the important part of jejunum mechanical barrier, composed of transporters

protein family, including Occludin and Claudin, which were mainly involved in enterocytes

proliferation differentiation, apoptosis and cell bypass permeability regulation. The damaged

mechanical barrier induced jejunal dysfunction, caused pathogens to infect enterocytes, it resulted in

tight junction structural damage, permeability increasing and inflammatory response [18]. According to

recent reports and our study results presented target proteins can be concluded to two predominant

themes: protein structure (Tight junction signaling pathway, Calcium signaling pathway and Signaling

pathways regulating pluripotency of stem cells) and inflammatory response (MAPK signaling pathway,

Chemokine signaling pathway and Cytokine-cytokine receptor interaction). GPR 41, a key activator of

jejunal mechanical barrier, participated in the regulation of body inflammatory response, and affect the

tight junction signaling pathway as a nutrient energy regulatory receptor [23]. The large concentrate

intake could start-up enterocytes gluconeogenesis process, activated the tight junction signaling

pathway, then regulated downstream related cell apoptosis pathways, which decreased the tight junction

proteins expressions, such as Occluding, Claudin and Zo-1. Meanwhile, the proliferation of Regulatory

T Cells to secrete many anti-inflammatory cytokines (IL-6, TNF-α, CCL5), promoted the inflammatory

response pathway (MAPK) related genes expression [24,25]. GPR41 was detected in the current study,

and presented significantly up-regulated in concentrate-fed group, it showed that long-term high dietary

concentrate feeding changed the normal jejunal metabolic pathways. Furthermore, several identified

candidate proteins were identified by label-free MS, followed by RT-PCR: Occluding, Claudin, Zo-1,

ATP1B3, HKDC1, IL-6, TNF-α and CCL5. Compared with forage-fed group, the expressing quantity

of Occluding, Claudin, Zo-1, ATP1B3 and HKDC1 in Yak jejunum presented significantly down-

regulated in concentrate-fed group, however, IL-6, TNF-α and CCL5 showed a contrary tendency, it

indicated that a large number of SCFAs in jejunum of the high-concentrate dietary could damage the

mechanical barrier, reduce the differentially expression proteins. At the same time, GPR41 was

activated to up-regulate the expression, and promoted the proinflammatory factors expression. In this

research, we found that feeding pattern result in Yak different jejunum physiological function,

morphology and protein composition. This fact was confirmed long-term high dietary concentrate

feeding can damage the jejunum epithelial morphology and structure. Therefore, we hold that the high

proportion concentrate dietary was only suitable for short-term fattening pattern, not cultivation of

reserved yak.

CONFLICT OF INTEREST

275 We certify that there is no conflict of interest with any financial organization regarding the material

276 discussed in the manuscript.

ACKNOWLEDGMENTS

278 The authors are grateful to all the participants who took part in this study.

279 This research was funded by the Agricultural Science and Technology Innovation Program (CAAS-

280 ASTIP-2014-LIHPS-01), the National Beef Cattle Industry Technology & System (CARS-38),

281 National Natural Science Foundation of China (31972561).

REFERENCES

283 1. Jia Jianlei, Qian Chen, Jing Zhao, et al. Differential Expression of Proteins in Datong Yak and
     Chaidamu Yellow Cattle Longissimus lumborum Muscles and Relation to Meat Water Holding
     Capacity[J]. Kafkas Univ Vet Fak Derg. 2018,24(5): 691-700. doi: 10.9775/kvfd.2018.19734

286 2. Duane E. Ullrey, Charlotte Kirk Baer and Wilson G. Pond. Encyclopedia of Animal science[M].
     Taylor & Francis Group, LLC. 2011. P1123-1124.

288 3. Long RJ, Ding ML, Guo XH, et al. The yak grazing system on the Qinghai-Tibetan plateau and its
     status[J]. Journal of the Australian Rangeland Society. 2008, 30(2): 241-246. doi:10.1071/rj08012

290 4. Huang Xiaodan, Huiyin Tan, André-Denis G Wright, et al. Comparison of methanogen diversity of
     yak (Bos grunniens) and cattle (Bos taurus) from the Qinghai-Tibetan plateau, China[J]. BMC
     Microbiology. 2012,237(12):1-10.doi: 10.1186/1471-2180-12-237

293 5. Wang Jinbao, Rui Zhang, Lu Zhang, et al. Histomorphometric Adaptation of Yak (Bos grunniens)
     Abomasum to the Qinghai-Tibetan Plateau Environment[J]. Int. J. Morphol. 2015, 33(2):764-776.
     doi: 10.4067/s0717-95022015000200055

6. Cao DX, Wang SL, Hao LZ, et al. Changes of β-carotene and retinol levels and BCO1 gene and
protein expressions in yak tissues at different nutritional seasons[J]. Journal of Animal and Feed
Sciences. 2019, 28:120-127. doi: 10.22358/jafs/109953/2019

7. Zhang Lihong, Xiong Jiang, Jiakui Li, et al. Characterization of the microbial community structure
in intestinal segments of yak (Bos grunniens) [J]. Anaerobe. 2020, 61: 102-115. doi:
10.1016/j.anaerobe.2019.102115

8. Moharrery A, Brask M, Weisbjerg MR. The effect of different physical forms of rapeseed as a fat
supplement on the activity of some enzymes in the duodenal chyme of dairy cows [J]. Journal
AnimalFeed Science. 2014, 23(4): 293-298. doi: 10.22358/jafs/65664/2014

9. Yang S (1993). Feed Analysis and Quality Test Technology [M]. Beijing Agriculture Press: Beijing.

10. Fan Xiaohong, Lizhuang Hao, Shujie Liu, et al. Effect of different additions of canola rapeseed on
the activity of digestive enzymes in jejunum and chyme[J]. Chinese Feed. 2017, 18(4):15-23. doi:
10.15906/j.cnki.cn11-2975/s.20171804

11. Jia Jianlei, Qiang Ding, Liping Zhang, et al. Establishment of the optimum two-dimensional
electrophoresis system of ovine ovarian tissue[J]. Genet. Mol. Res. 2014, 13(3): 6528-6538. doi:
10.4238/2014.August.26.3

12. Li Aoyun, Yaping Wang, Jiakui Li, et al. Probiotics isolated from yaks improves the growth
performance, antioxidant activity, and cytokines related to immunity and inflammation in mice[J].
Microb Cell Fact. 2019, 18:112-123. doi: 10.1186/s12934-019-1161-6

13. Ding Luming, Ruijun Long, Zhanhuan Shang, et al. Feeding behaviour of yaks on spring,
transitional, summer and winter pasture in the alpine region of the Qinghai–Tibetan plateau[J].
Applied Animal Behaviour Science. 2008, 111(3-4):373-390. doi: 10.1016/j.applanim.2007.06.008

14. Liu Yang, Caiyu Yan, Fujiang Hou, et al. Key sources and seasonal dynamics of greenhouse gas
fluxes from yak grazing systems on the Qinghai-Tibetan Plateau[J]. Scientific Reports. 2017, 7:1-
11. doi: 10.1038/srep40857

15. Ding Baoan, Shuangquan Ma, Stephen R. Madigosky, et al. Seasonal changes of rumen and intestine
morphology of the Qinghai yak (Bos grunniens)[J]. Veterinary World. 2018, 11(8): 1135-1138. doi:
10.14202/vetworld.2018.1135-1138

16. Guo Wei, Weiwei Wang, Ying Zhang, et al. Characterization of Anaerobic Rumen Fungal
Community Composition in Yak, Tibetan Sheep and Small Tail Han Sheep Grazing on the Qinghai-
Tibetan Plateau[J]. Animals. 2020, 10(1):144-156. doi: 10.3390/ani10010144

17. Xue Dan, Huai Chen, Fang Chen, et al. Analysis of the rumen bacteria and methanogenic archaea
of yak (Bos grunniens) steers grazing on the Qinghai-Tibetan Plateau[J]. Livestock Science. 2016,
188:61-71. doi: 10.1016/j.livsci.2016.04.009

18. Peng YS, M.A.Brown, Wu JP. Fatty Acid Profile in Milk Fat from Qinghai Plateau Yak at Different
Altitudes and Parities[J]. The Professional Animal Scientist. 2008, 24(5): 479-487. doi:

10.15232/s1080-7446(15)30890-1

19. Xue Chunxu. The impact of high concentrate diet on small intestinal fermentation, microbial
community and epithelial morphology of goats[D]. Nanjing Agricultural University.2017

20. Ma Li, Shixiao Xu, Hongjin Liu, et al. Yak rumen microbial diversity at different forage growth
stages of an alpine meadow on the Qinghai-Tibet Plateau[J]. PeerJ. 2019, 7:1-27. doi:
10.7717/peerj.7645

21. Li Yang. Effects of subacute rumen acidosis on the absorption of volatile fatty acids in rumen
epithelium of dairy goats and its mechanism research[D]. INNER MONGOLIA AGRICULTURAL
UNIVERSITY. 2019

22. Peng Along, Yaotian Fan, Hongrong Wang, et al. Effects of Thiamine on Maintaining Rumen
Epithelial Barrier Function in Ruminants[J]. Chinese Journal of Animal Nutrition. 2020, 32(6):1-6.
doi: 11.5461.S.20200303.1043.012

23. Zhang HM, Wang Y, Chang GJ, et al. Effect of high-concentrate diet on the inflammatory response
mediated by short-chain fatty acids-GPR41/43 in cecum of lactating goats[J]. Animal
Husbandry&Veterinary Medicine. 2018, 50(2):61-66. doi: 0529-5130(2018) 02-0061-06

24. Mamiko Kobayashi, Daisuke Mikami, Junsuke Uwada, et al. A short-chain fatty acid, propionate,
enhances the cytotoxic effect of cisplatin by modulating GPR41 signaling pathways in HepG2
cells[J]. Oncotarget. 2018, 9(59): 31342-31354. doi: 10.18632/oncotarget.25809

25. Mamiko Kobayashi, DaisukeMikami, HidekiKimura, et al. Short-chain fatty acids, GPR41 and
GPR43 ligands, inhibit TNF-α-induced MCP-1 expression by modulating p38 and JNK signaling
pathways in human renal cortical epithelial cells[J]. Biochemical and Biophysical Research
Communications. 2017, 486(2):499-505. doi: 10.1016/j.bbrc.2017.03.071

368 **Table 1** Ingredients and proximate nutrition content of concentrate

Composition	Content (%)	Nutritional composition	Content
Maize	67.0	ME b	3.01 Mcal/kg
Soybean meal	5.0	CP	14.88 %DM
Cottonseed meal	10.0	Ca	0.66 %DM
Wheat bran	14.0	P	0.43 %DM
Salt	0.5		
Sodium Bicarbonate	1.5		
Stone Powder	1.5		
Premix Feed a	0.5		

369 a The premix provided the following per kg of diets：VA 12000IU, VD 20 00IU, VE 30IU, Cu 12mg, Fe 64mg,

370 Mn 56mg, Zn 60mg, I 1.2mg, Se 0.4mg, Co 0.4mg;

371 b ME was calculation value (NRC,2007), others were measured value.

372   **Table 2** The alkaline phosphatase activity of yak jejunum in different feeding pattern

	AMS	LPS	CLS	TRS
forage-fed	0.176±0.043	0.218±0.072a	4.102±0.835	70.642±7.206a
concentrate-fed	0.121±0.071	0.542±0.053b	2.145±0.332	36.109±3.058b

374 **Table 3** The villous height, villous width, crypt depth, mucosa thickness and muscular thickness of yak

375   jejunum in different feeding pattern

	villous height	villous width	crypt depth	mucosa thickness	muscular thickness
forage-fed	563.33±11.48a	149.29±2.67a	178.84±6.59	808.38±15.52a	528.91±20.93a
concentrate-fed	408.31±10.46b	82.88±5.45b	194.61±8.37	939.18±14.14b	751.51±15.14b

Figure

Figure

Figure

Figure

Figure