Differences in cofactor, oxygen and sulfur requirements influence niche adaptation in deep-

sea vesicomyid clam symbioses

Corinna Breusing[1,*†], Maëva Perez[2,3†], Roxanne A. Beinart[1], C. Robert Young[2]

[1]University of Rhode Island, Graduate School of Oceanography, Narragansett, RI, USA

[2]National Oceanography Centre, Southampton, SO14 3ZH, UK

[3]University of Montreal, Department of Biological Sciences, Montreal, Quebec H3T 1J4, Canada

*Corresponding author: corinnabreusing@gmail.com

[†]Authors contributed equally

Running title: Niche adaptation in clam symbionts

Keywords: Vesicomyid clams, chemosynthetic endosymbionts, comparative genomics, vertical

transmission, reductive genome evolution, niche partitioning

Abstract

Vertical transmission of bacterial endosymbionts is accompanied by virtually irreversible gene

loss that can provide insights into adaptation to divergent ecological niches. While patterns of

genome reduction have been well described in some terrestrial symbioses, they are less

understood in marine systems where vertical transmission is relatively rare. The association

between deep-sea vesicomyid clams and chemosynthetic Gammaproteobacteria is one example

of maternally inherited symbioses in the ocean. Differences in nitrogen and sulfur physiology

between the two dominant symbiont groups, *Ca*. Ruthia and *Ca*. Vesicomyosocius, have been

hypothesized to influence niche exploitation, which likely affects gene content evolution in these

symbionts. However, genomic data are currently limited to confirm this assumption. In the

present study we sequenced and compared 11 vesicomyid symbiont genomes with existing

assemblies for *Ca*. Vesicomyosocius okutanii and *Ca*. Ruthia magnifica. Our analyses indicate

that the two vesicomyid symbiont groups have a common core genome related to

chemosynthetic metabolism, but differ in their potential for nitrate respiration and flexibility to

environmental sulfide concentrations. Moreover, *Ca*. Vesicomyosocius and *Ca*. Ruthia have

different enzymatic requirements for cobalamin and nickel and show contrasting capacities to

acquire foreign genetic material. Tests for site-specific positive selection in metabolic candidate

genes imply that the observed physiological differences are adaptive and thus likely correspond

to ecological niches available to each symbiont group. These findings highlight the role of niche

differentiation in creating divergent paths of reductive genome evolution in vertically transmitted

symbionts.

Introduction

Heritable symbioses with intracellular bacteria are observed across the eukaryotic domain of life [1]. These symbioses can have significant impacts on the host's niche through their effects on resource utilization, tolerance to stressors, and resistance to predators and pathogens [1–3]. Vertical transmission of bacterial lineages from parent to off-spring typically results in irreversible genome degradation due to bottleneck effects, relaxed selective pressures and restricted opportunities for horizontal gene transfer, limiting retention of genes to those that are essential to the functioning of the host-symbiont association [3]. Thus, differences in gene content among related symbionts can reveal how the host-symbiont pairs diverged in their niches over evolutionary time. For example, divergence in plant host use between insect species is evident in the biosynthetic pathways encoded in the genomes of their obligate endosymbionts [4]. Ultimately, niche differentiation driven by differential gene loss has the potential to be an important driver of host evolution through ecological speciation, as well as a significant factor in the structure of host communities through its effects on competition and habitat partitioning between host-symbiont pairs. Despite its importance for both ecological and evolutionary processes, there is still a significant gap in our understanding of the patterns of gene/pathway reduction in divergent vertically transmitted bacterial endosymbionts. This is especially true for the heritable endosymbionts of marine organisms, since vertical transmission is less common among aquatic symbioses [1].

Relatively strict vertical transmission of bacterial endosymbionts has been observed in hydrothermal vent- and cold seep-associated deep-sea clams of the family Vesicomyidae [5], providing an opportunity to examine patterns of genome reduction in a marine symbiosis. Vesicomyid clams are nutritionally dependent on chemosynthetic maternally inherited

Gammaproteobacteria, which derive chemical energy from the oxidation of reduced sulfur

compounds to produce food for their hosts [6–10]. The clam-bacteria association is obligate,

given that vesicomyid hosts have a highly reduced digestive system and cannot survive without

their symbionts [8]. Topological congruences between host mitochondrial and symbiont

phylogenetic trees suggest that symbionts co-evolve with their hosts [5], although disruptions of

these relationships have been reported due to infrequent horizontal transmission events that allow

for lateral gene transfer and homologous recombination between bacterial lineages [11–16].

Based on ribosomal sequence data, vesicomyid symbionts are classified into two phylogenetic

clades [17]. Clade I symbionts (a.k.a *Ca*. Vesicomyosocius) are typically associated with hosts of

the *gigas*-group, including the nominal genera *Akebiconcha*, *Archivesica*, *Laubiericoncha* and

Phreagena, whereas clade II symbionts (a.k.a. *Ca*. Ruthia) associate with diverse groups of

vesicomyid hosts [18]. Previous genomic comparisons between one representative symbiont

lineage from each clade (*Ca*. Ruthia magnifica and *Ca*. Vesicomyosocius okutanii) indicated that

the two groups differ in the amount of genome reduction [19–21]. This limited genomic data,

along with targeted PCR-based surveys, has suggested that *Ca*. Vesicomyosocius symbionts are

characterized by genomes with relatively low GC content that are deplete of critical genes for

nucleotide excision and recombination (e.g., *recA*, *mutY*, *uvrABCD*), indicating that they are in

an advanced stage of genome erosion. By contrast, *Ca*. Ruthia symbionts are thought to have

genomes with higher GC content that still contain intact homologs of these genes [17, 21, 22].

Variations in genome reduction and host affiliation between *Ca*. Vesicomyosocius and

Ca. Ruthia do not appear to be driven by adaptation to different broad-scale habitat types, as

clam species hosting lineages from both clades have been found at vents and seeps and often co-

occur at the same locality [8, 18, 23, 24]. However, the two symbiont clades appear to differ in

physiological characteristics related to nitrate reduction and sulfur metabolism, which may affect

microhabitat exploitation [8, 23], and could, thus, influence patterns of gene conservation

between clades. In fact, niche partitioning has been linked to patterns of gene loss and retention

in a variety of marine and freshwater bacteria [25, 26].

Genome sampling of multiple lineages within each symbiont clade is necessary to assess

whether the observed variation in nitrogen and sulfur physiology is a potential driver of niche

segregation between *Ca*. Vesicomyosocius and *Ca*. Ruthia. To address this issue and identify

other metabolic differences between the two symbiont clades, we sequenced the genomes of 11

Ca. Vesicomyosocius and *Ca*. Ruthia lineages and compared them with previous assemblies of

Ca. V. okutanii [20] and *Ca*. R. magnifica [19] (Table 1). Candidate metabolic genes that were

found to be differentially conserved between both symbiont clades were investigated for signals

of pervasive and episodic diversifying selection to assess their role in niche adaptation in these

symbionts.

Materials and Methods

SAMPLE COLLECTION AND DNA EXTRACTION

Clam specimens were collected from nine cold seep and hydrothermal vent sites in the Pacific

and Atlantic Ocean during research expeditions between 1994 and 2004 (Table 2). Upon

recovery of the submersibles, samples were dissected and frozen at −80°C. DNA was extracted

onshore from symbiont-bearing gill tissues with the DNeasy Blood & Tissue kit (Qiagen, Hilden,

Germany) following manufacturer's instructions. Clam host species were identified via

mitochondrial cytochrome-c-oxidase I (*COI*) sequencing using vesicomyid-specific primers [27].

For the remainder of this manuscript we will use the previously erected symbiont genus names

Ca. Vesicomysocius and *Ca*. Ruthia plus the host species name to define the different symbiont

lineages.

METAGENOMIC SEQUENCING, ASSEMBLY AND ANNOTATION

Barcoded 2x300 bp metagenomic libraries were prepared with the KAPA Hyperplus Library

Preparation kit (KAPA Biosystems, Wilmington, MA, US) and sequenced on an Illumina MiSeq

system at the National Oceanography Centre (Southampton, UK). After initial quality checks

with FASTQC v0.11.5 [28], reads were adapter-clipped with TRIMMOMATIC v0.36 [29] and

assembled *de novo* with VELVET v1.2.10 [30], SPADES v3.13.1 [31] or GENEIOUS v10.1.3

(http://www.geneious.com/) using manual optimizations of k-mer size distribution and read

depth. Symbiont contigs were binned based on GC content and read coverage profiles and

subsequently scaffolded with SSPACE v2.0 [32]. For assemblies that resulted in a single scaffold,

we tried to circularize the genomes by reassembling and overlapping the last 5000 bp of each

contig end with SPADES. The final symbiont genome assemblies were annotated with RAST v2.0

[33] and investigated for quality with QUAST v5.0.0 [34]. Positional homologs between

assemblies were identified by aligning the symbiont genomes with PROGRESSIVEMAUVE

v20150213 [35], using a minimum identity cutoff of 30% and a minimum coverage cutoff of

60%.

COMPARATIVE GENOMICS

We ran the ANVI'O v6.2 pangenomics workflow [36, 37] to assess core and lineage-specific

features of the *Ca.* Vesicomyosocius and *Ca.* Ruthia genomes and to determine phylogenomic

relationships based on 739 single copy gene clusters. We used the "--ncbi-blast" option to

calculate amino acid sequence similarities and the MCL algorithm to define protein clusters

based on the following settings: minbit = 0.5, mcl-inflation = 6, min-occurrence = 1. Gene

cluster annotations were based on the Cluster of Orthologous Groups (COG) database [38].

Complementarily, phylogenetic analyses of the symbiont *16S* rRNA gene were performed with

MRBAYES v3.2.7a [39] in CIPRES v3.3 [40] based on a GTR+I+G substitution model. Prior to

phylogenetic reconstruction, all sequences were aligned against the global SILVA *16S* rRNA

alignment with SINA v1.2.11 [41] to account for secondary structure of the *16S* rRNA. We ran

three heated chains and one cold chain for 1,100,000 generations, sampling every 100

generations and discarding the first 100,000 generations as burn-in. The symbiont *16S* rRNA

gene sequence of the *Bathymodiolus thermophilus* symbiont (CP024634) was used as outgroup

for tree rooting. MCMC convergence was assessed with TRACER v1.7.1 [42].

TESTS FOR ADAPTIVE EVOLUTION

Our comparative genomic analyses identified a number of differentially conserved candidate

genes that might affect niche adaptation in *Ca*. Ruthia and *Ca*. Vesicomyosocius: cysteine

dioxygenase (*cdo*), hydrogenase/urease accessory protein (*hupE*), methionine synthase (*metE*,

metH), respiratory nitrate reductase (*narGHIJ*), assimilatory nitrate reductase (*nasA*), and

sulfide:quinone oxidoreductase type I (*sqrI*). To examine whether sites within these genes are

subject to pervasive or episodic diversifying selection we applied Bayesian approximation and

mixed-effects maximum likelihood approaches implemented in FUBAR v2.2 [43] and MEME

v2.1.2 [44], respectively. Both programs infer per-site nonsynonymous (dN) and synonymous

(dS) substitution rates for a given coding nucleotide alignment and matching phylogenetic tree,

but aim to detect different forms of positive selection. FUBAR determines sites that show signals

of pervasive positive selection across the entire phylogeny [43], while MEME identifies sites that

are evolving under episodic positive selection in a proportion of branches [44]. Analyses were

performed using all available symbiont sequences for a gene, including positional homologs of

the phylogenetically related *Bathymodiolus thermophilus* symbiont (CP024634), to increase

statistical power. Input trees for model optimization were generated with FASTTREE v2.1.11 [45]

under a GTR substitution model. FUBAR analyses included 5 MCMC chains, with chain lengths

of 2000000, a burn-in of 1000000 and a sample size of 100, while MEME analyses were run with

default settings. Because site-level tests for positive selection are relatively conservative, we

chose recommended p-value thresholds of 0.1 for MEME and posterior probability thresholds of

0.9 for FUBAR to assess statistical significance [46].

DATA AVAILABILITY

The final genome assemblies including raw Illumina data have been uploaded to GenBank and

the Sequence Read Archive under BioProject number PRJNA641445. Host mitochondrial *COI*

sequences have been deposited in GenBank under accession numbers MT894120–MT894130.

Results

GENOME ASSEMBLY METRICS

Final assemblies for the eleven genomes comprised 1–39 contigs with a total size of 1.02–1.59

Mb, an N50 of 0.06–1.23 Mb, a GC content of 31.10–36.98%, and an average coverage of

49.20–153.10 (Table 1). Genomes of the different *Ca.* Vesicomyosocius and *Ca.* Ruthia lineages

were comparable in size to those of *Ca.* V. okutanii and *Ca.* R. magnifica, respectively [19, 20],

and were assembled into single circular chromosomes, with two exceptions. Genomes of the

different *Ca.* Ruthia lineages were less homogeneous in size and typically larger than those of

the *Ca.* Vesicomyosocius lineages. They also possessed higher gene density. The genome

assemblies of *Ca.* R. phaseoliformis and *Ca.* R. southwardae were the largest among all

symbiont genomes sequenced and were characterized by a relatively high amount of

fragmentation (Table 1).

COMPARATIVE GENOMICS AND PHYLOGENOMICS

Phylogenetic analyses of the *16S* rRNA gene and 739 orthologous single-copy gene clusters

identified the two previously described clades of vesicomyid symbionts [17], comprising the

nominal genera *Ca.* Ruthia and *Ca.* Vesicomyosocius (Figure 1). While the topologies of the *Ca.*

Vesicomyosocius lineages generally agreed between gene trees, the topologies of *Ca.* R.

magnifica, *Ca.* R. pliocardia, *Ca.* R. phaseoliformis and *Ca.* R. southwardae were incongruent

and poorly resolved based on the *16S* rRNA gene alone (Figure 1). Average nucleotide identities

(ANI) between lineages of the two clades were on average 82.93–83.85% in aligned nucleotide

regions, but only 42.49–66.97% if unaligned regions were considered (Table S1). ANIs within

clades were higher, with values of 93.55–99.99% (85.91–99.99%) between the different *Ca.*

Vesicomyosocius lineages, and 87.80–95.22% (52.43–76.59%) between the different *Ca.* Ruthia

lineages (Table S1). Based on conservative species-level ANI cutoffs of 95–96% [47], these

values suggest that most symbiont lineages represent distinct bacterial species, except for *Ca*. V.

diagonalis-extenta, *Ca*. V. okutanii-soyoae and possibly *Ca*. R. pacifica-rectimargo.

CORE GENOME CHARACTERISTICS

All vesicomyid symbiont genomes shared 822 core gene clusters (consisting of 10,931 genes),

the majority of which were assigned to five broader functional categories: (1) energy production

and conversion, (2) amino acid metabolism and transport, (3) coenzyme metabolism and

transport, (4) translation, and (5) posttranslational modification (Figure 2; Table S2, S3).

Energy metabolism

The genomes of all *Ca.* Ruthia and *Ca.* Vesicomyosocius lineages contained genes for the

oxidation of reduced sulfur compounds that serve as energy sources for chemoautotrophic

growth [9]. All genomes encoded the sulfur oxidation (SOX) multienzyme pathway (without

soxCD), the reversible dissimilatory sulfite reductase (rDSR) pathway as well as the adenosine

5'-phosphosulfate (APS) reductase pathway, indicating that these symbiont lineages are able to

oxidize sulfide, thiosulfate and/or sulfite for energy production (Table S2, S3) [48, 49]. In

addition, all genomes comprised genes for sulfide:quinone oxido-reductase type I and VI (SQR),

which can convert sulfide to sulfane sulfur [49]. With the exception of *Ca.* V. soyoae 2 and *Ca.*

V. okutanii, the *Ca.* Vesicomyosocius lineages contained two copies of the gene encoding SQR

type I, whereas this gene was present as a single copy in the *Ca.* Ruthia clade. In addition,

complex sulfur compounds may be made accessible via homologs of the polysulfide reductase

NrfD (Table S2, S3).

Based on their gene content, it is likely that all symbiont lineages can use a variety of

different enzymes to conserve energy via cross-membrane electron transport, including NADH-

ubiquinone oxidoreductase (Complex I), SQR, bacterial Rnf complex, cytochrome bc_1 complex

(Complex III), terminal cbb3-type cytochrome-c-oxidase (Complex IV) and F_0F_1-type ATP

synthase (Complex V) (Table S2, S3).

Inorganic carbon fixation and biosynthetic processes

Members of the *Ca.* Vesicomyosocius and *Ca.* Ruthia clades both encoded a form II ribulose

bisphosphate carboxylase (cbbM) and other key enzymes for carbon assimilation via the Calvin-

Benson-Bassham cycle as well as a complete gene set for the non-oxidative branch of the

pentose phosphate pathway (Table S2, S3). Both symbiont clades lacked the gene for

sedoheptulose-bisphosphatase and might instead rely on a reversible pyrophosphate-dependent

phosphofructokinase (PPi-PFK) to interconvert between sedoheptulose 1,7-bisphosphate and

sedoheptulose 7-phosphate. PPi-PFK is likely also used to catalyze the phosphorylation of

fructose-6-phosphate to fructose 1,6-bisphosphate during glycolysis, as the gene for its ATP-

dependent homolog was absent in all vesicomyid symbiont genomes (Table S2, S3).

All symbiont lineages have the potential to further metabolize glycolytic intermediates

and end products via a partial tricarboxylic acid (TCA) cycle and pentose phosphate pathway to

produce precursors for the generation of several macronutrients, coenzymes and nucleotides

(Table S2, S3). While functional gene copies of α-ketoglutarate decarboxylase (E1 component of

the α-ketoglutarate dehydrogenase complex) and fumarate reductase Fe-S subunit appeared to be

missing or degenerated in all symbiont genomes, we found intact genes coding for malate

dehydrogenase. Both the *Ca*. Ruthia and *Ca*. Vesicomyosocius genomes contained complete

gene sets for the independent biosynthesis of 19 amino acids and a variety of enzyme cofactors,

including most vitamins and their derivatives (e.g., coenzyme A, FAD, NAD^+), hemes and

sirohemes, porphyrins, molybdopterin, ubiquinone and glutathione. The gene encoding

homoserine kinase (*thrB*), an essential enzyme in threonine biosynthesis, was missing from all

symbiont genomes, although it is possible that its function might be performed by a

serine/threonine kinase that was present in genomes from both the *Ca.* Ruthia and *Ca.*

Vesicomyosocius clade. Similarly, a separate gene for histidinol phosphatase involved in

histidine biosynthesis was lacking from all symbiont genomes. However, the genomes of

symbionts from both clades contained homologs of the *hisB* gene, which encodes a bifunctional

imidazoleglycerol-phosphate dehydratase/histidinol-phosphatase. Pathways for the generation of

retinol, cobalamin, ascorbic acid, cholecalciferol, menaquinone and tocopherol were incomplete,

while protoheme biosynthesis appeared to occur through a novel form of protoporphyrinogen IX

oxidase (HemJ), which has so far mostly been described in cyanobacteria [50]. The *ubiD*/*ubiX*

gene complex for ubiquinone biosynthesis was absent in all symbiont lineages. The lack of

UbiD/UbiX might be compensated by UbiG via an alternative pathway that generates ubiquinone

through methylation of 3-polyprenyl-4,5-dihydroxybenzoate.

CLADE- AND LINEAGE-SPECIFIC GENOME CHARACTERISTICS

46 gene clusters (containing 321 genes) were specific to the *Ca.* Ruthia lineages, while six gene

clusters (containing 44 genes) occurred exclusively in the *Ca.* Vesicomyosocius clade (Figure 1;

Tables S2, S3). In both clades, many of these gene clusters had unknown or poorly characterized

functions, whereas others implied clade-specific differences in cell wall biogenesis, translation

and post-translational modification (Figure 2; Tables S2, S3). Genomes in the *Candidatus* Ruthia

clade also contained a variety of gene clusters associated with biosynthetic and transport

processes that were absent in genomes from the *Ca.* Vesicomyosocius clade (Figure 2; Tables

S2, S3). In several cases, genomes from the *Ca.* Ruthia and *Ca.* Vesicomyosocius clades shared

the same gene clusters but differed in the amount of degeneration in the associated gene loci

(Table S3). An overview of the main differences between the two symbiont clades is given in

Figure 3.

Methionine synthase

Candidatus Ruthia and *Ca.* Vesicomyosocius appear to use different enzymes for the synthesis

of methionine (Figure 3; Tables S2, S3). The gene for the cobalamin-dependent homocysteine

methyltransferase (*metH*) as well as genes for cobalamin (precursor) transport and conversion

(*btuM*, *btuR/cobA*) were conserved in genomes from the *Ca.* Vesicomyosocius clade but were

missing or degenerated in all of the *Ca.* Ruthia lineages, except for *Ca.* R. phaseoliformis and

Ca. R. southwardae. Conversely, the gene for the cobalamin-independent version of this enzyme

(*metE*) along with its transcriptional activator (*metR*) were exclusively found in the *Ca.* Ruthia

symbiont genomes. Notably, almost all genes for *de novo* cobalamin biosynthesis were absent

from the investigated symbiont genomes, with the exception of cobyrinic acid A,C-diamide

synthase (*cbiA*), adenosylcobalamin/alpha-ribazole phosphatase (*cobC*) (all genomes) and the

high affinity cobalamin transporter BtuB (*Ca*. V. gigas).

Nitrate reductase

An operon coding for the membrane-bound nitrate-reductase complex NarGHIJ was conserved

in all *Ca.* Vesicomyosocius symbiont genomes, but not in the *Ca.* Ruthia clade, which appeared

to contain non-functional remnants of this operon. Conversely, the symbionts of the *Ca.* Ruthia

clade encode the cytoplasmic assimilatory nitrate reductase NasA, which is degenerated in the

Ca. Vesicomyosocius clade (Figure 3; Tables S2, S3).

Cysteine dioxygenase and isocitrate lyase

The gene coding for cysteine dioxygenase type I, which catalyzes the conversion of L-cysteine to

cysteine sulfinic acid, was conserved in all *Ca.* Vesicomyosocius lineages, but absent or

degenerated in most *Ca.* Ruthia symbiont genomes (with the exception of *Ca.* R. phaseoliformis

and *Ca.* R. pliocardia). By contrast, only the *Ca*. Ruthia symbionts encode genes for isocitrate

lyase, a key enzyme of the glyoxylate cycle (Figure 3; Tables S2, S3).

Transcription, translation and post-translational modification

All vesicomyid symbiont genomes contained an operon for a Class Ia ribonucleotide reductase

(*nrdAB*), but only the *Ca.* Ruthia lineages appeared to also encode the gene for its transcriptional

repressor (*nrdR*). In addition, we found genes for several enzymes involved in protein

modification and response to cellular stress in the genomes of the *Ca*. Ruthia clade that were

absent in *Ca*. Vesicomyosocius (Figure 3; Table S2, S3). For instance, all *Ca.* Ruthia lineages

contained genes for the GTP-binding protein HflX (exception: *Ca*. R. pliocardia), and the peptide

methionine sulfoxide reductase MsrB, which play a role in dissociation of translationally arrested

ribosomes [51], and protein repair after oxidative damage, respectively. Likewise, most *Ca.*

Ruthia lineages encoded genes for GidB and TrmL methyltransferases and a tRNA-

dihydrouridine synthase (Dus), which are involved in RNA modification, as well as a gene for

the ribosomal silencing factor RsfS, which is an important translational regulator.

Cell wall and membrane biosynthesis

Ca. Ruthia and *Ca.* Vesicomoysocius differed in several genes that are involved in biogenesis of

the cellular envelope (Figure 3; Tables S2, S3). Although we found complete pathways for the

production of the common membrane lipid phosphatidylethanolamine in the genomes of all

vesicomyid symbiont lineages, genes for diacylglycerol kinase, which is necessary for

phospholipid recycling, was only present in the *Ca.* Ruthia clade. Similarly, all *Ca.* Ruthia

symbionts encoded a 1,6-anhydro-N-acetylmuramate kinase (AnmK) and an outer membrane

lipoprotein (SlyB), which are important for cell wall recycling and integrity, respectively. The

Ca. Ruthia lineages also contained a small-conductance mechanosensitive channel involved in

osmoregulation (MscS), a lipopolysaccharide (LPS) export system protein (LptA) involved in

LPS-translocation across the periplasm, and an N-acetyl-anhydromuramyl-L-alanine amidase

(AmpD) involved in cell wall degradation. Homologs of these genes were either completely

missing or pseudogenized in the *Ca.* Vesicomyosocius symbionts. Both symbiont clades

possessed genes for peptidoglycan biosynthesis, although MurD, MurF, MraY and MurG

enzyme functionalities might be impaired or altered by the presence of internal stop codons in

the case of *Ca.* V. diagonalis and *Ca.* V. extenta. In addition, both symbiont clades contained

glycosyltransferases related to LPS biosynthesis, but appeared to encode different isoforms of

this enzyme.

Transport across membrane

Multiple components of a type I secretion system (lapC, lapB, lapE, and the secreted agglutinin

RTX) were found in all of the *Ca.* Ruthia symbionts except for *Ca.* R. pliocardia. By contrast,

this locus was missing in the *Ca.* Vesicomyosocius clade (Figure 3; Table S3, S3). The *Ca.*

Ruthia symbionts also encoded a putative hydrogenase/urease accessory protein (HupE), which

is thought to be a nickel or cobalt transporter (Table S3) [52]. Although *hupE* is often associated

with operons coding for [NiFe] hydrogenases, we did not find genes encoding hydrogenase

subunits in any of the symbiont genomes. However, a gene encoding a nickel-dependent

glyoxalase I was present in the genomes of most *Ca*. Ruthia symbionts.

GENE LOSS

DNA repair and recombination

In agreement with Kuwahara et al. [17] and Shimamura et al. [22], genes of the nucleotide

excision repair pathway, *uvrA*, *uvrD*, *uvrD* paralog and *mfd*, were conserved in most symbiont

genomes, while *uvrB* and *uvrC* were degenerated in all *Ca*. Vesicomyosocius lineages (Figure 3;

Table S2, S3). Within the *Ca*. Ruthia clade, *uvrA*, *uvrB*, *uvrD* paralog and *mfd* were present in all

lineages, whereas *uvrC* was lost in *Ca*. R. pliocardia, and *uvrD* was lost in *Ca*. R. phaseoliformis

and *Ca*. R. southwardae. Most *Ca*. Ruthia lineages contained genes for repair of alkylated DNA

(*alkD*) and strand breaks (*radA*; exception: *Ca*. R. magnifica), while homologs of these genes

were absent from all *Ca*. Vesicomyosocius symbiont genomes. Furthermore, we found that

essential genes involved in SOS response to DNA damage, *recA*, *recFOR*, and *recX*, were lost in

the *Ca*. Vesicomyosocius clade and *Ca*. R. magnifica. In the other *Ca*. Ruthia lineages, these

genes were conserved with the exception of *recF*, which was degenerated in *Ca*. R. pacifica, *Ca*.

R. rectiomargo and *Ca*. R. pliocardia, and *recO*, which was degenerated in *Ca*. R.

phaseoliformis. Likewise, the gene coding for RuvC, an essential component of the last step of

the recF and recBCD pathways for homologous recombination [53], as well as the genes coding

for the XerCD recombinase system and the DNA recombination protein RmuC were lost in the

Ca. Vesicomyosocius clade, but conserved in most of the *Ca*. Ruthia lineages.

Mobile elements and defense against pathogens

The genomes of all vesicomyid symbionts are notably sparse in genes related to anti-viral

defense and transposition (Figure 3; Table S2, S3). Phage-related genes except for a putative

phage tape measure protein were completely missing in the *Ca.* Vesicomyosocius clade, while a

few transposases (e.g., InsA, InsE, InsO, Tra8, RayT), integrases and other phage-derived

proteins were found in some of the *Ca.* Ruthia lineages, in particular *Ca.* R. southwardae and *Ca.*

R. phaseoliformis. All genomes contained genes coding for the lambda lysogenization regulator

HflD. In addition, remnants of type I restriction-modification systems (HsdMRS) and mRNA-

degrading toxin-antitoxin systems (e.g., MazEF) were present in the genomes of *Ca.* R. pacifica,

Ca. R. rectimargo, *Ca.* R. phaseoliformis and *Ca.* R. southwardae, but lost in all other symbiont

genomes. *Candidatus* R. southwardae and *Ca.* R. phaseoliformis further contained degenerated

operons for the 5-methylcytosine-specific restriction endonuclease McrBC. In addition,

putatively defunct versions of Cascade complex genes that were previously part of a

CRISPR/Cas system were found in *Ca.* R. pliocardia (*cas1*, *cas2*) and *Ca.* R. southwardae (*cas1*,

cas3).

SITE-SPECIFIC POSITIVE SELECTION IN METABOLIC GENES

Site-specific tests for adaptive evolution identified signals of positive selection in all 10

investigated candidate genes, except for *narI* (Table 3). Bayesian approximation analyses based

on FUBAR estimated that 1–2 sites were under pervasive diversifying selection in *cdo*, *hupE*,

narG, *nasA* and *sqrI* across the entire phylogeny. MEME analyses corresponded well with results

obtained by FUBAR, confirming sites under widespread positive selection and detecting 1–8

additional sites under episodic diversifying selection in a proportion of branches for all genes

(with the exception of *narI*) (Table 3). In three cases, p-values calculated by MEME were

insignificant for positively selected sites identified by FUBAR (*narG*: 451, *nasA*: 451, *sqrI*: 4).

However, since p-values were relatively close to the significance threshold of 0.1, it is possible

that statistical significance would have been observed with the inclusion of additional sequences.

Discussion

Given that obligate endosymbionts experience limited opportunities for acquisition of new genes

through horizontal transfer, differences in gene loss between symbiont lineages have the

potential to permanently separate holobiont niches, which has consequences for ecological

processes, like habitat use, and evolutionary processes, like host speciation. Differential gene

loss in divergent lineages of obligate, vertically transmitted endosymbionts has been well-

recognized to be important in the ecology and evolution of sap-feeding insects. For example,

differences in symbiont gene content may differentiate use of plant species by their insect hosts

[3, 4]. This symbiont-induced specialization on particular plant species has implications for the

evolutionary diversification of insect populations and species [3, 54]. However, outside the well-

studied insect-bacteria symbiosis [55–58], assessments of differential gene loss across clades and

lineages of vertically transmitted endosymbionts have been fairly limited, despite the broad

significance of gene loss patterns to our understanding of holobiont ecology and evolution.

As one of the few known examples of vertically transmitted marine symbioses, deep-sea

vesicomyid clams represent an opportune system to study the process of incremental gene loss in

bacterial endosymbionts and its potential effect on niche differentiation in the ocean. In the

present study we compared the genomes of 13 lineages within the two dominant vesicomyid

symbiont clades, *Ca*. Ruthia and *Ca*. Vesicomyosocius, to assess commonalities and differences

in metabolic gene content and their eco-evolutionary implications. Similar to the vertically

transmitted endosymbionts of insects, which commonly display extreme stasis in a core genome

consistent with their primary role in nitrogen provisioning for their hosts [55–57], our analyses

reveal several shared features related to sulfur-based chemoautotrophic metabolism and energy

conversion (in line with [8]), which are fundamental to the nutritional services provided by the

endosymbionts of vesicomyid clams. By contrast, differences in gene loss among symbiont

lineages suggest a role for holobiont niche differentiation in shaping genome evolution. In the

endosymbionts of insects, differential gene loss is typically tied to variation in genes related to

components of the host diet, for example disparities in the availability of amino acids and

vitamins across food sources [3, 4, 58]. In the symbionts of vesicomyid clams, we observe

notable contrasts in cofactor, oxygen and sulfur requirements among clades. Several sites within

metabolic genes that were differentially conserved between *Ca*. Ruthia and *Ca*.

Vesicomyosocius appeared to be at least episodically under diversifying selection, suggesting

that genetic differences between the two symbiont clades reflect adaptations to different

ecological niches.

For instance, *Ca*. Ruthia and *Ca*. Vesicomyosocius encoded different, convergently

evolved types of methionine synthase [59], which vary in their dependence on vitamin B12 as

co-factor. While *Ca*. Ruthia contains the cobalamin-independent version MetE, *Ca*.

Vesicomyosocius comprises homocysteine methyltransferase MetH, which is cobalamin-

dependent. As both clades appear unable to synthesize cobalamin *de novo* (like their eukaryotic

hosts), these findings indicate that the environmental availability of vitamin B12 has the potential

to be an important factor influencing the distribution of these taxa. Cobalamin independence in

Ca. Ruthia may offer an obvious selective advantage, by allowing these symbioses to exploit

niches that would otherwise be inaccessible. By contrast, the requirement for exogeneous

vitamin B12 (or its derivatives) has the potential to limit the range of (micro)habitats *Ca*.

Vesicomyosocius-based associations can colonize, unless cobalamin is acquired from a

(currently unknown) secondary symbiont, as for example seen in some insect-bacteria symbioses

[60]. Despite this potential cost, the retention of a cobalamin-dependent methionine synthase in

Ca. Vesicomoysocius comes with an evolutionary benefit, given that MetH has a fifty fold

higher catalytic rate constant than MetE and thus enables faster growth [59]. The preservation of

cobalamin-dependent enzymes as a result of conferred physiological advantages appears to be

common across the eubacterial domain [61]. A recent genomic analysis by Shelton et al. [61]

showed that 86% of bacterial lineages have at least one cobalamin-dependent enzyme despite the

existence of a cobalamin-independent alternative, and that many of these lineages rely on

vitamin B12 production from other microbes in their environment. The importance of vitamin

B12 for the biology of the two symbiont groups is also evident in the fact that only *Ca*. Ruthia

encodes a transcriptional repressor (NrdR) for the ribonucleotide reductase NrdAB, a key

enzyme that controls the synthesis of DNA [62]. In *Ca*. Vesicomyosocius, expression of NrdAB

is probably regulated by cobalamin, which has been shown to repress NrdAB transcription

through riboswitches [63].

There is evidence that *Ca*. Vesicomyosocius and *Ca*. Ruthia differ in their requirements

for other enzyme cofactors, such as nickel. Only the *Ca*. Ruthia symbiont genomes encoded a

specific transporter for nickel uptake, and most of these lineages contained at least one

confirmed nickel-dependent enzyme, glyoxalase I [64]. By contrast, the ecological significance

of nickel for *Ca*. Vesicomyosocius is questionable. Although an intact gene for a putative

Mg/Co/Ni transporter (MgtE) was present in the genomes of most vesicomyid symbionts,

electrophysiological analyses indicate that MgtE is not capable of transporting nickel [65]. In

addition, unambiguous annotations for nickel-dependent enzymes were missing from the

genomes of the *Ca*. Vesicomyosocius symbionts. We found a gene encoding a protein of the

glyoxalase/bleomycin resistance protein/dioxygenase superfamily, but enzymes within this group

use a variety of different metals as cofactors and do not necessarily rely on nickel.

454   Apart from their contrasting dependencies on enzyme cofactors, *Ca.* Ruthia and *Ca.*

Vesicomyosocius exhibit important differences in nitrogen metabolism [8]. While all *Ca*.

Vesicomyosocius lineages encode genes for the membrane-bound dissimilatory nitrate reductase

NarGHIJ, which utilizes nitrate as terminal electron acceptor for anaerobic respiration, the *Ca*.

Ruthia symbionts contain genes for the assimilatory nitrate reductase NasA, which metabolizes

nitrate as nitrogen source for growth [66]. Despite the distinct physiological functions of these

enzymes, ammonium generated through dissimilatory nitrate reduction is likely also used for

biosynthetic purposes in *Ca*. Vesicomyosocius given the absence of an assimilatory equivalent.

As noted by Newton et al. [8], the use of NarGHIJ for nitrate reduction in *Ca*. Vesicomyosocius

might allow these symbioses to inhabit hypoxic environments, since the use of nitrate as an

electron acceptor would reduce the symbiont's requirement for oxygen and, consequently, also

reduce the oxygen requirement for the total holobiont. Segregation into differentially oxygenated

niches might further explain why several genes involved in oxidative stress response (e.g., *uvrA*,

recA, dsbC, msrB) were largely conserved in *Ca*. Ruthia but not in *Ca*. Vesicomyosocius.

468   The presence of a single (or predominant) metabolic pathway for methionine biosynthesis

and nitrate reduction in each symbiont group suggests a clearing of redundancies through

reductive genome evolution, but the differential loss between the groups does not appear to be

random. Indeed, we observe very different patterns of *narGHIJ* and *metH* pseudogenization even

between very closely related symbionts (e.g., *Ca.* R. pacifica and *Ca.* R. rectimargo), suggesting

that the inactivation of these genes has happened multiple times throughout the recent history of

the *Ca*. Ruthia clade. Elimination of functionally redundant genes is also apparent in the low

level of duplication in the genomes of all vesicomyid symbionts. Interestingly, almost all *Ca*.

Vesicomyosocius lineages, but none of the *Ca*. Ruthia symbionts contain two copies of SQR

type I. Previous studies have shown that SQRs have different substrate concentration optima,

with SQR type I being adapted to low sulfide concentrations in the micromolar range, and SQR

type VI being adapted to elevated sulfide concentrations in the millimolar range [67–69].

Perhaps the two SQR type I versions in *Ca*. Vesicomyosocius are necessary to maximize sulfide

oxidation at different micromolar H_2S concentrations, thereby optimizing the holobionts'

adaptability to fluctuations in environmental reductant levels. Variations in host and symbiont

metabolic plasticity and enzyme properties in relation to sulfur availability or processing might

further be important for avoiding competition between sympatric species [23, 24]. For example,

Goffredi and Barry [23] suggested that adaptation to different sulfide levels plays a role in niche

partitioning of co-occurring clam species in Monterey Bay. *Ca*. Vesicomysocius-based

associations have been found to occupy microhabitats with higher levels of sulfide, while *Ca*.

Ruthia-based symbioses inhabit zones with lower concentrations of this reductant [23].

Hydrogen sulfide can disrupt aerobic respiration and must be neutralized to avoid toxic effects

on the animal host. In a variety of symbiont-bearing vent and seep invertebrates, the non-

proteinogenic amino acid thiotaurine, a derivative of hypotaurine, has been shown to play a role

in sulfide detoxification [70–72]. Our analyses imply that in *Ca*. Vesicomyosocius-based

associations host H_2S tolerance might be amplified (or solely mediated) by symbiont-derived

thiotaurine, given that all lineages of the *Ca*. Vesicomyosocius clade encode genes for cysteine

dioxygenase, a key enzyme in taurine/hypotaurine biosynthesis. If *Ca*. Vesicomyosocius-based

symbioses are typically exposed to higher concentrations of sulfide, symbiont-driven sulfide

inactivation might be another adaptation to the specific ecological niches of these taxa.

The role of cysteine dioxygenase in bacteria is relatively poorly understood, although a

few functions have been suggested [73]. For example, its activity is likely relevant for

replenishing the internal sulfur and carbon pools, since its product cysteine sulfinic acid can be

transaminated to generate sulfite and pyruvate [73]. We suggest another potential physiological

role for this enzyme that might help to illuminate how *Ca*. Vesicomyosocius replenishes

metabolic intermediates of its partial TCA cycle, in particular succinate. While succinate

regeneration via the glyoxylate shunt has been proposed for *Ca*. Ruthia, the mechanism of

succinate recycling in *Ca*. Vesicomyosocius has so far remained unclear. Newton et al. [8]

hypothesized that succinate might be taken up from the host's cytoplasm via a putative sugar

transporter. Although we did find a gene for a potential di- and tricarboxylate transporter

(COG0471) in the genomes of all *Ca*. Vesicomyosocius lineages, it is also possible that succinate

might be restored through anaplerotic reactions resulting from the degradation of cysteine.

Cysteine is metabolized by cysteine dioxygenase to produce cysteine sulfinic acid, which

spontaneously oxidizes to L-cysteate. L-cysteate could be further decarboxylated to taurine via

glutamate decarboxylase and subsequently converted via taurine dioxygenase to yield sulfite,

succinate, CO_2 and aminoacetaldehyde. It is currently unknown whether this pathway occurs in

Ca. Vesicomyosocius, but it represents a viable hypothesis for a host-independent regeneration

of succinate.

In agreement with Kuwahara et al. [21], our analyses indicate that *Ca*. Ruthia and *Ca*.

Vesicomysocius differ in several enzymes that affect the structure and composition of the

bacterial cell wall and membrane. In particular, symbionts of the *Ca*. Ruthia clade contained a

variety of exclusive genes involved in cell envelope biosynthesis, which may correspond to

clade- or species-specific responses to pathogens and/or interactions with their clam hosts.

Compared to *Ca*. Vesicomyosocius, multiple lineages of the *Ca*. Ruthia group retain remnants of

genes involved in anti-viral defense and transposition, possibly due to increased selective

pressures or higher frequency of homologous recombination resulting from leaky vertical

transmission. Previous studies suggested that occasional horizontal transmission is present in

various vesicomyid symbiont species [11, 12, 15] and seems to be mostly a result of host-to-host

transfer [13, 14] or inter-specific hybridization [16]. Occasional horizontal acquisition creates

opportunities for recombination between symbiont lineages that decelerates the increasing

genome reduction and the ultimate evolutionary enslavement of these endosymbionts by their

hosts [9]. *Candidatus* R. southwardae and *Ca*. R. phaseoliformis exhibited the lowest degree of

genome erosion among all vesicomyid symbionts analyzed, which could imply that gene

salvaging through homologous recombination is particularly frequent in these lineages. *Ca*. R.

southwardae and *Ca*. R. phaseoliformis are associated with two *Abyssogena* species, which

exclusively occur at abyssal depths and belong to a more recently radiated group of vesicomyid

host taxa [18] where symbiont switching events have been observed [11, 15]. Thus, it is possible

that depth-related environmental stressors might select for gene retention, while a proclivity to

acquire foreign symbionts might favor recombination, thereby decreasing the rate of genome

reduction in these symbiont lineages. Increased taxonomic and population level sampling could

help to clarify the evolutionary processes and rates involved in gene content evolution in these

taxa.

The differential patterns of gene loss between *Ca*. Vesicomyosocius and *Ca*. Ruthia

reiterate that reductive genome evolution does not follow a universal trajectory but is a reflection

of the ecological and evolutionary context of the respective host-symbiont association [25, 26].

Convergent gene loss and pseudogenization imply common evolutionary pressures for some

genes, whereas selection on codons and lineage-specific gene retention imply niche-specific

adaptation in others. In both symbiont clades, physiological differences related to nickel,

cobalamin, oxygen and sulfur requirements appear to mediate microhabitat segregation and the

generation of divergent niche-specific gene pools. Future studies linking environmental data with

symbiont genomic information will be helpful to obtain further insights into the ecological basis

of reductive genome evolution in the symbionts of deep-sea vesicomyid clams.

Acknowledgements

We thank the able captains and crews of the R/Vs *Atlantis*, *Western Flyer* and *Point Lobos* as

well as the pilots of the submersibles *Alvin*, *Tiburon* and *Ventana* for supporting the sample

collections. We further thank N. Pratt and A. Baylay for their contributions to library preparation

and high-throughput sequencing at the National Oceanography Centre Genomics Facility. This

work was supported by grants of the David and Lucile Packard Foundation (to MBARI), the UK

Natural Environment Research Council (grant number NE/N006496/1 to C.R.Y.), the United

States National Science Foundation (grant number OCE-1736932 to R.A.B.) and National

Capability funding to the National Oceanography Centre (grant number NE/R015953/1). M.P.'s

contribution was funded through the Alexander Graham Bell Canada Graduate Scholarship and

the Michael Smith Foreign Study Supplements granted by the Natural Sciences and Engineering

Research Council of Canada.

Competing Interests

The authors declare no competing interests.

References

1. Russell SL. Transmission mode is associated with environment type and taxa across bacteria-
eukaryote symbioses: a systematic review and meta-analysis. FEMS Microbiol Lett. 2019;
366(3): fnz013.
2. Fisher RM, Henry LM, Cornwallis CK, Kiers ET, West SA. The evolution of host-symbiont
dependence. Nat Commun. 2017; 8: 15973.
3. Bennett GM, Moran NA. Heritable symbiosis: The advantages and perils of an evolutionary
rabbit hole. PNAS 2015; 112(33): 10169–10176.
4. Hansen AK, Moran NA. The impact of microbial symbionts on host plant utilization by
herbivorous insects. Mol Ecol. 2014; 23: 1473–1496.
5. Peek AS, Feldman RA, Lutz RA, Vrijenhoek RC. Cospeciation of chemoautotrophic bacteria
and deep-sea clams. PNAS 1998; 95(17): 9962–9966.
6. Childress JJ, Fisher CR, Favuzzi JA, Sanders NK. Sulfide and carbon dioxide uptake by the
hydrothermal vent clam, *Calyptogena magnifica*, and its chemoautotrophic symbionts.
Physiol. Zool. 1991; 64: 1444–1470.
7. Robinson JJ, Cavanaugh CM. Expression of form I and form II Rubisco in chemoautotrophic
symbioses: Implications for the interpretation of stable carbon isotope values. Limnol
Oceanogr. 1995; 40: 1496–1502.
8. Newton IL, Girguis PR, Cavanaugh CM. Comparative genomics of vesicomyid clam
(Bivalvia: Mollusca) chemosynthetic symbionts. BMC Genomics 2008; 9: 585.
9. Vrijenhoek RC. Genetics and evolution of deep-sea chemosynthetic bacteria and their
invertebrate hosts. In: Kiel S (ed). The Vent and Seep Biota. (Springer, Dordrecht,
Netherlands, 2010) pp 15–49.
10. Dubilier N, Bergin C, Lott C. Symbiotic diversity in marine animals: the art of harnessing
chemosynthesis. Nat Rev Microbiol. 2008; 6: 725–740.
11. Stewart FJ, Young CR, Cavanaugh CM. Lateral symbiont acquisition in a maternally
transmitted chemosynthetic clam endosymbiosis. Mol Biol Evol. 2008; 25: 673–687.
12. Stewart FJ, Young CR, Cavanaugh CM. Evidence for homologous recombination in
intracellular chemosynthetic clam symbionts. Mol Biol Evol. 2009; 26: 1391–1404.
13. Decker C, Olu K, Arnaud-Haond S, Duperron S. Physical proximity may promote lateral
acquisition of bacterial symbionts in vesicomyid clams. PLoS ONE 2013; 8: e64830.
14. Ikuta T, Igawa K, Tame A, Kuroiwa T, Kuroiwa H, *et al.* Surfing the vegetal pole in a small
population: extracellular vertical transmission of an 'intracellular' deep-sea clam symbiont. R
Soc Open Sci. 2016; 3: 160130.
15. Ozawa G, Shimamura S, Takaki Y, Takishita, Ikuta T, *et al.* Ancient occasional host
switching of maternally transmitted bacterial symbionts of chemosynthetic vesicomyid
clams. Genome Biol Evol. 2017; 9: 2226–2236.
16. Breusing C, Johnson SB, Vrijenhoek RC, Young CR. Host hybridization as a potential
mechanism of lateral symbiont transfer in deep-sea vesicomyid clams. Mol Ecol. 2019; 28:
4697–4708.

17. Kuwahara H, Takaki Y, Shimamura S, Yoshida T, Maeda T, *et al.* Loss of genes for DNA
recombination and repair in the reductive genome evolution of thioautotrophic symbionts of
Calyptogena clams. BMC Evol Biol. 2011; 11: 285.
18. Johnson SB, Krylova EM, Audzijonyte A, Sahling H, Vrijenhoek RC. Phylogeny and origins
of chemosynthetic vesicomyid clams. Syst Biodivers. 2017; 15(4): 346–360.
19. Newton IL, Woyke T, Auchtung TA, Dilly GF, Dutton RJ, *et al.* The *Calyptogena magnifica*
chemoautotrophic symbiont genome. Science 2007; 315: 998–1000.
20. Kuwahara H, Yoshida T, Takaki Y, Shimamura S, Nishi S, *et al.* Reduced genome of the
thioautotrophic intracellular symbiont in a deep-sea clam, *Calyptogena okutanii*. Curr. Biol.
2007; 17: 881–886.
21. Kuwahara H, Takaki Y, Yoshida T, Shimamura S, Takishita K, *et al.* Reductive genome
evolution in chemoautotrophic intracellular symbionts of deep-sea *Calyptogena* clams.
Extremophiles 2008; 12: 365–374.
22. Shimamura S, Kaneko T, Ozawa G, Matsumoto MN, Koshiishi T, *et al.* Loss of genes related
to Nucleotide Excision Repair (NER) and implications for reductive genome evolution in
symbionts of deep-sea vesicomyid clams. PLoS ONE 2017; 12: e0171274.
23. Goffredi SK, Barry JP. Species-specific variation in sulfide physiology between closely
related Vesicomyid clams. Mar Ecol Prog Ser. 2002; 225: 227–238.
24. Cruaud P, Decker C, Olu K, Arnaud-Haond S, Papot C, *et al*. Ecophysiological differences
between vesicomyid species and metabolic capabilities of their symbionts influence
distribution patterns of the deep-sea clams. Mar Ecol. 2019; 40: e12541.
25. Luo H, Huang Y, Stepanauskas R, Tang J. Excess of non-conservative amino acid changes in
marine bacterioplankton lineages with reduced genomes. Nat Microbiol. 2017; 2: 17091.
26. Baumgartner M, Roffler S, Wicker T, Pernthaler J. Letting go: bacterial genome reduction
solves the dilemma of adapting to predation mortality in a substrate-restricted
environment. ISME J. 2017; 11(10): 2258–2266.
27. Peek A, Gustafson R, Lutz R, Vrijenhoek R. Evolutionary relationships of deep-sea
hydrothermal vent and cold-water seep clams (Bivalvia: Vesicomyidae): Results from the
mitochondrial cytochrome oxidase subunit I. Mar Biol. 1997; 130: 151–161.
28. Andrews S. FastQC: a quality control tool for high throughput sequence data. 2010
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/
29. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics 2014; 30: 2114–2120.
30. Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn
graphs. Genome Res. 2008; 18: 821–829.
31. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, *et al*. SPAdes: a new genome
assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012;
19(5): 455–477.
32. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. Scaffolding pre-assembled contigs
using SSPACE. Bioinformatics 2010; 27(4): 578–579.
33. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, *et al.* The RAST Server: rapid
annotations using subsystems technology. BMC Genomics 2008; 9: 75.
34. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome
assemblies. Bioinformatics 2013; 29: 1072–1075.
35. Darling AE, Mau B, Perna NT. progressiveMauve: multiple genome alignment with gene
gain, loss and rearrangement. PLoS ONE 2010; 5: e11147.

36. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, *et al.* Anvi'o: an advanced analysis
and visualization platform for 'omics data. PeerJ 2015; 3: e1319.
37. Delmont TO, Eren AM. Linking pangenomes and metagenomes: the *Prochlorococcus*
metapangenome. PeerJ 2018; 6: e4320.
38. Tatusov RL, Galperin MY, Natale DA, Koonin EV. The COG database: a tool for genome-
scale analysis of protein functions and evolution. Nucleic Acids Res. 2000; 28(1): 33–36.
39. Huelsenbeck JP, Ronquist F. MrBAYES: Bayesian inference of phylogenetic trees.
Bioinformatics 2001; 17: 754–755.
40. Miller MA, Pfeiffer W, Schwartz T. Creating the CIPRES Science Gateway for inference of
large phylogenetic trees. In: Proceedings of the Gateway Computing Environments
Workshop (GCE). (New Orleans, LA, USA, 2010) pp 1–8.
41. Pruesse E, Peplies J, Glöckner FO. SINA: accurate high-throughput multiple sequence
alignment of ribosomal RNA genes. Bioinformatics 2012; 28: 1823–1829.
42. Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior summarization in
Bayesian phylogenetics using Tracer 1.7. Syst Biol. 2018; 67(5): 901.
43. Murrell B, Moola S, Mabona A, Weighill T, Sheward D, *et al.* FUBAR: A Fast,
Unconstrained Bayesian AppRoximation for inferring selection. Mol Biol Evol. 2013; 30(5):
1196–1205.
44. Murrell B, Wertheim JO, Moola S, Weighill T, Scheffler K, *et al*. Detecting individual sites
subject to episodic diversifying selection. PLoS Genet. 2012; 8: e1002764.
45. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately maximum-likelihood trees for
large alignments. PLoS ONE 2010; 5(3): e9490.
46. Spielman SJ, Weaver S, Shank SD, Magalis BR, Li M, *et al.* Evolution of viral genomes:
Interplay between selection, recombination, and other forces. In: Anisimova M (ed).
Evolutionary Genomics. Methods in Molecular Biology. (Humana, New York, NY, 2019) pp
427–468.
47. Richter M, Rosselló-Móra R. Shifting the genomic gold standard for the prokaryotic species
definition. PNAS 2009; 106: 19126–19131.
48. Nakagawa S, Takai K. Deep-sea vent chemoautotrophs: diversity, biochemistry and
ecological significance. FEMS Microbiol Ecol. 2008; 65: 1–14.
49. Klatt JM, Polerecky L. Assessment of the stoichiometry and efficiency of CO_2 fixation
coupled to reduced sulfur oxidation. Front Microbiol. 2015; 6: 484.
50. Kato K, Tanaka R, Sano S, Tanaka A, Hosaka H. Identification of a gene essential for
protoporphyrinogen IX oxidase activity in the cyanobacterium *Synechocystis* sp. PCC6803.
PNAS 2010; 107(38): 16649–16654.
51. Zhang Y, Mandava CS, Cao W, Li X, Zhang D, *et al*. HflX is a ribosome-splitting factor
rescuing stalled ribosomes under stress conditions. Nat Struct Mol Biol. 2015; 22(11): 906–
913.
52. Eitinger T, Suhr J, Moore L, Smith JAC. Secondary transporters for nickel and cobalt ions:
theme and variations. Biometals 2005; 18(4): 399–405.
53. Connolly B, Parsons CA, Benson FE, Dunderdale HJ, Sharples GJ, *et al*. Resolution of
Holliday junctions in vitro requires the *Escherichia coli ruvC* gene product. PNAS 1991;
88(14): 6063–6067.
54. Janson EM, Stireman III JO, Singer MS, Abbot P. Phytophagous insect–microbe mutualisms
and adaptive evolutionary diversification. Evolution 2008; 62(5): 997–1012.

55. Patiño-Navarrete R, Moya A, Latorre A, Peretó J. Comparative Genomics of *Blattabacterium*
cuenoti: The frozen legacy of an ancient endosymbiont genome. Genome Biol Evol. 2013;
5(2): 351–361.
56. Manzano-Marín A, Coeur d'acier A, Clamens A, Orvain C, Cruaud C, et al. Serial horizontal
transfer of vitamin-biosynthetic genes enables the establishment of new nutritional symbionts
in aphids' di-symbiotic systems. ISME J. 2020; 14: 259–273.
57. Chong R, Park H, Moran NA. Genome evolution of the obligate endosymbiont *Buchnera*
aphidicola. Mol Biol Evol. 2019; 36(7): 1481–1489.
58. Williams LE, Wernegreen JJ. Genome evolution in an ancient bacteria-ant symbiosis:
parallel gene loss among *Blochmannia* spanning the origin of the ant tribe Camponotini.
PeerJ 2015; 3: e881.
59. González JC, Banerjee RV, Huang S, Sumner JS, Matthews RG. Comparison of cobalamin-
independent and cobalamin-dependent methionine synthases from *Escherichia coli*: two
solutions to the same chemical problem. Biochemistry 1992; 31(26): 6045–6056.
60. McCutcheon JP, McDonald BR, Moran NA. Convergent evolution of metabolic roles in
bacterial co-symbionts of insects. PNAS 2009; 106(36): 15394–15399.
61. Shelton AN, Seth EC, Mok KC, Han AW, Jackson SN, et al. Uneven distribution of
cobamide biosynthesis and dependence in bacteria predicted by comparative genomics.
ISME J. 2019; 13: 789–804.
62. Torrents E. Ribonucleotide reductases: essential enzymes for bacterial life. Front Cell Infect
Microbiol. 2014; 4: 52.
63. Borovok I, Gorovitz B, Schreiber R, Aharonowitz Y, Cohen G. Coenzyme B12 controls
transcription of the *Streptomyces* class Ia ribonucleotide reductase nrdABS operon via a
riboswitch mechanism. J Bacteriol. 2006; 188(7): 2512–2520.
64. Boer JL, Mulrooney SB, Hausinger RP. Nickel-dependent metalloenzymes. Arch Biochem
Biophys. 2014; 0: 142–152.
65. Hattori M, Iwase N, Furuya N, Tanaka Y, Tsukazaki T, *et al*. Mg^{2+}-dependent gating of
bacterial MgtE channel underlies Mg^{2+} homeostasis. EMBO J. 2009; 28(22): 3602–3612.
66. Moreno-Vivián C, Cabello P, Martínez-Luque M, Blasco R, Castillo F. Prokaryotic nitrate
reduction: molecular properties and functional distinction among bacterial nitrate
reductases. J Bacteriol. 1999; 181(21): 6573–6584.
67. Marcia M, Ermler U, Peng G, Michel H. A new structure-based classification of
sulfide:quinone oxidoreductases. Proteins 2010; 78: 1073–1083.
68. Gregersen LH, Bryant DA, Frigaard NU. Mechanisms and evolution of oxidative sulfur
metabolism in green sulfur bacteria. Front Microbiol. 2011; 2: 116.
69. Shuman KE, Hanson TE. A sulfide:quinone oxidoreductase from *Chlorobaculum tepidum*
displays unusual kinetic properties. FEMS Microbiol Lett. 2016; 363: fnw100.
70. Pruski AM, Fiala-Médioni A. Stimulatory effect of sulphide on thiotaurine synthesis in three
hydrothermal-vent species from the East Pacific Rise. J Exp Biol. 2003; 206: 2923–2930.
71. Rosenberg NK, Lee RW, Yancey PH. High contents of hypotaurine and thiotaurine in
hydrothermal-vent gastropods without thiotrophic endosymbionts. J Exp Zool Pt A 2006;
305(8): 655–662.
72. Brand GL, Horak RV, Le Bris N, Goffredi SK, Carney SL, *et al*. Hypotaurine and thiotaurine
as indicators of sulfide exposure in bivalves and vestimentiferans from hydrothermal vents
and cold seeps. Mar Ecol. 2007; 28(1): 208–218.

73. Dominy Jr JE, Simmons CR, Karplus PA, Gehring AM, Stipanuk MH. Identification and
characterization of bacterial cysteine dioxygenases: a new route of cysteine degradation for
eubacteria. J Bacteriol. 2006; 188(15): 5561–5569.

Figure Legends

Figure 1 Bayesian phylogeny of the *16S* rRNA gene for vesicomyid symbionts (A) and pangenomic comparison based on 3,208 gene clusters organized according to their distribution across genomes. Each symbiont genome is represented as a circle layer, with *Ca*. Vesicomyosocius lineages shown in green and *Ca*. Ruthia lineages shown in red. Dark color shades within layers denote presence of a gene cluster, whereas light shades denote absence. Gene cluster groups representative of the core genome (grey), *Ca*. Vesicomyosocius-specific genome (green) and *Ca*. Ruthia-specific genome (red) are indicated. The top right dendrogram shows phylogenomic relationships among symbiont genomes based on 739 single copy gene clusters. Average nucleotide identities (80–100%) among genomes are presented below the dendrogram based on a color gradient, where darker shades indicate higher similarities.

Figure 2 (A) Core genome of vesicomyid symbiont based on categorization into Clusters of Orthologous Groups (COGs). (B) Functional categorization of gene clusters specific to *Ca*. Vesicomyosocius and *Ca*. Ruthia. Some gene clusters are functionally related to various metabolic pathways and are therefore classified into multiple categories.

Figure 3 Main differences in metabolic gene content between *Ca*. Vesicomyosocius and *Ca*. Ruthia and presence/absence of essential genes related to DNA repair, recombination and anti-viral defense. A detailed description is given in the manuscript text.

A

CP024634 *Bathymodiolus thermophilus* symbiont

CP060688 *Ca.* R. pliocardia

JACRUR000000000 *Ca.* R. phaseoliformis

CP000488 *Ca.* R. magnifica

JACRUS000000000 *Ca.* R. southwardae

CP060683 *Ca.* R. pacifica

CP060684 *Ca.* R. rectimargo

AP009247 *Ca.* V. okutanii

CP060687 *Ca.* V. soyoae 1

CP060686 *Ca.* V. soyoae 2

CP060680 *Ca.* V. diagonalis

CP060685 *Ca.* V. extenta

CP060681 *Ca.* V. gigas 1

CP060682 *Ca.* V. gigas 2

0.007

B

Ca. V. soyoae 2
Ca. V. soyoae 1
Ca. V. okutanii
Ca. V. gigas 2
Ca. V. gigas 1
Ca. V. extenta
Ca. V. diagonalis
Ca. R. southwardae
Ca. R. pliocardia
Ca. R. phaseoliformis
Ca. R. rectimargo
Ca. R. pacifica
Ca. R. magnifica

3,208 gene clusters

Phylogeny based on 739 single-copy gene clusters

Ca. R. magnifica
Ca. R. pacifica
Ca. R. rectimargo
Ca. R. phaseoliformis
Ca. R. pliocardia
Ca. R. southwardae
Ca. V. diagonalis
Ca. V. extenta
Ca. V. gigas 1
Ca. V. gigas 2
Ca. V. okutanii
Ca. V. soyoae 1
Ca. V. soyoae 2

Average Nucleotide Identity (80–100%)

Core gene clusters (822)

Ca. Ruthia clusters (46)

Ca. Vesicomyosocius clusters (6)

Table 1 General information about vesicomyid symbiont genomes compared in this study.

Symbiont	Accession No.	Genome size (Mb)	# Contigs	N50 (Mb)	GC (%)	Coverage (X)	# CDS	# RNAs	Reference
Ca. R. magnifica	CP000488	1.16	1	1.16	34.03	14.00	976	39	Newton et al. (2007)
Ca. R. pacifica	CP060683	1.18	1	1.18	36.58	140.20	1,456	38	This study
Ca. R. phaseoliformis	JACRUR000000000	1.53	8	0.37	36.93	69.48	2,210	39	This study
Ca. R. pliocardia	CP060688	1.23	1	1.23	36.98	113.20	1,643	39	This study
Ca. R. rectimargo	CP060684	1.19	1	1.19	36.69	90.80	1,476	40	This study
Ca. R. southwardae	JACRUS000000000	1.59	39	0.06	36.86	89.56	2,035	39	This study
Ca. V. diagonalis	CP060680	1.02	1	1.02	31.10	110.30	1,005	39	This study
Ca. V. extenta	CP060685	1.02	1	1.02	31.10	137.00	995	39	This study
Ca. V. gigas 1	CP060681	1.03	1	1.03	31.37	49.20	1,008	39	This study
Ca. V. gigas 2	CP060682	1.03	1	1.03	31.38	153.10	979	39	This study
Ca. V. okutanii	AP009247	1.02	1	1.02	31.59	9.00	937	38	Kuwahara et al. (2007)
Ca. V. soyoae 1	CP060687	1.02	1	1.02	31.63	89.40	992	39	This study
Ca. V. soyoae 2	CP060686	1.02	1	1.02	31.63	109.90	983	39	This study

Table 2 Sampling information for vesicomyid hosts.

Host species	Symbiont	Locality	Latitude	Longitude	Depth (m)	Habitat	Dive #[a]	Year
Abyssogena phaseoliformis	*Ca.* R. phaseoliformis	Aleutian Trench	54.3050	–157.2133	3550	Seep	–[b]	1996
Abyssogena southwardae	*Ca.* R. southwardae	Logatchev	14.7532	–44.9805	3038	Vent	A: 3133	1997
Archivesica diagonalis	*Ca.* V. diagonalis	Monterey Canyon	36.2274	–122.8796	3456	Seep	T: 488	2002
Archivesica gigas (1)	*Ca.* V. gigas 1	East Gorda Escarpment	40.3580	–125.0210	2094	Vent	T: 351	2001
Archivesica gigas (2)	*Ca.* V. gigas 2	Guaymas Transform Fault	27.3400	–111.2700	1754	Seep	T: 548	2003
Calyptogena magnifica	*Ca.* R. magnifica	East Pacific Rise	9.8480	–104.2935	2507	Vent	A: 3951	2004
Calyptogena pacifica	*Ca.* R. pacifica	Monterey Canyon	36.7739	–122.0488	650	Seep	V: 2555	2004
Calyptogena rectimargo	*Ca.* R. rectimargo	Monterey Canyon	36.6816	–122.1197	1540	Seep	V: 2338	2003
Phreagena extenta	*Ca.* V. extenta	Monterey Canyon	36.6088	–122.4366	2889	Seep	T: 406	2002
Phreagena okutanii	*Ca.* V. okutanii	Sagami Bay	37.9500	139.2000	1157	Seep	HD: 305	2004
Phreagena soyoae (1)	*Ca.* V. soyoae 1	Oregon Subduction Zone	44.6755	–125.1182	765	Seep	A: 2796	1994
Phreagena soyoae (2)	*Ca.* V. soyoae 2	Monterey Canyon	36.7762	–122.0842	985	Seep	V: 2059	2001
Pliocardia sp.	*Ca.* R. pliocardia	Blake Spur	32.4948	–76.1847	2155	Seep	A: 3710	2001

[a]Submersibles: A = *Alvin*, HD = *Hyper Dolphin*, T = *Tiburon*, V = *Ventana*
[b]Sampled with TV grab

Table 3 Sites under pervasive or episodic positive selection in 10 candidate genes based on FUBAR and MEME analyses. α = synonymous substitution rate at a site; $\beta+$ = non-synonymous substitution rate at a site for the positive/neutral evolution component; Pr = posterior probability (a value $\geqq 0.9$ indicates strong evidence for positive selection); q+ = very approximate proportion of branches evolving under positive selection; LRT = likelihood ratio test statistic for episodic diversification; p = p-value for episodic diversification (a value $\leqq 0.1$ indicates positive selection).

		FUBAR			MEME				
	Site	α	$\beta+$	Pr	α	$\beta+$	q+	LRT	p
cdo	65	1.4130	15.5050	0.9044	0.0000	94.6310	0.1380	4.8230	0.0413
	156	–	–	–	0.0000	1.2290	1.0000	3.3470	0.0890
hupE	39	2.7030	15.5090	0.9157	0.0000	33.0330	0.4180	5.2300	0.0335
	49	–	–	–	1.2160	261.8800	0.1010	5.7280	0.0259
	86	–	–	–	0.0000	6.1150	0.2910	4.5500	0.0476
	134	–	–	–	0.0000	4.7840	0.4770	5.1880	0.0342
	333	0.9250	8.8230	0.9314	0.0000	3.8540	1.0000	3.2000	0.0961
	373	–	–	–	1.1930	2433.1130	0.0970	4.3520	0.0527
metE	49	–	–	–	0.0000	126.9650	0.0970	5.3060	0.0322
	218	0.8680	8.8950	0.9377	0.0000	2.6500	1.0000	3.9300	0.0656
	234	–	–	–	0.7460	15.5340	0.2040	3.6460	0.0761
metH	123	–	–	–	0.0000	53.7870	0.0980	3.1960	0.0963
	129	–	–	–	0.0000	19.1500	0.3300	3.3170	0.0904
	615	–	–	–	0.6390	79.7860	0.0950	3.9620	0.0645
	621	–	–	–	0.4440	25.2060	0.1050	5.9500	0.0231
	654	–	–	–	0.0000	8.6840	0.1270	3.2490	0.0937
	749	–	–	–	0.4410	8.5390	0.3660	3.6720	0.0751

	902	–	–	–	0.0000	1.4390	0.3070	3.4000	0.0865
	953	–	–	–	0.3650	10.8580	0.1290	4.6050	0.0462
narG	138	–	–	–	0.0000	7.5290	0.2350	3.8440	0.0686
	451	4.2920	39.0210	0.9409	0.0000	21.5910	1.0000	2.4350	0.1441[†]
narH	54	–	–	–	0.0000	11.1380	0.2230	3.9820	0.0639
narI	–	–	–	–	–	–	–	–	–
narJ	211	–	–	–	0.0000	7.3140	0.2790	3.5150	0.0815
	216	–	–	–	0.0000	3.9790	0.3420	3.5750	0.0790
nasA	70	0.5960	12.9980	0.9580	0.0000	2.8050	1.0000	3.6770	0.0749
	79	–	–	–	0.0000	23.6900	0.1400	4.5030	0.0487
	451	1.4360	13.6900	0.9362	0.0000	3.6150	0.9990	2.1190	0.1707[†]
sqrI	4	0.6890	6.2540	0.9475	0.0000	1.5310	0.9970	2.9470	0.1098[†]
	168	–	–	–	3.2830	50.0800	0.0520	3.9720	0.0642
	366	–	–	–	0.6110	42.3580	0.1110	5.5300	0.0287
	428	–	–	–	0.0000	10000.0000	0.0830	3.7640	0.0715

[†] Test was not significant