Regular Paper

Subject Area (primary) Structure and Function of Cells; *Subject Area (secondary) Proteins, Enzymes and*

Metabolism

Glycosylphosphatidylinositol-anchoring is required for the proper transport and glycosylation of

classical arabinogalactan protein precursor in tobacco BY-2 cells

Daiki Nagasato[1], Yuto Sugita[1], Yuhei Tsuno[1], Rutsuko Tanaka[2], Maki Fukuda[3], Ken Matsuoka[1,3,4]

[1]Graduate School of Bioresource and Bioenvironmental Science, Kyushu University, Fukuoka, Japan

[2]Chikushigaoka High School, Fukuoka, Japan

[3]School of Agriculture, Kyushu University, Fukuoka, Japan

[4]Faculty of Agriculture, Kyushu University, Fukuoka, Japan

15
16 **Running Head:** Role of GPI-anchor on AGP transport and maturation
17
18

Correspondence to: Ken Matsuoka, Department of Bioscience and Biotechnology, Faculty of Agriculture,

Kyushu University, 744 Motooka, Nishi-ku, Fukuoka 819-0395, Japan.

Email: kenmat@agr.kyushu-u.ac.jp

Telephone and Fax: +81-92-802-4713

Number of black and white figures, 6; color figures, 3; and type and number of supplementary materials,

black and white figures, 3; color figure, 1; movie, 1; tables, 4; document, 2.

Abstract

Arabinogalactan proteins (AGPs) are extracellular proteoglycans with many O-linked glycan chains.

Precursors to many AGPs contain a C-terminal signal for the addition of a glycosylphosphatidylinositol

(GPI)-anchor, but the role of this modification has not been elucidated. NtAGP1, a tobacco precursor to AGP,

comprises a signal peptide, an AGP-coding region, and a GPI-anchoring signal, and it is classified as a

member of the classical AGP family. Using green fluorescent protein (GFP) and sweet potato sporamin (SPO)

as tags and tobacco BY-2 cells as the host, we analysed the transport and modification of NtAGP1. The fusion

protein of GFP or SPO and NtAGP1 expressed in BY-2 cells migrated as a large smear on

SDS-polyacrylamide gel. A confocal microscopic analysis indicated that the GFP and NtAGP1 fusion protein

localized to the plasma membrane (PM) and intracellular structures. Fractionation studies of microsomes

indicated that most of the fusion protein of SPO and NtAGP1 (SPO-AGP) localized to the PM. In contrast,

the expression of mutants without a GPI-anchoring signal yielded several forms. The largest forms migrating

as large smears on the gel were secreted into the culture medium, whereas other forms were recovered in the

endomembrane organelles. A comparison of the glycan structures of the microsomal SPO-AGP and the

secreted mutant SPO-AGP without a GPI-anchoring signal using antibodies against AGP glycan epitopes

indicated that the glycan structures of these proteins differ. These observations indicate that a GPI-anchoring

signal is required for both the proper transport and glycosylation of the AGP precursor.

Key words: AGP, arabinogalactan protein, GPI anchoring, transport pathway, glycan structure

Introduction

Arabinogalactan proteins (AGPs) are plant cell wall proteoglycans with diverse functions (Pereira et al. 2015; Hromadová et al. 2021). Most of them contain many arabinogalactan (AG)-type glycan chains attached to Hyp residues that are post-translationally generated by the action of prolyl hydroxylases (Showalter et al. 2010). These proteins are classified into at least five groups based on their domain composition (Showalter et al. 2010). One of these groups consists of precursors to the classical AGPs; each of these precursors is made up of a signal peptide for the translocation of the endoplasmic reticulum (ER) membrane, glycosylation domains with many proline residues, and a signal sequence for the attachment of a glycosylphosphatidylinositol (GPI) anchor (Showalter et al. 2010; Yeats et al. 2018). Some members of other classes of AGP precursor do not contain a GPI-anchoring signal (Showalter et al. 2010).

GPI anchors are a class of lipid anchors that retain proteins at the surface of the plasma membrane (PM), especially within the sphingolipid-sterol-rich lipid domain (Liu and Fujita 2020). Based on analyses of mammalian cells and baker's yeast, the biosynthesis of GPI anchors and their subsequent modification during transport to the PM proceed as follows. First, the biosynthesis of the GPI anchor starts at the ER, and then the pre-assembled GPI anchor is transferred from the ER to the carboxy-terminus of proteins that contain the GPI-anchoring signal to yield a GPI-anchored protein. With the assistance of p24 proteins, GPI-anchored proteins are efficiently sorted into coat protein complex II (COPII) vesicles at the ER exit site and transported to the Golgi apparatus (Bernat-Silvestre et al. 2020; Muñiz and Riezman 2016). During the subsequent transport to the PM by passing through the Golgi apparatus, the lipid part of the anchor undergoes remodeling and changes its structure (Liu and Fujita, 2020).

Most of the orthologs of genes that are indispensable for the biosynthesis and remodeling of GPI anchors in mammals and yeast are found in higher plants (Beihammer et al. 2020). Only a limited number of genes that are predicted to be involved in this step have been characterized to date (Bernat-Silvestre et al. 2021; Lin et al. 2022; Xu et al. 2022), in part because some of their null mutants are embryo-lethal (Yeats et al. 2018).

The biosynthesis of AG glycan on an AGP starts at the ER and *cis*-side of the Golgi apparatus, where

Pro residues are converted to Hyp residues by the action of prolyl-hydroxylases localized in the ER and the

cis-side of the Golgi apparatus (Yuasa et al. 2005; Velasquez et al. 2015; Parsons et al. 2018). The subsequent

addition of galactose to the Hyp residue by Hyp galactosyltransferases (Oka et al. 2010; Basu et al. 2013,

2015; Ogawa-Ohnishi and Matsubayashi 2015) and the building of the AG glycan on the galactose residue in

the Golgi apparatus allows the maturation of the AGP precursor (reviewed in Showalter and Basu 2106).

During or after maturation, glycosylated and GPI-anchored AGPs are transported to the PM, and some of

them are released to the extracellular space by the action of phospholipase in the cell wall. Several enzymes

involved in AGP glycan synthesis have been characterized (reviewed in Showalter and Basu 2016), but such

enzymes have not yet been characterized completely based on the structure of the AGP glycans from tobacco

BY-2 cells and Arabidopsis (Tan et al. 2004; Tryfona et al. 2015; Showalter and Basu, 2016).

As both the remodeling of the GPI anchors and the maturation of AGP glycans takes place in the

Golgi apparatus, there may be a linkage between these two modifications. The Golgi apparatus is an organelle

located in the endomembrane transport system, and this system directs proteins to several destinations

including the PM, extracellular space, and vacuoles. As such, GPI anchoring might play a role in the proper

transport of proteins. However, this possibility has not been adequately investigated, especially with respect

to the transport of AGP. It has been reported that GPI anchoring is indispensable for PM localisation of the

Citrin-fused FLA4 protein, a fasciclin-domain containing AGP in Arabidopsis (Xue et al. 2017). In this case,

most of the green fluorescence of the mutant Citrin-FLA4 lacking a GPI-anchoring signal appeared to be

retained in the ER, and only a part of the green fluorescence from this protein seemed to be secreted to the

extracellular space, although this partial secretion was sufficient to complement the mutant phenotype of fla4

(Xue et al. 2017). Although this observation indicates that GPI anchoring is required for the proper

localisation of this protein, this observation may not be applicable to other classes of AGP; e.g., classical AGP,

which does not contain a fasciclin domain, and almost all of the mature polypeptide region is predicted to be

heavily glycosylated with AG glycans (Showalter et al. 2010).

Here we investigated the role of GPI anchoring in the transport and glycosylation of tobacco NtAGP1

(GenBank no. BAU61512), which is a classical AGP expressed in tobacco BY-2 cells, by expressing two

different protein-tagged versions in tobacco BY-2 cells. We present evidence that GPI anchoring is required

for both the proper transport and the proper glycosylation of this protein.

Results

Expression of the GFP-NtAGP1 fusion protein and its mutant without a GPI-anchoring signal in tobacco

BY2-cells

The NtAGP1 was used as a model protein for classical AGP in this study. The amino acid sequence of

this protein is nearly identical to that of *Nicotiana alata*-style AGP (Du et al. 1994). NtAGP1 is encoded by

the expressed sequence tag (EST) clone BY28237 isolated from the mixture of cDNA prepared from various

hormone-treated BY-2 cells (Galis et al. 2006). Four EST clones that have identical or nearly identical

nucleotide sequences have been identified in our EST collections (Suppl. Table S1). Among them, three were

obtained from the cDNA library from the hormone-treated BY-2 cells (Galis et al. 2006), and the fourth was

from the mixture of cDNA prepared from lag-, log- and stationary growth-phase cells (Matsuoka et al. 2004).

An expression analysis using microarrays (Matsuoka et al. 2004) indicated that the expression level of all of

these clones was higher than average in both lag and log phases, with nearly equal levels of expression at the

stationary phase (Suppl. Table S1). A BLASTN search against tobacco EST sequences using the nucleotide

sequence for NtAGP1 identified 43 EST sequences (Suppl. Table S2). These sequences were obtained from a

number of different cDNA libraries from various tobacco organs, including a female gamete, two-celled

proembryo, leaf, flower, root, germinating seeds, and mixture of several tissues. NtAGP1 and its orthologous

genes are thus expressed in many tobacco organs and at major growth phases of tobacco BY-2 cells.

The precursor to NtAGP1 consists of 132 amino acids (aa): a 21-aa signal peptide, an 86-aa

glycosylation domain, and a 25-aa GPI-anchoring signal. Within the glycosylation region there are 23 Pro

residues, all of which are surrounded with amino acids that make up the targeting sites for prolyl

hydroxylases (Shimizu et al. 2005). Among these proline residues, 15 fit to the arabinogalactosylation site

(Shimizu et al. 2005), and the other eight are contiguous prolines that might form the site for the attachment

of oligo-arabinose (Xu et al. 2008). Because most of the distances between the proline residues in NtAGP1

are ≤5 aa, and because the minimum length of a peptide epitope is six aa residues (Singh et al. 2013), we

thought that it would be almost impossible to generate an antibody against the peptide sequence of NtAGP1

to detect the native protein in tobacco tissues and cells. We thus adopted the alternative approach of

expressing fusion proteins consisting of NtAGP1 and protein tags and characterising these proteins in tobacco

BY-2 cells.

We first generated a construct for the expression of a fusion protein containing green fluorescence

protein (GFP) between the signal peptide and the subsequent region of NtAGP1 (GFP-AGP). We also

generated a mutant of this protein without a GPI-anchoring signal (GFP-AGPΔC) and a signal peptide with

GFP only (GFP) (Fig. 1A). These proteins as well as the fusion proteins of the PM water channel and GFP

(PIP-GFP: Yamauchi et al. 2003) were expressed in tobacco BY-2 cells, and the resulting transformed cells

were cultured in suspension.

We then analysed the GFP fluorescence of proteins in these transformed cell cultures that were grown

for either 3 days (to the logarithmic growth phase) or 7 days (to the stationary growth phase) after the

separation of proteins by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and

fluorescence recording (Fig. 1B). The fluorescence of GFP-AGP migrated as a large smear at around the

200-kDa position, and the intensity of this signal was stronger at 7 days. In the case of GFP-AGPΔC, at least

three smear bands were observed: two bands that migrated to the 120-kDa and 48-kDa positions, and a

relatively sharp band that migrated to around the 37-kDa position, the size of which corresponded to the

non-glycosylated form of this approx. 35-kDa protein (Fig. 1B). Among these proteins, the largest form

migrated more slowly at 7 days than that at 3 days on the SDS-polyacrylamide gel. The control proteins, GFP

and PIP-GFP, migrated to the corresponding position on the SDS-polyacrylamide gel, as expected.

To determine the localisation of these GFP fusion proteins, cultures of the transformed cells were

separated into medium, cell wall, cellular membrane, and cellular soluble fractions. Proteins in these fractions

were separated, and the distribution of the GFP fusion proteins was analysed (Fig. 1C). GFP-AGP was

recovered predominantly in the membrane fraction and to a lesser extent in the soluble fraction, and trace

amounts were present in the cell walls and culture medium. In the case of GFP-AGPΔC, most of the large

smear was recovered in the medium, and a small amount of this form was also found in the membrane and

soluble fractions. In contrast, almost all of the 48-kDa form was recovered in the membrane fraction, and the

smallest 37-kDa form was recovered in the soluble fraction. In the case of PIP-GFP, which was used as the

fractionation control, most of this protein was recovered in the membrane fraction with a trace amount in the

cell walls, and faint signals of smaller forms, possibly truncated ones, were detected in the soluble fraction.

We next addressed the distribution of the GFP fluorescence from 7-day-old cells expressing either

GFP-AGP or GFP-AGPΔC by a confocal laser scanning microscopy (CLSM) examination (Fig. 2). Almost

all of the green fluorescence in the GFP-AGP-expressing cells was localized at the cell surface, possibly at

the PM, and a weak signal was observed in the vacuoles as well as in unidentified punctate structures (Fig.

2A, GFP-AGP). The intensities of fluorescence varied in the cells, and the cell-surface signal seemed

nonuniform. In contrast to the cell-surface pattern of GF-AGP signal, cells expressing GFP-AGPΔC did not

show such a pattern of fluorescence; they did exhibit weak green fluorescence in the vacuoles with an

unidentified punctate structure with green fluorescence close to the cell wall (Fig. 2, GFP-AGPΔC).

To determine whether the GFP-AGP signal was localized in the PM, we incubated the cells for 5 min

with FM 4-64 to visualize the plasma membrane, and both the GFP and FM 4-64 fluorescence were recorded

(Fig. 3A). The GFP and FM 4-64 fluorescence colocalized well at the edges of the cells. The quantification of

signal intensities on a line along the short axis at the middle of the cell (Fig. 3A, merged image, yellow line)

indicated that the majority of the GFP fluorescence merged with the FM 4-64 fluorescence (Fig. 3A graph),

although some GFP signal was also observed at the inside of the FM 4-64 signal.

To confirm whether a significant proportion of green fluorescence from GFP-AGP-expressing cells

was localized to the PM, we stained the cells with FM 4-64 and observed the cells under plasmolysis (Fig.

2B). The clear lines of a green fluorescence signal, which corresponded to Hechtian strands, were visible

between the PM and cell walls. At the same position, Hechtian strands with FM 4-64 fluorescence were

present. Some of the fluorescence also seems attached to the cell wall, and some was present in the cell.

These observations suggested that the majority of GFP-AGP was localized to the PM although some of it was

present in intracellular structures.

To further confirm whether the majority of the GFP-AGP signal at the edge of the cells indicates the

PM localisation, we stained the cytoplasm with 5- (and 6-) chloromethyl SNARF-1 acetate (SNARF) and

compared the localisation of the GFP fluorescence with that of SNARF (Fig. 3C). The majority of the

SNARF fluorescence localized close to the cell wall, in a pattern which was similar to that of GFP. However,

a close inspection of the merged images and the quantification of signal intensities on a line along the short

axis at the middle of the cell (Fig. 3B, merged image, yellow line) indicated that the peaks of the GFP

fluorescence were one or two pixels outward from the peak of SNARF fluorescence (Fig. 3B graph). These

observations further confirmed that a significant proportion of GFP-AGP was localized to the PM.

As described above, optical section images suggested that the GFP fluorescence of GFP-AGP on the

plasma membrane was not distributed evenly on the plasma membrane. To assess the non-uniform

localisation of GFP-AGP at the PM, we collected Z-stack images by CLSM and then reconstituted a 3D

movie (Suppl. Movie S1). The nonuniform distribution of GFP fluorescence was apparent in the reconstituted

3D movie. These data indicated that GFP-AGP was predominantly localized to the PM, with a nonuniform

distribution.

Although these data suggested that the majority of the NtAGP1 is localized to the PM, we could not confirm

this localisation based on the result of the expression of only one fusion protein, and it was difficult to further

characterize GFP-AGP, for several reasons. (1) The green fluorescence from GFP-AGP was unstable in the

stably transformed tobacco BY-2 cells, and within six weeks of subculture, almost all of the cells stopped

emitting green fluorescence. (2) The signal peptide-GFP fusion protein was not fully secreted to the culture

medium from tobacco BY-2 cells (Mitsuhashi et al. 2000). Thus, GFP alone may not be a good cargo for

secretion. (3) We have a fluorescence pulse-chase system that uses a photo-convertible fluorescence protein,

mKikGR (Habuchi et al. 2008) which has a higher-order structure that is similar to that of GFP, to monitor the

turnover of proteins (Abiodun and Matsuoka 2013). We thus attempted to apply this system to address the

transport of NtAGP1 by expressing the fusion protein of mKikGR and NtAGP1 in tobacco BY-2 cells.

Unfortunately, the fusion protein migrated with three distinct bands: a weak large smear that migrated at a

position similar to that of GFP-AGP, a predominant band around 60 kDa, and a weaker band around 50 kDa

(Suppl. Fig. S1). We therefore tested the transport and modification of NtAGP1 using a protein that is

structurally distinct from GFP as a tag in order to determine the localisation and transport of this protein and

its mutant lacking a GPI-anchoring signal.

Expression and localisation of the sporamin-NtAGP1 fusion protein and its mutant lacking an GPI-anchoring

signal in tobacco BY-2 cells

We used sweet potato sporamin (SPO) as another protein tag to address the transport, localisation, and

modification of NtAGP1 and its derivative lacking a GPI-anchoring signal. Sporamin is a monomeric soluble

and non-glycosylated storage protein of sweet potato (Maeshima et al. 1985) localized in the vacuoles

(Hattori et al. 1988). The expression of a mutant precursor to sporamin, which does not contain the propeptide

region but does contain the signal peptide for secretion, allowed the efficient secretion of this protein to the

medium   from transformed tobacco BY-2 cells (>90% of this pulse-labeled protein is secreted within 2 h of

chase) (Matsuoka and Nakamura 1991). The junction region between the propeptide and the mature sporamin

surrounding the 36th Pro residue consists of a cryptic Hyp O-glycosylation site for AG, and thus both a

wild-type precursor to sporamin as well as mutant precursors with a disrupted vacuolar targeting signal are

O-glycosylated when expressed in tobacco BY-2 cells (Matsuoka et al. 1995, Matsuoka and Nakamura 1999).

The efficiency of the secretion of mutants with a disrupted vacuolar targeting signal is comparable to

that of the mutant without the propeptide (Matsuoka and Nakamura 1999). In other words, the secretion

efficiency of sporamin is not affected by the presence or absence of O-glycosylation. We thus chose sporamin

as another protein tag to analyse the transport and modification of NtAGP1 and its mutant lacking a

GPI-anchoring signal.

The expression constructs for the signal peptide-sporamin-mature NtAGP1 with a GPI-anchoring

signal (SPO-AGP) were expressed in tobacco BY-2 cells, as were the expression constructs for the signal

peptide-sporamin-mature NtAGP1 without a GPI-anchoring signal (SPO-AGPΔC) (Fig. 4A). Cell and

medium fractions were prepared from stationary-grown transformed BY-2 culture, and then the intra- and

extra-cellular localisations of sporamin or its fusion proteins in these fractions were examined by

immunoblotting either directly without immunoprecipitation (Fig. 4B) or after target proteins were recovered

by immunoprecipitation (Fig. 4C). Immunoblotting without immunoprecipitation gave cross-reacted protein

bands in proteins from non-transformed cells (Fig. 4B, WT). Such signals were absent after

immunoprecipitation (Fig. 4C, WT). However, our comparison of the intensities of sporamin-related signals

on many immunoblots obtained under both these conditions indicated that the recovery by

immunoprecipitation varied among the proteins, especially SPO-AGP. We thus did not use

immunoprecipitation for the subsequent quantitative analysis, and we always included a negative control from

non-transformed cells.

SPO-AGP was detected as a smear band close to the top of the SDS-polyacrylamide gel (Fig. 4B, C).

This observation indicated that SPO-AGP was glycosylated. Most of the smear-migrating protein was

recovered in the cell fraction (Fig. 4B). In contrast, SPO-AGPΔC was detected as three distinct species with

different migration positions; a large form that migrated at the 90–120-kDa position, an intermediate form at

the 36-kDa position and a small one at the 22-kDa position (Fig. 4B, C). A specific band that migrated at the

front of the electrophoresis gel may have been the degradation product of SPO (Fig. 4B). A large form of

SPO-AGPΔC was detected in both the cell and medium fractions. Because the calculated molecular weight of

non-glycosylated SPO-AGPΔC is 27.5, this observation indicated that most of the large-form SPO-AGPΔC

was glycosylated and secreted into the extracellular space. The intermediate and small forms of SPO-AGPΔC

were detected in cell fractions. This observation suggests that they were localized in the cells. The small form

may be the degradation product, since its size (22 kDa) was similar to that of SPO (Fig. 4C) and smaller than

the calculated size of the non-glycosylated form of SPO-AGPΔC.

To reveal the intracellular localisation of SPO-AGP and SPO-AGPΔC, we performed subcellular

fractionation by differential centrifugation. Cell lysates were centrifuged at 1,000g and the precipitate,

defined as the P1 fraction (which contained unbroken cells, nuclei, cell walls and cell wall-attached PM) was

collected. The supernatant was centrifuged at 10,000g and the precipitate (P10), which contained most of the

mitochondria and plastids, was collected. The resulting supernatant was further centrifuged at 100,000g and

the obtained precipitate (P100) contained microsomal membranes. The supernatant, which was defined as the

S fraction, was also collected. SPO-AGP was detected in the P1, P100, and S fractions (Fig. 4D). In the case

of SPO-AGPΔC, the large and small forms were detected in the P1 and S fractions, while the intermediate

form was observed in all fractions (Fig. 4E).

To address the localisation of SPO-AGP, which was recovered in the P100 fraction, we prepared

microsomal membranes from transformed BY-2 cells expressing SPO-AGP using buffers containing either

MgCl$_2$ or EDTA, and we then separated the membranes by equilibrium sucrose density gradient

centrifugation in the presence or absence of Mg^{2+}, respectively. The distribution of SPO-AGP in these

separated fractions was examined by immunoblotting (Fig. 5A). The distributions of marker proteins were

also analysed by immunoblotting with specific antibodies: Sec61 for the ER (Yuasa et al. 2005), vacuolar

pyrophosphatase (V-PPase) for tonoplast (Toyooka et al. 2009), GLMT1 for the Golgi apparatus (Liu et al.

2015), plasma membranous ATPase (P-ATPase) for the PM (Toyooka et al. 2009) and the PM water channel

(PIP) for the PM (Suga et al. 2001). In the presence of Mg^{2+} (Fig. 5A, +MgCl$_2$), SPO-AGP migrated in

several fractions, mainly in the fractions corresponding to 41–44(w/w)% sucrose. Quantification of the

intensities of these immunoblot signals indicated that the migration pattern of SPO-AGP was similar to those

of P-ATPase and PIP, the markers of the PM. In the absence of Mg^{2+} (Fig. 5A, +EDTA), SPO-AGP also

migrated with a peak at around 41–43(w/w)% sucrose. This migration pattern was also similar to those of

P-ATPase and PIP. These fractionation data indicated that SPO-AGP was localized to the PM.

To confirm the PM localisation of SPO-AGP with a different method and to compare the localisation

of this protein with endogenous AGP, we fixed transformed cells expressing SPO-AGP with formaldehyde

and stained them with anti-sporamin or LM6 (Willats et al. 1998), which is a monoclonal antibody against

oligo-arabinose epitope in both AGP glycan and rhamnogalacturonan I in pectin (Fig. 5B). LM6 recognized

the glycan on SPO-AGP by immunoblotting (see below). For the control, non-transformed BY-2 cells were

fixed and stained as well (Fig. 5B). During the fixation most of the cells shrank and separated from the cell

walls, and thus the localisation at the cell wall and cells was easily distinguished, although some of the

plasma membrane proteins were cross-linked to cell walls during fixation.

The staining of SPO-AGP-expressing cells with anti-native SPO antibody showed a clear signal at the

cell wall, PM, and punctate structures in the cell, although a weak intracellular puncta signal was also

observed when non-transformed cells were stained with this antibody (Fig. 5B). The pattern of staining of the

SPO-AGP-expressing cells with anti-sporamin antibody resembled the pattern of staining of wild-type cells

with LM6 (Fig. 5B), although the SPO-AGP signal at the cell wall showed a more uneven distribution

compared to that of LM6. Specificities of the staining including SPO-AGP dependency were confirmed by

staining both SPO-AGP-expressing cells and wild-type cells with either and both antibodies (Suppl. Figs. S2,

S3). The double immunostaining of SPO-AGP-expressing cells with anti-native SPO antibody and LM6 (Fig.

5C) revealed that both signals were colocalized well, although the SPO-AGP signal in the cells was more

punctate compared to that of LM6. A comparison of signal intensities on a line along the short axis at the

middle of the cell (Fig. 5C, merged image, yellow line) further confirmed that the majority of the SPO-AGP

signal merged with the LM6 signal, especially at the cell wall and PM (Fig. 5C graph).

To address the stability of SPO-AGP, we measured the level of SPO-AGP after protein synthesis was

stopped in the presence of cycloheximide. Although the average intensity of the SPO-AGP signal under the

cycloheximide condition was lower than the control value, the difference was not significant (Suppl. Fig. S4).

These observations suggest that SPO-AGP is not an unstable protein and that is localized mainly at the PM.

Intracellular localisation of SPO-AGPΔC

We prepared protoplasts from transformed BY-2 cells expressing SPO-AGPΔC in order to determine

which forms of SPO-AGPΔC were localized outside of the PM (Fig. 6A). The large form of SPO-AGPΔC

was detected only in cells and was not detected in protoplasts. The intermediate-form protein was detected in

both cells and protoplasts, although the amount was smaller in protoplasts than in cells. The small form of

SPO-AGPΔC was detected in both fractions in nearly equal amounts. This result indicated that the large-form

SPO-AGPΔC was localized in the periplasm and cell wall; the intermediate SPO-AGPΔC was localized in

both the periplasm and the cells, and the small SPO-AGPΔC was localized almost exclusively in the cells.

To address whether some of the intracellular SPO-AGPΔC was localized in vacuoles, we separated

protoplasts into vacuoplast and cytoplast fractions and analysed the amounts of the different forms of

SPO-AGPΔC in these fractions. The vacuoplasts are composed of vacuoles surrounded by the PM, and

cytoplasts are composed of cytoplasm, nuclei, and PM. We measured the protein concentration and assessed

the α-mannosidase activity in the protoplast and cytoplast fractions. We then used SDS-PAGE to separate the

proteins that were present in the protoplast and cytoplast fractions in nearly equal amounts and the proteins

that were present in the protoplast and vacuoplast fractions in amounts that are almost equivalent to the levels

of α-mannosidase activity. The different forms of SPO-AGPΔC in these fractions were detected by

immunoblotting (Fig. 6B). Mitochondrial porin was used as a fractionation control. The intermediate

SPO-AGPΔC was detected in the protoplast and cytoplast fractions in nearly equal amounts relative to the

protein, whereas the small form was detected in both the protoplast and vacuoplast fractions in amounts

similar to the activity of α-mannosidase. This result suggests that the small SPO-AGPΔC was localized in

vacuoles and the intermediate SPO-AGPΔC was localized in other intracellular compartments.

To investigate the localisation of the intermediate SPO-AGPΔC in cells, we prepared microsomal

fractions from transformed BY-2 cells expressing SPO-AGPΔC in buffers containing either $MgCl_2$ or EDTA.

The microsomal membranes were separated by equilibrium sucrose density gradient centrifugation in the

presence or absence of Mg^{2+}, respectively. The distribution of the intermediate SPO-AGPΔC in these

separated fractions was examined by immunoblotting and compared to the distributions of marker proteins in

the secretory organelles (Fig. 6C). In the presence of Mg^{2+} (Fig. 6C, $+MgCl_2$), the intermediate SPO-AGPΔC

migrated in several fractions with a peak at around 39–42(w/w)% sucrose. This migration pattern was similar

to that of Sec61, a marker of the ER membrane. In the absence of Mg^{2+} (Fig. 6C, +EDTA), the intermediate

SPO-AGPΔC also migrated with a peak at around 31–34(w/w)% sucrose, and the peak shifted to a

lower-density position compared to that in the presence of Mg^{2+}. This shift corresponded to that of the

migration pattern of Sec61 (an ER marker). These fractions did not contain P-ATPase or PIP but did contain

GLMT1, a marker of the Golgi apparatus. These fractionation data suggest that the intermediate form of

SPO-AGPΔC is localized predominantly in the ER and partly in the Golgi apparatus.

NtAGP1 is a GPI-anchored protein

The NtAGP1 precursor contains a GPI-anchoring signal at its C-terminus. Our present analyses

demonstrated that GFP-AGP and SPO-AGP are localized to the PM (Figs. 2, 5). To determine whether the

GPI-anchoring signal on the NtAGP1 precursor is functional and whether the GPI anchor attached to

NtAGP1 was responsible for attaching this protein to the PM, we analysed the distribution of these fusion

proteins recovered in the microsomal fraction by two-phase separation using the nonionic detergent Triton

X-114. This nonionic detergent can be used to make a uniform solution at low temperature, which can then be

separated into two phases, such as an aqueous (aqu) phase and a detergent-rich (det) phase, at higher

temperature. By taking advantage of this characteristic of the nonionic detergent, a cell extract can be

separated into soluble and peripheral membrane proteins that can be recovered in the aqu phase, plus integral

membrane proteins as well as lipid-anchored proteins that can be recovered in the det phase (Bordier 1981).

Approximately half of the GFP-AGP (Fig. 6A, left) as well as almost all of the SPO-AGP (Fig. 6B,

top left) and almost all of the PIP-GFP (an integral membrane protein used as an experimental control; Fig.

6A, right), was recovered in the det fraction. Under the same separation condition, almost all of the

smear-migrating GFP-AGPΔC and SPO-AGPΔC were recovered in the aqu fraction. These observations

suggested that about half of the GFP-AGP and almost all of the SPO-AGP were localized to the PM with their

GPI anchors.

To address whether this membrane-association is actually mediated by the GPI anchor attached to

these proteins, we carried out two-phase separation experiments in the presence of

phosphatidylinositol-specific phospholipase C (PI-PLC), which can cleave the phosphodiester bond in a GPI

anchor (reviewed by Hopper 2001). In the presence of PI-PLC, almost all of the GFP-AGP in the microsomal

fraction was recovered in the aqu phase (Fig. 6C, right). Likewise, about half of the SPO-AGP that was

recovered in the TX-114 phase was recovered in the det phase (Fig. 6D, top). Under the same condition,

vacuolar pyrophosphatase, which is an integral membrane protein used as a control, was recovered almost

exclusively in the det phase (Fig. 6D, bottom). These observations indicated that (*i*) roughly half of both

GFP-AGP and SPO-AGP was anchored to the PM by a GPI anchor that was sensitive to PI-PLC, and (*ii*)

NtAGP1 is a GPI-anchored protein at the PM.

GPI anchoring is required for the proper assembly of AGP glcyan

We showed that the migration positions of the smear-migrating SPO-AGP and SPO-AGPΔC on

SDS-polyacrylamide gel were different (Fig. 4A, B). We considered that this difference might be due to the

difference in the AG glycan chains on these proteins. To address this possibility, we examined the reactivity

of anti-AG glycan monoclonal antibodies (mABs) to these proteins. Among the mABs that we tested,

SPO-AGPΔC was specifically recognized by LM2 (Fig. 7B). Both proteins were recognized by LM6 (Fig.

7C). Other antibodies, namely PN16.4B4 (Norman et al. 1986), CCRC-M7 (Steffan et al. 1995), and

MAC204 and MAC207 (Bradley et al. 1988), did not recognize either protein. Because LM2 recognizes

terminal β-glucuronic acid (β-GlcA) (Yates et al. 1996), the absence of the recognition of SPO-AGP by this

antibody indicated that such a structure was absent in the glycan in SPO-AGP. This result also suggested that

GPI anchoring is required for the proper assembly of AGP glycan.

Discussion

In this work we investigated the role of the GPI anchor in the transport and modification of NtAGP1,

a classical AGP in tobacco BY-2 cells. The expression of the GFP fusion protein allowed us to detect the

smear-migrating GFP fusion protein with low mobility on SDS-polyacrylamide gels (Fig. 1). This behavior of

the fusion protein was consistent with the typical migration pattern of classical AGP in many plant species

(e.g., Putoczki et al. 2007; Lind et al. 1994; Maurer et al. 2010). In contrast, the fusion protein formed using

another fluorescent protein, mKikGR (Habuchi et al. 2008), yielded one slow-migrating and two

fast-migrating smear bands (Suppl. Fig. S1). To determine which of the observed migration behaviors of these

fusion proteins represents the true nature of NtAGP1, we characterized another fusion protein with sporamin

and found that this fusion protein migrated as a slow-migration smear band on SDS-polyacrylamide gels (Fig.

4). We therefore concluded that the slow migration and formation of a smear band on SDS-polyacrylamide

gels are characteristic of NtAGP1. These observations as well as the observation of multiple bands of

GFP-fused *Arabidopsis* AGP4, which is a member of the classical AGP family that is transiently expressed in

Arabidopsis seedlings (Bernat-Silvestre et al. 2020) also suggested that differences in the non-glycosylated

region of AGP as well as the expression level may affect the degree of modification of classical AGPs.

Our analyses of the localisation of GFP-AGP and SPO-AGP indicated that these proteins are

predominantly localized to the PM (Figs. 2, 3, 5), and that both proteins are attached to the PM as

GPI-anchored forms (Fig. 7). In addition, some of these proteins were present in intracellular structures (Figs.

3, 5) and in the medium. These observations suggested that NtAGP1 is a GPI-anchored PM protein with a

small pool in an uncharacterized intracellular structure, possibly related to an as-yet uncharacterized

intermediate component of AGP synthesis (Poulsen et al. 2014). Some other population of NtAGP1 is also

released to the extracellular space following the cleavage of its GPI anchor, possibly via the actions of

phospholipase in the extracellular space (Fig. 9, upper). Similar low-level secretion of AGP or fluorescence

protein-tagged AGP was reported in *Arabidopsis* (Darjania et al. 2002; Bernat-Silvestre et al. 2021; Xue et al.

2017). Such a low-level secretion of AGP might be the result of a slow release of this protein into the medium,

since we observed that the cycloheximide treatment did not cause a significant decrease of cellular SPO-AGP

within 24 hr (Suppl. Fig. S4). The presence of LM6 signal in both the cell wall and intracellular structures

(Fig. 5) suggested that such a distribution is not specific to NtAGP1; rather it is common to other AGPs

expressed in tobacco BY-2 cells, although we cannot rule out a possibility that significant proportion of the

LM6 signals represent the presence of rhamnogalacturonan I.

In contrast to the fusion proteins with an intact AGP with a GPI-anchoring signal, we detected mutant

fusion proteins without a GPI-anchoring signal (i.e., GFP-AGPΔC and SPO-AGPΔC) as multiple bands at

different migration positions on SDS-polyacrylamide gels (Figs. 1, 4). The glycosylated forms of these

proteins that migrated the most slowly and the most readily formed smear bands were predominantly

recovered in the medium fraction (Figs. 1, 4). The fractionation analysis of cells expressing these proteins

indicated that the intermediate forms were recovered partly in the membranous organelles in the cells and

partly in the periplasmic space (Figs. 1, 4). The migration positions of these proteins on SDS-polyacrylamide

gels were slower than those of the calculated size of non-glycosylated forms of these proteins, suggesting that

these proteins are modified to some extent, probably with glycans. Purification of these proteins and analyses

of their sugar compositions will clarify whether such modification takes place.

Our fractionation study of microsomes from SPO-AGPΔC-expressing cells indicated that the

intermediate SPO-AGPΔC was localized in the ER with partial localisation in the Golgi apparatus (Fig. 6).

The smallest form of this protein, which was similar in size to SPO, was recovered in the soluble fraction

from the cells, which consisted of the cytoplasm and vacuolar sap. The smallest form of SPO-AGPΔC was

also recovered in the vacuoplast fraction. These observations indicated that some of the SPO-AGPΔC protein

was transported to the vacuoles, while most of the AGP part of these proteins was cleaved off or degraded in

the vacuoles (Fig. 9, lower).

These observations indicated that GPI anchoring is not only essential for the proper and efficient

transport of NtAGP1 to the plasma membrane or extracellular space but also required for the efficient

glycosylation. Some of the inefficient-modified forms, i.e., the intermediate forms, were retained in the ER,

and some of them could also be transported to the vacuoles for degradation (Fig. 9, lower). These

observations are partly consistent with the case of a Citrin-tagged and GPI-anchoring signal-deleted

fasciclin-like AGP mutant (Xue et al. 2017). In that case, most of the mutant protein was retained in the ER.

In our present experiments, however, vacuolar localisation was observed. This difference in vacuolar

localisation between the Xue et al. report and our present study might be attributable to either the different

detection systems or different proteins used.

It has been reported that the vacuoles are scavenger organelles that degrade damaged proteins and

organelles. In yeast, it has been shown that vacuolar targeting is a mechanism for degrading improperly

folded proteins in the ER, and this transport is mediated by a receptor for vacuolar targeting (Hong et al.

1996). In plants, targeting of soluble proteins to the lytic vacuoles (such as the central vacuoles in tobacco

BY-2 cells) is mediated by VSR proteins (reviewed in Shimada et al. 2018). Large and hydrophobic residues,

such as Ile and Leu, in the vacuolar-sorting determinants in proteins play a pivotal role in the targeting

(Ahmed et al. 2000; Brown et al. 2003; Matsuoka and Nakamura 1995, 1999; Paris et al. 1997). It is thus

interesting that inefficient glycosylation of SPO-AGPΔC may cause the exposure of a large and hydrophobic

side chain in the mature region of AGP, and this characteristic of SPO-AGPΔC may have allowed the

targeting of this protein to the vacuole after recognition by VSR proteins, as in the case of yeast (Hong et al.

1986). In fact, the SPLA sequence at aa positions 67–70 of the NtAGP1 precursor resembles the NPIR

vacuolar sorting determinants of sporamin and aleurain propeptides (Matsuoka and Nakamura 1995, 1999;

Paris et al. 1997), and this region may be a good candidate for a region that is exposed under inefficient

glycosylation.

Not only the mislocalisation but also alteration of the glycan moiety was observed in the secreted

SPO-AGPΔC from SPO-AGP (Fig. 8). Secreted SPO-AGPΔC was recognized by the mAB LM2, which was

shown to recognize β-GlcA residues in AG glycan (Yates et al. 1996). It has been shown that the structure of

the glycan moiety of artificial AGP secreted from tobacco BY-2 cells contains β-GlcA, which is further

modified partly by rhamnose (Tan et al. 2004, 2010). In addition, the β-GlcA moiety of AGP is sometimes

methylated by the action of DUF576-family methyltransferases (Temple et al. 2019). It is thus possible that

(*i*) the glycan structure of secreted SPO-AGPΔC is similar to that of the artificial AGPs, which contain a

terminal β-GlcA, whereas such a structure is absent in SPO-AGP, and (*ii*) this difference is achieved by an

efficient modification with rhamnose or methyl residues. Further analyses of the sugar compositions of the

purified SPO-AGP and secreted SPO-AGPΔC proteins are necessary to fully explore these possibilities.

What causes the difference in glycan structure between SPO-AGP and secreted SPO-AGPΔC? One

possible answer may be related to the difference in transport routes of these proteins in the late secretory

pathway, with different distributions of the enzymes responsible for modification for the AGP glycan. A study

of tobacco protoplasts using a GFP-tagged pectin methylesterase inhibitor protein and its mutant without a

GPI-anchoring signal revealed that the late transport pathways of these proteins are different (De Caroli et al.

2011). The DUF576-family methyltransferases that act against AGP glycan have been detected not only in the

Golgi apparatus but also in small punctate structures that are distinct from the Golgi apparatus (Temple et al.

2019). Similarly, a subset of glycosyltransferases involved in AGP glycan biogenesis was identified in a

non-Golgi and non-TGN (trans-Golgi network) punctate structure (Poulsen et al. 2015). An analysis of the

secretome of *Arabidopsis* indicated that there are at least two distinct secretory pathways in *Arabidopsis* leaf

cells (Uemura et al. 2019). One of these pathways, which depends on SYP4-type SNAREs, is involved in the

secretion of many hydrolases (Uemura et al. 2019). In an analysis focusing on AGP in the data generated and

described in that paper (PRIDE dataset identifier PXD009099 by H. Nakagami et al.), only a small fraction of

AGPs with a GPI-anchoring signal (18%) showed significantly high abundance in apoplasts in a wild-type

relative to a mutant with SYP4-type SNARE proteins (Suppl. Table S3).

In contrast, when the same approach was taken against two groups of secretory hydrolases, i.e.,

pectic-lyase like proteins without a GPI-anchoring signal and β-galactosidase, a greater proportion of these

proteins showed a high level of secretion in the wild-type compared to the mutant SYP4-type SNARE

proteins (35% and 38%, respectively). In *Arabidopsis*, the SYP4-type SNAREs are localized in not only the

TGN adjusting the Golgi apparatus but also a small punctate structure that is called the Golgi-independent

TGN or free TGN (Viotti et al. 2010; Kang et al. 2011; Uemura et al. 2014). A similar structure called a

secretory vesicle cluster (SVC), which is involved in pectin secretion, is present in tobacco BY-2 cells

(Toyooka et al. 2009). Analyses of the proteome and glycome of secretory vesicles marked with SYP61

indicated that these secretory vesicles are rich in the SYP4-type SNARE proteins as well as glycans that are

recognized by mABs against pectin (Drakakaki et al. 2012; Wilkop et al. 2019). Although glycans in the same

vesicle fraction are also recognized by mABs against AGP, this recognition is weaker than that in the cell wall

fraction, in contrast to the recognition by mABs that recognize pectin (Fig. 3 in Wilkop et al. 2019).

Collectively, these observations suggest that the GPI-anchored AGP precursor would be

predominantly transported from the TGN to an uncharacterized punctate organelle, where a subset of

glycosyltransferases and/or methyltransferases localize (Poulsen et al. 2015), and would then be transported

to the PM after efficient modification of the glycan in the compartment (Fig. 9 AGP). The absence of a GPI

anchor in the AGP precursor prevents taking this route from the TGN, and it passes through the default

secretion pathway; this mis-sorting limits the final maturation of AGP glycan.

Another possible explanation for the difference in glycan structure is that the accessibility to the

modification enzymes might not be efficient in the absence of a GPI anchor. Because most of the

glycosyltransferases and methyltransferases in the secretory pathway are integral membrane proteins whose

catalytic domains are located proximately to the phospholipid bilayer, the GPI anchoring of the AGP

precursor will bring the protein close to the catalytic sites of these enzymes. Without a GPI anchor, such

access would not be efficient, and this characteristic would prevent the efficient maturation of the glycan side

chain. A third possibility is that the transport speed is different between SPO-AGP and SPO-AGPΔC, and this

difference contributes to the difference in the glycan side chain. Future studies to address the pathways and

kinetics of transport and modification of AGP using SPO-AGP as a model will reveal which scenario or

combination of scenarios is most likely.

Materials and Methods

An outline of the methods and critical information about the materials used in this study are provided

next; more detailed information about the plasmid construction, cell fixation, cell fractionation, antibodies

and immunological methods, microscopy, two-phase separation, and other assays are described in the

Supplemental Materials and Methods.

Construction of plasmids

A tobacco cDNA clone, BY28237 (GenBank accession no. LC128049.1) encoding the NtAGP1

protein (GenBank accession no. BAU61512) was used for the construction of plasmids used herein. The

fusion protein of GFP and NtAGP1, designated as GFP-AGP, was designed using this clone and a mutant GFP,

i.e., sGFP(S65T) (Chiu et al. 1996). For the construction of sporamin fusions, the coding sequence for the

signal peptide and the mature sporamin fusion construct (Δpro sporamin; Matsuoka and Nakamura 1991) was

used for the signal peptide-sporamin region. All of the expression constructs were subcloned into plant

expression vector pMAT137 (Yuasa et al. 2005) and used for transformation. The detailed methods used for

the construction of plasmids are described in the Supplemental Materials and Methods.

Culture, transformation, fixation, and immunostaining of tobacco BY-2 cells

The cultures and transformation of tobacco BY-2 cells and their transformants were maintained as

described (Tasaki et al. 2014). For immunostaining, BY-2 cells or its transformants were fixed and

permeabilized and incubated with primary and secondary antibodies essentially as described (Yuasa et al.

2005). Seven-day-old cells (stationary phase) were used in all of the experiments except when stated

otherwise in the figure legends.

Confocal microscopy and image analysis

The localisations of GFP-AGP and GFP-AGPΔC expressed in transformed BY-2 cells were visualized

using a confocal laser scanning microscope (TCS SP8, Leica, Wetzlar, Germany) essentially as described

(Tasaki et al. 2014). In some cases, Z-stack images were collected and converted to a rotating movie. To

visualize the plasma membrane using FM 4-64 or cytoplasm using chloromethyl SNARF-1, cells were

stained with either FM4-64 or chloromethyl SNARF-1 acetate as described (Yuasa et al. 2005). Confocal

images of the red fluorescence of stained cells were collected as described above with adjustment of the

excitation and emission wavelengths.

Antibodies

Rabbit anti-mitochondrial porin antibserum was generated as described in the Supplemental Materials

and Methods. Antisera against SDS-denatured SPO (Matsuoka and Nakamura., 1991), affinity purified

anti-native SPO (Matsuoka et al. 1995), affinity purified anti-plant Sec61 (Yuasa et al. 2005), anti-V-PPase

(Toyooka et al. 2009), anti-PIP (Suga et al. 2001), and anti-GLMT1 (Liu et al. 2015) antibodies were used at

the appropriate dilutions. The monoclonal anti-AGP glycan antibodies LM2 and LM6 (Yates et al. 1996),

were purchased from PlantProbes (Leeds, UK). For the visualisation of the antigen-antibody complex by

immunoblotting or indirect immunostaining, Alexa Fluor 488 goat anti-rabbit IgG (H+L), Alexa Fluor 568

goat anti-rabbit IgG (H+L), Alexa Fluor 568 goat anti-mouse IgG (H+L), Alexa Fluor 568 goat anti-rat IgG

(H+L), Alexa Fluor 647 goat anti-rat IgG (H+L), and Alexa Fluor 647 goat anti-rat IgM (μ chain) were

purchased from Invitrogen (Eugene, OR, USA).

SDS-PAGE and detection of fusion proteins

For the estimation of the size and extent of modification of the fluorescent fusion proteins, the

proteins were separated by SDS-PAGE, and their fluorescence was recorded by direct scanning of the gel

using an image analyser (Typhoon 9600, GE Healthcare Bio-Science, Chicago, IL) as described (Tasaki et al.

2014). For the detection of SPO fusion proteins and AGP glycan, immunoblotting was carried out as

described (Oda et al. 2020) using appropriate primary and fluorescence-tagged secondary antibodies.

For the comparison of the reactivities of SPO-AGP and secreted SPO-AGPΔC against anti-glycan

antibodies, we performed an adjustment of the loading of SPO-AGP in the microsomes and secreted large

form of SPO-AGPΔC after semi-quantitative immunoblotting using anti-sporamin antibody. Thereafter the

volumes of the microsomes and the medium that gave nearly identical intensity were used for comparison by

immunoblotting using anti-glycan antibodies.

Immunoprecipitation

Immobilized anti-native sporamin was prepared as described (Matsuoka et al. 1995).

Immunoprecipitation was performed essentially as described by Matsuoka and Nakamura (1991), except that

the immobilized antibody was used instead of the anti-sporamin serum and protein A Sepharose.

Subcellular fractionation of cells

Cells and media were separated from the suspension culture by filtration. For the differential

centrifugation analyses, cells were mixed with buffer and disrupted using a Parr Cell Disruption Bomb (Parr

Instruments, Moline, IL). The disrupted suspension was subjected to differential centrifugation. The

precipitates after the 1,000g, 10,000g, and 100,000g centrifugations as well as the supernatant of the final

spin were designated as P1, P10, P100, and S fractions, respectively. For the fractionation of endomembrane

organelles and the plasma membrane, microsomes prepared from cells were separated by linear sucrose

density gradient centrifugation essentially as described by Matsuoka et al. (1997). Protoplasts were prepared

as described by Nagata et al. (1981). Vacuoplasts and cytoplasts were prepared from protoplasts as described

by Sonobe (1990).

Fractionation of GPI-anchored proteins by two-phase separation using Triton X-114

Precondensation of the nonionic detergent Triton X-114 and the fractionation of GPI-anchored

proteins were performed as described by Murata et al. (2012) with minor modifications. In some cases, 6.5 m

units of PI-PLC (phospholipase C, phosphatidylinositol-specific from *Bacillus cereus*) (Sigma-Aldrich, St.

Louis, MO) were included during the incubation.

Cycloheximide treatment

Twenty mL of the 7-day old SPO-AGP culture was transferred to a 100-mL Erlenmeyer flask, and 200

μL of 1 mM cycloheximide and was added to the flask. The resulting flask was shaken in the dark at 26°C at

130 rpm. At 0, 3, and 24 hr of incubation, 1 mL of the culture was collected from the flask. Cells were

collected by centrifugation, mixed with an equal amount of SDS-sample buffer, and lysed by sonication. After

heating, equal volumes of protein-solubilized solution were subjected to SDS-PAGE and immunoblotting. As

a control, water was used instead of cycloheximide solution.

Enzymes and protein assays

The α-mannosidase activity was measured as described (Boller and Kende, 1979) using *p*-nitrophenyl

phosphate as a substrate. Protein concentrations were measured using a DC protein assay kit (Bio-Rad,

Hercules, CA).

Accession number

NtAGP1 cDNA sequence: LC128049.

Funding information

This work was supported in part by a grant from the Japan Society for the Promotion of Science (JSPS)

KAKENHI (no. 26292194 to K.M.).

Acknowledgments

We thank Dr. Koji Yuasa of the RIKEN Plant Science Center for the production of recombinant mitochondrial

porin and Mr. Nweke A. Boniface for improving the manuscript.

Author contributions

YT, RT, and KM performed the experiments on GFP fusion proteins. DN, YS, MF, and KM performed the

experiments on sporamin-fusion proteins. DN, YT, YS, and KM designed the experiments and analysed the

data. KM conceived the project and wrote the article with contributions from all of the authors.

References

Abiodun, M.O. and Matsuoka, K. (2013) Evidence that proliferation of golgi apparatus depends on both de
novo generation from the endoplasmic reticulum and formation from pre-existing stacks during the
growth of tobacco BY-2 cells. Plant Cell Physiol 54: 541-554.

Basu, D., Liang, Y., Liu, X., Himmeldirk, K., Faik, A., Kieliszewski, M., et al. (2013) Functional
identification of a hydroxyproline-o-galactosyltransferase specific for arabinogalactan protein
biosynthesis in Arabidopsis. J Biol Chem 288: 10132-10143.

Basu, D., Wang, W., Ma, S., DeBrosse, T., Poirier, E., Emch, K., et al. (2015) Two Hydroxyproline
Galactosyltransferases, GALT5 and GALT2, Function in Arabinogalactan-Protein Glycosylation,
Growth and Development in Arabidopsis. PLoS One 10: e0125624.

Beihammer, G., Maresch, D., Altmann, F. and Strasser, R. (2020) Glycosylphosphatidylinositol-Anchor
Synthesis in Plants: A Glycobiology Perspective. Front Plant Sci 11: 611188.

Bernat-Silvestre, C., De Sousa Vieira, V., Sanchez-Simarro, J., Pastor-Cantizano, N., Hawes, C., Marcote,
M.J., et al. (2020) p24 Family Proteins Are Involved in Transport to the Plasma Membrane of
GPI-Anchored Proteins in Plants. Plant Physiol 184: 1333-1347.

Bernat-Silvestre, C., Sánchez-Simarro, J., Ma, Y., Montero-Pau, J., Johnson, K., Aniento, F., et al. (2021)
AtPGAP1 functions as a GPI inositol-deacylase required for efficient transport of GPI-anchored
proteins. Plant Physiol 187: 2156-2173.

Boller, T. and Kende, H. (1979) Hydrolytic enzymes in the central vacuole of plant cells. Plant Physiol 63:
1123-1132.

Bordier, C. (1981) Phase separation of integral membrane proteins in Triton X-114 solution. J Biol
Chem 256: 1604-1607.

Bradley, D., Wood, E., Larkins, A., Galfre, G., Butcher, G. And Brewin, N. (1988) Isolation of
monoclonal-antibodies reacting with peribacteroid membranes and other components of pea
root-nodules containing Rhizobium leguminosarum. Planta 173: 149-160.

Chiu, W., Niwa, Y., Zeng, W., Hirano, T., Kobayashi, H. and Sheen, J. (1996) Engineered GFP as a vital
reporter in plants. Curr Biol 6: 325-330.

Darjania, L., Ichise, N., Ichikawa, S., Okamoto, T., Okuyama, H. and Thompson, G. (2002) Dynamic
turnover of arabinogalactan proteins in cultured Arabidopsis cells. Plant Physiology and
Biochemistry 40: 69-79.

De Caroli, M., Lenucci, M.S., Di Sansebastiano, G.P., Dalessandro, G., De Lorenzo, G. and Piro, G. (2011)
Protein trafficking to the cell wall occurs through mechanisms distinguishable from default sorting in
tobacco. Plant J 65: 295-308.

Drakakaki, G., van de Ven, W., Pan, S., Miao, Y., Wang, J., Keinath, N.F., et al. (2012) Isolation and
proteomic analysis of the SYP61 compartment reveal its role in exocytic trafficking in
Arabidopsis. Cell Res 22: 413-424.

Du, H., Simpson, R.J., Moritz, R.L., Clarke, A.E. and Bacic, A. (1994) Isolation of the protein backbone of
an arabinogalactan-protein from the styles of Nicotiana alata and characterization of a corresponding
cDNA. Plant Cell 6: 1643-1653.

Habuchi, S., Tsutsui, H., Kochaniak, A.B., Miyawaki, A. and van Oijen, A.M. (2008) mKikGR, a monomeric
photoswitchable fluorescent protein. PLoS One 3: e3944.

Hong, E., Davidson, A.R. and Kaiser, C.A. (1996) A pathway for targeting soluble misfolded proteins to the
yeast vacuole. J Cell Biol 135: 623-633.

Hromadová, D., Soukup, A. and Tylová, E. (2021) Arabinogalactan Proteins in Plant Roots - An Update on
Possible Functions. Front Plant Sci 12: 674010.

Kang, B.H., Nielsen, E., Preuss, M.L., Mastronarde, D. and Staehelin, L.A. (2011) Electron tomography of
RabA4b- and PI-4Kβ1-labeled trans Golgi network compartments in Arabidopsis. Traffic 12:
313-329.

Lin, Z., Xie, F., Triviño, M., Zhao, T., Coppens, F., Sterck, L., et al. (2022) Self-incompatibility requires GPI
anchor remodeling by the poppy PGAP1 ortholog HLD1. Curr Biol 32: 1909-1923.e1905.

Lind, J.L., Bacic, A., Clarke, A.E. and Anderson, M.A. (1994) A style-specific hydroxyproline-rich
glycoprotein with properties of both extensins and arabinogalactan proteins. Plant J 6: 491-502.

Liu, J., Hayashi, K. and Matsuoka, K. (2015) Membrane topology of Golgi-localized probable
S-adenosylmethionine-dependent methyltransferase in tobacco (Nicotiana tabacum) BY-2
cells. Biosci Biotechnol Biochem 79: 2007-2013.

Liu, Y.S. and Fujita, M. (2020) Mammalian GPI-anchor modifications and the enzymes involved. Biochem
Soc Trans 48: 1129-1138.

Matsuoka, K., Higuchi, T., Maeshima, M. and Nakamura, K. (1997) A Vacuolar-Type H+-ATPase in a
Nonvacuolar Organelle Is Required for the Sorting of Soluble Vacuolar Protein Precursors in Tobacco
Cells. Plant Cell 9: 533-546.

Matsuoka, K. and Nakamura, K. (1991) Propeptide of a precursor to a plant vacuolar protein required for
vacuolar targeting. Proc Natl Acad Sci U S A 88: 834-838.

Matsuoka, K. and Nakamura, K. (1999) Large alkyl side-chains of isoleucine and leucine in the NPIRL
region constitute the core of the vacuolar sorting determinant of sporamin precursor. Plant Mol
Biol 41: 825-835.

Maurer, J.B., Bacic, A., Pereira-Netto, A.B., Donatti, L., Zawadzki-Baggio, S.F. and Pettolino, F.A. (2010)
Arabinogalactan-proteins from cell suspension cultures of Araucaria angustifolia. Phytochemistry 71:
1400-1409.

Murata, D., Nomura, K.H., Dejima, K., Mizuguchi, S., Kawasaki, N., Matsuishi-Nakajima, Y., et al. (2012)
GPI-anchor synthesis is indispensable for the germline development of the nematode Caenorhabditis
elegans. Mol Biol Cell 23: 982-995.

Muñiz, M. and Riezman, H. (2016) Trafficking of glycosylphosphatidylinositol anchored proteins from the
endoplasmic reticulum to the cell surface. J Lipid Res 57: 352-360.

Nagata, T., Okada, K., Takebe, I. and Matsui, C. (1981) Delivery of tobacco mosaic virus RNA into plant
protoplasts mediated by reverse-phase evaporation vesicles (liposomes). Molecular & General
Genetics 184: 161-165.

Norman, P., Wingate, V., Fitter, M. and Lamb, C. (1986) Monoclonal-antibodies to plant plasma-membrane
antigens. Planta 167: 452-459.

Oda, Y., Asatsuma, S., Nakasone, H. and Matsuoka, K. (2020) Sucrose starvation induces the degradation of
proteins in trans-Golgi network and secretory vesicle cluster in tobacco BY-2 cells. Bioscience
Biotechnology and Biochemistry 84: 1652-1666.

Ogawa-Ohnishi, M. and Matsubayashi, Y. (2015) Identification of three potent hydroxyproline
O-galactosyltransferases in Arabidopsis. Plant J 81: 736-746.

Oka, T., Saito, F., Shimma, Y., Yoko-o, T., Nomura, Y., Matsuoka, K., et al. (2010) Characterization of
endoplasmic reticulum-localized UDP-D-galactose: hydroxyproline O-galactosyltransferase using
synthetic peptide substrates in Arabidopsis. Plant Physiol 152: 332-340.

Paris, N., Rogers, S.W., Jiang, L., Kirsch, T., Beevers, L., Phillips, T.E., et al. (1997) Molecular cloning and
further characterization of a probable plant vacuolar sorting receptor. Plant Physiol 115: 29-39.

Parsons, H.T., Stevens, T.J., McFarlane, H.E., Vidal-Melgosa, S., Griss, J., Lawrence, N., et al. (2019)
Separating Golgi Proteins from Cis to Trans Reveals Underlying Properties of Cisternal
Localization. Plant Cell 31: 2010-2034.

Pereira, A.M., Pereira, L.G. and Coimbra, S. (2015) Arabinogalactan proteins: rising attention from plant
biologists. Plant Reprod 28: 1-15.

Poulsen, C.P., Dilokpimol, A. and Geshi, N. (2015) Arabinogalactan biosynthesis: Implication of
AtGALT29A enzyme activity regulated by phosphorylation and co-localized enzymes for nucleotide
sugar metabolism in the compartments outside of the Golgi apparatus. Plant Signal Behav 10:
e984524.

Poulsen, C.P., Dilokpimol, A., Mouille, G., Burow, M. and Geshi, N. (2014) Arabinogalactan
glycosyltransferases target to a unique subcellular compartment that may function in unconventional
secretion in plants. Traffic 15: 1219-1234.

Putoczki, T.L., Pettolino, F., Griffin, M.D., Möller, R., Gerrard, J.A., Bacic, A., et al. (2007) Characterization
of the structure, expression and function of Pinus radiata D. Don arabinogalactan-proteins. Planta 226:
1131-1142.

Sonobe, S. (1990) Cytochalasin B cytokinetic cleavage in miniprotoplast isolated from cultured tobacco
cells. Protoplasma 155: 239-242.

Shimizu, M., Igasaki, T., Yamada, M., Yuasa, K., Hasegawa, J., Kato, T., et al. (2005) Experimental
determination of proline hydroxylation and hydroxyproline arabinogalactosylation motifs in secretory
proteins. Plant J 42: 877-889.

Showalter, A.M. and Basu, D. (2016) Extensin and Arabinogalactan-Protein Biosynthesis:
Glycosyltransferases, Research Challenges, and Biosensors. Front Plant Sci 7: 814.

Showalter, A.M., Keppler, B., Lichtenberg, J., Gu, D. and Welch, L.R. (2010) A bioinformatics approach to
the identification, classification, and analysis of hydroxyproline-rich glycoproteins. Plant Physiol 153:
485-513.

Silva, J., Ferraz, R., Dupree, P., Showalter, A.M. and Coimbra, S. (2020) Three Decades of Advances in
Arabinogalactan-Protein Biosynthesis. Front Plant Sci 11: 610377.

Singh, H., Ansari, H.R. and Raghava, G.P. (2013) Improved method for linear B-cell epitope prediction
using antigen's primary sequence. PLoS One 8: e62216.

Steffan, W., Kováčč, P., Albersheim, P., Darvill, A. and Hahn, M. (1995) Characterization of a monoclonal
antibody that recognizes an arabinosylated (1-]6)-beta-D-galactan epitope in plant complex
carbohydrates. Carbohydrate Research 275: 295-307.

Suga, S., Imagawa, S. and Maeshima, M. (2001) Specificity of the accumulation of mRNAs and proteins of
the plasma membrane and tonoplast aquaporins in radish organs. Planta 212: 294-304.

Tan, L., Qiu, F., Lamport, D.T. and Kieliszewski, M.J. (2004) Structure of a hydroxyproline
(Hyp)-arabinogalactan polysaccharide from repetitive Ala-Hyp expressed in transgenic Nicotiana
tabacum. J Biol Chem 279: 13156-13165.

Tan, L., Varnai, P., Lamport, D.T., Yuan, C., Xu, J., Qiu, F., et al. (2010) Plant O-hydroxyproline
arabinogalactans are composed of repeating trigalactosyl subunits with short bifurcated side chains. J
Biol Chem 285: 24575-24583.

Tasaki, M., Asatsuma, S. and Matsuoka, K. (2014) Monitoring protein turnover during phosphate
starvation-dependent autophagic degradation using a photoconvertible fluorescent protein aggregate
in tobacco BY-2 cells. Front Plant Sci 5: 172.

Temple, H., Mortimer, J.C., Tryfona, T., Yu, X., Lopez-Hernandez, F., Sorieul, M., et al. (2019) Two
members of the DUF579 family are responsible for arabinogalactan methylation in Arabidopsis. Plant
Direct 3: e00117.

Toyooka, K., Goto, Y., Asatsuma, S., Koizumi, M., Mitsui, T. and Matsuoka, K. (2009) A mobile secretory
vesicle cluster involved in mass transport from the Golgi to the plant cell exterior. Plant Cell 21:
1212-1229.

Tryfona, T., Liang, H.C., Kotake, T., Tsumuraya, Y., Stephens, E. and Dupree, P. (2012) Structural
characterization of Arabidopsis leaf arabinogalactan polysaccharides. Plant Physiol 160: 653-666.

Uemura, T., Nakano, R.T., Takagi, J., Wang, Y., Kramer, K., Finkemeier, I., et al. (2019) A Golgi-Released
Subpopulation of the Trans-Golgi Network Mediates Protein Secretion in Arabidopsis. Plant
Physiol 179: 519-532.

Uemura, T., Suda, Y., Ueda, T. and Nakano, A. (2014) Dynamic behavior of the trans-golgi network in root
tissues of Arabidopsis revealed by super-resolution live imaging. Plant Cell Physiol 55: 694-703.

Velasquez, S.M., Ricardi, M.M., Poulsen, C.P., Oikawa, A., Dilokpimol, A., Halim, A., et al. (2015)
Complex regulation of prolyl-4-hydroxylases impacts root hair expansion. Mol Plant 8: 734-746.

Viotti, C., Bubeck, J., Stierhof, Y.D., Krebs, M., Langhans, M., van den Berg, W., et al. (2010) Endocytic and
secretory traffic in Arabidopsis merge in the trans-Golgi network/early endosome, an independent and
highly dynamic organelle. Plant Cell 22: 1344-1357.

Wilkop, T., Pattathil, S., Ren, G., Davis, D.J., Bao, W., Duan, D., et al. (2019) A Hybrid Approach Enabling
Large-Scale Glycomic Analysis of Post-Golgi Vesicles Reveals a Transport Route for
Polysaccharides. Plant Cell 31: 627-644.

Willats, W.G., Marcus, S.E. and Knox, J.P. (1998) Generation of monoclonal antibody specific to
(1-->5)-alpha-L-arabinan. Carbohydr Res 308: 149-152.

Xu, J., Tan, L., Lamport, D.T., Showalter, A.M. and Kieliszewski, M.J. (2008) The O-Hyp glycosylation
code in tobacco and Arabidopsis and a proposed role of Hyp-glycans in secretion. Phytochemistry 69:
1631-1640.

Xu, Z., Gao, Y., Gao, C., Mei, J., Wang, S., Ma, J., et al. (2022) Glycosylphosphatidylinositol anchor lipid
remodeling directs proteins to the plasma membrane and governs cell wall mechanics. Plant Cell.

Xue, H., Veit, C., Abas, L., Tryfona, T., Maresch, D., Ricardi, M., et al. (2017) Arabidopsis thaliana FLA4
functions as a glycan-stabilized soluble factor via its carboxy-proximal Fasciclin 1 domain. Plant
Journal 91: 613-630.

Yamauchi, N., Gosho, T., Asatuma, S., Toyooka, K., Fujiwara, T. and Matsuoka, K. (2013) Polarized
localization and borate-dependent degradation of the Arabidopsis borate transporter BOR1 in tobacco
BY-2 cells. F1000Res 2: 185.

Yates, E.A., Valdor, J.F., Haslam, S.M., Morris, H.R., Dell, A., Mackie, W., et al. (1996) Characterization of
carbohydrate structural features recognized by anti-arabinogalactan-protein monoclonal
antibodies. Glycobiology 6: 131-139.

Yeats, T.H., Bacic, A. and Johnson, K.L. (2018) Plant glycosylphosphatidylinositol anchored proteins at the
plasma membrane-cell wall nexus. J Integr Plant Biol 60: 649-669.

Yuasa, K., Toyooka, K., Fukuda, H. and Matsuoka, K. (2005) Membrane-anchored prolyl hydroxylase with
an export signal from the endoplasmic reticulum. Plant J 41: 81-94.

Zhou, K. (2022) The regulation of the cell wall by glycosylphosphatidylinositol-anchored proteins in. Front
Cell Dev Biol 10: 904714.

 Legends to figures

 Fig. 1. Expression of the GFP-AGP fusion protein and its mutant without a GPI-anchoring signal in tobacco

 BY-2 cells. **A:** Schematic representation of the GFP fusion constructs. **B:** Accumulation of GFP fusion

 proteins in the culture of transformed tobacco cells. Total cell lysates were prepared from cells from 3-day

 and 7-day cultures, then separated by SDS-PAGE, and the GFP fluorescence in the gel was recorded. Each

 lane contained 3.6 μg of protein. **C:** Fractionation of 7-day cultures and the distribution of the GFP fusion

 proteins. Each lane contained proteins corresponding to 40 μL of cell culture. M: medium fraction, CW: cell

 wall fraction, Me: membrane fraction from total cell lysate, So: soluble fraction from total cell lysate.

 Fig. 2. Confocal localisation study of GFP-AGP and GFP-AGPΔC in transformed tobacco BY-2 cells.

 Confocal image of 7-day-old transformed tobacco cells expressing GFP-AGP, GFP-AGPΔC, or

 non-transformed cells. *Left:* Confocal fluorescence image and (*right*) the corresponding DIC image. Bars: 30

 μm.

 Fig. 3. Comparison of the localisation of GFP-AGP and plasma membrane or cytosol markers. **A:**

 Comparison of GFP-AGP and plasma membrane marked with FM4-64. Cells expressing GFP-AGP were

 incubated for 5 min with FM 4-64, then the fluorescence of both GFP and FM 4-64 was recorded using

 CFLM. The intensities of the fluorescence signals along the *yellow line* in the merged image were quantified.

 Normalized (maximum as 100%) intensities are shown in the graph. **B:** Confocal image of cells expressing

 GFP-AGP after staining with FM4-64 and plasmolysis. Composite images from 23 Z-stack images

 encompassing 7.96 μm are shown for GFP-AGP and FM 4-64 signals. **C:** Comparison of GFP-AGP and

 cytoplasm marked with 5- (and 6-) chloromethyl SNARF-1. Cells expressing GFP-AGP were incubated with

 5- (and 6-) chloromethyl SNARF-1 acetate (SNARF) for 5 min; confocal images were collected, and the

 intensities of the signals were quantified as in panel A. Bars: 30 μm in A and C, and 10 μm in B.

Fig. 4. The expression, secretion, and subcellular localisation of the sporamin-AGP fusion protein and its

mutant lacking a GPI-anchoring signal in tobacco BY-2 cells. **A:** Schematic representation of the sporamin

fusion constructs. **B:** Expression and secretion of SPO-AGP and SPO-AGPΔC. Seven-day-old cultures (Cu)

of transformed BY-2 cells expressing SPO-AGP and SPO-AGPΔC and non-transformed BY-2 cells (WT)

were separated into cells (C) and medium (M). The proteins in these fractions, corresponding to 10 μL of cell

culture, were separated by SDS-PAGE, and sporamin-related polypeptides were detected by using an

anti-SDS-denatured sporamin antibody. *Closed arrowheads:* transgene-dependent polypeptides. **C:** Detection

of sporamin-related polypeptides after immunoprecipitation. Sporamin-related polypeptides were recovered

by immunoprecipitation using immobilized anti-native sporamin, and the recovered proteins were analysed

by immunoblotting as described in the Materials and Methods section. Each lane contained sporamin-related

polypeptides from 0.5 mL of 7-day-old culture. *Closed arrowheads:* the migration position of

transgene-dependent polypeptides detected in panel B. *Open arrowheads:* the calculated migration position of

non-glycosylated polypeptides. **D:** Subcellular fractionation study of SPO-AGP-expressing cells.

Fractionation of cells was carried out by differential centrifugation as described in the Materials and Methods,

and sporamin-related polypeptides in the fractions were recovered by immunoprecipitation, separated by

SDS-PAGE, and detected by immunoblotting. P1: the 1,000g precipitate containing cell walls and unbroken

cells, P10: the 10,000g precipitate containing most of the mitochondria, P100: the 100,000g precipitate

containing microsomes, S: the 100,000g supernatant containing soluble proteins in the cytoplasm, vacuoles

and periplasmic space. **E:** Subcellular fractionation study of SPO-AGPΔC-expressing cells. The subcellular

fractionation and detection of sporamin-related polypeptides were carried out as in panel D.

Fig. 5. Intracellular distribution of SPO-AGP. **A:** Microsomes were prepared from SPO-AGP-expressing cells

in the presence of Mg^{2+} or EDTA and subjected to isopycnic sucrose density gradient ultracentrifugation. The

resulting gradients were fractionated from the bottom into 17 fractions. The concentration of sucrose in the

gradient is shown at the top. The distribution of marker proteins was analysed by immunoblotting with

specific antibodies: P-ATPase and PIP for PM, Sec61 for the ER, GLMT1 for the Golgi apparatus, and

V-PPase for the vacuolar membrane. The distribution of SPO-AGP, which migrated close to the top of the gel,

was analysed by immunoblotting using an antibody against sporamin. *Middle panels:* the relative distribution

of SPO-AGP and marker proteins after quantification of the signals on blots. *Bottom panels:* the immunoblot

results. **B:** Immunostaining of SPO-AGP-expressing or non-transformed cells with anti-native SPO antibody

or anti-AGP glycan LM6. C: Colocalisation of SPO-AGP and anti-AGP glycan epitope.

SPO-AGP-expressing cells were stained with both anti-native SPO antibody and anti-AGP glycan LM6 with

appropriate secondary antibodies. *Left:* Confocal images of both stained and corresponding transmission

images. Intensities of the fluorescence signals along the *yellow line* in the merged images were quantified.

Normalized (maximum as 100%) intensities are shown in the graph.

Fig. 6. Distribution of different forms of SPO-AGPΔC in the cell. **A:** Distribution of different forms in cells

and protoplasts. Protoplasts were prepared from SPO-AGPΔC-expressing cells by digesting the cell walls as

described in the Materials and Methods. Proteins in cells and protoplasts were separated by SDS-PAGE, and

sporamin-related polypeptides were detected by immunoblotting. As a control, protoplasts were prepared

from non-transformed cells (WT), and cross-reactive polypeptides against anti-sporamin were analysed by

immunoblotting. Each lane contained 25 μg of protein. *Closed arrowheads:* the migration position of the

transgene-dependent polypeptides. *Asterisks:* a possible degradation product of sporamin that was

occasionally observed on immunoblots. **B:** The small form of SPO-AGPΔC was localized in the vacuoles. To

assess the localisation of the intermediate and small SPO-AGPΔC proteins, protoplasts were separated into

vacuoplasts and cytoplasts, and the distribution of SPO-AGPΔC proteins was analysed by immunoblotting. A

nearly equal amount of the activity of α-mannosidase, which is a vacuolar marker enzyme, was loaded in the

vacuoplast and protoplast lanes, and nearly equal amounts of proteins were loaded into the protoplast and

cytoplast lanes. As a fractionation control, mitochondrial porin was also detected by immunoblotting. **C:** The

distribution of the intermediate form of SPO-AGPΔC within endomembrane organelles. Microsomes were

prepared from SPO-AGPΔC-expressing cells in the presence of Mg^{2+} (*left*) or EDTA (*right*), subjected to

isopycnic sucrose density gradient ultracentrifugation, fractionated, and analysed for the distribution of

intermediate SPO-AGPΔC and marker proteins as described in the Fig. 5A legend.

Fig. 7. Both GFP-AGP and SPO-AGP are GPI-anchored proteins. **A:** Triton X-114 two-phase partition assay

of GFP-AGP, GFP-AGPΔC and PIP-GFP. Microsomes from GFP-AGP- or PIP-GFP-expressing cells and the

culture medium of GFP-AGPΔC-expressing cells was subjected to the two-phase separation assay as

described in the Materials and Methods. Proteins in these two phases were separated by SDS-PAGE, and the

fluorescence of GFP in the gel was recorded. Each lane contained proteins corresponding to an equal amount

of microsomes or medium. aqu: aqueous phase, det: detergent phase. **B:** Triton X-114 two-phase partition

assay of SPO-AGP and SPO-AGPΔC. Two-phase partition was carried out as described in the legend for

panel A. Proteins were separated by SDS-PAGE and detected by immunoblotting using anti-sporamin

antibody. **C:** PI-PLC digestion of GFP-AGP. The two-phase partition assay was carried out in the absence and

presence of PI-PLC. **D:** Two-phase partition assay of SPO-AGP. Microsomal proteins from

SPO-AGP-expressing cells were subjected to the second-round two-phase partition assay in the absence or

presence of PI-PLC. Both SPO-AGP and V-PPase, which is an integral membrane protein, were detected by

immunoblotting. Each lane in these figures contained proteins corresponding to an equal amount of

microsomes or medium.

Fig. 8. The large form of SPO-AGPΔC was specifically recognized by LM2 monoclonal antibody. **A:** The

detection by anti-sporamin antibody showed that nearly equal amounts of sporamin-fusion proteins were

loaded. **B:** Detection by the monoclonal antibody LM2, which recognizes a glycan epitope with b-linked

glucuronic acid. **C:** Detection by the monoclonal antibody LM6, which recognizes a glycan epitope

consisting of (1-5)-α-L-arabinosyl residues. AGP: microsomes from SPO-AGP-expressing cells, ΔC: culture

medium from SPO-AGPΔC expressing culture, WT: corresponding microsomes and medium from

non-transformed BY-2 cells with an equal amount of protein with either microsomes from SPO-AGP or

culture medium from SPO-AGPΔC.

Fig. 9. Proposed model of the transport and localisation of native NtAGP1 and its mutant lacking the

GPI-anchoring signal.

887
888
889

A

Signal peptide

GFP

GFP-AGP

GPI-anchoring signal

GFP-AGPΔC

PIP-GFP

B

	GFP		GFP-AGP		GFP-AGPΔC		PIP-GFP	
days	3	7	3	7	3	7	3	7

(kD)

250 —
150 —

100 —

50 —

37 —

C

	GFP-AGP				GFP-AGPΔC				PIP-GFP			
(kD)	M	CW	Me	So	M	CW	Me	So	M	CW	Me	So

250 —
150 —
100 —

50 —

37 —

GFP fluorescence | Transmission

GFP-AGP

GFP-AGP△C

Non-transformed
BY-2

A

B

C

D

E

A

Marker | SPO-AGPΔC | WT

[kDa]

P C   P C

75 —

25 —

*

B

SPO-AGPΔC

V P Cy

[kDa]

75 —

anti-sporamin

25 —

*

anti-Mt porin

WT

V P Cy

[kDa]

75 —

25 —

| α-Mannosidase (× 10^{-11} mol min^{-1}) | 2.23 | 2.23 | |
| Protein (µg) | | 23.5 | 23.4 |

| α-Mannosidase (× 10^{-11} mol min^{-1}) | 3.34 | 3.33 | |
| Protein (µg) | | 24.2 | 24.2 |

C

+MgCl$_2$

sucrose (w/w%)
relative intensity

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17

+EDTA

sucrose (w/w%)
relative intensity

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17

Fraction number

- ● SPO-AGPΔC
- ▲ Sec61
- ■ V-PPase
- □ GLMT1
- ● P-ATPase
- ○ PIP

SPO-AGPΔC
Sec61 (ER)
V-PPase (tonoplast)
GLMT1 (Golgi)
P-ATPase (PM)
PIP (PM)

A

Marker | microsome | medium
AGP | WT | ΔC | WT

75 —

anti-SPO

B

Marker | microsome | medium
AGP | WT | ΔC | WT

250 —
150 —
100 —
75 —

LM2

C

Marker | microsome | medium
AGP | WT | ΔC | WT

250 —
150 —
100 —
75 —

LM6